

CSN 2018 Site Report: Allen Park

AQS ID: 26-163-0001, POC 5 (42.2286, -83.2082) 1-in-3 Day Schedule

The Chemical Speciation Network (CSN) is a routine air monitoring network designed to complement the PM_{2.5} monitoring network; support the implementation of PM_{2.5} National Ambient Air Quality Standards (NAAQS); assist in developing and tracking emission control strategies; and provide data to aid in health studies. CSN sites are primarily located in urban areas and complement the largely rural Interagency Monitoring of PROtected Visual Environments (IMPROVE) network. The CSN target analytes are trace elements, ions, and carbon.


Percent of Samples Successfully Collected and Analyzed Per Year

2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018
85	91	97	96	96	94	99	98	97	99	98	99	97	98	99	96	99	95


Samples Successfully Collected and Analyzed in 2018 by Filter Type. PTFE: 116 (95.1%), Nylon: 112 (91.8%), Quartz: 120 (98.4%)

The plots below show temporal trends for site 26-163-0001 alongside network-wide CSN and IMPROVE average concentrations. The top plot shows the variability of the reconstructed fine mass (RFM) concentrations during 2018; RFM can only be calculated if all three filters collected on a sampling day are valid. The bottom plot illustrates the long-term trends of ambient concentrations; the gray shaded region represents the range of values measured each year at this site, illustrated using the 10th and 90th percentile values.

Daily Reconstructed Fine Mass in 2018



Long-Term Trends in Reconstructed Fine Mass Missing years are due to low number of RFM values.



More Information


To view and download CSN data: www3.epa.gov/airquality/airdata/

The EPA website with guidance documents and background information: https://www.epa.gov/amtic/chemical-speciation-network-csn EPA real-time air monitoring data: https://www.epa.gov/outdoor-air-quality-data

The Univ. of California, Davis website with information about current research and publications: https://aqrc.ucdavis.edu/csn The Colorado State Univ. website with data resources, literature, and visibility overviews: http://vista.cira.colostate.edu/improve/



The following plots summarize the chemical composition of particles collected at this site. The monthly averaged compositions calculated from 2014-2018 data are shown on the left while compositions for the day with the highest measured concentrations during 2018 are shown on the right.



Components	Calculation	Natural Sources	Anthropogenic Sources
Salt	$1.8 \cdot Chloride$	Ocean spray, dry lakebeds	Chemical manufacturing, lake consumption
Soil Dust	$2.2 \cdot Al + 2.49 \cdot Si + 1.63 \cdot Ca$	Soil resuspension, dust storms	Construction, agriculture, deforestation,
	$+2.42 \cdot Fe + 1.94 \cdot Ti$	long-range transport	unpaved roads
Soot	$Elemental\ Carbon$	Wildfires	Motor vehicles, wood burning, smoking
Organic Matter	$1.4 \cdot Organic\ Carbon$	Plants, animals, wildfires	Motor vehicles, cooking oils, household cleaners
Nitrate	$1.29 \cdot Nitrate$	Plants, animals	Fertilizer, stock yards, chemical manufacturing
Sulfate	$4.125 \cdot Sulfur$	Volcanism	Coal-fired power plants, chemical manufacturing

The following map shows the average RFM concentrations for nearby sites in both CSN and the rural IMPROVE Network. The point shapes indicate which network the sites are associated with. The color bar indicates the average annual RFM concentration (micrograms per cubic meter) measured at each site in 2018.

