

## CSN 2018 Site Report: Jerome Mack Middle School AQS ID: 32-003-0540, POC 5 (36.141895, -115.078725) 1-in-3 Day Schedule


The Chemical Speciation Network (CSN) is a routine air monitoring network designed to complement the  $PM_{2.5}$  monitoring network; support the implementation of  $PM_{2.5}$  National Ambient Air Quality Standards (NAAQS); assist in developing and tracking emission control strategies; and provide data to aid in health studies. CSN sites are primarily located in urban areas and complement the largely rural Interagency Monitoring of PROtected Visual Environments (IMPROVE) network. The CSN target analytes are trace elements, ions, and carbon.

Percent of Samples Successfully Collected and Analyzed Per Year


2010	2011	2012	2013	2014	2015	2016	2017	2018
90	87	99	92	96	97	95	94	89


Samples Successfully Collected and Analyzed in 2018 by Filter Type. PTFE: 121 (99.2%), Nylon: 121 (99.2%), Quartz: 84 (68.9%)

The plots below show temporal trends for site 32-003-0540 alongside network-wide CSN and IMPROVE average concentrations. The top plot shows the variability of the reconstructed fine mass (RFM) concentrations during 2018; RFM can only be calculated if all three filters collected on a sampling day are valid. The bottom plot illustrates the long-term trends of ambient concentrations; the gray shaded region represents the range of values measured each year at this site, illustrated using the 10<sup>th</sup> and 90<sup>th</sup> percentile values.



Long-Term Trends in Reconstructed Fine Mass





## More Information


To view and download CSN data: www3.epa.gov/airquality/airdata/

The EPA website with guidance documents and background information: https://www.epa.gov/amtic/chemical-speciation-network-csn EPA real-time air monitoring data: https://www.epa.gov/outdoor-air-quality-data

The Univ. of California, Davis website with information about current research and publications: https://aqrc.ucdavis.edu/csn The Colorado State Univ. website with data resources, literature, and visibility overviews: http://vista.cira.colostate.edu/improve/



The following plots summarize the chemical composition of particles collected at this site. The monthly averaged compositions calculated from 2014-2018 data are shown on the left while compositions for the day with the highest measured concentrations during 2018 are shown on the right.



Components	Calculation	Natural Sources	Anthropogenic Sources
Salt Soil Dust	$\begin{array}{c} 1.8 \cdot Chloride \\ 2.2 \cdot Al + 2.49 \cdot Si + 1.63 \cdot Ca \\ + 2.42 \cdot Fe + 1.94 \cdot Ti \end{array}$	Ocean spray, dry lakebeds Soil resuspension, dust storms	Chemical manufacturing, lake consumption Construction, agriculture, deforestation, unpaved roads
Soot Organic Matter	$Elemental\ Carbon$ 1.4 · Organic Carbon	long-range transport Wildfires Plants, animals, wildfires	Motor vehicles, wood burning, smoking Motor vehicles, cooking oils, household cleaners
Nitrate Sulfate	$\begin{array}{c} 1.29 \cdot Nitrate \\ 4.125 \cdot Sulfur \end{array}$	Plants, animals Volcanism	Fertilizer, stock yards, chemical manufacturing Coal-fired power plants, chemical manufacturing

The following map shows the average RFM concentrations for nearby sites in both CSN and the rural IMPROVE Network. The point shapes indicate which network the sites are associated with. The color bar indicates the average annual RFM concentration (micrograms per cubic meter) measured at each site in 2018.

