

GEOLOGIC SUMMARY

GEOLOGIC SETTING

The regional project area is located in the lower Ohio River Valley, and includes the confluence of the Green River with the Ohio River. The landscape of the map area is characterized by low-relief bedrock uplands separated by broad alluvial valleys. Although the area is south of the Pleistocene glacial limit, the Ohio River served as a major outlet for glacial meltwater and entrained sediment during glacial stages. Rapid accumulation of glacial outwash in the main Ohio River Valley and along the mouths of tributaries led to impoundment and extensive deposition of slackwater and lacustrine sediment in the tributary valleys. This lacustrine deposit has a complex and gradational transition with loess mantling adjacent uplands. The loess was primarily derived from the valley-bottom outwash. The uplands are underlain by Pennsylvanian relatively flat-lying, coal-bearing strata.

GEOLOGIC BEHAVIOR

The Quaternary deposits identified in the map area exhibit a wide range of grain size and geotechnical behaviors. Grain size distribution is one of the primary factors affecting the behavior of soils for geotechnical, hydrogeologic, and agricultural applications. The grain size distribution of unconsolidated sediments is dominantly controlled by the conditions under which the material was deposited. Low energy environments allow the deposition of fine-grained materials. High energy deposits limit deposition to only coarser grained materials. Eolian processes produce very well sorted (poorly graded) materials. Fluvial processes produce moderate sorting; colluvial processes produce poorly sorted deposits.

HAZARDS

Flooding is a nearly annual occurrence along the Ohio River. Floods in the late winter or early spring commonly inundate low-lying areas in the floodplain. Larger floods occur roughly every 10 to 20 years (eg. 1913, 1945, 1964, 1997), and cover parts of the low terraces. The maximum flood of record in the valley was in 1937, flooding river towns throughout the valley. Only structures on the highest outwash terraces and the lacustrine terrace (Qit) were spared flood damage. The impact of flooding is reflected in land-use patterns through the area. Older homes and businesses have survived on the lacustrine and high outwash terraces, and on the highest parts of low terraces (Qot1, Qot1a, Qot1g). Trailers and less expensively built homes are constructed on the low terraces. Only barns are found on the high parts of the floodplains (Qalfp). The floodplain and low parts of the low terraces are dominantly left to woodlands or used for row-crop agriculture. Most livestock husbandry in the alluvial valleys has been abandoned and is now restricted to upland areas above the 10- to 20-year flood zone. The low-relief lacustrine terrace is locally very poorly drained.

The silt soils that dominate the loess-mantled uplands are highly erodible. Soil piping and associated cover collapses are common hazards as ground water seeps through the silt and is commonly perched above fraggans. Great care must be taken during agricultural operations not to mobilize and lose this valuable resource.

The map area is proximal to the Wabash Valley Seismic Zone and the New Madrid Seismic Zone. Small earthquakes have been felt in the area relatively frequently. The significant thicknesses of unconsolidated sediment (locally as much as 140 feet in the regional map area) raise concerns about ground motion amplification of seismic waves and potential liquefaction. The variations in lithology and thickness between materials in different map units will likely cause different responses of these materials to seismic shaking.

EXPLANATION

QGS database, number indicate depth to bedrock in feet
S Landform observation and soil probe
B Landform observation

DISCLAIMER

Although these data have been processed successfully on a computer system at the Kentucky Geological Survey (KGS), no warranty, expressed or implied, is made by KGS regarding the utility of the data on any other system, nor shall the act of distribution constitute any such warranty.

KGS does not guarantee this map or digital data to be free of errors or inaccuracies. Some cultural features originate from data sources other than KGS, and may not align with geologic features on this map. KGS disclaims any responsibility or liability for interpretations from this map or digital data, or decisions based thereon.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

ACKNOWLEDGMENTS

This map was generated using new field mapping and compilation of unpublished and previously published data and was funded in part by the U.S. Geological Survey National Cooperative Mapping Program under the STATEMAP Program authorized by the National Geographic Mapping Act of 1992, Grant No. 06HQPA0003, and by the Kentucky Geological Survey.

Field mapping was completed by Michael L. Murphy from April 2006 to December 2006, with assistance from W. Andrews, R. Counts, M. Crawford, S. Martin, and S. Waininger (KGS).

Subsurface information was compiled from data on file at the Kentucky Geological Survey as well as data contributed by the Kentucky Transportation Cabinet and the U.S. Geological Survey.

REFERENCES


Converse H.T., Jr., and Cox, F.R., Jr., 1967, Soil survey of Henderson County, Kentucky. U.S. Department of Agriculture, 108p.

Jacobs, E.H., 1981, Soil survey of Union and Webster counties, Kentucky. U.S. Department of Agriculture, Soil Conservation Service, 126p.

Ray, L.L., 1965, Geomorphology and Quaternary geology of the Owensboro quadrangle, Indiana and Kentucky. U.S. Geological Survey Professional Paper 488, 31p.

Solis, M.P., and Venard, E.A., 2000, Spatial database of the Poole quadrangle, western Kentucky. Kentucky Geological Survey, ser. 12. Digitally Vectorized Geologic Quadrangle Data DVGQ-1088. Adapted from Fairer, G.M., 1972, Geologic map of the Poole quadrangle, western Kentucky. U.S. Geological Survey Geologic Quadrangle Map GQ 1088, scale 1:24,000.

Thies, C.V., 1929, The Geology of Henderson County, Kentucky. University of Cincinnati, 251p.



**QUATERNARY GEOLOGIC MAP OF THE POOLE
7.5-MINUTE QUADRANGLE, WESTERN KENTUCKY**
By Michael L. Murphy
2007