


CORRELATION OF MAP UNITS



DESCRIPTION OF MAP UNITS

- Qal Alluvium, modern (Holocene)**
Silty clay and sandy silt with minor sand and sparse gravel; thickness 10 to 30 feet (3 to 10 m); found along river banks and in floodplains of smaller streams; deposited by modern/historic stream processes; deposit is inset into adjacent map units; contact with adjacent units varies from sharp to poorly defined; locally inferred on the basis of topographic expression. Some streams in the mapped area have been rerouted for land-use purposes; locally, some Qal dredged from these streams has been extensively redistributed across adjacent fields and is unmappable.
- Qao Alluvium, natural levee deposits (Holocene)**
Sand and silt; deposited by levees; ridges or overwash terraces (Qot1); grades into adjacent floodplain deposits; typically sandier than adjacent floodplain deposits.
- Qas Alluvium, active modern floodplain sloughs (Holocene)**
Organic-rich, black and gray clayey silt, silty clay, and clay; found within low lying areas on floodplain (Qafp) and low overwash terrace (Qot1); serve as poorly drained pathways which channel water from the floodplain; areas that retain water year-round form bogs and cypress swamps.
- Qaf Alluvium, alluvial fans (Holocene)**
Silt, sand, and gravel; thickness uncertain; forms fan-shaped alluvial-colluvial aprons at mouths of small valleys; deposited by floods and debris flows from small tributary valleys developed in loess-mantled uplands; extent of unit mapped by topographic expression.
- Qc Colluvium (Holocene)**
Silt, sand, clay, and rock fragments; unsorted; which has been transported downslope under the influence of gravity; primarily mantles steep slopes.
- Qafp Alluvium, river floodplains (Holocene)**
Sand, silt, fine gravel, and clay; surface mantled by silty clay and sandy silt; surface forms the lowest well-developed terrace along major rivers; 30 to 45 feet (10 to 15 m) thick; overlies older unconsolidated deposits or bedrock; contact is sharp, drawn at scarp of next higher terrace; estimated to range in age up to 6,500 years.
- Qas1 Alluvium, abandoned Green River meander (Holocene)**
Organic-rich, black and gray clayey silt, silty clay, and clay; deposited within recently abandoned meander of Green River; can retain standing water for months; areas that retain water year-round form bogs and cypress swamps.
- Qat Alluvium, low terrace (Holocene)**
Silt, sand, and clay deposited by rivers; forms terrace above adjacent floodplain (Qafp); contact with adjacent units varies from sharp to poorly defined; locally inferred on the basis of topographic expression; distinguished by topographic expression from lower floodplain (Qafp), but found below Ohio River low overwash terrace (Qot1) and lacustrine terrace (Qlt).
- Qas2 Alluvium, abandoned Green River channel (Pleistocene - Holocene)**
Clayey silt, silty sand, and silty clay; 30 to 45 feet (10 to 15 m) thick; forms arcuate, low-lying trough; represents an abandoned channel of Green River as it migrated across the low terrace (Qot1g); overlies older overwash deposits (Qot2); contact sharp, identified by surface topography; floods frequently.
- Qot10 Alluvium, reworked outwash, Ohio River scrollwork terrace (Pleistocene - Holocene)**
Fine to coarse sand and gravel, with local lenses of silt and clay; gravel includes chert, quartzite, sandstone, siltstone, igneous and metamorphic rocks, limestone, and coal; lithologically similar to adjacent outwash terraces; surface mantled with alluvial silty sand and sandy silt; 30 to 45 feet (10 to 15 m) thick; surface forms well-developed, well-sorted topography on Ohio River valley; overlies older overwash deposits (Qot2); contact is approximate, inferred from surface topography.
- Qas3 Alluvium, abandoned Green River channel (Pleistocene - Holocene)**
Silty sand, clayey silt, and silty clay; 30 to 45 feet (10 to 15 m) thick; forms sinuous, low-lying trough (Katie Meadow slough); represents an abandoned channel of Green River as it migrated across the low terrace (Qot1g); overlies older overwash deposits (Qot2); contact sharp, identified by surface topography; floods frequently.

GEOLOGIC SUMMARY

GEOLOGIC SETTING
The regional project area is located in the lower Ohio River valley, near the confluence of the Wabash River with the Ohio River. The landscape of the map area is characterized by steep to low-relief bedrock uplands separated by broad alluvial valleys. Although the area is south of the Pleistocene glacial limit, the Ohio River served as a major outlet for glacial meltwater and entrained sediment during glacial stages. Rapid accumulation of glacial outwash in the main Ohio River valley and along the mouths of tributaries led to impoundment and extensive deposition of slackwater and lacustrine sediment in the tributary valleys. This lacustrine deposit has a complex and gradational transition with loess mantling adjacent uplands. The loess was primarily derived from the valley-bottom outwash. The uplands are underlain by Upper Mississippian carbonates and sands and Lower Pennsylvanian sands within the faulted Chalybeate Hills, and relatively flat-lying Upper Pennsylvanian coal-bearing strata.

GEOTECHNICAL BEHAVIOR
The Quaternary deposits identified in the map area exhibit a wide range of grain-size and geotechnical behaviors. Grain-size distribution is one of the primary factors affecting the behavior of soils for geotechnical, hydrogeologic, and agricultural applications. The grain size distribution of unconsolidated sediments is dominantly controlled by the conditions under which the material was deposited. Low-energy environments allow the deposition of fine-grained materials. High-energy environments limit deposition to only coarser grained materials. Eolian processes produce very well sorted (poorly graded) materials. Fluvial processes produce moderate sorting; colluvial processes produce poorly sorted deposits.

HAZARDS
Flooding is a nearly annual occurrence along the Ohio River. Floods in the late winter or early spring commonly inundate low-lying areas in the floodplain. Larger floods occur roughly every 10 to 20 years (e.g. 1913, 1945, 1964, 1997, 2008), and cover parts of the low terraces. The maximum flood of record in the valley was in 1937. Flooding river towns throughout the valley. Only structures on the highest overwash terraces and the lacustrine terrace (Qlt) were spared from flood damage. The impact of seasonal flooding is reflected in land-use patterns through the area. Older homes and businesses have survived on the lacustrine and high overwash terraces, and on the highest parts of low terraces. Trailers and low expensively built homes are constructed on the low terraces. Only barns are found on the high parts of the floodplain. The floodplain and low parts of the low terraces are dominantly left to woodlands or used for row-crop agriculture. Most livestock husbandry in the alluvial valleys has been abandoned and is now restricted to areas above the 10- to 20-year flood zone. The low-relief lacustrine terrace is locally very poorly drained.

The silt soils that dominate the loess-mantled uplands are highly erodible. Soil piping and associated cover collapses are common hazards as ground water seeps through the silt and is commonly perched above fragipans. Great care must be taken during agricultural operations not to mobilize and lose this valuable resource.

The map area is proximal to the Wabash Valley seismic zone and the New Madrid seismic zone. Small to moderate earthquakes (2008) have been felt in the area relatively frequently. The significant thicknesses of unconsolidated sediment (locally as much as 140 feet in the regional map area) raise concerns about ground motion amplification of seismic waves and potential liquefaction. The variations in lithology and thickness between materials in different map units will likely cause different responses of these materials to seismic shaking.

EXPLANATION

- Contact
- - - Approximate contact
- Inferred contact
- Fault
- Concealed fault
- 23 KGS database, number indicate depth to bedrock in feet
- KGS drilling
- Landform observation and soil probe
- Landform observation
- Abandoned sand, gravel, or clay pit

DISCLAIMER

Although these data have been processed successfully on a computer system at the Kentucky Geological Survey (KGS), no warranty, expressed or implied, is made by KGS regarding the utility of the data on any other system, nor shall the act of distribution constitute any such warranty.

KGS does not guarantee this map or digital data to be free of errors or inaccuracies. Some cultural features originate from data sources other than KGS, and may not align with geologic features on this map. KGS disclaims any responsibility or liability for interpretations from this map or digital data, or decisions based thereon.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

ACKNOWLEDGMENTS

This map was generated using new field mapping and compilation of unpublished and previously published data and was funded in part by the U.S. Geological Survey National Cooperative Mapping Program under the STATEMAP Program authorized by the National Geologic Mapping Act of 1992, Grant No. 07HQAG02, and by the Kentucky Geological Survey. Field mapping was completed by Scott Waininger & Ronald Counts from July 2007 to April 2008, with assistance from W. Andrews, M. Crawford, S. Martin, and M. Murphy (KGS). Subsurface information was compiled from data on file at the Kentucky Geological Survey as well as data contributed by the Kentucky Transportation Cabinet and the U.S. Geological Survey.

REFERENCES

Jacobs, E.H., 1981, Soil survey of Union and Webster counties, Kentucky: U.S. Department of Agriculture, Soil Conservation Service, 126p.

Jilton, W.R., 1943, The geology of Union County, Kentucky. An outline designed with special reference to oil and gas exploration: The Standard Printing Company, Inc., 83p.

Johnson, W.D., Jr., Smith, A.E., and Fairer, G.M., 1975, Geologic map of the Morganfield quadrangle, Union County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1269, scale 1:24,000.

Ray, L.L., 1965, Geomorphology and Quaternary geology of the Owensboro quadrangle, Indiana and Kentucky: U.S. Geological Survey Professional Paper 488, 71p.

Venard, E.A., and Solis, M.P., 2000, Spatial database of the Morganfield quadrangle, Union County, Kentucky: Kentucky Geological Survey, ser. 12. Digitally Vectorized Geologic Quadrangle Data DVQ-1269. Adapted from Johnson, W.D., Jr., Smith, A.E., and Fairer, G.M., 1975, Geologic map of the Morganfield quadrangle, Union County, Kentucky: U.S. Geological Survey Geologic Quadrangle Map GQ-1269, scale 1:24,000.

Universal Transverse Mercator projection, zone 16, North American Datum of 1927
Topographic base and cultural features are Kentucky Raster Graphics (KRG) from Kymartian.ky.gov/krgmaps/KRG of Morganfield Quadrangle

