

U. S. DEPARTMENT OF COMMERCE  
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION  
NATIONAL WEATHER SERVICE  
NATIONAL CENTERS for ENVIRONMENTAL PREDICTION

TECHNICAL NOTE

HIGH-RESOLUTION OCEAN SURFACE WIND ANALYSES  
USING SATELLITE DERIVED OCEAN SURFACE WINDS:  
ANALYSIS VALIDATION USING SYNTHETIC SATELLITE DATA

William H. Gemmill &  
Christopher A. Peters (\*)

ENVIRONMENTAL MODELING CENTER  
WASHINGTON, D. C.

THIS IS AN UNREVIEWED MANUSCRIPT, PRIMARILY  
INTENDED FOR INTERNAL EXCHANGE OF INFORMATION

(\*) Present Affiliation  
General Sciences Corporation  
Laurel, Maryland, 20707

---

OMB CONTRIBUTION NO. 147

JULY 1997

## OPC CONTRIBUTIONS

- No. 1. Burroughs, L. D., 1987: Development of Forecast Guidance for Santa Ana Conditions. National Weather Digest, Vol. 12 No. 1, 7pp.
- No. 2. Richardson, W. S., D. J. Schwab, Y. Y. Chao, and D. M. Wright, 1986: Lake Erie Wave Height Forecasts Generated by Empirical and Dynamical Methods -- Comparison and Verification. Technical Note, 23pp.
- No. 3. Auer, S. J., 1986: Determination of Errors in LFM Forecasts Surface Lows Over the Northwest Atlantic Ocean. Technical Note/NMC Office Note No. 313, 17pp.
- No. 4. Rao, D. B., S. D. Steenrod, and B. V. Sanchez, 1987: A Method of Calculating the Total Flow from A Given Sea Surface Topography. NASA Technical Memorandum 87799, 19pp.
- No. 5. Feit, D. M., 1986: Compendium of Marine Meteorological and Oceanographic Products of the Ocean Products Center. NOAA Technical Memorandum NWS NMC 68, 93pp.
- No. 6. Auer, S. J., 1986: A Comparison of the LFM, Spectral, and ECMWF Numerical Model Forecasts of Deepening Oceanic Cyclones During One Cool Season. Technical Note/NMC Office Note No. 312, 20pp.
- No. 7. Burroughs, L. D., 1987: Development of Open Fog Forecasting Regions. Technical Note/NMC Office Note. No. 323, 36pp.
- No. 8. Yu, T. W., 1987: A Technique of Deducing Wind Direction from Satellite Measurements of Wind Speed. Monthly Weather Review, 115, 1929-1939.
- No. 9. Auer, S. J., 1987: Five-Year Climatological Survey of the Gulf Stream System and Its Associated Rings. Journal of Geophysical Research, 92, 11,709-11,726.
- No. 10. Chao, Y. Y., 1987: Forecasting Wave Conditions Affected by Currents and Bottom Topography. Technical Note, 11pp.
- No. 11. Esteva, D. C., 1987: The Editing and Averaging of Altimeter Wave and Wind Data. Technical Note, 4pp.
- No. 12. Feit, D. M., 1987: Forecasting Superstructure Icing for Alaskan Waters. National Weather Digest, 12, 5-10.
- No. 13. Sanchez, B. V., D. B. Rao, and S. D. Steenrod, 1987: Tidal Estimation in the Atlantic and Indian Oceans. Marine Geodesy, 10, 309-350.
- No. 14. Gemmill, W. H., T. W. Yu, and D. M. Feit 1988: Performance of Techniques Used to Derive Ocean Surface Winds. Technical Note/NMC Office Note No. 330, 34pp.
- No. 15. Gemmill, W. H., T. W. Yu, and D. M. Feit 1987: Performance Statistics of Techniques Used to Determine Ocean Surface Winds. Conference Preprint, Workshop Proceedings AES/CMOS 2nd Workshop of Operational Meteorology, Halifax, Nova Scotia, 234-243.
- No. 16. Yu, T. W., 1988: A Method for Determining Equivalent Depths of the Atmospheric Boundary Layer Over the Oceans. Journal of Geophysical Research, 93, 3655-3661.
- No. 17. Yu, T. W., 1987: Analysis of the Atmospheric Mixed Layer Heights Over the Oceans. Conference Preprint, Workshop Proceedings AES/CMOS 2nd Workshop of Operational Meteorology, Halifax, Nova Scotia, 2, 425-432.
- No. 18. Feit, D. M., 1987: An Operational Forecast System for Superstructure Icing. Proceedings Fourth Conference Meteorology and Oceanography of the Coastal Zone. 4pp.
- No. 19. Esteva, D. C., 1988: Evaluation of Preliminary Experiments Assimilating Seasat Significant Wave Height into a Spectral Wave Model. Journal of Geophysical Research, 93, 14,099-14,105.
- No. 20. Chao, Y. Y., 1988: Evaluation of Wave Forecast for the Gulf of Mexico. Proceedings Fourth Conference Meteorology and Oceanography of the Coastal Zone, 42-49.

## 1. INTRODUCTION

The need for accurate high-resolution regional and coastal ocean surface wind analyses is increasing with the arrival of high-resolution ocean modeling capabilities for: waves, circulation (currents), and storm surges (water levels). Fortunately, high-resolution (or meso-scale) ocean surface wind data, from several satellite systems, are now available in sufficient quantity to provide such support. However, the science of converting parameters measured from satellites into ocean surface winds is far from exact. It is necessary to be able to assess the impact of data from various satellites on wind analyses; and, to determine how to blend the data from these different satellite systems to produce accurate meso-scale ocean surface wind fields.

Currently, most ocean surface wind fields are available from global data assimilation systems. Even when satellite surface wind data are used in these analyses, the satellite data are made into "super-obs" at the scale of the model. A "super-ob" is simply the average of observations within the immediate neighborhood of a model grid point. But since the model resolution is less than that of the observation, the meso-scale wind information is lost through this process. Mullen (1994) compared objectively generated surface analyses (NCEP global model) over the North Pacific ocean with subjectively drawn analyses to show that the objective analyses not only misplace the locations of storm systems but also consistently underestimate their intensity. Therefore, an analysis technique is developed and tested to provide objective analyses of high-resolution ocean surface wind fields over selected U. S. coastal regions. This method is based on a reanalysis procedure in an attempt to regain some additional information from the observations which has been lost in the larger-scale global analysis system. This analysis is an extension of an earlier ocean surface wind analysis program (Gemmill, 1991). Those wind analyses were initially based on the reanalysis of real-time ocean surface wind data using ships, fixed buoys and coastal stations, on a high-resolution grid, with the initial wind analysis interpolated from the global data assimilation system (GDAS) at 2.5 X 2.5 latitude and longitude. This paper presents the methodology of the current technique which is based on a higher resolution initial grid of GDAS (at 1.0 X 1.0 latitude/longitude) and the incorporation of satellite data. An updated version of the analysis system was presented by Gemmill & Peters (1996).

Bumke and Hasse (1989) showed that it was possible to improve the resolution of a low-resolution global ocean surface wind analysis directly by reanalyzing surface observations of sea level pressure and surface wind. Sanders (1990) has pointed out that obtaining an accurate detailed sea level pressure analysis is no easy task, even when sufficient surface data (ERICA drifting buoys) are available. However, those studies were based on only direct surface measurements (buoys and ships), whereas today we have high-resolution (25km-50km) real-time wind data available over wide swaths from satellite borne sensors. The data now are available from four DMSP satellites using the SSM/I sensor to provide wind speed data across a 1500 km swath, and the ERS2 satellite using the scatterometer sensor to provide wind vectors across a 500 km swath.



## 2. THE DATA

In this report we will deal with the impact of synthetic satellite data on analyses to gain insights into the importance of the different satellite data sets. But, first we will review characteristics of the various data sets available for ocean surface weather analyses.

It is well known that in-situ ocean surface wind measurements are sparse. These data have been studied in some detail to determine their accuracy by Wilkerson and Earle (1990) and Pierson (1990). The basic conclusion of these studies is that ship data when compared to buoy data are not adequate for accurate analysis of the ocean wind field. In fact, whether the wind measurement was made by an anemometer or by visual procedures, the quality was about the same. Gilhousen (1987) reports that the buoy wind data, although not without some problems, are within the NDBC error specifications of 1 m/s or 10% for speed and 20 degrees for direction. Coastal stations can be used for analysis, but the exposure must be known. Some of these platforms sit over the water, while others may be on high cliffs at the water's edge. Unfortunately, there is no uniform height for wind measurements over the ocean. Buoy heights range from 3.8 m to 14m, and ships somewhat higher at 20m, but that height depends on the size of the ship and its "load", and the problems with coastal stations have already been noted.

The DMSP satellite (SSM/I) provides wind speeds over periodic 102 minute polar orbits in swaths of 1500 km and at a horizontal resolution of 25 km and at a height of 20m above the sea. It is a passive microwave instrument (Table 1a). In mid-latitudes the satellite will pass over the same region twice a day on its ascending and descending passes. There are four SSM/I satellites (F10, F11, F13 & F14) which are in similar orbits. The "standard" SSM/I wind algorithm, which converts brightness temperatures to wind speed through linear regression was calibrated to meet the error requirement for speed of 2 m/s or 10% (Goodberlet, Swift and Wilkerson, 1989). But, these SSM/I wind speed data have two important limitations: winds cannot be measured 1) above 20 m/s and 2) for moderate moisture and rain events. Hence, an "all-weather" algorithm has been developed using the non-linear approach of neural networks (Krasnopolsky, Gemmill & Breaker, 1996). This algorithm gives improved wind speeds over the 5-25 m/s range for a wide range of moisture conditions.

The ERS2 satellite (with its scatterometer) provides wind vector data over periodic 102 minute polar orbits in swaths of 500 km, at a horizontal resolution of 50 km and at a height of 10m above the sea. This is an active microwave instrument (Table 1b). In mid-latitudes the satellite will pass over the same region twice a day on its ascending and descending passes. The "standard" scatterometer algorithm (which converts radar backscatter to wind speed through empirical functions) was calibrated to meet the error requirement for speed of 2 m/s or 10%, and for wind direction of 20 degrees. NCEP has been receiving the "fast-delivery" wind data from ESA in real-time for operational use. These wind speeds are of sufficient accuracy, but, the wind directions were found to lacking in accuracy when compared with buoys data. Since the raw radar backscatter data were also provided from ESA, NCEP processes these data to re-derive the wind speeds and directions (Peters et al, 1994) with substantial improvement in accuracy to the direction (Gemmill et al, 1994). In that study, we also compared various proposed transfer functions and showed that the transfer function (CMOD4) used by ESA was best for providing wind speeds.



Table 1. Satellite Specifications for SSM/I and Scatterometer Derived Ocean Surface Winds.

SATELLITE OCEAN SURFACE WIND SPECIFICATIONS

- POLAR ORBITER (102 MINUTES)

- TWICE DAILY AREAL COVERAGE  
(ONE ASCENDING & ONE DESCENDING PASS)

|                        | ERS1/2                                 | DMSP                                    |
|------------------------|--|---|
| SENSOR                 | SCATTEROMETER                          | SPECIAL SENSOR<br>MICROWAVE / IMAGER    |
| MODE                   | ACTIVE MICROWAVE<br>3 ANTENNA, 1-SIDED | PASSIVE MICROWAVE<br>BT's at 7 CHANNELS |
| DATA                   | WIND SPEED &<br>DIRECTION              | WIND SPEED                              |
| SWATH<br>No. OF POINTS | 500 km<br>19 CELLS                     | 1492 km<br>64 CELLS                     |
| FOOTPRINT              | 50 km                                  | 40 km                                   |
| WIND HEIGHT            | 10 m                                   | 20 m                                    |
| SPEED RANGE            | 4 -24 m/s                              | 3 - 25 m/s                              |
| SPEED ACCURACY         | +/- 2 m/s, for > 20 m/s (10%)          | +/- 2 m/s, for > 20 m/s (10%)           |
| DIRECTION ACCURACY     | +/- 20 deg                             | N / A                                   |

3. THE ANALYSIS PROCEDURE

The first guess for the wind field is provided the NCEP Global Data Assimilation System (Derber, Parrish and Lord, 1991) on a 1.0 X 1.0 degree latitude/longitude grid. These winds are obtained from the midpoint of the lowest sigma layer of the model (about 45 m above the ocean surface), are interpolated to the fine mesh grid which was chosen to be ½ degrees in longitude and 1/3 degrees in latitude and are reduced to 20m using the neutral log-profile for a constant flux layer and the winds. The test region for this analysis was selected to be the Northwestern Atlantic Ocean adjacent to the U. S. east coast. The technique used to reanalyze the winds is based on a conditional relaxation method, which is applied as follows. At each step the wind field is separated into "u" and "v" components, and each component is analyzed independently. The Laplacian of the first guess field component is calculated to be used as the forcing function. Wind data, by components, are used to correct the wind field at the nearest grid point, which then are set as fixed internal grid points.

Winds at non-corrected internal ocean grid points are determined by numerical relaxation against the forcing function. A conditional relaxation method is applied in two steps; first, after the ERS2 scatterometer wind vector data have been incorporated into the analysis, and then after the SSMI wind speed data have been incorporated for the final analysis. The SSMI wind speed is decomposed into “u” and “v” components using the directions from the nearest four neighbors after the ERS2 winds have been taken into account. The conditional relaxation method corrects the analysis to the data where data exist and spreads the influence of the data to the rest of the field by preserving the second derivative (Laplacian) of the first guess field. The use of the Laplacian as the forcing function essentially preserves the original first guess shape of the field at grid points where no data were available. See appendix A for a description of the conditional relaxation procedure.

The conventional surface data of buoys, ships and c-man stations are withheld from the analysis. Buoys are to be used for validation only.

During the reanalysis process, the satellite data has corrected the first guess surface wind field to reflect the new data. But now one would suspect that the first guess surface pressure analysis will not necessarily be consistent (in balance) with the reanalyzed wind field. Therefore, a "new" sea level pressure analysis is generated through the use of the diagnostic “balance equation” with the inclusion of friction terms, using the reanalyzed winds. Table 2 summarizes the flow of the analysis procedures.

Table 2. Overview For the Sequence of Steps to Perform Satellite Wind Analyses

| Field or Data Source                               | Operation Step   |
|--|--|
| The First Guess Surface Wind                       | Interpolate Global Wind Analysis to High-Resolution Grid           |
| Satellite Data                                     | Scatterometer Wind Vector Data is Retrieved                        |
| Intermediate High-Resolution Surface Wind Analysis | Execute Wind Analysis Program                                      |
| Satellite Data                                     | SSM/I Wind Speed Data is Retrieved                                 |
| New High-Resolution Surface Wind Analysis          | Execute Wind Analysis Program                                      |
| Balanced Sea Level Pressure Analysis               | Apply The Balance Equation   |
| Buoy Data  | Perform Validation of Surface Wind and Sea Level Pressure Analysis |



## 4. ANALYSIS METHOD VALIDATION

### a) Synthetic Data Tests

“Synthetic” data fields of pressures and winds are constructed which are assumed to represent the “true” state of the atmosphere. These fields provide the basis for evaluating the analysis procedure described in the previous section. A sea level pressure field is generated first by designing a mathematical formulation, which contains large scale and small scale information. This is designated as the “true” pressure field as shown in figure 1. For this case study, the “true” pressure analysis is specified to have a deep storm system with a central pressure of 992.6 mb located near 38N & 67W. To determine the “true” surface wind field, the gradient wind field is computed, then rotated and reduced to simulate the frictional effects of the boundary layer. For this study the wind directions are rotated 15 degrees to the left (toward low pressure) from the gradient (and geostrophic) wind direction, and gradient speeds reduced by 30% (see figure 2).

Now, the “first guess” sea level pressure and surface wind fields are estimated. The “first guess” sea level pressure field is created by removing the small scale information from the mathematical formulation of the “true” sea level pressure analysis, and shifting the location of major pressure systems to simulate the loss of detail created by smoothing and typical errors of first guess fields as shown in figure 3. For the first guess, the intensity of the storm is reduced to a central pressure of 1001.4 mb, (an increase of almost 9 mb higher than the “true” central pressure), and displaced to near 41N & 65W (a distance of almost 350 km to the northeast of the the “true” location). To determine the “first guess” surface wind field, the same operations are performed as above to determine the “true” wind field; the gradient wind field is computed, then rotated and reduced to simulate the frictional effects of the boundary layer (see figure 4).

Consider now the problem of introducing satellite wind measurements. It is assumed that the satellite wind data are free of any errors and are simply extracted from the “true” wind field (figure 2) to simulate the passes of one scatterometer (SCAT 1) and two SSMI (SSMI 1 & SSMI 2) sensors over the region at the time of the “true” analysis of the satellites. The swath of wind vector data (SCAT 1, figure 5) passes directly over the center of the storm, whereas the one swath of wind speed data (SSMI 1, figure 6) also passed over the center of the storm the but other swath of wind speed data (SSMI 2) passes far to the northeast (figure 7). In reality the time window used in analyses for satellite or conventional data is typically +/- 3 hours centered at the analysis time. For the sake of this analysis it is assumed that there are no errors in the data due to time differences between the analyses and data. An ideal set of synthetic buoy data (figure 8) are also extracted from the true wind field, with locations near the coastal areas to simulate available ground truth used for evaluation of the analyses.

We now have constructed a first guess wind and pressure field, representing the “real world”. We also have perfect satellite wind data, which allow can be used to determine directly the impact of on the analysis. The analysis procedure (described in section 3) is then executed using the perfect synthetic satellite wind data to correct the first guess wind field, and a new surface wind analysis is created. But since the wind field has been altered with the satellite data, the original sea level pressure first guess field will no longer be in balance with the new wind field. The new surface winds



## APPENDIX A: CONDITIONAL RELAXATION

Each of the components (u and v) is treated separately. Initially, the forcing function (the Laplacian) is computed from the appropriate component of the first guess which was obtained from the global analysis interpolated to the high-resolution grid;

$$F = \nabla^2 U_{fg} \quad (1a)$$

where  $\nabla^2$  is the Laplacian operator,  $U_{fg}$  is the appropriate first guess component.

The wind observations are separated by components, then checked for gross errors against the first guess, and suspect data is discarded. The "good" observations are used to correct the nearest grid point. Data are averaged if there more than one observation. All non-data grid points are relaxed to determine new values of the appropriate component. Grid points that are external boundaries, land, or have been corrected by data will held fixed through the relaxation. This process is continued until the difference between the Laplacian of corrected wind field component and its forcing function is nearly zero. This is done by solving iteratively (a2) at all non-data grid points to reduce the residual (R):

$$R = \nabla^2 U_n - F \quad (a2)$$

A new value for U is obtained by using the residual, R, and a relaxation coefficient (a) as follows,

$$U_{n+1} = U_n + R/a \quad (a3)$$

so that after n iterations:

$$R < \epsilon, \text{ for small } \epsilon \quad (a4)$$

at all grid points.



## APPENDIX B. THE BALANCE EQUATION

The balance equation can be written as:

$$\rho \nabla^2 p = f \frac{\partial v}{\partial x} - \frac{\partial u}{\partial y} - u \frac{\partial f}{\partial y} - 2 \frac{\partial u}{\partial y} \frac{\partial v}{\partial x} - \frac{\partial u}{\partial x} \frac{\partial v}{\partial y}$$

Where;  $\rho$  = specific density and  $f$  = coriolis parameter

One can solve for  $p$  by solving the balance equation, given  $u$  &  $v$  components of a wind field, by Simultaneous Over-Relaxation (SOR) using the subroutine published by Press et al (1989).

$$\nabla^2 p = F(x, y)$$

where  $F$  (the forcing function) contains only terms  $\rho$ ,  $f$ ,  $u$  &  $v$



REFERENCES:

- Bumke, K., and L. Hasse, 1989: An Analysis Scheme for Determination of True Surface Winds at Sea from Ship Synoptic Wind and Pressure Observations. *Boundary-Layer Meteorology*, 47, 295-308.
- Derber, J., D. Parrish, D. F. and S. J. Lord 1991: The New Global Operational Analysis System at the National Meteorological Center. *Weather and Forecasting* 6, 538-547.
- Gemmill, W. H. and C.A. Peters, 1996: The Use of Satellite Derived Wind Data in High-Resolution Regional Ocean Surface Wind Fields. Conference on Coastal Oceanic and Atmospheric Prediction, Jan 28- Feb 2, 1996 Atlanta, GA., AMS, pp 397-400.
- Gemmill, W., M. Woiceshyn, P., Peters, C. and V. Gerald, 1994: A Preliminary Evaluation of Scatterometer Wind Transfer Functions for ERS-1 Data. Tech. Note, OPC Contib. 97, Nat. Met. Center, Washington, D.C., 35pp
- Gemmill, W. H. 1991; High Resolution Regional Ocean Surface Wind Fields. Ninth Conference On Numerical Weather Prediction. 14-18 Oct., Denver, Co., AMS, pp 190-191
- Gilhousen, D., 1987: A Field Evaluation of NDBC Moored Buoy Winds,. *J. Atmos Oceanic Technol* 4, 94-104.
- Goodberlet, M., Swift, C. and J. Wilkerson, 1990: Ocean Surface Wind Speed Measurements of the Special Sensor Microwave/Imager(SSM/I). *IEEE Transactions on Geoscience and Remote Sensing* 24, 823-828.
- Krasnoplosky, V. M., W. H. Gemmill and L. C. Breaker, 1996: A New Transfer Function for SSM/I Based on an Expanded Neural Network Architecture. Tech. Note, OMB Contr No. 137, Nat. Center's for Environ. Prediction, Washington, D.C., 38 pp.
- Mullen, S.I. 1994: An Estimation of Systematic Error and Uncertainty in Surface Cyclone Analysis over the North Pacific Ocean: Some Forecasting Implications. *Weather & Forecasting*, 9, 221-227.
- Peters, C., V Gerald, P Woiceshyn & W. Gemmill, 1994: Operational Reprocessing ERS1 Scatterometer Winds: A Documentation, Tech. Note, OPC Contr No. 96, Nat Met Center, Washington, D.C., 12pp.
- Pierson, W., 1990: Examples of, Reason for, and Consequences of the Poor Quality of Wind Data From Ships for the Marine Boundary Layer: Implications for Remote Sensing. *J. of Geophysical Research* 95, No. C8, 13,313-340.
- Press, W.H., B.P. Flannery, S.A. Teukolsky and W.T. Vetterling, 1989: *Numerical Recipes*, Cambridge Univ Press, Cambridge, pp 652-659.
- Sanders, F., 1990: Surface Analysis Over the Oceans - Searching for Sea Truth. *Weather and Forecasting* 5, 596-612.
- Wilkerson, J. and M. Earle, 1990: A Study of Differences Between Environmental Reports by Ships in the Voluntary Observing Program and Measurements from NOAA Buoys. *J. of Geophysical Research* 95, No. C3, 3373-3385.





TABLE 3. SYNTHETIC DATA TEST CASE STUDIES:

- Evaluation Statistics of Analysis Errors Based On Gridded True Fields.

CASE 1: Wind Vector data swath across center of true storm (figures 5, 9 & 10).

CASE 2: Wind Speed data swath across center of true storm (figures 6, 11 & 12).

CASE 3: Wind Speed data swath northeast of true storm center (figures 7, 13 & 14).

CASE 4: All wind data sets

| PERFECT FLD  | CASE 1 | CASE 2 | CASE 3 | CASE 4 |
|--------------|--------|--------|--------|--------|
|              |        |        |        |        |
| SPEED - BIAS | (m/s)  |        |        |        |
| FG - TRUE    | -0.89  | -0.89  | -0.89  | -0.89  |
| NEW - TRUE   | 0.01   | 0.30   | -0.80  | 0.18   |
| SPEED - RMS  | (m/s)  |        |        |        |
| FG - TRUE    | 2.33   | 2.33   | 2.33   | 2.33   |
| NEW - TRUE   | 1.20   | 0.76   | 2.16   | 0.67   |
|              |        |        |        |        |
| SLP - BIAS   | (mb)   |        |        |        |
| FG - TRUE    | 0.60   | 0.60   | 0.60   | 0.60   |
| NEW - TRUE   | -0.34  | -0.43  | 0.48   | -0.50  |
| SLP - RMS    |        | (mb)   |        |        |
| FG - TRUE    | 1.99   | 1.99   | 1.99   | 1.99   |
| NEW - TRUE   | 0.87   | 1.36   | 1.76   | 0.86   |
|              |        |        |        |        |
| LOWEST SLP   | (MB)   |        |        |        |
| FG           | 1001.4 | 1001.4 | 1001.4 | 1001.4 |
| NEW          | 992.9  | 989.5  | 1000.6 | 990.3  |
| TRUE         | 992.6  | 992.6  | 992.6  | 992.6  |
|              |        |        |        |        |



TABLE 4. SYNTHETIC DATA TEST CASE STUDIES:

- Evaluation Statistics of Analysis Errors Using Synthetic Buoy Data (figure 8).

CASE 1: Wind Vector data swath across center of true storm (figure 5, 9 & 10).

CASE 2: Wind Speed data swath across center of true storm (figure 6, 11 & 12).

CASE 3: Wind Speed data swath northeast of true storm center (figure 7, 13 & 14).

CASE 4. All wind data

| BUOY DATA        | CASE 1 | CASE 2 | CASE 3 | CASE 4 |
|------------------|--------|--------|--------|--------|
| SPEED - BIAS     | (m/s)  |        |        |        |
| FG - DATA        | -3.16  | -3.16  | -3.16  | -3.16  |
| NEW - DATA       | -0.40  | 0.64   | -3.27  | 0.48   |
| SPEED - RMS      | (m/s)  |        |        |        |
| FG - DATA        | 4.56   | 4.56   | 4.56   | 4.56   |
| NEW - DATA       | 2.17   | 1.31   | 4.89   | 1.34   |
|                  |        |        |        |        |
| DIRECTION - BIAS | (Deg)  |        |        |        |
| FG - DATA        | -2     | -2     | -2     | -2     |
| NEW - DATA       | -3     | 3      | 1      | -3     |
| DIRECTION - RMS  | (Deg)  |        |        |        |
| FG - DATA        | 20     | 20     | 20     | 20     |
| NEW - DATA       | 5      | 22     | 19     | 7      |
|                  |        |        |        |        |
| SLP - BIAS       | (mb)   |        |        |        |
| FG - DATA        | 1.61   | 1.61   | 1.61   | 1.61   |
| NEW - DATA       | -0.91  | -2.12  | 0.88   | -1.27  |
| SLP - RMS        | (mb)   |        |        |        |
| FG - DATA        | 2.49   | 2.49   | 2.49   | 2.49   |
| NEW - DATA       | 1.42   | 2.57   | 1.80   | 1.87   |






Figure 1. The "true" sea level pressure analysis (mb).




Figure 2. The "true" surface wind analysis (knots).






Figure 3. The first-guess sea level pressure analysis (mb).




Figure 4. The first-guess surface wind analysis (knots).








Figure 5. Swath of synthetic scatterometer wind vector data (knots) passing over storm






Figure 6. Swath of synthetic SSM/I wind speed data (knots) passing over storm.




Figure 7. Swath of synthetic SSM/I wind speed data (knots) passing to the northeast of storm.






Figure 8. Locations of synthetic buoy wind data (knots).






Figure 9. Re-analyzed surface wind analysis (knots) using synthetic scatterometer wind vector data passing over the storm.




Figure 10. Sea level pressure analysis (mb) based on the balanced equation using the re-analyzed surface wind analysis generated from the synthetic scatterometer wind vector data passing over the storm.








Figure 11. Re-analyzed surface wind analysis (knots) using synthetic SSMI wind speed data passing over the storm.




Figure 12. Sea level pressure analysis based (mb) on the balanced equation using the re-analyzed surface wind analysis generated from the synthetic SSMI wind speed data passing over the storm.






Figure 13. Re-analyzed surface wind analysis (knots) using synthetic SSMI wind speed data passing to the northeast of the storm.




Figure 14. Sea level pressure analysis (mb) based on the balanced equation using the re-analyzed surface wind analysis generated from the synthetic SSMI wind speed data passing to the northeast of the storm.






Figure 15. Re-analyzed surface wind analysis (knots) using all synthetic wind data sets.




Figure 16. Sea level pressure analysis (mb) based on the balanced equation using the re-analyzed surface wind analysis generated from all synthetic wind data sets.



OPC CONTRIBUTIONS (Cont.)

- No. 21. Breaker, L. C., 1989: El Nino and Related Variability in Sea-Surface Temperature Along the Central California Coast. PACLIM Monograph of Climate Variability of the Eastern North Pacific and Western North America, Geophysical Monograph 55, AGU, 133-140.
- No. 22. Yu, T. W., D. C. Esteva, and R. L. Teboulle, 1991: A Feasibility Study on Operational Use of Geosat Wind and Wave Data at the National Meteorological Center. Technical Note/NMC Office Note No. 380, 28pp.
- No. 23. Burroughs, L. D., 1989: Open Ocean Fog and Visibility Forecasting Guidance System. Technical Note/NMC Office Note No. 348, 18pp.
- No. 24. Gerald, V. M., 1987: Synoptic Surface Marine Data Monitoring. Technical Note/NMC Office Note No. 335, 10pp.
- No. 25. Breaker, L. C., 1989: Estimating and Removing Sensor Induced Correlation from AVHRR Data. Journal of Geophysical Research, 95, 9701-9711.
- No. 26. Chen, H. S., 1990: Infinite Elements for Water Wave Radiation and Scattering. International Journal for Numerical Methods in Fluids, 11, 555-569.
- No. 27. Gemmill, W. H., T. W. Yu, and D. M. Feit, 1988: A Statistical Comparison of Methods for Determining Ocean Surface Winds. Journal of Weather and Forecasting, 3, 153-160.
- No. 28. Rao, D. B., 1989: A Review of the Program of the Ocean Products Center. Weather and Forecasting, 4, 427-443.
- No. 29. Chen, H. S., 1989: Infinite Elements for Combined Diffraction and Refraction. Conference Preprint, Seventh International Conference on Finite Element Methods Flow Problems, Huntsville, Alabama, 6pp.
- No. 30. Chao, Y. Y., 1989: An Operational Spectral Wave Forecasting Model for the Gulf of Mexico. Proceedings of 2nd International Workshop on Wave Forecasting and Hindcasting, 240-247.
- No. 31. Esteva, D. C., 1989: Improving Global Wave Forecasting Incorporating Altimeter Data. Proceedings of 2nd International Workshop on Wave Hindcasting and Forecasting, Vancouver, B.C., April 25-28, 1989, 378-384.
- No. 32. Richardson, W. S., J. M. Nault, and D. M. Feit, 1989: Computer-Worded Marine Forecasts. Preprint, 6th Symp. on Coastal Ocean Management Coastal Zone 89, 4075-4084.
- No. 33. Chao, Y. Y., and T. L. Bertucci, 1989: A Columbia River Entrance Wave Forecasting Program Developed at the Ocean Products Center. Technical Note/NMC Office Note 361.
- No. 34. Burroughs, L. D., 1989: Forecasting Open Ocean Fog and Visibility. Preprint, 11th Conference on Probability and Statistics, Monterey, Ca., 5pp.
- No. 35. Rao, D. B., 1990: Local and Regional Scale Wave Models. Proceeding (CMM/WMO) Technical Conference on Waves, WMO, Marine Meteorological of Related Oceanographic Activities Report No. 12, 125-138.
- No. 36. Burroughs, L.D., 1991: Forecast Guidance for Santa Ana conditions. Technical Procedures Bulletin No. 391, 11pp.
- No. 37. Burroughs, L. D., 1989: Ocean Products Center Products Review Summary. Technical Note/NMC Office Note No. 359. 29pp.
- No. 38. Feit, D. M., 1989: Compendium of Marine Meteorological and Oceanographic Products of the Ocean Products Center (revision 1). NOAA Technical Memo NWS/NMC 68.
- No. 39. Esteva, D. C., and Y. Y. Chao, 1991: The NOAA Ocean Wave Model Hindcast for LEWEX. Directional Ocean Wave Spectra, Johns Hopkins University Press, 163-166.
- No. 40. Sanchez, B. V., D. B. Rao, and S. D. Steenrod, 1987: Tidal Estimation in the Atlantic and Indian Oceans,  $3^\circ \times 3^\circ$  Solution. NASA Technical Memorandum 87812, 18pp.

OPC CONTRIBUTIONS (Cont.)

- No. 41. Crosby, D. S., L. C. Breaker, and W. H. Gemmill, 1990: A Definition for Vector Correlation and its Application to Marine Surface Winds. Technical Note/NMC Office Note No. 365, 52pp.
- No. 42. Feit, D. M., and W. S. Richardson, 1990: Expert System for Quality Control and Marine Forecasting Guidance. Preprint, 3rd Workshop Operational and Meteorological. CMOS, 6pp.
- No. 43. Gerald, V. M., 1990: OPC Unified Marine Database Verification System. Technical Note/NMC Office Note No. 368, 14pp.
- No. 44. Wohl, G. M., 1990: Sea Ice Edge Forecast Verification System. National Weather Association Digest, (submitted)
- No. 45. Feit, D. M., and J. A. Alpert, 1990: An Operational Marine Fog Prediction Model. NMC Office Note No. 371, 18pp.
- No. 46. Yu, T. W., and R. L. Teboule, 1991: Recent Assimilation and Forecast Experiments at the National Meteorological Center Using SEASAT-A Scatterometer Winds. Technical Note/NMC Office Note No. 383, 45pp.
- No. 47. Chao, Y. Y., 1990: On the Specification of Wind Speed Near the Sea Surface. Marine Forecaster Training Manual.
- No. 48. Breaker, L. C., L. D. Burroughs, T. B. Stanley, and W. B. Campbell, 1992: Estimating Surface Currents in the Slope Water Region Between 37 and 41°N Using Satellite Feature Tracking. Technical Note, 47pp.
- No. 49. Chao, Y. Y., 1990: The Gulf of Mexico Spectral Wave Forecast Model and Products. Technical Procedures Bulletin No. 381, 3pp.
- No. 50. Chen, H. S., 1990: Wave Calculation Using WAM Model and NMC Wind. Preprint, 8th ASCE Engineering Mechanical Conference, 1, 368-372.
- No. 51. Chao, Y. Y., 1990: On the Transformation of Wave Spectra by Current and Bathymetry. Preprint, 8th ASCE Engineering Mechanical Conference, 1, 333-337.
- No. 52. WAS NOT PUBLISHED
- No. 53. Rao, D. B., 1991: Dynamical and Statistical Prediction of Marine Guidance Products. Proceedings, IEEE Conference Oceans 91, 3, 1177-1180.
- No. 54. Gemmill, W. H., 1991: High-Resolution Regional Ocean Surface Wind Fields. Proceedings, AMS 9th Conference on Numerical Weather Prediction, Denver, CO, Oct. 14-18, 1991, 190-191.
- No. 55. Yu, T. W., and D. Deaven, 1991: Use of SSM/I Wind Speed Data in NMC's GDAS. Proceedings, AMS 9th Conference on Numerical Weather Prediction, Denver, CO, Oct. 14-18, 1991, 416-417.
- No. 56. Burroughs, L. D., and J. A. Alpert, 1993: Numerical Fog and Visibility Guidance in Coastal Regions. Technical Procedures Bulletin. No. 398, 6pp.
- No. 57. Chen, H. S., 1992: Taylor-Gelerkin Method for Wind Wave Propagation. ASCE 9th Conf. Eng. Mech. (in press)
- No. 58. Breaker, L. C., and W. H. Gemmill, and D. S. Crosby, 1992: A Technique for Vector Correlation and its Application to Marine Surface Winds. AMS 12th Conference on Probability and Statistics in the Atmospheric Sciences, Toronto, Ontario, Canada, June 22-26, 1992.
- No. 59. Yan, X.-H., and L. C. Breaker, 1993: Surface Circulation Estimation Using Image Processing and Computer Vision Methods Applied to Sequential Satellite Imagery. Photogrammetric Engineering and Remote Sensing, 59, 407-413.
- No. 60. Wohl, G., 1992: Operational Demonstration of ERS-1 SAR Imagery at the Joint Ice Center. Proceeding of the MTS 92 - Global Ocean Partnership, Washington, DC, Oct. 19-21, 1992.



OPC CONTRIBUTIONS (Cont.)

- No. 61. Waters, M. P., Caruso, W. H. Gemmill, W. S. Richardson, and W. G. Pichel, 1992: An Interactive Information and Processing System for the Real-Time Quality Control of Marine Meteorological Oceanographic Data. Pre-print 9th International Conference on Interactive Information and Processing System for Meteorology, Oceanography and Hydrology, Anaheim, CA, Jan. 17-22, 1993.
- No. 62. Breaker, L. C., and V. Krasnopolsky, 1994: The Problem of AVHRR Image Navigation Revisited. Int. Journal of Remote Sensing, 15, 979-1008.
- No. 63. Crosby, D. S., L. C. Breaker, and W. H. Gemmill, 1993: A Proposed Definition for Vector Correlation in Geophysics: Theory and Application. Journal of Atmospheric and Ocean Technology, 10, 355-367.
- No. 64. Grumbine, R., 1993: The Thermodynamic Predictability of Sea Ice. Journal of Glaciology, 40, 277-282, 1994.
- No. 65. Chen, H. S., 1993: Global Wave Prediction Using the WAM Model and NMC Winds. 1993 International Conference on Hydro Science and Engineering, Washington, DC, June 7 - 11, 1993. (submitted)
- No. 66. WAS NOT PUBLISHED
- No. 67. Breaker, L. C., and A. Bratkovich, 1993: Coastal-Ocean Processes and their Influence on the Oil Spilled off San Francisco by the M/V Puerto Rican. Marine Environmental Research, 36, 153-184.
- No. 68. Breaker, L. C., L. D. Burroughs, J. F. Culp, N. L. Gunasso, R. Tebouille, and C. R. Wong, 1993: Surface and Near-Surface Marine Observations During Hurricane Andrew. Technical Note/NMC Office Note #398, 41pp.
- No. 69. Burroughs, L. D., and R. Nichols, 1993: The National Marine Verification Program - Concepts and Data Management, Technical Note/NMC Office Note #393, 21pp.
- No. 70. Gemmill, W. H., and R. Tebouille, 1993: The Operational Use of SSM/I Wind Speed Data over Oceans. Pre-print 13th Conference on Weather Analyses and Forecasting, AMS Vienna, VA., August 2-6, 1993, 237-238.
- No. 71. Yu, T.-W., J. C. Derber, and R. N. Hoffman, 1993: Use of ERS-1 Scatterometer Backscattered Measurements in Atmospheric Analyses. Pre-print 13th Conference on Weather Analyses and Forecasting, AMS, Vienna, VA., August 2-6, 1993, 294-297.
- No. 72. Chalikov, D. and Y. Liberman, 1993: Director Modeling of Nonlinear Waves Dynamics. J. Physical, (To be submitted).
- No. 73. Woiceshyn, P., T. W. Yu, W. H. Gemmill, 1993: Use of ERS-1 Scatterometer Data to Derive Ocean Surface Winds at NMC. Pre-print 13th Conference on Weather Analyses and Forecasting, AMS, Vienna, VA, August 2-6, 1993, 239-240.
- No. 74. Grumbine, R. W., 1993: Sea Ice Prediction Physics. Technical Note/NMC Office Note #396, 44pp.
- No. 75. Chalikov, D., 1993: The Parameterization of the Wave Boundary Layer. Journal of Physical Oceanography, Vol. 25, No. 6, Par 1, 1333-1349.
- No. 76. Tolman, H. L., 1993: Modeling Bottom Friction in Wind-Wave Models. Ocean Wave Measurement and Analysis, O.T. Magoon and J.M. Hemsley Eds., ASCE, 769-783.
- No. 77. Breaker, L., and W. Broenkow, 1994: The Circulation of Monterey Bay and Related Processes. Oceanography and Marine Biology: An Annual Review, 32, 1-64.
- No. 78. Chalikov, D., D. Esteva, M. Iredell and P. Long, 1993: Dynamic Coupling between the NMC Global Atmosphere and Spectral Wave Models. Technical Note/NMC Office Note #395, 62pp.
- No. 79. Burroughs, L. D., 1993: National Marine Verification Program - Verification Statistics - Verification Statistics, Technical Note/NMC Office Note #400, 49 pp.

OPC CONTRIBUTIONS (Cont.)

- No. 80. Shashy, A. R., H. G. McRandal, J. Kinnard, and W. S. Richardson, 1993: Marine Forecast Guidance from an Interactive Processing System. 74th AMS Annual Meeting, January 23 - 28, 1994.
- No. 81. Chao, Y. Y., 1993: The Time Dependent Ray Method for Calculation of Wave Transformation on Water of Varying Depth and Current. Wave 93 ASCE.
- No. 82. Tolman, H. L., 1994: Wind-Waves and Moveable-Bed Bottom Friction. Journal of Physical Oceanography, 24, 994-1009.
- No. 83. Grumbine, R. W., 1993: Notes and Correspondence A Sea Ice Albedo Experiment with the NMC Medium Range Forecast Model. Weather and Forecasting, (submitted).
- No. 84. Chao, Y. Y., 1993: The Gulf of Alaska Regional Wave Model. Technical Procedure Bulletin, No. 427, 10 pp.
- No. 85. Chao, Y. Y., 1993: Implementation and Evaluation of the Gulf of Alaska Regional Wave Model. Technical Note, 35 pp.
- No. 86. WAS NOT PUBLISHED.
- No. 87. Burroughs, L., 1994: Portfolio of Operational and Development Marine Meteorological and Oceanographic Products. Technical Note/NCEP Office Note No. 412, 52 pp. [PB96-158548]
- No. 88. Tolman, H. L., and D. Chalikov, 1994: Development of a third-generation ocean wave model at NOAA-NMC. Proc. Waves Physical and Numerical Modelling, M. Isaacson and M.C. Quick Eds., Vancouver, 724-733.
- No. 89. Peters, C., W. H. Gemmill, V. M. Gerald, and P. Woiceshyn, 1994: Evaluation of Empirical Transfer Functions for ERS-1 Scatterometer Data at NMC. 7th Conference on Satellite Meteorology and Oceanography, June 6-10, 1994, Monterey, CA., pg. 550-552.
- No. 90. Breaker, L. C., and C. R. N. Rao, 1996: The Effects of Aerosols from the Mt. Pinatubo and Mt. Hudson Volcanic Eruption on Satellite-Derived Sea Surface Temperatures. Journal of Geophysical Research. (To be submitted).
- No. 91. Yu, T-W., P. Woiceshyn, W. Gemmill, and C. Peters, 1994: Analysis & Forecast Experiments at NMC Using ERS-1 Scatterometer Wind Measurements. 7th Conference on Satellite Meteorology and Oceanography, June 6-10, 1994, Monterey, CA., pg. 600-601.
- No. 92. Chen, H. S., 1994: Ocean Surface Waves. Technical Procedures Bulletin, No. 426, 17 pp.
- No. 93. Breaker, L. C., V. Krasnopolsky, D. B. Rao, and X.-H. Yan, 1994: The Feasibility of Estimating Ocean Surface Currents on an Operational Basis using Satellite Feature Tracking Methods. Bulletin of the American Meteorological Society, 75, 2085-2095.
- No. 94. Krasnopolsky V., L. C. Breaker, and W. H. Gemmill, 1994: Development of Single "All-Weather" Neural Network Algorithms for Estimating Ocean Surface Winds from the Special Sensor Microwave Imager. Technical Note.
- No. 95. Breaker, L. C., D. S. Crosby and W. H. Gemmill, 1994: The application of a New Definition for Vector Correlation to Problems in Oceanography and Meteorology. Journal of Applied Meteorology, 33, 1354-1365.
- No. 96. Peters, C. A., V. M. Gerald, P. M. Woiceshyn, and W. H. Gemmill, 1994: Operational Processing of ERS-1 Scatterometer winds: A Documentation. Technical Note.
- No. 97. Gemmill, W. H., P. M. Woiceshyn, C. A. Peters, and V. M. Gerald, 1994: A Preliminary Evaluation Scatterometer Wind Transfer Functions for ERS-1 Data. Technical Note.
- No. 98. Chen, H. S., 1994: Evaluation of a Global Ocean Wave Model at NMC. International Conference on Hydro-Science and Engineering. Beijing, China, March 22 - 26, 1995.

OPC CONTRIBUTIONS (Cont.)

- No. 99. Aikman, F. and D. B. Rao, 1994: NOAA Perspective on a Coastal Forecast System.
- No. 100. Rao, D. B. and C. Peters, 1994: Two-Dimensional Co-Oscillations in a Rectangular Bay: Possible Application to Water-Level Problems. Marine Geodesy, 18, 317-332.
- No. 101. Breaker, L. C., L. D. Burroughs, Y. Y. Chao, J. F. Culp, N. L. Gunasso, R. Tebouille, and C. R. Wong, 1994: Surface and Near-Surface Marine Observations During Hurricane Andrew. Weather and Forecasting, 9, 542-556.
- No. 102. Tolman, H. L., 1995: Subgrid Modeling of Moveable-bed Bottom Friction in Wind Wave Models. Coastal Engineering, Vol 26, pp 57-75.
- No. 103. Breaker, L. C., D. B. Gilhousen, H. L. Tolman and L. D. Burroughs, 1995: Initial Results from Long-Term Measurements of Atmospheric Humidity and Related Parameters the Marine Boundary Layer at Two Locations in the Gulf of Mexico. (To be submitted to Global Atmosphere and Ocean Systems).
- No. 104. Burroughs, L. D., and J. P. Dallavalle, 1995: Great Lakes Wind and Wave Guidance. Technical Procedures Bulletin No., (In preparation).
- No. 105. Burroughs, L. D., and J. P. Dallavalle, 1995: Great Lakes Storm Surge Guidance. Technical Procedures Bulletin No., (In preparation).
- No. 106. Shaffer, W. A., J. P. Dallavalle, and L. D. Burroughs, 1995: East Coast Extratropical Storm Surge and Beach Erosion Guidance. Technical Procedures Bulletin No., (In preparation)
- No. 107. WAS NOT PUBLISHED.
- No. 108. WAS NOT PUBLISHED.
- No. 109. WAS NOT PUBLISHED.
- No. 110. Gemmill, W. H, and C. A. Peters, 1995: The Use of Satellite Dervired Wind Data in High-Resolution Regional Ocean Surface Wind Fields. Conference on Coastal Oceanic and Atmospheric Prediction, Jan 28 - Feb 2, 1996, Atlanta, GA (accepted at preprint press).

OPC CHANGES TO OMB

- No. 111. Krasnopolsky, V. M, W. H. Gemmill, and L. C. Breaker, 1995: Improved SSM/I Wind Speed Retrievals at Higher Wind Speeds. Journal of Geophysical Research, (in press).
- No. 112. Chalikov, D., L. D. Breaker, and L. Loboeki, 1995: A Simple Model of Mixing in the Upper Ocean. Journal of Physical Ocean, (in press).
- No. 113. Tolman, H. L., 1995: On the Selection of Propagation Schemes for a Spectral Wind-Wave Model. NCEP Office Note No. 411.
- No. 114. Grumbine, R. W., 1995: Virtual Floe Ice Drift Forecast Model Intercomparison. NCEP Office Note. (To be submitted).
- No. 115. Grumbine, R. W., 1995: Sea Ice Forecast Model Intercomparison: Selecting a Base Model for NCEP Sea Ice Modelling. Technical Note.
- No. 116. Yu, T. W. and J. C. Derber, 1995: Assimilation Experiments with ERS-1 Winds: Part I - Use of Backscatter Measurements in the NMC Spectral Statistical Analysis System. Technical Note.
- No. 117. Yu, T. W., 1995: Assimilation Experiments with ERS1 Winds: Part II - Use of Vector Winds in NCEP Spectral Statistical Analysis System. Technical Note.
- No. 118. Grumbine, R. W., 1995: Sea Ice Drift Guidance. Technical Procedures Bulletin. (submitted)

OMB CONTRIBUTIONS (Cont.)

- No. 119. Tolman, H. L., 1996: Effects of Observation Errors in Linear Regression and Bin-Average Analyses. Quarterly Journal of the Royal Meteorological Society.
- No. 120. Grumbine, R. W., 1996: Automated Passive Microwave Sea Ice Concentration Analysis at NCEP. Technical Note.
- No. 121. Grumbine, R. W., 1996: Sea Ice Prediction Environment: Documentation. Technical Note.
- No. 122. Tolman, H. L and D. Chalikov, 1996: Source Terms in a Third-Generation Wind Wave Model. Journal of Physical Oceanography. Vol 26, pp 2497-2518.
- No. 123. Gemmill, W. H., V. Krasnopolsky, L. C. Breaker, and C. Peters, 1996: Developments to Improve Satellite Derived Ocean Surface Winds for use in Marine Analyses. Pre-print Numerical Weather Prediction Conference, Norfolk, VA, Aug. 19-23, 1996.
- No. 124. Breaker, L. C., D. B. Gilhousen, H. L. Tolman and L. D. Burroughs, 1996: Initial Results from Long-Term Measurements of Atmospheric Humidity and Related Parameters in the Marine Boundary Layer at Two Locations in the Gulf of Mexico. NCEP Office Note No. 414.
- No. 125. Yu, T. W., M. D. Iredell, and Y. Zhu, 1996: The Impact of ERS-1 Winds on NCEP Operational Numerical Weather Analyses and Forecast. Pre-print Numerical Weather Prediction Conference, Norfolk, VA, August 19-23, 1996.
- No. 126. Burroughs, L. D., 1996: Marine Meteorological and Oceanographic Guidance Products from the National Centers for Environmental Prediction. Mariners Weather Log, Vol. 40, No. 2, pp 1-4.
- No. 127. Loboeki, L., 1996: Coastal Ocean Forecasting System (COFS) System Description and User Guides. Technical Note.
- No. 128. WAS NOT PUBLISHED
- No. 129. Chaikov, D., 1996: A Global Ocean Model. Technical Note.
- No. 130. Yu, T.W., 1996: Applications of SSM/I Wind Speed Data to NCEP Regional Analyses. Technical Note.
- No. 131. Chaikov, D. and D. Sheinin, 1996: Direct Modeling of 1-D Nonlinear Potential Waves. Advances in Fluid Mechanics Series: Nonlinear Ocean Waves (submitted).
- No. 132. Krasnopolsky, V. M., W. H. Gemmill, L. C. Breaker, and V. Y. Raizer, 1996: SSM/I Wind Speed Retrieval Algorithm with Improved Performance at Higher Wind Speed. Remote Sensing of Environment (submitted).
- No. 133. Yu, T. W., 1996: The Effect of Drifting Buoy Data on NCEP Numerical Weather Forecast. Technical Note.
- No. 134. Krasnopolsky, V. M., 1996: A Neural Network Forward Model for Direct Assimilation of SSM/I Brightness Temperatures into Atmospheric Models. CAS/JSC Working Group on Numerical Experimentation (in press).
- No. 135. Krasnopolsky, V. M., W. H. Gemmill, and L. C. Breaker, 1996: A New Neural Network Transfer for SSM/I Retrievals. CAS/JSC Working Group on Numerical Experimentation (in press).
- No. 136. Krasnopolsky, V. M., 1996: NN Solutions for Forward & Inverse Problems in Satellite Remote Sensing. 1997 International Conference on Neural Networks (ICNN 97). (submitted).
- No. 137. Krasnopolsky, V. M., 1996: A New Neural Network Transfer Function for SSM/I Based on an Expanded Neural Network Architecture. Technical Note.
- No. 138. Chaikov, D. C., L. C. Breaker, and L. Loboeki, 1996: Parameterization of Mixing in Upper Ocean. Technical Note.
- No. 139. Chaikov, D. C., and D. Sheinin, 1996: Numerical Modeling of Surface Waves Based on Principal Equations of Potential Wave Dynamics. Technical Note.

OMB CONTRIBUTIONS (Cont.)

- No. 140. Krasnopolsky, V. M., 1997: A Neural Network-Based Forward Model for Direct Assimilation of SSM/I Brightness Temperatures. Technical Note.
- No. 141. Peters, C. A., 1997: Effects of Scatterometer Winds on the NCEP Global Model Analyses and Forecasts: Two Case Studies. Technical Note.
- No. 142. Kelley, J. G. W., F. Aikman, L. C. Breaker and G. L. Mellor, 1997: A Coastal Ocean Forecast System for the U.S. East Coast. Sea Technology.
- No. 143. Tolman, H. L., L. C. Bender and W. L. Neu, 1997: Comments on "The Goddard Coastal Wave Model. Part I: Numerical Method. Journal of Physical Oceanography.
- No. 144. Tolman, H. L., W. L. Neu and L. C. Bender, 1997: Comments on "The Goddard Coastal Wave Model. Part II: Kinematics. Journal of Physical Oceanography.
- No. 145. Breaker, L. C., D. B. Gilhousen, H. L. Tolman, and L. D. Burroughs, 1997: Initial Results from Long-Term Measurements Atmospheric Humidity and Related Parameters in the Marine Boundary Layer at Two Locations in the Gulf of Mexico. Journal of Marine Systems. (submitted)
- No. 146. Peters, C. A., 1997: Effects of Scatterometer Winds on the NCEP Global Model Analyses and Forecasts: Two Case Studies. Weather and Forecasting.
- No. 147. Gemmill, W. H. and C. A. Peters, 1997: High-Resolution Ocean Surface Wind Analyses Using Satellite Derived Ocean Surface Winds: Analyses Validation using Synthetic Satellite Data. Technical Note.

