

**U.S. Wheat and Barley Scab Initiative
 FY01 Final Performance Report (approx. May 01 – April 02)
 July 15, 2002**

Cover Page

PI:	Sanjaya Rajaram
Institution:	CIMMYT
Address:	Apdo. Postal 6-641 Mexico, D.F., Mexico 06600 MEXICO
Email:	s.rajaram@cgiar.org
Phone:	52-5804-2004
Fax:	52-5824-7559
Year:	FY2001 (approx. May 01 – April 02)
Grant Number:	59-0790-0-F076
Grant Title:	Fusarium Head Blight Research
FY01 ARS Award Amount:	\$ 136,285

Project

Program Area	Project Title	Requested Amount
Germplasm	Facilitating international germplasm and information exchange through CIMMYT	\$ 140,000
	Total Amount Requested	\$ 140,000

Principal Investigator

Date

Project 1: Facilitating international germplasm and information exchange through CIMMYT

1. What major problem or issue is being resolved and how are you resolving it?

The overall goal of this collaboration is to increase genetic resistance to Fusarium Head Blight (FHB) in commercially grown US wheat varieties and thus significantly increase the production and yield stability of wheat in the United States of America. Specifically, the objectives of the project area:

- To provide agronomically suitable FHB resistant germplasm to US collaborators through pre-breeding activities using synthetic wheat and major US cultivars;
- To conduct a world-wide search for and acquisition of suitable FHB resistant germplasm and to make this available to the US Wheat and Barley Scab Initiative;
- To test germplasm through the International Testing Nursery; and
- To provide elite germplasm pre-breeding activities, FHB resistant germplasm and International Nursery Testing for barley.

2. What were the most significant accomplishments?

Pre-breeding: Fifty-eight BC1F1 and F2 populations derived from crosses between five commercial USA recurrent spring wheat cultivars (Ivan, Reeder, Russ, Verde and Wheaton) with 12 scab resistant donor parents have been advanced. Screening at Toluca, Mexico with subsequent backcrossing is currently underway. In addition, over 2000 advanced lines screened during the past five years, during the reporting period the best 100 lines were characterized for resistance types I, II and IV, with the a sub-set of 72 lines further characterized for type III resistance. CIMMYT's collection of 800 D-genome synthetic hexaploids (SH) has been screened for type II resistance to FHB. Targeted for durum wheat improvement, sets of A-genome (AAAABB, 194 entries) and B-genome (AABBBB, 54 entries) SH were screened for FHB. Doubled haploids (DH) derived from FHB mapping populations Mayoor/SH(222)//Flycatcher (160) have been supplied to Rick Ward for molecular analysis. DH's derived from FHB mapping populations Frontana/Inia66 (117), Fukuho-komugi/Oligoculm 380 (110), Emblem/Saikai165 (104) and Catbird/Milan (100) are undergoing phenotypic resistance analysis (types I, II and III) in Toluca.

World-wide germplasm: FHB resistant wheat germplasm, which has undergone subsequent FHB resistance confirmation by CIMMYT scientists, has been obtained from: Argentina (127), Brazil (15) and Japan (15). Third-year resistance confirmation has been conducted for 37 CIMMYT-China shuttled derived bread wheat advanced lines. New *Aegilops tauschii* accessions have been obtained from Australia and WGRC to be used for primary SH production.

International Testing Nursery: The 6th International Scab Resistant Screening Nursery has been distributed to 30 international institutional cooperators in 19 countries. Targeted screening, under supervision of CIMMYT scientists, is being conducted in Argentina, Brazil, China, Mexico, Paraguay, Romania and Uruguay.

Barley: One hundred and five cross populations were generated using the following sources of FHB resistance were used in the ICARDA/CIMMYT barley breeding program during the reporting period: Six-row types: Chevron, Chamico/Tocte//Congona, PFC88209, MNBrite, MNS1. Two-row types: Atahualpa, Azafran, Gobernadora, Shyri, Svanhals, Zhedar1, Zhedar2.

Include below a list of the publications, presentations, peer-reviewed articles, and non-peer reviewed articles written about your work that resulted from all of the projects included in the grant. Please reference each item using an accepted journal format. If you need more space, continue the list on the next page.

- Ackermann, M. Diaz de and M. M. Kohli. 2001. Control Quimico de Fusariosis de la Espiga en Trigo. *Fitopatol. bras.* 26 (Suplemento), pp. 460. Agosto 2001.
- Cano S, R. Delgado and A. Mujeeb-Kazi. 2002. Contribution of alien germplasm for Fusarium head scab resistance in wheat. *Agronomy Abstracts, American Society of Agronomy*, IN PRESS.
- Capettini, F., H. Vivar , L. Gilchrist, and M. Henry. 2001. Building up Multiple Disease Resistance in Barley. Warren E, Kronstad Symposium. Cd. Obregon Sonora, Mexico. March 15, .2001
- Gilchrist, L., C. Velazquez, and H. Vivar. 1999. Evaluación de resistencia a Fusariosis de la espiga en la población de haploides doblados RECLA. Resúmenes del III Congreso Latinoamericano de Cebada . 5-8 de Octubre de 1999.
- Gilchrist, L., S. Rajaram and J. Crossa. 2000. New sources of scab resistance and breeding progress at CIMMYT. Proceedings of the International Symposium on Wheat Improvement for Scab Resistance. 5-11 May 2000 Suzhou and Nanjing, China.
- Gilchrist, L. and C. Velazquez .2000. Pathogenicity and virulences of eight *Fusarium graminearum* isolates originating in four regions of Mexico. 2000 National Fusarium Head Blight Forum. Cincinnati. December 10-12.
- Gilchrist, L. 2001. Perspectives on Fusarium Head Blight Resistance in Barley. In: H. Vivar and A. McNab (eds.). *Breeding Barley in the New Millennium: Proceedings of an International Symposium*. Mexico, D.F.:CIMMYT. pp: 61-71
- Gilchrist, L. Velazquez, C. and J. Crossa. 2001 Pathogenicity and Virulence of eight Fusarium graminearum isolates Originating in four Regions of Mexico. Warren E, Kronstad Symposium. Cd. Obregon Sonora, Mexico. March 15.2001
- He, Z.H., M. van Ginkel, L. Gilchrist and S. Rajaram. 2000. Progress of China /CIMMYT Shuttle Breeding and germplasm exchange aimed at combining high yield potential with scab resistance. 2000 National Fusarium Head Blight Forum. Cincinnati. December 10-12.
- He, Z.H., S. Rajaram, Z.Y. Xin and G.Z. Huang (eds.). 2001. *A History of Wheat Breeding in China*. Mexico, D.F.: CIMMYT.
- Mujeeb-Kazi, A., L. Gilchrist, R.L. Villarreal, and R. Delgado. 2000. New synthetic hexaploid and a set of bread wheat / synthetic hexaploid derivatives as sources for scab resistance. *Annual Wheat Newsletter*. 46: 81-82.
- Mujeeb-Kazi, A, A. Cortes, V. Rosas, S. Cano, J. Sanchez, L. Juarez and R. Delgado. 2001. Genetic diversity for improving scab resistance in wheat. In: Warren Kronstad Symposium, Cd. Obregon, Mexico, March 15, 2001, P. 126-129.
- Mujeeb-Kazi, A. 2001. Intergeneric hybrids in wheat : Current status. In: IVth International Triticeae Symposium, ed. Hernandez P, M. T. Moreno, J. I. Cubero and A. Martin, September 10-12, Cordoba, Spain, p 261-264.

- Mujeeb-Kazi, A. 2001. Synthetic hexaploids for bread wheat improvement. In: IVth International Triticeae Symposium, ed. Hernandez P, M. T. Moreno, J. I. Cubero and A. Martin, September 10-12, Cordoba, Spain, p 193-199.
- Mujeeb-Kazi, A., R. Delgado, S. Cano, L. Juarez and J. Sanchez. 2001. Scab resistance in bread wheat/synthetic hexaploid derivatives. Annual Wheat Newsletter 47:104-105.
- Mujeeb-Kazi, A., and R. Delgado. 2001. A second elite set of synthetic hexaploid wheats based upon multiple disease resistance. Annual Wheat Newsletter 47:114-115.
- Mujeeb-Kazi A., R. Delgado, L. Juarez, J. Sanchez and S. Cano. 2001. Scab resistance (Type II: Spread) in synthetic hexaploid germplasm. Annual Wheat Newsletter 47:118-120.
- Mujeeb-Kazi, A., and S. Rajaram. 2002. Production and utilization of synthetic hexaploid wheats for global biotic/abiotic stress constraints. Agronomy Abstracts, American Society of Agronomy, IN PRESS.
- Mujeeb-Kazi, A., and R. Delgado. 2002. D genome contribution to head scab resistance (Type II). Annual Wheat Newsletter, IN PRESS.
- Mujeeb-Kazi A. and R Delgado. 2002. Free threshing bread wheat/D genome synthetic hexaploids resistant to head scab. Annual Wheat Newsletter, IN PRESS.
- Rajaram, S., R.J. Peña, R.L. Villareal, A. Mujeeb-Kazi, R. Singh, and L. Gilchrist. 2001. Utilization of wild and cultivated emmer and of diploid wheat relatives in breeding . Israel Journal of Plant Science. Vol 49 : 93-104
- Velazquez, C., L. Gilchrist and J. Burgueño. 2000. Importance of spike growth stage for evaluating Fusarium Head Blight resistance in wheat. A supplement to Proceedings of the International Symposium on wheat improvement for scab resistance. 5-11 May 2000 Suzhou and Nanjing, China.
- van Ginkel, M., L. Gilchrist and C. Velazquez. 2000. Newly Accumulated Resistances in CIMMYT Bread Wheat Germplasm. 2000 National Fusarium Head Blight Forum. Cincinnati, OH. December 10-12, 2000.
- van Ginkel, M., and L. Gilchrist. 2001. New bread wheats for high rainfall environments: The package. Pp. 8-12 In: CIMMYT. 2001. Research Highlights of the CIMMYT Wheat Program, 1999-2000. Mexico, D.F.