

UNIFORM SOUTHERN SOFT RED WINTER WHEAT FUSARIUM HEAD BLIGHT SCREENING NURSERY

2003 NURSERY REPORT

J. P. Murphy
R. A. Navarro
D. A. Van Sanford

This is a joint progress report of cooperative investigations underway and funded by the State Agricultural Experiment Stations, private companies and the United States Department of Agriculture, Agricultural Research Service. This report contains preliminary data that have not been sufficiently confirmed to justify general release; interpretations may be modified with additional experimentation. Confirmed results will be published through established channels. The report is a tool for the use of the cooperators and their official staff and those persons having direct interest in the development of agricultural research programs.

This report is not intended for publication and should not be referred to in literature citations or quoted in publicity or advertising. Use of the data may be granted for certain purposes upon written request to the agency or agencies involved.

North Carolina State University
Department of Crop Science
Box 7629
Raleigh, NC 27695-7629

December 1, 2003

TABLE OF CONTENTS

Location Notes.....	3
List of Entries and Pedigrees.....	7
FHB Incidence	8
FHB Severity.....	9
FHB Severity, GGE Biplot Analysis.....	10
AUDPC Head Severity.....	11
FHB Index.....	12
FHB Index by Individual Isolates.....	13
Scabby Seed %	14
Scabby Seed %, GGE Biplot Analysis	15
Grain Quality.....	16
Vomitoxin (DON).....	17
Greenhouse Screening (Head Severity and Spread).....	18
Greenhouse Screening (Head Severity), GGE Biplot Analysis	19
Greenhouse Screening (Rachis Involvement)	20
Detached Leaf Evaluation.....	21
Heading Date.....	22
Plant Height	23
Leaf Diseases	24
Grain Yield	25
<i>Xgwm 493 and XBARC 133 Marker Genotypes</i>	26
Means Across Locations.....	27
Correlations Between Traits.....	28

LOCATION NOTES

Bay, Arkansas

- Cooperators: June Hancock, Luis Lazoanaya
Syngenta Seeds Inc.
- Reps: 2 Plot size: 4 rows 3 ft long
- Inoculation method: Infected corn on March 11 and infected wheat on April 22. Counts were made around May 12. Visual scores were rated on May 2.
- Precipitation during grain fill: Mainly natural moisture. From April 24 to May 19 we had almost 16 inches of rain.
- Avg temp. during grain fill: 65-70, very cool

Fayetteville and Kibler, Arkansas

Cooperators: Gene Milus, Peter Rohman, Sam Markell

University of Arkansas. Fayetteville (F) and Kibler (K).

Reps: 4 (field), 3 (greenhouse) Plot size: Two rows, five feet long (F). Three rows, five feet long (K).

Seed date: 10/15/02 (F) 10/23/02 (K) Harvest date: 6/09/03 (K), Fayetteville not harvested.

Fertilizer: 100 lbs N (F), 80 lbs N (K) split appl.

Inoculation method: FHB-infected corn kernels applied twice for total of 6 kernels/sq ft. Applied 4/04/03 and 04/14/03 (F). 03/26/03 and 04/07/03 (K).

Irrigation method: eight mist periods of 10 minutes between midnight and 8am on nine nights between 29 April and 13 May (F).

Two hour sprinkling periods three days per week from April to mid-May (K).

Precipitation during grain fill: 2.47" in April, 5.90" in May. Most occurred after irrigation ceased (F). 2.6" in April and 2.4" in May (K).

Date/Feekes growth stage when scored: 5/27/03 Feekes 11.2 (F). 5/19/03 Feekes 11.2 (K).

Comments: Three weeks of rainy weather after irrigation stopped led to severe FHB. No samples sent for DON analysis.

Urbana, Illinois

Cooperators: Fred Kolb, Larry Boze

University of Illinois, Urbana, IL.

Reps: 3 Plot size: 1 row x 3' Seed date: 09/27/02 Harvest date: 07/07/03

Fertilizer: 40 lb N/A; P and K okay; no spring topdress

Inoculation method: wheat grain spawn and corn fodder scattered in field and sprayed with inoculum.

Precipitation during grain fill: Little natural, misted 5:30-7 AM and 7:30-9 PM daily (0.12 in./hr).

Date/Feekes growth stage when scored: Feekes 10.5, 29 d after flowering. Longer than normal due to cool temperatures.

Comments: Cool weather delayed onset of symptoms.

Wooster, Ohio

Cooperators: Clay Sneller, Pat Lipps.

Ohio State University, Wooster, OH.

Reps: 3 Plot size: One 3' row.

Fertilizer: P, K according to soil test; 110 lbs N split applied.

Irrigation: Misted at sunrise and sunset for two weeks after anthesis.

Inoculation method: Infested corn kernels.

Rated 28-31 days after anthesis.

Comments: Cool temperatures extended the heading period and delayed disease onset.

Lexington, Kentucky

Cooperators: A. J. Stewart and D. Van Sanford

University of Kentucky, Lexington, KY.

Reps: 2 Plot size: Two 4' rows

Seed date: 10/22/021

Harvest date: 6/29/03

Fertilizer: P, K, acc. to soil tests, 110 lb N split application

Irrigation method: Evening/early morning irrigation.

Inoculation method: Scabby corn

Precipitation during grain fill: 8.59 in.

Avg temperature during grain fill: 61.7° F.

Date/Feekes growth stage when scored: 10.5 + 21 d

Blacksburg, Virginia

Cooperators: Carl A. Griffey, Julie Wilson, Daryoosh Nabati,

Virginia Tech, Blacksburg, VA.

Reps: 3, randomized complete block

Plot size: 4 x 5 ft (20 ft²)

Seed date: 10/9/02

Harvest date: 7/15/03.

Fertilizer: 25-70-60 applied 10/08/02 and 03/23/03.

Inoculation method: Field, conidial spray. Greenhouse, floret inoculation.

Precipitation during grain fill: 14.14 in.; mist irrigation also applied as needed

Avg temperature during grain fill: 62.7°F

Manhattan, Kansas

Cooperators: Gina Brown-Guedira.

USDA-ARS Genotyping Center

Kinston, North Carolina

Cooperators: Paul Murphy, Rene Navarro

North Carolina State University, Kinston, NC.

Reps: 2 Plot size: 4 rows x 3' long Seed date: 11/26/02 Harvest date: 6/16/03

Fertilizer: 130 lbs N

Inoculation method: Conidial suspension (3×10^4 spores/ml) sprayed on plots at anthesis; misted for 14 d

Precipitation during grain fill: 11.01in.

Avg temperature during grain fill: 62.7°F.

Comments: Very, very wet season. Late planting and constant moisture through winter resulted in poor stand establishment. Less than desirable plots. Although heavy natural FHB epidemic, disease development in nursery was poor.

Columbia, Missouri

Cooperator: Anne L. McKendry, Jessica Tremain

University of Missouri, Columbia, MO.

Reps: 4. RCBD. Plot size: 4 rows each 3' long. Planted: 10/01/02. Harvested: 01/14/03

Inoculation method (Field): Sprayed at 75% heading with a suspension of *Fusarium graminearum* macroconidia concentrated to 50,000 spores/ml. Plots were rated after 28 days which was later than usual due to cool temperatures.

Precipitation during grain fill: Overhead mist irrigation

Inoculation method (Greenhouse): Point inoculation of a single central floret at anthesis with 10 micro litres of a 50,000 spores/mL suspension of *Fusarium graminearum* macroconidia. Plants were misted for 72 hours and rated 21 days post inoculation.

Comments: Field conditions were cooler than usual this spring and we did not get normal amount of disease development. Greenhouse point inoculations were OK. All field data are means based on 2 random samples of 10 heads/plot in each of four replications (80 plants total). Greenhouse data are mean data for 8 plants/entry.

Fundulea, Romania

Cooperator: Marianna Ittu.

Agricultural Research Development Institute.

Reps: 3 Plot size: single 0.5m row. Planted: 10/23/02. Harvested 07/03/03.

Inoculation method: Injection of conidial suspension in midflower of 20 heads at anthesis.

Feekes growth stage when scored: 10.5 plus 10 and 20 days.

Average temperature during grain fill: 22.4°C.

Precipitation during grain fill: 35.8mm.

Comments: High rainfall at sowing time delayed planting. Long, severe winter led to poor survival of winter tender cultivars. Late spring and unusually hot and dry weather reduced plant height and grain yield.

Szeged, Hungary.

Cooperator: Akos Mesterhazy.
Cereal Research Institute.

Dublin, Ireland.

Cooperators: Roy Browne, Mike Cooke.
University College Dublin.

In vitro assay. Partial disease resistance components detected in a detached leaf assay as described by Browne and Cooke, (2004), European Journal of Plant Pathology,(in press).

Five isolates of *M. nivale* var. *majus* utilized. Five replicates over time. An experimental unit consisted of one Petri dish containing four leaves of an entry. A different isolate of *M. nivale* utilized in each of the five replications.

Entry List, 2003 Nursery

ENTRY NO	CULTIVAR/DESINATION	PEDIGREE	CONTRIBUTOR	IN NURSERY SINCE
1	ERNIE	PIKE /3/ STODDARD / BLUEBOY // STODDARD / D1707	CHECK (RES)	1999-00
2	COKER 9835	CK68-19 // CK61-19*3 / IN4946A4-18-2-10-2 /4/ Bb /3/ CK65-20*5 / W17-TRANS // TIFT /5/ P 2550	CHECK (SUS)	2000-01
3	AR 857-1-1	MADISON / YMI 6	BACON	2002-03
4	AR 857-1-2	MADISON / YMI 6	BACON	2002-03
5	AR 93019-2-1	VA88-52-69 / LOUVRIN 34	BACON	2002-03
6	AR93035-4-1	PIONEER 2548 / 4549-W1-2	BACON	2002-03
7	ARGE 97-1042-4-5	MASON / CATBIRD	MILUS	2002-03
8	ARGE 97-1033-3-5	FREEDOM / CATBIRD	MILUS	2002-03
9	ARGE 97-1048-3-6	MASON // SHA 3 / CATBIRD	MILUS	2002-03
10	ARGE 97-1038-3-5	MASON*2 // SHA3 / SUPER KAUZ	MILUS	2002-03
11	ARGE 97-1047-4-2	P2643 / 3 NING 7840 // PARULA / VEERY # 6	MILUS	2002-03
12	B980416	COKER 9543 / ABI 85-81	HANCOCK	2001-02
13	B980582	L881060 / L880436	HANCOCK	2001-02
14	B011066	COKER 9877 / 8268G1-18-4 // COKER 9474	HANCOCK	2002-03
15	B011117	YMI 6 / COKER 9877	HANCOCK	2002-03
16	GA 931587E53	84368 / FFR 518	JOHNSON	2002-03
17	GA 931630E48	FLEMING / P2580 // 84200-7	JOHNSON	2002-03
18	GA 94261E7	AGS 2000 // 841114 / FFR 518	JOHNSON	2002-03
19	GA 931233A24	GORE*2 / 83267	JOHNSON	2002-03
20	GA 941208E35	ROBERTS /4/ P 2580 // T83103*2 / HAMLET /3/ FLEMING	JOHNSON	2002-03
21	MV 5-46	VA91-54-222 / FFR 555W // VA 93-52-55, F5	COSTA	2002-03
22	MV 15-42	VA93-51-27 // VA91-54-343 / GA-GORE, F5	COSTA	2002-03
23	MV 27-78	VA90-52-26(LOV29/TYL/RCT*2/GAINES) / COKER 9835, F5	COSTA	2002-03
24	NC98-26192	P81401A1-42-1 / SALUDA /3/ P 2555 / COKER 9907 // MV14 / WAKEFIELD	MURPHY	2001-02
25	NC99-13296	STUCKY / MADISON // COKER 9474	MURPHY	2002-03
26	NC99-13308	STUCKY / MADISON // COKER 9474	MURPHY	2002-03
27	VAN98W342	COKER 983 // GA-ANDY / VA90-21-20(79IWWRN67//COKER65-20/ATR)	GRIFFEY	2002-03
28	VA00W562	PC-7(CHILL "S"/YM16:SCAB-RES) / PION2548 // PION2684	GRIFFEY	2001-02
29	VA00W566	PC-7(CHILL "S"/YM16:SCAB-RES) / PION2548 // PION2684	GRIFFEY	2001-02
30	VA01W461	PC-11(SHANGHAI4/CHILL"S":SCAB-RES)/3/ VA92-51-39 // FFR 555W / RCT /4/ COKER 9803	GRIFFEY	2001-02
31	VA02W713	NING 7840 / P2691 // ROANE	GRIFFEY	2002-03
32	VA02W732	PC-7(CHILL "S"/YM16:SCAB-RES)/3/ VA92-51-39 // CK9803 / RCT /4/ VA93-52-55	GRIFFEY	2002-03

FHB Incidence (1-100)

CULTIVAR/ DESIGNATION	BAY AR	COL'BIA MO	K'STON NC	B'BURG VA	URBANA IL	W'STER OH	LEX'TON KY	MEAN ALL LOC.
	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK
1 ERNIE	33 10	71 12	6 4	28 3	73 9	100 .	36 1	49.6 5
2 COKER 9835	70 30	100 32	43 26	70 28	100 31	100 .	62 15	77.8 32
3 AR 857-1-1	30 7	80 22	8 8	40 10	72 7	93 2	84 31	58.2 11
4 AR 857-1-2	22 3	63 7	12 12	33 7	68 4	93 2	46 3	48.3 3
5 AR 93019-2-1	50 21	60 5	13 15	62 24	82 14	98 8	52 11	59.6 15
6 AR93035-4-1	46 17	78 16	25 24	57 19	83 16	98 8	72 26	65.7 21
7 ARGE 97-1042-4-5	10 2	56 4	. .	23 2	77 11	100 .	47 4	44.4 2
8 ARGE 97-1033-3-5	8 1	51 3	13 16	30 4	60 3	100 .	48 6	44.3 1
9 ARGE 97-1048-3-6	28 6	48 2	8 7	43 13	72 8	100 .	62 17	51.5 7
10 ARGE 97-1038-3-5	32 9	63 6	8 9	40 11	77 12	98 8	65 19	54.7 10
11 ARGE 97-1047-4-2	24 4	70 10	4 1	33 8	70 5	100 .	54 12	50.7 6
12 B980416	48 18	44 1	10 10	32 6	73 10	93 2	65 20	52.1 9
13 B980582	32 8	65 9	7 6	22 1	70 6	100 .	48 5	49.1 4
14 B011066	50 20	71 11	17 19	43 14	82 15	98 8	55 13	59.4 14
15 B011117	40 14	64 8	22 22	40 12	55 1	92 1	50 8	51.8 8
16 GA 931587E53	72 31	95 30	7 5	85 31	99 29	100 .	68 23	75.1 30
17 GA 931630E48	52 24	80 21	. .	62 25	83 17	100 .	68 22	66.4 22
18 GA 94261E7	68 29	100 31	13 17	81 30	99 30	100 .	78 28	77.1 31
19 GA 931233A24	60 26	90 28	27 25	85 32	88 21	97 6	52 10	71.2 28
20 GA 941208E35	80 32	79 19	. .	33 9	98 27	100 .	81 30	70.9 27
21 MV 5-46	42 15	78 15	11 11	60 21	98 28	100 .	87 32	68.0 24
22 MV 15-42	58 25	80 20	. .	45 15	98 26	100 .	69 24	67.2 23
23 MV 27-78	52 23	94 29	23 23	63 26	92 22	100 .	71 25	70.7 26
24 NC98-26192	40 13	83 23	5 3	53 18	85 19	100 .	63 18	61.4 18
25 NC99-13296	52 22	86 25	. .	63 27	100 32	100 .	79 29	72.3 29
26 NC99-13308	38 12	85 24	12 13	47 17	94 25	100 .	49 7	60.7 17
27 VAN98W342	28 5	88 27	14 18	70 29	93 23	100 .	43 2	62.4 19
28 VA00W562	66 27	88 26	21 21	60 22	83 18	100 .	75 27	70.5 25
29 VA00W566	50 19	79 18	. .	30 5	78 13	100 .	62 16	58.8 13
30 VA01W461	68 28	75 14	19 20	45 16	57 2	93 2	51 9	58.3 12
31 VA02W713	34 11	74 13	5 2	60 23	85 20	97 6	66 21	60.1 16
32 VA02W732	44 16	79 17	12 14	58 20	93 24	100 .	59 14	63.7 20

Mean:	45	75	14	50	79	98	62	61.0
L.S.D. (0.05)	28	33	10	25	16	7	35	10.9
CV%	31.0	16.0	50.5	36.3	11.0	22.0	27.1	16.7

FHB Severity (1-100)


Cultivar/ Designation	BAY AR	F'VILLE AR	KIBLER AR	COL'BIA MO	K'STON NC	B'BURG VA	URBANA IL	W'STER OH	LEX'TON KY	MEAN ALL LOC.
	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK
1 ERNIE	12 8	44 16	40 17	8 14	2 3	16 12	19 6	58 20	13 2	23.5 13
2 COKER 9835	35 28	74 28	81 32	43 32	51 26	28 29	81 31	96 31	26 30	57.3 32
3 AR 857-1-1	28 25	24 4	25 4	9 17	2 4	16 13	12 3	19 2	18 11	16.9 3
4 AR 857-1-2	14 12	25 5	26 6	4 4	3 11	16 10	19 5	22 4	14 4	15.7 2
5 AR 93019-2-1	11 7	46 18	36 12	6 9	3 13	25 27	25 14	54 18	19 17	25.0 16
6 AR93035-4-1	18 22	38 14	53 21	10 20	4 20	24 25	24 11	46 15	23 27	26.7 18
7 ARGE 97-1042-4-5	4 1	30 10	38 13	5 5	. .	12 3	24 13	40 11	23 25	19.2 9
8 ARGE 97-1033-3-5	9 4	20 3	16 1	2 1	1 2	14 6	11 2	14 1	8 1	10.7 1
9 ARGE 97-1048-3-6	10 5	19 2	29 7	3 2	2 5	18 17	16 4	45 14	17 9	17.7 7
10 ARGE 97-1038-3-5	16 16	36 13	34 10	5 6	4 18	13 5	23 10	37 9	16 5	20.4 11
11 ARGE 97-1047-4-2	15 13	28 6	24 3	6 10	1 1	10 2	22 9	34 7	13 3	17.0 5
12 B980416	18 19	43 15	34 11	3 3	2 10	15 9	26 15	30 5	19 14	21.1 12
13 B980582	7 2	29 9	25 5	5 7	2 6	9 1	21 7	51 17	19 18	18.7 8
14 B011066	11 6	28 7	31 8	6 11	3 14	20 19	24 12	31 6	18 13	19.2 10
15 B011117	12 9	18 1	40 16	5 8	4 21	15 7	21 8	20 3	21 20	17.2 6
16 GA 931587E53	17 18	71 27	61 27	13 23	2 7	37 32	84 32	97 32	22 23	44.8 27
17 GA 931630E48	33 26	44 17	48 19	16 28	. .	24 26	43 22	89 25	19 15	36.6 23
18 GA 94261E7	27 24	68 25	63 28	38 31	3 15	28 30	81 30	94 29	27 31	47.6 31
19 GA 931233A24	33 27	76 29	73 30	19 30	21 25	34 31	47 25	92 27	22 21	46.3 29
20 GA 941208E35	36 29	76 30	75 31	14 25	. .	15 8	65 28	93 28	23 26	46.8 30
21 MV 5-46	44 32	77 31	53 22	13 24	3 16	23 24	63 27	86 24	46 32	45.3 28
22 MV 15-42	23 23	77 32	53 23	9 18	. .	17 14	74 29	74 21	25 29	41.1 25
23 MV 27-78	43 31	55 23	51 20	14 26	4 19	20 18	43 21	81 23	19 19	36.6 24
24 NC98-26192	18 20	57 24	39 15	7 12	2 8	21 20	42 20	35 8	17 7	26.4 17
25 NC99-13296	16 15	49 20	45 18	10 21	. .	25 28	46 24	89 26	24 28	35.2 22
26 NC99-13308	12 10	53 21	33 9	10 22	3 12	22 23	28 17	44 13	16 6	24.5 15
27 VAN98W342	9 3	48 19	55 24	9 19	5 24	22 22	44 23	58 19	19 16	29.8 20
28 VA00W562	16 17	53 22	65 29	14 27	3 17	21 21	40 19	50 16	17 8	31.1 21
29 VA00W566	18 21	33 11	61 26	8 15	. .	12 4	28 16	74 22	22 22	29.2 19
30 VA01W461	13 11	28 8	16 2	7 13	5 22	18 15	11 1	39 10	17 10	17.0 4
31 VA02W713	15 14	35 12	38 14	8 16	2 9	16 11	37 18	42 12	23 24	23.9 14
32 VA02W732	40 30	70 26	56 25	16 29	5 23	18 16	56 26	95 30	18 12	41.5 26

Mean:	20	46	44	11	5	20	35	57	20	29.1
L.S.D. (0.05)	14	15	13	5	6	10	13	20	16	10.2
CV%	35	28	27	34	75	36	20	22	37	37.4

FHB SEVERITY GGE BI PLOT ANALYSIS¹

Cultivar/ Designation	MEGA- ENVIRON.		W'STER	OH	MEAN	
	1	RANK			RANK	ALL LOC.
1 ERNIE	19.2	12	58	20	23.5	13
2 COKER 9835	52.4	32	96	31	57.3	32
3 AR 857-1-1	16.7	7	19	2	16.9	3
4 AR 857-1-2	14.9	6	22	4	15.7	2
5 AR 93019-2-1	21.4	14	54	18	25.0	16
6 AR93035-4-1	24.3	18	46	15	26.7	18
7 ARGE 97-1042-4-5	16.7	8	40	11	19.2	9
8 ARGE 97-1033-3-5	10.3	1	14	1	10.7	1
9 ARGE 97-1048-3-6	14.4	3	45	14	17.7	7
10 ARGE 97-1038-3-5	18.3	11	37	9	20.4	11
11 ARGE 97-1047-4-2	14.9	5	34	7	17.0	5
12 B980416	19.9	13	30	5	21.1	12
13 B980582	14.6	4	51	17	18.7	8
14 B011066	17.7	10	31	6	19.2	10
15 B011117	16.9	9	20	3	17.2	6
16 GA 931587E53	38.3	27	97	32	44.8	27
17 GA 931630E48	29.6	23	89	25	36.6	23
18 GA 94261E7	41.8	31	94	29	47.6	31
19 GA 931233A24	40.6	29	92	27	46.3	29
20 GA 941208E35	40.6	30	93	28	46.8	30
21 MV 5-46	40.2	28	86	24	45.3	28
22 MV 15-42	36.9	26	74	21	41.1	25
23 MV 27-78	31.1	24	81	23	36.6	24
24 NC98-26192	25.3	19	35	8	26.4	17
25 NC99-13296	27.9	21	89	26	35.2	22
26 NC99-13308	22.1	16	44	13	24.5	15
27 VAN98W342	26.3	20	58	19	29.8	20
28 VA00W562	28.7	22	50	16	31.1	21
29 VA00W566	23.2	17	74	22	29.2	19
30 VA01W461	14.3	2	39	10	17.0	4
31 VA02W713	21.7	15	42	12	23.9	14
32 VA02W732	34.7	25	95	30	41.5	26

Mean: 25.9 57 29.1
L.S.D. (0.05) 9.4 20 10.2
CV% 37.3 22 37.4



Although not strictly in a separate Mega-environment, the OH location had the largest within location standard deviation, and was investigated because of its discriminating ability relative to the other test locations. There was good agreement between the OH location and the mean severity over the other eight locations with respect to the most resistant genotypes. All seven of the entries not significantly different from the most resistant Entry 8 in OH were among the most resistant group over the remaining locations.

¹Yan et al., (2000). Crop Sci. 40:597-605.

**HEAD SEVERITY EXPRESSED AS AREA UNDER THE DISEASE PROGRESS CURVE (AUDPC).
FUNDULEA, ROMANIA**

Cultivar/ Designation	AUDPC ¹	RANK
1 ERNIE	140	10
2 COKER 9835	460	21
3 AR 857-1-1	.	.
4 AR 857-1-2	.	.
5 AR 93019-2-1	628	24
6 AR93035-4-1	671	26
7 ARGE 97-1042-4-5	168	13
8 ARGE 97-1033-3-5	52	1
9 ARGE 97-1048-3-6	73	2
10 ARGE 97-1038-3-5	209	15
11 ARGE 97-1047-4-2	467	22
12 B980416	518	23
13 B980582	77	3
14 B011066	128	9
15 B011117	.	.
16 GA 931587E53	244	18
17 GA 931630E48	165	12
18 GA 94261E7	123	8
19 GA 931233A24	.	.
20 GA 941208E35	264	19
21 MV 5-46	669	25
22 MV 15-42	230	17
23 MV 27-78	209	16
24 NC98-26192	92	7
25 NC99-13296	208	14
26 NC99-13308	149	11
27 VAN98W342	80	5
28 VA00W562	78	4
29 VA00W566	.	.
30 VA01W461	.	.
31 VA02W713	278	20
32 VA02W732	88	6

Mean: 249
L.S.D. (0.05) 102

FHB Index (1-100)

Cultivar/ Designation	BAY	COL'BIA	K'STON	B'BURG	URBANA	W'STER	LEX'TON	SZEGED ¹	MEAN									
	AR	MO	NC	VA	IL	OH	KY	HUN	ALL LOC.									
	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK									
1 ERNIE	3.5	8	6.1	14	0.1	5.0	7	14.3	8	58.1	20	6.0	2	18.0	7	13.9	9	
2 COKER 9835	24.6	31	43.1	32	21.7	.	20.7	29	80.7	31	96.3	31	24.5	28	79.7	32	48.9	32
3 AR 857-1-1	8.4	19	7.3	17	0.1	.	8.3	14	8.6	3	17.6	2	18.9	23	18.9	8	11.0	5
4 AR 857-1-2	2.9	6	2.4	4	0.4	.	6.3	10	13.1	6	20.3	4	8.3	3	26.6	9	10.0	3
5 AR 93019-2-1	5.4	13	3.9	9	0.5	.	17.3	27	20.6	15	52.8	18	12.6	9	41.9	21	19.4	16
6 AR93035-4-1	8.2	17	8.4	20	1.1	.	14.0	22	20.0	13	45.3	15	21.5	26	68.1	30	23.3	21
7 ARGE 97-1042-4-5	0.7	1	2.7	5	.	.	3.3	2	19.5	11	39.8	11	13.5	14	11.2	3	10.2	4
8 ARGE 97-1033-3-5	0.7	2	1.4	1	0.2	.	4.0	5	6.8	2	14.1	1	5.3	1	3.1	1	4.5	1
9 ARGE 97-1048-3-6	2.9	5	1.5	3	0.1	.	8.0	12	11.5	5	44.6	14	14.6	15	33.5	13	14.6	11
10 ARGE 97-1038-3-5	5.1	12	3.2	7	0.3	.	6.0	9	18.1	10	36.9	9	12.8	11	15.3	5	12.2	7
11 ARGE 97-1047-4-2	3.3	7	4.4	11	0.0	.	3.7	3	15.4	9	34.3	7	9.2	4	8.2	2	9.8	2
12 B980416	8.1	16	1.4	2	0.2	.	5.0	8	20.3	14	29.3	5	15.3	16	31.8	12	13.9	10
13 B980582	2.2	3	3.0	6	0.1	.	2.0	1	14.2	7	51.3	17	10.7	7	14.7	4	12.3	8
14 B011066	5.5	14	4.2	10	0.5	.	9.7	15	19.7	12	30.6	6	12.6	10	46.8	22	16.2	13
15 B011117	4.8	10	3.7	8	1.0	.	6.3	11	11.2	4	18.8	3	13.2	12	34.5	15	11.7	6
16 GA 931587E53	12.7	23	11.9	25	0.1	.	31.3	32	88.1	32	96.8	32	19.2	24	55.1	27	39.4	29
17 GA 931630E48	18.0	26	13.1	29	.	.	17.0	26	35.7	21	88.8	25	16.6	20	38.2	19	29.7	24
18 GA 94261E7	18.6	27	38.3	31	0.4	.	24.3	30	80.0	30	94.0	29	26.7	30	62.1	28	43.1	31
19 GA 931233A24	19.7	29	17.6	30	8.2	.	28.7	31	40.7	23	88.9	26	15.5	17	78.3	31	37.2	28
20 GA 941208E35	41.1	32	10.8	24	.	.	4.7	6	63.6	28	93.4	28	24.9	29	64.7	29	40.5	30
21 MV 5-46	19.3	28	9.8	23	0.3	.	17.0	24	61.8	27	86.3	24	50.8	32	39.3	20	35.6	27
22 MV 15-42	13.5	24	7.3	18	.	.	11.0	18	72.3	29	73.5	21	26.9	31	36.5	17	31.6	25
23 MV 27-78	22.3	30	13.0	28	0.8	.	13.7	21	39.1	22	80.5	23	16.9	21	37.9	18	28.0	22
24 NC98-26192	7.1	15	6.2	15	0.1	.	13.3	20	35.7	20	35.1	8	13.4	13	17.3	6	16.0	12
25 NC99-13296	8.3	18	8.8	21	.	.	18.7	28	46.0	25	89.3	27	24.4	27	28.1	10	29.1	23
26 NC99-13308	4.4	9	9.3	22	0.3	.	10.7	17	26.9	17	43.6	13	10.3	5	34.1	14	17.5	15
27 VAN98W342	2.6	4	7.8	19	0.7	.	17.0	25	40.8	24	58.0	19	10.4	6	30.8	11	21.0	18
28 VA00W562	10.4	22	13.0	27	0.8	.	15.0	23	33.8	19	49.6	16	16.1	18	36.0	16	21.8	19
29 VA00W566	9.0	21	6.3	16	.	.	3.7	4	21.7	16	74.3	22	17.6	22	46.9	23	22.8	20
30 VA01W461	8.9	20	5.7	12	0.9	.	8.0	13	6.1	1	37.2	10	11.5	8	54.5	26	16.6	14
31 VA02W713	5.0	11	6.0	13	0.1	.	10.0	16	31.8	18	41.0	12	21.4	25	50.5	25	20.7	17
32 VA02W732	17.7	25	12.0	26	0.6	.	13.0	19	52.4	26	95.4	30	16.5	19	47.2	24	31.9	26

Mean: 10.2 9.2 1.7 11.8 29.9 57.0 16.8 37.8 22.7
L.S.D. (0.05) 10.4 10.7 5.8 62.1 12.5 20.5 20.0 2.14 11.7
CV% 50.0 41.4 77.8 10.0 22.9 22.0 61.4 . 52.5

¹DATA BY INDIVIDUAL ISOLATES ON FOLLOWING PAGE.

**INDEX BY INDIVIDUAL ISOLATES,
SZEGED, HUNGARY**


Cultivar/ Designation	<i>F. gramin.</i> Isol. 12377	<i>F. gramin.</i> Isol. 44	<i>F. culmor.</i> Isol. 12375	<i>F. culmor.</i> Isol. 12551	MEAN	
	RANK	RANK	RANK	RANK	All Isol.	RANK
1 ERNIE	11	5	11	9	37	7
2 COKER 9835	75	32	70	32	91	31
3 AR 857-1-1	14	6	6	8	44	8
4 AR 857-1-2	23	10	15	9	45	9
5 AR 93019-2-1	38	20	32	17	36	22
6 AR93035-4-1	67	30	61	29	62	30
7 ARGE 97-1042-4-5	11	4	8	4	6	2
8 ARGE 97-1033-3-5	1	1	2	1	1	1
9 ARGE 97-1048-3-6	23	12	29	14	36	20
10 ARGE 97-1038-3-5	14	7	14	8	9	5
11 ARGE 97-1047-4-2	4	2	5	2	8	4
12 B980416	36	18	36	21	10	7
13 B980582	15	8	14	6	6	3
14 B011066	52	26	42	24	34	19
15 B011117	27	13	21	11	31	18
16 GA 931587E53	50	25	43	25	51	29
17 GA 931630E48	35	17	32	18	28	16
18 GA 94261E7	65	29	60	28	44	25
19 GA 931233A24	72	31	67	31	82	31
20 GA 941208E35	64	28	62	30	47	26
21 MV 5-46	31	14	29	15	30	17
22 MV 15-42	36	19	33	19	19	10
23 MV 27-78	38	21	33	20	22	12
24 NC98-26192	11	3	14	7	10	6
25 NC99-13296	21	9	18	10	23	13
26 NC99-13308	34	16	31	16	23	14
27 VAN98W342	32	15	22	12	22	11
28 VA00W562	23	11	23	13	42	24
29 VA00W566	42	22	40	23	39	23
30 VA01W461	57	27	40	22	51	28
31 VA02W713	45	23	43	26	48	27
32 VA02W732	45	24	48	27	37	22
Mean:	35	31	31	54	37.8	
L.S.D. (0.05)	2.1	

Scabby Seed %

Cultivar/ Designation	COL'BIA	K'STON	B'BURG	KIBLER	LEX'TON	FUN'LEA	MEAN
	MO	NC	VA	AR	KY	ROM	ALL LOC.
	RANK	RANK	RANK	RANK	RANK	RANK	RANK
1 ERNIE	1 1	18 13	46 11	71 8	9 10	20 12	27.6 8
2 COKER 9835	29 32	44 25	83 32	100 32	17 21	75 24	58.0 32
3 AR 857-1-1	2 4	18 12	66 23	70 7	5 2	. .	32.1 14
4 AR 857-1-2	1 1	7 3	52 13	53 1	9 9	. .	23.4 3
5 AR 93019-2-1	4 14	27 18	58 20	83 17	19 23	89 26	46.7 30
6 AR93035-4-1	4 14	38 22	67 24	90 25	13 15	86 25	49.6 31
7 ARGE 97-1042-4-5	2 4	. .	27 2	88 21	9 7	40 21	30.2 10
8 ARGE 97-1033-3-5	1 1	14 10	40 9	59 5	30 27	7 2	25.2 4
9 ARGE 97-1048-3-6	2 4	9 6	45 10	83 16	7 5	16 9	27.0 6
10 ARGE 97-1038-3-5	2 4	17 11	39 8	83 15	6 3	17 11	27.3 7
11 ARGE 97-1047-4-2	2 4	9 7	24 1	68 6	16 20	72 23	31.9 12
12 B980416	3 10	12 9	33 5	81 14	36 29	32 20	32.9 16
13 B980582	2 4	8 5	28 3	54 2	4 1	4 1	16.6 1
14 B011066	3 10	38 23	49 12	88 22	9 8	16 10	33.7 20
15 B011117	3 10	35 21	57 18	95 29	15 19	. .	40.2 23
16 GA 931587E53	12 30	30 20	79 30	88 24	51 32	12 7	45.4 27
17 GA 931630E48	7 27	. .	55 16	75 9	12 13	23 14	31.5 11
18 GA 94261E7	28 31	19 15	80 31	90 27	24 25	29 18	44.9 26
19 GA 931233A24	8 28	42 24	70 25	100 31	12 14	. .	45.6 29
20 GA 941208E35	4 14	. .	37 7	95 28	48 31	26 16	41.2 25
21 MV 5-46	5 23	19 17	63 22	83 19	18 22	57 22	40.9 24
22 MV 15-42	4 14	. .	29 4	78 10	13 17	22 13	26.4 5
23 MV 27-78	10 29	45 26	74 26	95 30	33 28	15 8	45.4 28
24 NC98-26192	4 14	19 16	53 14	80 13	13 16	12 6	30.1 9
25 NC99-13296	4 14	. .	57 19	78 12	6 4	29 19	32.1 13
26 NC99-13308	4 14	6 1	34 6	56 3	15 18	24 15	23.1 2
27 VAN98W342	5 23	6 2	79 29	88 23	10 12	11 5	33.2 17
28 VA00W562	5 23	18 14	77 27	85 20	20 24	7 3	35.4 21
29 VA00W566	3 10	. .	56 17	78 11	37 30	. .	38.8 22
30 VA01W461	6 26	30 19	54 15	56 4	25 26	. .	33.3 18
31 VA02W713	4 14	7 4	60 21	83 18	10 11	28 17	32.2 15
32 VA02W732	4 14	12 8	77 28	90 26	8 6	8 4	33.4 19
Mean:	5	21	55	81	18	30	34.8
L.S.D. (0.05)	7	26	19	13	.	10	15.2
CV%	91	62	26	.	45	.	38.1

Scabby Seed % GGE BI PLOT ANALYSIS¹

Cultivar/ Designation	MEGA- ENVIRON.		MEGA- ENVIRON.		MEAN ALL LOC. RANK
	1 RANK	2 RANK	1 RANK	2 RANK	
1 ERNIE	29	8	20	12	27.6 8
2 COKER 9835	55	32	75	24	58.0 32
3 AR 857-1-1	32	11	.	.	32.1 14
4 AR 857-1-2	24	4	.	.	23.4 3
5 AR 93019-2-1	38	21	89	26	46.7 30
6 AR93035-4-1	42	26	86	25	49.6 31
7 ARGE 97-1042-4-5	28	6	40	21	30.2 10
8 ARGE 97-1033-3-5	29	7	7	2	25.2 4
9 ARGE 97-1048-3-6	29	9	16	9	27.0 6
10 ARGE 97-1038-3-5	29	10	17	11	27.3 7
11 ARGE 97-1047-4-2	24	3	72	23	31.9 12
12 B980416	33	14	32	20	32.9 16
13 B980582	19	1	4	1	16.6 1
14 B011066	37	18	16	10	33.7 20
15 B011117	41	25	.	.	40.2 23
16 GA 931587E53	52	31	12	7	45.4 27
17 GA 931630E48	34	15	23	14	31.5 11
18 GA 94261E7	48	29	29	18	44.9 26
19 GA 931233A24	46	28	.	.	45.6 29
20 GA 941208E35	42	27	26	16	41.2 25
21 MV 5-46	38	20	57	22	40.9 24
22 MV 15-42	27	5	22	13	26.4 5
23 MV 27-78	52	30	15	8	45.4 28
24 NC98-26192	34	16	12	6	30.1 9
25 NC99-13296	33	12	29	19	32.1 13
26 NC99-13308	23	2	24	15	23.1 2
27 VAN98W342	38	19	11	5	33.2 17
28 VA00W562	41	24	7	3	35.4 21
29 VA00W566	40	23	.	.	38.8 22
30 VA01W461	34	17	.	.	33.3 18
31 VA02W713	33	13	28	17	32.2 15
32 VA02W732	38	22	8	4	33.4 19
Mean:	36.3	30	34.8		
L.S.D. (0.05)	13.2	10	15.2		
CV%	29.2	.	38.1		



Two Mega-environments were identified for Scabby Seed Percent:

Mega-environment 1: VA, AR (Kibler), NC, KY, MO.

Mega-environment 2: Romania alone.

Ten entries were not significantly different from B980582 (Entry 13), the most resistant line in Mega-environment 1. B980582 was the most resistant entry in Mega-environment 2, also.

The Va location exhibited the most discriminating ability within mega-environment 1.

GRAIN QUALITY

Cultivar/ Designation	GRAIN RATING ¹ URBANA IL	GRAIN QUALITY ² K'STON NC	RELATIVE GR. WT. ³ FUN'LEA ROM	1000 GR. WT. SZEGED HUN	TEST WEIGHT COL'BIA MO	GRAIN EVAL. ⁴ SZEGED HUN	%WET GLUTEN ⁵ SZEGED HUN	HARD- NESS ⁵ SZEGED HUN
1 ERNIE	3.7 3	1.3 4	85 12	35.8 15	58.3 18	vb5	27.1	5.0
2 COKER 9835	8.3 28	3.8 26	74 16	31.3 27	48.7 32	bsb4.5	.	.
3 AR 857-1-1	4.3 8	1.8 15	.	41.0 5	58.0 19	b4.5	34.7	15.2
4 AR 857-1-2	3.7 4	1.8 15	.	36.8 13	55.0 26	b4A1	33.8	14.1
5 AR 93019-2-1	7.0 23	2.3 19	43 24	34.8 18	58.5 16	b4A1	30.6	15.9
6 AR93035-4-1	4.3 9	2.0 17	24 26	34.2 20	58.0 20	bsb4	25.0	45.1
7 ARGE 97-1042-4-5	5.3 16	.	74 17	37.9 10	61.0 2	b4	34.3	33.0
8 ARGE 97-1033-3-5	4.7 12	1.5 8	99 1	31.2 29	59.5 8	bsb5A5	29.1	27.7
9 ARGE 97-1048-3-6	3.3 2	1.3 4	97 4	41.3 4	52.7 30	b4.5	33.0	30.2
10 ARGE 97-1038-3-5	3.7 5	1.5 8	86 11	35.2 16	59.6 5	b5	30.2	12.4
11 ARGE 97-1047-4-2	3.7 6	1.5 8	39 25	43.1 2	59.6 6	b5	32.6	12.1
12 B980416	6.7 22	1.5 8	63 20	27.9 31	59.0 12	b5	22.1	0.57
13 B980582	2.7 1	1.0 1	97 5	34.8 19	61.2 1	b5	.	.
14 B011066	4.3 10	3.0 22	90 7	30.8 30	59.6 7	b3-5	28.9	22.3
15 B011117	5.0 13	3.0 22	.	43.5 1	59.3 11	bsb4.5	33.7	44.4
16 GA 931587E53	8.3 29	2.0 17	89 8	34.0 23	55.8 24	vb4.5	27.5	6.92
17 GA 931630E48	6.3 19	.	81 13	37.1 12	56.6 23	bsb4-5	26.3	13.5
18 GA 94261E7	8.7 31	1.3 4	93 6	32.4 24	53.7 29	vb-sb5	29.8	4.27
19 GA 931233A24	7.0 24	3.0 22	.	31.7 25	54.6 27	vb5	29.6	17.2
20 GA 941208E35	8.3 30	.	62 21	26.5 32	58.8 14	vb5	27.5	13.4
21 MV 5-46	8.7 32	1.0 1	44 23	37.3 11	59.4 9	b4	28.3	28.8
22 MV 15-42	7.7 26	.	77 15	42.8 3	57.5 21	b5!	30.3	20.8
23 MV 27-78	6.3 20	2.3 19	81 14	36.0 14	53.9 28	bsb4.5A1	28.8	29.0
24 NC98-26192	6.3 21	1.5 8	67 19	38.8 7	58.5 17	vb5	.	.
25 NC99-13296	5.0 14	.	70 18	31.3 28	58.7 15	f5	26.2	0.55
26 NC99-13308	4.0 7	1.3 4	99 2	38.4 9	59.8 4	b5	28.9	2.74
27 VAN98W342	7.7 27	1.5 8	98 3	35.2 17	56.8 22	b4	26.0	6.69
28 VA00W562	6.0 17	2.8 21	89 9	34.2 21	55.2 25	b5	27.2	15.8
29 VA00W566	6.0 18	.	.	38.5 8	52.5 31	f5	29.2	21.3
30 VA01W461	4.3 11	3.0 22	.	40.9 6	59.4 10	bsb4.5	33.5	33.3
31 VA02W713	5.0 15	1.0 1	61 22	34.2 22	61.0 3	b5	21.1	20.9
32 VA02W732	7.0 25	1.5 8	87 10	31.5 26	58.9 13	bsb4.5A5	27.5	17.4

Mean: 5.2 1.9 76 36 57.5 . 29.1 18.3
L.S.D. (0.05) 1.8 1.1 13.2 . 5.8 . . .
CV% 21.7 29.9 . . 7.2 . . .

¹SCALE OF 0-9. ²SCALE OF 1-5. ³(INOCULATED / UNINOCULATED)*100.

⁴vb=LIGHT BROWN, b=BROWN, vb=DARK BROWN, bsb=BROWN-DARK-BROWN, f=WHITE. 1=VERY SHRIVELLED 5=NO SHRIVELLING.

⁵NIR.

**Vomitoxin (DON)*
(ppm)**

Cultivar/ Designation	BAY AR	K'STON NC	B'BURG VA	LEX'TON KY	URBANA IL	MEAN ALL LOC.
	RANK	RANK	RANK	RANK	RANK	RANK
1 ERNIE	2.2 16	0.0 1	6.5 18	17.3 26	32.5 22	11.7 22
2 COKER 9835	4.4 26	7.0 24	13.0 27	21.3 31	52.5 28	19.6 29
3 AR 857-1-1	3.5 23	0.5 11	3.8 8	10.0 7	8.5 2	5.2 2
4 AR 857-1-2	1.7 11	0.6 14	2.9 6	11.3 10	11.0 5	5.5 6
5 AR 93019-2-1	2.3 17	5.0 23	6.7 20	16.7 24	24.0 16	10.9 20
6 AR93035-4-1	2.2 15	4.4 22	5.3 13	16.7 23	15.5 10	8.8 13
7 ARGE 97-1042-4-5	1.0 3	. .	3.6 7	9.7 5	19.0 12	6.3 8
8 ARGE 97-1033-3-5	1.6 9	1.8 15	4.1 9	9.0 4	10.5 4	5.4 5
9 ARGE 97-1048-3-6	3.7 25	1.8 15	4.3 10	8.3 2	8.0 1	5.2 3
10 ARGE 97-1038-3-5	1.5 6	0.5 11	4.7 12	6.7 1	17.5 11	6.1 7
11 ARGE 97-1047-4-2	1.7 10	0.0 1	2.1 3	8.7 3	14.0 9	5.3 4
12 B980416	1.3 4	1.8 15	4.6 11	11.3 11	19.5 13	7.7 10
13 B980582	1.6 7	0.0 1	1.9 1	11.0 9	9.0 3	4.7 1
14 B011066	2.6 19	4.2 21	2.5 4	16.4 22	11.0 6	7.3 9
15 B011117	5.4 30	10.0 25	2.7 5	18.1 28	11.0 7	9.4 14
16 GA 931587E53	3.6 24	0.0 1	16.3 30	16.0 21	90.0 32	25.2 32
17 GA 931630E48	2.6 20	. .	14.3 29	18.3 29	40.0 27	16.7 28
18 GA 94261E7	2.5 18	2.0 18	16.3 31	22.7 32	72.5 31	23.2 31
19 GA 931233A24	8.7 32	11.0 26	13.0 28	17.7 27	30.0 21	16.1 26
20 GA 941208E35	8.4 31	. .	1.9 2	14.3 16	62.5 30	19.7 30
21 MV 5-46	2.1 14	0.0 1	9.7 25	15.0 20	52.5 29	15.9 25
22 MV 15-42	5.4 29	. .	5.7 15	18.7 30	35.0 24	14.1 24
23 MV 27-78	1.6 8	0.0 1	7.0 21	14.3 17	26.5 19	9.9 17
24 NC98-26192	0.6 1	0.5 11	7.8 22	17.0 25	23.5 15	9.8 16
25 NC99-13296	3.2 22	. .	18.8 32	13.0 12	37.5 26	16.1 27
26 NC99-13308	0.9 2	0.0 1	5.7 14	14.7 18	20.0 14	8.3 12
27 VAN98W342	2.1 13	0.0 1	12.3 26	9.7 6	24.5 17	9.7 15
28 VA00W562	4.6 27	2.8 19	6.5 19	14.7 19	25.0 18	10.7 19
29 VA00W566	5.3 28	. .	9.3 24	10.3 8	32.5 23	12.3 23
30 VA01W461	2.8 21	4.0 20	6.3 17	14.0 14	11.5 8	7.7 11
31 VA02W713	1.7 12	0.0 1	7.9 23	14.0 15	27.5 20	10.2 18
32 VA02W732	1.5 5	0.0 1	6.3 16	14.0 13	35.0 25	11.4 21
Mean:	2.9	2.2	7.3	14	26.1	11.3
L.S.D. (0.05)	3	.	54.9	11.5	11.1	10.8
CV%	51	.	5.5	4.1	21.0	76.9

*DON analysis conducted by Pat Hart's laboratory at Michigan State University.

Greenhouse Screening¹

Cultivar/ Designation	NC SEVERITY	VA SEVERITY	AR SEVERITY	KY SEVERITY	MO SEVERITY	MEAN ALL LOC.	NC SPREAD	MO SPREAD	MEAN SPREAD
	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK
1 ERNIE	9 9	4 1	18 12	19 1	23 14	14.6 2	1.4 8	2.3 10	1.8 6
2 COKER 9835	17 14	23 11	57 28	61 18	74 30	46.2 24	2.9 15	11.1 32	7.0 28
3 AR 857-1-1	16 12	38 24	57 29	30 3	11 5	30.5 14	2.3 11	1.2 2	1.7 4
4 AR 857-1-2	18 17	66 29	75 31	63 21	30 20	50.7 27	2.5 12	3.8 19	3.2 16
5 AR 93019-2-1	52 31	34 22	49 27	83 31	47 25	52.9 30	7.8 29	6.9 27	7.3 29
6 AR93035-4-1	17 13	29 19	34 20	68 23	19 11	33.3 18	3.3 16	2.8 13	3.0 15
7 ARGE 97-1042-4-5	38 27	7 3	9 2	40 5	21 12	22.9 3	6.9 28	3.0 15	4.9 23
8 ARGE 97-1033-3-5	15 11	6 2	6 1	38 4	6 1	14.3 1	3.6 18	0.9 1	2.3 10
9 ARGE 97-1048-3-6	4 1	38 25	12 6	52 12	9 2	23.0 5	0.8 2	1.2 3	1.0 1
10 ARGE 97-1038-3-5	4 2	26 16	32 19	92 32	21 13	35.0 22	0.8 3	3.1 16	1.9 7
11 ARGE 97-1047-4-2	9 8	64 28	35 21	70 24	44 24	44.3 23	1.5 9	5.6 22	3.6 19
12 B980416	4 3	18 5	10 3	57 14	26 16	23.0 4	0.6 1	3.5 18	2.1 8
13 B980582	22 20	25 15	22 16	40 6	13 6	24.2 6	3.5 17	1.8 7	2.7 11
14 B011066	22 21	20 9	15 9	51 11	13 7	24.4 8	4.0 21	1.9 8	3.0 13
15 B011117	5 5	75 30	11 4	70 25	11 4	34.4 20	0.9 4	1.5 5	1.2 2
16 GA 931587E53	18 16	27 17	17 11	46 8	48 27	31.2 15	3.8 19	5.9 24	4.8 21
17 GA 931630E48	49 29	19 6	41 25	73 27	73 29	51.0 28	8.9 31	10.4 30	9.6 32
18 GA 94261E7	17 15	31 20	48 26	57 13	82 32	46.9 25	2.8 13	9.1 29	5.9 25
19 GA 931233A24	74 32	84 32	81 32	60 17	53 28	70.6 32	10.9 32	7.1 28	9.0 31
20 GA 941208E35	49 30	47 26	60 30	63 20	48 26	53.4 31	6.5 26	5.6 23	6.1 26
21 MV 5-46	19 19	78 31	23 17	62 19	77 31	52.1 29	4.3 22	10.6 31	7.4 30
22 MV 15-42	26 23	34 23	20 14	23 2	27 18	26.2 11	4.4 23	3.8 20	4.1 20
23 MV 27-78	38 26	48 27	39 23	73 28	42 23	47.9 26	6.8 27	6.4 26	6.6 27
24 NC98-26192	29 25	23 12	11 5	67 22	36 21	33.3 17	5.1 25	4.6 21	4.9 22
25 NC99-13296	47 28	10 4	16 10	73 26	10 3	31.3 16	8.6 30	1.3 4	5.0 24
26 NC99-13308	25 22	32 21	20 15	59 16	14 8	30.0 13	3.9 20	1.5 6	2.7 12
27 VAN98W342	19 18	19 7	20 13	58 15	26 17	28.6 12	2.8 14	3.2 17	3.0 14
28 VA00W562	11 10	24 14	13 7	46 7	28 19	24.3 7	1.5 10	2.9 14	2.2 9
29 VA00W566	6 7	22 10	40 24	77 30	25 15	34.0 19	1.1 7	2.4 11	1.8 5
30 VA01W461	5 4	20 8	38 22	49 10	16 10	25.5 10	0.9 5	2.5 12	1.7 3
31 VA02W713	26 24	23 13	14 8	47 9	15 9	25.0 9	4.6 24	1.9 9	3.3 17
32 VA02W732	5 6	28 18	26 18	76 29	38 22	34.7 21	0.9 6	5.9 25	3.4 18

Mean:

22

34

30

58

32

35

3.8

4.2

4.0

L.S.D. (0.05)

.

.

30

29

.

20

.

4.5

CV%

.

.

22

.

46

.


57.2

¹Severity data based on the percentage of infected spikelets/total spikelets 21 to 28 days post inoculation.

Spread = total number of diseased spikelets in a head.

Greenhouse Head Severity GGE Biplot Analysis¹

Cultivar/ Designation	MEGA- ENVIRON.		MEGA- ENVIRON. RANK	MEAN ALL LOC. RANK
	1	2		
1 ERNIE	14	1	16	9
2 COKER 9835	47	23	45	26
3 AR 857-1-1	42	17	14	6
4 AR 857-1-2	68	31	24	18
5 AR 93019-2-1	55	28	49	29
6 AR93035-4-1	44	19	18	11
7 ARGE 97-1042-4-5	19	3	29	22
8 ARGE 97-1033-3-5	17	2	11	4
9 ARGE 97-1048-3-6	34	14	7	1
10 ARGE 97-1038-3-5	50	24	12	5
11 ARGE 97-1047-4-2	56	29	26	19
12 B980416	28	7	15	7
13 B980582	29	9	17	10
14 B011066	29	8	18	12
15 B011117	52	25	8	2
16 GA 931587E53	30	10	33	24
17 GA 931630E48	44	20	61	31
18 GA 94261E7	45	21	50	30
19 GA 931233A24	75	32	64	32
20 GA 941208E35	57	30	49	28
21 MV 5-46	55	27	48	27
22 MV 15-42	26	4	27	20
23 MV 27-78	54	26	40	25
24 NC98-26192	34	13	33	23
25 NC99-13296	33	12	28	21
26 NC99-13308	37	16	19	14
27 VAN98W342	32	11	23	17
28 VA00W562	28	5	19	13
29 VA00W566	46	22	16	8
30 VA01W461	36	15	10	3
31 VA02W713	28	6	20	15
32 VA02W732	44	18	22	16
Mean:	40	27	30.2	
L.S.D. (0.05)	24.9	31.7	20.0	
CV%	38.8	59.5	45.6	



Two Mega-environments were identified in the Greenhouse Head Severity evaluations:

Mega-environment 1: AR (Bay), KY and VA.

Mega-environment 2: NC and MO.

The most resistant entry in Mega-environment 1 (Ernie) was ranked ninth in Mega-environment 2.

The most resistant entry in Mega-environment 2 (Entry 9) was ranked fourteenth in Mega-environment 1.

But both entries were within the most resistant groups in their respective Mega-environments.

The largest change in rankings between the two Mega-environments were for Entries 7, 10, 15 and 22

Greenhouse (Rachis Involvement)¹

Cultivar/ Designation	COL'BIA MO	RALEIGH NC
1 ERNIE	1.0	0.5
2 COKER 9835	0.9	0.0
3 AR 857-1-1	0.9	0.0
4 AR 857-1-2	0.9	0.3
5 AR 93019-2-1	0.9	0.0
6 AR93035-4-1	0.9	0.3
7 ARGE 97-1042-4-5	0.9	0.3
8 ARGE 97-1033-3-5	0.8	0.3
9 ARGE 97-1048-3-6	0.6	0.5
10 ARGE 97-1038-3-5	0.8	0.3
11 ARGE 97-1047-4-2	0.9	0.5
12 B980416	1.0	0.0
13 B980582	0.3	0.0
14 B011066	0.6	0.3
15 B011117	0.4	0.5
16 GA 931587E53	0.9	0.3
17 GA 931630E48	1.0	0.8
18 GA 94261E7	1.0	0.3
19 GA 931233A24	0.9	0.3
20 GA 941208E35	1.0	0.3
21 MV 5-46	1.0	0.3
22 MV 15-42	0.8	0.1
23 MV 27-78	0.8	0.3
24 NC98-26192	0.5	0.5
25 NC99-13296	0.5	0.3
26 NC99-13308	0.9	0.5
27 VAN98W342	0.9	0.5
28 VA00W562	0.8	0.3
29 VA00W566	0.9	0.3
30 VA01W461	0.6	0.3
31 VA02W713	1.0	0.3
32 VA02W732	0.8	0.3

Mean: 0.80 0.28

¹Records whether the rachis on an individual head had disease where 0=no rachis involvement and 1=rachis showed symptoms.

DETACHED LEAF EVALUATIONS
DUBLIN, IRELAND

Cultivar/ Designation	Incubation Period ¹ (days)	Latent Period ¹ (days)	RANK
1 ERNIE	2.0	5.8	32
2 COKER 9835	2.7	6.3	27
3 AR 857-1-1	2.6	8.3	4
4 AR 857-1-2	2.5	7.9	7
5 AR 93019-2-1	3.6	6.7	19
6 AR93035-4-1	3.4	6.4	25
7 ARGE 97-1042-4-5	2.6	5.9	30
8 ARGE 97-1033-3-5	3.3	6.5	22
9 ARGE 97-1048-3-6	2.3	6.4	24
10 ARGE 97-1038-3-5	2.6	7.2	14
11 ARGE 97-1047-4-2	2.8	6.8	18
12 B980416	3.2	7.5	13
13 B980582	2.7	6.5	23
14 B011066	2.2	6.0	29
15 B011117	2.5	9.6	1
16 GA 931587E53	3.4	6.4	26
17 GA 931630E48	2.7	6.1	28
18 GA 94261E7	3.2	7.8	10
19 GA 931233A24	2.9	7.9	8
20 GA 941208E35	3.1	7.1	15
21 MV 5-46	2.9	6.7	20
22 MV 15-42	2.7	8.0	6
23 MV 27-78	3.3	6.6	21
24 NC98-26192	2.8	5.8	31
25 NC99-13296	2.4	7.0	16
26 NC99-13308	2.3	8.6	2
27 VAN98W342	3.5	7.8	9
28 VA00W562	3.3	7.6	12
29 VA00W566	3.5	8.2	5
30 VA01W461	3.0	8.6	3
31 VA02W713	2.6	6.9	17
32 VA02W732	2.4	7.7	11
MDV11-52	4.0	9.4	.
VA01W476	1.7	8.7	.
VA00W562	3.6	8.1	.
McCORMICK	3.1	7.2	.
NC-NEUSE	3.3	7.3	.
PAT	3.5	7.1	.
ROANE	2.6	7.7	.
Mean	2.9	7.3	
LSD. (0.05)	0.8	1.7	
C.V. (%)	21.4	18.6	

¹ Based on inoculation with five isolates of *M. nivale* var. *majus* as described by Browne and Cooke (2004), European J. Plant. Path. (in press).

Heading Date (Julian Days*)

CULTIVAR/ DESIGNATION	BAY AR	COLUMBIA MO	KINSTON NC	B'BURG VA	URBANA IL	WOOSTER OH	LEX'TON KY	FUN'LEA ROM	SZEGED HUN	MEAN ALL LOC.
	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK
1 ERNIE	120 5	129 1	116 13	129 12	131 1	143 20	126 2	137 1	131 2	129 2
2 COKER 9835	124 29	133 27	119 22	134 31	137 31	147 31	130 29	140 5	134 20	133 29
3 AR 857-1-1	123 26	131 10	118 18	129 13	137 28	143 21	127 3	. .	134 21	132 25
4 AR 857-1-2	122 22	133 22	119 20	130 20	136 22	145 27	127 4	. .	134 22	132 26
5 AR 93019-2-1	123 28	133 28	121 24	133 29	137 29	148 32	128 15	143 10	136 32	134 30
6 AR93035-4-1	122 23	133 23	119 23	131 28	137 26	145 28	128 16	143 11	134 23	132 27
7 ARGE 97-1042-4-5	118 1	133 24	. .	128 4	132 3	141 1	128 17	144 14	132 8	130 9
8 ARGE 97-1033-3-5	121 19	133 26	119 21	130 24	135 19	143 22	128 18	145 16	132 9	132 23
9 ARGE 97-1048-3-6	121 13	131 12	118 19	130 25	137 27	142 14	127 5	145 17	134 24	132 22
10 ARGE 97-1038-3-5	121 20	131 7	115 8	128 7	134 10	142 15	128 19	140 6	131 3	130 8
11 ARGE 97-1047-4-2	121 14	131 13	112 1	124 1	133 6	142 16	125 1	143 12	131 4	129 1
12 B980416	122 24	132 18	118 16	129 17	135 18	143 23	128 20	137 2	132 10	131 16
13 B980582	120 8	131 8	117 14	129 14	133 7	146 30	129 25	146 23	132 11	131 20
14 B011066	126 31	133 29	123 26	135 32	137 30	144 26	129 26	145 18	134 25	134 31
15 B011117	126 32	133 30	121 25	133 30	142 32	143 24	130 30	151 27	135 31	135 32
16 GA 931587E53	120 9	132 15	114 5	129 18	133 8	142 17	127 6	139 3	133 14	130 6
17 GA 931630E48	119 4	130 3	. .	128 8	132 4	142 18	130 31	146 24	131 5	130 12
18 GA 94261E7	121 15	133 31	115 11	129 9	133 9	141 2	127 7	. .	133 15	131 14
19 GA 931233A24	124 30	132 19	117 15	130 22	135 16	145 29	129 27	147 25	134 26	133 28
20 GA 941208E35	123 27	129 2	. .	127 2	131 2	143 25	127 8	139 4	132 12	130 3
21 MV 5-46	121 16	130 4	113 2	129 10	134 13	141 3	127 9	140 7	133 16	130 5
22 MV 15-42	120 10	130 5	. .	130 23	135 17	141 4	128 21	145 19	133 17	131 18
23 MV 27-78	121 21	132 17	115 9	130 21	136 25	141 5	127 10	144 15	134 27	131 19
24 NC98-26192	120 6	132 16	114 6	129 15	134 11	141 6	127 11	140 8	131 6	130 4
25 NC99-13296	118 2	131 14	. .	128 5	134 14	141 7	131 32	140 9	130 1	130 7
26 NC99-13308	120 11	132 20	115 10	129 19	134 12	141 8	129 28	145 20	131 7	131 17
27 VAN98W342	120 12	132 21	113 3	129 16	135 20	141 9	127 12	145 21	133 18	131 15
28 VA00W562	121 17	133 32	115 12	130 26	135 21	142 19	128 22	150 26	132 13	132 24
29 VA00W566	121 18	130 6	. .	127 3	136 23	141 10	127 13	. .	134 28	130 13
30 VA01W461	122 25	133 25	118 17	130 27	136 24	141 11	128 23	. .	133 19	132 21
31 VA02W713	120 7	131 11	113 4	129 11	132 5	141 12	127 14	143 13	134 29	130 10
32 VA02W732	118 3	131 9	114 7	128 6	134 15	141 13	128 24	145 22	134 30	130 11
Mean:	121	132	116	130	135	143	128	143	133	131
L.S.D. (0.05)	3	3	2	2	2	.	2	.	.	2
CV%	1	2.8	1	1	1	.	1	.	.	1

*Days after December 31, 2001

Plant Height (in)

CULTIVAR/ DESIGNATION	COL'BIA	B'BURG	LEX'TON	SZEGED	FUN'LEA	MEAN
	MO	VA	KY	HUN	ROM	ALL LOC.
	RANK	RANK	RANK	RANK	RANK	RANK
1 ERNIE	37 12	31 8	34 10	30 7	21 7	31 8
2 COKER 9835	30 1	31 5	33 5	30 4	21 6	29 3
3 AR 857-1-1	44 32	42 30	39 32	35 19	.	38 31
4 AR 857-1-2	43 30	43 31	39 29	34 15	.	37 30
5 AR 93019-2-1	43 31	43 32	39 30	45 32	29 24	40 32
6 AR93035-4-1	38 21	33 20	35 19	36 20	27 22	34 20
7 ARGE 97-1042-4-5	41 26	32 14	35 18	40 25	26 19	35 22
8 ARGE 97-1033-3-5	43 29	36 27	38 26	42 31	25 18	37 28
9 ARGE 97-1048-3-6	42 28	35 24	37 24	42 30	29 26	37 29
10 ARGE 97-1038-3-5	39 23	33 21	37 22	41 28	27 23	35 24
11 ARGE 97-1047-4-2	39 24	37 28	38 27	40 26	26 20	36 26
12 B980416	36 9	33 17	34 11	35 16	24 15	32 14
13 B980582	38 22	35 25	36 21	38 23	26 21	35 21
14 B011066	42 27	33 22	35 14	40 24	29 25	36 25
15 B011117	38 17	31 7	35 16	38 22	.	33 16
16 GA 931587E53	37 13	32 15	35 12	32 9	22 9	32 11
17 GA 931630E48	38 20	32 16	38 28	37 21	24 14	34 19
18 GA 94261E7	37 14	33 18	35 13	32 10	21 8	32 12
19 GA 931233A24	38 18	34 23	37 23	33 11	.	33 17
20 GA 941208E35	35 4	29 2	32 1	31 8	22 10	30 4
21 MV 5-46	35 5	32 10	32 3	29 2	25 16	31 7
22 MV 15-42	37 15	31 9	38 25	33 12	24 11	33 15
23 MV 27-78	36 6	32 11	34 9	30 6	18 2	30 6
24 NC98-26192	38 16	30 4	32 2	34 13	20 4	31 9
25 NC99-13296	36 7	32 12	32 4	34 14	20 5	31 10
26 NC99-13308	39 25	38 29	39 31	40 27	25 17	36 27
27 VAN98W342	35 3	29 3	34 8	28 1	17 1	29 2
28 VA00W562	37 11	31 6	33 7	30 5	19 3	30 5
29 VA00W566	31 2	28 1	33 6	29 3	.	28 1
30 VA01W461	36 10	36 26	35 15	42 29	.	35 23
31 VA02W713	38 19	33 19	36 20	35 18	24 13	33 18
32 VA02W732	36 8	32 13	35 17	35 17	24 12	32 13

Mean: 38 33 35 35 24 33
 L.S.D. (0.05) 1 2 5 . 3 4
 CV% 1 4 7 . . 6

Leaf Diseases

CULTIVAR/ DESIGNATION	% STRIPE RUST F'VILLE ¹ AR	%SEPTORIA LEAF BLOTCH F'VILLE ¹ AR	% GREEN LEAVES KIBLER ² AR	POWDERY MILDEW KINSTON ³ NC SZESEG ⁴ HUN.	LEAF RUST SZESEG HUN.
1 ERNIE	19	50	14	4.0	mst
2 COKER 9835	31	50	26	3.0	0
3 AR 857-1-1	4	31	60	7.0	s30
4 AR 857-1-2	5	30	69	6.0	s20
5 AR 93019-2-1	1	15	83	4.0	ms30
6 AR93035-4-1	1	9	55	3.0	s60
7 ARGE 97-1042-4-5	0	19	50	.	0
8 ARGE 97-1033-3-5	0	10	60	4.5	mrt
9 ARGE 97-1048-3-6	5	21	78	7.0	0
10 ARGE 97-1038-3-5	0	28	76	4.5	mrt
11 ARGE 97-1047-4-2	0	35	83	4.0	0
12 B980416	8	35	56	2.5	mrt
13 B980582	4	23	44	3.5	0
14 B011066	13	19	45	4.0	0
15 B011117	2	35	65	7.0	0
16 GA 931587E53	0	87	21	1.5	0
17 GA 931630E48	0	35	60	.	0
18 GA 94261E7	8	36	41	2.0	0
19 GA 931233A24	0	28	78	3.0	0
20 GA 941208E35	0	61	78	.	0
21 MV 5-46	6	68	50	0.0	s50
22 MV 15-42	5	88	40	.	0
23 MV 27-78	0	82	10	0.5	0
24 NC98-26192	0	38	81	4.0	0
25 NC99-13296	0	13	51	.	ms5
26 NC99-13308	0	19	80	3.0	0
27 VAN98W342	1	89	5	.	s100
28 VA00W562	0	89	7	1.5	0
29 VA00W566	0	60	18	.	mst
30 VA01W461	0	19	87	1.0	mst
31 VA02W713	11	55	60	0.0	s50
32 VA02W732	0	73	81	0.0	0
Mean:	4	42	53	3	.
LSD (0.05)	4	17	22	.	.

¹Percentage of flag leaf diseased. Rated 5-28. ²Septoria tritici blotch was principle disease together with some stripe rust.

³0=Resistant, 9=Fully Susceptible.

⁴For example 'ms30.5' indicated genotype was moderately susceptible with 30% of flag-1 covered with pustules.

1=lower leaves, 3=flag-2, 5=flag-1, 7=flag leaf, 9=head.

GRAIN YIELD

CULTIVAR/ DESIGNATION	GRAIN YIELD (bu/ac))		PLOT YIELD (kg/plot)		WINTER SURVIVAL (%) SZEGED HUN
	COLOMBIA MO	RANK	SZEGED HUN	RANK	
1 ERNIE	62.4	15	2.6	26	60 20
2 COKER 9835	42.5	32	2.4	29	30 30
3 AR 857-1-1	64.8	12	1.2	31	15 31
4 AR 857-1-2	55.0	25	1.1	32	15 32
5 AR 93019-2-1	58.3	20	2.9	20	95 1
6 AR93035-4-1	58.3	21	2.9	16	80 3
7 ARGE 97-1042-4-5	83.4	1	2.8	22	70 10
8 ARGE 97-1033-3-5	70.9	8	2.6	27	50 25
9 ARGE 97-1048-3-6	53.6	26	2.7	23	40 29
10 ARGE 97-1038-3-5	73.2	6	3.3	3	80 4
11 ARGE 97-1047-4-2	67.0	10	3.0	10	70 11
12 B980416	79.3	3	3.5	2	70 12
13 B980582	82.4	2	3.3	4	90 2
14 B011066	74.4	5	2.9	17	70 13
15 B011117	62.3	16	3.2	5	70 14
16 GA 931587E53	57.8	22	3.1	8	80 5
17 GA 931630E48	51.3	28	2.9	18	50 26
18 GA 94261E7	49.6	30	3.0	12	60 21
19 GA 931233A24	56.8	24	3.0	14	70 15
20 GA 941208E35	67.9	9	3.0	15	80 6
21 MV 5-46	64.3	13	3.0	11	50 27
22 MV 15-42	58.5	19	3.1	6	50 28
23 MV 27-78	50.0	29	2.9	21	70 16
24 NC98-26192	57.3	23	3.0	13	80 7
25 NC99-13296	72.5	7	2.9	19	60 22
26 NC99-13308	63.5	14	2.2	30	60 23
27 VAN98W342	61.3	18	2.6	28	70 17
28 VA00W562	52.4	27	2.7	24	70 18
29 VA00W566	47.1	31	3.0	9	60 24
30 VA01W461	61.4	17	2.7	25	75 9
31 VA02W713	78.2	4	3.7	1	80 8
32 VA02W732	65.8	11	3.1	7	70 19

Mean:

62.6

2.8

64

***Xgwm 493* and *XBARC 133* MARKER GENOTYPES**
(These two SSR loci flank the FHB resistance QTL on chromosome 3BS)

CULTIVAR/ DESIGNATION	<i>Xgwm 493</i>	<i>XBARC 133</i>
1 ERNIE	137	122
2 COKER 9835	137,160,188	120
3 AR 857-1-1	137,164	120
4 AR 857-1-2	137,156,188	114,120,125
5 AR 93019-2-1	137,156,188	120,125
6 AR93035-4-1	137,190	120
7 ARGE 97-1042-4-5	137,188	114,126
8 ARGE 97-1033-3-5	137,186	125
9 ARGE 97-1048-3-6	137,154,190	112,119
10 ARGE 97-1038-3-5	135,155,192	112,119
11 ARGE 97-1047-4-2	135,157,192	114,120
12 B980416	137,155,188	126
13 B980582	137,192	119
14 B011066	137,192	119
15 B011117	137	119
16 GA 931587E53	137,156,188	126
17 GA 931630E48	137,188	114
18 GA 94261E7	137,156,188	113,120
19 GA 931233A24	137,190	114,116
20 GA 941208E35	137,190	114,120
21 MV 5-46	137,194	113
22 MV 15-42	137,156,192	122
23 MV 27-78	135	NO DATA
24 NC98-26192	137,160	NO DATA
25 NC99-13296	135,192	113,119
26 NC99-13308	135,156	119,125
27 VAN98W342	137,156,190	113,119
28 VA00W562	NO DATA	120
29 VA00W566	137,190	119
30 VA01W461	137,156,188	120
31 VA02W713	137,192	119
32 VA02W732	137,194	114
Sumai 3	137,194	122
Ning 7840	137,194	122

Means Across Locations

CULTIVAR/ DESIGNATION	FHB Incidence (0-100)	FHB Severity (0-100)	FHB Index (0-100)	Scabby Seed %	Vomitoxin DON (ppm)	Heading Date (Julian)	Plant Height (in)	Greenhse Severity (0-100)
	RANK	RANK	RANK	RANK	RANK	RANK	RANK	RANK
1 ERNIE	49.6	5	23.5	13	13.9	9	27.6	8
2 COKER 9835	77.8	32	57.3	32	48.9	32	58.0	32
3 AR 857-1-1	58.2	11	16.9	3	11.0	5	32.1	14
4 AR 857-1-2	48.3	3	15.7	2	10.0	3	23.4	3
5 AR 93019-2-1	59.6	15	25.0	16	19.4	16	46.7	30
6 AR93035-4-1	65.7	21	26.7	18	23.3	21	49.6	31
7 ARGE 97-1042-4-5	44.4	2	19.2	9	10.2	4	30.2	10
8 ARGE 97-1033-3-5	44.3	1	10.7	1	4.5	1	25.2	4
9 ARGE 97-1048-3-6	51.5	7	17.7	7	14.6	11	27.0	6
10 ARGE 97-1038-3-5	54.7	10	20.4	11	12.2	7	27.3	7
11 ARGE 97-1047-4-2	50.7	6	17.0	5	9.8	2	31.9	12
12 B980416	52.1	9	21.1	12	13.9	10	32.9	16
13 B980582	49.1	4	18.7	8	12.3	8	16.6	1
14 B011066	59.4	14	19.2	10	16.2	13	33.7	20
15 B011117	51.8	8	17.2	6	11.7	6	40.2	23
16 GA 931587E53	75.1	30	44.8	27	39.4	29	45.4	27
17 GA 931630E48	66.4	22	36.6	23	29.7	24	31.5	11
18 GA 94261E7	77.1	31	47.6	31	43.1	31	44.9	26
19 GA 931233A24	71.2	28	46.3	29	37.2	28	45.6	29
20 GA 941208E35	70.9	27	46.8	30	40.5	30	41.2	25
21 MV 5-46	68.0	24	45.3	28	35.6	27	40.9	24
22 MV 15-42	67.2	23	41.1	25	31.6	25	26.4	5
23 MV 27-78	70.7	26	36.6	24	28.0	22	45.4	28
24 NC98-26192	61.4	18	26.4	17	16.0	12	30.1	9
25 NC99-13296	72.3	29	35.2	22	29.1	23	32.1	13
26 NC99-13308	60.7	17	24.5	15	17.5	15	23.1	2
27 VAN98W342	62.4	19	29.8	20	21.0	18	33.2	17
28 VA00W562	70.5	25	31.1	21	21.8	19	35.4	21
29 VA00W566	58.8	13	29.2	19	22.8	20	38.8	22
30 VA01W461	58.3	12	17.0	4	16.6	14	33.3	18
31 VA02W713	60.1	16	23.9	14	20.7	17	32.2	15
32 VA02W732	63.7	20	41.5	26	31.9	26	33.4	19

Mean:	61	29.1	22.7	34.8	11.3	131	33	35
L.S.D. (0.05)	10.9	10.2	11.7	15.2	10.8	2	4	20
CV%	16.7	37.4	52.5	38.1	76.9	1	6	46

Correlations Between Traits Over Locations.

	SEVERITY	INDEX	SCABBY SEED	VOMITOXIN DON	G'HOUSE TYPE 2	HEADING DATE	PLANT HEIGHT
INCIDENCE	0.88	0.91	0.67	0.82	0.48	ns	-0.59
SEVERITY		0.98	0.63	0.88	0.55	ns	-0.62
INDEX			0.68	0.90	0.54	ns	-0.57
SCABBY SEED				0.61	0.53	ns	-0.37
VOMITOXIN (DON)					0.44	ns	-0.54
G'HOUSE TYPE 2						ns	ns
HEADING DATE							ns