## **USDA-ARS/** U.S. Wheat and Barley Scab Initiative FY07 Final Performance Report (approx. May 07 – April 08) July 15, 2008

# **Cover Page**

| PI:                    | Robert Bowden  |  |
|------------------------|--|--|
| Institution:           | USDA-ARS   |  |
| Address:               | Plant Science and Entomology Research Unit (PSERU)           |  |
|                        | 4008Throckmorton Hall  |  |
|                        | Manhattan, KS 66506  |  |
|                        | USA  |  |
| E-mail:                | rbowden@ksu.edu  |  |
| Phone:                 | 785-532-2368   |  |
| Fax:                   | 785-532-6167   |  |
| Fiscal Year:           | 2007   |  |
| USDA-ARS Agreement ID: | NA   |  |
| USDA-ARS Agreement     | Test of a Novel Control Method against Fusarium Head Blight. |  |
| Title:                 |  |  |
| FY07 ARS Award Amount: | \$ 45,000  |  |

# **USWBSI Individual Project(s)**

| USWBSI            |  | ARS Adjusted |
|-------------------|--|--------------|
| Research          |  | Award        |
| Area <sup>*</sup> | Project Title  | Amount       |
| PGG               | Test of Sex Pheromones as a Novel Control Method against | \$45,000     |
|                   | Fusarium Head Blight.                                    | \$45,000     |
|                   | Total Award Amount                                       | \$ 45,000    |

| Principal Investigator | Date |
|------------------------|------|

HGG - Host Genetics & Genomics

IIR – Integrated/Interdisciplinary Research

PGG – Pathogen Genetics & Genomics

VDUN – Variety Development & Uniform Nurseries

<sup>\*</sup> CBCC – Chemical, Biological & Cultural Control

EEDF – Etiology, Epidemiology & Disease Forecasting

FSTU – Food Safety, Toxicology, & Utilization of Mycotoxin-contaminated Grain GET – Genetic Engineering & Transformation

HGR – Host Genetics Resources

FY07 (approx. May 07 – April 08)

PI: Bowden, Robert

USDA-ARS Agreement #: NA

**Project 1:** Test of Sex Pheromones as a Novel Control Method against Fusarium Head Blight.

## 1. What major problem or issue is being resolved and how are you resolving it?

The germination of conidia and ascospores of *Fusarium graminearum* (*Gibberella zeae*) is inhibited by fungal sex pheromones under laboratory conditions. This suggests that pheromone peptides might be able to prevent germination and penetration of plants by *F. graminearum* under natural conditions. For example, a transgenic plant expressing pheromone in the florets could exhibit a novel form of Type I resistance to initial infection. There are many hurdles to overcome, but the first priority is to test the feasibility of this strategy. This proposal aims to determine optimum conditions for spore inhibition by the pheromone and then conduct infection inhibition tests by applying pheromones onto wheat spikes.

2. List the most important accomplishment and its impact (how is it being used?). Complete all three sections (repeat sections for each major accomplishment):

## **Accomplishment:**

We found that germination of conidia in the laboratory was inhibited for 24 hours at concentrations of 40  $\mu$ mol of the ppg1 pheromone peptide from G. zeae. The ppg1 pheromone peptide from  $Neurospora\ crassa$  was more effective and inhibited germination for 24 hr at a concentration of 10  $\mu$ mol. Synthetic pheromone peptides were produced by substitution of amino acids from the pheromone peptide from Neurospora into the peptide from G. zeae. Some of the synthetic recombinant peptides were more effective than the natural peptides and were able to inhibit germination for 24 hr at 1  $\mu$ mol. In growth chamber pathogenicity tests, 100  $\mu$ mol of G. zeae pheromone peptide co-inoculated with conidia into single florets was ineffective at preventing infection.

#### **Impact:**

As a result of this work, we know that sex pheromones have important functions in the sexual cycle of *G. zeae*, the fungus that causes Fusarium head blight. Therefore, sex pheromones are a potential target for disease control. Initial attempts to inhibit conidial germination and prevent disease infection with sex pheromones were not successful. Further work is needed to define conditions affecting pheromone inhibition of germination of conidia.

# As a result of that accomplishment, what does your particular clientele, the scientific community, and agriculture as a whole have now that they didn't have before?:

The functions of sex pheromones in *G. zeae* have been determined. A variety of natural and synthetic fungal sex pheromone peptides have been shown to inhibit germination of conidia and ascospores in the laboratory.

FY07 (approx. May 07 – April 08)

PI: Bowden, Robert

USDA-ARS Agreement #: NA

Include below a list of the publications, presentations, peer-reviewed articles, and non-peer reviewed articles written about your work that resulted from all of the projects included in the grant. Please reference each item using an accepted journal format. If you need more space, continue the list on the next page.

#### REFEREED JOURNALS

- 1. Lee, J., J. E. Jurgenson, J. F. Leslie & R. L. Bowden. 2008. Alignment of genetic and physical maps of *Gibberella zeae*. Appl. Environ. Microbiol. 74:2349-2359.
- 2. Lee, J., J. F. Leslie & R. L. Bowden. 2008. Expression and function of sex pheromones and receptors in the homothallic ascomycete *Gibberella zeae*. Eukaryot. Cell 7:1211-1221.

### **ABSTRACTS**

- 1. Lee, J., R. L. Bowden & J. F. Leslie. 2007. Pheromone functions in *Gibberella zeae*. *Fungal Genetics Newsletter* 54(Suppl.):134.
- 2. Lee, J., J. F. Leslie & R. L. Bowden. 2007. Functions of the sex pheromones of Gibberella zeae. Proceedings of the 2007 National Fusarium Head Blight Forum (Kansas City, Missouri):30.