

**USDA-ARS/  
U.S. Wheat and Barley Scab Initiative  
FY08 Final Performance Report (approx. May 08 – April 09)  
July 15, 2009**

**Cover Page**

<b>PI:</b>	William Bockus
<b>Institution:</b>	Kansas State University
<b>Address:</b>	Department of Plant Pathology 4733 Throckmorton Hall Manhattan, KS 66506
<b>E-mail:</b>	bockus@ksu.edu
<b>Phone:</b>	785-532-1378
<b>Fax:</b>	785-532-5692
<b>Fiscal Year:</b>	2008
<b>USDA-ARS Agreement ID:</b>	59-0790-6-058
<b>USDA-ARS Agreement Title:</b>	Development of Scab Resistant Wheat Cultivars for Kansas.
<b>FY08 USDA-ARS Award Amount:</b>	\$ 26,157

**USWBSI Individual Project(s)**

<b>USWBSI Research Category*</b>	<b>Project Title</b>	<b>ARS Adjusted Award Amount</b>
HW-CP	Development of Scab Resistant Wheat Cultivars for Kansas.	\$26,157
	<b>Total Award Amount</b>	<b>\$ 26,157</b>

\_\_\_\_\_  
Principal Investigator

\_\_\_\_\_  
Date

\* MGMT – FHB Management

FSTU – Food Safety, Toxicology, & Utilization of Mycotoxin-contaminated Grain

GDER – Gene Discovery & Engineering Resistance

PBG – Pathogen Biology & Genetics

BAR-CP – Barley Coordinated Project

HW-CP – Hard Winter Wheat Coordinated Project

VDHR – Variety Development & Uniform Nurseries – Sub categories are below:

SPR – Spring Wheat Region

NWW – Northern Winter Wheat Region

SWW – Southern Winter Wheat Region

Michigan State University, 380 Plant & Soil Sciences Building, East Lansing, MI 48824-1325

Phone: (517) 355-0271 ext. 1183 ■ Fax: (517) 353-3955

Email: [scabusa@scabusa.org](mailto:scabusa@scabusa.org) ■ Website: <http://www.scabusa.org>

**U.S. Wheat & Barley Scab Initiative Networking & Facilitation Office**

**Project 1:** *Development of Scab Resistant Wheat Cultivars for Kansas.*

**1. What major problem or issue is being resolved relevant to Fusarium head blight (scab) and how are you resolving it?**

Serious scab (*Fusarium* head blight) epidemics occurred in Kansas in 1982, 1990, 1993, 1995, and 2008 and annual losses average \$2.1 million. Approximately one million acres of wheat in the eastern part of Kansas are annually at risk from scab where rainfall is higher during heading and corn residue is more prevalent. Since 1980, wheat acreage in the eastern one quarter of Kansas has declined by two thirds and a major cause of the decline has been farmer aversion to the risk of scab epidemics. Therefore, the availability of cultivars with resistance to scab is highly desirable for eastern Kansas. Additionally, there are three main reasons that scab has the potential to become much more prevalent in central Kansas where wheat is the traditional dominant crop (about 5 million additional wheat acres). First, there is a trend for increasing cultivation of corn in that part of the state, the main reservoir of inoculum of the scab pathogen. Second, there is a strong trend for decreasing tillage, which enhances spore production and release from the residue. Third, the prevalent varieties in central Kansas are susceptible to scab. Resistant wheat cultivars adapted to this area of Kansas would also be highly desirable. Genetic resistance offers the best hope for economic management of this disease. The long-term goal of this research is to develop hard red and hard white winter wheat cultivars adapted for Kansas with improved resistance to scab. Short term objectives are to: 1) test existing local cultivars for resistance, 2) test advanced breeding lines for resistance, 3) test exotic germplasm lines for resistance, 4) test the Hard Winter Wheat (Kansas, Nebraska, South Dakota) Scab Nursery for reaction to scab, and 5) incorporate new sources of scab resistance into the Kansas wheat breeding program. Testing will be done in misted field nurseries using soil-applied infested corn grain inoculum and in the greenhouse using single-floret inoculations. Visual disease evaluation methods will be used to rate the percentage spikelets infected by the pathogen and ground grain samples will be analyzed for the toxin DON. Data will be disseminated to wheat producers and used by wheat breeders as they make selections for future Kansas cultivars.

**2. List the most important accomplishment and its impact (i.e. how is it being used) to minimize the threat of Fusarium head blight or to reduce mycotoxins. Complete both sections (repeat sections for each major accomplishment):**

**Accomplishment:**

Until involvement in the USDA Scab Initiative, there was virtually no effort to identify sources of scab resistance in Kansas breeding programs. The Initiative has resulted in the development of accurate and efficient greenhouse and field testing nurseries that are providing useful ratings for current cultivars in Kansas, advanced breeding lines, and allow participation in the regional scab nurseries. Respectively, these nurseries allow dissemination of information to growers on the reaction of current commercial cultivars, selection by breeders for scab resistance in their breeding lines, and identification of additional sources of resistance from other breeding efforts in the region that can be incorporated into Kansas breeding lines. Kansas has also taken the lead in organizing a Hard Winter Wheat Scab Screening Nursery for the hard red winter wheat breeding programs of Kansas, Nebraska, and South Dakota. This latter nursery provides

PI: Bockus, William

USDA-ARS Agreement #: 59-0790-6-058

valuable data on the reaction of hard red winter wheat cultivars to scab in their area of adaptation. The long-term goal of the research is to develop, deploy, and advertise winter wheat cultivars adapted for Kansas with improved levels of resistance to scab.

**Impact:**

Two commercial cultivars in Kansas (Hondo and Heyne) were identified in 2000 (and confirmed in later years) as having good levels of scab resistance (3 and 4 on the 1-9 scale where 1=immune and 9=highly susceptible). During the past few years, these cultivars have averaged only 12 and 15% scab, respectively compared with about 50% in susceptible cultivars. Similarly, the cultivar Lakin has shown moderate levels of resistance with 22-34% scab. Six other commercial cultivars have also displayed moderate levels of resistance equal to, or better than, Lakin. Therefore, we have identified a few sources of scab resistance already present in cultivars adapted to Kansas that can be used by producers and may be potential sources of resistance for the development of future cultivars. Both KSU wheat breeders and the USDA wheat geneticist have been involved in the project by having their breeding lines evaluated for resistance to scab. Several breeding “populations” are tested each year from which the breeders make selections of promising lines showing resistance. Also, there are approximately 40 advanced breeding lines (The Kansas Intrastate Nursery) that are tested each year. Because of the scab testing efforts, a new column for reaction to “Head Scab” was added to the popular KSU extension publication *Wheat Variety Disease and Insect Ratings* for the fall, 2000 issue and has been updated in each subsequent year. For the first time, this has allowed producers in Kansas to use the reaction to scab to help select cultivars for planting. Similarly, data produced from nurseries funded by the Scab Initiative have recently been incorporated into another popular extension publication (*Kansas Performance Tests with Winter Wheat Varieties*). Both publications are available as “hard copy” or online. The involvement of breeders and the wheat geneticist has resulted in significant progress to improve the level of resistance to scab in future commercial wheat cultivars. This research has resulted in a germplasm release in 2004 from Kansas State University with resistance to scab derived from *Triticum timopheevii* ssp. *armenicum*. It normally takes 10-12 years to produce a new cultivar from the time initial crosses are made. Right on schedule, the first Kansas scab-resistant cultivar produced directly from the activity of the Initiative is due for release in Fall 2009, 10 years after beginning to receive funding from the Initiative.

**Include below a list of the publications, presentations, peer-reviewed articles, and non-peer reviewed articles written about your work that resulted from all of the projects included in the grant. Please reference each item using an accepted journal format. If you need more space, continue the list on the next page.**

1. Bockus, W. W., Baenziger, P. S., and Ibrahim, A. M. H. 2009. Reaction of Kansas, Nebraska, and South Dakota winter wheat accessions to Fusarium head blight (FHB), 2008. Plant Disease Management Reports (online). Report 3:CF006. DOI:10.1094/PDMR03 The American Phytopathological Society, St. Paul, MN.
2. Bockus, W. W., and Davis, M. A. 2009. Reaction of selected winter wheat accessions to Fusarium head blight, 2008. Plant Disease Management Reports (online). Report 3:CF007. DOI:10.1094/PDMR03. The American Phytopathological Society, St. Paul, MN.
3. Bockus, W. W. De Wolf, E., and Wegulo, S. N. 2009. Host resistance correlated with the amount of DON reduction with fungicide on winter wheat, 2008. Plant Disease Management Reports (online). Report 3:CF008. DOI:10.1094/PDMR03. The American Phytopathological Society, St. Paul, MN.
4. Bockus, W. W., Fritz, A. K., and Martin, T. J. 2009. Reaction of the 2008 Kansas Intrastate Nursery to Fusarium head blight. Plant Disease Management Reports (online). Report 3:CF009. DOI:10.1094/PDMR03. The American Phytopathological Society, St. Paul, MN.
5. Wegulo, S. N., Hernandez Nopsa, J., and Bockus, B. 2009. Effects of ProSaro fungicide and cultivars on Fusarium head blight (FHB) of wheat, 2008. Plant Disease Management Reports (online). Report 3:CF028. DOI:10.1094/PDMR03. The American Phytopathological Society, St. Paul, MN.
6. Appel, J. A., De Wolf, E., Bockus, W. W., Bowden, R. L., and Todd, T. 2008. Preliminary 2008 Kansas Wheat Disease Loss Estimates. Kansas Cooperative Plant Disease Survey Report. ([http://www.ksda.gov/includes/document\\_center/plant\\_protection/](http://www.ksda.gov/includes/document_center/plant_protection/))
7. Wegulo, S. N., Nopsa, J. H., and Bockus, W. W. 2008. Effects of fungicide treatments and cultivars on FHB and DON in winter wheat. Proceedings of the Annual Fusarium Head Blight Forum p. 70.
8. Bockus, W. W., Davis, M. A., De Wolf, E., and Wegulo, S. N. 2008. Host resistance correlated with the amount of DON reduction achieved with fungicides. Proceedings of the Annual Fusarium Head Blight Forum p. 9.
9. Bockus, W. W. 2009. Determining and reporting the reaction of Kansas commercial wheat varieties to wheat diseases. Poster at the meeting of the Kansas Wheat Conference, Hays.

**If your FY08 USDA-ARS Grant contained a VDHR-related project, include below a list all germplasm or cultivars released with full or partial support of the USWBSI. List the release notice or publication. Briefly describe the level of FHB resistance. If this is not applicable (i.e. no VDHR-related project) to your FY08 grant, please insert 'Not Applicable' below.**

There were no germplasm releases issued during the FY08 award.