

**USDA-ARS/
U.S. Wheat and Barley Scab Initiative
FY12 Final Performance Report
July 16, 2013**

Cover Page

PI:	Laura Sweets
Institution:	University of Missouri
Address:	Department of Plant Microbiology and Pathology 210 Waters Hall Columbia, MO 65211
E-mail:	SweetsL@missouri.edu
Phone:	573-884-7307
Fax:	573-882-1467
Fiscal Year:	FY12
USDA-ARS Agreement ID:	59-0206-9-078
USDA-ARS Agreement Title:	Managing FHB Through Integrated Practices and within-field Inoculum Sources.
FY12 USDA-ARS Award Amount:	\$ 13,743*

USWBSI Individual Project(s)

USWBSI Research Category**	Project Title	ARS Award Amount
MGMT	Evaluation of Integrated Management Strategies for Fusarium Head Blight.	\$ 11,696
MGMT	Effects of Local Corn Debris Management on FHB and DON Levels (Year Two).	\$ 2,047
	Total ARS Award Amount	\$ 13,743

Principal Investigator

Date

* Partial funding for this research is under ARS agreement # 59-0206-9-077

** MGMT – FHB Management

FSTU – Food Safety, Toxicology, & Utilization of Mycotoxin-contaminated Grain

GDER – Gene Discovery & Engineering Resistance

PBG – Pathogen Biology & Genetics

BAR-CP – Barley Coordinated Project

DUR-CP – Durum Coordinated Project

HWW-CP – Hard Winter Wheat Coordinated Project

VDHR – Variety Development & Uniform Nurseries – Sub categories are below:

 SPR – Spring Wheat Region

 NWW – Northern Soft Winter Wheat Region

 SWW – Southern Soft Red Winter Wheat Region

Project 1: *Evaluation of Integrated Management Strategies for Fusarium Head Blight.*

1. What major problem or issue is being resolved relevant to Fusarium head blight (scab) and how are you resolving it?

This research project addresses the need of identifying the best management methods for FHB/DON or good farming practices for FHB/DON management through integrated management studies. Needs addressed also include the evaluation of potential disease reductions through combinations of host resistance and fungicides, the documentation of the impact of crop sequence on disease risk and its potential role as part of the integrated management of FHB/DON and the development of outreach materials and opportunities for exchange of information with clientele.

The proposed research project very clearly mirrors Goal #1 of the FHB Management Action Plan, i.e. “Validate integrated management strategies for FHB and DON”. It also contributes to the goals of developing the next generation of management tools for FHB/DON control and enhancing communication and end user education/outreach by providing valuable research results on best management practices to clientele.

2011-2012 was the sixth year for the Integrated Management of Fusarium Head Blight and Deoxynivalenol study in Missouri. The recommended treatments were applied to five soft red winter wheat varieties. Ratings were made at the appropriate times and data for the six years of the study has been analyzed and summarized.

2. List the most important accomplishment and its impact (i.e. how is it being used) to minimize the threat of Fusarium head blight or to reduce mycotoxins. Complete both sections (repeat sections for each major accomplishment):

Accomplishment:

Weather conditions were conducive for the development of FHB in all five varieties in both crop sequence trials during the 2012 season. In the corn residue trial all main and interaction effects were statistically significant for both yield and DON. Yields in the corn residue were lower than in the soybean residue and DON levels were higher in the corn residue than in the soybean residue. These results confirm the importance of crop rotation and variety selection as management tools for both FHB and DON. Fungicide application improved yield and reduced DON levels with a greater impact on moderately resistant varieties.

Impact:

This research has shown the importance of crop sequence or residue type on the level of FHB and DON in the subsequent wheat crop. Crop rotation as a management tool for both FHB and DON management needs to be stressed. In general, the greatest reductions in FHB intensity and DON accumulation were observed when moderately resistant varieties were used with crop rotation and fungicide application. Resistant varieties had lower levels of

both FHB and DON in both residue types and with or without fungicide application. Under high disease pressure, a three tier management approach of crop rotation with a non-host, moderately resistant to resistant varieties and fungicide application was required to achieve < 2ppm DON.

This research project has provided data that can be used by producers to manage both FHB and DON levels in wheat. The importance of crop rotation, variety selection and the use of fungicide applications as the wheat flowers if weather conditions are conducive to the development of FHB are viable options for producers trying to manage FHB and reduce DON levels.

Project 2: *Effects of Local Corn Debris Management on FHB and DON Levels (Year Two).*

1. What major problem or issue is being resolved relevant to Fusarium head blight (scab) and how are you resolving it?

Reduction or elimination of within-field sources of inoculum of *Fusarium graminearum* is the basis for cultural control measures such as crop rotation sequences in which cereals follow non-cereal crops. The goal of this USWBSI research project is to provide realistic estimates of ‘DON reduction’ that can be expected from cultural controls that reduce within-field inoculum sources. We utilized moldboard plowing of corn debris as a proxy for planting after a non-cereal crop to compare directly with wheat planted no-till into corn debris in commercial-scale wheat fields planted following grain corn harvest in Illinois, Kentucky, Michigan, Missouri, Nebraska, New York, and Vermont. Following corn harvest in 2010, replicated wide (60 ft) strips were moldboard plowed or left non-plowed prior to sowing wheat over the entire field with a no-till drill. Wheat in each strip was monitored for FHB and sampled for laboratory quantification of head infection by *F. graminearum* and contamination of grain by DON in 2011. FHB symptoms at soft dough stage were low to moderate at every location except Missouri. Yet, at crop maturity, a high percentage of wheat heads was found to be infected by *F. graminearum* in all locations except Nebraska and Vermont. Measurable DON was found in grain from every environment and the levels were lowest in Vermont and highest in Kentucky and Nebraska. It is interesting that the Nebraska site showed the lowest disease index and lowest incidence of head infection, but the highest average toxin level. Moldboard plowing resulted in a significant decrease in FHB index in four environments (IL, MO, NY, MI), though the magnitude of the difference was large only in Missouri. In Nebraska, FHB index was significantly higher in the moldboard-plowed treatment in which the wheat crop matured earlier than in the no-till corn debris treatment. Moldboard plowing was associated with a small but significant decrease in recovery of *F. graminearum* from mature heads in three environments (IL, MI, NY). There was no significant effect of plowing on DON level in five environments (IL, KY, MO, NY, VT) and there were small, but significant decreases in toxin in moldboard-plowed compared to no-till strips in two environments (MI and NE). An additional treatment of minimum tillage (chisel plow) was added in the Michigan experiment; DON levels in the minimum-till plots were intermediate between moldboard and no-till but not significantly different from no-till. There is a strong trend in year one data suggesting that inoculum from area atmospheric sources exerts a far greater effect than inoculum from in-field corn residue on the level of DON contamination.

2. List the most important accomplishment and its impact (i.e. how is it being used) to minimize the threat of Fusarium head blight or to reduce mycotoxins. Complete both sections (repeat sections for each major accomplishment):

Accomplishment:

There is a strong trend in two years’ data suggesting that inoculum from area atmospheric sources exerts a far greater effect than inoculum from in-field corn residue on the level of DON contamination in agricultural-scale strip plots, yet there were small, but significant

decreases in toxin in moldboard-plowed compared to no-till strips in two environments (MI and NE).

Impact:

Based on year one data, it appears that localized corn debris management prior to wheat planting results, on average, in fairly small reductions in FHB and in DON contamination in wheat. Regional atmospheric inoculum appears to play a far greater role in corn-dominated landscapes, thus prioritizing the need for resistant wheat varieties and effective fungicides as the main pillars of integrated management in corn production regions. A second year of experimentation in seven additional wheat environments in 2012 provided increased evidence of the magnitude of the effect of localized corn residue management on DON reduction.

Include below a list of the publications, presentations, peer-reviewed articles, and non-peer reviewed articles written about your work that resulted from all of the projects included in the grant. Please reference each item using an accepted journal format. If you need more space, continue the list on the next page.

Refereed Publication:

Willyerd, K.T., C. Li, L.V. Madden, C.A. Bradley, G.C. Bergstrom, L.E. Sweets, M. McMullen, J.K. Ransom, A. Grybauskas, L. Osbourne, S.N. Wegulo, D.E. Hershman, K. Wise, W.W. Bockus, D. Groth, R. Dill-Mackey, E. Milus, P.D. Esker, K.D. Waxman, E.A. Adee, S.E. Ebelhar, B.G. Young and P.A. Paul. 2012. Efficacy and stability of integrating fungicide and cultivar resistance to manage Fusarium head blight and deoxynivalenol in wheat. *Plant Disease*, Vol. 96 No. 7, pp. 957-967 (5% contribution) (published).

Proceedings:

Bergstrom, G.C., J.A. Cummings, K.D. Waxman, C.A. Bradley, A.L. Hazelrigg, D.E. Hershman, M.Nagelkirk, L.E. Sweets and S.N. Wegulo. 2012. Effects of local corn debris management on FHB and DON levels in fourteen U.S. wheat environments in 2011 and 2012. *Proceedings of the 2012 National Fusarium Head Blight Forum, Orlando, FL, December 4-6, 2012. P. 5-6. Abstract in proceedings and poster at forum. (10% contribution) (published).*

Sweets, Laura E. 2012. Integrated management strategies for Fusarium head blight of soft red winter wheat in Missouri: summarization of six years of trial data. *Proceedings of the 2012 National Fusarium Head Blight Forum, Orlando, FL, December 4-6, 2012. P.27-28. Abstract in proceedings and poster at forum. (100% contribution) (published).*

Willyerd, K., K. Ames, G. Bergstrom, C. Bradley, J. Cummings, P. Gross, L. Madden, J. Ransom, K. Ruden, J.D. Salgado, L. Sweets, K. Wise, and P. Paul. 2012. Uniform FHB integrated management trials: a summary from 2012. *Proceedings of the 2012 National Fusarium Head Blight Forum, Orlando, FL, December 4-6, 2012. P.29-33. Abstract in proceedings and poster at forum. (10% contribution) (published).*

Extension Manuals:

Bradley, Kevin, Laura E. Sweets, Wayne C. Bailey, and J. Allen Wrather. 2012. 2013 Missouri pest management guide: corn, grain sorghum, soybean, winter wheat, rice and cotton. University of Missouri Extension publication M171. 200 pp.