

Efficacy of Miravis Ace® for FHB and DON Management across Environments and Grain Market Classes: A progress report

J.D. Salgado¹, G. Bergstrom², C. Bradley^{3,4}, K. Bowen⁵, E. Byamukama⁶, A. Byrne¹⁰, A. Collins⁷, C. Cowger⁸, J. Cummings², V. Chapara⁹, M.I. Chilvers¹⁰, E. De Wolf²¹, R. Dill-Macky¹², H.M. Darby¹³, P.D. Esker⁷, A. Friskop⁹, J. Halvorson⁹, N. Kleczewski^{3,14}, L.V. Madden¹, J. Marshall¹⁵, H. Mehl¹⁶, M. Nagelkirk^{10,11}, J. Starr², J. Stevens¹⁷, D. Smith¹⁸, M. Smith¹², S. Wegulo¹⁷, K. Wise^{4,19}, D. Yabwalo⁶, H.M. Young-Kelly²⁰, and P.A. Paul^{1*}

¹The Ohio State University/OARDC, Wooster 44691; ²Cornell University, Ithaca, NY 14853; ³University of Illinois, Urbana, IL 61801; ⁴University of Kentucky, Lexington, KY 40506; ⁵Auburn University, Auburn, AL 36849; ⁶South Dakota State University, Brookings, SD 57007; ⁷The Pennsylvania State University, University Park, PA 16802; ⁸North Carolina State University/USDA-ARS, Raleigh, NC 27695; ⁹North Dakota State University, Fargo, ND 58102; ¹⁰Michigan State University Extension, Sandusky, MI 48471; ¹¹Michigan State University, East Lansing, MI 48824; ¹²University of Minnesota, St. Paul, MN 55108; ¹³The University of Vermont, Burlington, VT 05405; ¹⁴The University of Delaware, Newark, DE 19719; ¹⁵University of Idaho, Idaho Falls, ID 83402; ¹⁶Virginia Tech, Suffolk VA 23437; ¹⁷University of Nebraska-Lincoln, NE 68588; ¹⁸University of Wisconsin, Madison, WI 53706; ¹⁹Purdue University, West Lafayette, IN 47907; ²⁰The University of Tennessee at Knoxville, Jackson, TN 38301. ²¹Kansas State University, Manhattan, KS 66506.

*Corresponding Author: PH: 330-263-3842; Email: paul.661@osu.edu

OBJECTIVES

1. Evaluate the integrated effects of fungicide treatment and genetic resistance on FHB and DON in all major grain classes, with emphasis on a new fungicide, Miravis Ace.
2. Compare the efficacy of Miravis Ace when applied at heading or at anthesis to that of a standard anthesis application of Prosaro or Caramba.

INTRODUCTION

Results from previous uniform fungicide trials (UFTs) and management coordinated projects (IM_CP) showed that Demethylation Inhibitors (DMI) fungicides such as Prosaro® and Caramba® are the most effective against FHB and DON. Applications made at or up to 6 days after anthesis to moderately resistant cultivars provide more than 70% reduction of both FHB index and DON, relative to a non-treated, susceptible check (2,4,7). Results from UFTs also showed that Quinone Outside Inhibitor (QoI) fungicides consistently led to an increase in DON accumulation in harvested grain, particularly when applied close to anthesis (3). Based on these findings, DMIs are the most widely used fungicides for FHB and DON management, but this is not a good fungicide resistance management strategy. Isolates of *Fusarium graminearum* with resistance to tebuconazole have been reported (1,5), and a recent quantitative synthesis of 20 years of data from UFTs (Madden et al. *personal communication*) suggests that the efficacy of this and other DMIs may be declining in some areas. Preliminary results from a few pilot studies showed that Miravis Ace, a new Succinate Dehydrogenase Inhibitor + propiconazole fungicide, provides comparable levels of FHB and DON reduction to that of Prosaro and Caramba when applied at anthesis (Feekes 10.5.1). Moreover, this new fungicide is reported to be just as effective as the industry standards when applied at Feekes 10.3.

MATERIALS AND METHODS

Two sets of field experiments were conducted in 11 US wheat-growing states in 2018 to evaluate the performance of Miravis Ace. **For objective #1 (IM_CP)**, fungicide treatments (sub-plot) consisting of a non-treated check (**CK**), Prosaro at 6.5 fl. oz./A applied at 50% early anthesis (**I**), and Miravis Ace at 13.7 fl. oz./A applied at 50% early anthesis (**II**) or at 50% heading (Feekes 10.3, **III**) were applied to replicate plots (3-6 blocks) of susceptible (S), moderately susceptible (MS), and moderately resistant

(MR) cultivars (whole-plot). **For objective #2 (UFT)**, plots of a susceptible cultivar were planted in 3-6 replicate blocks and subjected to various combinations of the following fungicide programs: **1)** a non-treated check; **2)** Miravis Ace at 11.5 fl. oz./A (low rate) at Feekes 10.3-10.5; **3)** Miravis Ace at 13.7 fl. oz./A (high rate) at Feekes 10.3-10.5; **4)** Miravis Ace at 11.5 fl. oz./A at anthesis; **5)** Miravis Ace at 13.7 fl. oz./A at anthesis; **6)** Prosaro at 6.5 fl. oz./A at anthesis; and **7)** Caramba at 13.5 fl. oz./A at anthesis. In addition, we evaluated two-treatment programs consisting of an application of Miravis Ace (at low or high rate) at anthesis followed by an application of Prosaro at 6.5 fl. oz./A (**8**), Caramba at 13.5 fl. oz./A (**9**), or Folicur at 4 fl. oz./A (**10**) at 3-5 days post anthesis. Separate post-anthesis (3-5 days) applications of Prosaro at 6.5 fl. oz./A (**11**) and Miravis Ace at 13.7 fl. oz./A (**12**), and a pre-anthesis (Feekes 10.3-10.5) application of Prosaro (**13**) were also included in some trials.

All plots were artificially inoculated with either *F. graminearum*-colonized grain spawn or a spore suspension of the fungus sprayed approximately 24-36 hours after the anthesis treatments were applied. Some locations were naturally infected. FHB index (IND) was rated or calculated as previously described (6) on 60-100 spikes per plot at approximately Feekes 11.2. Grain samples were sent to a USWBSI-supported laboratory for mycotoxin analysis. Linear mixed models (multi-location) were fitted to the pooled arcsine square root-transformed IND and log-transformed DON data to evaluate management program and treatment effects.

RESULTS AND DISCUSSION

DON data were not available for some environments at the time of this report. For **objective 1**, trial-level mean IND and DON in the non-treated susceptible check (S_CK) ranged from 2.3 to 46.4 % and 0.5 to 8.6 ppm, respectively. All cultivar x fungicide treatment combinations had significantly ($p < 0.05$) lower mean IND (**Fig. 1A**) and DON (**Fig. 1B**) than S_CK. When applied at 13.7 fl. oz./A at Feekes 10.5.1 (**II**) or at Feekes 10.3-10.5 (**III**), Miravis Ace was not significantly different from Prosaro at 6.5 fl oz/A applied at anthesis (**Fig. 1A and B**).

For **objective 2**, trial-level mean IND and DON in the non-treated check (CK) ranged from 0.22 to 42.1 % and 1.3 to 38.7 ppm, respectively. All fungicide programs that included an anthesis application resulted in significantly ($p < 0.05$) lower mean IND and DON than the check (**Fig. 2**). For IND, means were not significantly different among Prosaro, Caramba, and Miravis Ace, and applications made at Feekes 10.3-10.5 were not significantly different from those made at 50% Feekes 10.5.1, or 4 days after 50% early anthesis, in the case of Miravis Ace. Two-treatment programs consisting of Miravis Ace applied at 50% early anthesis followed by Prosaro, Caramba, or Folicur applied 4 days after resulted in the lowest overall mean IND, with the Miravis Ace + Folicur and Miravis Ace + Caramba combinations having significantly lower means than the anthesis-only Prosaro or Caramba treatments (**Fig. 2A**). Two-treatment programs also outperformed all other tested programs in terms of mean DON reduction, however, treatments applied at between Feekes 10.3 and 10.5 (MIR_H and PRO_H) were not significantly different from the non-treated check (**Fig. 2B**). In particular, the anthesis and post-anthesis applications of Miravis Ace (MIR_A and MIR_A+4) had significantly lower mean DON than MIR_H. Although based on a relatively small sample size (6-16 trials for the MGMT_CP and 2-12 for the UFT, depending on the treatment and the response), these results are very promising. When applied at anthesis, Miravis Ace was just as effective as Prosaro and Caramba in terms of FHB and DON reduction. However, when applied between Feekes 10.3 and 10.5, the performance of this new fungicide was not consistent between the MGMT_CP and the UFT in terms of efficacy against DON. A second year of data will be collected in 2019, and a more complete set of analyses will be performed.

ACKNOWLEDGEMENTS AND DISCLAIMER

This material is based upon work supported by the U.S. Department of Agriculture, under Agreement Nos. 59-0206-4-018, 59-0206-4-006, 59-0206-5-007, 59-0206-5-005, 59-0206-4-016, 59-0206-4-035, 59-0206-4-012, 59-0206-4-036, 59-0206-4-040, 59-0206-6-010, 59-0206-4-037, 59-0206-6-012, 59-0206-6-014, 59-0206-4-017 and 59-0206-6-009. This is a cooperative project with the U.S. Wheat & Barley Scab Initiative. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the authors and do not necessarily reflect the view of the U.S. Department of Agriculture.

REFERENCES

1. Klix, M. B., Verreet, J.-A., and Beyer, M. 2007. Comparison of the declining triazole sensitivity of *Gibberella zeae* and increased sensitivity achieved by advances in triazole fungicide development. *Crop Prot.* 26:683-690.
2. Paul, P. A., Bradley, C. A., Madden, L. V., Dalla Lana, F., Bergstrom, G. C., Dill-Macky, R., Wise, K. A., Esker, P., McMullen, M. P., Grybauskas, A., Kirk, W., Milus, E. A., and Ruden, K. 2018. Effects of pre- and post-anthesis applications of demethylation inhibitor fungicides on *Fusarium* head blight and deoxynivalenol in spring and winter wheat. *Plant Dis.* 102: <https://doi.org/10.1094/PDIS-03-18-0466-RE>.
3. Paul, P. A., Bradley, C. A., Madden, L. V., Dalla Lana, F., Bergstrom, G. C., Dill-Macky, R., Esker, P. D., Wise, K. A., McMullen, M., Grybauskas, A., Kirk, W. W., Milus, E., and Ruden, K. 2018. Meta-analysis of the effects of QoI and DMI fungicide combinations on *Fusarium* head blight and deoxynivalenol in wheat. *Plant Dis. in press* - <https://doi.org/10.1094/PDIS-02-18-0211-RE>
4. Salgado, J. D., Ames, K., Bergstrom, G., et al. 2017. A Multi-state coordinated project to evaluate integrated management strategies for *Fusarium* head blight and deoxynivalenol in wheat. *Phytopathology* 107: Abstract.
5. Spolti, P., Del Ponte, E. M., Dong, Y., Cummings, J. A., and Bergstrom, G. C. 2014. Triazole sensitivity in a contemporary population of *Fusarium graminearum* from New York wheat and competitiveness of a tebuconazole-resistant isolate. *Plant Dis.* 98:607-613.
6. Stack, R. W., and McMullen, M. P. 1998. A visual Scale to estimate severity of *Fusarium* head blight in wheat. NDSU Extension Service: Small Grains Publications. Online Publication/PP-1095.
7. Willyerd, K. T., Li, C., Madden, L. V., Bradley, C. A., Bergstrom, G. C., Sweets, L. E., McMullen, M., Ransom, J. K., Grybauskas, A., Osborne, L., Wegulo, S. N., Hershman, D. E., Wise, K., Bockus, W. W., Groth, D., Dill-Macky, R., Milus, E., Esker, P. D., Waxman, K. D., Adey, E. A., Ebelhar, S. E., Young, B. G., and Paul, P. A. 2012. Efficacy and stability of integrating fungicide and cultivar resistance to manage *Fusarium* head blight and deoxynivalenol in wheat. *Plant Dis.* 96:957-967.