

NOAA Technical Memorandum NWS WR-139

AIDS FOR FORECASTING MINIMUM TEMPERATURE IN THE
WENATCHEE FROST DISTRICT

Robert S. Robinson

National Weather Service Office
Wenatchee, Washington
April 1979

UNITED STATES
DEPARTMENT OF COMMERCE
Juanita M. Kreps,
Secretary

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION
Richard Frank,
Administrator

NATIONAL WEATHER
SERVICE
Richard E. Hallgren,
Director



This Technical Memorandum has been
reviewed and is approved for
publication by Scientific Services
Division, Western Region.

A handwritten signature in black ink, appearing to read "L. W. Snellman". The signature is written in a cursive style with a long, sweeping tail that extends to the right.

L. W. Snellman, Chief
Scientific Services Division
Western Region Headquarters
Salt Lake City, Utah

CONTENTS

	<u>Page</u>
Tables and Figures	iv
I. Introduction	1
II. Procedure	1
III. Results	2
IV. Summary	3
V. References	3

TABLES AND FIGURES

	<u>Page</u>
Table 1. Summary of Regression Estimate of Minimum Temperature for 105 Nights, 1964 - 1975	4
Table 2. Verification on Three Years of Data, 79 Nights, 1976-77-78	4
Table 3. Summary of Regression Estimate of Minimum Temperatures for 99 Nights, 1972 - 1976	5
Table 4. Verification on 1977 and 78 Seasons, 63 Nights	5
Table 5. LFM 1000- to 500-mb Thickness Forecasts, 12Z 24-Hour Prog, 1976-77.	5
Figure 1. Lowest Minimum Temperature Forecast Nomogram for the District on the Next Morning using 1600 PST Dew Point at Omak or Wenatchee and the 1600 PST Interpolated 1000-500 mb Thickness	6
Figure 2. Lowest Minimum Temperature Forecast Nomogram for the District on the Next Morning Using Cos (2 x day of year) and 0400 PST 1000-500 mb Thickness at Quillayute	7

AIDS FOR FORECASTING MINIMUM TEMPERATURE IN THE WENATCHEE FROST DISTRICT

Robert S. Robinson
Weather Service Office
Wenatchee, Washington

I. INTRODUCTION

As in other fruit-frost districts, the Hygrometric Formula (Ellison, 1928) has been used in Wenatchee as an aid to forecasting minimum temperatures during the spring. However, this formula was developed for the Wenatchee key station and there is a wide variation in minimum temperatures across the district, averaging about 12°F between the warmest and coldest key stations during April and May. More importantly, due to the elevation differences and rugged terrain, there are no consistent relationships between frost key stations. While National Meteorological Center (NMC) guidance in the form of MOS minimum temperature forecasts is available for Yakima, experience has shown that it is not reliable enough for use as an aid in the fruit-frost forecasting in north-central Washington. Because of these factors, a semiobjective may-typing system has been used which is usually very helpful in determining minimum temperature distribution across the district and somewhat useful in obtaining absolute values. To complement this scheme, an objective aid was sought which would estimate the coldest minimum temperature to be expected among the key stations in the district on a given night.

II. PROCEDURE

As in other minimum-temperature studies, it was decided in the beginning to eliminate nights in which cloudiness was a factor. A review of past records indicated that cirrus of any amount, classified as thin, did not have any significant effect on minimum temperatures. Also, any cloud coverage of one-half or less was usually not enough to prevent at least one or two key stations in the district from achieving good nighttime radiation conditions. An examination of nights during five past seasons showed that 41 percent of all nights met this high-radiation criterion during April and May. Of these nights, 84 percent were cold (32°F or less at any key station) during April and 69 percent during May. Therefore, efforts were directed toward developing an aid which would indicate the coldest temperature in the district on nights with clear to scattered clouds or thin cirrus only. The sky-cover observations for 06Z, 09Z, and 12Z at Pangborn Field (EAT) near Wenatchee were used to measure the degree of cloudiness over the district.

Two types of forecasts are issued during the frost season for the Wenatchee district. Individual key station forecasts are made in the evening. However, a preliminary estimate of the lowest key station temperature expected in the district is issued in the morning for the following morning. Two aids were developed to utilize information available at

these times of day. A number of variables which have been used in other minimum temperature forecast schemes (Dickey) and (Oard, 1977) were examined in developing the aid for use in the evening. These included 700- and 850-mb temperatures, maximum temperature, surface temperature at 00Z, relative humidity, 1000- to 500-mb thickness, 700- to 1000-mb thickness, dew point at 00Z, and $\cos N X$ day of the year (where N was any of a few tested integers*). The stepwise multiple regression technique used by Oard would have been the ideal method to handle the correlation of these variables and arrive at the most efficient regression equation. However, since the necessary computer facility was not available, selection of variables was accomplished by manual computation of linear correlation coefficients on a small sample. While some of these variables showed good correlation with minimum temperature, most were eliminated because they were also highly correlated with either 1000- to 500-mb thickness or dew point. The independent variables finally chosen were a locally derived 1000- to 500-mb thickness and surface dew point, both taken at 00Z. The thickness was computed from the average of the 00Z surface pressures at EAT and Omak (OMK) and an average 500-mb height over the district. The dew point used was the lowest of Wenatchee WSO and OMK at 00Z. The $\cos 2 X$ day of the year was added to try to account for the shortening length of night during the season. The improvement in multiple correlation coefficient between variables and minimum temperature, reduction of variance, and standard error of estimate due to the addition of successive variables chosen is shown in Table 1. These were computed for 105 nights, 1964-1975, using a WANG programmable calculator on loan from Seattle WSFO. Verification on 79 nights during 1976, 1977, and 1978 (Table 2) showed no significant improvement due to the addition of $\cos 2 X$ day of the year (nor did any of the other values for N). Therefore, the final regression equation was formed using only the thickness and dew point as independent variables.

Selection of variables for an aid for the morning preliminary forecast followed the same procedure as outlined above. Variables tested were mainly from 12Z data in April and May 1972-1976. The final variables chosen were the 1000- to 500-mb thickness at Quillayute (UIL) at 12Z and $\cos 2 X$ day of the year (Table 3). Verification of the regression equation developed from these variables on the independent sample (1977-78) is shown in Table 4.

III. RESULTS

Test results in Tables 2 and 4 show that both aids can provide useful forecasts of minimum temperature providing the amount of cloudiness can be properly assessed. Graphical solutions to the regression equations were prepared (Figures 1 and 2) for operational use. The utility of the morning forecast aid has been increased by recognizing the ability of the LFM model to forecast with reasonable accuracy the 1000- to 500-mb

*Editor's Note: Normally $\cos N X$ day of year, where $N=1$, is used in such studies. This study, however, is only over a two-month period (April and May), not the full 360-day period of the complete cycle of $\cos 1 X$ day of year, so other N values were tested as well.

thickness in the vicinity of UIL. Based on experience with the facsimile output of the LFM thickness prog on the 12Z run, the accuracy is less than useful beyond 24 hours. As indicated in Table 5, however, the thickness forecast by the LFM at 24 hours is very good. The average error in the thickness forecast over UIL during the 1976 and 77 seasons was less than 25 meters during April and May. When this is applied to the morning forecast aid, it results in a variation of less than one degree F. This should allow a forecaster to make a more accurate assessment of the potential district minimum temperature about 48 hours in advance.

Several shortcomings have also been identified. Strong warm advection causes both equations to forecast temperatures too high. When an increase in the 1000- to 500-mb thickness occurs at this time of year, it more often acts as a stabilizing factor, capping cooler air at low levels. Strong cold advection on the other hand doesn't seem to introduce any significant bias. While wind is a factor governing minimum temperatures, the topographical makeup of the district is such that wind lulls will occur during the early morning allowing temperatures to fall to near values forecast by the aids at some station within the district. One exception to this occurs when the amount of cloudiness is just within the specified limits on windy nights. Verification of equation 1 indicated that on the average the mean 1000- to 500-mb thickness at 00Z between UIL and Spokane (GEG) may be used in place of the locally derived value without greatly impairing accuracy. Sufficient data of this kind were not available for the development of the equation.

IV. SUMMARY

Regression equations and their graphical solution were developed as aids for forecasting the lowest minimum temperature among frost key stations in the Wenatchee frost district on a given night during the spring. Two aids were developed, one based on morning data for the preliminary frost forecast and one based on evening data for the primary forecast. Both require that nights have no more than scattered clouds or thin cirrus. Verification on independent data showed both equations to be quite useful. To further improve their usefulness, more study is needed to objectively determine whether or not sufficient cloudiness will be present to affect minimum temperatures.

V. REFERENCES

- Dickey, W. W.: An Aid for Forecasting the Minimum Temperature During Spring at Yakima, Washington. Unpublished local study, Seattle WSFO, Washington.
- Ellison, E. S., 1928: A Critique on the Use of Minimum-Temperature Formulas. Monthly Weather Review, Vol. 56, No. 12, pp 485-495.
- Oard, M. J., 1977: A Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. NOAA Technical Memorandum NWS WR 113, NOAA, U. S. Dept. Commerce, Washington, D. C.

TABLE 1
Equation 1

SUMMARY OF REGRESSION ESTIMATE OF MINIMUM TEMPERATURE FOR 105
NIGHTS, 1964 - 1975

VARIABLES USED	MULTIPLE CORRELATION COEFFICIENT	REDUCTION OF VARIANCE	STD. ERROR OF ESTIMATE
1000- to 500-mb Thickness	0.7802	0.6087	2.5672
1000- to 500-mb Thickness and 00Z Dew Point	0.8408	0.7070	2.3222
1000- to 500-mb Thickness 00Z Dew Point, and cos 2 X day of year	0.8631	0.7450	2.1336

TABLE 2
Equation 1

VERIFICATION ON THREE YEARS OF DATA, 79 NIGHTS, 1976-77-78

VARIABLES USED	MEAN ABSOLUTE ERROR	STD. ERROR OF ESTIMATE
1000- to 500-mb Thickness	2.0534	2.5715
1000- to 500-mb Thickness and 00Z Dew Point	1.7020	2.1250
1000- to 500-mb Thickness, 00Z Dew Point, and cos 2 X day of year	1.6959	2.2201

TABLE 3

Equation 2

SUMMARY OF REGRESSION ESTIMATE OF MINIMUM TEMPERATURES FOR 99
NIGHTS, 1972 - 1976

VARIABLES USED	MULTIPLE CORRELATION COEFFICIENT	REDUCTION OF VARIANCE	STD. ERROR OF ESTIMATE
UIL 1000- to 500-mb Thickness	0.6981	0.4874	3.1725
UIL 1000- to 500-mb Thickness, and cos 2 X day of year	0.7563	0.5719	2.9133

TABLE 4

Equation 2

VERIFICATION ON 1977 and 78 SEASONS, 63 NIGHTS

VARIABLES USED	MEAN ABSOLUTE ERROR	STD. ERROR OF ESTIMATE
UIL 1000- to 500-mb Thickness	2.3657	3.1027
UIL 1000- to 500-mb Thickness, and cos 2 X day of year	2.001	2.5947

TABLE 5

LFM 1000- to 500-MB THICKNESS FORECASTS, 12Z 24-HOUR
PROG - 1976-77 (Decameters)

VARIABLES USED	MEAN ABSOLUTE ERROR	STD. ERROR OF ESTIMATE
April	2.4400	3.1559
May	2.1452	2.8933
COMBINED	2.2768	3.0133




FIGURE 1. LOWEST MINIMUM TEMPERATURE FORECAST NOMOGRAM FOR THE DISTRICT ON THE NEXT MORNING USING 1600 PST DEW POINT AT OMAK OR WENATCHEE (WHICHEVER IS LOWER) AND THE 1600 PST INTERPOLATED 1000-500 MB THICKNESS.




FIGURE 2. LOWEST MINIMUM TEMPERATURE FORECAST NOMOGRAM FOR THE DISTRICT ON THE NEXT MORNING USING $\cos(2 \times \text{DAY OF THE YEAR})$ AND 0400 PST 1000-500 MB THICKNESS AT QUILLAYUTE.

NOAA Technical Memoranda NWSR: (Continued)

- 92 Snake Management in the Willamette Valley. Earl W. Bates, May 1974. (COM-74-11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM-74-11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM-74-11555/AS)
- 96 Map Type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM-75-10423/AS)
- 97 Eastern Pacific Cut-off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB-250-711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM-75-10719/AS)
- 99 A Study of Flash Flood Susceptibility--A Basin in Southern Arizona. Gerald Williams, August 1975. (COM-75-11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuff, October 1975. (PB-246-902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB-253-053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB-252-366/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB-254-650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John G. Plankinton, Jr., March 1976. (PB-254-649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB-259-594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB-260-437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, Sept. 1976. (PB-273-677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Danney, November 1976. (PB-264-653/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB-265-941/AS)
- 115 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Gard, February 1977. (PB-273-694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB-273-676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB-268-347)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-value. R. F. Quiring, April 1977. (PB-272-351)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB-268-740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Barry Randerson, June 1977. (PB-271-290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R. F. Quiring, June 1977. (PB-271-704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB-271-742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB-272-661)
- 123 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB-273-155/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Nielke, February 1978. (PB-281-337/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB-283-030/AS)
- 129 Fire Whirls. David W. Coens, May 1978. (PB-283-866/AS)
- 130 Flash-Flood Procedure. Ralph G. Hatch and Gerald Williams, May 1978. (PB-286-014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mallin and David E. Olsen, September 1978. (PB-289-916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-740 Weather Radar. R. G. Pappas, R. Y. Lee, and B. W. Finke, October 1978. (PB-289-767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB-291-317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB-291-195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB-292-716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB-292-247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979.
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Hefflick and James R. Fors, February 1979.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

NOAA, the *National Oceanic and Atmospheric Administration*, was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth, and to assess the socioeconomic impact of natural and technological changes in the environment.

The six Major Line Components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS — Important definitive research results, major techniques, and special investigations.

TECHNICAL REPORTS — Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS — Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

CONTRACT AND GRANT REPORTS — Reports prepared by contractors or grantees under NOAA sponsorship.

TECHNICAL SERVICE PUBLICATIONS — These are publications containing data, observations, instructions, etc. A partial listing: Data serials; Prediction and outlook periodicals; Technical manuals, training papers, planning reports, and information serials; and Miscellaneous technical publications.

ATLAS — Analysed data generally presented in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, atmospheric conditions, etc.



Information on availability of NOAA publications can be obtained from:

**ENVIRONMENTAL SCIENCE INFORMATION CENTER
ENVIRONMENTAL DATA SERVICE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE**

**3300 Whitehaven Street, N.W.
Washington, D.C. 20235**