

NOAA Technical Memorandum NWS WR-144

ARIZONA COOL SEASON CLIMATOLOGICAL SURFACE WIND
AND PRESSURE GRADIENT STUDY

Ira S. Brenner*
National Weather Service Forecast Office
Phoenix, Arizona

May 1979

*Present affiliation: National Hurricane Center
Miami, Florida

UNITED STATES
DEPARTMENT OF COMMERCE
Juanita M. Kreps,
Secretary

NATIONAL OCEANIC AND
ATMOSPHERIC ADMINISTRATION
Richard Frank,
Administrator

NATIONAL WEATHER
SERVICE
Richard E. Hallgren,
Director



This Technical Memorandum has been reviewed and is approved for publication by Scientific Services Division, Western Region.

A handwritten signature in black ink, appearing to read "L. W. Snellman". The signature is written in a cursive style with a long, sweeping tail that extends to the right.

L. W. Snellman, Chief
Scientific Services Division
Western Region Headquarters
Salt Lake City, Utah

CONTENTS

	<u>Page</u>
Figures	iv
Abstract	1
I. Introduction	1
II. Developmental Data Analysis	2
III. Conclusions	5
IV. Acknowledgments	5
V. References	5

FIGURES

	<u>Page</u>
Figure 1a.	Average Sea-Level Pressure Gradients that Produce Average Sustained Surface Winds of 3515G25 at Yuma, Arizona, for Six Consecutive Hours 6
Figure 1b	Same as Above for Winds of 3510-3515 at Yuma, Arizona 6
Figure 2a	Same as Above for Winds of 0815 at Phoenix, Arizona 7
Figure 2b	Same as Above for Winds of 0808-0813 at Phoenix, Arizona 7
Figure 3a	Same as Above for Winds of 0512G25 at Flagstaff, Arizona 8
Figure 3b	Same as Above for Winds of 0510-0515 at Flagstaff, Arizona 8
Figure 4a	Same as Above for Winds of 1215G25 at Tucson, Arizona 9
Figure 4b	Same as Above for Winds of 1210-1215 at Tucson, Arizona 9
Figure 5a	Same as Above for Winds of 0918G30 at Douglas, Arizona 10
Figure 5b	Same as Above for Winds of 0910-0915 at Douglas, Arizona 10
Figure 6	Same as Above for Winds of 3618G30 at Needles, California 11
Figure 7	Same as Above for Winds of 3415G25 at Blythe, California 11
Figure 8	Graphs of Pressure Gradient vs. Surface Winds for Needles and Blythe, California, and Flagstaff, AZ . 12
Figure 9	Graphs of Pressure Gradient vs. Surface Winds for Tucson, Douglas, and Yuma, Arizona 12
Figure 10	Graphs of Pressure Gradient vs. Surface Winds for Phoenix, Arizona 13

ARIZONA COOL SEASON CLIMATOLOGICAL SURFACE WIND
AND PRESSURE GRADIENT STUDY

Ira S. Brenner*
Weather Service Forecast Office
Phoenix Arizona

ABSTRACT. The average sea-level pressure gradients which produce sustained surface winds above 8 kts for at least six consecutive hours during the cool season at predetermined key stations in or adjacent to Arizona are investigated. Only wind directions from northerly or easterly components are included in the developmental data sample. Graphs are provided in the developmental data sample. Graphs are provided which relate the derived pressure gradients to varying surface wind speeds at each key station.

I. INTRODUCTION

Surface wind direction and sustained speed forecasts during the cool season are not only essential to statewide aviation interests, but also to surface travelers and numerous outdoor enthusiasts. Model Output Statistic (MOS) products forecast trends of surface wind speeds reasonably well, but frequently fall well short of acceptable levels of accuracy in many critical cases (Grayson and Tuft, 1978). Since sustained wind speeds of less than 8 kts are generally not a problem, this study concentrated only on sustained speeds of 8 kts or greater at key stations. Gustiness of 25 kts or greater was also included provided the 8 kts sustained threshold was satisfied. In order that these winds be included in the developmental data sample, they had to persist at the key station for 6 consecutive hours without a major shift in direction.

Winds from a southerly component, as are common in advance of an upper-level trough and associated cold front, were not considered in this study. It was felt that Arizona forecasters, as well as MOS, handle these wind direction and speed forecasts reasonably well.

During the period of study, differences in sea-level pressure in millibars from selected stations to each key station were tabulated for each six-hour period used and averaged in each of six predetermined sustained 6-hour wind-speed categories. The categories selected were 8-12 kts, 13-17 kts, 23-27 kts, 28-32 kts, and greater than 32 kts. Wind directions at the key station for each category were also tabulated and averaged. The number of cases per-wind category were additionally recorded.

Following this procedure, for each surrounding station to each key station, the combined average wind speed and the corresponding average direction and pressure gradient were determined and plotted on a map for that particular key station.

*Present affiliation: National Hurricane Center, Miami, Florida.

The key stations used in this study and the respective wind directions investigated for use in the developmental sample were as follows:

1. Yuma, AZ (YUM) 330° to 030°
2. Phoenix, AZ (PHX) 060° to 100°
3. Flagstaff, AZ (FLG) 030° to 070°
4. Tucson, AZ (TUS) 100° to 150°
5. Douglas, AZ (DUG) 070° to 120°
6. Needles, CA (EED) 330° to 030°
7. Blythe, CA (BLH) 320° to 020°.

Additional stations which were utilized to compute the sea-level pressure differences to the key stations included:

1. Las Vegas, NV (LAS)
2. Cedar City, UT (CDC)
3. Blanding, UT (BDG or 4BL)
4. Gallup, NM (GUP)
5. Deming, NM (DMN)
6. Winslow, AZ (INW).

II. DEVELOPMENTAL DATA ANALYSIS

The cool season was defined as the months of October through April. The period of study commenced October 1976 and ended April 1978. Figures 1-7 display the combined averaged wind direction, speed, and pressure gradients yielded by the developmental data for each of the key stations. The average sea-level pressure differences are in whole and tenths of a millibar. The difference, of course, at the key station being analyzed is always zero. The averaged wind direction and speed corresponding to the plotted pressure differences is indicated below the key station name.

In the cases of Figure 1 (YUM), Figure 2 (PHX), Figure 3 (FLG), Figure 4 (TUS), and Figure 5 (DUG), the 6-hour averages of wind speeds appeared to fall into two distinct categories and were so separated.

For all the key stations, the number of mutually exclusive 6-hour periods (N) that comprised the developmental data sample is indicated near the name of the key station. A total of four 6-hour periods is possible per day and 424 days were involved in the study. This yielded a maximum of 1696 six-hour periods possible in the study for each key station. Therefore, during this particular period of study, the winds which were used to develop Figures 1-7 were observed the following percentages of the total data sample:

STATION	AVERAGE WIND	N	PERCENT OF TOTAL SAMPLE	COMBINED PERCENT
YUM	3515G25	67	4%	9%
YUM	3510 to 3515	84	5%	
PHX	0815	32	2%	5%
PHX	0808 to 0813	56	3%	
FLG	0512G25	34	2%	6%
FLG	0510 to 0515	72	4%	
TUS	1215G25	45	3%	7%
TUS	1210 to 1215	71	4%	
DUG	0918G30	52	3%	8%
DUG	0910 to 0915	86	5%	
EED	3618G30	202	12%	12%
BLH	3415G25	192	11%	11%

At first glance, one might judge that these percentages are too small to be considered very meaningful. However, it must be remembered that these are percentages where moderate to strong sustained or gusty winds persisted for 6 consecutive hours. Aviators, boaters, and drivers of high profile vehicles would consider these sustained winds for this long a period very significant.

Combining the percentages where stronger and weaker winds were indicated (YUM, PHX, FLG, TUS, and DUG), and comparing these values with the remaining two stations, EED and BLH, provides a few additional points of interest. It would seem that the Colorado River region of western Arizona is rather susceptible to moderate to strong northerly winds. The key station farthest north within this river valley, EED, had the greatest percent frequency (12%) of the three stations analyzed, with YUM (southernmost) having the least (9%). This is borne out by Figures 1, 6, and 7 which show that the highest surface pressure for this pattern is normally located in the region between CDC and the Nevada border due west of CDC. The pressure gradients are indicated to be tighter between LAS and EED than between EED and YUM on each of these figures.

DUG, with a frequency of 8% for easterly winds was the fourth highest of the key stations. A review of the data indicated that this was largely in response to a southward plunge of high pressure into western New Mexico. This is commonly associated with a "backdoor-coldfront" situation. The frequency of 7% at TUS for southeast winds can be attributed to much the same reasoning as that applied to DUG. Figures 4 and 5 both show the southward plunge of higher pressure into western New Mexico. The tight pressure gradient indicated from Grand Canyon (GCN) to between FLG and INW and south-eastward toward ShowLow (SOW) and DMN suggests the presence of a frontal boundary.

The northeast-to-east winds at PHX and FLG, as displayed by Figures 2 and 3 appear to be associated with an intermediate position of the high-pressure area between that shown by Figures 1, 6, and 7 and that of Figures 4 and 5. At this time, winds begin to decrease considerably along the Colorado River, while increasing in southeastern Arizona.

Further examination of Figures 1-7 gives insight as to which of the surrounding stations give the best correlation to each key station in terms of pressure difference and resulting surface wind speeds. These are listed as follows:

KEY STATION	BEST CORRELATING STATION
YUM	LAS
PHX	INW
FLG	CDC
TUS	DMN
DUG	DMN
EED	CDC
BLH	LAS

At this point, an attempt was made to provide prognostic utility from this study. The original averaged pressure gradients and wind directions for each of the predetermined sustained 6-hour wind-speed categories were plotted in graphical form for the above key and correlating station combinations. The mid-point, rather than the actual range of each wind-speed category, was labeled along the abscissa with the average pressure gradient along the ordinate. The graphs are shown in Figures 8-10. The combined averaged data for each key station are also on the graph close to the respective curve for that station. Near each plotted point, the wind direction in degrees and the number of cases are indicated.

For all the graphs, a definite "S" configuration resulted. The crest of the "S" was located near 20 kts in each case. These curves indicate that wind speed does not increase in a simple linear fashion with increasing pressure gradient. The slopes of each of the curves between 15 kts and 20 kts are larger in all cases than elsewhere on that same curve. This means that it takes more pressure gradient to cause an increase in the winds in this range than for other wind-speed ranges.

Experience has shown that with steadily rising pressure to the north or east, actual surface winds do indeed frequently increase rapidly up to about 15 kts, then level off for awhile, before increasing sharply once again. Perhaps an explanation is that the full potential of the surface pressure gradient wind is generally realized at approximately 15 kts under these conditions. Beyond that, further significant increases in surface winds apparently are delayed until turbulent mixing brings down the stronger upper-level winds. This appears to occur on the average at about 20 kts. The EED, BLH, and DUG curves suggest that once this 20-kt threshold is achieved, surface winds continue to increase quite rapidly with the continuing increase of pressure gradient.

III. CONCLUSIONS

Discussion with other forecasters at PHX WSFO indicates that the results of this study correspond exceptionally well to what has been subjectively analyzed over past years. Although this is really more of a diagnostic rather than prognostic tool, it does sustain objective operational utility--not only as a potential forecast tool, but from a familiarization standpoint for new Arizona forecasters as well.

IV. ACKNOWLEDGMENTS

Special thanks are in order, once again, for my two favorite secretaries, Mrs. Tommie McCabe and Mrs. Evelyn Allan.

V. REFERENCES

Grayson, T., and W. Tuft, 1978: Does MOS Underforecast Wind Speeds? Maybe Yes, Maybe No! Western Region NWS Technical Attachment No. 78-27, National Weather Service Western Region, Salt Lake City, Utah.




FIGURE 1A. AVERAGE SEA-LEVEL PRESSURE GRADIENTS THAT PRODUCE AVERAGE SUSTAINED SURFACE WINDS OF 3515025 AT YUMA, ARIZONA, FOR SIX CONSECUTIVE HOURS.




FIGURE 1B. SAME AS FIGURE 1A BUT FOR WINDS OF 3510-3515 AT YUMA, ARIZONA.




FIGURE 2A. AVERAGE SEA-LEVEL PRESSURE GRADIENTS THAT PRODUCE AVERAGE SUSTAINED SURFACE WINDS OF 0815 AT PHOENIX, ARIZONA, FOR SIX CONSECUTIVE HOURS.




FIGURE 2B. SAME AS FIGURE 2A BUT FOR WINDS OF 0808-0813 AT PHOENIX, ARIZONA. -7-




FIGURE 3A AVERAGE SEA-LEVEL PRESSURE GRADIENTS THAT PRODUCE AVERAGE SUSTAINED SURFACE WINDS OF 0512G25 AT FLAGSTAFF, ARIZONA, FOR SIX CONSECUTIVE HOURS.




FIGURE 3B SAME AS FIGURE 3A BUT FOR WINDS OF 0510-0515 AT FLAGSTAFF, ARIZONA.




FIGURE 4A AVERAGE SEA-LEVEL PRESSURE GRADIENTS THAT PRODUCE AVERAGE SUSTAINED SURFACE WINDS OF 1215025 AT TUCSON, ARIZONA, FOR SIX CONSECUTIVE HOURS.




FIGURE 4B SAME AS FIGURE 4A BUT FOR WINDS OF 1210-1215 AT TUCSON, ARIZONA.




FIGURE 5A AVERAGE SEA-LEVEL PRESSURE GRADIENTS THAT PRODUCE AVERAGE SUSTAINED SURFACE WINDS OF 0918030 AT DOUGLAS, ARIZONA, FOR SIX CONSECUTIVE HOURS.




FIGURE 5B SAME AS FIGURE 5A BUT FOR WINDS OF 0910-0915 AT DOUGLAS, ARIZONA.




FIGURE 6 AVERAGE SEA-LEVEL PRESSURE GRADIENTS THAT PRODUCE AVERAGE SUSTAINED SURFACE WINDS OF 361830 AT NEEDLES, CALIFORNIA, FOR SIX CONSECUTIVE HOURS.




FIGURE 7. AVERAGE SEA-LEVEL PRESSURE GRADIENTS THAT PRODUCE AVERAGE SUSTAINED SURFACE WINDS OF 3415025 AT BLYTHE, CALIFORNIA, FOR SIX CONSECUTIVE HOURS.




FIGURE 8. GRAPHS OF PRESSURE GRADIENT VS. SURFACE WINDS FOR NEEDLES AND BLYTHE, CALIFORNIA, AND FLAGSTAFF, ARIZONA.




FIGURE 9. GRAPHS OF PRESSURE GRADIENT VS. SURFACE WINDS FOR TUCSON, DOUGLAS, AND YUMA, ARIZONA.

PRESSURE GRADIENT VS. SURFACE WINDS (SUSTAINED)

DDD AVERAGE WIND DIRECTION FOR THE CORRESPONDING WIND SPEED
 NNN NUMBER OF CASES FOR THE CORRESPONDING WIND SPEED
 INTERSECTION POINT OF PRESSURE GRADIENT AND RESULTING SURFACE WIND




FIGURE 10. GRAPHS OF PRESSURE GRADIENT VS. SURFACE WINDS FOR PHOENIX, ARIZONA.

NOAA Technical Memoranda NWSNR: (Continued)

- 82 Smoke Management in the Willamette Valley. Earl W. Bates, May 1974. (OO-74-11277/AS)
- 83 An Operational Evaluation of 200-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (OO-74-11487/AS)
- 94 Conditional Probability of Visibility Loss from One-Half Mile to Radiation Fog at Fresno, California. John D. Thomas, August 1974. (OO-74-1555/AS)
- 96 Map Type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (OO-75-10428/AS)
- 97 Eastern Pacific Cut-off Low of April 21-23, 1974. William J. Alder and George R. Miller, January 1976. (PB-258-711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (OO-75-10712/AS)
- 99 A Study of Flash Flood Susceptibility—A Basin in Southern Arizona. Gerald Williams, August 1975. (OO-75-1380/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuff, October 1975. (PB-245-902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB-255-035/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB-252-366/AS)
- 105 Forecasting the Mono Wind. Charles P. Rascha, Jr., February 1976. (PB-254-650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John G. Plankinton, Jr., March 1976. (PB-254-349)
- 107 Map Types as Aids in Using MOS Maps in Western United States. Ira S. Brenner, August 1976. (PB-259-594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB-260-457/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, Sept. 1976. (PB-273-677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Benney, November 1976. (PB-264-655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB-265-941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Caré, February 1977. (PB-275-394/AS)
- 114 Tropical Cyclone Kahlögen. James R. Fors, February 1977. (PB-275-576/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Selman, April 1977. (PB-268-847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-value. R. F. Quiring, April 1977. (PB-272-631)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB-268-740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Barry Rinderson, June 1977. (PB-271-290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R. F. Quiring, June 1977. (PB-271-304/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Mobb, Jr., June 1977. (PB-271-742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Loffwich, August 1977. (PB-272-661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Loffwich and Charles J. Neumann, August 1977. (PB-273-553/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB-281-367/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Guésgat, April 1978. (PB-283-036/AS)
- 129 Fire Whirls. David W. Coens, May 1978. (PB-283-866/AS)
- 130 Flash-Flood Procedure. Ralph G. Hatch and Gerald Williams, May 1978. (PB-286-014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olson, September 1978. (PB-289-016/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-740 Weather Radar. R. G. Pappas, R. Y. Lee, and B. W. Flinke, October 1978. (PB-289-767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB-291-517/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB-291-195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB-292-716/AS)
- 136 Basic Synoptic Principles. Thomas L. Dietrich, January 1979. (PB-292-247/AS)
- 137 LFM 24-hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Rascha, Jr., January 1979.
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979.
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979.
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather District. James McLeamb, April 1979.
- 141 Comparison of LFM and WRF Precipitation Guidance for Nevada During Boreas. Christopher Hill, April 1979.
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Coray, April 1979.
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

NOAA, the *National Oceanic and Atmospheric Administration*, was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth, and to assess the socioeconomic impact of natural and technological changes in the environment.

The six Major Line Components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS — Important definitive research results, major techniques, and special investigations.

TECHNICAL REPORTS—Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS — Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

CONTRACT AND GRANT REPORTS—Reports prepared by contractors or grantees under NOAA sponsorship.

TECHNICAL SERVICE PUBLICATIONS—These are publications containing data, observations, instructions, etc. A partial listing: Data serials; Prediction and outlook periodicals; Technical manuals, training papers, planning reports, and information serials; and Miscellaneous technical publications.

ATLAS—Analysed data generally presented in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.



Information on availability of NOAA publications can be obtained from:

**ENVIRONMENTAL SCIENCE INFORMATION CENTER
ENVIRONMENTAL DATA SERVICE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE**

3300 Whitehaven Street, N.W.
Washington, D.C. 20235