


NOAA Technical Memorandum NWS WR-179



STRATIFIED MAXIMUM TEMPERATURE RELATIONSHIPS BETWEEN SIXTEEN
ZONE STATIONS IN ARIZONA AND RESPECTIVE KEY STATIONS

Salt Lake City, Utah
June 1983

**U.S. DEPARTMENT OF
COMMERCE**

/ National Oceanic and
Atmospheric Administration

/ National Weather
Service



NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P. O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U. S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22151. Prices vary for all paper copy; \$3.50 microfiche. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operational Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
 - 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
 - 28 Weather Extremes. R. J. Schmidli, April 1968 (revised February 1981).
 - 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB-178425)
 - 30 Numerical Weather Prediction and Synoptic Meteorology. Capt. Thomas D. Murphy, U.S.A.F., May 1968. (AD-673365)
 - 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB-179084)
 - 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB-179289)
 - 35 Joint ESSA/FAA ARTC Radar Weather Surveillance Program. Herbert P. Benner and DeVon B. Smith, December 1968 (revised June 1970). AD-681857)
 - 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (PB-183055)
 - 37 Disposal of Logging Residues without Damage to Air Quality. Owen P. Cramer, March 1969. (PB-183057)
 - 39 Upper-Air Lows over Northwestern United States. A. L. Jacobson, April 1969. (PB-184296)
 - 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970's. L. W. Snellman, August 1969. (PB-185068)
 - 42 Analysis of the Southern California Santa Ana of January 15-17, 1966. Barry B. Aronovitch, August 1969. (PB-185670)
 - 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB-185762)
 - 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB-187763)
 - 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB-190476)
 - 47 Statistical Analysis as a Flood Routing Tool. Robert J. C. Burnash, December 1969. (PB-188744)
 - 48 Tsunami. Richard P. Augulis, February 1970. (PB-190157)
 - 49 Predicting Precipitation Type. Robert J. C. Burnash and Floyd E. Hug, March 1970. (PB-190962)
 - 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB-191743)
 - 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB-193102)
 - 52 Sacramento Weather Radar Climatology. R. G. Pappas and C. M. Veliouette, July 1970. (PB-193347)
 - 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
 - 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB-194394)
 - 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB-194389)
 - 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB-194710)
 - 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM-71-00017)
 - 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM-71-00016)
- NOAA Technical Memoranda (NWS WR)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon. Arthur W. Fritz, October 1970. (COM-71-00120)
 - 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM-71-00349)
 - 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
 - 65 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM-71-00829)
 - 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM-71-00956)
 - 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM-72-10433)
 - 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM-72-10554)
 - 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM-72-10707)
 - 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM-72-11140)
 - 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM-72-11136)
 - 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM-72-11146)
 - 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM-72-10920)
 - 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM-72-10021)
 - 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM-73-10150)
 - 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (revised March 1978). (COM-73-10251)
 - 83 A Comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM-73-10669)
 - 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM-73-11264)
 - 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert V. G. Lee, June 1973. (COM-73-11276)
 - 89 Objective Forecast Precipitation over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, Sept. 1973. (COM-73-11946/3AS)
 - 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM-73-10465)
 - 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM-74-11277/AS)
 - 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM-74-11407/AS)
 - 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM-74-11555/AS)
 - 96 Map Type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM-75-10428/AS)
 - 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB-250-711/AS)
 - 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM-75-10719/AS)
 - 99 A Study of Flash Flood Susceptibility--A Basin in Southern Arizona. Gerald Williams, August 1975. (COM-75-11360/AS)
 - 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB-246-902/AS)
 - 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB-253-053/AS)
 - 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB-252-866/AS)
 - 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB-254-650)
 - 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB-254-649)
 - 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB-259-594)
 - 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB-260-437/AS)
 - 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB-273-677/AS)
 - 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB-264-655/AS)
 - 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB-265-941/AS)
 - 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB-273-694/AS)
 - 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB-273-676/AS)
 - 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB-268-847)
 - 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R. F. Quiring, April 1977. (PB-272-831)
 - 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB-268-740)
 - 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB-271-290/AS)

NOAA Technical Memorandum NWS WR-179

STRATIFIED MAXIMUM TEMPERATURE RELATIONSHIPS BETWEEN SIXTEEN
ZONE STATIONS IN ARIZONA AND RESPECTIVE KEY STATIONS

Ira S. Brenner

National Weather Service Forecast Office
Phoenix, Arizona
June 1983


UNITED STATES
DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary

National Oceanic and
Atmospheric Administration
John V. Byrne, Administrator

National Weather
Service
Richard E. Hallgren, Director



This publication has been reviewed
and is approved for publication by
Scientific Services Division,
Western Region.

A handwritten signature in cursive script that reads "Glenn E. Rasch".

Glenn E. Rasch, Chief
Scientific Services Division
Western Region Headquarters
Salt Lake City, Utah

TABLE OF CONTENTS

	<u>Page</u>
I. Background Information.....	1
II. Data Tabulation.....	2
III. Data Analysis and Results.....	3
IV. Applications.....	5
V. Conclusions.....	6
VI. Acknowledgments.....	7
VII. References.....	7
Figures 1 through 4.....	8-12
Data Tables 1 through 4.....	13-18

STRATIFIED MAXIMUM TEMPERATURE RELATIONSHIPS BETWEEN SIXTEEN
ZONE STATIONS IN ARIZONA AND RESPECTIVE KEY STATIONS

Ira S. Brenner
National Weather Service Forecast Office
Phoenix, Arizona

I. BACKGROUND INFORMATION

This study involved an extensive tabulation of temperature relationships between each of sixteen zone stations in Arizona and respective key stations. Key stations were defined as the five cities in Arizona for which Model Output Statistics (MOS) temperature forecasts are generated, plus the city of Prescott. Each of these cities was considered to be meteorologically representative of the climatic zone in which it is located (Figure 1). For example, Prescott would be the key station for the central basin and northwest zone of Arizona (average elevations of the zone stations 3000 to 6000 feet). Similarly, Flagstaff would be the key station for the central mountain zone (elevations above 6000 feet).

This research used a temperature and season dependent stratification to expand upon the basic principle of using normal monthly maximum and minimum temperature deviations between various zone stations and the appropriate key station as a guide to forecasting temperatures at each of the zone stations. Normal monthly maximum and minimum temperatures at all twenty two of the stations for which temperature forecasts are prepared in Arizona are readily available at the National Weather Service Forecast Office in Phoenix, Arizona (WSFO PHX). Thus, the average monthly differences from each of the sixteen zone stations to the respective key station have been computed for both maximums and minimums. The differences, or deviations, are used as guidance in the preparation of the Arizona community and recreational area forecasts of maximum and minimum temperatures for the zone stations.

However, the use of normals which have been derived for an entire month at a given station was theorized to have inherent weaknesses. Normals are made up of extremes; and extremes are typically masked when normals are tabulated over a large period of record. In Arizona, extremes in the temperatures at a given station during a given month can be common, and can be induced by small scale changes just as easily as by adjustments of the larger scale features. For example, during the cool season, the southern latitude of Arizona can typically allow strongly and rapidly rebounding temperatures following an unusually cold outbreak. Similarly, very cold outbreaks can typically follow a period of unusually warm readings. During the summer monsoon season (July and August), clouds and thunderstorms can cause wide day-to-day variances in temperatures. In general, varying degrees of winds, clouds, humidity, precipitation (or lack of), and snow cover, can induce considerable day-to-day variances and anomalies in either the maximum or minimum temperature at a given station, or even within a given area of the state.

It was felt that the eventual value of the observed maximum or minimum temperature should be a direct byproduct of the overall character of the concurrent synoptic regime. For example, a high temperature of only fifty five degrees at Tucson in the middle of September can generate rather strong inferences about the existing synoptic weather regime. During a "normal" synoptic regime for the same time of year, the average

maximum temperature at Douglas is five degrees cooler than Tucson. When Tucson has an anomalous high temperature of fifty five degrees in the middle of September, Douglas averages two degrees warmer, as opposed to the "normal" five degrees cooler, than Tucson.

It is not so important to know the exact cause of the unusually low maximum at the key station in this example. The important point is that the majority of similar low maximums which have occurred in September at Tucson in the past were likely caused by similar or related synoptic regimes. And in the majority of those similar regimes, Douglas has typically reacted by averaging two degrees warmer than Tucson.

Therefore, this study set out to substantiate the above idea by deriving a set of relationships between observed key and zone station temperatures that are temperature as well as seasonally dependent. It was hoped that the outcome of this stratification would ultimately provide a more representative means for relating the forecasted temperatures at key stations to those of the zone stations during anomalous weather regimes.

II. DATA TABULATION

The period selected as the data base was from July 1971 through August 1981. Six key stations were selected (Figure 2). These stations included those normally transmitted as the coded cities forecasts by the Automation of Field Operations and Services (AFOS) system (under the heading PHXCCFPHX). The key stations chosen were Phoenix (PHX), Tucson (TUS), Flagstaff (FLG), Winslow (INW), Yuma (YUM), and Prescott (PRC). Shown as well in Figure 2 are the sixteen zone stations for which routine temperature and precipitation probability forecasts are prepared by Phoenix WSFO (under the AFOS heading PHXRECPHX).

Five characteristically similar seasonal periods were selected for the purpose of data stratification. The periods were grouped as follows:

1. November through February
2. March through April
3. May through June
4. July through August
5. September through October.

Figure 3 represents a sample of the tabulation forms used to record the maximum temperature data during the period of study. The key station Winslow is used as an example for the seasonal period March through April. The observed maximum temperatures at Winslow for each day of every March and April during the period of study were stratified according to the appropriate five degree temperature range. As each daily maximum was identified with the proper temperature range, the corresponding maximum temperatures at each of the two Winslow zone stations for that same day were recorded within the appropriate data entry square.

During the course of this data tabulation, it became obvious that "reset" maximum temperatures were going to partially contaminate the data sample. These resets occasionally occur at stations which read and reset the maximum thermometers only once daily at 4 p.m. MST. Should the next day be cooler, the official high temperature for that day will be reported as the 4 p.m. temperature from the previous day. In some of these cases,

subjective adjustments to these resets could be made by noting a consistent amount of 24 hour maximum temperature drop at several surrounding stations that reset their thermometers twice a day. In all but the most obvious cases, adjustments were not made, and the reset temperature was not included in the data tabulation.

Returning to the Winslow example, temperature data at each zone station for each March and April during the entire period were recorded. Tabulations were then made of the mean of these temperatures, the standard deviation, and the number of entries for each zone station within each five degree Winslow temperature range. Finally, the difference or deviation of each mean temperature from the midpoint of each temperature range for Winslow was determined for each individual zone station.

The above process, using the appropriate maximum temperature data, was repeated for the other key stations of Tucson, Flagstaff, and Prescott. The key station Yuma did not have any related zone stations. The key station Phoenix only had one affiliated zone station at Coolidge. Since maximum temperatures at Coolidge are consistently very similar to those of Phoenix, the stratification process seemed to provide little additional advantage over that of simply using the normal monthly deviation. Therefore, it was decided to continue to use the normal monthly deviation of the Coolidge maximum from Phoenix.

A similar stratification process was also attempted for minimum temperature data during the period of study for all key stations. However, this procedure was discarded near the midpoint of the data tabulation. The less conservative nature of minimum temperatures rapidly became apparent due to the wide variation of minimum temperatures being recorded for each zone station within each appropriate key station temperature range. Therefore, the use of average monthly deviation of minimum temperatures at each zone station from the corresponding key station is preferred over the results of a separate stratification process for minimum temperatures.

III. DATA ANALYSIS AND RESULTS

This discussion will be confined to the results of the stratification process of maximum temperatures only. Data Tables 1 through 4 display the tabulated results of the data collection for the key stations Tucson, Prescott, Flagstaff, and Winslow. For each five degree temperature range, the mean temperature $[\bar{x}]$, the deviation $[\text{dev}]$ of that mean temperature from the midpoint of the temperature range, the standard deviation $[s]$, and the number of cases $[n]$ used to compile the above values, are listed by key station and seasonal period.

One of the more positive results of this tabulation is the low standard deviations $[s]$. Figure 4 is a summarized analysis of $[s]$ from the study. In Figure 4, the $[s]$ values for each key station temperature range in the data tables were combined and averaged within each seasonal period by zone station. Only those $[s]$ values that were derived from a data sample of $[n]$ greater than or equal to 10 for that particular key station temperature range were included in the averaging process. The results of the averaging were grouped by range of $[s]$ for each zone station in the first portion of Figure 4. Following this, individual $[s]$ values for zone stations were averaged to determine a zone average for

each key station within each seasonal period. These results are listed by ranges in the second portion of Figure 4.

The fact that the majority of these station averages of $[s]$ were 3.50 degrees or less supports a strong relationship between the maximum temperature of the key station and the corresponding maximum temperature at the respective zone stations. Further, these zone station standard deviation averages were actually overwhelmingly in the 2.50 to 3.00 degree grouping for the Tucson and Prescott zones. Certainly, if the maximum temperatures at the individual zone stations were not correlated to the maximum temperatures at the corresponding key station, these standard deviation averages would have been much larger. Note as well that for each key station, it should be expected that the largest values of these zone station averages of $[s]$ would occur during the rather changeable November through February seasonal period. Despite this, the averages within this period were still quite good, being near or below 4.00 degrees.

The individual zone station averages of standard deviation in Figure 4 also show which stations had the strongest or weakest correlations with their key station. The zone stations Bisbee (BIB), Globe (GLB), McNary (MNY), and Page (PGA) appear to consistently exhibit the weakest relationships, while Nogales (NOG) and Payson (OE4) have relatively strong relationships. The stations Safford (E74), Kingman (IGM), Show Low (E03), and Canyon de Chelly (CNC) also seem to trend towards an overall weak relationship to the key station, while Fort Huachuca (FHU), Douglas (DUG), Grand Canyon (GCN), Cottonwood (COT), and Sedona (SED) lean in the direction of a somewhat stronger relationship.

A considerable amount of discussion could be generated as to the reason for a given degree of relationship between a zone station and the corresponding key station. These relationships could be controlled by a number of factors ranging perhaps from the distance to the key station, to differences in elevation, to station exposure and local effects. Nevertheless, the varying degrees of these individual station relationships can be utilized in the positive sense as a confidence factor by the forecaster. The important point is that the results of the standard deviation analysis indicate that overall, the relationships between the key stations and their respective zone stations are sound.

Perhaps the foremost positive result of this study is portrayed by the distributions of the deviations $[dev]$ of the mean temperature $[x]$ at a given zone station from the midpoint of the appropriate key station temperature range. The primary hypothesis which provided the foundation for this study was that the monthly temperature deviations at a given zone station were not representative during anomalous weather patterns. Data Tables 1 through 4 clearly validate this hypothesis. For nearly all zone stations, a substantial variance in $[dev]$ occurs as one propagates away from the key station temperature ranges containing the maximum $[n]$.

Three separate patterns were observed relating to the manner in which the values of $[dev]$ changed at the zone stations within each seasonal period. The patterns are summarized in Figure 5. Whenever a particular pattern was identified at the majority of the zone stations affiliated with a given key station, that key station was entered in Figure 5 for that pattern under the appropriate seasonal period.

The most common pattern observed is that the $[dev]$ becomes increasingly less positive (or more negative) from the coldest anomalies for a

given seasonal period toward the warmest anomalies. Of those key stations which fall within this category, about half exhibit a change in sign of the value of [dev] from positive to negative.

The second pattern, which surfaced rather infrequently, is a small change in [dev] of 3 degrees or less from the coldest to the warmest anomalies. Occurrences of this pattern favor the transition months of March/April and September/October.

The final pattern, which only occurs once, is where the values of [dev] become increasingly negative toward each of the coldest and warmest anomalies. The single occurrence is noted for the Tucson zone stations during the November/February seasonal period. Applying the most common pattern, where [dev] becomes less positive (or more negative) from the coldest to the warmest anomaly, one would have expected different values of [dev] for the 36 to 40 degree range, as well as the 41 to 45 degree interval. These should have ranged from about plus two at Nogales to minus one at Safford and Douglas, to minus three at Fort Huachuca, and minus four at Bisbee. These projected values would have been quite reasonable when considering the effects of winter temperature inversions in the Tucson valley. These temperature inversions will typically limit the amount of difference between the maximum temperature at Tucson and the maximums at other Tucson zone stations which are at higher elevations and generally above the inversion level. Yet, at the coldest anomalies, the gap between the Tucson maximum and the maximums at the Tucson zone stations begins to widen.

A possible explanation could rest with "backdoor coldfronts" and other incidences of low level easterly flow generated by a buildup of surface pressure over the Southern Rockies and Southern Plains. During these not-so-infrequent occurrences, colder air in the lower levels pushes into southeast Arizona, typically generating a cold anomalous situation. However, the topography of southeast Arizona is such that this air can modify considerably, due to downslope effects, by the time it reaches the Tucson valley. This downslope condition tends to negate the effects of the cold air at Tucson, while the remainder of the Tucson zone stations undergo rather strong low-level cold-air advection. This could possibly account for the uniqueness of this particular [dev] pattern.

IV. APPLICATIONS

The results of this study are routinely applied to the development of the community and recreational forecasts of temperatures and probabilities of precipitation for sixteen zone stations in Arizona (AFOS heading PHXRECPHX). The appropriate AFOS applications program is initiated by entering the run line command, "RUN:RECS AAA BBB CCC D E F ". The first, second, and third period temperature forecasts for Prescott are entered for AAA, BBB, and CCC, respectively. The single digit precipitation probabilities are also entered for D, E, and F. This probability portion of the run line will also accept a "-" or A "+" as well in lieu of five percent and one hundred percent, respectively. The program then reads the coded cities forecast product (PHXCCFPHX) to obtain the temperature and probability of precipitation forecasts for the remainder of the key stations.

The seasonal period and the hour of the forecast being prepared (early morning or afternoon) are determined within the program using the computer clock. Therefore, given these parameters, the forecasted temperatures at each key station from the PHXCCFPHX product, and the Prescott information from the run line, a series of searches and calculations commences. For the periods which involve maximum temperature forecasts, the results of the stratification process are applied toward the calculations. For minimum temperatures, calculations using normal monthly deviations from zone station to key station are performed. Probabilities at the key station are simply assigned to each zone station related to that key station.

Within seconds, a completed PHXRECPHX product can be displayed at, and even transmitted from, the AFOS console. It is at this time, however, that any adjustments that might be necessary can be made to the AFOS-generated PHXRECPHX product.

Generally, it has been found that the results of the calculations, as displayed, need only minor adjustments, if at all. This is particularly true during nonchanging regimes where unusual local effects are not operative. However, the need for subjective adjustments should be considered when the weather conditions at the key station and the appropriate zone stations are expected to differ significantly enough to upset the basic dependency relationship. Among examples of situations that could be considered are local fog or low clouds, separate areas with snow on the ground, local winds, isolated areas of precipitation, and air mass boundaries. Similarly, situations where a portion of a zone has clouds (for example, lingering clouds behind an exiting storm), while the remainder of the zone is clear, should be included for consideration.

During persistent regimes when subjective adjustments would not normally be needed, it has proven beneficial to compute a quick verification of the maximum temperature forecasts from several previous days. This check can occasionally identify a temporary bias in the stratification process at a given zone station, which could then be applied to the current forecast, provided the prevailing persistent regime is expected to continue.

V. CONCLUSIONS

The results of this study support the original hypothesis that normal monthly deviation relationships of maximum temperature between a given zone station and its key station are largely unrepresentative during anomalous weather regimes. A much stronger relationship was identified using a temperature and seasonally dependent mean difference or deviation between the maximum temperature at a given zone station, and that of the respective key station.

The observed maximum temperature at a key station is accepted as generally being a function of the overall synoptic regime operative at the time. The study validated the premise that, in general, maximum temperatures at each zone station are dependent upon the actual value of the maximum temperature at the respective key station. It follows that the maximum temperatures at the zone stations can be related indirectly to the character of the overall existing synoptic regime as well.

Therefore, it is concluded that the wide range of variation in the magnitude of the mean deviation of the maximum temperature between each zone and respective key station is indeed a result of sensitivity and dependency upon the overall characteristic of the synoptic regime which produced the observed key station maximum. Similar synoptic regimes can thus be expected to typically produce similar mean deviations between a given zone and key station. However, subjective adjustments to the mean deviations must be considered during situations where the basic dependency relationship is expected to be altered.

In the case of minimum temperature relationships, it appears that the basic dependency relationship is too easily overcome by small-scale effects. The resulting less conservative nature of minimum temperatures therefore increases the possible need for a regression analysis at each zone station to assist in determining the average effects of each small-scale parameter on the observed minimum temperature. The results of this analysis, if satisfactory, could perhaps then be applied to the value obtained at the zone station from using the normal monthly difference in minimum temperatures between the key and zone station.

VI. ACKNOWLEDGMENTS

Special thanks are in order for Dave Toronto. Without his programming expertise, I would still be trying to automate the results of this study.

VII. REFERENCES

- [1] Monthly Normals of Temperature, and Heating and Cooling Degree Days 1951-80 -- Arizona. Climatography of The United States No. 81, National Climatic Center, Asheville, North Carolina, pp. 1-5, Sept. 1982.
- [2] Sellers, William D., and Richard H. Hill (editors). Arizona Climate 1931-72, University of Arizona Press, Tucson, 54 pp., 1974.




FIGURE 1

KEY STATIONS AND CORRESPONDING ZONE STATIONS

KEY STATION	#1 PHOENIX (PHX)	#2 TUCSON (TUS)	#3 PRESCOTT (PRC)	#4 FLAGSTAFF (FLG)	#5 WINSLOW (INW)	#6 YUMA (YUM)
ZONE STATION AND NUMBER	#1 COOLIDGE (COL)	#2 NOGALES (NOG)	#7 KINGMAN (IGM)	#12 GRAND CYN (GCN)	#15 CYN DE (CNC)	
		#3 SIERRA VISTA (FHU)	#8 GLOBE (GLB)	#13 SHOW LOW (E03)	#16 PAGE (PGA)	
		#4 BISBEE (BIB)	#9 COTTONWOOD (COT)	#14 MCNARY (MNY)		
		#5 DOUGLAS (DUG)	#10 SEDONA (SED)			
		#6 SAFFORD (E74)	#11 PAYSON (0E4)			

FIGURE 2

MONTHS-MARCH/APRIL KEY STATION-INW

KEY STATION TEMP. RANGE	36- 40	41- 45	46- 50	<--- TO ---> <--- TO --->	81- 85

1972					
CYN DE CHELLY	44	44	47, 49, 50, 51		78, 79, 76, 78, 77, 73, 76
.....					
PGA	47	46	54, 54, 50, 53		83, 81, 81, 82, 81, 83, 81

1973					
CYN DE CHELLY	45	46, 46	48, 47, 47		77, 73, 77, 76 77
.....					
PGA	45	48, 48	50, 54, 50		75, 78, 79, 79 78

↓	↓	↓	↓	↓	↓

1981					
CYN DE CHELLY	44	46, 46	48, 47, 48		79, 81, 81
.....					
PGA	46	48, 49	50, 52, 54		73, 78, 76

FIGURE 3

KEY STATION AND SEASONAL PERIOD	INDIVIDUAL ZONE STATION AVERAGES OF STANDARD DEVIATION					ZONE AVERAGE OF STANDARD DEVIATION			
	2-2.50	2.50-3	3-3.50	3.50-4	4-4.50	2.50-3	3-3.50	3.50-4	4-4.50
TUS									
NOV - FEB			NOG/FHU DUG/E74		BIB		3.44		
MAR - APR			NOG/FHU DUG/E74	BIB		2.87			
MAY - JUN	FHU		NOG/DUG BIB/E74			2.86			
JUL - AUG	NOG		FHU/BIB DUG/E74			2.71			
SEP - OCT			NOG FHU/BIB DUG/E74			3.13			

KEY STATION AND SEASONAL PERIOD	INDIVIDUAL ZONE STATION AVERAGES OF STANDARD DEVIATION					ZONE AVERAGE OF STANDARD DEVIATION			
	2-2.50	2.50-3	3-3.50	3.50-4	4-4.50	2.50-3	3-3.50	3.50-4	4-4.50
PRC									
NOV - FEB		0E4	COT/SED	IGM/GLB			3.43		
MAR - APR			COT/SED 0E4	IGM/GLB		2.92			
MAY - JUN			IGM/COT SED/0E4	GLB		2.86			
JUL - AUG			IGM/GLB COT/SED 0E4			2.72			
SEP - OCT			IGM/COT SED/0E4	GLB		2.99			

KEY STATION AND SEASONAL PERIOD	INDIVIDUAL ZONE STATION AVERAGES OF STANDARD DEVIATION					ZONE AVERAGE OF STANDARD DEVIATION			
	2-2.50	2.50-3	3-3.50	3.50-4	4-4.50	2.50-3	3-3.50	3.50-4	4-4.50
FLG									
NOV - FEB				GCN	E03/MNY				4.02
MAR - APR			GCN	E03/MNY			3.57		
MAY - JUN			GCN/E03	MNY		3.35			
JUL - AUG		GCN	E03/MNY			3.03			
SEP - OCT			GCN/E03	MNY		3.49			

KEY STATION AND SEASONAL PERIOD	INDIVIDUAL ZONE STATION AVERAGES OF STANDARD DEVIATION					ZONE AVERAGE OF STANDARD DEVIATION			
	2-2.50	2.50-3	3-3.50	3.50-4	4-4.50	2.50-3	3-3.50	3.50-4	4-4.50
INW									
NOV - FEB				CNC	IPGA		3.96		
MAR - APR			CNC	IPGA		3.30			
MAY - JUN			CNC	IPGA			3.61		
JUL - AUG			CNC/PGA			3.20			
SEP - OCT			CNC	IPGA		3.47			

FIGURE 4

PATTERNS IN THE CHANGE IN [DEV]

1. LESS POSITIVE FROM COLDEST ANOMALY TO WARMEST ANOMALY
 (AN * MEANS A CHANGE OF SIGN TO NEGATIVE AT THE
 MAJORITY OF THE ZONE STATIONS)

NOV/FEB	MAR/APR	MAY/JUN	JUL/AUG	SEP/OCT
PRC	TUS	TUS*	TUS	TUS*
FLG*	INW*	PRC	PRC	FLG*
INW*		FLG	FLG*	
		INW	INW*	

2. LITTLE CHANGE (3 DEGREES OR LESS ON THE AVERAGE) FROM
 THE COLDEST TO THE WARMEST ANOMALY

NOV/FEB	MAR/APR	MAY/JUN	JUL/AUG	SEP/OCT
	PRC			PRC
	FLG			INW

3. LESS POSITIVE TOWARD THE COLD AND WARM ANOMALIES

NOV/FEB	MAR/APR	MAY/JUN	JUL/AUG	SEP/OCT
TUS				

FIGURE 5

KEY STN.	KEY STN. RANGE FOR MAXIMUMS	6-10	11-15	16-20	21-25	26-30	31-35	36-40	41-45	46-50	51-55	56-60	61-65	66-70	71-75	76-80	81-85	86-90	91-95	96-100	101-105	106-110	
		DEV																					
		X																					
TUS		S																					
(JUL-AUG)	NOG	N																					
		DEV																					
		X																					
		S																					
	FHU	N																					
		DEV																					
		X																					
		S																					
	BIB	N																					
		DEV																					
		X																					
		S																					
	DUG	N																					
		DEV																					
		X																					
		S																					
	E74	N																					
		DEV																					
		X																					
TUS		S																					
(SEP-OCT)	NOG	N																					
		DEV																					
		X																					
		S																					
	FHU	N																					
		DEV																					
		X																					
		S																					
	BIB	N																					
		DEV																					
		X																					
		S																					
	DUG	N																					
		DEV																					
		X																					
		S																					
	E74	N																					

KEY STN.	KEY STN. RANGE FOR MAXIMUMS		6-10	11-15	16-20	21-25	26-30	31-35	36-40	41-45	46-50	51-55	56-60	61-65	66-70	71-75	76-80	81-85	86-90	91-95	96-100	101-105	106-110				
PRC (JUL-AUG)	IGM	DEV																									
		\bar{X}																									
		S																									
		DEV																									
		\bar{X}																									
		S																									
	GLB	DEV																									
		\bar{X}																									
		S																									
	COT	DEV																									
		\bar{X}																									
		S																									
	SED	DEV																									
		\bar{X}																									
		S																									
	OEA	DEV																									
		\bar{X}																									
		S																									
PRC (SEP-OCT)	IGM	DEV																									
		\bar{X}																									
		S																									
	GLB	DEV																									
		\bar{X}																									
		S																									
	COT	DEV																									
		\bar{X}																									
		S																									
	SED	DEV																									
		\bar{X}																									
		S																									
	OEA	DEV																									
		\bar{X}																									
		S																									

KEY STN.	KEY STN. RANGE FOR MAXIMUMS	6-10	11-15	16-20	21-25	26-30	31-35	36-40	41-45	46-50	51-55	56-60	61-65	66-70	71-75	76-80	81-85	86-90	91-95	96-100	101-105	106-110
	DEV			+6.00	+3.82	+1.62	-.84	-1.39	-2.87	-3.65	-4.01	-3.85	-4.19	-4.04	-3.61	-4.08						
	X			24.00	26.82	29.62	32.16	36.61	40.13	44.35	48.99	54.15	58.81	63.96	69.39	73.92						
INW (NOV-FEB)	GDC	S		3.04	3.46	4.27	4.33	4.33	4.38	4.38	4.36	4.31	4.20	3.23	2.46	2.37						
	N			5	11	33	61	86	141	183	227	159	117	74	36	12						
	DEV			+9.95	+7.40	+4.65	+2.00	-.48	-2.11	-3.78	-5.42	-6.51	-7.79	-8.35	-8.89	-10.09						
	X			27.95	30.40	32.65	35.00	37.52	40.89	44.22	47.58	51.49	55.21	59.65	64.11	67.91						
	PGA	S		3.11	3.44	4.40	4.16	4.42	4.48	4.46	4.42	4.36	4.35	3.91	3.26	3.21						
	N			4	12	33	58	84	131	173	204	152	113	71	35	11						
	DEV							+4.37	+ .76	-2.46	-4.36	-4.43	-4.35	-3.89	-3.53	-4.56	-5.64					
	X							42.37	43.76	45.54	48.64	53.57	58.65	64.11	69.47	73.44	77.36					
INW (MAR-APR)	GDC	S						1.83	3.02	3.35	3.16	3.47	3.39	3.32	3.17	2.50	1.99					
	N							6	14	37	64	65	72	79	81	85	28					
	DEV							+3.40	+1.45	-.12	-1.65	-2.95	-3.03	-2.93	-2.89	-3.51	-4.52					
	X							41.40	44.45	47.88	51.35	55.05	59.97	65.07	70.11	74.49	78.48					
	PGA	S						2.88	2.94	2.70	2.96	3.42	3.15	4.14	4.29	4.28	4.12					
	N							5	11	32	60	59	62	73	75	76	22					
	DEV																					
	X																					
INW (MAY-JUN)	GDC	S																				
	N																					
	DEV																					
	X																					
	PGA	S																				
	N																					
	DEV																					
	X																					
INW (MAY-JUN)	GDC	S																				
	N																					
	DEV																					
	X																					
	PGA	S																				
	N																					
	DEV																					
	X																					
INW (JUL-AUG)	GDC	S																				
	N																					
	DEV																					
	X																					
	PGA	S																				
	N																					
	DEV																					
	X																					
INW (SEP-OCT)	GDC	S																				
	N																					
	DEV																					
	X																					
	PGA	S																				
	N																					

- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R. F. Quiring, June 1977. (PB-271-704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB-271-742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB-272-661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB-273-155/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB-281-387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB-283-080/AS)
- 129 Fire Whirls. David W. Goens, May 1978. (PB-283-866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB-286-014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB-289-916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R. G. Pappas, R. Y. Lee, B. W. Finke, October 1978. (PB289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., Jan. 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott HeFlick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather District. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 145 On the Use of Solar Radiation and Temperature Models to Estimate the Snap Bean Maturity Date in the Willamette Valley. Earl M. Bates, August 1979. (PB80-160971)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80-155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80-160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80-174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80-220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80-196033)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins, June 1980. (PB80-225493) (Out of print.)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80-225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB81-108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. TenHarkel, July 1980. (PB81-106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81-102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochiai, August 1980. (PB81-108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81-134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81-133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81-205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81-224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J. C. Burnash and Larry Terra, April 1981. (PB81-224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. Lt. Colonel Keith R. Gnoy, April 1981. (PB81-225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81-23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82-230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82-127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981. (revised September 1981). (PB82-196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82-127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981.
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82-153883)
- 171 Verification of 72-Hour 500-mb Map-Type Predictions. R. F. Quiring, November 1981. (PB82-158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82-177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82-196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82-193103)
- 175 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1981. Emil B. Gunther and Staff, June 1982. (PB82-252420)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82-253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidli, April 1969 (revised March 1983).
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E. B. Gunther, June 1983.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS — Important definitive research results, major techniques, and special investigations.


CONTRACT AND GRANT REPORTS — Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS — Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS — Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS — Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS — Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

**ENVIRONMENTAL SCIENCE INFORMATION CENTER (D822)
ENVIRONMENTAL DATA AND INFORMATION SERVICE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE**

**6009 Executive Boulevard
Rockville, MD 20852**