



NOAA Technical Memorandum NWS WR-208

**METEOROLOGICAL FACTORS CONTRIBUTING TO THE
CANYON CREEK FIRE BLOWUP
SEPTEMBER 6 AND 7, 1988**

David W. Goens

**Weather Service Office
Missoula, Montana
June 1990**

**U.S. DEPARTMENT OF
COMMERCE**

/ National Oceanic and
Atmospheric Administration

/ National Weather
Service



NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P.O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidli, April 1968 (Revised March 1986). (PB88 177672/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB 179084)
- 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. (PB 184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 1969. (PB 185068)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)
- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 65 Climate of Sacramento, California. Tony Martini, April 1990. (Fifth Revision) (PB89 207781/AS)
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)

- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251)
- 83 A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS)
- 96 Map Type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Out-Of-Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuf, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 269 594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAIN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 115 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 116 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 126 Climate of San Francisco. E. Jan Null, February 1978. Revised by George T. Pericht, April 1988. (PB88 208624/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)

NOAA Technical Memorandum NWS WR-208


**METEOROLOGICAL FACTORS CONTRIBUTING TO THE
CANYON CREEK FIRE BLOWUP
SEPTEMBER 6 AND 7, 1988**

**David W. Goens
Weather Service Office
Missoula, Montana
June 1990**

UNITED STATES
DEPARTMENT OF COMMERCE
Robert A. Mosbacher, Secretary

National Oceanic and
Atmospheric Administration
*John A. Knauss, Under Secretary
and Administrator*

National Weather Service
*Elbert W. Friday, Jr., Assistant
Administrator for Weather Services*



This publication has been reviewed
and is approved for publication by
Scientific Services Division,
Western Region

Kenneth B. Mielke

Kenneth B. Mielke, Chief
Scientific Services Division
Salt Lake City, Utah

TABLE OF CONTENTS

| | PAGE |
|--|------|
| TABLE OF FIGURES | iv |
| I. INTRODUCTION | 1 |
| II. TOPOGRAPHY | 1 |
| III. CLIMATOLOGY AND LONG-RANGE FORECASTS | 1 |
| IV. SETTING THE SCENE - METEOROLOGICALLY | 5 |
| V. "BLACK TUESDAY AND ASH WEDNESDAY", SEPTEMBER 6 & 7, 1988 | 9 |
| VI. SUMMARY | 15 |
| VII. CONCLUSIONS | 18 |
| VIII. ACKNOWLEDGEMENTS | 20 |
| IX. REFERENCES CITED | 20 |

TABLE OF FIGURES

| | | |
|-----|--|----|
| 1. | CANYON CREEK FIRE GROWTH MAP | 2 |
| 2. | PALMER DROUGHT CHART, MAY 14, 1988 | 4 |
| 3. | SURFACE ANALYSIS, 1200 UTC, SEPTEMBER 1, 1988 | 6 |
| 4. | 500 MB CHART, 1200 UTC, SEPTEMBER 1, 1988 | 7 |
| 5. | FIRE PERIMETER, SEPTEMBER 5, 1988 | 8 |
| 6. | 850 MB CHART, 0000 UTC, SEPTEMBER 6, 1988 | 9 |
| 7. | SURFACE CHART, 0000 UTC, SEPTEMBER 7, 1988 | 10 |
| 8. | WIND PROFILE, SPOKANE, WASHINGTON, 0000 UTC, SEPTEMBER 7, 1988 | 11 |
| 9. | OBSERVED WINDS, MOISTURE, AND BENCHMARK RAWS SITES, SEPTEMBER 6 AND 7, 1988 | 12 |
| 10. | SURFACE ANALYSIS, 0600 UTC, SEPTEMBER 7, 1988 | 13 |
| 11. | SURFACE ANALYSIS, 0900 UTC, SEPTEMBER 7, 1988 | 14 |
| 12. | 500 MB CHART, 0000 UTC, SEPTEMBER 7, 1988 | 15 |
| 13. | 500 MB CHART, 1200 UTC, SEPTEMBER 7, 1988 | 16 |
| 14. | GREAT FALLS ATMOSPHERIC SOUNDING, 1200 UTC, SEPTEMBER 7, 1988 | 17 |
| 15. | SATELLITE PHOTOGRAPH, 1800 UTC, SEPTEMBER 7, 1988 | 18 |
| 16. | OBSERVED WIND, GREAT FALLS, MONTANA, SEPTEMBER 6-7, 1988 | 19 |
| 17. | OBSERVED WIND, MISSOULA, MONTANA, SEPTEMBER 6-7, 1988 | 19 |
| 18. | CANYON CREEK FIRE, FINAL FIRE SIZE | 21 |

METEOROLOGICAL FACTORS CONTRIBUTING TO THE
CANYON CREEK FIRE BLOWUP
SEPTEMBER 6 AND 7, 1988

I. INTRODUCTION

The Canyon Creek Fire was ignited by lightning on June 25, 1988, within the Scapegoat Wilderness Area of Montana's Lolo National Forest. The fire was initially allowed to burn unconstrained in accordance with guidelines set forth in the Scapegoat-Danaher Fire Management Plan (1982). The National Weather Service's Fire Weather Office in Missoula, Montana, provided weather information for the initial decision process, allowing the fire to play its natural role in the wilderness, and throughout the summer and fall for management decisions concerning this fire.

During the life of the Canyon Creek Fire, four major wind events resulted in rapid fire growth (See Fig. 1). The fire increased in size to 10,000 acres on July 22. Strong winds on August 9 drove the fire to 33,000 acres. On August 29, the fire increased to 51,200 acres as it was again fanned by gusty winds. The most significant run began on the evening of September 6. By daybreak on September 7, the fire had quadrupled in size and encompassed more than 240,000 acres. The firestorm had moved through 180,000 acres of private, state, and federal lands in one 16-hour period, mostly at night.

The drought of 1988 conditioned the wildlands of the northern Rockies, providing an environment conducive to large fire growth. Meteorological events that produce periodic strong winds occur fairly frequently during the summer in Montana. The weather systems that pushed the Canyon Creek Fire's first three major runs were not atypical. The last and most dramatic run occurred under an uncommon and unusual

combination of meteorological events. This paper will document those events.

II. TOPOGRAPHY

The Scapegoat Wilderness Area is mountainous and rugged with elevations ranging from around 4,500 feet to over 8,500 feet. The area is mostly timbered with various coniferous tree species. The Continental Divide cuts through the eastern third of the wilderness. The timberline is generally around 7,200 feet, with much of the Continental Divide above the timberline.

A portion of the Dearborn River drainage was unburned during the major fire run of September 6 & 7, providing a conspicuous void in the burn pattern (see Fig. 1). The Dearborn River is contained within a major drainage on the east slopes of the Continental Divide. This drainage is flanked to the south and west through much of its upper reaches by a predominant ridge line which is part of the Continental Divide. The unburned area was spared by a combination of sheltering effects, primarily by the high ridge line to the south, and the previously burned area up-wind that provided a partial fire break.

III. CLIMATOLOGY AND LONG-RANGE FORECASTS:

1. Climatology

A. Winds


Winds throughout Montana west of the Continental Divide are normally light





Mont.



Canyon Creek Fire Growth Map
Fig. 1

2



| | | | |
|----------|-------|-------|---|
| 7/22 | 4000 | ha |  |
| 8/9 | 13200 | ha |  |
| 8/29 | 20480 | ha |  |
| 9/6-7 | 96000 | ha |  |
| Cont Div | | 00000 | |

2

during the summer and early fall. The average wind speed in Missoula is 6.6 mph in August, and 6.0 mph in September. Along the east slopes of the Continental Divide winds are stronger throughout the year. Average wind speed in Great Falls is 10.3 mph during August, and 11.4 mph in September.

Normally, strongest surface winds occur during the afternoon hours from spring through summer and fall in the Northern Rockies. This is a result of a number of atmospheric processes, including solar induced convective forces, and turbulent mixing in the lower atmosphere. Winds usually decrease at night, even under unusual weather situations.

Whenever the synoptic scale circulation pattern provides discontinuities in the air masses, i.e., fronts and/or low-level jet streams, then very often the synoptic scale features will overpower local processes. The duration of strong wind events associated with frontal passages is typically in the range of a few hours before to a few hours after frontal passage. When systems slow down or become stationary, it is usually because of a weakening in the pressure gradient and/or an achieved balance in other dynamic forces, and winds are neither persistently strong nor gusty.

B. Precipitation and Lightning

The climate of the Scapegoat and contiguous Wilderness areas is characterized by cold, snowy winters, cool, wet springs, and warm, dry summers. Average precipitation in the area ranges from 40 to 60 inches per year (USDA-SCS, 1977). Snowfall is heavy, especially in areas along and west of the Continental Divide. Snowfall accumulation is distributed fairly uniformly from November through March. Rainfall in May and June provides about 25 percent of the annual precipitation.


The area west of the Continental Divide receives more moisture as it is primarily affected by weather systems of Pacific origin. Areas east of the Continental Divide are drier as much of the moisture from Pacific systems is precipitated out in the upslope flow on the west side.

Thunderstorms are common during the summer months (six to ten per month), with most storms being accompanied by some precipitation. Dry lightning, though not a common event, occurs most often during July and August. Lightning is responsible for an average of only three detectable fire starts annually within the Scapegoat-Danaher Fire Management Zone (Bailey, 1989).

C. Drought

The Palmer Drought Severity Index (PDSI) was developed to identify the onset, severity, and duration of a drought expressed as a departure of weather conditions from the norm (Palmer, 1965). It was initially derived to describe moisture conditions in the Great Plains, but has subsequently been applied to all regions of the United States. It is essentially a soil moisture accounting algorithm, calibrated to relatively homogenous hydrologic areas. The PDSI seems to provide an indication of potential fire severity as it indirectly reflects fine fuel moisture and the moisture content of large dead fuels.

The northern Rocky Mountains experienced drought conditions in 1987 and 1988. Mid-May (1988) PDSI charts (Fig. 2) indicated the northern Rocky Mountains were in severe to extreme drought. Drought conditions intensified by early July, and the moisture content of wildland fuels dried to unseasonably low levels.



Palmer Drought Chart, May 14, 1988

Fig. 2

Extended drought periods have an important impact on wildfire danger, and the occurrence and severity of large fires (Davis, 1959). The most significant effect may be the critical drying of large dead forest fuels. When the large dead fuels are extremely dry, their contribution may create fire intensity sufficient to cause soil damage, contribute to crown fires, and sustain fire during occasional moist periods. The moisture content in the smaller size class of fuels (less than one inch diameter) is important to the initiation and spread of a fire. During the long days of summer, lack of precipitation allows smaller fuels to dry to critical levels, thereby providing receptive fuel beds that easily kindle and sustain ignitions.

Normally, much of the western United States experiences a break in the warm, dry summer weather during the second or third week of August. This break occurs as low-pressure systems move southeast out of the Gulf of Alaska and across the Pacific Northwest and northern Rockies, bringing wetting rains and cooler temperatures. This break has been referred to as the "August Singularity" (Christopherson, 1980) and occurs with notable frequency. Climatological records for Missoula show it has occurred 23 times in the last 30 years. This weather event normally signals the end of the summer fire season. Fuels moistened by the wetting rains are prevented from drying to dangerous levels, primarily because of the shorter days (fewer hours of sunshine) of late summer. Many of the significant fire years in the northern

Rockies (1967, 1973, 1979, 1984, and 1988) occurred when this break in the weather did not develop until the last few days in August or early September. The break in 1988 did not come until after September 7.

2. Long-Range Forecasts

On July 11, a "Fire Analysis Team" comprised of U.S. Forest Service officials met to evaluate the risks and benefits of allowing the Canyon Creek Fire to continue in prescription status. The Fire Weather Unit at the National Weather Service's Missoula office provided long-range weather forecasts for both the 30-day and 90-day outlook periods for western Montana. These forecasts were based upon guidance prepared by the Climate Analysis Center Prediction Branch and were one of many inputs that contributed to the decision to allow the fire to continue unconstrained.

The 30-day outlook for July indicated near normal precipitation and near or a little below normal temperatures. The 90-day outlook for July through September indicated below normal temperatures and near or slightly above normal precipitation. There are many factors that are considered in the formulation of the extended-range forecasts, including climatology, sea surface temperatures, etc. (Harnack, 1986). Because the long-range forecasts were indicating a trend of cooler temperatures with more moisture, which is consistent with the normal August break in the weather, Missoula Fire Weather Forecasters felt comfortable advising the "Fire Analysis Team" based on this trend.

IV. SETTING THE SCENE - METEOROLOGICALLY

On August 29, a relatively dry Pacific front moved across Montana. West winds


in excess of 30 mph (reported by field observers) pushed the fire across a constructed containment line and out of the Scapegoat Wilderness Area. Aerial and infrared mapping on the 30th of August showed the fire had grown to over 51,000 acres. The fire had exceeded prescription parameters and was declared a WILDFIRE during the early morning hours.

By September 1, high pressure in the upper troposphere had developed over northern Alberta, Canada and extended south into Nevada. On the surface, high pressure was building over eastern Montana, with a thermally induced low-pressure area developing northward from California into Oregon (Fig. 3 and 4). This established an east-to-west pressure gradient force over western Montana.

The east winds induced by this pattern had two significant effects on the fire. First, the east side of the fire was being pushed back onto itself, thereby aiding suppression efforts. On the western sections of the fire, control problems escalated. The east winds pushed the fire into unburned islands within the fire line, and also spread fire to the west and south outside of previously established control lines.


The east winds persisted for three days, but on September 4, the weather pattern began to change. As the surface high pressure center moved east into the Dakotas, pressures began to fall along the lee slopes of the Continental Divide. The development of the "Lee-Side Trough" decreased the pressure gradient forces in the fire area. Winds were generally light and primarily slope-induced for nearly 24 hours.

The fire perimeter for the morning of September 5, (Fig. 5) shows the effects of three days of east winds, most significantly on the western portions of the fire. By mid-day on the 5th, west




Surface Analysis, 1200 UTC, September 1, 1988

Fig. 3



500 mb Chart, 1200 UTC, September 1, 1988

Fig. 4



Fire Perimeter, September 5, 1988


Fig. 5

winds began to increase on the western portion of the fire. East of the Continental Divide, winds were still out of the east, but were light.

By the evening of September 5, a complex pattern of weather events was evolving. A surface wave began to form from a low pressure center over northern Alberta. This can be seen on the 850 mb chart for 0000 UTC, September 6 (Fig. 6). The fire area was in the warm sector, and low-level (850 mb) southwesterly winds increased to over 30 kts during the night. The air mass became progressively more unstable as cool air advection aloft began to override the warm, dry air at the surface. These meteorological factors,

combined with physical factors of the fire environment, were developing into a "Blow-Up" situation for the Canyon Creek Fire. At 0500 UTC on September 6, (11:00 p.m. on September 5) a field observer on the fires west zone reported 9% relative humidity in a drainage bottom. Areas of unburned timber within the fire ignited, exhibiting extreme fire behavior throughout the evening and into the early morning hours of the 6th.

The weather pattern during the first five days of September was of extreme significance to the events that followed.



850 mb Analysis, 0000 UTC, September 6, 1988


Fig. 6

V. "BLACK TUESDAY AND ASH WEDNESDAY" -- SEPTEMBER 6 AND 7

By the morning of September 6, the fire was estimated at a little over 57,000 acres... a modest increase of around 6,000 acres in five days. A large percentage of this increased acreage was on the southwest side of the fire. Equally

significant, however, was the amount of area within the fire perimeter that was hot due to reburn during the three-day period of east winds and the wind reversal of September 5.

The surface weather chart for 0000 UTC on September 7, (6:00 p.m. on September 6) (Fig. 7) shows two frontal systems moving toward western Montana from the



Surface Chart, 0000 UTC, September 7, 1988


Fig. 7

northwest. The first was apparent only in the dew-point field; the second, a cooler continental push. In the upper levels, winds were quite strong and increasing. Winds measured on the 0000 UTC,

September 7, atmospheric sounding at Spokane, Washington (Fig. 8) indicated a low-level jet of nearly 45 knots near 7,000 ft. MSL. At 24,000 feet (400 mb), winds were 75 knots.

SPOKANE, WA

00 UTC 7 SEP 88



Wind Profile, Spokane, Washington, 0000 UTC, September 7, 1988

Fig. 8


On September 6, westerly surface winds increased on the fire's east zone during the mid-afternoon, and on the west zone by late afternoon. The trough, in the lee of the Rockies, moved east and pressure

gradient forces increased. Surface winds in portions of the east zone of the fire were reported as erratic, 20 to 30 mph, with gusts up to 50 mph.


The first front moved across the fire area during the evening hours of September 6. This front was difficult to track due to the lack of a good mesoscale data network, as well as the usual problems associated with frontal movement in mountainous terrain (Saucier, 1965). Reports from the Monture and Benchmark Remote Automatic Weather Stations (RAWS) and Silver King Lookout were consistent, all reporting strong west winds throughout the night. Winds averaged 15 to 20 mph with gusts between 40 and 60 mph (Fig. 9).

Characteristic air mass changes at the surface were not apparent with the first frontal passage. This is most likely due to active mixing caused by the wind, with contributions from the tremendous energy release in the lower atmosphere by the fire. During the 16-hour blowup, an average of 160 acres a minute were consumed. Temperatures at the RAWS sites remained above 60 F and humidity below 40 percent throughout the night. Closer to the fire, observers reported temperatures in the mid-70s (F) and humidity in the mid-teens at 0900 UTC on September 7 (3:00 a.m. MDT).

MONTURE RAWS, 9/6-7
AVG WND + GSTS




BNCHMRK RAWS, 9/6-7
AVG WND + GSTS



Observed Winds, Monture and Benchmark RAWS Sites
September 6 and 7, 1988

Fig. 9




Surface Analysis, 0600 UTC, September 7, 1988

Fig. 10

The surface analysis for 0600 UTC on September 7, (12:00 a.m. MDT) (Fig. 10) showed the second (polar) frontal system moving into northwestern Montana. By 1:00 a.m. MDT, ash began falling on the Great Falls, Montana airport, over 50 miles east of the fire. By 0900 UTC (3:00 a.m. MDT), the front had moved over the fire area (Fig. 11). Visibility was restricted at Great Falls to five miles by smoke and falling ash through the remainder of the night. Rapid pressure rises, along with a slight wind shift, reported in the 7:00 a.m. MDT surface weather observation signaled the weak frontal passage at Great Falls.

The 500 mb chart for 0000 UTC September 7, (Fig. 12) shows the polar jet stream moving south with a 50 knot speed maximum approaching northern Idaho. The 1200 UTC, September 7, 500 mb chart (Fig. 13) indicated a wind speed maximum of nearly 100 knots approaching the fire area.

The 1200 UTC (6:00 a.m. MDT) September 7 sounding from Great Falls (Fig. 14) shows a shear layer between 12,000 and 14,000 feet with winds increasing from 25 knots to 50 knots. Below this shear layer, a lower level wind maximum of 35 knots existed between



Surface Analysis, 0900 UTC, September 7, 1988


Fig. 11

8,000 and 9,000 feet, near mountaintop level. A subsidence inversion was just above the shear zone, and most likely associated with the right front quadrant of the jet stream (Reiter, 1963). This inversion was significant as it helped substantiate, on a much larger scale, the unusually warm and dry conditions that existed throughout the night over the fire.

The subsidence-type of inversion just above mountaintop level has also been identified as fundamental in the development of "Foehn" type strong

mountain downslope winds (Klemp and Lilly, 1976). The compressional warming associated with the strong downslope flow on the east slopes of the Continental Divide contributed to both the rapid rate of spread and intensity of the fire.

The secondary frontal boundary oriented on an east-west line remained nearly stationary over the fire area through midday on September 7. Strong westerly surface winds continued along the east slopes of the Continental Divide. Visible imagery from a polar-orbiting satellite (Fig. 15) showed the frontal boundary



500 mb Chart, 0000 UTC, September 7, 1988


Fig. 12


across the fire area and a spiral smoke plume downwind of the fire. The interesting spiral configuration of the smoke plume is indicative of a core of strong winds near the surface, i.e., a low-level jet stream. The satellite photograph showed fires further south in Idaho and in Yellowstone Park being influenced by the strong, westerly winds south of the polar front and jet core.

By 1200 UTC (6:00 a.m. MDT) on September 7, the fire had concluded its major run. Although the winds were still strong on the surface and aloft, the fuels along the eastern head of the fire were so sparse and discontinuous that further spread was limited (Morris, 1989).

VI. SUMMARY

The factors of fuels, topography, and weather all contributed to the Canyon Creek Fire's momentous spread on September 6 and 7. An in-depth evaluation of the role of topography and fuel supply on this fire is best left to other subject-matter experts. It is sufficient to say that fuels were available and ready to burn, both as a result of the age and class of trees, condition of the forest environment, and the precedent dry weather conditions. The complex topography played a significant role in the fire growth, and the final fire shape, size, and burn pattern.





Great Falls Atmospheric Sounding, 1200 UTC, September 7, 1988

Fig. 14


surface winds increased significantly. In addition, the jet provided a further contribution to fire activity through the warming (resulting from active low-level mixing) and the compression effects as the winds were forced over the terrain. The effects in the wind pattern can be seen in recorded data from nearby observation sites:

1. Great Falls (Fig. 16): The average wind speed of 19 mph on September 6, was nearly 175 percent of the monthly average of 11.3 mph. From midday on the 6th to midday on the 7th, the average wind speed was about twice the monthly average. This is especially noteworthy because of the persistence of the strong winds throughout the night.

2. Missoula (Fig. 17): The average wind speed of 10.2 mph (170 percent of normal) on September 6, was significant, especially when we note that winds were

less than 5 mph until after midday. The average wind speed for Missoula in September is only 6.0 mph. On September 7, the average wind speed was 16.7 mph, approaching nearly three times the monthly average. Again, this is even more meaningful because the strong winds persisted throughout the night.

Topography, both west and east of the Continental Divide, played an important role in the wind's influence on the fire. West of the Divide, the rough, broken mountainous terrain was a factor in increasing the mixing depth at night (due to terrain induced turbulence), bringing the stronger winds aloft to the surface. The winds were most likely increased by terrain forced convergence of the airflow. Finklin (1973) noted a similar behavior in the wind field at night as it effected the dramatic Sundance Fire run in North Idaho on September 1, 1967. The "downslope" effect east of the Divide is a



Satellite Photograph, 1800 UTC, September 7, 1988

Fig. 15

well-documented occurrence and was most likely amplified by the location and strength of a low-level jet stream.


VII. CONCLUSIONS

Wind-driven fires can be easily recognized by their characteristic shape (Anderson, 1983). It is obvious by a casual examination of the fire growth map (Fig. 18) for September 6 and 7, that the Canyon Creek fire was driven by the wind. Topographic discontinuities were factors in the unburned area within the

Dearborn River drainage on the east half of the fire.


The meteorological events that contributed to the explosive overnight growth of the fire were singularly significant and collectively unique. The four major events were:

1. Long-term drought unrelieved by usual late summer rains;
2. Three days of east winds in an area that has predominate west winds;



Observed Wind, Great Falls, Montana, September 6-7, 1988

Fig. 16



Observed Wind, Missoula, Montana, September 6-7, 1988

Fig. 17

3. The passage of two dry frontal systems within twelve hours accompanied by persistent strong west winds;

4. The occurrence of a low-level jet stream that moved over the fire area and remained stationary for over 12 hours.

The combination of these meteorological events imposed on an environment ripe with fuels, and in favorable topography, produced and enhanced the spectacular and devastating fire growth of the Canyon Creek Fire on September 6 and 7, 1988.

VIII. ACKNOWLEDGEMENTS

The author wishes to thank Mr. Jerry Williams of the Lolo National Forest for his patience and persistent encouragement. Mr. Byron Bonny (USFS, Clearwater NF), provided technical consultation through his direct involvement in the Canyon Creek Fire from ignition to extinction. A special debt of gratitude is owed to Mr. Hudson Garvin of WSO Missoula for his assistance in analyzing the surface charts. Ms. Brenda Graham, also of WSO Missoula, deserves credit for her review of this paper through various drafts. Brenda's comments and suggestions helped the author keep the "Horse and Cart" in proper alignment.

IX. REFERENCES CITED

Anderson, Hal E., 1983: Predicting Wind-Driven Wildland Fire Size and Shape. *USDA Forest Service Research Paper INT-305*. p. 26 Intermt. Forest and Range Exp. Stn., Ogden, Utah

Average Annual Precipitation for Montana 1941-1970, 1977. USDA Soil Conservation Service, Portland, Oregon.

Bailey, Dan, 1989: Personal Conversation, data extracted from the Fire Weather

Data Library, USDA Forest Service, Fort Collins, Colorado.

Christopherson, J.A., 1980. The August Singularity. Unpublished report of file, National Weather Service Office, Missoula, Montana.

Davis, K.P., 1959. *Forest Fire Control and Use*. pp. 97-98. McGraw Hill Book Co., New York, New York.

Finklin, A.I., 1973. Meteorological Factors in the Sundance Fire Run. *USDA Forest Service Gen. Tech. Rep. INT-6*, p. 46 Intermt. Forest and Range Exp. Stn., Ogden, Utah.

Harnack, R.P., 1986. *Principles and Methods of Extended Forecasting in the U.S.* p. 35. Meteorological Monographs, National Weather Association.

Klemp, J.B. and D.K. Lilly, 1976. The Dynamics of Wave Induced Downslope Winds, pp 320-329. *Journal of Atmospheric Sciences*, #32.

Morris, Ed, 1989. Personal Conversation. USDA Forest Service, Region 1. Deputy Incident Commander, Canyon Creek Fire.

Palmer, W.C., 1965. Meteorological Drought. *US Department of Commerce, Weather Bureau Research Paper No. 45*. p. 58. National Weather Service, Suitland, Maryland.

Reiter, E.R., 1963. *Jet Stream Meteorology*. p. 515. University of Chicago Press, Chicago, Illinois.

Saucier, W.J., 1955. *Principles of Meteorological Analysis*. p 433. University of Chicago Press, Chicago, Illinois

Scapegoat-Danaher Fire Management Plan, 1982. *USDA Forest Service Region-1*. p. 25., Lolo National Forest, Missoula, Montana.



Canyon Creek Fire, Final Fire Size

Fig. 18

- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins (Retired) and Alexander R. Smith. Fourth Revision, March 1989. (PB89 180624/AS)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82 158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82 177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidli, April 1969 (Revised December 1986). (PB87 142063/AS)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteries, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85 110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85 132694)
- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 1878887AS)
- 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lussky, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lussky, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Radid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
- 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
- 199 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Lussky, April 1987. (PB87 185229/AS)
- 200 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke, September 1987. (PB88 110895/AS)
- 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson and George Clark, September 1987. (PB88 114038/AS)
- 202 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88 101935/AS)
- 203 An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern California. John P. Monteverdi and Scott A. Braun, October 1988. (PB89 121297/AS)
- 204 Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Carven Scott, November 1988. (PB89 128649/AS)
- 205 Forecast Guidelines For Fire Weather and Forecasters - How Nighttime Humidity Affects Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS)
- 206 A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarters, Scientific Services Division, August 1989. (PB89 230833/AS)
- 207 The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, June 1990.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.


CONTRACT AND GRANT REPORTS--Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS--Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161