



**NOAA Technical Memorandum NWS WR-234**

---

**SOME CLIMATOLOGICAL AND SYNOPTIC ASPECTS  
OF SEVERE WEATHER DEVELOPMENT IN THE  
NORTHWESTERN UNITED STATES**

**Eric C. Evenson and Robert H. Johns  
National Severe Storms Forecast Center  
Kansas City, Missouri**

**October 1995**

---

**U.S. DEPARTMENT OF  
COMMERCE**

/ National Oceanic and  
Atmospheric Administration

/ National Weather  
Service



NOAA TECHNICAL MEMORANDA  
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P.O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Luciane Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidli, April 1968 (Revised March 1966). (PB86 177672/AS). (Revised October 1991 - PB92-115062/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belsky, July 1968. (PB 179084)
- 32 Probability Forecasting—A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento—Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. PB 184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 1969. (PB 185068)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)
- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donald W. Kuehl, August 1970. (PB 194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 65 Climate of Sacramento, California. Tony Martini, April 1990. (Fifth Revision) (PB89 207781/AS)
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)

- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972. (Revised March 1978). (COM 73 10251)
- 83 A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS)
- 96 Map Type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Anderson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Lefwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Lefwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 126 Climate of San Francisco. E. Jan Null, February 1978. Revised by George T. Pericht, April 1988. (PB88 208624/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudgeal, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Molner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summer-time Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)

**NOAA Technical Memorandum NWS WR-234**

**SOME CLIMATOLOGICAL AND SYNOPTIC ASPECTS  
OF SEVERE WEATHER DEVELOPMENT IN THE  
NORTHWESTERN UNITED STATES**


**Eric C. Evenson and Robert H. Johns  
National Severe Storms Forecast Center  
Kansas City, Missouri**

**October 1995**

*UNITED STATES  
DEPARTMENT OF COMMERCE  
Ronald H. Brown, Secretary*

*National Oceanic and  
Atmospheric Administration  
D. James Baker, Under Secretary  
and Administrator*

*National Weather Service  
Elbert W. Friday, Jr., Assistant  
Administrator for Weather Services*



This publication has been reviewed  
and is approved for publication by  
Scientific Services Division,  
Western Region



Delain A. Edman, Chief SSD  
Scientific Services Division  
Salt Lake City, Utah

## TABLE OF CONTENTS

---

|  |    |
|--|----|
| I. INTRODUCTION .....  | 1  |
| II. METHODOLOGY .....  | 1  |
| III. CLIMATOLOGY .....   | 2  |
| IV. SYNOPTIC AND THERMODYNAMIC CONDITIONS ASSOCIATED<br>WITH SSWEs ..... | 4  |
| V. CONCLUSION .....  | 9  |
| VI. REFERENCES .....   | 10 |

## TABLE OF FIGURES AND TABLE

---

- Figure 1: a) Area within solid, black lines denotes study area and is defined in this paper as the northwestern United States while b) shows important topographical features within the study area.
- Figure 2: Plot of all severe weather reports during each March through September period from 1955 through 1993. Triangles represent tornadoes while straight lines indicate tornado tracks. Dark circles indicate hail reports while cross symbol represents wind gusts or damage. Diamond shapes indicate hail and wind damage reported at the same location.
- Figure 3: Type and number of severe weather reports in the northwestern United States for the March-September period from 1955 through 1993.
- Figure 4: Severe weather reports in three 13 year intervals for the March-September period from 1955 through 1993.
- Figure 5: Monthly distribution of SSWEs during the period from 1955 through 1993.
- Figure 6: Mean composite chart at 0000 UTC for Pattern A SSWEs affecting Idaho, Montana, and Wyoming. Dotted line denotes 850 mb thermal ridge. Frontal boundary is position of 700 mb front. Long dashed lines labeled H7, H5, and H3 indicate trough axis positions at 700, 500, and 300 mb. Thin line with arrow indicates the jet axis at 500 mb while thick line with arrow represents the jet axis at 300 mb. Broad zigzag line shows an area of 500 and 300 mb diffluence while 500 and 300 mb ridge axes are denoted by long, north-south oriented narrow zigzag line.
- Figure 7: As in figure 6, except for Pattern A SSWEs affecting Oregon and Washington.
- Figure 8: As in figure 6, except for Pattern B SSWEs affecting Idaho, Montana, and Wyoming.
- Figure 9: Skew-T log p sounding for Spokane, Washington for August 1991 at a) 1200 UTC 5th, b) 0000 UTC 6th, c) 1200 UTC 6th, and d) 0000 UTC 7th.
- Figure 10: Average 500 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern A SSWEs in Idaho, Montana, and Wyoming.

- Figure 11: Average 850 mb temperatures in degrees Celsius at 0000 UTC during Pattern A SSWEs in Idaho, Montana, and Wyoming.
- Figure 12: Average 700 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern A SSWEs in Idaho, Montana, and Wyoming.
- Figure 13: Plot of all severe weather reports for the 24 hour period beginning at 1200 UTC 30 April 1987. Dark circles indicate hail reports while cross symbol represents wind gusts or damage.
- Figure 14: Composite chart for 0000 UTC 1 May 1987. Symbols same as used in figure 6.
- Figure 15: Upper air analyses at a) 850 mb b) 700 mb c) 500 mb d) 300 mb for 0000 UTC 1 May 1987.
- Figure 16: Average 500 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern B SSWEs in Idaho, Montana, and Wyoming.
- Figure 17: Average 700 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern B SSWEs in Idaho, Montana, and Wyoming.
- Figure 18: Average 850 mb temperatures in degrees Celsius at 0000 UTC during Pattern B SSWEs in Idaho, Montana, and Wyoming.
- Figure 19: Plot of all severe weather reports for the 24 hour period beginning at 1200 UTC 10 May 1989. Dark circles indicate hail reports while cross symbol represents wind gusts or damage. Diamond shapes indicate hail and wind damage reported at the same location.
- Figure 20: Composite chart for 0000 UTC 11 May 1989. Symbols used are same as in figure 6.
- Figure 21: Upper air analyses at a) 850 mb, b) 700 mb, c) 500 mb, and d) 300 mb for 0000 UTC 11 May 1989.
- Figure 22: Average 500 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern A SSWEs in Washington and Oregon.

Figure 23: Average 700 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern A SSWEs in Washington and Oregon.

Figure 24: Plot of all severe weather reports over Washington and Oregon for the 24 hour period beginning at 1200 UTC 6 August 1991. Dark circles indicate hail reports while cross symbol represents wind gusts or damage. Diamond shapes indicate hail and wind damage reported at the same location.

Figure 25: Composite chart for 0000 UTC 7 August 1991. Symbols used are same as in figure 6.

Figure 26: Upper air analyses at a) 850 mb, b) 700 mb, c) 500 mb, and d) 300 mb for 0000 UTC 7 August 1991.

Table 1: Significant Severe Weather Episodes



# SOME CLIMATOLOGICAL AND SYNOPTIC ASPECTS OF SEVERE WEATHER DEVELOPMENT IN THE NORTHWESTERN UNITED STATES

*Eric C. Evenson and Robert H. Johns*

## Abstract

*This study examines severe convective storms (those producing either 3/4 inch diameter or greater hail, or wind gusts equal to or greater than 50 knots, or tornadoes) occurring in that portion of the northwestern United States extending from Washington and Oregon through Idaho into the western portions of Montana and Wyoming. The climatology of severe weather events in this region reveals that the frequency of occurrence is greatest near the Continental Divide with the frequency progressively decreasing towards the Washington and Oregon coasts. Compared with other portions of the United States (e.g. the Great Plains), the frequency is relatively low. However, the data reveal that within the region, significant severe weather episodes (SSWEs) that are particularly destructive, and/or affect relatively large areas occur about twice a year. To aid forecasters in the anticipation of SSWEs in the northwestern United States, composite charts displaying common patterns and parameter values have been prepared.*

## I. INTRODUCTION

Severe convective storms<sup>1</sup> are not as common over the northwestern United States as in other parts of the country (e.g., the Great Plains region). However, severe convective storms do occur in this region, and in certain situations, can be quite widespread and destructive. Determining days when severe thunderstorms are a significant threat in this region is a challenge for operational meteorologists. Thus, a better understanding of the synoptic and thermodynamic conditions that are associated with significant episodes of severe weather in the

northwestern United States is needed to aid meteorologists in forecasting such events. This study examines the climatology of severe weather events occurring in the northwestern United States. Further, significant severe weather events are identified, and the associated meteorological conditions are examined to determine common synoptic patterns and parameter values.

## II. METHODOLOGY

For this study, the northwestern United States is defined as the following area: Washington, Oregon, Idaho, and the western portions of Montana and Wyoming (Fig. 1). The eastern boundary across Montana and Wyoming is roughly 50 to 100 miles east of the Continental Divide, and was selected to allow for those severe

---

<sup>1</sup> Severe convective storms are defined as those that produce hail 3/4 of an inch in diameter or greater, wind gusts of 50 knots or greater, or convectively induced wind damage, and/or tornadoes.

weather episodes that commence west of the Divide but produce some reports immediately to the east. In general, the eastern boundary was placed near the divide to delineate between different convective environments. For example, the eastern portions of Montana and Wyoming are typically affected by higher concentrations of low-level moisture, and the upslope flow lifting mechanism (Doswell 1980) is usually more clearly defined.

To aid in developing a severe weather climatology for the study region, both Storm Data and the SVRLOT data analysis program (Hart 1993) were utilized to collect and analyze severe weather events for the period from 1955 through 1993. Further, events from this dataset were systematically examined to identify significant severe weather episodes (SSWEs).

For this study an SSWE is defined as any of the following:

A) A severe weather episode where 10 or more severe weather events occur in the study area during a 24 hour period beginning at 1200 UTC.

B) A severe weather episode with 5 or more severe weather events in the study area during a 24 hour period beginning at 1200 UTC, including at least one tornado of F3 or greater intensity.

---

<sup>2</sup> This time period was chosen since 98 percent of all severe weather reported in the region occurred during these months. Only 52 reports (2 percent) of severe weather occurred during the fall and winter months from October through February.

C) A severe weather episode in which the Storm Data description suggests a widespread severe weather event had occurred in the study area even though the specific severe weather report criteria in either A or B are not met (e.g., a generalized entry indicating that numerous trees were blown down and/or large hail had occurred or over portions of several states).

Johns and Doswell (1992) have noted that composite analyses are useful in identifying basic meteorological patterns and parameters related to the development of severe local storms. Two types of composite charts have been constructed for use in this study. The parameter field composite chart displays the composite pattern of a single parameter field (e.g., 700 mb temperatures) for a number of cases. A more complex composite analysis is the "severe weather mean composite chart". By using symbols, this chart displays the mean position of multiple meteorological features at various levels of the atmosphere from a number of cases. This synthesis allows one to obtain a three-dimensional picture of the conditions associated with severe weather development.

### III. CLIMATOLOGY

#### a. All severe weather events

During the period March-September<sup>2</sup> from 1955 through 1993 over two thousand (2133) severe weather events were reported in the study area (Fig. 2). During this 39 year period, there were 817 reports of severe weather in Idaho and 766 in western Montana. The number of reported events is considerably less in Oregon (216 reports), western Wyoming (184 reports), and Washington (150 reports). The low number of reports in Wyoming can partly

be attributed to the fact that this is the smallest subdivision of the study area. Examination of the distribution of reports in Fig. 2 indicates that, in general, the frequency of severe weather decreases as one progresses westward from the Continental Divide to the Washington and Oregon coasts.

Local variations in report frequency on Fig. 2 suggests that the highest concentrations of reports are closely correlated with the highest density of population (McNulty 1981). For example, in Idaho the primary concentration is across the southern portion of the state in the Snake River valley (Fig. 1B). Other concentrated areas include the Lewiston and Coeur d'Alene areas of northern Idaho. Note there is a distinct minimum in frequency over central Idaho (Fig. 2), where population is limited due to the existence of the Sawtooth Mountains (Fig. 1B). Mountainous terrain exists in other areas of the study region and these areas also show a relative minimum of severe weather reports.

Three quarters of the total number of reports in the dataset occur during the summer months of June (27%), July (29%), and August (19%). A state by state breakdown of the type and number of severe weather reports in the northwestern United States for the March-September period from 1955 through 1993 is illustrated in Fig. 3. The most common type of event is convective wind gusts (or damage) which make up over half (53%) of the total reports. Large hail comprises 34% of all reports with tornadoes accounting for 13%. The frequency of large hail appears to decrease rapidly as one goes west from the Continental Divide region to the Pacific Ocean. Over one-half (57%) of all hail reports occurred over Montana and Wyoming with 28% over Idaho and only 15% over Washington and

Oregon. This frequency distribution is likely associated with greater amounts of moisture and instability that usually exist further east toward the Continental Divide during the warm season.

A comparison of the number of severe weather reports in three 13 year intervals (1955-1967, 1968-1980, and 1981-1993) for the March-September period is shown in Fig. 4. Note that over 50% of all reports in each state were reported in the latest 13 year period, 1981-1993. Wyoming showed the greatest increase with 68 percent of all severe weather events reported in the latest period.

#### b. Significant Severe Weather Episodes (SSWEs)

Using the criteria defined in Section 2, twenty seven SSWEs were identified during the 39 year period (Table 1). Of this total, 22 cases met criterion I while 3 episodes met criterion III. Only 2 episodes, which were associated with the Vancouver, Washington F3 tornado on April 5, 1972 (Hales 1994) and the Teton Wilderness F4 tornado in Wyoming on July 21, 1987 (Fujita 1989) met criterion II.

The dataset suggests that SSWEs are confined to the spring and summer months (April-September; see Fig. 5). Further, one-third of the SSWEs (9) were reported during the month of June. The distribution of SSWEs for the three 13 year periods discussed earlier shows that 1 SSWE was reported between 1955-1967, 5 SSWEs from 1968-1980, and 21 SSWEs from 1981-1993. Therefore, about eighty percent (78%) of the SSWEs were reported in the last 13 year period! The higher number of severe weather reports and resultant SSWEs during the latest 13 year period are likely the result of several factors including increased population, a heightened meteorological awareness from

the public, and improved National Weather Service (NWS) verification and storm reporting procedures (Hales, 1987). Given this evolution in improved reporting, it appears the data supports a frequency of SSWEs in the study area of close to two per year. With the improved detection capabilities of the WSR-88D radar currently being installed throughout the region, it is likely to be found that the actual frequency distribution is greater than two per year.

#### **IV. SYNOPTIC AND THERMODYNAMIC CONDITIONS ASSOCIATED WITH SSWEs**

Meteorological features associated with each of the individual SSWEs in the dataset were examined to determine common patterns. This examination suggests two common synoptic patterns based on mid- and upper-level trough orientation: Pattern A - the "negative tilt" pattern and Pattern B - the "trough axis" pattern. To better assess the relative locations of meteorological features associated with these patterns to the area of severe weather occurrence, the study region was divided into two subregions: 1) Idaho, western Montana, and western Wyoming (ID/MT/WY), and 2) Oregon and Washington (OR/WA). Pattern A is the most common synoptic pattern associated with SSWEs occurring in the northwestern United States (21 cases) and it has been observed in both subregions (Figs. 6 and 7). Pattern B is less frequent (5 cases) and it has been observed in the ID/MT/WY subregion only (Fig. 8). Meteorological features associated with one SSWE, the tornado episode of 5 April 1972, which occurred in the OR/WA subregion, fits neither Pattern A nor Pattern B. Further, it is the only case in the OR/WA subregion that affected areas west of the Cascade

Mountain range. This case is discussed further in Section 4b.

In the interior northwestern United States (i.e., east of the Cascade Mountains) during the warmer months of the year moisture values are usually low and inverted-V soundings are typically present (Beebe 1955, Barnes and Newton 1986). However, the air mass associated with SSWEs in this region is often modified with generally higher moisture values in the lower portions of the troposphere. Surface dew points are typically 45F or greater in the area of occurrence before the event.

In many of the SSWE cases, moisture values increase downward from the midlevels with time (e.g., Fig. 9). This moisture increase appears to result from both advection and mixing. Northward transport of moisture into the region typically occurs in the southerly to southwesterly flow ahead of the upper trough and is typically concentrated in the vicinity of the frontal band at 700 mb. At times, particularly later in the warm season, this moisture transport appears to be a northward extension of the southwest monsoon (Hales 1974). The northward transport is greatly facilitated by the strongly backed (southerly) flow ahead of a negatively tilted upper trough (Pattern A). This helps to explain why Pattern A is the most common upper air pattern associated with severe thunderstorm development in the northwestern United States.

Because of the high mountain ranges in the western United States, the northward transport of moisture into the region is most effective at midlevels. Once midlevel moisture has advected into the region, a significant contribution to low-level moisture appears to result from midlevel moisture mixing downward into the relatively deep mixed layer typical of the region. One of the authors (Evenson) and

forecasters in the region (personal communication) have also noted that precipitation from high based convection (cloud bases generally between 600 and 500 mb) commonly occurs in the area during the 24 hr period preceding an SSWE event. This precipitation is usually light (less than one-quarter in at the surface), but can serve to redistribute moisture into the lower portions of the troposphere, enhancing potential instability.

a. Idaho, western Montana, and western Wyoming

Examination of meteorological features associated with the 22 SSWEs affecting this subregion reveals that 17 cases are associated with Pattern A, or the "negative tilt" pattern. The remaining 5 cases are associated with Pattern B, or the "trough axis" pattern.

1) Pattern A (negative tilt pattern) in ID/MT/WY subregion

For the ID/MT/WY subregion cases, this pattern typically features a long-wave trough position in the eastern Pacific Ocean (not shown) with a mid-level ridge axis extending from central Saskatchewan southward into eastern Wyoming (Fig. 6). A negatively-tilted short-wave trough<sup>3</sup> extending from 700 mb up through 300 mb is lifting northeastward into the subregion from the west central United States. Significant 300 mb 12 hour height falls, as large as 80 to 100 meters, are associated with the short-wave trough. South to

<sup>3</sup> A negatively tilted trough is one whose axis is not meridionally oriented, but leans toward the west with increasing latitude (Bluestein 1992).

subregion ahead of the trough with pronounced diffluence indicated at both 500 and 300 mb levels. A 500 mb jet axis of 40 to 60 kt winds typically extends along a band from west central Nevada into west central Montana while a southwesterly flow aloft prevails across the jet axis of 60 kt or greater winds at 300 mb extends from western Nevada north northeastward into northern Idaho. In the nearly half of the cases the 300 mb wind maxima were between 80 and 100 kt.

The precursor 500 mb thermal field (at 1200 UTC) displays a warm axis across central Montana with a cold axis somewhere off the west coast (Fig. 10). Colder temperatures (less than -15C) are confined to an area from the central portions of Washington and Oregon westward. During the 12 h period between 1200 and 0000 UTC temperatures cool slightly (1 to 2C) over central and southern Idaho as a weak thermal trough axis moves northeastward into the area. However, during the same 12 h period temperatures warm slightly, on average, over western Montana and western Wyoming. This suggests that destabilization owing to cooling aloft is typically not a major factor with these cases.

Probably the most significant contributions to destabilization are 1) the increase in moisture values in the lower half of the troposphere from moisture transport, mixing, and precipitation moistening, and 2) diurnal heating. Surface dew points are typically in the 50 to 60F range ahead of the cold front with 45F being the lower limit. A relatively warm north northwest-south southeast oriented 850 mb thermal ridge (temperatures in the 18-20C range) is present ahead of the cold front (Fig. 11) and strong diurnal heating typically adds to destabilization in the warm sector. Surface based CAPE values are typically between 1000 and 2000 Jkg<sup>-1</sup> in the subregion at

0000 UTC, and in some instances values reach  $2500 \text{ Jkg}^{-1}$ . Surface based lifted index values (SBLIs) are relatively stable in the precursor soundings (at 1200 UTC), ranging from +2 to -2. However, by 0000 UTC instability has increased significantly. SBLI values by that time typically range from -3 to -6, with a few values reaching -8.

All of the 17 Pattern A cases in this subregion are associated with a cold front, with severe thunderstorm development typically occurring in the vicinity or ahead of the front. However, surface reflections of fronts in the western United States are usually ill-defined. These boundaries can be more clearly identified by examining the thermal pattern changes at the 850 mb, and especially the 700 mb levels (Williams 1972). The composite 850 and 700 mb temperature fields associated with Pattern A cases in ID/MT/WY subregion (Figs. 11 and 12) suggest that a cold front usually extends from northwestern Montana across central Idaho into northeastern Nevada at 0000 UTC (Fig. 6).

## 2) ID/MT/WY subregion Pattern A case study - 30 April 1987

On the afternoon of 30 April 1987 severe thunderstorms producing primarily damaging winds struck western and northern Idaho and northwest Montana (Fig. 13). Winds estimated between 80 and 100 mph in northwestern Montana caused extensive damage. Damaging winds were also reported across southern Idaho as well and into portions of western Wyoming.

The severe weather composite chart for the 30 April 1987 case (Fig. 14) appears to fit Pattern A relatively well. The locations of most features associated with this case are close to those in the mean severe weather composite chart for Pattern A in the ID/MT/WY subregion (Fig. 6). There are a few variations, however. For

example, in the 0000 UTC 1 May 1987 case the 500 and 300 mb jet axes downstream from the short-wave trough, are farther west than the mean and there is an easterly component to the flow (Fig. 15). Further, the short-wave trough is moving more northward than northeastward, in part because of the strong ridge over south central Canada. Despite these variations in jet structure and trough movement, large 12 h (1200 to 0000 UTC) 300 mb height falls are associated with the short-wave trough (from 90 meters at Boise, Idaho to 140 meters at Salem, Oregon). Further, an area of pronounced diffluence at the mid- and upper-levels is nearly coincident with the primary area of severe weather occurrence, a signal common to Pattern A cases.

Also typical of Pattern A, the 500 mb temperature changes over the subregion between 1200 and 0000 UTC in the 30 April 1987 case are relatively small. Temperatures remained constant or warmed slightly in Montana and Wyoming, with modest cooling taking place in southwestern Idaho (-2C at Boise). Diurnal heating contributed to destabilization, particularly over the western portions of Montana and Wyoming where afternoon temperatures reached upper 70s and lower 80s F.

## 3) Pattern B (trough axis pattern) in ID/MT/WY subregion

Pattern B or the "trough axis" pattern differs from Pattern A in that a positively tilted long wave trough (at 700, 500, and 300 mb) is propagating eastward across the northwestern states. Since there are only five Pattern B cases in the dataset, the composite charts of associated features (Figs. 8, 16, 17, and 18) should be used with caution. Generally, many of the features and their locations are similar to those in Pattern A. These include the

850 mb thermal axis, the 700 mb and 500 mb thermal patterns, the 500 mb wind speeds, the location of the cold front (at 700 mb), and the upper ridge axis over the northern Plains. Further, cooling aloft over the subregion between 1200 and 0000 UTC is minimal. The severe weather composite chart suggests that there is typically some upper level diffluence over the subregion, but it appears to be less pronounced than it is with Pattern A. Also, there may be a closed low associated with the mid- and upper-level trough in Pattern B cases.

Within the subregion, Pattern B cases are typically associated with weaker instability than Pattern A cases. CAPE and SBLI values at 0000 UTC in Pattern B cases range from 700 to 1500  $\text{Jkg}^{-1}$  and -2 to -5, respectively. The weaker instability associated with this pattern results largely from lower values of moisture than is the case with Pattern A cases. Surface dew points ahead of the cold front range from the mid 40s F to the lower 50s F.

#### 4) ID/MT/WY subregion Pattern B case study - 10 May 1989

Several clusters of severe thunderstorms producing damaging winds and isolated large hail affected portions of western Montana, southeastern Idaho, and western Wyoming on 10 May 1989 (Fig. 19). The storms became severe during the late morning hours, continued through the afternoon, and diminished during the evening. Wind gusts reaching 70 and 80 mph were reported in portions of western Montana and southeast Idaho, damaging buildings and felling trees and power poles. Winds blew the roof off a commercial building and hail accumulated to 5 inches in depth in the south central Idaho town of Jerome.

The mid- and upper-level flow pattern in this case is basically Pattern B with a positively tilted long wave trough moving eastward in the western United States (Figs. 20 and 21). However, it does vary from the Pattern B severe weather composite in that there are two distinct short-wave troughs that have come into phase. Consequently, there are two branches to the mid- and upper-level jets, and the severe thunderstorm development appears to occur near and to the left of the southern branch 300 mb jet. Any diffluence in the subregion appears to be weak. Further, mid- and upper-level wind speeds in the subregion appear to be weaker than the mean.

The low-level (850 mb) thermal ridge remained nearly stationary during the day and extended from southern Saskatchewan southward across eastern Montana into central Wyoming at 0000 UTC (Fig. 21A). A north-south trough axis at 700 mb, initially along the Oregon/Washington coast at 1200 UTC, shifted eastward into the intermountain region during the day and a strong thermal gradient representing the 700 mb front extended from northwestern Montana into southwestern Idaho at 0000 UTC (Fig. 21B). The thermal gradient and 12 h changes at 500 mb, however, were generally quite weak (Fig. 21C). The only significant cooling occurred at Great Falls, Montana where the 500 mb temperature fell 3C in the 12 h period ending at 0000 UTC.

#### b. Oregon and Washington

Examination of the meteorological features associated with the 5 SSWEs affecting this subregion reveals that 4 cases display Pattern A characteristics. The remaining case (5 April 1972) has some characteristics that fit neither Pattern A nor Pattern B. Because of this, it is discussed separately

later in this section and is not included in the composite.

1) Pattern A (negative tilt pattern) in OR/WA subregion

Generally, the Pattern A severe weather composite chart for the OR/WA subregion resembles that of the ID/MT/WY subregion, but with the features shifted westward (Fig. 7). The mid- and upper-level short-wave troughs are typically offshore at 1200 UTC and move northeastward into the Oregon/northern California region by 0000 UTC. Because of the sparsity of data offshore, satellite imagery is critical in assessing the orientation and timing of the short-wave trough in these instances.

As the short-wave trough moves into the subregion, temperature changes in the midlevels are typically not large. In the 12 h period ending at 0000 UTC, some cooling may occur over the western portions of both Washington and Oregon while minor warming is likely over eastern Washington (Figs. 22 and 23). Changes are typically less than 2C at the 500 mb level. The cooling is greater at 700 mb, averaging more than 2C along coastal sections of Washington and Oregon.

At 850 mb, a thermal ridge extends from Utah across Idaho and into southeast British Columbia. Temperature values at Boise, Idaho (BOI) are greater than 25C and at Spokane, Washington (GEG) greater than 18C. Maximum afternoon surface temperatures are typically in the 80s and lower 90s F. Surface based lifted index values of -4 or less and CAPE values of 1000 and 2000 Jkg<sub>1</sub> are common in the area of severe weather occurrence.

2) OR/WA subregion Pattern A case study - 6 August 1991

Severe thunderstorms producing damaging winds and large hail developed in north central Oregon just east of the Cascade Mountains early on the afternoon of 6 August 1991 (Fig. 24). This activity developed northward into central Washington by late afternoon and spread northeastward over eastern Washington and portions of northeast Oregon during the late afternoon and evening hours. Convective gusts as high as 70 to 90 mph and golfball sized hail were reported at several locations. A gust to 100 mph was recorded at the Umatilla Army Depot in northeast Oregon. Five people were injured when a tent collapsed at the Umatilla County Fair in Hermiston, Oregon and a trailer was picked up and turned upside down by thunderstorm winds in Omak, Washington.

The composite chart for the 6 August 1991 case (Fig. 25) resembles the OR/WA Pattern A mean composite chart (Fig. 7). The primary differences are that in the 6 August case, the short-wave trough is a bit farther west and the upper ridge is farther east than on the mean composite.

Other characteristics of the 6 August 1991 case include, relatively dry warm axes at 850 and 700 mb that extended northward across Idaho and northwestern Montana at 0000 UTC 7 August (Figs. 26A and B). A moist axis can be noted along the 700 mb thermal gradient from central Washington southward into northwestern California. A moist axis is also noted at 850 mb, but the axis appears to lag behind (west) the 850 mb cold front. Surface temperatures during the afternoon reached the mid-80s to lower 90s F over much of Washington and Oregon east of the Cascades. Surface dew points were greater than 50F from east central Oregon northward. Some early morning precipitation may have contributed to higher low-level moisture values in portions of the region. Surface



based lifted index values reached -4 or less and surface based CAPE values averaged around 1000 to 1500  $Jkg_1$  at 0000 UTC. These values are somewhat lower than was found with the other case studies, but were still sufficient for SSWE development.

At the 500 and 300 mb levels (Figs. 26C and D), relatively strong southerly winds prevail. An area of diffluence is noted over the eastern portions of Washington and Oregon. Cooling aloft was minimal and confined to areas west of the Cascade mountains.

3) The unusual OR/WA subregion case of 5 April 1972

About midday on April 5 1972 severe weather developed in northwest Oregon and extreme southwestern Washington west of the Cascades and spread northeastward during the afternoon hours into eastern Washington before ending. Five tornadoes were reported, including two of F3 intensity and two of F2 intensity. One of the F3 intensity tornadoes killed 6 people and injured 300 others in Vancouver, Washington.

Although this case was associated with southwesterly flow aloft and a negatively tilted short-wave trough (Pattern A characteristics), other features differ significantly from those typically found in Pattern A cases. The episode commenced on the west side of the Cascades and was associated with a surface trough, well behind the cold front. Further, temperatures aloft were quite cold when compared to the other cases and 500 mb temperatures were generally less than -20C. These conditions closely resemble those that Hales (1985) found to be associated with cool season tornado development in coastal portions of southern California. For additional details concerning the meteorology and evolution

of this case the reader is referred to Hales (1994).

## V. CONCLUSION

The results of this study have shown that severe weather occurrence in the northwestern United States is generally restricted to spring and summer (98% of events reported during March through September) and is most common during the summer months of June through August (75% of all reports). Generally the frequency of occurrence decreases as one goes west from the continental divide to the Pacific coast.

The most common type of event reported is damaging convective winds and many of these events are likely associated with isolated microbursts and downbursts that are common in the western United States during the warm season. On occasion, however, when the wind fields are relatively strong and instability and forcing are sufficient, organized deep convection develops over the northwestern United States and more widespread severe weather results. In recent years these more widespread and significant severe weather episodes (SSWEs) have been reported about twice a year.

To assist forecasters in anticipating SSWE development, mean composite charts of those pertinent meteorological features and parameter values associated with SSWEs occurring within that portion of the region east of the Cascades were prepared. Although the number of cases (26) available to construct these charts was limited, several common features appear to stand out. All of the cases are associated with a long wave trough to the west of the area of occurrence and south to southwesterly flow at mid- and upper-levels prevails over the area of occurrence. Also,

all cases are associated with a short-wave trough moving into the region.

In most cases the short-wave trough is negatively tilted (80% of the cases) which appears to facilitate moisture transport into the region. This is probably a primary reason why those cases (20%) that are associated with a positively tilted short-wave troughs are restricted to the eastern portions of the region where moisture was more readily available.

Other conditions common to the composite cases are the relatively strong winds associated with the mid- and upper-level jets entering the region (40 to 60 kt at 500 mb and 50 to 100 kt at 300 mb), and the lack of any strong cooling above the region at midlevels (500 mb). Severe weather development is typically associated with a diffluent zone aloft and usually takes place along to well ahead of the boundary layer cold front. Because of mountainous terrain, the boundary layer cold front is usually most easily identified by examining the 700 mb thermal field. The front is typically located near or just east of the leading edge of the stronger thermal gradient at 700 mb.

Instability typically reaches moderate values in Pattern A (negatively tilted trough) cases with surface based lifted index values of -3 to -6 and CAPE values of 1000 to 2000  $\text{Jkg}^{-1}$ . However, in a few cases lifted index values may reach -8 and CAPE values may reach 2500  $\text{Jkg}^{-1}$ . In Pattern B cases instability is typically weak to moderate with surface based lifted index values of -2 to -5 and CAPE values of 700 to 1500  $\text{Jkg}^{-1}$ . This degree of instability is achieved in part through diurnal heating and moisture transport. Moisture is most efficiently transported into the region by the mid-level flow and likely mixes downward through precipitation, evaporation, and other means.

This study demonstrates that there are some common meteorological characteristics associated with most significant severe weather episodes occurring in the northwestern United States. Further, recent trends in severe weather reporting for the area suggests that such episodes occur on average about twice a year or more. With continued trends in increased awareness in the region and with vastly increased capabilities for detection with the installation of the WSR-88D radar network, it may be found that SSWEs in the region are even more common than current statistics would indicate. For this reason it is important for forecasters to have effective tools to aid in forecasting these larger scale events. It is hoped that this study will lead to improved forecast techniques concerning SSWE development in the northwestern United States.

## VI. REFERENCES

- Barnes, S. L., and C. W. Newton, 1985: *Thunderstorms in the synoptic setting. Thunderstorms: A Social, Scientific, and Technological Documentary*. Vol. 2: Thunderstorm morphology and Dynamics. (E. Kessler, Ed.), p. 75-111.
- Beebe, R. G., 1955: Types of airmasses in which tornadoes occur. *Bull. Amer. Meteor. Soc.*, **36**, 349-350.
- Bluestein, H. B., 1992: *Synoptic-Dynamic Meteorology in Midlatitudes: Volume 1 - Principles of Kinematics and Dynamics*. Oxford University Press, Inc. 431 pp.
- Doswell, C. A. III, 1980: Synoptic-scale environments associated with High Plains severe thunderstorms. *Bull. Amer. Meteor. Soc.*, **61**, 1388-1400.

Fujita, T. T., 1989: The Teton-Yellowstone Tornado of 21 July 1987. *Mon. Wea. Rev.*, *Amer. Meteor. Soc.*, **117**, **9**, 1913-1940.

Hales, J. E., Jr., 1974: Southwestern United States summer monsoon source - Gulf of Mexico or Pacific Ocean. *J. Appl. Meteor.*, **12**, 331-342.

Hales, J. E., Jr., 1985: Synoptic features associated with Los Angeles tornado occurrences. *Bull. Amer. Meteor. Soc.*, **66**, 657-662.

Hales, J. E., Jr., 1987: An examination of the National Weather Service severe local storm warning program and proposed improvements. NOAA Technical Memorandum NWSS NSSFC-15, U.S. Department of Commerce.

Hales, J. E., Jr., 1994: The Vancouver WA tornado on April 5, 1972 as a benchmark for west coast significant tornado episodes. Preprints, 6th Conference on Mesoscale Processes, Portland, Oregon, *Amer. Meteor. Soc.*, 134-137.

Hart, J. A., 1993: SVR PLOT: A new method of accessing and manipulating the NSSFC severe weather database. Preprints, 17th Conference on Severe Local Storms, St. Louis, Missouri, *Amer. Meteor. Soc.*, 40-41.

Johns, R. H., and C. A. Doswell III, 1992: Severe local storms forecasting. *Weather and Forecasting*, *Amer. Meteor. Soc.*, **7**, **4**, 588-612.

McNulty, R. P., 1981: Tornadoes west of the divide: A climatology. *Natl. Wea. Dig.*, **6**, **2**, 26-30.


Palmen, E., and Newton, C.W., 1969: *Atmospheric Circulation Systems*. Academic Press, New York. 603 pp.

Williams, P. Jr., 1972: Western region synoptic analysis - problems and methods. NOAA Technical Memorandum NWS WR-71, 71 pp.1. Introduction


**Table 1**  
**Significant Severe Weather Episodes**

| No. | Date    | States      | Reports<br>T/W/A | D/I   | Significant Events  |
|-----|---------|-------------|------------------|-------|---|
| 1   | 9/4/60  | OR/ID/MT    | -                | 0/0   | Widespread wind damage  |
| 2   | 6/27/70 | ID/MT/WY    | 0/16/1           | 0/0   | 80-100 mph winds in Montana                                     |
| 3   | 4/5/72  | WA/OR       | 5/0/0            | 6/301 | Two F3 tornadoes in Washington                                  |
| 4   | 6/22/73 | ID/MT       | -                | 0/0   | Widespread wind damage  |
| 5   | 6/23/75 | ID/MT       | 1/-/-            | 0/1   | 1 F2 tornado and wind damage                                    |
| 6   | 7/14/75 | WA/ID       | 0/3/11           | 0/0   | 2-3" diameter hail near Reubens, Id                             |
| 7   | 6/28/82 | ID/MT       | 1/4/6            | 0/0   | 30 million dollars damage in Helena, MT due to 3" diameter hail |
| 8   | 8/9/82  | WA/OR       | 1/9/2            | 0/0   | Winds estimated to 100 mph in NE OR                             |
| 9   | 8/11/82 | ID/MT       | 0/8/8            | 0/0   |   |
| 10  | 7/6/83  | ID/MT/WY    | 0/14/1           | 1/2   | 80-100 mph winds reported in ID, MT, and WY                     |
| 11  | 7/9/83  | ID          | 0/7/7            | 0/0   |   |
| 12  | 8/24/84 | ID/MT       | 1/8/6            | 0/0   |   |
| 13  | 4/30/87 | ID/MT/WY    | 0/16/2           | 0/0   | 80-100 mph winds in western MT                                  |
| 14  | 6/15/87 | ID/MT/WY    | 1/27/2           | 0/6   | \$10 million worth of damage in Nampa, ID                       |
| 15  | 7/21/87 | OR/ID/WY    | 2/5/2            | 0/0   | F4 tornado in the Teton Wilderness                              |
| 16  | 6/28/88 | MT          | 0/0/13           | 0/0   | Baseball size hail in western MT                                |
| 17  | 5/10/89 | ID/MT/WY    | 0/11/1           | 0/0   |   |
| 18  | 9/17/89 | ID/MT       | 1/13/0           | 0/4   | Extensive wind damage in SE ID                                  |
| 19  | 8/20/90 | WA/OR/ID/MT | 0/5/7            | 0/0   |   |
| 20  | 8/6/91  | WA/OR       | 0/9/7            | 0/5   | 80-100 mph winds in NE OR with 2" diameter hail                 |
| 21  | 9/10/91 | ID/WY       | 1/3/6            | 0/0   |   |
| 22  | 4/17/92 | ID          | 0/24/1           | 0/0   |   |
| 23  | 6/12/92 | ID/MT       | 1/9/2            | 0/0   |   |
| 24  | 6/28/92 | WA/OR/ID    | 0/10/3           | 0/0   |   |
| 25  | 7/22/92 | OR/ID/MT    | 2/6/3            | 0/0   |   |
| 26  | 5/3/93  | OR/ID       | 1/10/0           | 0/4   | Extensive wind damage in SE ID                                  |
| 27  | 6/21/93 | OR/ID/MT    | 3/11/0           | 0/1   | 97 mph winds at Choteau, MT                                     |

Note - Reports column denotes number of tornadoes/wind/hail events while D/I column denotes number of deaths/injuries.



A



B

Figure 1: a) area within solid, black lines denotes study area and is defined in this paper as the northwestern United States while b) shows important topographical features within the study area




Figure 2: Plot of all severe weather reports during each March through September period from 1955 through 1993. Triangles represent tornadoes while straight lines indicate tornado tracks. Dark circles indicate hail reports while cross symbol represents wind gusts or damage. Diamond shapes indicate hail and wind damage reported at the same location.

# Severe Weather in the Northwest U.S. 1955-1993




Figure 3: Type and number of severe weather reports in the northwestern United States for the March-September period from 1955 through 1993.

# Northwest U.S. Severe Weather Reports 13 Year Intervals




Figure 4: Severe weather reports in three 13 year intervals for the March-September period from 1955 through 1993.



# NW U.S. SSWE Monthly Climatology

## 27 Episodes




Figure 5: Monthly distribution of SSWEs during the period from 1955 through 1993.




Figure 6: Mean composite chart at 0000 UTC for Pattern A SSWEs affecting Idaho, Montana, and Wyoming. Dotted line denotes 850 mb thermal ridge. Frontal boundary is position of 700 mb front. Long dashed lines labeled H7, H5, and H3 indicate trough axis positions at 700, 500, and 300 mb. Thin line with arrow indicates the jet axis at 500 mb while thick line with arrow represents the jet axis at 300 mb. Broad zigzag line shows an area of 500 and 300 mb diffluence while 500 and 300 mb ridge axes are denoted by long, north-south oriented narrow zigzag line.




Figure 7: As in figure 6, except for Pattern A SSWEs affecting Oregon and Washington.







Figure 8: As in figure 6, except for Pattern B SSWEs affecting Idaho, Montana, and Wyoming.




A



B



C



D

Figure 9: Skew-T log p sounding for Spokane, Washington for August 1991 at a) 1200 UTC 5th, b) 0000 UTC 6th, c) 1200 UTC 6th, and d) 0000 UTC 7th.




Figure 10: Average 500 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern A SSWEs in Idaho, Montana, and Wyoming.




Figure 11: Average 850 mb temperatures in degrees Celsius at 0000 UTC during Pattern A SSWEs in Idaho, Montana, and Wyoming.




Figure 12: Average 700 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern A SSWEs in Idaho, Montana, and Wyoming.






Figure 13: Plot of all severe weather reports for the 24 hour period beginning at 1200 UTC 30 April 1987. Dark circles indicate hail reports while cross symbol represents wind gusts or damage.







Figure 14: Composite chart for 0000 UTC 1 May 1987. Symbols same as used in figure 6.




A



B



C



D

Figure 15: Upper air analyses at a) 850 mb b) 700 mb c) 500 mb d) 300 mb for 0000 UTC 1 May 1987.




Figure 16: Average 500 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern B SSWEs in Idaho, Montana, and Wyoming.




Figure 17: Average 700 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern B SSWEs in Idaho, Montana, and Wyoming.




Figure 18: Average 850 mb temperatures in degrees Celsius at 0000 UTC during Pattern B SSWEs in Idaho, Montana, and Wyoming.




Figure 19: Plot of all severe weather reports for the 24 hour period beginning at 1200 UTC 10 May 1989. Dark circles indicate hail reports while cross symbol represents wind gusts or damage. Diamond shapes indicate hail and wind damage reported at the same location.






Figure 20: Composite chart for 0000 UTC 11 May 1989. Symbols used are same as in figure 6.





A






Figure 22: Average 500 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern A SSWEs in Washington and Oregon.





Figure 23: Average 700 mb temperatures in degrees Celsius at 1200 UTC (dashed) and 0000 UTC (solid) during Pattern A SSWEs in Washington and Oregon.



0600CST 08/06/91 - 0600CST 08/07/91

SEVERE WEATHER REPORTS

Figure 24: Plot of all severe weather reports over Washington and Oregon for the 24 hour period beginning at 1200 UTC 6 August 1991. Dark circles indicate hail reports while cross symbol represents wind gusts or damage. Diamond shapes indicate hail and wind damage reported at the same location.







Figure 25: Composite chart for 0000 UTC 7 August 1991. Symbols used are same as in figure 6.




A



B



C



D

Figure 26: Upper air analyses at a) 850 mb, b) 700 mb, c) 500 mb, and d) 300 mb for 0000 UTC 7 August 1991.

- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298999/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins (Retired) and Alexander R. Smith, Fifth Revision, July 1992. (PB92 220177)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leitwich and Gail M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leitwich, Jr., October 1981. (PB82195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82 158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82 177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 190664)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidli, April 1969 (Revised December 1986). (PB87 142063/AS)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249004)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteris, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85 110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85 132694)
- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 187887AS)
- 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lussky, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lussky, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Radii Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
- 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
- 199 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Lussky, April 1987. (PB87 185229/AS)
- 200 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke, September 1987. (PB88 110895/AS)
- 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson and George Clark, September 1987. (PB88 114038/AS)
- 202 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88 101935/AS)
- 203 An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern California. John P. Monteverdi and Scott A. Braun, October 1988. (PB89 121297/AS)
- 204 Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Carven Scott, November 1988. (PB89 128649/AS)
- 205 Forecast Guidelines For Fire Weather and Forecasters - How Nighttime Humidity Affects Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS)
- 206 A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarters, Scientific Services Division, August 1989. (PB89 230833/AS)
- 207 The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, June 1990. (PB90-240268)
- 208 Meteorological Factors Contributing to the Canyon Creek Fire Blowup, September 6 and 7, 1988. David W. Goens, June 1990. (PB90-246085)
- 209 Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow Whitlatch, December 1990. (PB91-129239)
- 210 Hydrotools. Tom Egger, January 1991. (PB91-151787/AS)
- 211 A Northern Utah Soaker. Mark E. Struthwolf, February 1991. (PB91-168716)
- 212 Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 1991. (PB91-208439)
- 213 Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Mollner, July 1991. (PB91-227405/AS)
- 214 Emergency Operational Meteorological Considerations During an Accidental Release of Hazardous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424)
- 215 Weather-Tools. Tom Egger, October 1991. (PB93-184950)
- 216 Creating MOS Equations for RAWS Stations Using Digital Model Data. Dennis D. Gettman, December 1991. (PB92-131473/AS)
- 217 Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 1992. (PB92-196104)
- 218 NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 1992. (PB93-146785)
- 219 A Case Study of the Operational Usefulness of the Sharp Workstation in Forecasting a Mesocyclone-Induced Cold Sector Tornado Event in California. John P. Monteverdi, March 1993. (PB93-178697)
- 220 Climate of Pendleton, Oregon. Claudia Bell, August 1993. (PB93-227536)
- 221 Utilization of the Bulk Richardson Number, Helicity and Sounding Modification in the Assessment of the Severe Convective Storms of 3 August 1992. Eric C. Evenson, September 1993. (PB94-131943)
- 222 Convective and Rotational Parameters Associated with Three Tornado Episodes in Northern and Central California. John P. Monteverdi and John Quadros, September 1993. (PB94-131943)
- 223 Climate of San Luis Obispo, California. Gary Ryan, February 1994. (PB94-162062)
- 224 Climate of Wenatchee, Washington. Michael W. McFarland, Roger G. Buckman, and Gregory E. Matzen, March 1994. (PB94-164308)
- 225 Climate of Santa Barbara, California. Gary Ryan, December 1994. (PB95-173720)
- 226 Climate of Yakima, Washington. Greg DeVoir, David Hogan, and Jay Neher, December 1994. (PB95-173688)
- 227 Climate of Kalispell, Montana. Chris Maier, December 1994. (PB95-169488)
- 228 Forecasting Minimum Temperatures in the Santa Maria Agricultural District. Wilfred Pi and Peter Felsch, December 1994. (PB95-171088)
- 229 The 10 February 1994 Oroville Tornado-A Case Study. Mike Staudenmaier, Jr., April 1995.
- 230 Santa Ana Winds and the Fire Outbreak of Fall 1993. Ivory Small, June 1995.
- 231 Washington State Tornadoes. Tresté Huse, July 1995.
- 232 Fog Climatology at Spokane, Washington. Paul Frisbie, July 1995.
- 233 Storm Relative Isentropic Motion Associated with Cold Fronts in Northern Utah. Kevin B. Baker, Kathleen A. Hadley, and Lawrence B. Dunn, July 1995

## NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

*The National Oceanic and Atmospheric Administration* was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

**PROFESSIONAL PAPERS**--Important definitive research results, major techniques, and special investigations.


**CONTRACT AND GRANT REPORTS**--Reports prepared by contractors or grantees under NOAA sponsorship.

**ATLAS**--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

**TECHNICAL SERVICE PUBLICATIONS**--Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

**TECHNICAL REPORTS**--Journal quality with extensive details, mathematical developments, or data listings.

**TECHNICAL MEMORANDUMS**--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

**NATIONAL TECHNICAL INFORMATION SERVICE**

**U. S. DEPARTMENT OF COMMERCE**

**5285 PORT ROYAL ROAD**

**SPRINGFIELD, VA 22161**