



**NOAA TECHNICAL MEMORANDUM
NWS WR-238**

**SIGNIFICANT WEATHER PATTERNS AFFECTING
WEST-CENTRAL MONTANA**

**Joe Lester
Dave Bernhardt
Ken Mielke**

**Nexrad Weather Service Forecast Office
Great Falls, MT**

May 1996

**U.S. DEPARTMENT OF
COMMERCE**

National Oceanic and
Atmospheric Administration

National Weather
Service



NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, 125 South State Street - Rm 1210, Salt Lake City, Utah 84138-1102. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-PP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17900)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-175240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidl, April 1968 (Revised March 1986). (PB86 177672/AS). (Revised October 1991 - PB92-115062/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB 179084)
- 32 Probability Forecasting—A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento—Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. PB 184296
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 1969. (PB 185068)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)
- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 65 Climate of Sacramento, California. Tony Martini, April 1990. (Fifth Revision) (PB89 207781/AS)
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)

- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)
- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251)
- 83 A Comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11284)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
- 91 Arizona "Eddy" Tomadoes. Robert S. Ingram, October 1973. (COM 73 10466)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS)
- 96 Map type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 126 Climate of San Francisco. E. Jan Null, February 1978. Revised by George T. Pericht, April 1988. (PB88 208624/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudgel, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Molner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 28977/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)

**NOAA TECHNICAL MEMORANDUM
NWS WR-238**

**SIGNIFICANT WEATHER PATTERNS AFFECTING
WEST-CENTRAL MONTANA**

**Joe Lester
Dave Bernhardt
Ken Mielke**


**Nexrad Weather Service Forecast Office
Great Falls, MT**

May 1996


UNITED STATES
DEPARTMENT OF COMMERCE
Mickey Kantor, Secretary

National Oceanic and
Atmospheric Administration
D. James Baker, Under Secretary
and Administrator

National Weather Service
Elbert W. Friday, Jr., Assistant
Administrator for Weather Services



This publication has been reviewed
and is approved for publication by
Scientific Services Division,
Western Region



Delain A. Edman, Chief
Scientific Services Division
Salt Lake City, Utah

TABLE OF CONTENTS

I.	INTRODUCTION	1
II.	DATA	1
III.	METHODOLOGY	1
IV.	HEAVY RAIN	2
V.	HEAVY SNOW	2
VI.	STRONG WINDS	4
VII.	SEVERE WEATHER	5
VIII.	CONCLUSIONS	5
IX.	REFERENCES	5

FIGURES

FIGURE 1	500mb Heights for Heavy Rain of 1.50 Inches or more	7
FIGURE 2	Sea Level Pressures for Heavy Rain of 1.50 Inches or more	8
FIGURE 3	250mb Winds for Heavy Rain of 1.50 Inches or more	9
FIGURE 4	500mb Heights and 250mb Winds for Heavy Rain of Two Inches or more	10
FIGURE 5	Sea Level Pressures and 500mb Heights for Case 1 of Heavy Snow of Six Inches or more	12
FIGURE	Sea Level Pressures, 500mb Heights, and 850mb Heights/Temps for Case 2 of Heavy Snow of Six Inches or more	13
FIGURE 7	Sea Level Pressures, 500mb Heights, and 850mb Heights/Temps for Case 3 of Heavy Snow of Six Inches or more	14
FIGURE 8	Sea Level Pressures, 500mb Heights, and 250mb Winds for Case 2 of Heavy Spring Snow of Eight Inches or more	15
FIGURE 9	Sea Level Pressures, 850mb Heights/Temps, and 500mb Heights for Strong Sustained Winds of 30mph or greater in Winter	17
FIGURE 10	Sea Level Pressures, 850mb Heights/Temps, 700mb Heights/Temps, and 500mb Heights for Strong Sustained Winds of 25mph or greater in Summer	19
FIGURE 11	500mb Heights, Sea Level Pressures, 850mb Heights/Temps, and 250mb Winds for Severe Summer Weather	21

TABLES

TABLE 1	Great Falls Precipitation of 1.50 Inches or more in 24 Hours (1949-1989)	6
TABLE 2	Great Falls Snowfalls of Six Inches or more in 24 Hours (1950-1989)	11
TABLE 3	Great Falls Mean Daily Winds of 30 mph or greater from November through March (1963-1989).....	16
TABLE 4	Great Falls Mean Daily Winds of 25 mph or greater from May through September (1950-1989)	18
TABLE 5	Great Falls Thunderstorms with Hail (1965-1989)	20

COMPOSITES OF SIGNIFICANT WEATHER PATTERNS AFFECTING WEST-CENTRAL MONTANA

*Joe Lester, Dave Bernhardt, and Ken Mielke
NWSFO Great Falls, MT*

I. Introduction

As anyone from west-central Montana will admit, severe or extreme weather conditions are not uncommon to that region of the state. Excessive rain, heavy snow, strong winds, and severe thunderstorms have frequented the area through the years, and their accurate forecast is of vital importance. After identifying the dates of the most extreme weather events to occur in Great Falls, Montana since 1949, and examining their corresponding synoptic patterns, the development of composite fields for each scenario was possible, thereby giving an "average" field for each weather phenomenon. With the development of synoptic composites comes a better understanding of these significant weather events. Furthermore, they may serve as a tool for new forecasters, as well as a refresher for current ones.

II. Data

A compact disc, jointly produced by the National Center for Atmospheric Research (NCAR) and the University of Washington, and distributed by the National Climatological Data Center (NCDC), contains archived gridded data for much of the northern hemisphere. A program named COMP (University of Washington, 1990) allows for the compositing of several grid fields of various dates by producing an output file containing mean values and standard deviations. NMCDRAW (Leblang, 1992) contours and displays the gridded analyses over a North America map background, thus making it possible to view synoptic graphics from specific dates. Furthermore, the software allows the user to view the composited gridded data for specific

synoptic situations. Finally, LASNMC (Leblang, 1992) allows for the printing of the graphics to a laser printer.

Much information is available on the compact disc. Data includes: gridded files of sea level pressure; 850mb heights, temps and winds; 700mb heights and temps; 500mb heights; 250mb winds; and 200mb heights. Most of this data is given daily at 00UTC and 12UTC. The sea level and 500mb fields from 1946 to 1989 are available, whereas the other fields date back to the early 1960s. Once displayed, editing of the graphics can be done by smoothing out the contours, changing the contour interval, and overlaying fields (although only one field can be overlaid on another and no editing of them is allowed).

III. Methodology

Prior to the creation of composite maps, considerable research was performed. The initial task was to define the significant weather phenomena; rain, snow, wind, and thunderstorms were chosen. The data in this particular study was identified and gathered from the Local Climatological Data (LCD) for Great Falls, MT (NCDC). Daily data from the desired time period was easily obtained from the LCDs. After defining the weather phenomena and finding the dates on which these occurred, the gridded data were looked at carefully in hopes of recognizing specific patterns. For example, flow from a certain direction and basic surface or upper-air features could be distinguishing factors. When these

patterns were found, the dates of the significant weather events were grouped accordingly. Without proper sorting, composites would be vague and generally unrepresentative of the situation. Meaningful composites were then produced and analyzed.

IV. Heavy Rain

To begin with, occurrences of heavy rain in Great Falls in a 24 hour period were investigated, with a lower threshold of 1.50 inches chosen. These are fairly rare events, as only 25 cases have met this criteria from 1949 to 1989 (Table 1). Furthermore, 17 of the 25 cases occurred in the spring months of April through June. The latter part of the summer in west-central Montana tends to be warm and dry. The greatest 24 hour rainfall occurred on May 24-25, 1980, when 3.42 inches fell. Incredibly, three of the top eight events happened during an 11 day stretch in 1953. From May 24, 1953 to June 3, 1953 nearly ten inches of rain fell at the Great Falls International Airport, or roughly two-thirds of the annual average. Also, in six of the 24 storms, the air temperature was cool enough to support some rain/snow mix. However, nearly all of the precipitation in these cases was in the form of rain and thus included in this study.

Examination of the 500mb flow at the approximate time of the rain led to the placement of 21 cases into a similar category, with the mean pattern showing a low in northeast Oregon and the trough extending to its south (Fig. 1c). One can also see how this upper-level low develops from the west in the day before the rain and moves over southwest Montana 12 hours after the rain event (Fig. 1a, 1b, 1d). Furthermore, the 500mb field is characterized by a negative tilt trough and southwest diffluent flow over Montana leading up to the rain event. As an approximation, the "12 hours before" composite could represent the onset of the rain and the "12 hours after," the end. The dynamics are favorable for heavy precipitation. The mean sea level pressure field shows a well-developed low-

pressure system over northwest South Dakota (Fig. 2c), which weakens and moves off to the east in the day after the rain (Fig. 2d). Prior to the rain, the low intensifies as it moves out of the central Rockies and toward the northeast (Fig. 2a, 2b). The 250mb wind pattern shows a strong jet streak moving down the Pacific coast, prior to the onset of the precipitation, signifying the strengthening of the upcoming storm system and the deepening of the accompanying trough (Fig. 3a). At the beginning of the rain event, the winds continue to be strong on the back side of the trough, creating a slightly negative tilt (Fig. 3b). This increase in storm energy is coupled with diffluence aloft, which shows up clearly on the 250mb wind field in the left-front quadrant of the jet streak. Upward vertical motion is likely enhanced by this diffluence. Twelve hours later, or during the rain event, these strong upper-air features show a deep trough over the western states with a jet max located in southern Nevada (Fig. 3c).

Rain amounts of over two inches in 24 hours are very rare in Great Falls, as there have been only 11 in the 40-year period. Composites of ten of these storms were produced and subsequently compared to the previous study above. The results are similar, with the most noticeable difference being a slightly deeper 500mb trough developing to the west as early as 72 hours before the heavier rain period (Fig. 4a, 4b). This is possibly due to the elimination of the storms of less than two inches from the composite, as some may have had weaker dynamics. In addition, the 250mb winds composite (Fig. 4c) shows slightly stronger winds on the back side of the trough during the heavier rain, signifying the storms' greater intensity.

V. Heavy Snow

The next situation investigated was extreme snowfall amounts in a 24-hour period. A cut-off of six inches was chosen and 54 such cases were found from 1950 to 1989 (Table 2), thus storms of this caliber generally occur a little more than once per year. The largest 24-hour snowfall in Great Falls occurred on April 20, 1973, when 16.8 inches fell.

Of the 54 snow events, 16 have surface features characterized by an arctic front passing southward through Montana, followed by a large area of high pressure originating in the Yukon Territory, Canada. The remaining 38 events are the result of an intense storm system moving out of the central Rockies. These 38 storms can also be divided into two subsets, one with a strong ridge of high pressure over the Mississippi and Ohio River valleys and the other with a strong high to the north in Alberta and Manitoba, Canada. These two cases contain 19 and 18 events, respectively, and will be looked at separately.

Case 1

The 16 arctic front snowstorms (Case 1) are primarily mid-winter events, as all but one occurred between November and March. In addition, they are the least common of the three observed cases and contain the lightest snow amounts, with only four events of more than eight inches. The sea level pressure chart (Fig. 5a) shows very clearly the large area of high pressure, centered over the Yukon, and the arctic frontal boundary cutting approximately through central Montana. Twenty four hours later (Fig. 5b), the high has pushed southward into Montana and the northern plains. Temperature research has further shown that the high is associated with extremely cold air, as temperatures plummeted after the frontal passage. This type of snowstorm is often coined an "Alberta Clipper." The 500mb flow shows a massive low-pressure system located over Hudson Bay and strong northwest flow aloft over

western Canada and the Pacific Northwest (Fig. 5c). This type of pattern is typical for bringing Alberta Clippers and cold, arctic air to west-central Montana.

Case 2

Case 2, composed of 19 snowstorms, is characterized by an intense surface low-pressure system moving out of the central Rockies and then being steered into eastern Montana and the western Dakotas by a fairly strong ridge of high pressure over the eastern half of the country. It is primarily a springtime event, as only three of this type occurred prior to February. The heavier snow events fall into this category, including the greatest 24-hour total, 16.8 inches. Of the 19 storms, eight are of 10 inches or more. The sea level pressure composite (Fig. 6a) shows a low centered over southeast Wyoming / north-central Colorado and a high pressure to its east. This scenario produces strong upslope conditions at the surface in west-central Montana, which is perhaps prolonged due to the ridge to the east serving as a blocking feature. The 500mb field (Fig. 6b) shows the trough over the western states and the ridge over the Mississippi and Ohio River valleys. The 850mb chart (Fig. 6c) shows a temperature of approximately -2 Celsius over west-central Montana. At an elevation of 1116m, Great Falls is not much below that level, which has a height of about 1390m. Thus, an 850mb temperature of -2 Celsius shows air cold enough to support snow but also warm enough to hold abundant moisture.

Case 3, with 18 storms, involves a similar low-pressure system centered over the Wyoming / Colorado border but strong high pressure exists in Canada as well. This case had only two storms of at least 10 inches but had seven storms over eight inches, making it more potent than the

Case 1 situation. The sea level pressure composite (Fig. 7a) shows a strong gradient over the north central U.S. and upslope conditions in west-central Montana. Case 3 storms tend to move east quickly, whereas the Case 2 events move more slowly and to the northeast. The 500mb field (Fig. 7b) verifies this characteristic by showing zonal flow across the eastern two-thirds of the country, as compared to Case 2, where the 500mb flow contains higher amplitude waves. Similar to before, the 850mb 0 Celsius isotherm runs through central Montana (Fig. 7c).

In west-central Montana, spring snow storms are often the most severe of the season. During the 40-year span of this study, there were 12 storms of eight inches or more in a two month period from March 15 through May 15. Of these 12 events, nine were Case 2 types. Following is a discussion of these nine spring snow storms.

The sea level pressure composite (Fig. 8a) shows a very intense storm system centered over the Colorado / Wyoming / Nebraska area with strong upslope conditions over west-central Montana. This is similar to Fig. 6a except that the low is more intense. Figure 8b shows a deeper 500mb trough as well. One will notice that, although it has weakened somewhat, the low-pressure center at the surface remains in the same position 12 hours later (Fig. 8c). It is possible that this slow storm motion is responsible for prolonged upslope in the west-central Montana region and, therefore, the high precipitation totals. Furthermore, examination of the 250mb flow (Fig. 8d) shows very strong diffluence aloft, giving added lift.

VI. Strong Winds

The next area of interest was surface winds. Although the very strong, gusty winds associated with severe thunderstorms are common, they are generally not represented on a synoptic scale. It was important to look at sustained winds throughout a given day. Therefore, average daily sustained wind speeds of 30 miles per hour or

greater was chosen as the criteria. Because the strongest average winds in Great Falls occur during the winter, the months of November through March were used for this study. From 1963 to 1989, there were 26 such days (Table 3), or roughly one per year. The mean sea level pressure pattern shows a strong gradient between the lower pressures to the north and the higher pressures to the south (Fig. 9a). This pattern, along with the location of the Rockies to the west, results in strong, southwest winds in the Great Falls area. These downslope winds are called chinook winds and are very common in the winter months, with most examples not even considered here as only mean daily winds in excess of 30 miles per hour were investigated. The warming that may occur as a result of the chinook can be shown in the progression of the 850mb heights / temperatures composites (Figs. 9b-c). It is interesting to discover that the 500mb field (Fig. 9d) is very similar to that of Case 1 of the snowstorms (the Alberta Clipper type), and both occur during the winter with huge lows near Hudson Bay. However, crucial differences do exist. Whereas, with the Alberta Clipper, the 500mb flow is from the north with cyclonic curvature to the immediate west of Montana, the chinook winds type has flow from the west, anticyclonic curvature, and much greater heights (giving rise to warm, dry conditions).

An investigation of strong summer-time winds was also undertaken. Average winds are not as strong during the summer so the daily average cutoff was lowered to 25 miles per hour and the year span extended back to 1950. The months of May through September were considered. In the 40-year period, 28 days met this criteria (Table 4). Of the 28 days, only seven occurred during June, July, or August. The high wind occurrence is

primarily a late spring or early fall phenomenon. Also, no thunderstorms occurred on any of the high wind days. Thus, the average speeds are not contaminated by thunderstorm gusts. Seventeen of the 28 days had similar synoptic patterns and were composited. The surface composite shows a low pressure to the north and a high pressure over the Pacific Ocean, resulting in a strong pressure gradient from central Washington to central Montana (Fig. 10a). The structure of the isobars implies the passage of a cold front, stretching through eastern Montana to eastern Colorado. Although at first glance it might appear that the winds should be westerly or northwesterly, terrain in the Great Falls area promotes prevailing southwest winds at the surface. The 850mb and 700mb fields show the colder air moving in behind the front, as well as a tight height gradient in the region (Figs. 10b-c). The 500mb composite (Fig. 10d) shows west to southwest flow with a low to the north. Also evident is the shortwave that had passed through west-central Montana shortly before, and this is confirmed through examination of the 500mb fields before and after the winds (not given).

VII. Severe Weather

The final area of study was severe summer weather, and days with thunderstorms accompanied by hail were chosen. From the year 1965 until 1989, 58 such occurrences were recorded at the Great Falls International Airport, 41 of which fall into a similar category (Table 5). The 500mb composite of these 41 cases shows a trough along the Pacific coast, resulting in southwest flow over central Montana (Fig. 11a). The mean sea level composite features a region of lower pressures to the south, stretching from Wyoming to southern Nevada and into Mexico (Fig. 11b). At the time of the severe weather, the pressure in the Great Falls area was at its lowest, and the low in Wyoming was at its most intense stage. Looking at the 850mb temps/heights composite (Fig. 11c), one will notice very warm air at that level, near 20 Celsius over west-central Montana. This is due to strong

warm air advection leading up to the severe weather, with cooling afterward. The 250mb winds field shows slight diffluence aloft, thus enhancing the upward vertical motion (Fig. 11d).

VIII. Conclusions

Although no two storm systems are alike, composite maps give a good representation of the general features associated with a specific type of weather. Consequently, they act as an effective tool for forecasters to use. Thanks to the easy access to an abundance of historical gridded data, the development of composites is simple.

In Great Falls, Montana, the changes in a variety of weather conditions come frequently and abruptly. This fact necessitates a solid understanding of each situation, which will unquestionably be aided through the interpretation of meaningful composites.

XI. References

- COMP. University of Washington-Atmospheric Sciences Department, 1990.
- Fors, J.R., Leblang, R.S. and Turner, R.G., 1993: *Synoptic Composites of Three Significant Weather Types in North Dakota Using NMC Gridded Data on CD-ROM*. American Meteorological Society Weather Analysis and Forecasting Conference, Vienna, VA.
- Leblang, Richard, 1992: LASNMC. National Weather Service Forecast Office, Bismarck, ND.
- Leblang, Richard, 1992: NMCDRAW. National Weather Service Forecast Office, Bismarck, ND.

National Climatological Data Center: Local
Climatological Data, Great Falls, MT, 1949-
1989.

National Center for Atmospheric Research
and University of Washington, 1990:
*National Meteorological Center Grid Point
Data Set. CDROM, Version II.*

TABLE 1: Great Falls rain of 1.50 inches or more in 24 hours (1949-1989) #

1)	3.42"	May 24-25, 1980	+
2)	3.20"	May 24-25, 1953	
3)	2.74"	August 24-25, 1989	*
4)	2.74"	June 7-8, 1964	
5)	2.43"	April 30 - May 1, 1951	+
6)	2.40"	July 9-10, 1983	
7)	2.40"	May 29, 1953	
8)	2.37"	June 2-3, 1953	
9)	2.29"	June 25-26, 1969	
10)	2.11"	June 1-2, 1954	+
11)	2.09"	June 28-29, 1982	
12)	2.00"	August 15-16, 1985	^
13)	1.94"	May 20-21, 1957	
14)	1.89"	June 24-25, 1965	
15)	1.82"	September 26-27, 1982	+
16)	1.82"	August 19-20, 1974	
17)	1.81"	May 20-21, 1962	
18)	1.73"	September 11-12, 1978	
19)	1.72"	August 9, 1950	
20)	1.67"	May 16-17, 1949	+
21)	1.66"	May 18-19, 1959	
22)	1.64"	May 26-27, 1955	
23)	1.60"	April 25-26, 1986	+
24)	1.51"	July 12-13, 1989	*

All data recorded at the airport

* Data was not available for composite

+ Some snow mixed with rain

^ Did not fit composite pattern




Fig. 1a: 500mb HEIGHTS 24 hours before the rain event. A trough is roughly along the west coast. Winds aloft are from the SSW over Montana.




Fig. 1b: 500mb HEIGHTS 12 hours before the rain event. The trough has not moved at this point, however, the low has intensified. SSW winds aloft continue. This approximately represents the onset of the rain.




Fig. 1c: 500mb HEIGHTS. The trough is now located over the Great Basin and the low has dug further to the south, bringing south winds aloft. Note the slight negative tilt.




Fig. 1d: 500mb HEIGHTS 12 hours after the rain event. This approximately represents the end of the 24 hour rain period. The low is located over southwest Montana and the trough is no longer negatively tilted.

FIGURE 1: 500mb HEIGHTS (dam) for rain events of 1.50 inches or more.




Fig. 2a: SEA LEVEL PRESSURES 24 hours before the rain event. The system is located over Utah and is not yet very well organized.




Fig. 2b: SEA LEVEL PRESSURES 12 hours before the rain event. The low has intensified and is moving out of the central Rockies.




Fig. 2c: SEA LEVEL PRESSURES. The low is centered over northwest South Dakota, giving rise to upslope conditions in west-central Montana.




Fig. 2d: SEA LEVEL PRESSURES 24 hours after the rain event. The low has weakened considerably and moved east.

FIGURE 2: SEA LEVEL PRESSURES (mb) for rain events of 1.50 inches or more.




Fig. 3a: 250mb WINDS 24 hours before the rain event. A jet streak is located off of the Pacific northwest coast, on the back side of the trough. This is a sign of the strengthening of the upcoming storm system.




Fig. 3b: 250mb WINDS 12 hours before the rain event. Winds are still strong on the back side of the trough, creating a slightly negative tilt. Diffuence exists over west-central Montana.




Fig. 3c: 250mb WINDS. A deep trough sets over the western states, and a jet maximum is located over southern Nevada. There is still diffuence aloft, enhancing the lift.

FIGURE 3: 250mb WINDS (m/s) for rain events of 1.50 inches or more.




Fig. 4a: 500mb HEIGHTS 72 hours before the heavy rain event. A trough is developing along the west coast. There is SW flow aloft over west-central Montana.




Fig. 4b: 500mb HEIGHTS. The trough is slightly deeper than in Figure 1c for 1.50 inches or greater rainfall.




Fig. 4c: 250mb WINDS. Winds on the back side of the trough are stronger than those in Figure 3c for 1.50 inches or greater rainfall.

FIGURE 4: 500mb HEIGHTS (dam) and 250mb WINDS (m/s) for rain events of 2.00 inches or more.

TABLE 2: Great Falls snowfalls of six inches or more in 24 hrs (1950-1989) #

CASE 1:

1)	10.2"	January 15-16, 1984	9)	6.9"	January 7, 1962
2)	10.2"	February 21-22, 1982	10)	6.9"	December 3, 1958
3)	8.9"	February 16, 1959	11)	6.8"	April 11, 1986
4)	8.0"	November 11, 1959	12)	6.8"	November 24, 1983
5)	7.4"	December 1-2, 1972	13)	6.6"	March 17, 1950
6)	7.4"	January 2-3, 1966	14)	6.1"	December 23, 1984
7)	7.3"	January 21, 1964	15)	6.1"	December 23, 1977
8)	7.0"	November 26, 1955	16)	6.0"	March 4, 1951

CASE 2:

1)	16.8"	April 20, 1973	11)	8.8"	April 7, 1975
2)	11.5"	March 19-20, 1987	12)	8.6"	May 9, 1983
3)	11.0"	March 28-29, 1977	13)	8.3"	October 3, 1957
4)	11.0"	February 21, 1951	14)	8.1"	February 27, 1953
5)	11.0"	June 7-8, 1950	15)	7.9"	April 2, 1955
6)	10.2"	April 6-7, 1982	16)	6.8"	October, 14, 1975
7)	10.0"	April 25-26, 1976	17)	6.3"	March 27, 1979
8)	10.0"	March 29-30, 1967	18)	6.2"	November 15, 1952
9)	9.0"	February 10, 1978	19)	6.0"	April 7, 1950
10)	9.0"	April 19, 1967			

CASE 3:

1)	11.6"	May 28-29, 1989	10)	7.7"	April 28, 1970
2)	10.3"	April 27, 1989	11)	7.5"	March 19, 1982
3)	9.2"	April 23, 1960	12)	7.1"	October 28, 1972
4)	8.7"	April 5, 1967	13)	6.9"	March 17, 1968
5)	8.6"	January 10, 1988	14)	6.7"	February 26, 1958
6)	8.4"	March 13, 1984	15)	6.4"	October 6, 1985
7)	8.1"	January 6, 1989	16)	6.4"	March 27, 1961
8)	7.9"	February 16, 1952	17)	6.2"	May 10, 1967
9)	7.8"	January 28, 1959	18)	6.1"	September 28, 1954

Note: Gridded data for September 17-18, 1988 snowstorm (8.4") was not available.

All data recorded at the airport




Fig. 5a: SEA LEVEL PRESSURES. An arctic frontal boundary cuts through southcentral Montana. A large high pressure region is to the north.




Fig. 5b: SEA LEVEL PRESSURES 24 hours after the Case 1 snow event. An Alberta Clipper has passed through Montana and high pressure is invading the northern plains, as well as much colder temperatures.




Fig. 5c: 500mb HEIGHTS. A large low is located over Hudson Bay, resulting in WNW flow over west-central Montana.

FIGURE 5: SEA LEVEL PRESSURES (mb) and 500mb HEIGHTS (dam) for Case 1 snow events of six inches or more.




Fig. 6a: SEA LEVEL PRESSURES. Upsloping in west-central Montana is the result of a well developed storm system centered on the CO/WY border. Higher pressure exists over the Mississippi and Ohio River valleys.




Fig. 6b: 500mb HEIGHTS. A trough is located over the western states and a ridge is to the east, resulting in SSW flow aloft.




Fig. 6c: 850mb HEIGHTS (solid) and TEMPERATURES (dashed). The temperature over west-central Montana is about -2°Celsius. A low is centered in Wyoming.

FIGURE 6: SEA LEVEL PRESSURES (mb), 500mb HEIGHTS (dam), and 850mb HEIGHTS (dam) & TEMPS (°C) for Case 2 snow events of six inches or more.




Fig. 7a: SEA LEVEL PRESSURES. A strong low lies on the CO/WY border and a broad high is to the north in Canada. There is a strong pressure gradient in the northcentral U.S. and upslope conditions in west-central Montana.




Fig. 7b: 500mb HEIGHTS. A trough lies over the western U.S., with zonal flow over the eastern two-thirds of the country.




Fig. 7c: 850mb HEIGHTS (solid) and TEMPERATURES (dashed). Similar to Figure 6c, the 0° Celsius isotherm cuts through central Montana. A low is centered in northwest South Dakota.

FIGURE 7: SEA LEVEL PRESSURES (mb), 500mb HEIGHTS (dam), and 850mb HEIGHTS (dam) & TEMPS (°C) for Case 3 snow events of six inches or more.




Fig. 8a: SEA LEVEL PRESSURES. An intense low pressure system is located over the CO/WY border. Strong upslope conditions exist in west-central Montana.




Fig. 8b: 500mb HEIGHTS. The trough is deep, with a low centered in southwest Idaho. A high amplitude ridge is the main feature to the east.




Fig. 8c: SEA LEVEL PRESSURES 12 hours after the heavy Case 2 spring snow event. The low has weakened somewhat but remains in the same position as in Figure 8a. Upslope conditions continue in west-central Montana.




Fig. 8d: 250mb WINDS. A jet maximum is located over southern Arizona and strong diffluence can be found over west-central Montana.

FIGURE 8: SEA LEVEL PRESSURES (mb), 500mb HEIGHTS (dam), and 250mb WINDS (m/s) for heavy Case 2 snow events of eight inches or more from March 15 through May 15.

**TABLE 3: Great Falls mean daily winds of 30mph or greater
from November through March (1963-1989) #**

DATE	AVERAGE SPEED(mph) & DIRECTION	FASTEST MILE(mph)
1) January 30, 1989	33.3 SW	42
2) December 6, 1984	31.2 SW	37
3) January 2, 1984	31.4 SW	42
4) December 11, 1980	33.2 SW	48
5) November 7, 1978	30.1 SW	43
6) January 6, 1977	30.5 SW	34
7) February 11, 1976	32.2 SW	45
8) February 8, 1976	34.1 SW	42
9) January 27, 1976	30.8 SW	47
10) December 21, 1974	30.9 SW	51
11) March 5, 1974	33.8 SW	49
12) January 29, 1974	40.0 SW	52
13) January 15, 1974	32.8 SW	55
14) January 13, 1974	32.5 SW	37
15) December 10, 1973	31.2 SW	36
16) December 26, 1972	30.3 SW	50
17) November 30, 1972	32.1 SW	42
18) January 15, 1972	33.8 SW	36
19) January 8, 1972	31.6 SW	45
20) January 4, 1972	33.2 SW	43
21) December 8, 1971	31.6 SW	37
22) November 19, 1971	31.5 SW	42
23) February 3, 1970	30.3 SW	56
24) December 3, 1968	30.5 SW	47
25) December 23, 1963	31.0 SW	42
26) January 19, 1963	31.3 SW	36

All data recorded at the airport




Fig. 9a: SEA LEVEL PRESSURES. A Great Basin high and lower pressures to the north bring a strong gradient to west-central Montana. Southwest, downslope winds result.




Fig. 9b: 850mb HEIGHTS (solid) and TEMPERATURES (dashed). The isotherms show warming in west-central Montana due to the SW winds.




Fig. 9c: 850mb HEIGHTS (solid) and TEMPERATURES (dashed) 12 hours after the strong winter winds. The isotherms show further warming along the east slopes of the Rockies.




Fig. 9d: 500mb HEIGHTS. A large low is over Hudson Bay. Westerly flow aloft is prominent from the Pacific northwest to British Columbia.

FIGURE 9: SEA LEVEL PRESSURES (mb), 850mb HEIGHTS (dam) & TEMPS (°C), and 500mb HEIGHTS (dam) for strong winter winds of 30mph or greater from November through March.

**TABLE 4: Great Falls mean daily winds of 25mph or greater
from May through September (1950-1989) #**

<u>DATE</u>	<u>AVERAGE SPEED(mph) & DIRECTION</u>	<u>FASTEST MILE(mph)</u>
1) May 19, 1989	29.3 SW	43
2) September 6, 1984	29.7 SW	49
3) May 14, 1971	25.0 SW	38
4) September 22, 1970	26.9 SW	42
5) September 25, 1958	28.0 SW	62
6) September 19, 1958	32.9 SW	57
7) September 14, 1958	25.7 WSW	47
8) September 13, 1958	31.3 SW	56
9) June 28, 1958	26.3 SW	47
10) September 7, 1957	25.3 SW	56
11) July 3, 1957	25.6 W	51
12) September 21, 1956	27.0 WSW	38
13) July 7, 1955	25.8 WSW	40
14) May 8, 1955	27.7 WNW	61
15) September 19, 1954	27.6 WSW	49
16) September 18, 1954	25.8 SW	43
17) June 17, 1954	26.8 WSW	50
18) June 16, 1954	25.6 WSW	46
19) September 28, 1953	26.3 WSW	59
20) May 10, 1953	28.5 NW	38
21) May 7, 1953	25.4 SSW	62
22) June 3, 1953	25.6 WNW	57
23) June 22, 1952	30.5 SW	54
24) May 29, 1952	25.7 SW	45
25) May 25, 1951	29.6 W	52
26) May 24, 1951	25.6 WSW	45
27) May 2, 1951	25.6 SW	42
28) May 22, 1950	26.3 SW	56

All data recorded at the airport
Note: On none of these days did thunderstorms occur




Fig. 10a: SEA LEVEL PRESSURES. A strong pressure gradient exists from eastern Washington to west-central Montana. A cold front stretches from eastern Montana to eastern Colorado.




Fig. 10b: 850mb HEIGHTS (solid) and TEMPERATURES (dashed). The isotherms show colder air moving into west-central Montana behind the cold front. The height gradient is strong.




Fig. 10c: 700mb HEIGHTS (solid) and TEMPERATURES (dashed). As in the 850mb field, the height gradient is strong and cold air advection is occurring in west-central Montana.




Fig. 10d: 500mb HEIGHTS. The flow aloft is from the WSW in west-central Montana.

FIGURE 10: SEA LEVEL PRESSURES (mb), 850mb HEIGHTS (dam) & TEMPS (°C), 700mb HEIGHTS (dam) & TEMPS (°C), and 500mb HEIGHTS (dam) for strong summer winds of 25 mph or greater from May through September.

TABLE 5: Great Falls thunderstorms with hail (1965-1989) #

<u>DATE</u>	<u>TOTAL PRECIP</u>	<u>SIZE OF HAIL</u>
1) May 23, 1989	.23"	1/4"
2) July 5, 1988	.92"	1/2"
3) July 3, 1988	.30"	1/2"
4) May 24, 1988	.02"	1/4"
5) July 24, 1987	.08"	1/2"
6) July 2, 1987	.22"	1/2"
7) June 6, 1987	.45"	1/4"
8) July 26, 1986	.38"	3/8"
9) June 29, 1986	.30"	1/4"
10) June 17, 1986	.05"	3/8"
11) August 7, 1985	.07"	1/4"
12) May 28, 1985	1.12"	3/4"
13) July 25, 1983	.25"	1/4"
14) July 20, 1983	.15"	
15) August 10, 1982	.21"	1/4"
16) June 29, 1982	.76"	1/2"
17) June 28, 1982	1.33"	1/4"
18) August 3, 1981	.06"	1/8"
19) July 6, 1981	.41"	1/8"
20) June 25, 1980	.30"	1/2"
21) June 12, 1980	.64"	1/4"
22) June 30, 1979	.28"	1/4"
23) June 16, 1979	.89"	1/4"
24) August 22, 1978	.30"	1/2"
25) June 29, 1978	.17"	3/8"
26) June 22, 1978	.14"	1/4"
27) August 27, 1977	.15"	1/4"
28) August 14, 1977	.37"	1/4"
29) July 12, 1976	.39"	1/4"
30) July 11, 1976	1.00"	1/8"
31) July 6, 1976	.43"	1/2"
32) June 10, 1976	.07"	3/4"
33) June 3, 1976	.28"	3/4"
34) June 30, 1975	.91"	1/2"
35) August 31, 1971	.32"	3/8"
36) June 27, 1970	.56"	1/4"
37) August 6, 1967	.11"	1/2"
38) July 2, 1966	.77"	1/8"
39) July 11, 1965	.11"	1/4"
40) July 9, 1965	.17"	1/4"
41) June 24, 1965	.70"	3/4"

All data recorded at the airport and only those storms used in composites are listed




Fig. 11a: 500mb HEIGHTS. A trough over the west coast brings SW flow aloft in west-central Montana.




Fig. 11b: SEA LEVEL PRESSURES. Nothing is well-organized, however, lower pressures lie to the south of Montana.




Fig. 11c: 850mb HEIGHTS (solid) and TEMPERATURES (dashed). Very warm air, up to 20 degrees Celsius, lies in west-central Montana. Colder air moves in after the severe weather as the wave moves east.




Fig. 11d: 250mb WINDS. Strong SW flow can be seen aloft, as well as slight diffluence.

FIGURE 11: 500mb HEIGHTS (dam), SEA LEVEL PRESSURES (mb), 850mb HEIGHTS (dam) & TEMPS ($^{\circ}$ C), and 250mb WINDS (m/s) for severe weather events.

- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033)
- 152 Climate of Salt Lake City, Utah, William J. Alder, Sean T. Buchanan, William Cope (Retired), James A. Cisco, Craig C. Schmidt, Alexander R. Smith (Retired), Wilbur E. Figgins (Retired) April 1996 - Sixth Revision)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82 158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82 177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidl, April 1969 (Revised December 1986). (PB87 142063/AS)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteris, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85 110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85 132694)
- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 1878887AS)
- 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lussky, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lussky, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Rapid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
- 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
- 199 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Lussky, April 1987. (PB87 185229/AS)
- 200 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke, September 1987. (PB88 110895/AS)
- 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson and George Clark, September 1987. (PB88 114038/AS)
- 202 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88 101935/AS)
- 203 An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern California. John P. Monteverdi and Scott A. Braun, October 1988. (PB89 121297/AS)
- 204 Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Carven Scott, November 1988. (PB89 128649/AS)
- 205 Forecast Guidelines For Fire Weather and Forecasters - How Nighttime Humidity Affects Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS)
- 206 A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarters, Scientific Services Division, August 1989. (PB89 230833/AS)
- 207 The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, June 1990. (PB90-240268)
- 208 Meteorological Factors Contributing to the Canyon Creek Fire Blowup, September 6 and 7, 1988. David W. Goens, June 1990. (PB90-245085)
- 209 Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow Whitlatch, December 1990. (PB91-129239)
- 210 Hydrotools. Tom Egger, January 1991. (PB91-151787/AS)
- 211 A Northern Utah Soaker. Mark E. Struthwolf, February 1991. (PB91-168716)
- 212 Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 1991. (PB91-208439)
- 213 Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Mollner, July 1991. (PB91-227405/AS)
- 214 Emergency Operational Meteorological Considerations During an Accidental Release of Hazardous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424)
- 215 WeatherTools. Tom Egger, October 1991. (PB93-184950)
- 216 Creating MOS Equations for RAWFS Stations Using Digital Model Data. Dennis D. Gettman, December 1991. (PB92-131473/AS)
- 217 Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 1992. (PB92-196104)
- 218 NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 1992. (PB93-146785)
- 219 A Case Study of the Operational Usefulness of the Sharp Workstation in Forecasting a Mesocyclone-Induced Cold Sector Tornado Event in California. John P. Monteverdi, March 1993. (PB93-178697)
- 220 Climate of Pendleton, Oregon. Claudia Bell, August 1993. (PB93-227536)
- 221 Utilization of the Bulk Richardson Number, Helicity and Sounding Modification in the Assessment of the Severe Convective Storms of 3 August 1992. Eric C. Evenson, September 1993. (PB94-131943)
- 222 Convective and Rotational Parameters Associated with Three Tornado Episodes in Northern and Central California. John P. Monteverdi and John Quadros, September 1993. (PB94-131943)
- 223 Climate of San Luis Obispo, California. Gary Ryan, February 1994. (PB94-162062)
- 224 Climate of Wenatchee, Washington. Michael W. McFarland, Roger G. Buckman, and Gregory E. Matzen, March 1994. (PB94-164308)
- 225 Climate of Santa Barbara, California. Gary Ryan, December 1994. (PB95-173720)
- 226 Climate of Yakima, Washington. Greg DeVoy, David Hogan, and Jay Neher, December 1994. (PB95-173688)
- 227 Climate of Kalispell, Montana. Chris Maier, December 1994. (PB95-169488)
- 228 Forecasting Minimum Temperatures in the Santa Maria Agricultural District. Wilfred Pi and Peter Felsch, December 1994. (PB95-171088)
- 229 The 10 February 1994 Oroville Tornado-A Case Study. Mike Staudenmaier, Jr., April 1995. (PB95-241873)
- 230 Santa Ana Winds and the Fire Outbreak of Fall 1993. Ivory Small, June 1995. (PB95-241865)
- 231 Washington State Tornadoes. Trestle Huse, July 1995. (PB96-107024)
- 232 Fog Climatology at Spokane, Washington. Paul Frisbie, July 1995. (PB96-106604)
- 233 Storm Relative Isentropic Motion Associated with Cold Fronts in Northern Utah. Kevin B. Baker, Kathleen A. Hadley, and Lawrence B. Dunn, July 1995. (PB96-106596)
- 234 Some Climatological and Synoptic Aspects of Severe Weather Development in the Northwestern United States. Eric C. Evenson and Robert H. Johns, October 1995. (PB96-112958)
- 235 Climate of Las Vegas, Nevada. Paul H. Skrbac and Scott Cordero, December 1995. (PB96-135553)
- 236 Climate of Astoria, Oregon. Mark A. McInerney, January 1996.
- 237 The 6 July 1995 Severe Weather Events in the Northwestern United States: Recent Examples of SSWEs. Eric C. Evenson, April 1996.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS--Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS -- Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161