



**NOAA TECHNICAL MEMORANDUM
NWS WR-253**

**INFERENCE OCEANIC KELVIN/ROSSBY WAVE
INFLUENCE ON NORTH AMERICAN WEST COAST
PRECIPITATION**

**Martin E. Lee
National Weather Service
Northwest River Forecast Center
Portland, Oregon**

**Dudley Chelton
College of Oceanic and
Atmospheric Sciences
Oregon State University
Corvallis, OR**

April 1998

**U.S. DEPARTMENT
OF COMMERCE**

National Oceanic and
Atmospheric Administration

National Weather
Service



NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, 125 South State Street - Rm 1210, Salt Lake City, Utah 84138-1102. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.


ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidli, April 1968 (Revised March 1985). (PB86 177672/AS). (Revised October 1991 - PB92-115062/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB 179084)
- 32 Probability Forecasting—A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento—Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A. L. Jacobson, April 1969. PB 184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L. W. Snellman, August 1969. (PB 185068)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)
- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 65 Climate of Sacramento, California. Richard Honton and Tony Martini (Retired), August 1996. (Fifth Revision) (PB89 207781/AS)
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM

- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251)
- 83 A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS)
- 96 Map type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 126 Climate of San Francisco. E. Jan Null, February 1978. (Revised by George T. Pericht, April 1988 and January 1995). (PB88 208624/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Molner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric



**NOAA TECHNICAL MEMORANDUM
NWS WR-253**

**INFERENCED OCEANIC KELVIN/ROSSBY WAVE
INFLUENCE ON NORTH AMERICAN WEST COAST
PRECIPITATION**

**Martin E. Lee
National Weather Service
Northwest River Forecast Center
Portland, Oregon**

**Dudley Chelton
College of Oceanic and
Atmospheric Sciences
Oregon State University
Corvallis, OR**

April 1998

UNITED STATES
DEPARTMENT OF COMMERCE
William M. Daley, Secretary

National Oceanic and
Atmospheric Administration
D. James Baker, Under
Secretary and Administrator

National Weather Service
John J. Kelly, Jr., Assistant
Administrator for Weather Services

**This publication has been reviewed
and is approved for publication by
Scientific Services Division,
Western Region**



**Delain A. Edman, Chief
Scientific Services Division
Salt Lake City, Utah**

TABLE OF CONTENTS

I. Introduction	1
II. Significant 1994-1996 Trans-Pacific Ocean/Atmosphere Patterns	2
III. Oceanic Rossby Wave Observations and Theory	5
IV. Tropical Air-Sea Interaction	7
V. Proposed Ocean-Atmosphere Conceptual Model	9
VI. Conclusion	11
VII. Acknowledgments	14
VIII. References	15

Inferenced Oceanic Kelvin/Rossby Wave Influence on North American West Coast Precipitation

Martin E. Lee *
•Dudley Chelton **

Abstract

The complex interaction of coupled ocean-atmosphere processes is still far from being completely understood, and documented. This Technical Attachment documents a nonclassic, oceanic Kelvin/Rossby wave conceptual model to partially explain above normal 1994-1996 precipitation patterns along the North American west coast. The proposed conceptual model focuses on an eastern Pacific post-El Niño heat transfer mechanism associated with oceanic Kelvin/Rossby waves that potentially affect preferential atmospheric forcing, resulting in selectively focused offshore storm track and moisture advection trajectories. Synergistic components of this model thereby have the potential to enhance (or reduce) U.S. west coast interannual precipitation. The magnitude of U.S. west coast flood damages incurred during the winters of 94/95 and 95/96 imply that analysis and assimilation of this proposed conceptual oceanic Kelvin/Rossby wave interaction may be worth considering in order to improve accuracy in forecast expectations for timing and placement of both above and below normal west coast annual and interannual precipitation. Other implications resulting from this research also include: 1) the imminent necessity of incorporating a multidisciplinary, combined earth systems approach in west coast precipitation forecasting; and 2) a concluding argument regarding the continued necessity for in situ oceanic observation programs beyond tropical latitudes.

I Introduction

Accurate prediction of local precipitation is important for early flood warning and reservoir management, especially in mountainous areas of the western United States where severe flooding during heavy precipitation occurs in steep terrain and narrow valleys (Miller and Kim 1996). The National Weather Service (NWS),

recognizing the economic and social value of accurate precipitation forecasts, dedicates significant resources to increase lead times of watches and warnings for flood and flash flood events (Olson et al. 1995). However, abrupt and vast data sparse upstream conditions commencing along a dominant land-sea boundary, marked by the west coast of the United States and the Pacific Ocean,

* National Weather Service, Northwest River Forecast Center, Portland, Oregon.
Corresponding author address: Martin E. Lee, NWS NWRFC, 5241 NE 122nd Ave., Portland, OR 97230. E-mail: mlee@nwrfc.noaa.gov.

** College of Oceanic and Atmospheric Sciences, Oregon State University, Corvallis, OR 97331; also a TOPEX/POSEIDON Science Working Team Principal Investigator.

introduce pronounced complexity to precise forecasting of significant precipitation. Upstream, air-sea interaction has the potential to modify both atmospheric temperature and wind fields that may then alter even global-scale atmospheric and oceanic circulations (Minwei and Dayong 1988). The ability to produce reliable forecasts in NWS Western Region locations of the United States is thus dependant on an appreciation and conceptual understanding of significant upstream ocean-atmosphere interactions.

During the winter of 1994-95, California received unusually large amounts of precipitation, with the Russian River Basin being among the hardest hit areas in January 1995, reporting over an estimated 800 million dollars in flood-related damages from just three consecutive strong storms between 7 and 11 January (Miller and Kim 1996). In the following winter (1995-96), western Oregon and Washington were inundated with numerous record-breaking winter floods. What substantive mechanism/s contributed to this sequence of interannual events? In general terms, variation in Pacific coast precipitation is the result of "annual variation in the interaction between the maritime westerlies from the North Pacific and the mountains of the U.S. West Coast" (Chen et al. 1996).

This Technical Memorandum considers hypothetical "nonclassic" (Hobbs et al. 1996) upstream ocean-atmosphere interactions responsible for producing excessive winter (1994-95, and 1995-96) west coast precipitation. Although not necessarily in the respective order given here, the dominant conceptual

considerations in this analysis can be summarized by the following questions: 1) What are oceanic Rossby waves? 2) Did an interannual (1994-1996) heavy precipitation pattern shift northward along the west coast of the United States in association with a propagating oceanic Rossby wave associated with a pronounced but concluding El Niño event, trailing off with diminishing propagation speed at increasingly high latitudes? 3) What is the surface manifestation of oceanic Rossby waves as pertains to air-sea interaction that may affect precipitation? 4) Are sea surface temperature (SST) anomalies associated with oceanic Rossby waves significant enough to alter preferred low-mid latitude moisture advection trajectories, localized atmospheric forcing, and resultant interannual west coast precipitation patterns? and 5) Is it cost-effective to even consider oceanic Rossby waves as a relevant factor in North American west coast operational forecasting? and 6) What are the short-long term implications if the answer to question number 5 is "yes"?

II Significant 1994-1996 Trans-Pacific Ocean/Atmosphere Patterns

The Spring of 1995 was marked by below-normal equatorial eastern Pacific SSTs, and was followed by an active western Pacific Typhoon season in the fall of 1995. This indicated that warm equatorial surface water had shifted west of the date line. The spring and summer of 1996 provided further confirmation of this in a disastrous manner. During this period Typhoons Ann, Frankie, Gloria, and Herb produced severe damage in

Eastern Asia. For example, early-mid July 1996 severe, typhoon related weather and flooding in southern-central China was responsible for approximately 1,108 deaths, and an estimated \$11.23 billion in economic losses across Anhui, Guangxi, Guizhou, Hubei, Hunan, Jiangsu, Jiangxi, and Zhejiang provinces. In these provinces, flooding was responsible for damage to approximately 2.8 million homes and the collapse of another 810,000 homes, leaving millions of people homeless, and also destroying over 5 million tons of grain (Reuters Limited, Beijing: July 25, 1996). Before moving into Eastern China, Typhoon Gloria was responsible for an estimated 50 deaths in the Philippines (Philippine News Agency, Manila: July 29, 1996), and \$21 million in crop damages across southern Taiwan (Reuters Limited, Hong Kong: August 2, 1996). Typhoon Herb then caused another 39 deaths in the Philippines (Reuters Limited, Manila: July 29, 1996), also before clipping Taiwan and moving into Eastern China.

Remnant decaying tropical storms advecting to higher latitudes then resulted in 68 deaths and \$246 million in flood damage across South Korea (Reuters Limited, Seoul: July 31, 1996). This included the loss of an estimated four tons of ammunition (hand grenades, mortar and artillery shells, and land mines) swept away by flood waters. Another estimated 860 tons of ammunition stored near the North Korean Border was buried by flood-induced landslides. During this stormy season in the western Pacific, Typhoon Bart typified a fully recurving storm (Fig. 1), with its decaying remnants entraining into a trans-Pacific westerly jet stream, eventually resulting in U.S. west coast precipitation interception during May

1996. However, this U.S. west coast precipitation was minor in significance when compared to cumulative U.S. west coast winter 1994-95 and 1995-96 precipitation.

Halpert et al. (1996) note that abnormally high 1994-95 winter season precipitation was observed in California and the southwestern United States. This pattern, especially from December 1994 to February 1995, was associated with an eastward extension of a jet stream core across the Pacific and into California. During January 1995, the jet stream core and storm track aimed at California was displaced approximately 18° of latitude south of their typical location. Resulting significantly above-normal precipitation steered into California at that time, producing excessive flooding throughout the state. During January 1995, the southern Cascades, the northern Sierra Nevadas, and several central-coastal California locations observed 950-1050 mm of precipitation, and October-April totals in parts of southern California reached at least two times the normal seasonal total records (Fig. 2). According to the Products and Services Division of the National Climate Data Center, over 3.0 billion dollars in flood damage was assessed from January-March 1995 flooding in California – including 27 deaths.

In the following winter (1995-96), severe river flooding (especially west of the Cascades) occurred in western Washington and northwestern Oregon, where 200-500 mm of heavy rain fell from November-December 1995 (Fig. 3). Enhanced warm, moist, southwesterly flow facilitated this excessive precipitation pattern anomaly. As noted above, March-

May 1995 marked the end of El Niño/Southern Oscillation (ENSO) conditions across the Pacific. For example, below normal SSTs developed from the South American coast and the equatorial Pacific to 120°W, where SSTs reached -1.0°C in many areas (Fig. 4). Central Pacific maximum positive SST anomalies had significantly diminished, and central equatorial Pacific atmospheric convection returned to normal. Near normal low-level equatorial easterlies also reappeared, and western Pacific SST anomalies exceeded 0.5°C above normal between 120°E to 150°E.

El Niño processes, by classic definition, were therefore relinquishing control of weather patterns over the eastern Pacific. Yet, according to the Products and Services Division of the National Climate Data Center, February 1996 Oregon uninsured flood damage alone reached 400 million dollars. Over 300 million dollars damage to Oregon highways and waterway infrastructures was also reported. Overall flood damage to the Pacific Northwest is now estimated to approach \$1 billion – including 14 fatalities. Thus, one of the driving questions becomes: “What other mechanisms have we learned enough about to explain this above normal, non El Niño, winter precipitation pattern over western North America?”

Surprising structure and symmetry in eastern Pacific March-May 1996 positive SST anomalies (Fig. 4) provide potential clues to answering the previously raised question. For example, a coupled, northern and southern hemisphere, poleward "banana-shaped" positive SST anomaly structure is evident over the eastern Pacific in Fig. 4, symmetric about

the equator. Subsidiary clues are derived from an apparent northwestward shift of eastern Pacific positive SST anomalies along the U.S. west coast, from March-May 1995 to September-November 1995. This northern hemisphere, northwestward SST positive anomaly shift is puzzling in the context of first order eastern Pacific semi-permanent atmospheric and ocean current circulation patterns. For example, clockwise circulation around semi permanent eastern Pacific high pressure results in mean equatorward surface wind components along the southwestern U.S. coast. Similarly, eastern Pacific mean ocean currents along the western U.S. coast are characterized by equatorward flow (Fig. 5).

A direct correlation exists between the organized eastern Pacific positive SST anomaly behavior described above and positive Sea Surface Height (SSH) anomaly structure (Fig. 6). Additionally, this 1995-1996 ocean system phenomenon has symmetry matching strong November 1995 and February 1996 storm tracks – routed from near the Hawaiian Islands, along northward curving subtropical jet streams, often referred to as a “Pineapple Express.” These associations eventually resulted in the formulation of a conceptual model that can explain the previously discussed northwestward shift of SST anomalies. The positive, symmetric SSH anomaly structure in Fig. 6 correspond to positive SST anomalies (Bob Cheney 1996, personal communication) via a selective redistribution of heat energy (Dudley Chelton 1997, personal communication). Both the positive SST and SSH offshore anomalies are associated with oceanic Rossby waves. In the following sections, this association is developed into a

conceptual model that can be used to inferentially explain how and why oceanic Kelvin and Rossby waves may be linked directly to: 1) poleward shifts of eastern Pacific positive SST anomalies; and 2) selectively focused precipitation patterns along the U.S. west coast. Concluding sections of this text also explore the plausibility and ramifications of this proposed model, as well as subsidiary planning, design, and management issues.

III Oceanic Rossby Wave Observations and Theory

Technological advancement in observation sampling capabilities has made it possible to improve our understanding of ocean circulations that play critical roles in varying climate over a wide range of time scales: from seasonal to interannual (e.g., El Niño phenomenon) to decades and even centuries - as revealed in recent ice core analyses (Johnson et al. 1992). Our ability to identify ever smaller scale ocean circulation processes is also increasing. For example, the decline in intensity of the eastward flowing Kuroshio Extension and its westward recirculation gyre has been linked to energy transfers from the mean current flow to an eddy field in a region south of the Kuroshio Extension (Qiu 1995). In the quest to further understand these increasingly evident complex ocean system processes, remote sensing has become the most practical approach to observe seasonal-to-interannual ocean variability on global scales with adequate space and time resolution; and recent advancements in satellite observing systems have significantly improved our ability to monitor global ocean circulation.

A precision radar altimeter aboard the TOPEX/POSEIDON satellite provides the most accurate, currently available altimeter data for studying general ocean circulation and its variability (Fu et al. 1994; Fu and Cheney 1996). This is possible using altimetric measurement of sea surface elevation, because low-frequency sea level variations (i.e., with tidal variation removed) are indicative of motions throughout much of the water column - extending to the sea floor in certain instances (Pedlosky 1979). TOPEX is the U.S. acronym for Ocean Topography Experiment, a joint mission with France. POSEIDON, the mythical Greek god of the sea, is the acronym for the French contribution to this mission. The TOPEX/POSEIDON satellite is a modified Multimission Modular Spacecraft configured with the first spaceborne dual-frequency (5.3 and 13.6 GHz) radar altimeter. The radar altimeter was designed and constructed by the Applied Physics Laboratory of the Johns Hopkins University (APL/JHU), and the 2,500 kg satellite was launched into a highly stable orbit on August 10, 1992, by an Ariane 42P rocket from the European Space Agency's Guiana Space Center in French Guiana (Fu et al. 1994). The launch vehicle was contributed to this project by the "Centre National d'Etudes Spatiales (CNES)," or the National Center of Space Studies, France's equivalent of NASA.

The TOPEX/POSEIDON satellite has a 66° inclination orbit ~ 1,336 km above the Earth's surface (Fig. 7). Sea surface height is observed by a TOPEX radar signal reflected off the ocean's surface. The dual-frequency altimeter allows for a correction for ionospheric effects on the 2-way travel time of radar pulses. The time calculated for a standard local sea level

radar return is compared to the measured return time. If the return signal takes less time, the ocean surface is above its standard reference sea level. Insertion into orbit was so successful that sufficient fuel remains aboard the satellite to sustain its mission significantly beyond the year 2000.

Satellite altimetry, provided via projects like the TOPEX/POSEIDON mission, is probably the only way that global oceanic circulations can be studied relative to the sea surface as a reference level, because the geostrophic surface velocity can be determined from the satellite derived surface height (Fu and Smith 1996). Ocean surface currents are initially forced by momentum transfer from the atmosphere. Yet, once free from direct forcing by the wind, these nearly geostrophic currents tend to move slowly and nearly frictionlessly under pressure gradient and Coriolis force influences. Thus, sea surface height anomaly data across these currents facilitates theoretical current velocity estimations. A positive 1 cm sea surface height change may generate an elevated slab containing the equivalent of several millions of cubic meters of water per second, a rate of discharge greater than all the rivers in the world (from the JPL TOPEX/ POSEIDON Mission HomePage, <http://topex-www.jpl.nasa.gov/>). Although lesser in scope, significant responses also occur at smaller than global-hemispheric scales, where other quasigeostrophic processes are at work. Oceanic Rossby waves fall within this category. The theoretical oceanic Rossby wave length scale L_r (~ the Rossby deformation radius) is equal to the product of a characteristic buoyancy frequency and its vertical scale divided by the Coriolis parameter. At ~ 30° latitude,

$L_r \sim 70$ km, compared to the analogous atmospheric Rossby length scale, $L_A \sim 1900$ km. Conversely, the time scale for $L_r \sim 60$ days, while the time scale for $L_A \sim 2$ days (Stewart 1985). Local vertical, latitudinal variation of the Earth's angular rotation vector – resulting from curvature of the Earth's surface – provides the restoring force for oceanic Rossby waves (Pedlosky 1979). These waves are considered to play a key role in modern large-scale ocean circulation theory. They are also considered an important mechanism in the westward intensification of large-scale circulation gyres (Anderson and Gill 1975; Anderson et al. 1979).

Wind and thermal buoyancy forcing, at eastern ocean boundaries and interior regions, generate oceanic Rossby waves; and El Niño events are also oceanic Rossby wave generators (Barnett et al. 1991; Neelin et al. 1994; Battisti and Sarachik 1995). Eastward propagating oceanic equatorial Kelvin waves, forced in the western Pacific by wind anomalies associated with El Niño events, generate poleward propagating coastally trapped Kelvin waves when they reach the eastern boundary of the Pacific, resulting in positive sea surface height anomalies that have been observed to propagate as far north as the Gulf of Alaska. These forced, positive sea surface height anomalies generate westward propagating oceanic Rossby waves along the west coast of the Americas. Model results suggest that these westward propagating waves may extend well into high latitudes, covering the entire North Pacific, remaining largely undamped to 160° W longitude, and continuing into the western Pacific a decade later (Jacobs et al. 1994). Jacobs et al. (1994) also note that El Niño generated oceanic Rossby waves may be

linked with North Pacific mid latitude ocean circulation anomalies a decade later. They have hypothesized that these transient ocean circulations may have a non-trivial impact on North American weather patterns. The simplistic theoretical impact these Rossby waves have on eastern Pacific meteorology is that they appear to facilitate a complex exchange process. The 1-10 cm sea surface height signatures of oceanic Rossby waves are associated with much greater thermocline depth variations (~ 3 orders of magnitude, and opposite in sign). Thus, oceanic Rossby wave signals represent large thermal variations in the upper ocean.

Initially, oceanic Rossby waves provide transient ocean circulation adjustment in response to large-scale atmospheric forcing. The resulting anomalous change in ocean circulation may then be capable of influencing climate. Despite their importance in time varying oceanic processes, Rossby wave surface signatures are difficult to detect, with height variations of less than 10 cm over wavelengths ranging from *hundreds* to *thousands* of kilometers. However, the TOPEX/ POSEIDON altimeter is capable of continuous measurement accuracy of ~ 3 cm since becoming fully operational in October 1992, and amplitudes as small as ~ 1 cm can be detected in large-scale sea level signals over the entire world ocean, after measurement error filtering is performed. The TOPEX/POSEIDON altimeter is thus able to detect oceanic Rossby waves unambiguously (Schlax and Chelton 1994; Parke et al. 1987; Fu et al. 1994). The remainder of this Technical Memorandum examines the associative correlations of these oceanic Rossby waves, via their sea surface

height and SST anomalies, with potential 1994-1996 eastern Pacific and western North American weather modification.

IV Tropical Air-Sea Interaction

Bocheng and Qiyu (1988) remark that "the western tropical Pacific Ocean is the largest heat and moisture source on the earth...." As a result, it has a significant impact on Asian, as well as world climate. Variations in western tropical Pacific current systems are intimately linked to El Niño events. This air-sea interaction is the subject of intensive research that has resulted in our increased appreciation of how both oceanic and atmospheric responses are inextricably dependent upon forcings and exchanges between these two complicated systems. For example, Hendon and Salby (1996) note that atmospheric frictional wave-CISK (conditional instability of the second kind) explains eastward frequency selections in western Pacific tropical to subtropical atmospheric circulation and anomalous convection interactions. In the western Pacific this interaction is highly correlated with equatorial frictional convergence that shifts towards warm SST anomalies, and is associated with dynamical responses involving forced Kelvin and Rossby wave disturbances. Shifting toward positive temperature anomalies results in a net tropical eddy available potential energy gain that facilitates amplification of eastward convective components, and the frequency of these eastward components is dominated by unstable tropical Kelvin wave structure in the atmosphere. However, frictional wave-CISK correlations (assuming a zonal SST variation) become insignificant in the eastern Pacific - because typically colder SSTs do not provide sufficient

climatological convection. This distinction of eastern Pacific tropical-subtropical atmospheric processes, different from those in the western Pacific, is another manifestation of how important it is to understand air-sea interactions over the Pacific.

Tropical Pacific Ocean SSTs are relatively uniform, in that cold, dense water characteristically lies below a thin, wind-mixed layer (~ 100 m thick) of much warmer water. The highly stratified region separating the weakly stratified shallow and deep layer is called the thermocline (Stewart 1985). However, at the onset of the 1986-1988 El Niño/Southern Oscillation (ENSO) episode, anomalous eastward transport of warm surface mixed layer water was observed equatorward of 10° N latitude. In response to the eastward export of surface mixed layer water away from the western Pacific, the thermocline in the tropical western Pacific essentially "shoaled," coinciding with observed decreases in dynamic height over the upper (300-400 m) tropical western Pacific Ocean (Toole et al. 1988). Eastward export of warm surface layer water also resulted in colder temperatures observed over the surface of the tropical western Pacific Ocean.

According to Chan and Lau (1988), SSTs are one of the most significant factors in the generation and maintenance of tropical convective cloud processes. These convective processes have been correlated with selective variance around tropical SSTs of 28°C, and outgoing longwave radiation values of ~ 240 Wm⁻². The SST relationships between a convectively active, relatively warm western Pacific versus a convectively suppressed, relatively cooler eastern

Pacific may vary by only a few degrees C. Hansen (1988) categorized ~ 1°C anomalies observed during 1986 in the Pacific as "large." Various moist process cycles, resulting in eastward propagation of convection toward the eastern Pacific (including El Niño/Southern Oscillation (ENSO) events) may be attributed to these smaller warm SST anomalies through their enhancement of unstable conditions conducive to sustained atmospheric convection (Chan and Lau 1988).

This unstable air-sea interaction is largely driven by turbulent flux exchange across the ocean-atmosphere interface, transforming SST anomalies into diabatic heating that results in forcing of the free atmosphere (Min wei and Dayong 1988). In the tropics, significant positive and/or negative SST anomalies can also be frequently correlated with observed changes in atmospheric boundary layer wind regimes (Dayong et al. 1988). In the tropical eastward on-set of an El Niño event, positive SST anomalies directly result in surface wind convergence and associated precipitation fields (Barnett et al. 1991). Barnett et al. (1991) caveat that the direct effect of heat exchange from the ocean to the atmosphere is relatively weak in forcing of the main troposphere. However, the air-sea temperature difference has a very important indirect role in the thermodynamics of atmospheric circulation: the low-level surface wind convergence associated with the positive SST forcing provides critical initiation of vertical mass and water vapor flux above the planetary boundary layer (PBL), thus determining where heat from condensation in the troposphere occurs. This tropospheric heat of condensation is estimated to be four times greater than

sensible heat flux from the ocean into the lower PBL, but its positioning is tied to the positive SST anomaly feedback.

Selective subtropical to global scale temperature modifications result at the hemispheric scale from Hadley circulation of net poleward energy flux, and the Hadley circulation of this energy flux is largely forced by the addition of low latitude surface latent heating (Palmén and Newton 1969; Min wei and Dayong 1988). On the scale of tropical Kelvin wave processes in the eastern Pacific, Hendon and Salby (1996) conclude that convective disturbances "can be understood as the propagating response to transient heating that is localized." Using numerical simulations, Wendell (1988) also found that even midlatitude SST warming significantly impacted cyclogenesis in boundary layer interaction with cyclones, via increasing planetary boundary layer (PBL) "convergence and redistribution of surface heat to the midtroposphere...." Jacobs et al. (1994) have observed high-latitude SST anomalies across the North Pacific of the same magnitude and temporal-spatial extent as those in equatorial latitudes during El Niño events. Despite relatively minor magnitudes of positive SST anomalies (i.e., compared to magnitudes of typical frontal inversions in the atmosphere), the microscale effects of converging low-level moist air also contribute to further destabilizing of the PBL (Stull 1988). Collectively, these low-mid latitude dynamic processes thus have the potential to re-define higher latitude storm tracks, primarily as a result of mean zonal temperature field modification (Lee and Mak 1996).

V Proposed Ocean-Atmosphere Conceptual Model

Fig. 6 is a global map of sea level constructed from TOPEX/ POSEIDON altimeter data acquired in April 1996. The positive SSH anomalies of ≥ 3 cm over the eastern Pacific, in both the northern and southern hemispheres, have a structure consistent with that expected for an oceanic Rossby wave crest: the positive SSH anomaly gradually curves from mid-high latitude far eastern Pacific coastal boundaries towards western Pacific tropical latitudes. Positive SST anomalies in Fig. 4 have similar structure to that of the positive SSH anomalies in Fig. 6. These two positive anomaly structures are directly associated, and the mechanism responsible for this correlated structure is the latitudinal variation of oceanic Rossby waves initiated nearly simultaneously along the eastern boundary of the Pacific (Chelton and Schlax 1996). These latitudinally dependent oceanic Rossby waves are thought to be initiated by passage of coastally trapped, poleward propagating baroclinic oceanic Kelvin waves. Oceanic Kelvin waves have been observed in current meter and hydrographic data along the west coast of South America (Smith 1978; Huyer 1980), and Dorman (1985) has presented evidence of trapped, internal oceanic Kelvin wave existence along the coast of California.

Coastally trapped, oceanic Kelvin waves, propagating poleward along the west coasts of North and South America, are balanced directly along the coast by pressure gradient and Coriolis forces (Dorman 1985). Allowing poleward propagating, trapped oceanic Kelvin waves originating off Central America to

reach the Gulf of Alaska, this relatively undamped ageostrophic force balance is directly analogous to trapped barrier jets in the atmosphere. Trapped oceanic Kelvin waves decay exponentially offshore over a length scale equivalent to the Rossby radius of deformation. The initiation of a westward propagating oceanic Rossby wave by a trapped, poleward propagating oceanic Kelvin wave is illustrated in Fig. 8. Because of the " β -effect" of latitudinal variation of the Coriolis parameter, SSH anomalies associated with coastal Kelvin waves propagate westward as oceanic Rossby waves. As time elapses and trapped, oceanic Kelvin waves continue to propagate poleward and oceanic Rossby waves are initiated at increasingly higher latitudes.

The phase delay in the initiation of oceanic Rossby waves at higher latitudes contributes to the increasing lag of more poleward Rossby wave initiation. Also, the phase speed of oceanic Rossby waves decreases rapidly with increasing latitude at midlatitudes (Fig. 9). These two factors result in wave crests and troughs that are approximately symmetric about the equator, as previously discussed in terms of the positive SST and SSH anomaly symmetry in Figs. 4 and 6. The associations presented in the previous section suggest the potential atmospheric forcing effects that this oceanic Rossby wave crest-front may have on eastern Pacific organized precipitation processes. In summary, this ocean-atmosphere interaction is linked by geographically focused boundary layer latent heat flux that is selectively forced by the structured, positive SST patterns associated with oceanic Rossby waves. Could this selective enhancement of

upstream precipitation along preferential storm tracks have resulted in heavy, post El Niño west coast precipitation (Figs. 2-3, 8, and 10)?

A simple simulation of trapped oceanic Kelvin and Rossby wave propagation along the North American west coast was developed to further study the plausibility of this conceptual model. A constant phase speed of 50 cm/s was used for poleward propagating trapped, oceanic coastal Kelvin waves propagating poleward along the west coast of Central and North America. Observed Pacific oceanic Rossby wave phase speeds, from Chelton and Schlax (1996), were selected for use in this simulation (Fig. 9). Results from this simulation, using these component oceanic Kelvin and Rossby wave properties, are illustrated in Fig. 11. These simulation results reveal critical details regarding the organized oceanic Rossby wave crests observed in nature. For example, a localized minimum in the westward propagation distance (between $\sim 15^\circ\text{N}$, $102\text{-}121^\circ\text{W}$) clearly appears between 100-250 days after Kelvin waves propagate poleward of 8°N latitude. This is primarily a result of the orientation of the Central American coastline. This coastal effect results in Rossby waves at higher latitude later time-steps to actually exceed the westward longitudinal position achieved by previously initiated lower latitude Rossby waves. This simulation result is extremely useful in explaining TOPEX/POSEIDON positive SSH anomaly patterns off the west coast of Central America in Fig. 12.

Oceanic Rossby wave crests identified in TOPEX/POSEIDON data, between April 1994-February 1995 and December 1994-May 1996, were tracked from $\sim 14^\circ\text{N}$ to

30°N (Fig. 11). These two dominant wave crests coincided in time, respectively, with the abnormally wet 1994-1996 winter seasons experienced in the Western U.S. The mean, estimated oceanic Kelvin wave phase speeds for the 94/95 winter case was ~ 38.3 cm/s (over a 294 day period), while for the 95/96 winter case, it was estimated to be ~ 51.0 cm/s (over a 392 day period). The combined, mean oceanic Kelvin phase speed for both cases was ~ 44.6 cm/s (within 12% of the assumed Kelvin phase speed of 50 cm/s, selected for use in the Fig. 11 simulation result). The estimated phase speed differences may possibly be attributed to a change of the initial phasing of the two analyzed oceanic Kelvin/Rossby wave events (an ~ 13 cm/s difference in 1994-1996 freely propagating oceanic Kelvin wave phase speed would require substantial changes in ocean density structure). Regardless of the specific reason for the differences, this raises critical questions concerning the seasonal positioning of dominant oceanic Rossby wave crests.

Did more southeastward positioning of estimated 94/95 winter season dominant oceanic Rossby wave front (the 350-400 day Rossby crest solution in Fig. 11) result in selective aiming of more preferential, heavy precipitation storm tracks into southern California? Similarly, in the following 95/96 winter season, did a more northwestward positioned dominant oceanic Rossby wave crest (the 500-600 day Rossby crest solution in Fig. 11) provide sufficient forcing to result in more westward positioned, transient "pineapple express" storm tracks to intersect the North American west coast further north, toward Oregon and Washington (Figs. 2-3, and 8)? A multitude of other questions

remains regarding these simulation results, in conjunction with our increasing empirical knowledge of ocean science via observational data from sources such as the TOPEX/POSEIDON project. For example, if the conceptual model proposed here is realized in nature, what are the implications regarding precipitation/storm track re-adjustment 2-6 years following the demise of a significant El Niño event in the eastern Pacific (for example, in the 800-2400 day Rossby crest solutions in Fig. 11)? Without the oceanic Kelvin-Rossby wave conceptual model component described above (Figs. 8 and 11), it is difficult to make concise analytic inferences regarding the origins of observed sea surface anomaly patterns, and potentially our west coast heavy winter precipitation patterns, too.

VI Conclusion

Our understanding of large to local scale coupled ocean-atmosphere processes is far from complete, but knowledge is accumulating, for example, from studies such as those conducted by the Climate Diagnostic Center to examine the role of air-sea interactions in climate and climate change. Results from one of these studies illustrate the significant upper-level ocean (surface to 400 m) thermal structure interannual to decadal variability that is known to occur in the central North Pacific (Fig. 13). Westward propagating oceanic Rossby waves, capable of significantly altering thermal properties in this slab of surface layer water, have now been observed throughout much of the world's oceans (Schlax and Chelton 1994). The slow propagation speeds of midlatitude oceanic Rossby waves, in conjunction with associated SST anomalies, and previously discussed

inferences concerning possible resultant atmospheric responses, suggest that a significant, and difficult to forecast relationship between oceanic Kelvin/Rossby waves and North American west coast weather patterns may exist. This relationship may be a critical component leading to unusual interannual west coast weather patterns. However, the results of this analysis are not conclusive, and do not constitute absolute proof of the hypothesized links between oceanic Rossby waves and atmospheric processes.

From preliminary inspection, oceanic Rossby wave forcing of the atmosphere appears unable to produce continuous storm track modification, either intraseasonally (time scales of 10-100 days) or interannually. Other dominant, larger scale synoptic-hemispheric scale processes often diminish the signal level of oceanic Rossby wave forcing (Trenberth 1997). However, if the associative links suggested in this Technical Memorandum (ie., significant, positive SST anomalies with geographical patterns organized by oceanic Kelvin/Rossby wave processes – intersecting upstream trans-Pacific jet stream tracks) reach a critical feedback threshold, then it may be possible to make the following inferences. Meridionally skewed crests and troughs of dominant warm core, eastern Pacific oceanic Rossby waves near the west coast of the United States, particularly following El Niño events transitioning into La Niña episodes, may be highly correlated with the geographical distribution and intensity of above normal, west coast winter precipitation (Fig. 8). Thus, recognition of this ocean system process, occasionally resulting in transient resonant, phase-

locking with atmospheric processes, may significantly improve medium-long range forecasting accuracy of heavy seasonal west coast precipitation interception. For example, the mean 1996 Pacific SSH anomaly pattern (Fig. 14) has evident correlation with placement of significantly above normal December 1996 precipitation (Fig. 15). Conversely, drought forecasts for the west coast may also be improved – by anticipating the development of prolonged high pressure ridging over Washington, Oregon, and California. This drought scenario may result as the SST patterns associated with oceanic Rossby waves steer above-normal precipitation into British Columbia and Alaska (refer to Fig. 11 wave crest and trough locations after 800 days). Or, perhaps west coast scenarios suggestive of below-normal precipitation may also be correlated with cold SST patterns associated with oceanic Rossby wave troughs (Fig. 16). For example, Fig. 17 places below-normal precipitation over Washington and northern Oregon, aligned along the upstream orientation of a dominant oceanic Rossby wave trough in Fig. 16.

Regarding the consideration of oceanic Rossby waves in west coast drought forecasting, mid Pacific model results imply oceanic Rossby waves of low-latitude origin in association with El Niño events influence the location of the Kuroshio Extension between 35-40°N (Jacobs et al. 1994). However, the role of larger scale ocean-atmosphere processes – such as the eastward extension of a warm northerly Kuroshio return current – may be more important to North American west coast forecasting. The cumulative potential west coast meteorological and hydrologic scenario

results postulated in the previous paragraph nonetheless suggest that consideration of pronounced eastern Pacific oceanic Rossby wave activity may be warranted. Organized oceanic Rossby wave thermal forcing of the atmosphere appears to be more of a microscale influence, but over a sufficiently large and consistent subregion. Repetitive tracking of storms following northward along the eastern U.S. are often linked to the baroclinic zone established along the coast, defined by contrasting cold continental air over run by warmer, moist marine air masses advected over the Gulf Stream (Carlson 1991). Although not energetically capable of producing similar extreme contrasts, eastern Pacific oceanic Rossby waves may be instrumental in establishing a much weaker yet still analogous baroclinic atmospheric tracking zone. Using this conceptual model, forecasters may be more able to make large-scale process inferences in sparse and/or remote data sets (Figs. 14-17). What other existing, coupled conceptual models do forecasters presently use to explain the patterns in Figs. 14 - 17? Keeping in mind the billions of dollars in west coast flood damage between 1994-1996, it seems prudent to consider the potential association/s proposed in the multifaceted conceptual model introduced above?

If this conceptual model is considered a potential mechanism to partially explain variations in heavy west coast precipitation patterns, how do we systematically make the necessary associations suggested in this Technical Memorandum? Slow oceanic Rossby wave propagation speeds (on the order of 10 cm/s) facilitate interrupted analysis and tracking, for example, when other

operations take prolonged precedence, and/or relevant analysis data availability is temporarily limited. Observing the presence of dominant oceanic Rossby waves may entail a consistent, periodic analysis of remotely sensed eastern Pacific SST and SSH anomalies, in conjunction with possibly correlated DMSP microwave precipitation and preferential GOESS-9 water vapor moisture advection tracks from low to mid latitudes. Drifting buoy data is another source of supplementary, in situ SST anomaly observations to correlate with SSH data.

This integrated approach is probably the only presently available operational technique to use in conceptually identifying the potential occurrence of pronounced oceanic Rossby wave activity at the field office level. Because of oceanic Rossby wave process subtleties, development of a working conceptual model that incorporates key evolution and structure details will be necessary before predictive skills can be evolved (McGinley 1986).

TOPEX/POSEIDON altimeter data capable of resolving oceanic Kelvin/Rossby waves has only been available since October 1992 (Fu et al. 1994). This initial date of TOPEX/POSEIDON data availability occurred in the midst of the last prolonged El Niño episode, extending over a 5 year period between 1990-1994 (Halpert et al. 1995). As a result, we still have much to learn about what this new type of data can tell us regarding ocean-atmosphere interactions, both preceding and following El Niño episodes. Many questions remain to be answered regarding the relevancy of this forecast problem. For example, are

oceanic Kelvin/Rossby waves deterministic? Do the effects of zonal atmospheric Walker circulations over the Pacific play a more critical role than oceanic Rossby and Kelvin waves in selectively altering SSTs (Liu 1997)? Yet, as Trenberth (1997) suggests, a key step in predicting interannual ENSO cycle onset and evolution entails antecedent identification of how oceanic Kelvin and Rossby waves redistribute warm, tropical water, in concert with atmospheric wind forcing.

Answers to the plethora of remaining questions will be discovered as we cumulatively increase our understanding over the full range of significant ocean-atmosphere feed-back mechanisms occurring across the Pacific Ocean, as well as over the global scale ocean-atmosphere system. Hopefully, the analysis discussed in this Technical Memorandum demonstrates that our understanding of nonclassic North American west coast applied meteorology can be advanced significantly via forecaster's conceptual recognition, and reporting of potentially significant, coupled, upstream-offshore feedback processes. Conversely, it is clear that identification of these complex earth system processes will necessitate an interdisciplinary approach, both in terms of professional training (Mass 1996), and operational interaction. This type of approach to improving North American west coast forecasting will accelerate the conceptual intersection of valuable functional knowledge in, but by no means limited to, oceanography, geodetic and thermographic remote sensing, terrestrial hydrologic implications, as well as the atmospheric sciences. However, in this endeavor, it is also evident, from

previously detected systematic satellite observation data errors, that remote sensing of subtle but critical ocean-atmosphere interactions will require ground-truth calibration with, for example, long term in situ buoy data.

VII Acknowledgments

Vital support included numerous data sets made available via the Internet by: 1) the Jet Propulsion Laboratory's TOPEX/POSEIDON Global Ocean-Monitoring Mission;¹ 2) the NOAA/ERL Climate Diagnostics Center (CDC);² 3) as well as the U.S. Department of Commerce, NOAA, Climate Prediction Center (CPC);³ and 4) the NOAA Laboratory for Satellite Altimetry. The CPC Internet posting of Halpert et al. (1996)⁴ proved to be extremely valuable throughout this project. Mr. Charles Orwig, a NWS Northwest River Forecast Center (NWRFC) Senior Hydrologist, provided much appreciated encouragement throughout this research effort, and contributed to the final review and editing of this manuscript. Special thanks to Steven M. Taylor, from the U.S. Naval Postgraduate School, for generating the May 1996 Pacific Ocean surface

¹ At <http://topex-www.jpl.nasa.gov/>

² At http://www.cdc.noaa.gov:80/~map/maproom/text/climat_products.shtml

³ At http://wrcc.sage.dri.edu/recent_climate.html

⁴ At http://nic.fb4.noaa.gov:80/products/assessments/assess_95

current plot. Regards to Ryan Smith, from NOAA's Global Drifter Center,⁵ for providing selected Pacific drifting buoy data plots; and to Bob Cheney, Chief of NOAA's Geosciences Laboratory Satellite and Ocean Dynamics Branch,⁶ for answering numerous questions regarding oceanic Rossby waves. The GMS image of Typhoon Bart was made available by Taikan Oki, from the Institute of Industrial Science, University of Tokyo, Japan. The author is grateful to the numerous other unnamed individuals who contributed to this analysis (primarily via the Internet). Critical feedback regarding this research was obtained at the February 1997 Pacific Northwest Weather Workshop, and the August 1997 First Pacific Rim Hydroclimate Workshop; support essential to attend these workshops was made possible by the efforts of Dr. Thomas Potter, Director, NWS Western Region Headquarters, Donald Laurine, NWRFC Developmental and Operations Hydrologist, and Andy Edman, Scientific Services Division, NWS Western Region Headquarters. Invaluable reference materials supporting this research were obtained from the NCAR Foothills Laboratory Library, Boulder, Colorado, and the NOAA Central Library, Silver Spring, Maryland (email reference@nodc.noaa.gov).

⁵ At

<http://www.aoml.noaa.gov/phod/dac/dacdata.html>

⁶ At

<http://ibis.grdl.noaa.gov/SAT/SAT.html>

VIII References

- Anderson, D.L.T., K.Bryan, A.E. Gill, and C. Pacanowski, 1979: The Transient Response of the North Atlantic: Some Model Studies. *J. Geophys. Res.*, **84**, 4795-4815.
- Anderson, D.L.T., and A.E. Gill, 1975: Spin-up of a stratified ocean, with applications to upwelling. *Deep-Sea Res.*, **22**, 583-595.
- Barnett, T.P., L. Latif, E. Kirk, and E. Roeckner, 1991: On ENSO Physics. *J. Clim.*, **4**, 487-515.
- Battisti, D.S., and E.S. Sarachik, 1995: Understanding and predicting ENSO. *Rev. Geophys.*, **33** (suppl.), 1367-1376.
- Bocheng L., and Q. Liao, 1988: The Variations in Current System[s] in the Western Tropical Pacific and Its Relations[hip] to the Wind Field. *Proceedings of US-PRC International TOGA Symposium*, Beijing, China, 33-39.
- Carlson, T.N., 1991: *Mid-Latitude Weather Systems*. HarperCollins Academic, 507 pp.
- Chelton, D.B., and M.G. Schlax, 1996: Global Observations of Oceanic Rossby Waves. *Science*, **272**, 234-238.
- Chen T-C., J-M. Chen, and C.K. Wikle, 1996: Interdecadal Variation in U.S. Pacific Coast Precipitation over the Past Four Decades. *Bull. Amer. Meteor. Soc.*, **77** (6), 1197-1205.

- Dayong J., Y. Meiyu, and Min wei, 1988: A GCM Study on the Response of the Atmosphere Surface Wind Fields to Tropical SST Anomalies. *Proceedings of US-PRC International TOGA Symposium*, Beijing, China, 433-439.
- Dorman, C.E., 1985: Evidence of Kelvin Waves in California's Marine Layer and Related Eddy Generation. *M. Weather Rev.*, **113** (5), 73-85.
- Fu L-L., E.J. Christensen, C.A. Yamarone Jr., M. Lefebvre, Y. Ménard, M. Dorrer, and P. Escudier, 1994: TOPEX/POSEIDON mission overview. *J. Geophys. Res.*, **99**, 24369-24381.
- Fu L-L., and R.E. Cheney, 1996: Application of satellite altimetry to ocean circulation studies: 1987-1994. *Rev. Geophys.*, **77** (suppl.), 213-223.
- Fu L-L., and R.D. Smith, 1996: Global Ocean Circulation from Satellite Altimetry and High-Resolution Computer Simulation. *Bull. Amer. Meteor. Soc.*, **77** (11), 2625-2636.
- Halpert, M.S., G.D. Bell, V.E. Kousky, and C.F. Ropelewski, 1996: Climate Assessment for 1995. *Bull. Amer. Meteor. Soc.*, **77** (5), S1-S44.
- Hansen, D.V., 1988: Surface Current and Temperature Patterns in the Western Tropical Pacific. *Proceedings of US-PRC International TOGA Symposium*, Beijing, China, 121-127.
- Heburn, G.W., and R.C. Rhodes, 1987: Numerical simulations of wind forced seasonal and interannual variability of the transport through the Caribbean Sea. *Trans. Amer. Geophys. Union*, **68**, 337 pp.
- Hendon, H.H., and M.L. Salby, 1996: Planetary-Scale Circulations Forced by Intraseasonal Variations of Observed Convection. *J. Atmos. Sci.*, **53**, 1751-1758.
- Hobbs, P.V., J.D. Locatelli, and J.E. Martin, 1996: A New Conceptual Model for Cyclones Generated in the Lee of the Rocky Mountains. *Bull. Amer. Meteor. Soc.*, **77** (6), 1169-1178.
- Hogan T. F., and T.E. Rosmond, 1991: The Description of the Navy Operational Atmospheric Prediction System's Spectral Forecast Model. *Mon. Wea. Rev.*, **119**, 1786-1815.
- Huyer, A., 1980: The offshore structure and subsurface expression of sea level variations off Peru, 1976-1977. *J. Phys. Oceanogr.*, **10**, 1755-1768.
- Ingmanson, D.E., W.J. Wallace, 1979: *Oceanography: An Introduction*. Wadsworth Pub. Co., 357 pp.
- Jacobs, G.A. et al., 1994: Decade-scale trans-Pacific propagation and warming effects of an El Niño anomaly. *Nature*, **370**, 360-370.
- Johnson, S.J., and Coauthors, 1992: Irregular glacial interstadials recorded in a new Greenland ice core. *Nature*, **359**, 311-312.

- Lee, W.-J., and M. Mak, 1996: The Role of Orography in the Dynamics of Storm Tracks, *J. Atmos. Sci.*, **53**, 1737-1750.
- Liu, Z., 1996: Oceanic Regulation of the Atmospheric Walker Circulation. *Bull. Amer. Meteor. Soc.*, **78** (3), 407-411.
- Mass, C.F., 1996: Are We Graduating Too Many Atmospheric Scientists?, *Bull. Amer. Meteor. Soc.*, **77** (6), 1255-1267.
- McGinley, J., 1986: Nowcasting Mesoscale Phenomena. *Mesoscale Meteorology and Forecasting*, Edited by Ray, P.S., American Meteorological Society, 793 pp.
- Min wei, and J. Dayong, 1988: The Characteristics of Solar Radiation and Energy Fluxes over the Western Pacific Ocean: Observations of US-PRC Cruises 2 and 4. *Proceedings of US-PRC International TOGA Symposium*, Beijing, China, 237-250.
- Neelin, J.D., M. Latif., and F.-F. Jin, 1994: Dynamics of Coupled Ocean-Atmosphere Models: The Tropical Problem, *Rev. Fluid Mech.*, **26**, 617-659.
- Nuss, W., 1988: Boundary Layer Interaction in Oceanic Midlatitude Cyclones. *The 7th Conf. On Ocean-Atmos. Interaction*, Anaheim, CA., Amer. Meteor. Soc., 216-217.
- Olson D.A., N.W. Junker, and B. Korty, 1995: Evaluation of 33 Years of Quantitative Precipitation Forecasting at the NMC. *Wea. Forecasting.*, **10** (3), 498-511.
- Palmén, E., and C.W. Newton, 1969: *Atmospheric Circulation Systems*. Academic Press, 603 pp.
- Parke, M.E., D.L. Stewart, D.L. Farless, and D.E. Cartwright, 1987: On the Choice of Orbits for an Altimetric Satellite to Study Ocean Circulation and Tides. *J. Geophys. Res.*, **92**, 11693-11707.
- Pedlosky, J., 1979: *Geophysical Fluid Dynamics*. Springer-Verlag, 624 pp.
- Qiu, B., 1995: Variability and energetics of the Kuroshio Extension and its recirculation gyre from the first two-year TOPEX data. *J. Phys. Oceanogr.*, **25**, 1827-1842.
- Schlag, M.G., and D.B. Chelton, 1994: Aliased tidal errors in TOPEX/POSEIDON sea surface height data, *J. Geophys. Res.*, **99**, 24761-24775.
- Smith, R.L., 1978: Poleward propagating perturbations in currents and sea levels along the Peru coast. *J. Phys. Oceanogr.*, **83**, 6083-6092.
- Stewart, R.H., 1985: *Methods of Satellite Oceanography*. University of California Press, 360 pp.
- Stull, R.B., 1988: An Introduction to Boundary Layer Meteorology. Kluwer Academic Publishers, 666 pp.
- Trenberth, K.E., 1997: Short-Term Climate Variations: Recent Accomplishments and Issues for Future Progress. *Bull. Amer. Meteor. Soc.*, **78** (6), 1081-1096.




Fig. 1. GMS image of Typhoon Bart recurving northeastward off the northeast coast of Luzon on May 15, 1996, with upper level storm outflow entraining into the mid-latitude westerlies (image courtesy of the Institute of Industrial Science, University of Tokyo, Japan).

PERCENT OF NORMAL PRECIPITATION

1994/1995 Wet Season

October 1994 - April 1995




Fig. 2. Concentration of 200% above normal Winter 1994-95 wet season precipitation anomalies focused over southern coastal California (from Halpert et al. 1996).




Fig. 3. November-December Winter 1995 ≥ 100 mm precipitation anomalies over Northwestern Oregon and Western Washington, using departures from 1961-90 base period means (from Halpert et al. 1996).




Fig. 4. Composite northern hemisphere SST and precipitation anomalies positionally correlated, and indicating the presence of a Spring-Fall 1995 west-northwest shifting positive anomaly structure in the eastern Pacific (from Halpert et al. 1996).




Fig. 5. The United States Navy Fleet Numerical Oceanographic Center (FNOOC) has oceanographic programs dependent on the Navy Operational Global Atmospheric Prediction System (NOGAPS), and one of these programs provides ocean current predictions (Hogan and Rosmond 1991; Heburn and Rhodes 1987). The plot displayed above is a NOGAPS surface Pacific Ocean currents analysis for May 1996 (courtesy of Steven M. Taylor, from the U.S. Naval Postgraduate School, Monterey, California).




Fig. 6. Ten days of consecutive TOPEX/POSEIDON data are used to generate each of the sea surface height anomaly images included in this paper (images originally produced at the Jet Propulsion Laboratory (JPL) in Pasadena, California using software codeveloped with the University of Colorado, Boulder). Time shown at the top of this image, in a “yy/mm/dd” format, represents the “center-time” of a 10 day observation blend periods. The center time of this image was April 22, 1996. The dominant, positive eastern Pacific SSH anomalies have axial symmetry similar to positive SST anomalies in Fig. 4. This symmetry reelected across the equator results from the intersection of individual dominant oceanic Rossby waves along an organized Rossby wave front.




Fig. 7. TOPEX/POSEIDON satellite and orbit placement (originally a JPL image, modified to incorporate subsidiary information regarding functionality of the satellite).




Fig. 8. Proposed conceptual model. Several details are exaggerated to separate features for illustrative clarity. For example, oceanic Kelvin waves actually slope directly away from the west coast in a non-gaussian manner. There is also more of a continuous transfer process than illustrated, as Kelvin waves dissipate offshore and transition directly to oceanic Rossby waves.

Observed Oceanic Pacific Rossby Wave Phase Speed, from Chelton and Schlax (1996)




Fig. 9. Observed northern hemisphere Pacific oceanic Rossby wave phase speeds (cm/s), from Chelton and Schlax (1996).




Fig. 10. December 1995 mean tropospheric temperature anomalies. Positive temperature anomalies, derived from microwave sounding data, occur along a track similar to eastern Pacific oceanic Rossby wave fronts (from Halpert et al. 1996; original data contributed by the University of Alabama at Huntsville).

MODELED NE PACIFIC OCEANIC ROSSBY WAVE PROGRESSION




Fig. 11. Oceanic Rossby/Kelvin wave propagation simulation results.




Fig. 12. Relative Pacific Ocean Rossby wave front speed minima off the coast of Central America evident in TOPEX/POSEIDON February 2, 1995 data (provided by JPL).




Fig. 13. Upper ocean (surface - 400 m) interannual-decadal SST anomaly variability observed in the eastern Pacific between 30N-42N, 176W-144W (courtesy of the National Climatic Data Center (NCDC)).

T/P Sea Level Anomaly Year 96



NOAA / Laboratory for Satellite Altimetry





Fig. 14. Mean SSH anomalies for 1996 indicating presence of organized oceanic Rossby wave fronts across the Pacific in the northern and southern hemispheres (provided by the NOAA Laboratory for Satellite Altimetry).

Percent of Normal Precipitation, December, 1996



Percent of Normal Precipitation




Fig. 15. December 1996 winter west coast precipitation anomalies, with significant coincidence of oceanic Rossby wave front alignment in fig. 14 to the > 300 percent above normal precipitation track reaching into southwestern Montana (courtesy of NCDC).




Fig. 16. 10 day mean TOPEX/POSEIDON SSH anomalies centered around January 15, 1993, during the 1992-93 winter season, with relative oceanic Rossby wave front trough evident in the eastern Pacific (image provided by JPL).

PRECIPITATION ANOMALIES (MM)
BASE PERIOD 1951-1980





Fig. 17. Winter 1992-93 precipitation anomalies, with generally below normal precipitation indicated over a majority of the Pacific Northwest (provided by NCDC).

DRIFTER DATA ASSEMBLY CENTER

DEC 30, 1996

by Swenson

12/30/96



-3.0 -1.5 -0.5 0.5 1.5 3.0

SST anomaly [Reynolds & Smith, 1995]

- △ = REGULAR BUOY
- = BAROMETER
- = ACML-SPAP
- ▽ = SALINITY
- ▽ = BODEGA
- = BAR/WHO
- = CANADIAN EXP (M71)
- (UNFILLED SYMBOL = (UN)DROGLED BUOY)

Fig. 18. December 30, 1996 SST anomaly data (courtesy of NOAA's Drifter [Buoy] Data Assembly Center). A sub-region of eastern Pacific 1.5 to 3.0 C SST anomalies is highlighted in yellow.




Fig. 19. Satellite derived SST anomaly satellite data for the period of September 22-28, 1996 (courtesy of NCDC). Inferred eastern Pacific oceanic Rossby wave fronts, north and south of the equator, along the dashed white lines.

- Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033)
- 152 Climate of Salt Lake City, Utah. William J. Alder, Sean T. Buchanan, William Cope (Retired), James A. Cisco, Craig C. Schmidt, Alexander R. Smith (Retired), Wilbur E. Figgins (Retired), February 1998 - Seventh Revision (PB98-130727)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intra-seasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82 195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82-158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82-177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidli and Austin Jamison, April 1969 (Revised July 1996). (PB96-191614)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteris, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85-110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85-132694)
- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 1878887AS)
- 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lusk, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project. Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lusk, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Radid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
- 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
- 199 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Lusk, April 1987. (PB87 185229/AS)
- 200 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke, September 1987. (PB88 110895/AS)
- 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson George Clark, September 1987. (PB88 114038/AS)
- 202 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88-101935/AS)
- 203 An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern Califor John P. Monteverdi and Scott A. Braun, October 1988. (PB89 121297/AS)
- 204 Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Car Scott, November 1988. (PB89 128649/AS)
- 205 Forecast Guidelines For Fire Weather and Forecasters - How Nighttime Humidity Affects Wild: Fuels. David W. Goens, February 1989. (PB89 162549/AS)
- 206 A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarte Scientific Services Division, August 1989. (PB89 230833/AS)
- 207 The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, J 1990. (PB90-240268)
- 208 Meteorological Factors Contributing to the Canyon Creek Fire Blowup, September 6 and 7, 19 David W. Goens, June 1990. (PB90-245085)
- 209 Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow Whitlat December 1990. (PB91-129239)
- 210 Hydrotools. Tom Egger, January 1991. (PB91-151787/AS)
- 211 A Northern Utah Soaker. Mark E. Struthwolf, February 1991. (PB91-168716)
- 212 Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 199 (PB91-208439)
- 213 Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Moliner, July 19 (PB91-227405/AS)
- 214 Emergency Operational Meteorological Considerations During an Accidental Release of Hazardo Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424)
- 215 WeatherTools. Tom Egger, October 1991. (PB93-184950)
- 216 Creating MOS Equations for RAWFS Stations Using Digital Model Data. Dennis D. Gettma December 1991. (PB92-131473/AS)
- 217 Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 199 (PB92-196104)
- 218 NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 199 (PB93-146785)
- 219 A Case Study of the Operational Usefulness of the Sharp Workstation in Forecasting Mesocyclone-Induced Cold Sector Tornado Event in California. John P. Monteverdi, March 199 (PB93-178697)
- 220 Climate of Pendleton, Oregon. Claudia Bell, August 1993. (PB93-227536)
- 221 Utilization of the Bulk Richardson Number, Helicity and Sounding Modification in the Assessme of the Severe Convective Storms of 3 August 1992. Eric C. Evenson, September 199 (PB94-131943)
- 222 Convective and Rotational Parameters Associated with Three Tornado Episodes in Northern a Central California. John P. Monteverdi and John Quadros, September 1993. (PB94-131943)
- 223 Climate of San Luis Obispo, California. Gary Ryan, February 1994. (PB94-162062)
- 224 Climate of Wenatchee, Washington. Michael W. McFarland, Roger G. Buckman, and Gregory Matzen, March 1994. (PB94-164308)
- 225 Climate of Santa Barbara, California. Gary Ryan, December 1994. (PB95-173720)
- 226 Climate of Yakima, Washington. Greg DeVoir, David Hogan, and Jay Neher, December 199 (PB95-173688)
- 227 Climate of Knappton, Montana. Chris Maier, December 1994. (PB95-169488)
- 228 Forecasting Minimum Temperatures in the Santa Maria Agricultural District. Wilfred Pi and Pet Felsch, December 1994. (PB95-171088)
- 229 The 10 February 1994 Oroville Tornado-A Case Study. Mike Staudenmaier, Jr., April 199 (PB95-241873)
- 230 Santa Ana Winds and the Fire Outbreak of Fall 1993. Ivory Small, June 1995. (PB95-241865)
- 231 Washington State Tornadoes. Tresté Huse, July 1995. (PB96-107024)
- 232 Fog Climatology at Spokane, Washington. Paul Frisbie, July 1995. (PB96-106604)
- 233 Storm Relative Isentropic Motion Associated with Cold Fronts in Northern Utah. Kevin B. Bake Kathleen A. Hadley, and Lawrence B. Dunn, July 1995. (PB96-106596)
- 234 Some Climatological and Synoptic Aspects of Severe Weather Development in the Northwest United States. Eric C. Evenson and Robert H. Johns, October 1995. (PB96-112958)
- 235 Climate of Las Vegas, Nevada. Paul H. Skrbac and Scott Cordero, December 199 (PB96-135553)
- 236 Climate of Astoria, Oregon. Mark A. McInerney, January 1996.
- 237 The 6 July 1995 Severe Weather Events in the Northwestern United States: Recent Examples of SSWEs. Eric C. Evenson, April 1996.
- 238 Significant Weather Patterns Affecting West Central Montana. Joe Lester, May 1996 (PB96-178751)
- 239 Climate of Portland, Oregon. Clinton C. D. Rockey, May 1996. (PB96-17603)
- 240 Downslop Winds of Santa Barbara, CA. Gary Ryan, July 1996. (PB96-191697)
- 241 Operational Applications of the Real-time National Lightning Detection Network Data at the NWS(Tucson, AZ. Darren McCollum, David Bright, Jim Meyer, and John Glueck, September 199 (PB97-108450)
- 242 Climate of Pocatello, Idaho. Joe Heim, October 1996. (PB97-114540)
- 243 Climate of Great Falls, Montana. Matt Jackson and D. C. Williamson, December 1996. (PB97 126684)
- 244 WSR-88D VAD Wind Profile Data Influenced by Bird Migration over the Southwest United States Jesus A. Haro, January 1997. (PB97-135263)
- 245 Climatology of Cape for Eastern Montana and Northern Wyoming. Heath Hockenberry and Keit Meier, January 1997. (PB97-133425)
- 246 A Western Region Guide to the Eta-29 Model. Mike Staudenmaier, Jr., March 1997. (PB97 144075)
- 247 The Northeast Nevada Climate Book. Edwin C. Clark, March 1997. (First Revision - January 199 - Andrew S. Gorelow and Edwin C. Clark - PB98-123250)
- 248 Climate of Eugene, Oregon. Clinton C. D. Rockey, April 1997. (PB97-155303)
- 249 Climate of Tucson, Arizona. John R. Glueck, October 1997.
- 250 Northwest Oregon Daily Extremes and Normals. Clinton C. D. Rockey, October 1997.
- 251 A Composite Study Examining Five Heavy Snowfall Patterns for South-Central Montana. Jonathan D. Van Ausdall and Thomas W. Humphrey, February 1998. (PB98-125255)
- 252 Climate of Eureka, California. Alan H. Puffer, February 1998. (PB98-130735)

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS---Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS -- Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161