



**NOAA TECHNICAL MEMORANDUM
NWS WR-282**

HAZARDOUS WEATHER CLIMATOLOGY FOR ARIZONA

by:
Craig Shoemaker
Jeffrey T. Davis

February, 2008

And is approved for publication by
Scientific Services Division
Western Region

Andy Edman, Chief
Scientific Services Division
Salt Lake City, UT

Hazardous Weather Climatology for Arizona

By

Craig Shoemaker
and
Jeffrey T. Davis

National Weather Service
Tucson, Arizona

February 28, 2008

Table of Contents

List of Figures	v
I. Introduction	1
II. Methodology	1
A. Data	1
B. Data Biases and Discrepancies	1
C. Climate Divisions.....	2
III. Overview of Results	2
A. Role of Thunderstorms in Hazardous Weather across Arizona.....	2
B. Ranking of Hazardous Weather in Arizona	3
IV. Severe Weather Events.....	3
A. Tornadoes.....	4
B. Hail.....	5
C. Wind.....	6
V. Significant Weather Events	7
A. Flash Floods and Floods	7
B. Dust Storms.....	8
C. Lightning	8
D. Dust devils and Gustnadoes	9
VI. Other Weather Related Events	9
A. Drought	9
B. Wildfires	10
C. Snow and Ice	10
VII. Summary	11
VIII. References	13

List of Figures

- Figure 1:** Example of *Storm Data*, historical, and spotter reports skewed toward cities and major highways.
- Figure 2:** Example of the *Storm Data* discrepancy for the 1970 Labor Day Weekend storm. No deaths were reported in the *Storm Data* publication for this event.
- Figure 3:** Arizona climate regions as defined by Sellers and Hill (1974)
- Figure 4:** Injures and fatalities in Arizona sub-divided by weather event for the period of 1955 to 2004
- Figure 5:** Total tornadoes in each county from 1950 to 2004
- Figure 6:** Mogollon Rim Convergence Zone (MRCZ)
- Figure 7:** Tornado frequency by F scale from 1950 to 2004
- Figure 8:** Tornado frequency by month and F scale from 1950 to 2004
- Figure 9:** Tornado frequency by month from 1950 to 2004
- Figure 10:** Tornado frequency by month and climate region from 1950 to 2004
- Figure 11:** Tornado frequency by hour from 1950 to 2004
- Figure 12:** Secondary peak in the annual tornado frequency for the Southwest and Northeast climate regions for the period of 1950 to 2004
- Figure 13:** The percentage of hail events by County from 1955 to 2004
- Figure 14:** Hail frequency by month and size from 1955 to 2004
- Figure 15:** Hail frequency by hour from 1955 to 2004
- Figure 16:** Total thunderstorm wind events by climate region from 1955 to 2004
- Figure 17:** Thunderstorm wind frequency by month from 1955 to 2004
- Figure 18:** Thunderstorm wind frequency by hour from 1955 to 2004
- Figure 19:** Thunderstorm wind frequency by hour and climate region from 1955 to 2004
- Figure 20:** Flash flood and flood frequency by month from 1955 to 2004
- Figure 21:** Flash flood and flood frequency by hour from 1955 to 2004
- Figure 22:** Flash flood and flood frequency by month and climate region from 1955 to 2004
- Figure 23:** The areal extent of some historically significant floods in Arizona
- Figure 24:** Tropical system tracks over the southwestern U.S. and eastern Pacific from 1958-2003
- Figure 25:** Frequency of dust storm fatalities by month from 1955 to 2004
- Figure 26:** Average number of thunderstorm days per year for the United States
- Figure 27:** The ten year flash density for the U.S from 1996-2005
- Figure 28:** Lightning Fatalities and Injuries by month from 1955 to 2004
- Figure 29:** Rate of lightning casualties (deaths and injuries combined) per population ordered by state from 1959 to 1994
- Figure 30:** Rate of lightning fatalities per population ordered by state from 1997 to 2006
- Figure 31:** A close up of the ten year flash density for Arizona from 1996-2005
- Figure 32:** Flash density key

- Figure 33:** Flash densities (flashes per 100 km² per monsoon season) for 20 km x 20 km boxes during all hours for 1985-1990
- Figure 34:** National Palmer Drought Severity Index 1895-1995
- Figure 35:** Drought Extent, Length and Composite Palmer Drought Severity Indexes for Arizona 1950-1995
- Figure 36:** Aerial extent of historical droughts in Arizona
- Figure 37:** Arizona statewide water year precipitation, 1896-2005
- Figure 38:** Wildfire Hazard Areas Map
- Figure 39:** Fire Hazard is based upon factors such as fuel types and condition, natural fire regimes, and topography and the influence of structures.
- Figure 40:** Fire hazard key
- Figure 41:** Mean annual snowfall for Arizona for the period 1961-1990
- Figure 42:** Key for the mean annual snowfall for Arizona for the period 1961-1990
- Figure 43:** Mean annual snowfall days for Arizona for the period 1961-1990
- Figure 44:** Key for the mean annual snowfall days for Arizona for the period 1961-1990
- Figure 45:** Snow and Ice by month from 1955 to 2004

I. Introduction

Providing expedient severe weather warnings to save lives and property is the most important mission provided by National Weather Service offices in Arizona. This study discusses numerous hazardous weather types including tornadoes, hail, wind, flash floods and floods, lightning, dust storms, dust devils, extreme temperatures, drought, wildfires and winter storms. A comprehensive understanding of the state's hazardous weather climatology will better prepare forecasters in anticipating the timing, strength, location and nature of hazardous weather. This climatological information will also be useful to emergency managers and private citizens in understanding the characteristics of severe weather across Arizona. Therefore, the purpose of this study is to quantitatively describe the hazardous weather climatology for Arizona using reports from 1950 to 2004 for tornadoes and 1955 to 2004 for other events.

II. Methodology

A. Data

This paper is a compilation of nearly five decades of meteorological data formed from multiple hazardous weather data sources. Tornadoes, hail, high winds, lightning, flooding, severe winter-weather cases, extreme heat and cold outbreaks are included. Seven sources of information were used to compile the severe weather information:

- National Weather Service's (NWS) Storm Prediction Center (SPC) Database from 1950 to 2004
- National Climatic Data Center's (NCDC) *Storm Data* (NOAA 1955-2004) NCDC Storm Events Database
- National Highway Traffic Safety Administration's (NHTSA) Fatality Analysis Reporting System (FARS) from 1994 to 2004
- University of Arizona Storm Database
- Arizona Department of Health Services' (ADHS) Division of Public Health Services Report on Injury Mortality among Arizona residents (1990-2000)
- Arizona Climate: *The First Hundred Years* (Sellers, Hill and Sanderson-Rae, 1985).

Data analysis using ArcView GIS version 3.1 allowed for the plotting of the above mentioned databases to determine trends of events with regards to topography along with seasonal and diurnal frequencies.

B. Data Biases and Discrepancies

The low population density (45.2 people per sq mi) in Arizona (U.S. Census, 2000) has likely resulted in many severe weather events not being reported, especially those which occur away from the high density population centers of the larger cities in Arizona. Additionally, there is a significant bias to reports along roadways in predominantly rural areas. Therefore, it is important to note that the events from which statistics in this study are generated are actually a subset of all significant events which have occurred in Arizona. Figure 1 illustrates the bias of wind events to population centers and along or near roads.

Many database discrepancies were encountered as the study progressed. One example of this is with the Labor Day weekend storm of 1970, which was a flash flood and flood event caused by tropical storm Norma. The official *Storm Data* report shows no deaths or injuries as depicted in Figure 2. However, the Sellers et al. (1985) report shows 23 deaths caused by the system. Further investigation shows that indeed there were 23 deaths. Yet, another discrepancy can be seen in the event reporting methods used in official *Storm Data* entries over the years. When severe weather occurs on a statewide scale, it is often reported in broad terms (central, southern, northern) instead of the preferred county break down (Pima, Yuma, Maricopa). This is especially the case for events prior to the NWS modernization and associated restructuring in the 1990's. The numerous data sources used in this study have sorted out such discrepancies and lead to a more comprehensive dataset which is summarized in this paper.

C. Climate Divisions

Arizona is the sixth largest state in the United States at just over 114,000 sq mi. and elevations varying from under 300 ft to over 12,000 ft (Sellers and Hill, 1974). This topographical variation over a vast spatial area yields an extreme collection of climate regimes with extraordinarily diverse weather. Therefore, for this study Arizona has been broken down into six climate regions, originally developed by Sellers and Hill (1974). Figure 3 shows a map of the six different climate regions used in this study. These climate divisions better highlight the seasonal and diurnal frequencies that would be normally hidden by the typical county or the NWS County Warning Area (CWA) breakdown.

The Northeast region encompasses the area from the Little Colorado River northeast, including the Painted Desert. The Plateau zone stretches the length of the Mogollon rim from the Kaibab Plateau and Grand Canyon to the White Mountains. The Northwest region covers the Colorado River valley over the northwest part of the state. The Central region includes areas along the southern rise to the Mogollon rim. The Southwest region encompasses the lower desert areas including Phoenix and Yuma. The Southeast region covers eastern Pima and Pinal counties, eastward to the New Mexico border. Each of these regions is affected differently by synoptic and meso-scale systems and the resulting monsoonal and winter-time precipitation.

III. Overview of Results

A. Role of Thunderstorms in Hazardous Weather across Arizona

Thunderstorms are responsible for a large percentage of the hazardous weather observed in Arizona. The thunderstorm season, which is driven by the North American Monsoon, encompasses the majority of annual thunderstorm activity. The North American Monsoon is a regional-scale circulation that develops over southwest North America during the months of July through September. It is associated with a dramatic increase in moisture that occurs over what is normally an arid region of North America, as a shift in the mid-level flow from westerly to easterly, occurs (Adams and Comrie, 1997).

The thunderstorm season in Arizona is fairly well defined, although it varies considerably between northern and southeastern sections of the state due to elevation. The thunderstorm season generally begins over the southeast portion of the state in late June to early July (Tucson's average start date is around July 3rd and Douglas is the last week in June). Elevations

above 5,000 feet generally receive their first thunderstorms a week earlier than the above mentioned dates. A gradual decrease in thunderstorm activity occurs in September and October, with some seasonal variability due to tropical activity. In contrast, across northern Arizona (including Flagstaff) the thunderstorm season typically does not become established until mid or late July. Over central Arizona, including the Plateau region and Mogollon rim, the season usually gets started in early July, with the average at Phoenix around the 7th of July.

Thunderstorms also have different characteristics within the state. Across the higher elevations (above 5,000 feet) hail, generally small, and heavy rains are most prevalent. As the thunderstorms progress off the mountains into valley locations, high winds and dust storms become a major concern. This is a result of the elevation change, which allows for a large amount of the rainfall to evaporate before reaching the surface, causing more momentum as rain-cooled air accelerates to the surface. This process can result in numerous micro bursts as these thunderstorms progress into valleys. The drastically warmer temperatures over the valley areas cause the majority of hail stones to melt before reaching the surface. Flash flooding does occur due to heavy rainfall with valley thunderstorms, especially over urban areas. A great flood threat also results when heavy rain falling at higher elevation flows into lower elevation rivers and washes.

B. Ranking of Hazardous Weather in Arizona

Although thunderstorms produce a large percentage of weather hazards, the greatest hazard in Arizona is extreme temperature which is responsible for the largest number of fatalities. Figure 4 shows the frequency of injuries and fatalities in Arizona by weather type for the period of 1955 to 2004. For the extreme heat/cold category, the highest component is extreme heat. It is very likely that heat is an even greater killer than the data depicts. Untold numbers of illegal immigrant deaths go unreported yearly, so the fatalities may be significantly higher (Keim et al, 2006). Available data is only over a ten year period (1990-2000), making it even more impressive compared to other weather events. Extreme heat will not be examined in later sections of this paper due to the unreliability of casualty statistics before 1990, and the vague cause of death reports associated with heat victims.

Floods and flash floods are the next most dangerous hazards when considering injuries and fatalities, followed closely by dust storms. Tornadoes and thunderstorm winds are responsible for a number of injuries across the state, however only three fatalities due to tornadoes have been reported. The greatest number of fatalities outside of extreme heat occurs from May to September, primarily caused by dust storms, flash floods and lightning. Most injuries occur during the months of May through September, with thunderstorm wind in addition to dust storms, flash floods and lightning as the major contributors to the totals.

IV. Severe Thunderstorm Events

A. Tornadoes

Tornadoes are a more common occurrence in Arizona than most people are aware, primarily over the Central valleys and Plateau regions. For the state, the average of reported tornadoes is approximately 4 per year. Figure 5 shows the distribution of reported tornadoes by county from 1950 to 2004. These statistics reveal that the lower valleys of Maricopa, Pinal and Pima

counties, along with the little Colorado River valley, are well-defined maxima for tornadoes. Tornado development in Arizona is often complicated, since they rarely form under textbook conditions. Instead, many Arizona tornadoes are of the non-supercell type; Arizona tornadoes are not associated with low or mid-level mesocyclones, but are formed by thunderstorms interactions with preexisting boundary layer circulations. An example of this would be the stretching of a preexisting dust-devil, gustnado or lesser circulation into the vertical by a thunderstorms updraft. This causes the circulation to tighten and spin faster, allowing for a brief spin up of tornadic wind speeds. Several documented cases of this type have come to light over the past several years, with the onset of digital cameras and camcorders. One such well documented area for non-supercell tornadoes is the region between the Little Colorado River Valley and the Mogollon Rim, known as the Mogollon Rim Convergence Zone (MRCZ) (Blanchard, 2000). This terrain driven convergence is a region of enhanced thunderstorm development and vertical vorticity. Although, MRCZ has been documented by a recent study (Figure 6), many smaller zones of this type exist across the state. As a result tornado detection and verification is often a distorted practice across the state.

Despite the 196 reported tornadoes in Arizona, only 3 fatalities have been reported since 1950, while 139 injuries were documented. The low number of fatalities can be in part explained by the strength of tornadoes found across the state. Wind speeds in tornadoes vary from 40 mph to 318 mph (F0 to F5 on the Fujita scale). Since the mid 1980's, tornado strength has been measured using the Fujita Scale, or F scale (Abbey and Fujita, 1975). This scale ranks tornadoes by damage intensity, with F0 being the weakest and F5 the strongest. This scale was developed to create a relationship between tornado wind strength and the expected structural damage accompanying this wind. Since there is a highly varying degree of structural integrity of buildings across Arizona, the application of the Fujita scale is difficult to apply. In Figure 7, the reported Arizona tornadoes are listed by F scale frequency, and Figure 8 shows the distribution by month and F scale. Certainly, the strength of many tornadoes is not known, but less than 7% of the tornadoes with a known intensity are in the strong category (F2 or F3). Given this information, a high percentage of reported tornadoes in this state are weak, either F0 or F1.

Figure 9 depicts the monthly frequency of all reported tornadoes and Figure 10 shows the distribution by month and region. Over 95% of tornadic storms occur in May through September, with about 60% occurring in July through September. Tornadoes were reported in all months, F2 tornadoes were mainly recorded May through October, except for one occurrence in January. Figure 11 shows the frequency of tornadoes by hour, with the majority of tornadic storms occurring between 11:00 AM and 11:00 PM MST. The number of tornadoes that occur after dark (8:00 PM MST) is very small, with no reports of tornadoes between 2:00 AM and 7:00 AM MST.

Secondary tornado peaks occur for Southwest and Northeast climate regions as shown in Figure 12. In the Southwest region, this secondary peak occurs from January through March in association with strong late winter storm systems. Over the Northeast region two peaks show up outside of the monsoon season: one from April through June, and the other September and October. To further substantiate these peaks, data from the Four Corners region of New Mexico, Colorado and Utah are included in Figure 12 as well. Over the Northeast Arizona zone

this peak is attributed to the interaction of residual monsoonal moisture with late summer-early fall storm systems (Adams and Comrie, 1997). These transitional events are often characterized by high shear and marginally unstable environments (Blanchard, 2006).

B. Hail

The NWS defines a hail event as severe when large hail is at least 0.75 of an inch in diameter. Hail ranks only as the third most frequent hazardous weather type, but is still responsible for a considerable percentage of property and crop damage. The state averages 12 severe hail events per year. Hail is not a major threat for bodily harm in the state; no fatalities and seven injuries have been reported due to hail. These numbers are the lowest of any severe weather type. High freezing levels tend to diminish the threat of large hail across the state, since a significant amount melting occurs as hail stones fall to the surface. High cloud bases also lead to some sublimation of the stones, further reducing the size seen at ground level (Hsin-mu Lin, et al. 2005). Frequent small hail occurrences are seen at higher elevations, but are generally below severe levels, due to weaker updraft strengths with mountain thunderstorms. The percentage of hail events by county for Arizona is depicted in Figure 13.

A significant bias towards population centers is seen with hail reports. This bias should not be a surprise given the visual conformation needed for hail measurements. In unpopulated areas, either the hail melts before a measurement can be made, or the damage from the melted hail is attributed to another severe weather element such as wind. Although these figures show the highest severe hail frequencies over Maricopa, Pinal, Pima and Coconino counties it is likely many of the lower frequency counties such as Graham and Greenlee counties have an equal or greater severe hail frequency.

Figure 14 shows the number of hail reports by month and by size. Damaging or severe hail (0.75 to 1.50 inches) is most common in June through August, as is very large hail (over 1.50 inches); although a significant number of very large hail reports also occur in September. The size of some hail reports are not known, but were included in *Storm Data* because they caused significant crop and/or property damage or had a significant accumulated depth. These reports are included in this study due to their likelihood of being severe hail. Figure 15 reveals damaging hail is most likely to fall in the afternoon and early evening, as is the case with tornadoes. Just over 70% of the reported severe hail events fell between 2:00 PM and 6:00 PM MST. The largest hail stone reported in the state is 4.50 inches, which is softball size. This occurred in Mayer, Yavapai County, on September 28, 1995 (*Storm Data*).

C. Wind

A wind event is considered severe when speeds of 58 mph or greater are measured or estimated, or produce significant damage. Arizona averages 29 severe wind events per year, with the greatest concentration over the southern most climate regions. The number of reported thunderstorm wind events by climate region is shown in Figure 16. Thunderstorms moving off the higher terrain into the hot, dry desert tend to produce stronger downdrafts due to extreme evaporation rates of precipitation and greater descent distance. Therefore, severe downbursts and outflow boundaries are more common over these regions. Additionally, wet microbursts also occur over these areas, especially during the mid and late monsoon season. However, isolated events can occur with strong baroclinic systems outside of the monsoon season. These

events are typically most common over the west and northwest. Dry microbursts occur when rain falls through an extremely dry atmosphere and evaporates, causing rapid cooling of the air. This air rapidly descends and accelerates as it nears the surface. Very little if any rainfall reaches the ground; instead the cooled air spreads out in the form of strong and occasionally damaging winds. Wet microbursts occur during very heavy precipitation, typically in warm and moist atmospheres. They rely on drag of precipitation to accelerate the air towards the surface. The air then spreads out, as it impacts the ground, in the form of strong and occasionally damaging winds.

Arizona thunderstorm wind events are mainly clustered over a two month period: July and August, shown in Figure 17. This should not be a surprise given this is the peak of the monsoon. The slightly higher occurrence rate in September is predominantly over the first half of the month when the monsoon is generally still active. The diurnal cycle of thunderstorm wind events is clustered from 3:00 PM to 9:00 PM MST, with the peak at 7:00 PM MST as shown in Figure 18. This timeframe is to be expected, given the diurnal nature of monsoonal thunderstorms and their dependence on solar heating. Figure 19 shows that hourly events are not uniform over the state. Instead, severe wind events have clearly defined time ranges for each climate region. The time offsets are due to how daily heating and moisture interact with the complex terrain of the state. This diurnal difference is not surprising since thunderstorms form over the higher terrain earlier in the day. Convection then typically shifts into the valleys and lower desert areas in the late afternoon and evening. The most intense and direct solar heating occurs over the mountains during the morning hours. This differential heating produces sufficient buoyancy for earlier thunderstorm development over the mountains. Intense surface heating during the afternoon, allows convection and accompanying damaging winds to transition into valleys and lower deserts by evening, providing adequate moisture is in place. The highest thunderstorm associated wind speed measured in the state is 175 knots. This occurred in Spring Valley which is in Yavapai County on September 6, 1996 at 11:15AM MST (*Storm Data*).

V. Significant Weather Events

A. Flash Floods and Floods

Flash floods are defined by the NWS as a quick response to heavy rainfall in streams, rivers, or any flood prone area, six hours or less after the heavy rain started. Flood events are defined as longer lasting events, with rivers rising above flood stage over 6 hours to many days. For this study flash floods and floods will be combined, due to the broad grouping of numerous flood events with flash floods before NWS modernization in the mid 1990's. A great number of events prior to modernization are simply recorded as statewide flooding, leaving no way to differentiate between flash and river flooding.

In Arizona, the greatest numbers of flood events during the monsoon season are of the flash flood type. Therefore, flood events occurring during the monsoon period (July-September) will be referred to as flash floods for this study. In Arizona, flooding ranks second to extreme temperatures in causing fatalities. Over the period 1959-1999, Arizona ranked tenth in the nation in flooding deaths per million people (NCDC). In the western United States, the monthly distribution of flood events displays a summer maximum due to thunderstorms, and a

secondary winter maximum (January-March) due to winter storms. The monthly distribution of floods in Arizona is shown in Figure 20, and reveals a strong association with the summertime North American Monsoon, with nearly three fourths of floods occurring from July through September. Unlike much of the western United States, the secondary maximum is much smaller during the winter months. Figure 21 shows the distribution of floods by hour. Most of the flash floods occur between 4:00 PM and 7:00 PM similar to the peak hours of severe hail and tornadoes.

Intense rainfall over a short period of time can result in large amounts of water surging through canyons in the mountains of Arizona; however, the valleys receive numerous flash floods as well. Of particular concern in valley locations are normally dry washes, which become raging rivers in a matter of minutes due to intense rainfall. These dry washes are very common over desert terrain and crisscross many urbanized areas, exacerbating flash flood dangers. In Figure 22, the number of reported flash floods and floods for each climate region and the monthly distribution is shown. Flash flood frequencies are greatest over the Southeast followed by the Southwest climate region. These two regions contain the majority of Arizona's population, with the greatest flash flood occurrences in the Tucson and Phoenix metropolitan areas over the July through September period. The Southeast climate zone has a flash flood frequency nearly equal to all other zones combined for the months of July and August. Finally, slight secondary maxima can be seen over the Northwest, Plateau, Central and Southwest climate zones' January through March due to winter storms. Although, these events tend to be less frequent, they are typically more widespread and quite destructive as they usually involve main stem river flooding. Some historically significant aerial flood events and recurrence intervals are shown in Figure 23.

Concealed in the data is flooding associated with tropical systems. Tropical systems are infrequent over the state, occurring only once every five years, as west coast troughs typically lack the amplitude to pull the systems northward. The low proportion of tropical systems impacting the southwestern United States is illustrated in Figure 24.

However, when west coast troughs have the strength to pull tropical systems north; the effects can be devastating. During these events flash flooding and main stem river flooding often occur simultaneously. The deadliest natural disaster in Arizona's history was Tropical Storm Norma on Labor Day 1970. Flooding from Norma resulted in 23 deaths as reported by Sellers et al. (1985).

B. Dust Storms

Dust storms are often underestimated as significant weather events. They rank only behind the extreme heat/cold and flood categories for causing deaths and injuries. Most of these deaths are due to multi-car pileups on roadways and interstates. In Arizona, several different weather regimes are associated with major dust storms. Brazel and Nickling (1986) identified four of them: (1) frontal, (2) thunderstorm, (3) tropical disturbance, and (4) upper level trough and/or cut-off low.

Early season monsoon thunderstorms are the most notorious for producing deadly dust storms. July is clearly the most deadly dust storm month across the state, with nearly double the

fatalities compared to other months depicted in Figure 25. July is the transition month from dry season (March-June) to monsoon season. Thunderstorms during the first few weeks of the monsoon produce very little rainfall, as much of the lower atmosphere remains dry. Much of the rain with early monsoon season thunderstorms evaporates before reaching the surface. This leads to dry downburst winds, which organize into larger and strong outflows. These strong winds pick up the dry soil, dust and sediments pre-conditioned by the dry season, often producing long lasting dust storms with visibilities near zero. Typically, these severe storms frequently referred to as “haboobs” (hə-'büb), affect the Phoenix metropolitan area and surrounding desert areas during the monsoon season.

C. Lightning

Thunderstorms and lightning are a common occurrence in Arizona, mainly during the summer months. Arizona has an annual thunderstorm frequency comparable to many perceived thunderstorm rich Central Plains states. Total thunderstorm days range from fewer than 10 days along the lower Colorado River valley to 80 days over the White Mountains Figure 26. However, the majority of this thunderstorm activity is compacted into a three-month period (July-September). Similarly, thunderstorm lightning flash density shows a favorable comparison to many Plains States; with a maximum over the Mogollon rim and much of southeast Arizona as shown in Figure 27. Arizona lightning fatalities and injuries fall into this timeframe as well (July-September), with August the peak month as shown in Figure 28. The three month period of February-April along with December is devoid of any lightning casualties. Despite the relatively high flash density, Arizona ranks in the middle of the pack nationally for lightning casualties per capita Figure 29 and Figure 30. A major reason for this is the major population centers of Phoenix and Yuma are in lower flash density areas, unlike much of the rest of the state Figure 31 and Figure 32. Of the major population centers Tucson is located in the most lightning flash dense zone (2-8 flashes/sq. km/yr) as depicted in Figure 33. Still, the 80 fatalities and 200 injuries that have occurred over the state are quite significant.

D. Dust Devils and Gustnadoes

Although some may find it odd and even a bit humorous to include dust devils in a hazardous weather climatology, facts point to a destructive side of the often fair weather, benign vortex. Winds of dust devils can reach up to 70 miles per hour, which is certainly strong enough to do minor to moderate damage. Officially, four dust devil reports are listed in *Storm Data*; unfortunately no dust devils have been listed before 1994 or after 1997. Many dust devils in *Storm Data* are cross listed, combined, or misclassified as dust storms, making it nearly impossible to differentiate between the two. Of the four listed, two caused considerable damage in Tucson.

Dust devils are generally small vortexes not associated with thunderstorms, which are made visible by rotating clouds of dust or debris. The key here is that they are not associated with thunderstorms and more specifically do not extend upward to the cloud base of a thunderstorm. Dust devils form due to strong surface heating during hot weather. Strong heating from the sun causes the surface to become much warmer than air above the surface. Additionally, uneven heating of the surface occurs, certain terrain and vegetation type's heat up more and faster than others. As a result, the warm air has a tendency to rise in bubbles. As these bubbles of warm air

rise, cooler air from above and to the sides of the bubble rushes around and below to fill the void. This inrush of air can happen in a somewhat uneven manner, and as it reaches the center of the void filled by the bubble of rising air, can begin to spin (Renno et al. 1998). The combination of lower air pressure and heating causes the air, dust and debris inside the dust devil to rise. Wind speeds near the centers of dust devils can range from just a few miles per hour to perhaps as much as 70 miles per hour.

In Arizona gustnadoes are probably quite common among the strongest thunderstorm-produced gust fronts. Gustnadoes are short-lived, ground-based, shallow, vortices that develop on a gust front associated with either thunderstorms or showers. They may only extend from 30 to 300 feet above the surface. Again, the key here is there will be no connection with the cloud base. They may be accompanied by rain, but usually are only visible as a debris cloud or dust whirl at or near the ground. Wind speeds can reach 60 to 80 mph, resulting in significant damage, similar to that of a F0 or F1 tornado. However, gustnadoes are not considered tornadoes. Gustnadoes are not associated with storm-scale rotation (they do not extend up into the cloud) that is involved with true tornadoes. They are more likely to be associated visually with a shelf cloud that is found on the forward side of a thunderstorm near strong outflow winds.

VI. Other Weather Related Events

A. Drought

Drought is defined most succinctly as insufficient water to meet needs (Redmond, 2002). Many people are unaware that it is possible to have a drought in the desert; which is an arid environment. Therefore, Arizona droughts seem to take people by surprise, even though they have been worsening for years. The composite Palmer Drought Severity Index (1895-1995 in Figure 34 and 1950-1995 in Figure 35) shows that drought across the state has been steadily worsening over the past several decades. A drought is often looked upon as a long term problem until it affects us in the short term and our everyday life. Figure 36 shows the areal extent of historic droughts and their recurrence interval in years. A federal emergency declaration (allows government aid) was issued for the 1973-77 drought by Federal Emergency Management Agency (FEMA). Figure 37 shows the Arizona statewide water year precipitation from 1896 to 2005. The red highlighted regions are periods of low precipitation. Notice how they correspond favorably to historic droughts. It is difficult to measure state drought effects in monetary or casualty statistics since its effects are subtle, gradual and often linked to some other hazardous event (such as heat and wildfires). However, it is almost certain that when considering a drought's entire duration, its damaging effects are comparable or even greater than other more visible hazardous weather types.

B. Wildfires

Wildfires pose an increasingly significant threat to life, property and the economy of Arizona. High impact wildfires have been far more common over the last decade than previous several decades. In fact, three of the most serious wildfires, Rodeo-Chediski in 2002, Aspen in 2003, and Cave Creek in 2005 have occurred over the past five years. Both the Rodeo-Chediski and Aspen fires were given the dubious distinction of being declared major disasters by FEMA. These are the only two fires to be given this classification in the state's fire history. The increase can be attributed to accumulation of living and dead fuels coupled with extended

drought conditions. Another factor is the population boom of the past several decades as the urban-forest interface merges. Overall, the greatest threat is at higher elevation where fuels are more supportive of extreme fires. Figure 38, Figure 39 and Figure 40 show wildfire hazard severity and risk, notice that the greatest threat is at higher elevation, where fuels are more supportive of extreme fires.

C. Snow and Ice

Arizona winters are highly variable with regards to temperature and snowfall from one year to the next. The impact of any given Arizona winter is highly dependant on the type of storm track that becomes established. Ideally for wet winters, a track from the Pacific Ocean, south to San Diego and across the state develops. Cold, dry winters typically see a storm track along the western Rockies from the Great Basin southeast to New Mexico, although this track can also provide snowfall for northern Arizona. The variability in storm track and subsequent Arizona winters are influenced by ENSO patterns. El Nino is an abnormal warming of the equatorial Pacific Ocean waters and tends to strengthen the southern jet stream. This jet stream develops an orientation from the central Pacific eastward across Arizona. Pacific storm systems and associated moisture follow this track, bringing precipitation, often as heavy snowfall, to the state. Conversely, a La Nina pattern develops when there is abnormal cooling of the ocean waters near the equatorial Pacific. This leads to a northward shift of the jet stream with an orientation over the Northwest United States. Under this pattern Arizona is typically influenced by a ridge of high pressure which blocks moisture and incoming systems leading to warm wintertime temperatures. However, many winters are not easily classified as El Nino or La Nina and show high variability within the winter year itself.

The main time period for winter weather in Arizona is from November through March, with the peak in January and February. The heaviest snow typically occurs along the Mogollon rim where yearly snowfall amounts in excess of 100 inches are common. Figure 41 and Figure 42 shows the yearly snowfall across the state. Notice the extreme snowfall variability, with amounts from zero over the southwest part of the state to up to 200 inches over the Mogollon rim. Figure 43 and Figure 44 depict the mean annual snowfall days over the state, which further illustrates just how different winters are from one part of the state to the next.

Cold temperatures are also quite common over the northern portion of the state and at elevations above 7000 feet in the southeast. In these locations low temperatures fall to well below zero. Exposure to the cold temperatures is a major contributor to the total fatalities statistics in the Heat/Cold hazardous weather category. Cold temperatures often act in unison with heavy snowfall events that can strand and isolate people, allowing exposure to the cold to take hold. This is especially the case for those engaged in outdoor activities from November through March.

Travel in Arizona's diverse terrain can also be hazardous during storms. Many highways that traverse the state see elevation changes of several thousand feet over a short distance. These changes in elevation can lead to rapid changes in visibility and deteriorating road conditions catching motorists off guard. Vehicle accidents during winter storms account for a large percentage of the fatalities that fall in the Snow/Ice hazardous weather category. Figure 45 shows the snow and ice storm frequency by month over the state.

VII. Summary

The results of this study can be summarized in the following:

- The extremely low population density (45.2 people per sq mi) of much of Arizona has likely resulted in many hazardous weather events not being reported, especially those which occur away from population centers. Additionally, there is a significant bias to reports along roadways in rural areas.
- The greatest weather hazard in Arizona is exposure to extreme temperatures (Extreme Heat/Cold category) which is responsible for the largest number of fatalities. The highest component is extreme heat and this is actually an even greater killer than the data depicts.
- Arizona has an annual thunderstorm frequency comparable to many Plains States.
- Tornadoes are a more common occurrence in Arizona than most people are aware, primarily over the central valleys and plateau regions. For the state, the average number of reported tornadoes is approximately 4 per year.
- A high percentage of reported tornadoes in this state are weak, either F0 or F1.
- Hail ranks only as the third most frequent hazardous weather type, but is still responsible for a considerable percentage of property and crop damage. The state averages 12 severe hail events per year.
- Arizona averages 29 severe wind events per year, with the greatest concentration over the southern most climate regions.
- Arizona thunderstorm wind events are mainly clustered over a two month period: July and August.
- In Arizona, flooding ranks second to extreme heat in the number of reported fatalities. Arizona ranks tenth in the nation in flooding deaths per capita. The Southeast climate zone has a flash flood frequency nearly equal to all other zones combined for the months of July and August.
- Dust storms are often underestimated as significant weather events and rank only behind extreme heat/cold and flooding for deaths and injuries.
- Despite a relatively high flash density, Arizona ranks in the middle of the pack nationally for lightning casualties per capita.

- Dust devils in Arizona can have a surprisingly destructive side, with winds up to 70 miles per hour. Dust devils are often cross-listed with dust storms making it hard to obtain accurate information on historically dust devil damage.
- It is difficult to measure state drought effects in monetary or casualty form, since its effects are subtle, gradual, and often linked to some other hazardous event (heat and wildfires).
- High impact wildfires have been far more common over the last decade than previous several decades.
- The main Arizona wintertime hazardous weather aspect is exposure to the cold temperatures and heavy snowfall over the higher terrain of northern and eastern parts of the State.

VIII. References

- Abbey, R. F., Jr., and Fujita, T. T., 1975. Use of tornado path lengths and gradations of damage to assess tornado intensity probabilities, in Preprints, 9th Conference on Severe Local Storms, Norman: American Meteorological Society, Boston, p. 286- 293.
- Adams, D.K. and A.C. Comrie, 1997: The North American Monsoon. *Bull. Amer. Meteor. Soc.*, 78, 2197–2213.
- ADHS' Division of Public Health Services: Injury Mortality among Arizona residents. [Available online at <http://www.azdhs.gov/plan/report/im/imindex.html>]
- Arizona State Land Department: Forestry Division. [Available online at http://www.azsf.az.gov/fire_managment/communities_at_risk/default.asp]
- Blanchard, D.O., 2000: Forecasting Severe Weather along the Mogollon Rim Convergence Zone. *Preprints, 20th Conf. on Severe Local Storms*, Orlando, Meteor. Soc., 563-566.
- _____, 2006: A Cool Season Severe Weather Episode in Northern Arizona. [Available online at <http://ams.confex.com/ams/pdfpapers/115214.pdf>]
- Brazel, A.J. and W.G. Nickling, 1986: The Relationship of Weather Types to Dust Storm Generation in Arizona (1965-1980). *Journal of Climatology*, 6, 255-275.
- CLIMAS Drought Planning and Mitigation. [Available online at <http://www.ispe.arizona.edu/climas/research/drought/Figures/fig1.html>]
- Climate Prediction Center (CPC). Composite Palmer Drought Severity Index 1950-1995 [Available online at <http://www.cpc.noaa.gov/products>]
- Colorado Climate Center, Colorado State University. Mean annual snowfall for the United States for the period 1961-1990. Mean number of days per year with one inch or more of snow on the ground for the period 1961-1990. [Available online at <http://ccc.atmos.colostate.edu/snowbooklet.php#Snmaps>]
- Corbosiero, K. L., The Contribution of Eastern North Pacific Tropical Cyclones to the Warm Season Rainfall Climatology of the Southwest United State [Available online at <http://www.atmos.ucla.edu/~kristen/monsoon/monsoon.html>]
- Curran, E.B., R.L. Holle, and R.E. López, 2000: Lightning Casualties and Damages in the United States from 1959 to 1994. *Journal of Climate*, 13, 3448-3453.

- FEMA: Arizona Disaster History. [Available online at http://www.fema.gov/news/disasters_state.fema?id=4]
- Hsin-mu Lin, Pao K. Wang and Robert E. Schlesinger, Three-dimensional nonhydrostatic simulations of summer thunderstorms in the humid subtropics versus High Plains, *Atmospheric Research* Volume 78, Issues 1-2, , November 2005, Pages 103-145.
- Keim, S.M., M.Z. Mays, B. Parks, E. Pytlak, R.M. Harris, and M.A. Kent, 2006: Estimating the Incidence of Hear-Related Deaths among Immigrants in Pima County, Arizona. *J. of Immigrants and Minority Health*, **8**, No. 2, 185-191.
- National Drought Mitigation Center. National Palmer Drought Severity Index (1895-1995) [Available online at <http://www.drought.unl.edu/whatis/palmer/pdsihist.htm>]
- National Oceanic and Atmospheric Administration, 1959-2004: *Storm Data*. National Climatic Data Center, Asheville, NC, **1-40**, Nos. 1-12.
- NCDC's Storm Events Database. [Available online at <http://www4.ncdc.noaa.gov/cgi-win/wwcgi.dll?wwEvent~Storms>]
- NHTSA's Fatality Analysis Reporting System (FARS). [Available online at <http://www.-fars.nhtsa.dot.gov>]
- NWS' Storm Prediction Center Database. [Available online at <http://www.spc.noaa.gov/wcm>]
- Oklahoma Climatological Survey. Map of average number of thunderstorm days in U.S. [Available online at <http://www.ocs.ou.edu>]
- Paulson, R.W., E.B. Chase, R.S. Roberts, and D.W. Moody, compilers, 1991: National Water Summary 1988-89: Hydrologic Events and Floods and Droughts. U.S. Geological Survey Water-Supply, Paper 2375, 591.
- Redmond, K.T., 2002: The Depiction of Drought: A Commentary. *Bull. Amer. Meteor. Soc.*, **83**, 1143–1147.
- Renno, N. O., M. L. Burkett, and M. P. Larkin, A simple thermodynamical theory for dust devils, *J. Atmos. Sci.*, **55**, 3244– 3252, 1998.
- Sellers, W.D., and R.H. Hill, 1974: *Arizona Climate (1931-1972)*. University of Arizona Press.
- _____, R.H. Hill, and M. Sanderson-Rae, 1985: *Arizona Climate: The first Hundred Years (1885-1985)*. Tucson, University of Arizona.
- University of Arizona's College of Agriculture Arizona Storm Database. [Available online at <http://ag2.calsnet.arizona.edu/egi-bin/storms.egi>]
- United States Census Bureau, 2000: [Available online at <http://www.census.gov>]
- Vaisala NLDN.(Vaisala). Vaisala's Ten year flash density for the U.S. from 1996-2005. Rate of lightning fatalities per population ordered by state from 1997 to 2006 [Available online at <http://www.vaisala.com>]
- Watson, A.I., R.L. Holle, and R.E. Lopez, 1994: Diurnal cloud-to-ground lightning patterns in Arizona during the southwest monsoon. *Mon. Wea. Rev.*, **122**, 1716-1725.




Figure 1. Example of *Storm Data*, historical, and spotter reports skewed toward cities and major highways for wind events.

Arizona Climate Regions

from Sellers and Hill (1974)





Figure 3. Arizona climate regions used in this study as defined by Sellers and Hill (1974).

Arizona Deaths and Injuries by Hazardous Weather Type



Data Sources:

- Arizona Climate- *The First Hundred Years (Sellers, Hill and Sanderson-Rae)*
- NCDC Storm Data 1955-2004
- SPC Events Database 1950-2004
- NCDC Storm Events Database: Online- www4.ncdc.noaa.gov/cgi-win/wwcgi.dll?wwEvent~Storms
- DOT Fatality Analysis Reporting System: Online- www-fars.nhtsa.dot.gov
- University of Arizona Storm Database: Online- ag2.calsnet.arizona.edu/cgi-bin/storms.cgi
- Injury Mortality Among Arizona Residents, 1990-2000 Report (March 2002)


Figure 4. Injures and fatalities in Arizona sub-divided by weather event for the period of 1955 to 2004

Arizona Tornadoes

1950-2004

Number of Tornadoes Reported By County

Apache:	6
Cochise:	12
Coconino:	20
Gila:	3
Graham:	0
Greenlee:	0
La Paz:	0
Mohave:	14
Maricopa:	57
Navajo:	13
Pima:	17
Pinal:	25
Santa Cruz:	0
Yavapai:	19
Yuma:	10



Arizona Averages 4 Tornadoes per Year

Figure 5. Total tornadoes in each county from 1950 to 2004

Mogollon Rim Convergence Zone (MRCZ)

A region of non-supercell tornadoes

Blanchard (2000)




Figure 6. Mogollon Rim Convergence Zone (MRCZ) noted by Blanchard (2000)




Figure 7. Tornado frequency by F scale from 1950 to 2004




Figure 8. Tornado frequency by month and F scale from 1950 to 2004




Figure 9. Tornado frequency by month from 1950 to 2004




Figure 10. Tornado frequency by month and climate region from 1950 to 2004




Figure 11. Tornado frequency by hour (Time MST) from 1950 to 2004

Arizona Tornadoes by Month and Region 1950-2004





Figure 12. Secondary peak in the annual tornado frequency for the Southwest and Northeast climate regions from 1950 to 2004. The frequency for the four corners region of the United States is included for comparison with the Arizona Northeast climate region.

Arizona Hail

1955-2004

Percentage of Hail Events By County



Arizona Averages 12 Hail Events per Year

Figure 13. The percentage of hail events by County from 1955 to 2004




Figure 14. Hail frequency by month and size from 1955 to 2004





Figure 15. Hail frequency of hour from 1955 to 2004

Arizona Thunderstorm Wind 1955-2004

Total Number of Reported Events by Region



Arizona Averages 29 Thunderstorm Wind Events per Year

Figure 16. Thunderstorm wind events by climate region from 1955 to 2004




Figure 17. Thunderstorm wind frequency by month from 1955 to 2004




Figure 18. Thunderstorm wind frequency by hour from 1955 to 2004

Arizona Thunderstorm Wind

Frequency by Hour and Region

1955-2004




Figure 19. Thunderstorm wind frequency by hour and climate region from 1955 to 2004




Figure 20. Flash flood and flood frequency by month from 1955 to 2004




Figure 21. Flash flood and flood frequency by hour from 1955 to 2004




Figure 22. Flash flood and flood frequency by month and climate region from 1955 to 2004

Areal Extent of Floods




Figure 23. Areal extent of some historical floods in Arizona (U.S Geological Survey)




Figure 24. Tropical system tracks over the Pacific and southwest U.S. 1958-2003. Red denotes tracks over the southwest states, black are tracks over the Pacific and dissipation over Mexico. (Corbosiero)




Figure 25. Frequency of dust storm fatalities by month from 1955 to 2004




Figure 26. Average number of thunderstorm days (Oklahoma Climatological Survey)




Figure 27. The ten year flash density for the U.S from 1996-2005 (Vaisala Inc.)




Figure 28. Lightning Fatalities and Injures by month from 1955 to 2004




Figure 29. Rate of lightning casualties (deaths and injuries combined) per population ordered by state from 1959 to 1994 (Curran, et. al, 2000)

State	Fatality Rate Per Million People from 1997 to 2006	Rank of Fatality Rate Per Million	State	Fatality Rate Per Million People from 1997 to 2006	Rank of Fatality Rate Per Million
Wyoming	1.21	1	Virginia	0.13	26
Colorado	0.70	2	Missouri	0.13	27
Montana	0.55	3	Texas	0.12	28
Utah	0.49	4	Ohio	0.11	29
Florida	0.44	5	Maryland	0.11	30
Arkansas	0.41	6	West Virginia	0.11	31
South Dakota	0.40	7	New Jersey	0.11	32
Mississippi	0.39	8	Iowa	0.10	33
Alabama	0.36	9	Michigan	0.10	34
Louisiana	0.36	10	Indiana	0.10	35
Vermont	0.33	11	Pennsylvania	0.10	36
South Carolina	0.32	12	Illinois	0.09	37
Rhode Island	0.29	13	Puerto Rico	0.08	38
Georgia	0.26	14	Kansas	0.07	39
North Carolina	0.24	15	Connecticut	0.06	40
Nebraska	0.23	16	Nevada	0.05	41
Oklahoma	0.23	17	Minnesota	0.04	42
Tennessee	0.21	18	New York	0.04	43
Arizona	0.18	19	Massachusetts	0.03	44
Kentucky	0.17	20	Oregon	0.03	45
New Mexico	0.16	21	California	0.02	46
Maine	0.16	22	Alaska	0	47
North Dakota	0.16	23	Delaware	0	48
Idaho	0.15	24	D.C.	0	49
Wisconsin	0.15	25	Hawaii	0	50
			New Hampshire	0	51
			Washington	0	52

Figure 30. Rate of lightning fatalities per population ordered by state from 1997 to 2006. Data and image provided courtesy of Vaisala NLDN (Vaisala Inc.)




Figure 31. A close up of the ten year flash density for Arizona from 1996-2005 (Vaisala Inc.)




Figure 32. Flash density key (Vaisala Inc.)




Figure 33. Flash densities (flashes per 100 km² per monsoon season) for 20 km x 20 km boxes during all hours for 1985-1990 (Watson, et. al 1993).




Figure 34. National Palmer Drought Severity Index (1895-1995)
(National Drought Mitigation Center)

DROUGHT EXTENT and LENGTH





Figure 35. Drought Extent, Length and Composite Palmer Drought Severity Indexes for Arizona (1950-1995) (Climate Prediction Center)

Areal Extent of Droughts




Figure 36. Areal extent of some historical droughts in Arizona (Paulson, et al.)





Figure 37. Arizona statewide water year precipitation, 1896-2005 (CLIMAS). Red highlights time periods of low precipitation.



Wildfire Hazard Severity

- Extreme
- High
- Medium
- Water (not included)
- Agriculture (not included)
- Barren (not included)

Note: Based on Urban-Wildland Fire Interface Code, IFCI 2000.

Figure 38. Wildfire Hazard Areas Map (Arizona State Land Department: Forestry Division)




Figure 39. Fire Hazard is based upon factors such as fuel types and condition, natural fire regimes, and topography and the influence of structures.
 (Arizona State Land Department: Forestry Division)



Figure 40. Fire hazard key (Arizona State Land Department: Forestry Division)




Figure 41. Mean annual snowfall for Arizona for the period 1961-1990
(Colorado Climate Center)




Figure 42. Key for mean annual snowfall for Arizona for the period 1961-1990
(Colorado Climate Center)




Figure 43. Mean annual snowfall days for Arizona for the period 1961-1990 (Colorado Climate Center)




Figure 44. Key for mean annual snowfall days for Arizona for the period 1961-1990 (Colorado Climate Center)




Figure 45. Snow and Ice by month from 1955 to 2004

**NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region
Subseries**

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, 125 South State Street - Rm 1311, Salt Lake City, Utah 84138-1102. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidl, April 1968 (Revised March 1986). (PB86 177672/AS). (Revised October 1991 - PB92-115062/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB 179084)
- 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. PB 184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 1969. (PB 185068)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)

- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 65 Climate of Sacramento, California. Laura Masters-Bevan. NWSO Sacramento, November 1998 (6th Revision). (PB99-118424)
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)
- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251)
- 83 A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS) (Revised August 2002 3rd Revision)
- 96 Map type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)

- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 126 Climate of San Francisco. E. Jan Null, February 1978. (Revised by George T. Pericht, April 1988 and January 1995). (PB88 208624/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033)
- 152 Climate of Salt Lake City, Utah. William J. Alder, Sean T. Buchanan, William Cope (Retired), James A. Cisco, Craig C. Schmidt, Alexander R. Smith (Retired), Wilbur E. Figgins (Retired), February 1998 - Seventh Revision (PB98-130727)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 103779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82-158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82-177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidt and Austin Jamison, April 1969 (Revised July 1996). (PB96-191614)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteris, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85-110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85-132694)

- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 1878887AS)
- 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lussky, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lussky, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Rapid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl R. Anderson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
- 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
- 199 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Lussky, April 1987. (PB87 185229/AS)
- 200 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke, September 1987. (PB88 110895/AS)
- 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson and George Clark, September 1987. (PB88 114038/AS)
- 202 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88-101935/AS)
- 203 An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern California. John P. Monteverdi and Scott A. Braun, October 1988. (PB89 121297/AS)
- 204 Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Carven Scott, November 1988. (PB89 128649/AS)
- 205 Forecast Guidelines For Fire Weather and Forecasters -- How Nighttime Humidity Affects Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS)
- 206 A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarters, Scientific Services Division, August 1989. (PB89 230833/AS)
- 207 The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, June 1990. (PB90-240268)
- 208 Meteorological Factors Contributing to the Canyon Creek Fire Blowup, September 6 and 7, 1988. David W. Goens, June 1990. (PB90-245085)
- 209 Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow Whitlatch, December 1990. (PB91-129239)
- 210 Hydrotools. Tom Egger, January 1991. (PB91-151787/AS)
- 211 A Northern Utah Soaker. Mark E. Struthwolf, February 1991. (PB91-168716)
- 212 Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 1991. (PB91-208439)
- 213 Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Mollner, July 1991. (PB91-227405/AS)
- 214 Emergency Operational Meteorological Considerations During an Accidental Release of Hazardous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424)
- 215 WeatherTools. Tom Egger, October 1991. (PB93-184950)
- 216 Creating MOS Equations for RAWS Stations Using Digital Model Data. Dennis D. Gettman, December 1991. (PB92-131473/AS)
- 217 Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 1992. (PB92-196104)
- 218 NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 1992. (PB93-146785)
- 219 A Case Study of the Operational Usefulness of the Sharp Workstation in Forecasting a Mesocyclone-Induced Cold Sector Tornado Event in California. John P. Monteverdi, March 1993. (PB93-178697)
- 220 Climate of Pendleton, Oregon. Claudia Bell, August 1993. (PB93-227536)
- 221 Utilization of the Bulk Richardson Number, Helicity and Sounding Modification in the Assessment of the Severe Convective Storms of 3 August 1992. Eric C. Evenson, September 1993. (PB94-131943)
- 222 Convective and Rotational Parameters Associated with Three Tornado Episodes in Northern and Central California. John P. Monteverdi and John Quadros, September 1993. (PB94-131943)
- 223 Climate of San Luis Obispo, California. Gary Ryan, February 1994. (PB94-162062)
- 224 Climate of Wenatchee, Washington. Michael W. McFarland, Roger G. Buckman, and Gregory E. Matzen, March 1994. (PB94-164308)
- 225 Climate of Santa Barbara, California. Gary Ryan, December 1994. (PB95-173720)
- 226 Climate of Yakima, Washington. Greg DeVoir, David Hogan, and Jay Neher, December 1994. (PB95-173688)
- 227 Climate of Kalispell, Montana. Chris Maier, December 1994. (PB95-169488)
- 228 Forecasting Minimum Temperatures in the Santa Maria Agricultural District. Wilfred Pi and Peter Felsch, December 1994. (PB95-171088)
- 229 The 10 February 1994 Oroville Tornado--A Case Study. Mike Staudenmaier, Jr., April 1995. (PB95-241873)
- 230 Santa Ana Winds and the Fire Outbreak of Fall 1993. Ivory Small, June 1995. (PB95-241865)
- 231 Washington State Tornadoes. Tresté Huse, July 1995. (PB96-107024)
- 232 Fog Climatology at Spokane, Washington. Paul Frisbie, July 1995. (PB96-106604)
- 233 Storm Relative Isentropic Motion Associated with Cold Fronts in Northern Utah. Kevin B. Baker, Kathleen A. Hadley, and Lawrence B. Dunn, July 1995. (PB96-106596)
- 234 Some Climatological and Synoptic Aspects of Severe Weather Development in the Northwestern United States. Eric C. Evenson and Robert H. Johns, October 1995. (PB96-112958)
- 235 Climate of Las Vegas, Nevada. Paul H. Skrbac and Scott Cordero, December 1995. (PB96-135553)
- 236 Climate of Astoria, Oregon. Mark A. McInerney, January 1996.
- 237 The 6 July 1995 Severe Weather Events in the Northwestern United States: Recent Examples of SSWEs. Eric C. Evenson, April 1996.
- 238 Significant Weather Patterns Affecting West Central Montana. Joe Lester, May 1996. (PB96-178751)
- 239 Climate of Portland, Oregon. Clinton C. D. Rockey, May 1996. (PB96-17603) - First Revision, October 1999
- 240 Downslop Winds of Santa Barbara, CA. Gary Ryan, July 1996. (PB96-191697)
- 241 Operational Applications of the Real-time National Lightning Detection Network Data at the NWSO Tucson, AZ. Darren McCollum, David Bright, Jim Meyer, and John Glueck, September 1996. (PB97-108450)
- 242 Climate of Pocatello, Idaho. Joe Heim, October 1996. (PB97-114540)
- 243 Climate of Great Falls, Montana. Matt Jackson and D. C. Williamson, December 1996. (PB97-126684)
- 244 WSR-88D VAD Wind Profile Data Influenced by Bird Migration over the Southwestern United States. Jesus A. Haro, January 1997. (PB97-135263)
- 245 Climatology of Cape for Eastern Montana and Northern Wyoming. Heath Hockenberry and Keith Meier, January 1997. (PB97-133425)
- 246 A Western Region Guide to the Eta-29 Model. Mike Staudenmaier, Jr., March 1997. (PB97-144075)
- 247 The Northeast Nevada Climate Book. Edwin C. Clark, March 1997. (First Revision - January 1998 - Andrew S. Gorelow and Edwin C. Clark - PB98-123250)
- 248 Climate of Eugene, Oregon. Clinton C. D. Rockey, April 1997. (PB97-155303)
- 249 Climate of Tucson, Arizona. John R. Glueck, October 1997
- 250 Northwest Oregon Daily Extremes and Normans. Clinton C. D. Rockey, October 1997
- 251 A Composite Study Examining Five Heavy Snowfall Patterns for South-Central Montana. Jonathan D. Van Ausdall and Thomas W. Humphrey. February 1998. (PB98-125255)
- 252 Climate of Eureka, California. Alan H. Puffer. February 1998. (PB98-130735)
- 253 Inferred Oceanic Kelvin/Rossby Wave Influence on North American West Coast Precipitation. Martin E. Lee and Dudley Chelton. April 1998. (PB98-139744)
- 254 Conditional Symmetric Instability--Methods of Operational Diagnosis and Case Study of 23-24 February 1994 Eastern Washington/Oregon Snowstorm. Gregory A. DeVoir. May 1998. (PB98-144660)
- 255 Creation and Maintenance of a Comprehensive Climate Data Base. Eugene Petrescu. August 1998. (PB98-173529)
- 256 Climate of San Diego, California. Thomas E. Evans, III and Donald A. Halvorson. October 1998. (PB99-109381)
- 257 Climate of Seattle, Washington. Dana Felton. November 1998. (PB99-113482)
- 258 1985-1998 North Pacific Tropical Cyclones Impacting the Southwestern United States and Northern Mexico: An Updated Climatology. Armando L. Garza. January 1999. (PB99-130502)
- 259 Climate of San Jose, California. Miguel Miller. April 1999. (PB99-145633)
- 260 Climate of Las Vegas, Nevada. Paul H. Skrbac. December 1999
- 261 Climate of Los Angeles, California. David Bruno, Gary Ryan, with assistance from Curt Kaplan and Jonathan Slemmer. January 2000
- 262 Climate of Miles City, Montana. David A. Spector and Mark H. Strobin. April 2000
- 263 Analysis of Radiosonde Data for Spokane, Washington. Rocco D. Pelatti. November 2000
- 264 Climate of Billings, Montana. Jeffrey J. Zeltwanger and Mark H. Strobin. November 2000
- 265 Climate of Sheridan, Wyoming. Jeffrey J. Zeltwanger, Sally Springer, Mark H. Strobin. March 2001
- 266 Climate of Sacramento, California. Laura Masters-Bevan. December 2000 (7th Revision)
- 267 Sulphur Mountain Doppler Radar: A Performance Study. Los Angeles/Oxnard WFO. August 2001
- 268 Prediction of Heavy Snow Events in the Snake River Plain Using Pattern Recognition and Regression Techniques. Thomas Andretta and William Wojcik. October 2003
- 269 The Lewis and Clark Expedition 18-03-1806, Weather, Water and Climate, Vernon Preston, Pocatello Idaho, December 2004.

- 270 Climate of San Diego, California, Emmanuel M. Isla, September 2004 (2nd Edition)
- 271 Climate of Las Vegas, Nevada, Andrew S. Gorelow, January 2005, (2nd Edition)
- 272 Climate of Sacramento, California, Revised by: Laura A. Bevan and George Cline, June 2005
- 273 Climate of Flagstaff, AZ 4th Revision. Mike Staudenmaier, Jr, Reginald Preston(R) Paul Sorenson (R) , August 2005
- 274 Climate of Prescott, AZ, Bob Fogarty, Mike Staudenmaier Jr., Flagstaff WFO, AZ, August 2005.
- 275 Climate of San Diego, CA, 3rd Revision. Noel M. Isla, Jennifer Lee, March 2006
- 276 Climate of Reno, NV, Brian Ohara, Reno, NV October 2006
- 277 Forecaster's Handbook for Extreme Southwestern California Based On Short Term Climatological Approximations: Part I - The Marine Layer and Its Effect On Precipitation and Heating Ivory J. Small, October 2006
- 278 Forecaster's Handbook for Extreme Southwestern California Based On Short Term Climatological Approximations: Part II – Wind Effects on Terrestrial and Marine Environments Ivory J. Small, December 2006
- 279 Effects of Wildfire in the Mountainous Terrain of Southeast Arizona: An Empirical Formula to Estimate 5-Year Peak Discharge from Small Post-Burn Watersheds, William B. Reed and Mike Schaffner , June 2007
- 280 Climate of Fresno, California, Chris Stachelski, Gary Sanger, February 2008
- 281 Climate of Bakersfield, California, Chris Stachelski, Gary Sanger February 2008
- 282 Hazardous Weather Climatology for Arizona - Craig Shoemaker, Jeffrey T. Davis. February, 2008

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS--Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

TECHNICAL SERVICE PUBLICATIONS -- Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.

Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161