

A UNITED STATES  
DEPARTMENT OF  
**COMMERCE**  
PUBLICATION



# NOAA Technical Memorandum NWS WR87

U.S. DEPARTMENT OF COMMERCE  
National Oceanic and Atmospheric Administration  
National Weather Service

## A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon.

ROBERT F. Y. LEE

Western Region

SALT LAKE CITY,  
UTAH

June 1973

NOAA TECHNICAL MEMORANDA  
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 23 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS.

Papers 1 to 23, except for 5 (revised edition) and 10, are available from the National Weather Service Western Region, Scientific Services Division, P. O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84111. Papers 5 (revised edition), 10, and all others beginning with 24 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Bldg., 5285 Port Royal Road, Springfield, Va. 22151. Price: \$3.00 paper copy; \$0.95 microfiche. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda

- WRTM 1 Some Notes on Probability Forecasting. Edward D. Diemer, September 1965. (Out of print.)  
WRTM 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.  
WRTM 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965 to February 20, 1966. Edward D. Diemer, March 1966.  
WRTM 4 Use of Meteorological Satellite Data. March 1966.  
WRTM 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (revised November 1967, October 1969). (PB-178000)  
WRTM 6 Improvement of Forecast Wording and Format. C. L. Glenn, May 1966.  
WRTM 7 Final Report on Precipitation Probability Test Programs. Edward D. Diemer, May 1966.  
WRTM 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (revised January 1967). (Out of print.)  
WRTM 9 A Collection of Papers Related to the 1966 NMC Primitive-Equation Model. June 1966.  
WRTM 10 Sonic Boom. Loren Crow (6th Weather Wing, USAF, Pamphlet), June 1966. (Out of print.) (AD-479366)  
WRTM 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.  
WRTM 12 A Comparison of Fog Incidence at Missoula, Montana, with Surrounding Locations. Richard A. Dightman, August 1966. (Out of print.)  
WRTM 13 A Collection of Technical Attachments on the 1966 NMC Primitive-Equation Model. Leonard W. Snellman, August 1966. (Out of print.)  
WRTM 14 Application of Net Radiometer Measurements to Short-Range Fog and Stratus Forecasting at Los Angeles. Frederick Thomas, September 1966.  
WRTM 15 The Use of the Mean as an Estimate of "Normal" Precipitation in an Arid Region. Paul C. Kangieser, November 1966.  
WRTM 16 Some Notes on Acclimatization in Man. Edited by Leonard W. Snellman, November 1966.  
WRTM 17 A Digitalized Summary of Radar Echoes Within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.  
WRTM 18 Limitations of Selected Meteorological Data. December 1966.  
WRTM 19 A Grid Method for Estimating Precipitation Amounts by Using the WSR-57 Radar. R. Granger, December 1966. (Out of print.)  
WRTM 20 Transmitting Radar Echo Locations to Local Fire Control Agencies for Lightning Fire Detection. Robert R. Peterson, March 1967. (Out of print.)  
WRTM 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.  
WRTM 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.  
WRTM 23 "K" Chart Applications to Thunderstorm Forecasts Over the Western United States. Richard E. Hambidge, May 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- WBTM 24 Historical and Climatological Study of Grinnell Glacier, Montana. Richard A. Dightman, July 1967. (PB-178071)  
WBTM 25 Verification of Operational Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)  
WBTM 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)  
WBTM 27 Objective Minimum Temperature Forecasting for Helena, Montana. D. E. Olsen, February 1968. (PB-177827)  
WBTM 28 Weather Extremes. R. J. Schmidl, April 1968 (revised July 1968). (PB-178928)  
WBTM 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB-178425)  
WBTM 30 Numerical Weather Prediction and Synoptic Meteorology. Capt. Thomas D. Murphy, U.S.A.F., May 1968. (AD-673365)  
WBTM 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB-179084)  
WBTM 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB-179289)  
WBTM 33 Objective Forecasting. Philip Williams, Jr., August 1968. (AD-680425)  
WBTM 34 The WSR-57 Radar Program at Missoula, Montana. R. Granger, October 1968. (PB-180292)  
WBTM 35 Joint ESSA/FAA ARTC Radar Weather Surveillance Program. Herbert P. Benner and DeVon B. Smith, December 1968 (revised June 1970). (AD-681857)  
WBTM 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (Out of print.) (PB-183055)  
WBTM 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB-183057)  
WBTM 38 Climate of Phoenix, Arizona. R. J. Schmidl, P. C. Kangieser, and R. S. Ingram. April 1969. (Out of print.) (PB-184295)  
WBTM 39 Upper-Air Lows Over Northwestern United States. A. L. Jacobson, April 1969. (PB-184296)  
WBTM 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L. W. Snellman, August 1969. (PB-185068)  
WBTM 41 High Resolution Radiosonde Observations. W. S. Johnson, August 1969. (PB-185673)  
WBTM 42 Analysis of the Southern California Santa Ana of January 15-17, 1966. Barry B. Aronovitch, August 1969. (PB-185670)  
WBTM 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB-185762)  
WBTM 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB-187763)  
WBTM 45/1 Precipitation Probabilities in the Western Region Associated with Winter 500-mb Map Types. Richard A. Augulis, December 1969. (PB-188248)

U. S. DEPARTMENT OF COMMERCE  
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION  
NATIONAL WEATHER SERVICE

NOAA Technical Memorandum NWSTM WR-87

A REFINEMENT OF THE USE OF K-VALUES IN FORECASTING  
THUNDERSTORMS IN WASHINGTON AND OREGON

Robert F. Y. Lee  
Weather Service Forecast Office  
Portland, Oregon

WESTERN REGION  
TECHNICAL MEMORANDUM NO. 87

SALT LAKE CITY, UTAH  
JUNE 1973



TABLE OF CONTENTS

	<u>Page</u>
List of Figures and Tables	iii-iv
Abstract	i
I. Introduction	1-2
II. Thunderstorm-Day Frequency	2-4
III. Results from Dependent Data	4
IV. Modification of the Objective Probabilities	5-6
V. Conclusion	6
VI. Acknowledgments	6
VII. References	6-7

## LIST OF FIGURES AND TABLES

	<u>Page</u>
Figure 1. Location of Areas A and B and Radiosonde Stations Bounding Each Area	8
Figure 2. Scatter Diagram Relating Frequency of Thunderstorm Days in Area A to Average 850-mb Temperatures and Average K-Values Computed from Radiosonde Observations at Boise, Salem, Medford, and Winnemucca	9
Figure 3. Relationship Between Computed Values of $\hat{Y}$ and Relative Frequency of Thunderstorm Days for Area A	10
Figure 4. Same as Figure 2, Except Lines Have Been Adjusted to Reflect Parabolic Relationship Shown in Figure 3	11
Figure 5. Scatter Diagram Relating Frequency of Thunderstorm Days in Area B to Average 850-mb Temperature and Average K-Value Computed from Radiosonde Observations at Boise, Salem, Quillayute, and Spokane	12
Figure 6. Relationship Between Computed Values of $\hat{Y}$ and Relative Frequency of Thunderstrom Days for Area B	13
Figure 7. Same as Figure 5, Except Lines Have Been Adjusted to Reflect Parabolic Relationship Shown in Figure 6	14
Figure 8. Mean Number of Thunderstorm Days in U. S. National Forests in Washington and Oregon from 1 July to 15 September 1950-1966 and 1968	15
Figure 9. Reliability Graph Relating "Forecast" Probabilities for Area A to Observed Relative Frequency of Occurrence of Thunderstorm Days	16
Figure 10. Reliability Graph Relating "Forecast" Probabilities for Area B to Observed Relative Frequency of Occurrence of Thunderstorm Days	17
Table 1. Percent Frequency of Occurrence of Thunderstorm Days in Area A in Categories of $\hat{Y}$	18
Table 2. Percent Frequency of Occurrence of Thunderstorm Days in Area B in Categories of $\hat{Y}$	18

List of Figures and Tables (Continued)

	<u>Page</u>
Table 3. Percent Frequency of Occurrence of Thunderstorm Days in Area A by Categories of P	19
Table 4. Percent Frequency of Occurrence of Thunderstorm Days in Area B by Categories of P	19
Table 5. List of Equations	20
Table 6. Scores for Probability "Forecasts" from Dependent Data	21
Table 7. Scores for Categorical "Forecasts" from Dependent Data $P \geq 40\%$ Used as Forecast of Thunderstorm Event	21

# A REFINEMENT OF THE USE OF K-VALUES IN FORECASTING THUNDERSTORMS IN WASHINGTON AND OREGON

## ABSTRACT

Average 1200 GMT K-values and 850-mb temperatures were used to develop an objective aid for making probability forecasts for afternoon and evening lightning occurrences in two target areas over Oregon and Washington. Thunderstorm and lightning reports from airway and Forest Service stations were used in this study.

The aid was developed on three summer seasons (1970-72 inclusive). Results on dependent data indicate that the objective forecasts can be expected to be better than climatological probabilities for the two target areas, and should be useful in preparing the short-range forecast of probability of thunderstorm occurrence issued during the early morning. Some qualitative synoptic features are discussed for modifying the objective forecast and to localize the concentration of thunderstorm occurrence in the target areas.

## 1. INTRODUCTION

Hambidge [1] has indicated that George's [2] technique of forecasting thunderstorm probability occurrence is applicable to western United States. George's results were applicable to southeastern United States. George's technique requires a scalar analysis of three parameters: K-value, 850-mb and 700-mb height summation, and winds below 700 mb. For air-mass thunderstorms, George used the K-value distribution to classify the number of thunderstorms for an area, modified by confluence or diffluence, and wind speeds below 700 mb. Hambidge used K-values to forecast probability of thunderstorm occurrence over western United States, modified also by confluence or diffluence.

In Washington and Oregon, fire-weather forecasters are required to forecast lightning probability for a number of federal and state protected areas. The forecaster is required to forecast a numerical probability of lightning occurrence. It is felt that a distribution of K-values combined with other numerical parameters will produce a useful objective method for forecasting lightning probabilities.

Wadsworth [3] and George both stress that parameters used in an objective forecast method should incorporate both static and dynamic atmospheric processes. The K-value is indicative of static conditions relative to thunderstorm occurrence. The 850-mb temperature can be a good dynamic predictor for thunderstorm occurrence. Generally, summer thunderstorms over Washington and Oregon occur west of the axis of the 500-mb ridge and ahead of a short-wave trough or upper low with colder air aloft moving over warmer air at lower levels. Thus, after a hot spell in the Pacific Northwest and before the on-rush of low-level marine air, summer thunderstorms generally tend to occur in this transitional period.

Through a limited investigation with thunderstorm occurrences, it was discovered that the 850-mb temperature was a good parameter to screen the K-value array in deriving probabilities of thunderstorm occurrences. Using the relationship of the 850-mb temperature with the K-value produced a better probability distribution of lightning occurrence than the relationship of the K-value with the 850-mb dew point, the 700-mb dew-point depression, the 500-mb temperature or the Showalter stability index. Although the 850-mb temperature is used twice in this objective method, the K-value masks the absolute value of 850-mb temperature. Since the 850-mb temperature is considered as a good dynamic predictor and is related to the synoptic patterns, it was felt that the array of the 850-mb temperature had to be used as a predictor.

## II. THUNDERSTORM-DAY FREQUENCY

Two forecast areas were selected and are shown in Figure 1. Area A includes about 30,000 square miles in Oregon and is bounded approximately by four upper-air sounding stations: Salem, Medford, Boise, and Winnemucca. Area B includes about 54,000 square miles in Oregon and Washington and is bounded approximately by Salem, Spokane, Boise, and Quillayute. Predictors from 1200 GMT soundings from these six upper-air stations and observed lightning or thunderstorm occurrences at airway and Forest Service stations in Oregon and Washington from June 1 to September 30, 1970, 1971 and 1972 were used to develop frequency distributions.

The following conditions were specified in deriving the objective forecast: 1) Thunderstorm and lightning occurrences are considered to be identical; 2) A thunderstorm day was defined as a day when one or more thunderstorms occurred in the given area between 1000 and 2200 Pacific Daylight Time; 3) Average K-values and average 850-mb temperatures were computed from four soundings surrounding the given area. The K-value equals the algebraic summation of the 850-mb temperature and dew point, minus the 700-mb dew-point depression, and minus the 500-mb temperature (degrees Celsius):

$$K = T_{850} - T_{500} - (DP.D.)_{700} + DP_{850} .$$

Figure 2 shows the joint relationship between average K-value, average 850-mb temperature and the occurrence of lightning in Area A. A linear least squares regression equation of the following form was computed for these data.

$$Y = A X_1 + b X_2 + C$$



where  $Y = 100$  if one or more thunderstorms occurred in the area;  $0$  if no thunderstorms occurred

$X_1$  = Average 850-mb temperature

$X_2$  = Average K-value

The resulting equation was:

$$\hat{Y} = 2.59 X_1 + 1.99 X_2 - 30.90. \quad (1)$$

The lines shown on Figure 2 are the graphical representation of equation 1. A value of  $\hat{Y}$  was computed for each case and the relative frequency of thunderstorm days determined for categories of  $\hat{Y}$ , as shown in Table 1, and plotted as crosses connected by dashed lines in Figure 3. The relationship of  $\hat{Y}$  to the frequency of occurrences of thunderstorm days obviously is not linear, and strongly suggest a parabolic relationship. Therefore, a parabola of the form:

$$P = A + B Y + C Y^2$$

was fitted to the points, resulting in:

$$P = 3.204 + .2647Y + .0124Y^2 \quad (2)$$

$$P = 0 \text{ for all values of } Y \leq -15$$

$$P = 100 \text{ for all values of } Y \geq 79.$$

The plot of the parabola is shown as the dotted curve on Figure 3. Making use of this relationship the lines on Figure 2 were adjusted to reflect the parabolic relationship between the 850-mb temperature, K-value and percent frequency (probability) of occurrences of a thunderstorm day. These adjusted lines are shown in Figure 4.

Thus, by merely entering Figure 4 with values of average 850-mb temperature and K-value, the probability (to the nearest percent) of a thunderstorm day in Area A is obtained by interpolation between the lines on Figure 4. Alternatively, of course, it is a simple matter to program the solution of equations 1 and 2 for a desk programmable calculator or for a minicomputer. An unbiased categorical forecast of thunderstorms is obtained by forecasting thunderstorms if  $P \geq 40$ , no thunderstorm if  $P < 40\%$ .

A similar analysis was performed for Area B as shown in Figure 5, 6 and 7 and Table 2. The linear regression equation relating average 850-mb temperatures and average K-values to occurrence of thunderstorm is:

$$Y = 2.36 X_1 + 2.04 X_2 - 19.91 \quad (3)$$

and the parabolic equation relating Y to the percent frequency of occurrence of thunderstorm days is:

$$P = 5.2123 + .4181Y + .0084Y^2$$

(P = 0 for all values of Y  $\leq$  -10)

(P = 100 for all values of Y  $\geq$  84).

### III. RESULTS FROM DEPENDENT DATA

Listed in Table 5 are a number of formulas for scores used to assess the "goodness" of probability and categorical forecasts. Applying these formulas to "forecasts" made from the dependent data resulted in the scores listed in Tables 6 and 7. Climatological probabilities for the given area were computed from the highest average thunderstorm days based on 18 years of record from the Forest Service shown in Figure 8.

Results are somewhat better for Area A than for Area B. This could possibly be due to the much larger size of Area B, and perhaps due to the inclusion of the Quillayute (UIL) sounding which is rather far removed from the eastern portion of Area B which has by far the greater frequency of thunderstorm days. Based on these scores, independent operational forecasts using this objective aid alone could be expected to show an improvement over a climatological probability forecast of about 30% in Area A and near 25% in Area B.

An indication of the resolution of the probability forecasts can be obtained by noting the average forecast probability of near 50% on thunderstorm days and an average forecast probability of 20 - 25% on no-thunderstorm days. Certainly, greater resolution than this is desirable, and perhaps the forecaster can improve upon this resolution subjectively following the ideas discussed in the following section.

The reliability of probability forecasts based on this aid can be expected to be good (on a reasonably large sample of independent data). Note the close agreement of the "forecast" probabilities and the observed percent frequency of occurrence of thunderstorms in Figures 9 and 10, with an average deviation of 4% for both areas.

A summary of categorical "forecasts" is presented in Table 7. The least biased categorical forecast is obtained for both areas from the probability forecasts by using P = 40% as the discriminant value. The prefigurance, postagreement and threat scores are very similar for both areas, and are quite respectable.

#### IV. MODIFICATION OF THE OBJECTIVE PROBABILITIES

By means of Figures 4 and 7 and daily 1200Z soundings which surround the given area, a forecaster has a means of deriving objectively a first approximation of lightning probabilities for afternoon and evening thunderstorms. The average lead time for the forecast of lightning occurrence is about seven hours. This is based on the idea that from 0600 to 0700 PDT the forecaster has ample time and information to use the frequency distribution and that thunderstorms tend to begin on the average about 1400 PDT. The maximum lead time is about 11 hours, assuming that the majority of thunderstorms begin after 1800 PDT.

Since this method does not have a true dynamic parameter, the forecaster can subjectively modify the first approximation of thunderstorm probability by using 500-mb types and sea-level pressure patterns. From examining the 1200 GMT 500-mb patterns for ten thunderstorm days in the summer of 1971, two types of 500-mb patterns were noted. It is realized that ten days are a small sample but the patterns correlate well with the general features of thunderstorm occurrences in the Pacific Northwest, based on long experience of forecasters in this area.

The first pattern involves a slow eastward progression of both a deep trough and a large ridge, oriented north to south. And the second involves a steady, extensive, southerly current. In the first situation, the trough-axis is located near 130 to 140 degrees west longitude, while the ridge-axis is located near 110 to 120 degrees west longitude. Under this first situation, one can subjectively increase the lightning probability. In the second situation, areas of confluence and speed convergence are embedded in the southerly current, and are associated with a minor perturbation or a weak cut-off low. Under the second situation, one can increase the objectively obtained first estimation of lightning probability under and ahead of areas of confluence and speed convergence.

In reference to the sea-level pressure pattern, position and eastward movement of the thermal trough are significant in the climatological probability for the occurrence of thunderstorms. Surface winds on either side of the surface thermal trough indicate a convergence region and a line of discontinuity. It is suspected that often the thermal trough is a low-level trigger mechanism for thunderstorms. Ward [4] has indicated that 67% of 625 days of thunderstorm occurrence over Oregon and Washington National Forests occurred when the thermal trough was along or east of the Cascade Range at 0100 GMT. Ward used a 10-year period from May to September 1925-1934. With an approaching upper trough or weakening cold front, and the eastward movement of the surface thermal trough, one should increase the objectively determined probability along and ahead of the expected thermal trough position during the afternoon and evening.

A forecaster can also subjectively modify the objective lightning probability by noticing the thunderstorms occurring upstream under steady flow that is expected to persist over the given area, and by anticipating kinematic convergence or divergence on the 700-mb map.

In localizing thunderstorm occurrences and probabilities to areas smaller than Areas A and B, one can use the surface thermal trough position at 1200 GMT. Sullivan's [5] localization of thunderstorm occurrence in Alaska with respect to the thermal trough applies in this area. Instead of an east-to-west thermal trough as in Alaska, there is a tendency for thermal troughs to be aligned north to south in Oregon and Washington, probably due to the Cascade Range's orientation. Sullivan indicated that nearly all thunderstorm occurrences were within 150 miles of the thermal trough-axis. In this analysis of the ten thunderstorm days in the summer of 1971, the elongated areas of thunderstorms tended to be parallel to the 1200 GMT thermal trough-axis and concentrated in a band from 30 miles west of the axis to 130 miles east, under a slow eastward progression of a deep trough and large ridge. When there is a perturbation embedded in a southerly flow, the concentration of thunderstorms is within 50 to 70 miles east and west of the 1200 GMT surface thermal trough-axis, with a secondary concentration at 100 to 120 miles east of the thermal trough-axis.

## V. CONCLUSION

The use of the mean K-value and the mean 850-mb temperature can produce a useful objective thunderstorm probability of occurrence as a first approximation in making a short-range forecast. However, other considerations such as present and future dynamic conditions of the atmosphere for the given area should be used to subjectively modify the first approximation to arrive at the final thunderstorm probability for the short-range forecast.

## VI. ACKNOWLEDGMENTS

Appreciation is expressed to Mr. Harold S. Ayer, Fire Weather Supervising Forecaster, Portland, Oregon, and to Scientific Services Division, Western Region Headquarters, for their reviews and suggestions. Special thanks are due Mr. Woodrow W. Dickey for his suggestions and aid in analysis of the data.

## VII. REFERENCES

- [1] HAMBIDGE, RICHARD E. *"K" Chart Application to Thunderstorm Forecasts Over the Western United States*. ESSA TM, WRTM 23, 1967.
- [2] GEORGE, JOSEPH J. *Weather Forecasting for Aeronautics*, pp. 407-415. Academic Press, New York and London, 1960.
- [3] WADSWORTH, GEORGE P. *Compendium of Meteorology*, pp. 849-855. American Meteorological Society, Boston, Mass., 1951.

- [4] WARD, ROBERT ALLEN. *Pressure Distribution in Relation to Thunderstorm Occurrence on Oregon and Washington National Forests*. Monthly Weather Review, Vol. 64, pp. 37-43, 1936.
- [5] SULLIVAN, WILLIAM G. *Low-Level Convergence and Thunderstorms in Alaska*. Monthly Weather Review, Vol. 91, pp. 89-92, 1963.
- [6] PANOFSKY, HANS A. and GLENN W. BRIER. *Some Applications of Statistics to Meteorology*, pp. 198-200. Pennsylvania State University, University Park, Penn., 1958.
- [7] ROBERTS, C. F., J. M. PORTER and G. F. COBB. *Report on the Forecast Performance of Selected Weather Bureau Office for 1966-1967*. ESSA TM, WBTM FCST-9, p. 8, December 1967.




Figure 1. Location of Areas A and B and Radiosonde Stations Bounding Each Area.




FIGURE 2. SCATTER DIAGRAM RELATING FREQUENCY OF THUNDERSTORM DAYS IN AREA A TO AVERAGE 850-MB TEMPERATURES AND AVERAGE K-VALUES COMPUTED FROM RADIOSONDE OBSERVATIONS AT BOISE, SALEM, MEDFORD, AND WINNEMUCCA. LINES REPRESENT SOLUTION OF LINEAR REGRESSION EQUATION:  $\hat{Y} = 2.59 X_1 + 1.99 X_2 - 30.90$ . CROSS REPRESENTS THUNDERSTORM DAY; DOT NONTHUNDERSTORM DAY.




FIGURE 3. RELATIONSHIP BETWEEN COMPUTED VALUES OF  $\hat{Y}$  AND RELATIVE FREQUENCY OF THUNDERSTORM DAYS (CROSSES AND DASHED LINE) FOR AREA A. DOTTED CURVE REPRESENTS PARABOLA FITTED TO CROSSES.






FIGURE 4. SAME AS FIGURE 2, EXCEPT LINES HAVE BEEN ADJUSTED TO REFLECT PARABOLIC RELATIONSHIP SHOWN IN FIGURE 3.




FIGURE 5. SCATTER DIAGRAM RELATING FREQUENCY OF THUNDERSTORM DAYS IN AREA B TO AVERAGE 850-MB TEMPERATURE AND AVERAGE K-VALUE COMPUTED FROM RADIOSONDE OBSERVATIONS AT BOISE, SALEM, QUILLAYUTE, AND SPOKANE. LINES REPRESENT SOLUTION OF LINEAR REGRESSION EQUATION:  $\hat{Y} = 2.36 X_1 + 2.04 X_2 - 19.91$ . CROSS REPRESENTS THUNDERSTORM DAY; DOT NONTHTUNDERSTORM DAY.




FIGURE 6. RELATIONSHIP BETWEEN COMPUTED VALUES OF  $\hat{Y}$  AND RELATIVE FREQUENCY OF THUNDERSTORM DAYS (CROSSES AND DASHED LINE) FOR AREA B. DOTTED CURVE REPRESENTS PARABOLA FITTED TO CROSSES.




FIGURE 7. SAME AS FIGURE 5, EXCEPT LINES HAVE BEEN ADJUSTED TO REFLECT PARABOLIC RELATIONSHIP SHOWN IN FIGURE 6.




FIGURE 8. MEAN NUMBER OF THUNDERSTORM DAYS IN U. S. NATIONAL FORESTS IN WASHINGTON AND OREGON FROM 1 JULY TO 15 SEPTEMBER 1950-1966 AND 1968.




FIGURE 9. RELIABILITY GRAPH RELATING "FORECAST" PROBABILITIES FOR AREA A TO OBSERVED RELATIVE FREQUENCY OF OCCURRENCE OF THUNDERSTORM DAYS. NUMBER BESIDE EACH POINT IS NUMBER OF CASES IN FORECAST PROBABILITY CATEGORY (SEE TABLE 3).




FIGURE 10. RELIABILITY GRAPH RELATING "FORECAST" PROBABILITIES FOR AREA B TO OBSERVED RELATIVE FREQUENCY OF OCCURRENCE OF THUNDERSTORM DAYS. NUMBER BESIDE EACH POINT IS NUMBER OF CASES IN FORECAST PROBABILITY CATEGORY (SEE TABLE 4).

TABLE 1

PERCENT FREQUENCY OF OCCURRENCE OF THUNDERSTORM DAYS  
IN AREA A IN CATEGORIES OF  $\hat{Y}$

<u><math>\hat{Y}</math></u>	<u>Total</u>	<u>No. of TSTM Days</u>	<u>%</u>
$\leq -15$	13	0	0
-14 to +14	67	4	5
15 to 24	55	6	11
25 to 34	58	11	18
35 to 44	51	15	29
45 to 54	49	23	47
55 to 64	22	17	77
65 to 74	23	19	83
$\geq 75$	6	6	100

TABLE 2

PERCENT FREQUENCY OF OCCURRENCE OF THUNDERSTORM DAYS IN AREA  
B IN CATEGORIES OF  $\hat{Y}$

<u><math>\hat{Y}</math></u>	<u>Total Cases</u>	<u>No. of TSTM Days</u>	<u>%</u>
$\leq -10$	10	0	0
-9 to +9	31	3	10
10 to 19	33	5	15
20 to 29	56	8	14
30 to 39	77	24	31
40 to 49	46	20	43
50 to 59	52	28	54
60 to 69	30	23	77
70 to 79	15	12	80
$\geq 80$	3	3	100



TABLE 3

PERCENT FREQUENCY OF OCCURRENCE OF THUNDERSTORM DAYS  
IN AREA A BY CATEGORIES OF P

<u>P(%)</u>	<u>Total</u>	<u>No. of TSTM Days</u>	<u>% Freq.</u>
0	11	0	0
1-6	45	5	11
7-14	63	4	6
15-24	61	10	16
25-34	45	12	27
35-44	35	12	34
45-54	33	16	48
55-64	15	13	74
65-74	8	4	
75-84	15	13	86
85-94	7	6	
≥95	6	6	100

TABLE 4

PERCENT FREQUENCY OF OCCURRENCE OF THUNDERSTORM DAYS  
IN AREA B BY CATEGORIES OF P

<u>P</u>	<u>Total Cases</u>	<u>No. of TSTM Days</u>	<u>% Freq.</u>
0	10	0	0
1-6	16	1	6
7-14	39	5	13
15-24	62	10	16
25-34	80	24	30
35-44	33	11	33
45-54	48	26	54
55-64	28	20	71
65-74	19	14	74
75-84	9	8	80
85-94	6	4	
≥95	3	3	100

TABLE 5

LIST OF EQUATIONS\*

- N = Total cases  
 TO = Number of occurrences of thunderstorm days  
 TF = Number of forecasts of thunderstorm days  
 TC = Number of correct thunderstorm forecasts  
 H = Total correct forecasts (thunderstorm and non-thunderstorm days)  
 P = Probability  
 E = 0 for non-thunderstorm event  
 E = 1 for thunderstorm event  
 C = Long-term climatological probability of TSTM days  
 CS = Sample climatology of thunderstorm day (observed frequency of occurrence of thunderstorm days in data sample)  
 BF = Brier Score for forecasts  
 BC = Climat Brier Score  
 BCS = Sample Climat Brier Score  
 S = Improvement over climatology  
 PF = Prefigurance  
 PA = Post Agreement  
 TS = Threat Score  
 % Cor. = Percent of categorical forecast correct

Equations:

$$1. \quad BF = \sum \frac{(P - E)^2}{N}$$

$$5. \quad \% \text{ Correct} = \frac{H}{N} (100)$$

$$2. \quad BC = (C - CS)^2 + CS(1 - CS)$$

$$6. \quad PF = \frac{TC}{TO}$$

$$3. \quad BCS = CS (1 - CS)$$

$$7. \quad PA = \frac{TC}{TF}$$

$$4. \quad S = \frac{BC - BF}{BC} (100) \text{ Improvement over climat}$$

$$8. \quad TS = \frac{TC}{TF + TO - TC}$$

$$S = \frac{BCS - BF}{BCS} (100) \text{ Improvement over sample climat}$$

\*See [6] and [7]

TABLE 6

SCORES FOR PROBABILITY "FORECASTS" FROM DEPENDENT DATA

<u>Probability Forecasts</u>	<u>Area A</u>	<u>Area B</u>
Total Cases	344	353
No. TSTM Days	101	126
Brier Score	.1450	.1765
Climat Brier Score	.2148	.2316
Sample Climat Brier Score	.2074	.2295
Improvement over Climat	32.50%	23.79%
Improvement over Sample Climat	30.74%	23.09%
Average Fcst Prob. for TSTM Days	51%	49%
Average Fcst Prob. for Non-TSTM Days	21%	27%

TABLE 7

SCORES FOR CATEGORICAL "FORECASTS" FROM DEPENDENT DATA  
 $P_{\geq 40\%}$  USED AS FORECAST OF TSTM EVENT

	<u>AREA A</u>			<u>AREA B</u>			
	<u>Forecast</u>			<u>Forecast</u>			
	TSTM	No-TSTM	TOTAL	TSTM	No-TSTM	TOTAL	
Obs   TSTM	65	36	101	Obs   TSTM	83	43	126
Obs   No-TSTM	35	208	243	Obs   No-TSTM	50	177	227
TOTAL	100	244	344	TOTAL	133	220	353

% Correct =  $273/344(100) = 79.3\%$

Prefiguration =  $65/101 = .643$

Post Agreement =  $65/100 = .650$

Threat Score =  $65/136 = .477$

Bias =  $100/101 = .990$

% Correct =  $260/353(100) = 73.6\%$

Prefiguration =  $83/126 = .658$

Post Agreement =  $83/133 = .624$

Threat Score =  $83/176 = .471$

Bias =  $133/126 = 1.055$

Western Region Technical Memoranda: (Continued)

- No. 45/2 Precipitation Probabilities in the Western Region Associated with Spring 500-mb Map Types. Richard P. Augulis. January 1970. (PB-189434)
- No. 45/3 Precipitation Probabilities in the Western Region Associated with Summer 500-mb Map Types. Richard P. Augulis. January 1970. (PB-189414)
- No. 45/4 Precipitation Probabilities in the Western Region Associated with Fall 500-mb Map Types. Richard P. Augulis. January 1970. (PB-189435)
- No. 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates. December 1969. (PB-190476)
- No. 47 Statistical Analysis as a Flood Routing Tool. Robert J. C. Burnash. December 1969. (PB-188744)
- No. 48 Tsunami. Richard A. Augulis. February 1970. (PB-190157)
- No. 49 Predicting Precipitation Type. Robert J. C. Burnash and Floyd E. Hug. March 1970. (PB-190962)
- No. 50 Statistical Report of Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona 1969. Wayne S. Johnson. April 1970. (PB-191743)
- No. 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell. July 1970. (PB-193102)
- No. 52 Sacramento Weather Radar Climatology. R. G. Pappas and C. M. Veliquette. July 1970. (PB-193347)
- No. 53 Experimental Air Quality Forecasts in the Sacramento Valley. Norman S. Benes. August 1970. (PB-194128)
- No. 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch. August 1970.
- No. 55 Application of the SSARR Model to a Basin Without Discharge Record. Vail Schermerhorn and Donald W. Kuehl. August 1970. (PB-194394).
- No. 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck. September 1970. (PB-194389)
- No. 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote. September 1970. (PB-194710)
- No. 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson. October 1970. (COM-71-00017)
- No. 59 Application of P.E. Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman. October 1970. (COM-71-00016)

NOAA Technical Memoranda NWS

- No. 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon. Arthur W. Fritz, October 1970. (COM-71-00120)
- No. 61 Relationship of Wind Velocity and Stability to SO<sub>2</sub> Concentrations at Salt Lake City, Utah. Werner J. Heck, January 1971. (COM-71-00232)
- No. 62 Forecasting the Catalina Eddy. Arthur L. Eichelberger, February 1971. (COM-71-00223)
- No. 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM-71-00349)
- No. 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- No. 65 Climate of Sacramento, California. Wilbur E. Figgins, June 1971. (COM-71-00764)
- No. 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM-71-00829)
- No. 67 Precipitation Detection Probabilities by Los Angeles ARTC Radars. Dennis E. Ronne, July 1971. (COM-71-00925)
- No. 68 A Survey of Marine Weather Requirements. Herbert P. Benner, July 1971. (COM-71-00889)
- No. 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM-71-00956)
- No. 70 Predicting Inversion Depths and Temperature Influences in the Helena Valley. David E. Olsen, October 1971. (COM-71-01037)
- No. 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM-72-10433)
- No. 72 A Paradox Principle in the Prediction of Precipitation Type. Thomas J. Weltz, February 1972. (COM-72-10432)
- No. 73 A Synoptic Climatology for Snowstorms in Northwestern Nevada. Bert L. Nelson, Paul M. Fransioli, and Clarence M. Sakamoto, February 1972. (COM-72-10338)
- No. 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM-72-10554)
- No. 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM-72-10707)
- No. 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM-72-11140)
- No. 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM-72-11136)
- No. 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM-72-11146)
- No. 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM-72-10920)
- No. 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM-72-10021)
- No. 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM-73-10150)
- No. 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972. (COM-73-10251)
- No. 83 A Comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM-73-10669)
- No. 84 Southwestern United States Summer Monsoon Source--Gulf of Mexico or Pacific Ocean? John E. Hales, Jr., March 1973.
- No. 85 Range of Radar Detection Associated with Precipitation Echoes of Given Heights by the WSR-57 at Missoula, Montana. Raymond Granger, April 1973.
- No. 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973.