

Inhomogeneous Responses of Ionospheric Electron Density to the May 2024 Superstorm

Dong L. Wu¹, Nimalan Swarnalingam^{1,2}, C. C. Jude H.
Salinas^{1,3}, Daniel J. Emmons⁴, Kyle E. Fitch⁴, and Jae N.
Lee^{1,3}

1. NASA – Goddard Space Flight Center, Greenbelt, Maryland
2. Dept. of Physics, The Catholic University of America, Washington, DC
3. The University of Maryland Baltimore County, Baltimore, Maryland
4. Dept. of Engineering Physics, Air Force Institute of Technology,
Wright-Patterson AFB, Ohio



Outline

Motivation

- Superstorm: A rare event in 20 years
- Structural evolution of ionospheric perturbations?

GNSS Limb (RO, POD) Data and Sampling

- 20k not enough for spatiotemporal sampling of fast processes (e.g., E/F-region dynamo)
- Strong demands on sampling from science apps:
 - Temporal: < hourly
 - Spatial: < 5 deg

Storms Observations

- Polar region: Spire + FY3
- 2-hourly zonal mean for E-region (GNSS-RO)
- 4-hourly zonal mean for F-region (GNSS-POD)




SuperStorms with $Dst \sim -400$ nT



17 Z May 10




May 2024



Lee et al. (2024, in preparation)

Sept 2003


CHAMP
SAC-C



Mannucci et al. (2005)



CSDA/CDAAC/ FengYun-3	# of Daily Profiles (May 2024)	
	E-region (V5)	F-region (V6p)
CSDA/Spire	7000	1800
COSMIC-2	6000	7200
FY3-D	450	350
FY3-E	900	950
FY3-F	1400	1200
FY3-G	1500	30
PlanetiQ	~1200	~1900
MetOp-B	~480	-
MetOp-C	~480	-
Kompsat-5	~230	-
TSX	~230	-
TDX	?	-
PAZ	?	-




CHAMP
vTEC ($z > 400\text{km}$)




Mannucci et al. (2005)

CSDA/Spire GNSS-RO Sampling



Wu (2024, GPSWorld)

gps240510g: Lat = [-90 89], Lon = [-180 179], UTC = 0:0:0




TIDs = Traveling ionospheric disturbances

Swarnalingam et al. (2024, in preparation)




Electron Density N_e from GNSS Limb Sounding

D/E-Region (GNSS-RO Link)




F-Region (GNSS-POD Link)



Optimal Estimation Inversion: $y = y_0 + K \cdot (x - x_0) + \epsilon_y$ $\hat{x} = [S_a^{-1} + K^T S_y^{-1} K]^{-1} [S_a^{-1} a + K^T S_y^{-1} y]$

Wu et al. (2022, v4 algorithm)
 (D/E-region N_e , vTEC, new v5)
 Shaver et al. (2023, N_e vs ionosonde)
 Salinas et al. (2024, empirical model)
 Salinas et al. (IROWG-10 poster)

Wu et al. (2023, v6p algorithm)
 Swarnalingam et al. (2023, NmF2, hmF2)
 Swarnalingam et al. (2024, in review)
 (Comparisons with IRI, NeQuick2)




Daytime 4-Hourly F-Region Ne





- Spire + FY3 sampling helps
- Large polar and tropical depletion in F-region
- Stronger and earlier in the NH than in the SH
- Reduced Ne in the NH subtropics during the recovery
- Enhanced Ne in the SH subtropics during the recovery

Spire only



Spire + FY3






Nighttime 4-Hourly F-Region Ne



- Spire + FY3 sampling helps
- Significant polar and tropical depletion
- Reduced Ne in the NH subtropics during the recovery

Spire only


Night, , Z=290km, Ne ($\times 10^{11} \text{ e}^-/\text{m}^3$)



17 Z May 10 (Day 131)

Spire + FY3

Night, , Z=290km, Ne ($\times 10^{11} \text{ e}^-/\text{m}^3$)



17 Z May 10 (Day 131)




Daytime 2-Hourly (Spire + FY3) E-Region Ne




- Consistent polar depletion between RO vTEC and F-region Ne
- Unlike F-region, stronger E-region depletion in the SH
- Small E-region changes during the recovery


Uncalibrated vTEC from RO (20-50km)



17 Z May 10 (Day 131)



17 Z May 10 (Day 131)




Nighttime 2-Hourly E-Region Ne




- Weak enhancement in RO vTEC
- Unlike F-region, significant nighttime E-region enhancements
 - Associated with auroral electron precipitation
- Small E-region changes during the recovery

Night, TEC from L1 Slope (TECU)



17 Z May 10 (Day 131)

Night, , Z=110km, Ne ($\times 10^{10} e^-/m^3$)




17 Z May 10 (Day 131)




9-Hourly UTC (00Z-09Z) Daily E-Region Ne Maps at 90 km (Spire + FY3 + COSMIC2)



Aurora Borealis
over the U.S.
from VIIRS DNB mosaic



Courtesy of:
CSU/CIRA & NOAA




$\Delta\text{lon}=20^\circ, \Delta\text{lat}=10^\circ$




Nighttime Polar E-Region Ne Maps at 90 km (Spire + FY3 + COSMIC2)



Aurora Borealis
over the U.S.
from VIIRS DNB mosaic



Courtesy of:
CSU/CIRA & NOAA




- Insufficient nighttime polar coverage from Spire + FY3




Daytime Polar E-Region Ne Maps at 90 km (Spire + FY3 + COSMIC2)


Aurora Borealis
over the U.S.
from VIIRS DNB mosaic




Courtesy of:
CSU/CIRA & NOAA



- Better daytime polar coverage from Spire + FY3



Daytime Polar F-Region Ne Maps at 350 km (Spire + FY3 + COSMIC2)




- Polar daytime Ne was depleted in the F-region shortly after the storm hits.
- Recovery took ~2 days.
- Russia-Ukraine war has a significant impact on sampling.



Summary

- GNSS-RO and GNSS-POD soundings provided unprecedented observations of the May 10 superstorm impacts on the global E- and F-region ionosphere.
- Hemispheric differences are evident in F-region responses:
 - Stronger and earlier NH Ne reduction
 - Enhanced Ne in the SH subtropics during the recovery
- E-region nighttime Ne enhancement extended to mid-latitudes, consistent with worldwide auroral reports.
- The 20k of daily GNSS-RO samples remain insufficient for observing rapid global ionospheric disturbances.




Acknowledgments


- UCAR COSMIC Data Analysis and Archive Center (CDAAC) for C2 L1B data
- NASA's Commercial Smallsat Data Acquisition (CSDA) and NOAA's Commercial Data Program (CDP) for Spire L1B data
- China Meteorological Administration (CMA) National Satellite Meteorological Center (NSMC) for FY3 L1B data
- Support from NASA's Living With a Star (LWS) program




Backup Slides



May 8



May 10






Sampling Density Maps

NH

2024d125–136 Average Sampling)

Missing
Data in
Ukraine-Ru
ssia



SH

2024d125–136 Average Sampling)

