
Wildfires

Identification

1. Indicator Description

This indicator tracks wildfire frequency, total burned acreage, burn severity, and the seasonality of wildfire activity in the United States from 1983 to 2021 or 2022 (depending on the metric). Evidence suggests that the incidence of large forest fires in the western United States and Alaska has increased since the 1980s, and it is projected to increase further in those regions as the climate changes (Ostoja et al., 2023). Although wildfires occur naturally and play a long-term role in the health of ecosystems, climate change threatens to increase the frequency, extent, and severity of fires through increased temperatures and drought. Earlier spring melting and reduced snowpack result in decreased water availability during hot summer conditions, which in turn contributes to an increased risk of wildfires, allowing fires to start more easily and burn hotter. The literature suggests that wildfires have undergone an increase in not just severity but also season length. Due to such factors as low fuel moisture, earlier warmer temperatures, and earlier snowmelt, the wildfire season has lengthened in many systems, and this trend is predicted to continue (Fill et al., 2019; Westerling, 2016).

Thus, while climate change is not the only factor that influences patterns in wildfire, the many connections between wildfire and climate make this indicator a useful tool for examining a possible impact of climate change on ecosystems and human well-being. Wildfires are also relevant to climate because they release carbon dioxide into the atmosphere, which in turn contributes to additional climate change.

Components of this indicator include:

- Wildfire frequency (Figure 1).
- Burned acreage from wildfires (Figure 2).
- Wildfire burn severity (Figure 3).
- Burned acreage from wildfires by state over time (Figures 4 and 5).
- Monthly burned acreage from wildfires over time nationwide (Figure 6).
- Monthly burned acreage from wildfires over time in the East and West (Figure 7).

2. Revision History

May 2014:	Indicator published.
June 2015:	Updated Figures 1 and 2 with data through 2014. Updated Figures 3 and 4 with data through 2013. Split Figure 4 into Figures 4 and 5.
April 2016:	Updated Figures 1 and 2 with data through 2015.
August 2016:	Updated Figures 3, 4, and 5 with data through 2014.
April 2021:	Updated Figures 1 and 2 with data through 2020 and Figures 3, 4, and 5 with data through 2018. Added Figures 6 and 7 (monthly patterns).
July 2022:	Updated Figures 1 and 2 with data through 2021 and Figures 3, 4, 5, 6, and 7 with data through 2020.

June 2024: Updated Figures 1 and 2 with data through 2022 and Figures 3, 4, 5, 6, and 7 with data through 2021.

Data Sources

3. Data Sources

Wildfire data come from three sources:

- Summary data for wildfire frequency and burned acreage from 1983 through 2022 (Figures 1 and 2) are provided by the National Interagency Coordination Center (NICC), housed within the National Interagency Fire Center (NIFC).
- For comparison in Figures 1 and 2, EPA obtained a data set called the United States Department of Agriculture (USDA) Forest Service Wildfire Statistics, which provides annual frequency and burned acreage totals through 1997 based on a different counting approach.
- Burn severity (Figure 3), state-by-state burn acreage (Figures 4 and 5), and monthly burned acreage (Figures 6 and 7) data were obtained from the Monitoring Trends in Burn Severity (MTBS) project, sponsored by the Wildland Fire Leadership Council (WFLC). The MTBS is a joint project of the USDA Forest Service Remote Sensing Applications Center (RSAC) and the United States Geological Survey (USGS) Earth Resources Observation and Science (EROS) Center. Other collaborators include the National Park Service, other USGS and USDA research facilities, and various academic institutions. The project provides data on individual wildfire incidents that meet certain size criteria ($\geq 1,000$ acres in the western United States or ≥ 500 acres in the eastern United States). These data were available from 1984 to 2021.

The analysis in Figures 4 and 5 normalizes wildfire extent by the land area of each state. Land areas come from the U.S. Census Bureau.

4. Data Availability

NIFC data for annual trends in wildfire frequency and acreage are available from the NIFC website at: www.nifc.gov/fireInfo/fireInfo_statistics.html. These NIFC data are also mirrored in the Wildland Fire Annual Reports from 2000 through 2023 at: www.nifc.gov/fire-information/statistics. NIFC totals are based on raw fire incidence data reported via the Incident Command System (ICS) Incident Status Summary Reports (ICS-209 forms). Some raw ICS-209 forms are available for individual viewing at: www.wildfire.gov/application/sit209.

The USDA Forest Service Wildfire Statistics represent a complementary approach to compiling fire occurrence and extent data. These statistics come from annual Forest Service reports officially known as annual “Wildland Fire Statistics,” but more commonly called “Smokey Bear Reports.” These compilation reports are based on reports submitted to the Forest Service by individual state and federal agencies, covering land within each agency’s jurisdiction. Smokey Bear Reports were provided to EPA by Forest Service researcher Karen Short. The Smokey Bear Report extent totals that appear in Figure 2 have also been published in Short (2015).

MTBS project analyses use raw ICS-209 form data from 1984 to 2021 as the basis for further processing. Summary data are publicly available at: <https://mtbs.gov/direct-download>. This online database search tool provides detailed and comprehensive records, including the location of initial fire occurrence as a GIS point file, burned area as a GIS polygon shape file, and burned area severity classification as a raster file for each individual fire incident. Detailed records for this indicator were downloaded from the MTBS website.

The U.S. Census Bureau has published official land areas for each state in the Statistical Abstract of the United States, available online at: www.census.gov/library/publications/time-series/statistical_abstracts.html.

Methodology

5. Data Collection

This indicator presents four measures of wildfires over time reported on an annual basis: (1) the total number of wildfires, (2) acreage burned by wildfires, (3) the burn severity of those fires, and (4) acreage burned by month. For the purposes of this indicator, a wildfire is a fire that fits the following nearly synonymous definitions:

“An unplanned, unwanted wildland fire including unauthorized human-caused fires, escaped wildland fire use events, escaped prescribed fire projects, and all other wildland fires where the objective is to put the fire out” (<https://gacc.nifc.gov/gbcc/dispatch/wy-tdc/home/sites/default/files/site-files/Master%20Glossary%20of%20Terms.pdf>)

“A wildland fire originating from an unplanned ignition, such as lightning, volcanos, unauthorized and accidental human caused fires, and prescribed fires that are declared wildfires” (NWCG, 2023)

NWCG (www.nwcg.gov/publications/pms205) defines a wildland as “an area in which development is essentially non-existent, except for roads, railroads, powerlines, and similar transportation facilities. Structures, if any, are widely scattered.” Fire severity is defined as the “degree to which a site has been altered or disrupted by fire; loosely a product of fire intensity and residence time.” The data for this indicator cover all 50 states.

Figures 1 and 2. Wildfire Frequency and Extent in the United States, 1983–2022

Wildfire frequency and burn acreage data are based upon local-, state-, and national-level reporting of wildland fire incidents submitted to NIFC via the ICS-209 form (Fire and Aviation Management and Predictive Services, 2009). The data captured in these forms can also be submitted to NIFC using the Incident Management Situation (SIT)-209 reporting application. The ICS-209 guidelines require that large fires (100+ acres in timber and 300+ acres in grasslands) must be reported, but they do not set a minimum fire size for reporting. Thus, the data set includes small fires, including some that may have burned just a few acres or less.


Supplementary data come from annual Smokey Bear Reports, which are based on annual reports submitted to the Forest Service by individual state and federal agencies. These original reports describe

fires taking place on land within each reporting agency’s fire protection jurisdiction. The USDA Forest Service stopped compiling Smokey Bear Reports after 1997.

Figure 3. Damage Caused by Wildfires in the United States, 1984–2021

MTBS uses satellite imagery to map burn severity and perimeters of large fires ($\geq 1,000$ acres in the western United States or ≥ 500 acres in the eastern United States). These thresholds are applied based on the “West” and “East” regions shown in Figure TD-1.

Figure TD-1. Region Boundaries for MTBS Size Threshold Application and Monthly Burned Acreage Analysis



MTBS starts primarily from ICS-209 reports and solicits additional data from the states if inclusion in ICS-209 is unclear. Other sources for fire occurrence data include federal data, the National Fire Plan Operations and Reporting System (NFPORS), and InciWeb. These records are compiled into a standardized project database. MTBS identifies corresponding imagery using the Global Visualization Image Selection (GLOVIS) browser developed by the USGS EROS Center. ArcGIS shapefiles and scene-specific Advanced Very High Resolution Radiometer (AVHRR) greenness plots are incorporated into the viewer to aid scene selection and determination of peak periods of photosynthetic activity. Pre-fire and post-fire images are selected for each incident. Wildfires are analyzed on the scale of individual incidents, but the data can also be aggregated at other spatial scales for analytical purposes.

Figures 4 and 5. Average Annual Burned Acreage and Change in Burned Acreage by State, 1984–2021

Figures 4 and 5 are based on acreage data for large fires as compiled by the MTBS program through the analytical steps described above for Figure 3. These numbers are based on ICS-209 reports and additional state data compiled by MTBS.

Figures 6 and 7. Comparison of Monthly Burned Area Due to Wildfires in the United States Between 1984–2002 and 2003–2021

Figures 6 and 7 are based on the same MTBS acreage data for large fires that were used for Figures 3, 4, and 5. The MTBS data set identifies each fire’s date of ignition, which this indicator uses to assign each fire to a month.

6. Indicator Derivation

Figures 1 and 2. Wildfire Frequency and Extent in the United States, 1983–2022

NIFC compiles local, state, and national reports to create the annual summary statistics published online. Data are aggregated to provide national, state and local statistics. EPA aggregated state-by-state totals in the annual Smokey Bear Reports to generate additional measures of annual wildfire frequency and extent.

Figure 3. Damage Caused by Wildfires in the United States, 1984–2021

Burn severity is a qualitative measure describing the degree to which a site has been altered by fire (NWCG, 2023). MTBS uses the Normalized Burn Ratio (NBR) to measure burn severity. NBR is a normalized index that uses satellite imagery from Landsat 5 and/or Landsat 7 TM/ETM bands 4 (near-infrared) and 7 (mid-infrared) to compare photosynthetically healthy and burned vegetation. Pre- and post-fire NBR are calculated to compare vegetation conditions before and after each wildfire.

The difference between pre- and post-fire NBRs is the Differenced Normalized Burn Ratio (dNBR). Calculated dNBR values are grouped into severity classes to give a qualitative assessment of the effects of fire damage. Exact classification thresholds may vary by site, based on site-specific ecological characteristics and expert judgment. Further discussion of the NBR and dNBR calculation methodology can be found at: www.mtbs.gov/mapping-methods and the glossary links therein.

Selected satellite images are also filtered through a complex sequence of data pre-processing, perimeter delineation, and other data quality assurance techniques. These procedures are documented in full on the MTBS website at: www.mtbs.gov/mapping-methods.

The timing of the satellite imagery selected for analysis depends on the type of assessment that is conducted for a given fire. The optimal assessment type is selected based on the biophysical setting in which each fire occurs. MTBS conducts two main types of assessments:

- Initial Assessments compare imagery from shortly before and shortly after the fire, typically relying on the first available satellite data after the fire—on the scale of a few days. These assessments focus on the maximum post-fire data signal and are used primarily in ecosystems that exhibit rapid post-fire vegetation response (i.e., herbaceous and particular shrubland systems).
- Extended Assessments compare “peak green” conditions in the subsequent growing season with “peak green” conditions in the previous growing season, prior to the fire. These assessments are designed to capture delayed first-order effects (e.g., latent tree mortality) and dominant

second-order effects that are ecologically significant (e.g., initial site response and early secondary effects).

MTBS occasionally conducts a Single Scene Assessment, which uses only a post-fire image (either “initial” or “extended”), when limited by factors such as data availability.

Figure 3 was created by filtering MTBS’s database output to remove any fires not meeting MTBS’s size criteria—although most such fires would not have been processed by MTBS anyway—and removing fires classified as “prescribed,” “wildland fire use,” or “unknown.” The resulting analysis is therefore limited to fires classified as true “wildfires.”

The total acreage shown in Figure 3 (the sum of the stacked burn severity sections) does not match the total acreage in Figure 2 because the burn severity analysis in Figure 3 is limited to fires above a specific size threshold ($\geq 1,000$ acres in the western United States and ≥ 500 acres in the eastern United States) and because the graph does not include acreage classified as “outside perimeter” or “non-processing area mask,” the latter of which denotes areas within the fire perimeter that could not be assessed for burn severity because the imagery was affected by clouds, cloud shadows, or data gaps. The “Key Points” text that describes the proportion of high severity acreage is based on high severity as a percentage of total assessed acreage (i.e., the total acreage after non-processing area has been excluded). The size threshold resulted in five states that did not have any reported fires for the time period: Connecticut, Illinois, New Hampshire, Rhode Island, and Vermont. Puerto Rico is included in Figure 3 totals.

Figure 4. Average Annual Burned Acreage by State, 1984–2021

To create this map, EPA divided the annual acreage burned in each state by the state’s total land area. After doing this for all years during the period of record (1984–2021), EPA calculated an average value and plotted it on the map. The same five states without fires meeting the size threshold for Figure 3 are colored gray to indicate insufficient data.

Figure 5. Change in Annual Burned Acreage by State Between 1984–2002 and 2003–2021

To create this map, EPA calculated each state’s average annual acreage burned per square mile for the first half of the record (1984–2002) and the average for the second half (2003–2021). EPA found the difference between these values and expressed it as a percentage difference (e.g., average annual acreage during the second half of the record was 10 percent higher than average annual acreage burned during the first half). The same five states without fires meeting the size threshold for Figure 3 are colored gray to indicate insufficient data. Changes have been characterized using this method rather than measuring a slope over time (e.g., a linear regression) because of the length and shape of the data set. Visual inspection of the NIFC line in Figure 2 (burned acreage across all states) suggests periods of relative stability punctuated by a noticeable jump in acreage that starts during the late 1990s. This jump coincides with a period of transition in certain natural climate oscillations that tend to shift every few decades—notably, a shift in the Pacific Decadal Oscillation (PDO) around 1998 (Peterson & Schwing, 2003; Rodionov & Overland, 2005). This shift—combined with other ongoing changes in temperature, drought, and snowmelt—may have contributed to warmer, drier conditions that have fueled wildfires in parts of the western United States (Kitzberger et al., 2007; Westerling, 2016). With nearly 40 years of data punctuated by a phase transition, and with research strongly suggesting that the PDO and other decadal-scale oscillations contribute to cyclical patterns in wildfires in the western United States, EPA

determined that linear regression is not an appropriate method of describing changes over time in this particular indicator. Instead, EPA chose to simply compare two sub-periods in a manner that considers all years of data and avoids inferring an annual rate of change. Without a nuanced statistical analysis to define a break point between two sub-periods, EPA chose to simply break the record into two periods of equal length: 1984–2002 and 2003–2021.

EPA plans to investigate opportunities for a more robust interpretation of state-level trends over time in future editions of this indicator.

Figures 6 and 7. Comparison of Monthly Burned Area Due to Wildfires in the United States Between 1984–2002 and 2003–2021


EPA created Figures 6 and 7 by starting from the filtered data set described above for Figure 3 (i.e., excluding any fires not meeting MTBS’s size criteria). EPA divided the period of record into two equal halves (1984–2002 and 2003–2021) to support analysis of change over time. Changes have been characterized using this comparative method rather than measuring a slope over time (e.g., a linear regression) because of the variability in the data set. The timing of the burn acreage was highly variable from year to year (even more so when viewed within specific regions of the country). Instead, EPA chose to simply compare two sub-periods in a manner that considers all years of data and avoids inferring an annual rate of change. Without a nuanced statistical analysis to define a break point between two sub-periods, EPA chose to simply break the record into two 19-year periods. As more data are added in future years, the “halfway” break point will move accordingly.

To create the graphs, EPA calculated the average annual acreage burned each month across the United States for each half of the period of record. The date of fire occurrence used in this analysis corresponds to the ignition date of the fire as defined in the MTBS data set. For this figure, fires are aggregated by month. For Figure 7, the calculation was conducted separately for fires in the East and West, as defined by the MTBS program and shown in Figure TD-1.

Comparison of Sources

Figure TD-2 compares total wildfire extent estimates from NIFC, Smokey Bear Reports, and MTBS. This graph shows that MTBS estimates follow the same pattern as the NIFC data set but are always somewhat lower than NIFC’s totals because MTBS excludes small fires. The graph also shows how the most recent MTBS estimates compare with the MTBS data release used in the previous update to this indicator. As expected, the data show evidence of revisions to historical data, but the changes are not extensive.

Figure TD-2. Comparison of Wildfire Extent from Three Data Sources, 1983–2021



Indicator Development

EPA added Figures 6 and 7 to this indicator in 2021 as a result of a new indicator development effort that focused on wildfire season characteristics. EPA used wildfire activity data because it was nationally available with a reasonable data record length (consistently collected since the early 1980s). However, there are many analytical approaches and definitions for examining the wildfire season that have been used in the peer-reviewed scientific literature, most of which have been specific to regional conditions including ecosystem type, vegetation, and fire regimes. EPA plans to continue to research methods and data for examining long-term changes in the seasonality of wildfires (onset, peak, end, and duration).

NIFC’s website provides data from 1960 through 2022, and Smokey Bear Reports are available from 1917 to 1997. The data available prior to the early 1980s, however, provide incomplete geographic coverage, as fire statistics at the time were not compiled from the full extent of “burnable” lands. Thus, Figures 1 and 2 of this indicator begin in 1983, which was the first year of nationwide reporting via ICS-209 reports. Figures 3 through 7 begin in 1984, which was the first year for which the MTBS project conducted its detailed analysis. MTBS depends on aerial imagery and the level of detail captured consistently in ICS reports. Thus, while a longer period of record would be desirable when analyzing long-term changes in a climatological context, EPA could not extend this indicator with pre-1983 data without introducing inconsistencies and gaps that would preclude meaningful comparisons over time and space.


For more discussion regarding the availability, coverage, and reliability of historical wildfire statistics, see the authoritative discussions in Short (2014) and Short (2015). Based on these sources, Table TD-1 summarizes the available data sets, their coverage, and their underlying sources. Pre-1983 estimates have been published by NIFC in the past; they derive from the Smokey Bear Reports (Short, 2015). These reports are the only underlying nationwide source of pre-1983 wildfire statistics.

Table TD-1. Comparison of Historical Wildfire Data Sources

Data set	Variables	Temporal range	Resolution	Geographic coverage	Underlying sources
NIFC <i>(Figures 1 and 2)</i>	Acreage and incidence (number of fires)	1983–2022	Annual	National	ICS incident reports
NIFC pre-1983	Acreage and incidence	1960–1982	Annual	National with gaps	Smokey Bear Reports, which are based on estimates submitted by various agencies
Smokey Bear Reports <i>(recent data in Figures 1 and 2)</i>	Acreage and incidence	1917–1997	Annual	National with gaps	Estimates submitted by various agencies
MTBS <i>(Figures 3 through 7)</i>	Burn severity; acreage by state	1984–2021	Annual	National	ICS incident reports

A fundamental shift in wildfire reporting took place in the early 1980s with the onset of the ICS reporting system. Prior to this time, reports were submitted to the USDA Forest Service by selected state and federal agencies, covering land within each agency’s jurisdiction. Many of these reports were limited to fires on land with “protected” status (i.e., land designated for cooperative fire control). Fires occurring on “unprotected” land would not necessarily be fought, and they would not be counted in the statistics either. Figure TD-3 below, based on data obtained from the USDA Forest Service (Short, 2015), demonstrates how the reporting area was well below the total nationwide “burnable” acreage until the 1980s. Increases in the reporting area occurred when additional agencies joined the reporting program. For example, the Bureau of Land Management began reporting around 1964, which accounts for the noticeable jump in the blue line in Figure TD-3.

Figure TD-3. “Smokey Bear Reports” Reporting Area Over Time



The Smokey Bear Reports achieved essentially complete participation and coverage by the early 1980s. They continued to be compiled even with the advent of the ICS system around 1983, until being phased out in the late 1990s. Thus, the Smokey Bear Reports and the ICS reports provide complementary coverage for much of the 1980s and 1990s. During the first few years of ICS data collection—particularly 1983 and 1984—the Smokey Bear Reports showed much higher fire occurrence than NIFC’s ICS-derived statistics. The USDA Forest Service attributes this difference to the ramp-up and gradual adoption of ICS nationwide. Other studies such as Dennison et al. (2014) also describe the advantages of using recent, robust data sets instead of longer, less complete fire databases previously used for trend analysis.

The MTBS program’s burned area and burn severity mapping methods have evolved over time (Picotte et al., 2020), though any significant methodological changes are applied to the whole historical dataset to ensure consistency. MTBS notes that remapped MTBS products can be significantly different from original mappings. Per MTBS’s recommendation, EPA always downloads the most recent data release to update this indicator. The latest major update to mapping methods and procedures occurred with MTBS’s November 24, 2020, data release, which included a comprehensive reprocessing of the entire MTBS archive to introduce consistent vector attribution and metadata updates, plus mapping of additional fires and revisions to previously mapped fires. This upgrade contributed to some visible changes in Figures 4 and 5 between EPA’s 2021 and 2022 web updates for this indicator, but they did not change the overall patterns, trends, or statistical significance of what this indicator shows.

7. Quality Assurance and Quality Control

The ICS-209 form provides clear guidelines for wildfire reporting for state, local, and federal agencies. These guidelines are accessible on the NIFC website at: www.nifc.gov/nicc/predictive-services/intelligence. The information in the ICS-209 forms is compiled by the SIT program to provide

daily situation reports that summarize wildfire conditions at state and national levels. This compiled information forms the basis for NIFC summary statistics. NIFC does not provide details on how it counts, measures acreage, or filters out the double reporting of fires, fires that split, fires that merge, incomplete forms, or other potential data irregularities. The frequency of these confounding factors, however, is likely limited and may not seriously compromise the quality of the data presented in this indicator.

MTBS standardizes and corrects raw fire incidence data. The project avoids editing source data with the exception of correcting a record's geospatial coordinates if (1) the coordinates are clearly incorrect, and (2) a correction can be made with confidence. All selected scenes from the database are ordered and processed following existing USGS-EROS protocols. Data obtained from MTBS were also cross-checked prior to conducting analyses for this indicator.

Analysis

8. Comparability Over Time and Space

NIFC methods and statistics have not changed since 1983, and they can be compared on an annual basis at national scales. The sole exception is the NIFC fire count and burned acreage data points for the year 2004, which are missing totals from state lands in North Carolina. Thus, these two points slightly compromise the comparability over both time and space. Smokey Bear Reports also used consistent methods from 1983 to 1997, and they covered the full extent of "burnable" U.S. lands throughout this period. MTBS has used consistent methods to classify burn severity from 1984 through 2021, allowing for annual comparisons through time and allowing for spatial comparisons among states. MTBS is based on a type of satellite imagery that has been collected consistently with sufficient resolution to support this analysis throughout the period of record. Although MTBS has updated its methods over time (Picotte et al., 2020), program staff go back and apply any new methods to the full historical record to ensure continued comparability. The most recent large reanalysis occurred in late 2020 and was conducted over the entire timeframe of available data.

Figures 3 through 7 were derived from an MTBS data set that uses different size thresholds for different states. This data set includes fires $\geq 1,000$ acres in the western United States and ≥ 500 acres in the eastern United States. Thus, the indicator might undercount small fires in the West, compared with the East. However, because total burned area is dominated by large fires, this is not expected to heavily affect the calculations underlying Figures 3 through 7. In addition, these thresholds have held consistent for each region over time, which lends validity to the analysis of regional trends over time.

9. Data Limitations

Factors that may impact the confidence, application, or conclusions drawn from this indicator are as follows:

1. Wildfire activity can be influenced by a variety of other factors besides climate. Examples include changes in human activities and land management strategies over time, particularly changes in fire suppression and fire management practices, which (among other things) can potentially contribute to more damaging fires in the future if they result in a buildup of fuel in the understory. Resources available to fight and manage wildfires can also influence the amount

of area burned over time. Grazing activities can also influence the amount and type of vegetation in the landscape, and changes in land cover and land use—for example, forest to non-forest conversion—can affect the extent and type of “burnable” land. Thus, further analysis is needed before an apparent change in wildfire activity can necessarily be attributed to climate change.

2. The dominant drivers of wildfire activity can vary by region. Contributing factors may include (but are not limited to) temperatures in specific seasons (particularly spring), drought, and precipitation that contributes to vegetation growth. As described in Section 6, wildfire trends in some regions have been linked with certain phases of multi-year and multi-decadal climate oscillations (Kitzberger et al., 2007; Westerling et al., 2006). Climate patterns that lead to more wildfire activity in some parts of the United States may lead to a simultaneous decrease in activity in other regions (e.g., the Northwest versus the Southwest). Reconstructions based on tree rings can provide hundreds of years of context for understanding such patterns and how they vary regionally (e.g., Swetnam & Betancourt, 1998).
3. While this indicator is officially limited to “wildland” fires, it includes fires that encroach on—or perhaps started in—developed areas at the wildland-urban interface (WUI). Encroachment of the WUI over time into previously wild lands could influence trends in wildfire frequency and extent (Radeloff et al., 2005).
4. NIFC data, which are derived from government entities of varying scope or jurisdiction, can be limited by inconsistencies across how data are reported through ICS-209 forms. With aggregation from so many potential sources, wildfire incidence data, particularly historical data, may be redundant or erroneous. Data aggregation among sources may result in variability in reporting accuracy and protocol.
5. The MTBS program depends on certain conditions to make accurate measurements of burn severity:
 - Accurate fire location coordinates that match burn scars visible via satellite.
 - Accurate fire size information that ensures that fires meeting the MTBS size criteria are properly included.
 - Accurate date of ignition and out date that guide the appropriate selection of imagery, particularly for baseline assessments.
 - Pre-fire and post-fire images that are cloud-free to avoid visual obscuration of the fire area.
6. Some fires of very low severity may not be visible in the satellite imagery and therefore impossible to delineate or characterize. Cloud cover, cloud shadow, or data gaps can also preclude damage assessment. To account for all of these limitations, the MTBS project includes a burn severity classification of “non-processing area mask.” This classification accounts for approximately 2.0 percent of the total wildfire acreage from 1984 through 2022.

10. Sources of Uncertainty

Uncertainties in these data sets have not been quantified. The most likely sources of uncertainty relate to initial data collection methods. Federal land management agencies have varying standards for content, geospatial accuracy, and nomenclature. Duplicate records occur due to reporting of a given

incident by multiple agencies, such as redundant reports from local, state, or federal entities. In any given year, as much as three-quarters (or more) of all fire incidents are reported by non-federal state and local agencies (for example, see NICC [2023]). Cases of gross geospatial inaccuracies may also occur. Similar inconsistencies occur within state databases; however, the MTBS project addresses issues such as duplicates and nomenclature during pre-processing. Given the comprehensive reporting infrastructure that has been in place over the past several decades, it is unlikely that any large fires would have been missed.

11. Sources of Variability

Forest conditions, and therefore wildfire incidents, are highly affected by climate conditions. In addition to year-to-year variations, evidence suggests that wildfire patterns in the western United States are influenced by multi-year and multi-decadal climate oscillations such as the PDO (www.ncei.noaa.gov/access/monitoring/pdo) and the Atlantic Multidecadal Oscillation (www.aoml.noaa.gov/phod/amo_faq.php). For example, see Kitzberger et al. (2007) and Westerling (2016) for discussion of warmer, drier conditions that have contributed to increases in wildfire activity in certain regions.

Changes in the frequency of wildfire triggers (e.g., lightning, negligent or deliberate human activity) could also affect wildfire frequency and the timing of wildfire activity. Burn severity is affected by local vegetation regimes and fuel loads. Finally, improvements or strategic changes in firefighting and fire management may affect wildfire prevalence, size, and resulting damages. Forest management practices have changed over time from complete fire suppression to controlled burns. These varied approaches and the extent to which they are applied on state or regional levels can influence the wildfire data presented in this indicator.

12. Statistical/Trend Analysis

As described in Section 6, the nature of this topic and the length and shape of the time series suggest that linear regression is not a suitable tool for characterizing long-term wildfire trends and their significance. Thus, the figures and key points do not report regression results. Ordinary least-squares linear regressions from the NIFC data set have been calculated here, just for reference. Regression slopes and p-values are indicated in Table TD-2 below.

Table TD-2. Wildfire Regression Statistics

Indicator component	Regression slope	P-value
NIFC fire frequency (Figure 1)	+10 fires/year	0.966
NIFC burn acreage (Figure 2)	+169,952 acres/year	< 0.001

EPA used a two-tailed paired t-test to compare the monthly burned acreage time series (the lines in the graphs in Figures 6 and 7) between the two time periods. This testing treats each period as a series of 12 monthly means (the 12 months—each representing the mean from a sample set of 19 years’ worth of data); the paired aspect means that the test compares the January mean from the first period to the January mean from the second period, then does the same for February and so on. The paired t-test offers a simple assessment of whether wildfire extent (burned acreage) over the course of the year has increased or decreased to a statistically significant degree. Differences were significant to a 95 percent

level ($p < 0.05$) in the United States as a whole (Figure 6; p -value of 0.018 for the comparison) and the West (Figure 7; $p = 0.017$). The East had an insignificant difference between the two periods ($p = 0.823$).

Use of a simple t-test inevitably requires some assumptions about the shape and underlying characteristics of the data. This test is used to provide initial approximations about the statistical attributes of this indicator. One strength of the testing described here is that it identifies regions where there has been an increase in fire activity through much of the year. One weakness is that it may fail to identify differences in regions where fire activity is concentrated in just a few months of the year, because most months are zero or near-zero and thus show very little difference between the two time periods.

References

- Dennison, P. E., Brewer, S. C., Arnold, J. D., & Moritz, M. A. (2014). Large wildfire trends in the western United States, 1984–2011. *Geophysical Research Letters*, 41(8), 2928–2933. <https://doi.org/10.1002/2014GL059576>
- Fill, J. M., Davis, C. N., & Crandall, R. M. (2019). Climate change lengthens southeastern USA lightning-ignited fire seasons. *Global Change Biology*, 25(10), 3562–3569. <https://doi.org/10.1111/gcb.14727>
- Fire and Aviation Management and Predictive Services. (2009). *Interagency wildland fire incident information reporting application (SIT-209)*. www.wildfire.gov/application/sit209
- Kitzberger, T., Brown, P. M., Heyerdahl, E. K., Swetnam, T. W., & Veblen, T. T. (2007). Contingent Pacific–Atlantic Ocean influence on multicentury wildfire synchrony over western North America. *Proceedings of the National Academy of Sciences*, 104(2), 543–548. <https://doi.org/10.1073/pnas.0606078104>
- NICC (National Interagency Coordination Center). (2023). *Wildland fire summary and statistics annual report 2022*. www.nifc.gov/sites/default/files/NICC/2-Predictive%20Services/Intelligence/Annual%20Reports/2022/annual_report.2.pdf
- NWCG (National Wildfire Coordinating Group). (2023). *NWCG glossary of wildland fire, PMS 205* (updated August 2023). www.nwcg.gov/publications/pms205
- Ostoja, S. M., Crimmins, A. R., Byron, R. G., East, A. E., Méndez, M., O’Neill, S. M., Peterson, D. L., Pierce, J. R., Raymond, C., Tripathi, A., & Vaidyanathan, A. (2023). Focus on western wildfires. In USGCRP (U.S. Global Change Research Program), *Fifth National Climate Assessment*. <https://doi.org/10.7930/NCA5.2023.F2>
- Peterson, W. T., & Schwing, F. B. (2003). A new climate regime in northeast Pacific ecosystems. *Geophysical Research Letters*, 30(17), 2003GL017528. <https://doi.org/10.1029/2003GL017528>
- Picotte, J. J., Bhattarai, K., Howard, D., Lecker, J., Epting, J., Quayle, B., Benson, N., & Nelson, K. (2020). Changes to the Monitoring Trends in Burn Severity program mapping production procedures and data products. *Fire Ecology*, 16(1), 16. <https://doi.org/10.1186/s42408-020-00076-y>

- Radeloff, V. C., Hammer, R. B., Stewart, S. I., Fried, J. S., Holcomb, S. S., & McKeefry, J. F. (2005). The wildland-urban interface in the United States. *Ecological Applications*, 15(3), 799–805. <https://doi.org/10.1890/04-1413>
- Rodionov, S., & Overland, J. E. (2005). Application of a sequential regime shift detection method to the Bering Sea ecosystem. *ICES Journal of Marine Science*, 62(3), 328–332. <https://doi.org/10.1016/j.icesjms.2005.01.013>
- Short, K. C. (2014). A spatial database of wildfires in the United States, 1992-2011. *Earth System Science Data*, 6(1), 1–27. <https://doi.org/10.5194/essd-6-1-2014>
- Short, K. C. (2015). Sources and implications of bias and uncertainty in a century of US wildfire activity data. *International Journal of Wildland Fire*, 24(7), 883–891. <https://doi.org/10.1071/WF14190>
- Swetnam, T. W., & Betancourt, J. L. (1998). Mesoscale disturbance and ecological response to decadal climatic variability in the American Southwest. *Journal of Climate*, 11(12), 3128–3147. [https://doi.org/10.1175/1520-0442\(1998\)011<3128:MDAERT>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<3128:MDAERT>2.0.CO;2)
- Westerling, A. L. (2016). Increasing western US forest wildfire activity: Sensitivity to changes in the timing of spring. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371(1696), 20150178. <https://doi.org/10.1098/rstb.2015.0178>
- Westerling, A. L., Hidalgo, H. G., Cayan, D. R., & Swetnam, T. W. (2006). Warming and earlier spring increase western U.S. forest wildfire activity. *Science*, 313(5789), 940–943. <https://doi.org/10.1126/science.1128834>