


Advances in the assessment and monitoring of salinization and status of biosaline agriculture

Report of an expert consultation held in Dubai,
United Arab Emirates, 26–29 November 2007



Cover photographs:

Salt-affected soils in Uzbekistan.

Courtesy of T. Friedrich (FAO).

Map:

Salt-affected soils around the world.

Courtesy of F. Nachtergaele (FAO).

Copies of FAO publications can be requested from:

SALES AND MARKETING GROUP

Communication Division

Food and Agriculture Organization of the United Nations

Viale delle Terme di Caracalla

00153 Rome, Italy

E-mail: publications-sales@fao.org

Fax: +39 06 57053360

Web site: <http://www.fao.org>

Advances in the assessment and monitoring of salinization and status of biosaline agriculture

WORLD
SOIL
RESOURCES
REPORTS

104

Report of an expert consultation held in Dubai,
United Arab Emirates, 26–29 November 2007

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

Advances in the assessment and monitoring of salinization and status of biosaline agriculture Reports of expert consultation held in Dubai, United Arab Emirates, 26–29 November 2007.
World Soil Resources Reports No. 104. FAO, Rome.

ISBN: 978-92-5-106439-9

All rights reserved. Reproduction and dissemination of material in this information product for educational or other non-commercial purposes are authorized without any prior written permission from the copyright holders provided the source is fully acknowledged. Reproduction of material in this information product for resale or other commercial purposes is prohibited without written permission of the copyright holders. Applications for such permission should be addressed to:

Chief

Publishing Management Service

Information Division

FAO

Viale delle Terme di Caracalla, 00153 Rome, Italy

or by e-mail to:

copyright@fao.org

Contents

Acknowledgements	vii
Part 1	
Executive summary	1
Rationale for an expert consultation	1
Objectives	1
Format of the SPUSH consultation and INBA meeting	2
Attendance	2
General recommendations	4
Recommendations for further applied research	5
Part 2	
Abstracts according to programme	7
Inaugural session	7
Welcome	7
Introduction to the SPUSH Network	8
A. M. MASHALI	
Session 1: Assessment and monitoring of salt-affected soils at field, landscape and irrigation district levels	9
Assessing and monitoring the risk of salinization in a Sicilian vineyard using the Geonics EM38	10
G. CRESCIMANNO, K. B. MARCUM, F. MORGA AND C. REINA	
Use of an aboveground electromagnetic induction meter for assessing salinity changes in natural landscapes and agricultural fields	11
J. ARRIOLA MORALES ¹ AND J. BATLLE SALES ²	
Primary soil salinity, sodicity and alkalinity status of different water management areas in South Africa: The quest for baseline data	12
J. P. NELL, M. C. DE VILLIERS AND M. VAN DER WALT	

Using satellite imagery and electromagnetic induction to assess soil salinity, drainage problems and crop yield in the Río Fuerte Irrigation District	14
L. PULIDO MADRIGAL AND J. GONZALEZ MERAZ	
Session 2: Assessment and monitoring of salt-affected soils at national and regional levels	15
Soil salinity monitoring and assessment in irrigated Arab agriculture	16
G. ABDELGAWAD	
Advances in the assessment of salt-affected soils for mapping, monitoring and management strategies in India	17
G. SINGH AND A. K. MANDAL	
Salt-affected soils in Thailand: Assessment and monitoring of salinization	19
R. IM-ERB AND S. SUKCHAN	
An overview of the salinity problem in Iran: Assessment and monitoring technology	21
S. A. M. CHERAGHI, Y. HASHEMINEJHAD AND M. H. RAHIMIAN	
Advances in the assessment and monitoring of soil salinization for managing salt-affected habitats in Egypt	22
M. H. GOMAA	
Recent evolution of soil salinization and its driving processes in China	23
J. YANG	
Assessment and management of salt-affected soils in Sudan	24
A. A. EL-MOBARAK	
Session 3: Modelling for salinity/sodicity development	26
Overview of salinity modelling approaches at different spatial-temporal scales	27
J. BATLLE SALES	
SMSS: a soil-plant model describing the impact of irrigation on soil and runoff water salinity	28
M. LAHLOU AND M. BADRAOUI	
Session 4: Mapping and interpretation of spatial data	29
Emerging challenges addressing the characterization and mapping of salt-induced land degradation	30
M. QADIR AND E. DE PAUW	

Monitoring, predicting and quantifying soil salinity, sodicity and alkalinity in Hungary at different scales: Past experiences, current achievements and outlook with special regard to European Union initiatives	32
T. TÓTH	
Salinization and sodification of irrigated soils in Eastern Kenya	34
S. N. WANJOGU AND P. T. GICHERU	
Nature and distribution of salts in the Upper Wami-Ruvu Plains, Morogoro, Tanzania: Implications for land management methods	35
M. KILASARA AND E. L. KABONI	
Salt-affected soils in Romania	37
M. DUMITRU, C. SIMOTA, D. RADUCU, E. DUMITRU, I. CALCIU AND N. FLOREA	
Session 7: sustainable biosaline agricultural systems	38
Biosaline agriculture: Prospects and potential within global and regional context	39
S. ISMAIL AND F. TAHA	
Greywater use for irrigation of home gardens in peri-urban areas of Jordan	40
M. BINO	
Saline agriculture: Pakistan scenario	41
R. H. QURESHI AND J. AKHTAR	
Extent of salt-affected land in Central Asia: Biosaline agriculture and the utilization of salt-affected resources	42
K. TODERICH, S. ISMAIL, I. MASSINO, M. WILHELM, S. YUSUPOV, T. KULIEV AND S. RUZIEV	
Extent and utilization of salt-affected lands: Biosaline agriculture and marginal resources in Tajikistan	45
S. SANGINBOY	
The salinity problem in the Sultanate of Oman: Past, present and future perspectives	46
S. B. A. R. AL-RASBI	
Water shortage in western US and saline recycled water use for urban irrigation	47
K. B. MARCUM	

Part 3**Findings and reports from working groups and field visit 49**

Findings 49

Reports from working groups 51

Field visit 55

Annex I**Information on the networks 57**

The global network on salinization prevention and productive use of salt-affected habitats (SPUSH) 57

The Inter-Islamic Network on Biosaline Agriculture (INBA) 57

Annex II**SPUSH expert consultation and INBA meeting programme 59****Annex III****List of participants 67**

Acknowledgements

The Food and Agriculture Organization of the United Nations and its Plant Production and Protection Division, express their gratitude to the International Center for Biosaline Agriculture (ICBA) for hosting the SPUSH Expert Consultation, as well as for contributing with their time and expertise to the organizing committee. They also express their gratitude to ICBA and the Inter-Islamic Network on Biosaline Agriculture (INBA) for providing generous funds for the events. In addition, the contribution from the following institutes to fund the participation of their experts are highly appreciated: Arizona State University, Department of Applied Biological Sciences, USA; International Center for Agricultural Research in the Dry Areas/International Water Management Institute (ICARDA/IWMI), Syrian Arab Republic; the OIC Standing Committee on Scientific and Technological Cooperation (COMSTECH); Università di Palermo, Dipartimento ITAF, Italy; United Nations Educational, Scientific and Cultural Organization (UNESCO); Universidad de Valencia, Departamento de Biología Vegetal, Spain.

The organizing committee also acknowledges the scientific and technical contribution of the following participating countries: Algeria, China, Egypt, Hungary, India, Iran (Islamic Republic of), Italy, Jordan, Kenya, Mexico, Morocco, Oman, Pakistan, Romania, Saudi Arabia, South Africa, Spain, the Sudan, the Syrian Arab Republic, Tajikistan, the United Republic of Tanzania, Thailand, United Arab Emirates, the United States of America and Uzbekistan.

The Organizing Committee thanks the following participants for contributing to the preparation of this report: Dr. Giuseppina Crescimano, Palermo University, Italy; Dr. Patrick Gicheru, National Agricultural Research Laboratories, Kenya; Dr. Shoaib Ismail, ICBA, United Arab Emirates; Dr. Method Kilasara, Department of Soil Science, Sokoine University of Agriculture, the United Republic of Tanzania; Dr. Amin Mashali, FAO; Dr. J. P. Nell, ARC-Institute for Soil, Climate and Water, South Africa; Dr. M. Qadir, ICARDA/IWMI, the Syrian Arab Republic; Riaz H. Qureshi, Higher Education Commission, Pakistan; Dr. Gurbachan Singh, Central Soil Salinity Research Institute, India; and Dr. Sanginov Sanginboy, Tajikistan Academy of Agricultural Sciences, Republic of Tajikistan.

The Organizing Committee thanks Dr. Amin Mashali for the work carried out for the preparation of the expert consultation, the selection of papers and the writing of these proceedings. It also acknowledges the substantial work of Dr. Rhodri P. Thomas for the technical review, edition and formatting of the proceedings.

The publication of these proceedings was coordinated by Dr. Clemencia Licona Manzur, Soil Reclamation and Development Officer, Plant Production and Protection Division.

The Secretariat of the Global Network on Salinization Prevention and Productive Use of Salt-affected Habitats (SPUSH) Network is hosted by the Plant Production and Protection Division of FAO.

Part 1

Executive summary

RATIONALE FOR AN EXPERT CONSULTATION

Soil salinization and sodication have been identified as major processes of land degradation and loss of agricultural production. The high costs of measuring salinization and sodication, as well as inconsistencies in data collection and reporting methods, have resulted in incomplete – and often contradictory – information on the extent and distribution of salt-affected soils at country, regional and global levels. Nevertheless, scientists and government authorities consider that the problem is extensive and growing.

The causes of salinity/sodicity, which vary between countries and regions, need to be identified, assessed and monitored carefully so that they can be managed and controlled. The few maps that show the extent of salt-affected soils are often based on old data and do not give a picture of the current extent of salinity, as data to determine the rate of change of salinization need to be updated regularly. Therefore, there is a need for practical and cost-effective methodologies for assessing, monitoring and mapping the extent and distribution of salt-affected soils; for assessing the causes and sources of the problem; and for choosing management options and evaluating the effectiveness of those options.

In order to collect up-to-date information on the methods for assessment and monitoring of salinization and sodication, the Food and Agriculture Organization of the United Nations (FAO), through the Global Network on Salinization Prevention and Productive Use of Salt-affected Habitats (SPUSH), along with the Center for Biosaline Agriculture (ICBA) and the Inter-Islamic Network on Biosaline Agriculture (INBA), organized the SPUSH Expert Consultation on *Advances in Assessment and Monitoring of Salinization for Managing Salt-affected Habitats*. This was held in Dubai from 26 to 29 November 2007. (More information on SPUSH and INBA Networks is included in Annex I.)

Holding the Expert Consultation in ICBA constituted an opportunity to learn about its work, as well as that of the INBA Network. Therefore, the organizers decided to allocate an additional day for holding the Meeting on *Status and Progress of Biosaline Agriculture of the Inter-Islamic Network on Biosaline Agriculture* (INBA), in order to promote further exchange of information between the Networks and explore areas for collaboration.

OBJECTIVES

The objectives of the SPUSH Consultation and the INBA Meeting were to:

- exchange experiences with data collection and analysis for the assessment and monitoring of salinity and sodicity, with particular emphasis on practical applications at local, national, regional and global levels;
- collect up-to-date official country statistics on the extent of different types of salt-affected soils;
- reactivate the SPUSH Network, introduce new topics and identify priorities for action;

- present the work of the host institution and that of INBA members in order to explore points of convergence, strengthen information exchange and explore potential areas of collaboration.

The SPUSH Expert Consultation and the INBA Meeting were a step towards collaboration between two international institutions to bring experts together for the purpose of developing a consensus on various aspects of the problem of soil salinity and, in particular, its assessment and monitoring. Another objective of this meeting was to provide a common perspective to various players engaged in research and development efforts in the cross cutting areas, to tackle this important problem.

FORMAT OF THE SPUSH CONSULTATION AND INBA MEETING

The Expert Consultation consisted of presentations by participants, group discussions and a field visit to ICBA facilities. The topics covered were:

- assessment and monitoring of salt-affected soils (at field, landscape and irrigation district levels);
- assessment and monitoring of salt-affected soils at national and regional levels;
- modelling for salinity-sodic development;
- mapping and interpretation of spatial data;
- reactivation of the SPUSH Network and future work;
- status and progress of biosaline agriculture with examples of work carried out by INBA members.

The detailed programme is included in Annex II. Abstracts of papers presented are included in Part 2; the findings and recommendations of working groups are included in Part 3; and full papers are included in the CD-ROM accompanying this publication.

ATTENDANCE

The Expert Consultation and the INBA Meeting were attended by senior soil and water scientists and authorities representing national institutes, members of the SPUSH and INBA Networks from different regions:

- Africa: Kenya, South Africa and Tanzania;
- Asia: China, India, Pakistan, Thailand;
- Europe: Italy, Romania, Spain, Tajikistan and Uzbekistan;
- Near East: Egypt, Iran, Jordan, Morocco, Oman, Saudi Arabia, Sudan, United Arab Emirates and United States of America.

Officers from the following international institutions also participated in the SPUSH Consultation and INBA Meeting: the Food and Agriculture Organization of the United Nations (FAO); the International Center for Agricultural Research in the Dry Areas/ International Water Management Institute (ICARDA/IWMI), the International Center for Biosaline Agriculture (ICBA), the International Union of Soil Sciences (IUSS) Working Group of Salt-affected Soils, the Islamic Development Bank (IDB), the OIC Standing Committee on Scientific and Technological Cooperation (COMSTECH) and the United Nations Educational, Scientific and Cultural Organization (UNESCO).

A total of 40 participants attended the events (the list of participants is included in Annex III). The organizers also received abstracts from member countries and international institutions that could not participate, either due to unforeseen circumstances or lack of funds. As the organizers had a limited amount of funds available, they received more abstracts than could be funded.

CONCLUSIONS

The following conclusions include the ideas and concepts presented in papers as well as the results of group discussions:

- Soil salinity/sodicity, waterlogging and low soil fertility are serious problems at the global level and are present in the countries participating in the Consultation to varying extent and severity. Due to the high cost of measurements, the lack of data and the inconsistencies between data provided by various sources, there is incomplete information on the extent, distribution, rate and degree of salinity development for most of participating countries. In some countries, the existence of these soils was discovered only because of the pressing demand for agricultural utilization in specific areas. Few countries have up-to-date data on the extent of salt-affected soils.
- Researchers and field technicians often lack funds, other resources and government support to enable them to work on issues related to salt-affected soils including assessment, monitoring and mapping.
- Human resources required to assess, monitor and map salt-affected soils are limited in most participating countries.
- Many countries lack systematic national procedures to monitor secondary salinization and sodication. Studies are done on an *ad hoc* basis. Reliable data to establish baseline conditions are also lacking.
- Depending on the scale, the different methods required for assessing and monitoring salt-affected soils in participating countries varied from simple field surveys, sampling and laboratory measurements, including electrical conductivity (EC) and exchangeable sodium percentage (ESP), to the use of expert assessment, salinity sensors, electromagnetic sensors, remote sensing, Geographic Information Systems (GIS) and modelling.
- Models require a very intensive effort for data collection and are not commonly used in most participating countries. However, some countries do have standardized assessments, modelling and mapping methodologies and procedures that could be shared with other SPUSH members to improve classification and mapping of their salt-affected soils.
- Policies that relate to the management of salt-affected soils in the participating countries are often not effective. Most countries do not have a strategy or policy at national level for assessing and monitoring salt-affected soils.
- The available maps on the extent, severity and rate of salinization/sodication and waterlogging in participating countries need to be updated as they are based on old data. In this regard, SPUSH may develop guidelines and a database on resource surveys and land use planning. This database must be maintained and updated regularly.
- The Expert Consultation participants acknowledged the efforts made by the Organizing Committee to make possible the SPUSH Consultation and INBA meetings and to bring together different experts to discuss common problems. Participants also hoped that the efforts made will lead to further strengthening and development of cooperation between members of the two Networks.

GENERAL RECOMMENDATIONS

The following is a summary of the recommendations from the SPUSH Expert Consultation and the INBA Meeting; many of them relate to the future activities of the SPUSH Network. Detailed recommendations for each topic covered are included in Part 3:

- There is a need to strengthen research and implementation of assessment, monitoring, mapping and modelling of salt-affected soils in participating countries. The SPUSH Network could be fundamental for the promotion of collaboration between different interested national and international organizations.
- Efforts to explore possible financial support from other agencies or national programmes to strengthen the Network activities should be intensified. The SPUSH secretariat should explore funding possibilities with FAO, ICBA, UNESCO, EU, UNEP, ICARDA and Gulf Cooperation Countries (GCC) to keep the Network active.
- The SPUSH Network should be expanded to other countries having problems with salt-affected soils. In this context, further collaboration should continue between the SPUSH and INBA Networks and other relevant networks.
- An expert group needs to be constituted within the SPUSH Network to address technical issues such as improving remote sensing methods, standardizing definitions, considering the assessment of waterlogged saline soils; studying the cost-effectiveness of microwave and thermal lines for the assessment of salt-affected soils.
- SPUSH Network countries are encouraged to establish benchmark sites for periodic monitoring of salinity, sodicity, water quality; develop and use early warning mechanisms to undertake preventive measures; develop farmer-friendly salinity assessment and monitoring systems.
- Countries that have standardized assessment, modelling and mapping methodologies are encouraged to share expertise with other SPUSH members in order to assist them to update existing data on the extent and distribution of salt-affected soils. Improving the data of SPUSH members will facilitate not only management at national level but also the preparation of a new global salinity map, to increase awareness of the problem and potentially to obtain funding to combat problems.
- Several countries have identified promising germplasm of trees, shrubs, grasses and crops (as well as fish) that can contribute to the management of salt-affected soils. Network members are encouraged to share, as far as possible, their knowledge in this area with other members of the SPUSH and INBA Networks.
- Members should seek to formulate joint network project/programmes related to assessment, mapping, monitoring and modelling of salt-affected soils, to benefit from members' expertise and undertake more effective fund raising.
- The SPUSH Network should serve as a focal point to create national and international awareness, to exchange information on sustainable and environmentally-sound use of salt-affected soils and related issues such as assessment, mapping, monitoring, modelling, and impacts of climate change. It is recommended that the Network considers coordinated field experiments, education, advisory services, publications for different audiences and through different information media.
- The Network will promote the development of kits on assessment and monitoring for field technicians and farmers. This material should be available free of charge. China offered the technical facilities to develop such kits and ICARDA, Iran, Jordan and Tanzania showed interest in contributing.

- The SPUSH member countries should be encouraged to publish good quality papers. The Network should also distribute guidelines for different publics. Papers and presentations in the Expert Consultation should be published as proceedings after editing.
- The secretariat of SPUSH should continue to maintain the Network website which should be modified to allow direct contribution and regular updates from members. Quality publications, maps and available data should be included in the website.
- A task force to ensure the continuity of the Network was established. The initial appointed members were India, Iran, Italy, Pakistan, ICARDA and FAO.
- Participants recommended that the next Expert Consultation should be organized as soon as funds are available. Hungary and Spain offered to host the next meeting, while Iran offered to be the host after Hungary or Spain. It is suggested that the topic of the next Consultation could be “Impacts of climate change on salinity development”. However, many of the member countries may not have enough related information in their countries, as climate change is a complex issue. The topic suggested as an alternative is “Sustainable management of salt-affected soils under changing climatic conditions”, which could include biosaline agriculture, in which case INBA could participate. In all cases, the national institutes are the focal points to participate in the Expert Consultation and not individuals.

RECOMMENDATIONS FOR FURTHER APPLIED RESEARCH

- There is a need to design appropriate monitoring networks. Monitoring of irrigation schemes is critical.
- There is also a need for developing methodologies for assessing different kinds of salinity and crop damage based on integrated use of remote sensing and plant physiology.
- The development of salt-tolerant crop varieties will become fundamental (breeding with the application of gene-surgery, selection, production of salt-tolerant plants). The use of biosaline agriculture could also be further advocated; e.g. the use of *Salicornia*, *Atriplex*, *Salvadora* and *Prosopis* has been identified as a good alternative in salt-affected areas.
- The classification of salt-affected soils should be harmonized and it should consider their potential land use and include soils rich in magnesium.
- Land use classification and decision support systems need to be further developed.
- There is a need for research on the implementation of sustainable irrigation technologies using low quality water, including saline water and wastewater, considering different impacts to human and ecosystem health.
- Unfavourable changes in groundwater quality due to over-exploitation need to be further studied.
- More effort should be made on investigating the impact of climate change and salt dynamics in soils.
- There is a need to devise early warning systems that use simple indicators.
- Researchers should contribute to the development of simple kits for farmers to assess and monitor salinity in the field.
- Assessment and monitoring of salinity should include associated salts/metals, such as like boron, iron, aluminium, manganese, arsenic, selenium, nitrates, etc.
- More studies are needed to show if salt-affected soils and poor quality waters may also be an asset for seaweed cultivation for biomass/hormone production and growing halophytes for energy or salt production.

- There is a need for advice on assessment and management technologies that incorporate socio-economic and environmental factors.
- There is a need for models that better represent field conditions. For this reason, further parameters should be considered.

Part 2

Abstracts according to programme

INAUGURAL SESSION

Welcome

The Inaugural Session took place at the Headquarters of the International Center for Biosaline Agriculture (ICBA), Dubai, United Arab Emirates (UAE). Mr. Shawki Barghouti, Director General, Mr. Fawzi Al-Sultan, Chairman, and Dr. Faisal Taha, Director Technical Programme, ICBA, talked about the mandate and on-going activities of ICBA as a scientific research institution focusing on development and promoting systems that facilitate crop production in areas characterized by saline water or soils.

Mr. Mohammed Toure, Islamic Development Bank (IDB), and Dr. Anwar Nasim, OIC Standing Committee on Scientific and Technological Cooperation (COMSTECH), addressed the Consultation with a brief presentation of the role, funding programmes, activities and achievements of their institutions.

Mr. Kayan Jaff, FAO Representative in the UAE, described FAO's mandate and the work carried out in the region to improve agricultural production and presented FAO's work on issues related to soil degradation, including salt-affected soils and related subjects within the framework for combating global land degradation. He also highlighted FAO's collaboration with several institutions.

Dr. Amin Mashali, on behalf of Dr. Clemencia Licona Manzur, Soil Reclamation and Development Officer (FAO), emphasized FAO's role in providing advice on preventing soil degradation and managing problem soils. He also discussed the reactivation and expansion of the SPUSH Network to cover aspects of prevention of salinization as well as the management of salt-affected soils.

Introduction to the SPUSH Network

A. M. Mashali

Food and Agriculture Organization of the United Nations (FAO), Viale delle Terme di Caracalla, 00153 Rome, Italy.

E-mail: amin.mashali@fao.org

Soil salinization has been identified as a major process of land degradation. As the nature of the problem is dynamic, the cost of measurement is high and there is a lack of data required to classify a soil as salt-affected. To date there is incomplete information on the extent, distribution and degree of salinity development for all countries affected.

According to various estimates, the extent of salt-affected soils in the world differs considerably. This is due to the lack of systematic surveys, the continuous change in the extent of salinization due to secondary salinization/sodicitation or seawater intrusion and differences between countries' approaches for detecting and classifying salt-affected soils.

Many experts have tried to map salt-affected soils. However, figures and available maps are questionable and need to be updated, since they may be based on data and maps collected more than 30 years ago. There are differences between the countries in the diagnosis, classification and criteria they use to identify salt-affected soils, e.g. saline soils can overlap with sodic soils, acid sulphate soils, mangrove soils.

Therefore, practical methodologies for monitoring and mapping of salt-affected soils is a requisite for determining the extent and magnitude of salt-affected soils; for assessing the causes and sources of the problem as well as the effectiveness and appropriateness of irrigation and drainage practices; and for the effective management strategies.

To avoid the fragmentation of technical research and development efforts, FAO has recognized the need to reactivate and expand the Global Network on Salinization Prevention and Productive Use of Salt-affected Habitats (SPUSH).

This paper provides a brief overview of the causes of salinity and mapping efforts of salt-affected soils, as well as the previous work of the SPUSH Network.

**SESSION 1: ASSESSMENT AND MONITORING OF SALT-AFFECTED SOILS
AT FIELD, LANDSCAPE AND IRRIGATION DISTRICT LEVELS**

Assessing and monitoring the risk of salinization in a Sicilian vineyard using the Geonics EM38

G. Crescimanno¹, K. B. Marcum², F. Morga¹ and C. Reina¹

¹Università di Palermo, Dipartimento ITAF – Sezione Idraulica, Viale delle Scienze 13, 90128 Palermo, Italy.

E-mail: gcrescim@unipa.it

²Arizona State University, Department of Applied Biological Sciences, Mesa, AZ, U.S.A.

E-mail: kenneth.marcum@asu.edu

In many arid and semi-arid countries, as well as in Sicily, the increasing scarcity of good quality waters, coupled with the intensive use of soil under semi-arid to arid climatic conditions, results in irrigation with saline waters, leading to secondary salinization.

Salinization is closely associated with the process of desertification. Salinity may have direct negative effects on crop yields by reducing the ability of plant roots to take up water. The reduced availability of water to the plant is due to soluble ions and molecules causing an osmotic pressure effect. Threshold relationships between the soil electrical conductivity (EC) and crop yield have been empirically determined for several crops and can be used to evaluate the influence of saline irrigation water on agricultural production.

The salinity of irrigation water is defined as the total sum of dissolved inorganic ions and molecules. Soil salinity can be measured in the soil by determining the EC of the soil solution. The EC measured in the saturated extract (EC_e) is used as an expression of salinity. The USSL (1954) developed the saturation extract technique, a way to estimate soil salinity that uses a reference water content. However, methods suitable for a rapid assessment of soil salinity are necessary for surveying large areas susceptible to degradation and for prevention of desertification.

Application of an electromagnetic induction sensor (EM38) makes possible, after calibration, rapid surveys of large areas for identification of the places with the greatest risk of salinization, where detailed investigation is necessary to develop countermeasures and strategies suitable to control desertification. Application of this technique provides a measurement of the EC of the “bulk soil” (EC_a). Since EC_a is influenced not only by the chemical and physical properties of the soil solution, but also by those of the solid phase (soil texture, mineralogical composition of the soil), a preliminary calibration is necessary in order to convert the field measured EC_a into the saturated electrical conductivity (EC_e).

A number of international and national projects have been developed in Sicily since 1998, with the objective of developing integrated approaches for sustainable management of irrigation. This investigation is part of the “Evolution of cropping systems as affected by climate change” (CLIMESCO) project (2007–2009), and was carried out within the framework of a bilateral agreement for scientific cooperation between the University of Palermo and Arizona State University.

The objective of this paper was to show the results of using the Geonics EM38 probe (Geonics Limited, Mississauga, Ontario, Canada) for monitoring salinization in Sicily, where irrigation with saline water is increasingly practised and a risk of salinization and desertification is envisaged. Maps showing the spatial and temporal variability of salinity determined by two different irrigation treatments are analysed and the effect of using two irrigation waters of different salinity are discussed.

Use of an aboveground electromagnetic induction meter for assessing salinity changes in natural landscapes and agricultural fields

J. Arriola Morales¹ and J. Batlle Sales²

¹Facultad de Ingeniería Ambiental, Benemérita Universidad Autónoma de Puebla, Mexico.

E-mail: janette.arriola@icbuap.buap.mx, Janette.Arriola@uv.es

²Dept. Biología Vegetal, Universidad de Valencia, Spain.

Salinity can be considered as a property that changes quickly in time, influenced by lateral water movement through a landscape and vertical water fluxes in soils, due to infiltration and evapotranspiration. An irrigation event or the rainfall associated with an intense storm can change the amount and distribution of salts in a soil profile. For large areas, soil conditions can differ from one site to another, imposing local differences in management practices.

The nature of soil salinity/alkalinity is best studied through soil sampling and analysis of the saturated paste extracts and other soil properties. Within a framework of modern land management, some decisions are more rationally based if some soil geostatistical information is available. The collection of numerous field samples and subsequent analysis in the laboratory is a costly activity for assessing/monitoring soil salinity. On the other hand, the decision about the necessary number of samples to provide a uniform area cover, providing statistical information, as well as the decision about the sample location, poses a difficult question that should be answered after a previous salinity screening of the field. The ratio of the desired number of samples to the associated cost of sampling can be optimised through an adequate sampling design.

Electromagnetic induction (EM) is a useful non-invasive technique for taking quick measurements. It can provide raw information about soil components (salts concentration, soil moisture and texture). The EM signal response can be related to the apparent soil electrical conductivity (ECa) at particular depths, through statistical calibration. In a conceptual model, the EM signal response can be described as a complex function of soil solution conductivity, soil moisture, temperature and amount and type of clay, among other factors, each soil depth contributing unevenly to total signal response.

The two examples presented show the way to derive soil salinity information from above-ground electromagnetic induction measurements. The first example is a multitemporal monitoring of salinity in a small agricultural field with uniform texture and topography, with sodium chloride-type salinity, using a regular-grid surveying scheme. Two sensor geometries are used and, based on the signal treatment, quasi three-dimensional multitemporal maps of bulk soil salinity can be drawn and trends of soil salinity recognized.

The second example is an automated longitudinal survey along a landscape, designed for discovering differences in salinity type, soil texture and total moisture content. Soil sampling sites are selected from the profiling and statistical treatment of response signal.

Keywords: electromagnetic induction, soil salinity, salts zonation.

Primary soil salinity, sodicity and alkalinity status of different water management areas in South Africa: The quest for baseline data

J. P. Nell, M. C. de Villiers and M. van der Walt

ARC-Institute for Soil, Climate and Water, Private Bag X79, Pretoria, 0001 South Africa.
E-mail: pnell@arc.agric.za

The objective of this paper was to describe and quantify the primary salinity, sodicity and alkalinity status of soils in terms of the 19 water management areas of South Africa by using data from more than 26 520 soil samples. South Africa's salt-affected soils, derived from complex geological formations and soil forming processes, comprise almost 32 percent of the country's surface area.

The data were derived from soil survey reports for irrigation and environmental planning and the South African Land Type Survey of the ARC-Institute for Soil, Climate and Water. The minimum requirements for inclusion in the data set were: (a) the profiles should have comprehensive chemical and physical analyses, and the preference was given to data sets where soil analyses followed the methods of the *Non-Affiliated Soil Analyses Working Group* (1990) and where the analyses were done in the ARC-ISCW laboratory; (b) accurate profile location information should be available; (c) only primarily data could be used – no human-induced salinization or sodication; (d) soil profile description should have been done according to *Soil Classification: A Binomial System for South Africa* (1977) or *Soil Classification: a Taxonomic System for S.A.* (1991). Although data verification was carried out on most samples previously, much effort was devoted to data cleaning. From the original data (in excess of 40 000 data points), only 26 520 data points were used due to the stringent cleaning protocol.

After a countrywide process of public consultation, 19 water management areas covering the entire country were established. The boundaries of the water management areas lie mostly along the divides between surface water catchments and do not coincide with the administrative boundaries that define the areas of jurisdiction of provincial and local government authorities. The water quality data that were used in the assessment of fitness for use of South Africa's surface water resources for domestic and irrigated agricultural use were collected as part of the National Chemical Monitoring Programme. This programme has been in operation since the early 1970s and samples are collected regularly at approximately 1 600 monitoring stations at a frequency that varies from weekly to monthly sampling. The data collected is stored on the Department of Water Affairs and Forestry's database and information management system, namely the Water Management System. For this study, a suitable water quality sample site was considered to be one with an adequate level of sampling (not too infrequent or sparse) over the study period.

The problems of primary soil salinity, soil sodicity and soil alkalinity are most widespread in the arid and semi-arid water management areas of South Africa, but salt-affected soils also occur extensively in sub-humid and humid water management areas, particularly in the coastal areas where the intrusion of seawater through estuaries, rivers and groundwater causes problems. In South Africa, where the rainfall is approximately five to ten times less than the potential evaporation, salts derived from rock weathering, bio-cycling and atmospheric deposition may accumulate to relatively high levels in the soil.

Currently, the secondary salinity and sodicity of South African waters and soils are on the increase due to mining, urban, industrial and agricultural developments and the re-use of water resources. Irrigated agriculture is not only at the receiving end of water quality deterioration, but it is itself a major contributor to the observed water quality deterioration in many rivers. The use of such water poses a future threat for soils in areas of South Africa where leaching is limited.

In South Africa, many water bodies are naturally high in dissolved salts, especially where rivers flow over marine sediments. Irrigation farmers are often accused of causing much of the salinization, sodication and alkalization of water in South Africa, because they are the biggest users of water. Leaching of primarily mineral salt from the surrounding geology after rainfall, as well as pollution from industrial and mining activities are, incorrectly, not considered as also being important. Soil salinity seems to be largely under control in South Africa at present, but this situation could change dramatically as a result of industrial, municipal and mining effluents used on agricultural land. Primary soil sodicity and alkalinity are a bigger problem than primary salinity in South Africa.

Using satellite imagery and electromagnetic induction to assess soil salinity, drainage problems and crop yield in the *Río Fuerte* Irrigation District

L. Pulido Madrigal and J. Gonzalez Meraz

Instituto Mexicano de Tecnología del Agua. Paseo Cuauhnáhuac 8532, Progreso 62550, Jiutepec, Morelos, México.

E-mail: lpulido@tlaloc.imta.mx

The survey of soil salinity and drainage deficient areas, as well as the estimation of yields in the *Río Fuerte* Irrigation District, northwest Mexico, were carried out by applying satellite imagery, a portable electromagnetic-type sensor and a GPS unit. Soil and plant samples were obtained in selected salt-affected fields grown with wheat (*Triticum aestivium*), cotton (*Gossypium hirsutum*), sorghum (*Sorghum bicolor*) and maize (*Zea mays*), which were considered as the reference crops, since all together covered most of the cropped area of the irrigation district. Spectral values (TM2, TM3 and TM4 bands) were extracted from Landsat TM images. The images were obtained during the flowering stage of each crop. Salinity and spectral data were analyzed and multiple regression models were obtained to estimate the salinity status of the cropped areas, together with its correspondent crop yield. Salinity and yield maps for each crop were digitized and classified on the Landsat image using the regression models. The non-referenced area of the district (fields planted with any other crops, non-cropped, fallowed, or abandoned) was mapped *in-situ* using an EM38 electromagnetic sensor, along with a GPS unit to locate the geographic coordinates of each of the sites. Both Landsat- and EM38-based salinity maps were joined and a final map covering all the Irrigation District area was obtained. The total mapped area was 319 976 ha and it was estimated a salt-affected area of 138 345 ha (43 percent, $EC >4 \text{ dS m}^{-1}$). Poorly drained areas were also obtained based on water table monitoring for the most critical period of the year. This information was also added to the salinity map previously obtained, so salt-affected areas with shallow water table were determined for reclamation and planning purposes. Yield maps were also combined with the salinity map to analyze the effect of the various existing salinity levels on crop yield. The economical analysis performed showed this method is highly cost and time effective, compared to the traditional one (extensive soil sampling and laboratory analysis).

Keywords: satellite imagery, salinity, drainage, yield, regression model, electromagnetic sensor.

**SESSION 2: ASSESSMENT AND MONITORING OF SALT-AFFECTED SOILS
AT NATIONAL AND REGIONAL LEVELS**

Soil salinity monitoring and assessment in irrigated Arab agriculture

G. Abdelgawad

Scientific Adviser to the General Director, the Arab Center for the Studies of Arid Zones and Drylands (ACSAD), Damascus, Syria.

E-mail: a-abdulgawad@acsad.org, abdelgaw@scs-net.org, gmhimed@yahoo.uk

In the Arab world, irrigated agriculture contributes about 70 percent of the total crop production commodities. Irrigated agriculture in the Arab world is about 10.7 Mha. The irrigation systems used for irrigating such land are flood, sprinklers and localized. About 87.6 percent for the irrigated areas is irrigated with different flood irrigation practices, 11 percent with sprinkler and 1.4 percent localized. The efficiencies of the irrigation systems used are not more than 50 percent. These low efficiencies of irrigation systems lead to increase in soil salinity and waterlogging. In addition, due to the shortage of good quality water, several Arab countries have large reservoirs of low quality ($3\text{--}11\text{ dS m}^{-1}$) underground or agricultural drainage water, as well as sewage drainage water. Use of such water is possible if it is properly managed. The use of low quality water must be used with extensive planning and proper infrastructure of drainage networks.

Good agriculture practices will minimize salinity build-up in the soil profile. These agricultural practices are: good irrigation scheduling; addition of organic manures; deep ploughing; and no tillage practices. Addition of soil inorganic and organic amendment has been used to prevent soil alkalization. Low quality water has been used by the Arab Center for the Studies of Arid Zones and Drylands (ACSAD) in different Arab states for irrigating winter crops such as barley, wheat and potato. This supplementary irrigation practice improved the income of poor farmers without degrading their fields by salinity. This finding was determined by salinity monitoring for seven years in more than 80 farmers' fields and in ACSAD experimental stations over fourteen years. The salts in these fields are leached out as far as 3.5 m below the soil surface in sandy and sandy loam soils. Measurement of salts in the soil profile indicated that salts moved down to drainage systems in clay, clay loam and silt clay.

Different methods were used by ACSAD for monitoring and assessment of soil salinity. These included field methods, such as EM38, salinity sensors and traditional methods of collecting soil samples and determining the salinity.

Each method has its limitations and some of them gave wrong the interpretation and correlation with yield increase or decrease if not carried out correctly. It was found that the method that correlated most closely with yield increase or decrease was when the soil solutions were collected by ceramic cups.

The salt balance for irrigated Mediterranean coastal soils shows most salts, which accumulate in summer crops, were leached out by rain in winter, when the soil profiles were more liable to leaching.

Advances in the assessment of salt-affected soils for mapping, monitoring and management strategies in India

G. Singh and A. K. Mandal

Central Soil Salinity Research Institute, Karnal - 132 001, India.

E-mail: dr.gurbachan@gmail.com, director@cssri.ernet.in

Salt-affected soils in India have variable distribution patterns and occur mostly in belts or patches that make them difficult to map and monitor. The first mapping of salt-affected soils in India began in 1902, when Leather delineated several *Usar* (alkali) patches enriched with sodium carbonates in the fields of the Etah district in Uttar Pradesh. After independence in 1947, several workers across the country conducted traditional soil surveys including field traversing, profile studies and analysis of soils and water samples to produce reports and maps on salt-affected soils falling under various projects and irrigation schemes. Such surveys produced fragmented and localized results. They differed in methodology and criteria; hence the information produced could not be extrapolated to other areas. The Committee on Natural Resources of Planning Commission, in 1966, estimated 6 Mha of salt-affected soils and depicted its distribution in a map, based on standard surveys using aerial photo interpretation. The map showed occurrence of salt-affected soils in compact areas of the Indo-Gangetic Plain and in scattered blocks in other parts. After its establishment in 1969 at Karnal, the Central Soil Salinity Research Institute (CSSRI) carried out a countrywide assessment of the problem and made a first estimation about the nature and extent of salt-affected soils. By focusing on soil characteristics and the feasibility of reclamation, two main classes of salt-affected soils, namely alkali and saline, were recognized. Abrol and Bhumbra upgraded the estimate to 7 Mha in 1971 and Bhumbra, in 1975, depicted six categories of salt-affected soils on a map, i.e. alkali soils, saline soils, potentially saline soils, coastal saline soils, deltaic saline soils and acid sulphate soils. In 1980, Murthy compiled and synthesized benchmark profiles of salt-affected soils from all over India and classified them into associations of major groups. The mapping legend consisted of 12 associations. The absence of systematic surveys and a reconnaissance map showing the distribution and extent of salt-affected soils prompted the individuals to report on their extent. The calculated figures, as expected, ranged from 3.3 to 26.1 Mha. The wide variations in the figures reflect the degree of concern and perception authors had for the problem.

The recent mapping methodology includes visual (manual) and digital (computer) interpretation of satellite data supported by field and laboratory investigations. The first systematic mapping of salt-affected soils of the entire country was made in 1996 by the National Remote Sensing Agency (NRSA) in Hyderabad, in association with other national and state level organizations including the CSSRI and the National Bureau of Soil Survey and Land Use Planning (NBSS and LUP of ICAR). A total of 125 false colour composite prints of the Landsat TM satellite images were used in mapping salt-affected soils at a 1:250 000 scale. The methodology consisted of development of a nationwide mapping legend, interpretation of satellite data, ground truth collection, analysis of soil samples, post field interpretation and reconciliation and area estimation. In order to accommodate salt-affected soils occurring in different parts of the country, a common legend was evolved after extensive discussions with the collaborating partners engaged in either conventional soil survey or those working with remote sensing techniques. The 125 map sheets showed a 6.73 Mha salt-affected area in the country. However, it did not differentiate between saline and sodic soils, the two major classes for which distinct sets of reclamation techniques are evolved under Indian conditions.

By abstracting information from the mapping legend – kind of soils, clay minerals, climatic conditions and physiographic setting of various regions – the area of 6.73 Mha of salt-affected soils was divided into two major classes (saline and sodic). Moderate and highly saline soils were easily identified due to the presence of white salts on the surface. A combination of red and infra red bands helped in separating saline and sodic soils. Integration of thermal band interpretation with false colour composite helped in resolving the problem of spectral similarity between saline soils and sand dunes.

The use of GPS in surveys of salt-affected soils has improved the quality of comparable studies like improvement in soil properties in the post reclamation phase, precise detection of hot spots of salinity emergence, expansion, identification and establishment of benchmark sites of salt-affected soils. Furthermore, the CSSRI has developed sustainable and cost-effective technologies for the reclamation and management of salt-affected soils.

The state-wide distribution and extent of saline and sodic soils is presented in the paper. A brief account of these technologies and their impact in terms of productivity gains, socio-economic dimensions and environment is also discussed.

Salt-affected soils in Thailand: Assessment and monitoring of salinization

R. Im-Erb and S. Sukchan

Land Development Department, Chatuchak, Bangkok, Thailand.

E-mail: rungsun@ldd.go.th

Agriculture has long been important for the socio-economic growth of Thailand. About 21 Mha of the land area is currently under cultivation, 13 Mha are forests and the remaining 17 Mha are composed of urban development, public areas, sanitation, swamp land, railroads, highways, real estate and others. In terms of agricultural production, land has been used to provide for the rapidly increasing food demand, which increased along with the population growth. This has led to diverse use of the land for many purposes, eventually resulting in degradation of land, particularly anthropogenic soil salinization of arable land.

Salinity is the oldest and one of the most important environmental problems. The problem of salinization is the presence of excessive salts on the top layer of the soil, resulting in the deterioration of its chemical and physical properties. It is a form of land degradation that has become a major cause of low agricultural productivity in the Northeastern region, the Central Plain and along the coastal belt of the country.

Salt-affected soils cover approximately 3.5 Mha. These soils are classified broadly into two groups, i.e. inland saline soils and coastal saline soils. These two broad categories of saline soils cover approximately 3.0 and 0.58 Mha, respectively. These naturally occurring saline soils normally expand to cover wider areas with time. Anthropogenic soil salinization has also become an important part of the rapid increase in soil salinization.

The assessment and monitoring of salinization for managing salt-affected land in Thailand has been greatly facilitated by various methods. The goal of this process is to assess the influence of salinity, preventing distribution of salinity on arable land and to manage the improvement and sustainable utilization of land resources. The criteria in defining salt-affected areas are based on an electrical conductivity (EC) value of 2 dS m^{-1} , while water has an EC value of 0.75 dS m^{-1} . Methods were developed to complement the soil salinity classification system, to evaluate and predict the existing salt-affected areas and to develop appropriate soil management guidelines. The conventional method was used in 1963 to assess and monitor soil salinity. This new method applies integrated technologies that have been assessed in terms of measurement of EC and laboratory analysis of soil and water samples, including determination of vegetation and plant growth.

Since 1989, advanced technologies for the assessment and monitoring of soil salinity have been used to provide spatial coverage faster than conventional methods. Image analysis of remote sensing data (Landsat) using visual and digital methods is carried out and verified by ground truth data and laboratory investigations. False colour composite bands 5, 3, 1 (blue, green and red) showed promising data for visual interpretation. Ground based surveys, using electromagnetic induction (EM34 and EM38), were conducted to measure bulk soil conductivity to a depth of approximately 7, 15 and 30 m, while EM38 measured conductivity over a depth of approximately 1.5 m. Piezometer installation on the salt-affected area was also used to observe and monitor water quality, depth of water table and flow.

The Land Development Department has been implementing the assessment and monitoring of salinization using the above-mentioned methods in different areas for several decades. The prototype of soil survey and monitoring, remote sensing and some

computer models, were applied to generate maps of salt-affected areas and to obtain manuals and reports on appropriate prevention of soil salinization and improvement of technologies and approaches, database of soil resources and salinity management guidelines (including assessment and prediction of soil salinization and distribution). Outputs have been used to support stakeholders in the salt-affected areas, e.g. maps of salt-affected areas in the Northeastern region and the Central Plain provinces, a map of reforestation areas for soil salinity control, a report on the impact of the tsunami in the West Coast area, the impact of shrimp farming on arable land and monitoring of reforestation to prevent soil salinization.

An overview of the salinity problem in Iran: Assessment and monitoring technology

S. A. M. Cheraghi, Y. Hasheminejhad and M. H. Rahimian

National Salinity Research Center, Azadegan Blvd., Nahalestan Ave., Yazd, Iran.
E-mail: cheraghi@insrc.org

Agriculture plays a vital role in the national economy of Iran. It accounts for 27 percent of the GNP and 23 percent of the labour force of the country. Scarcity of water is the major constraint for agricultural development in the country and based on projections for 2025, Iran will be in a water scarcity situation. Considering the growing demand for water from other sectors and the fact that 93 percent of renewable water resources are already allocated to the agricultural sector, allocation of more water to the sector is not foreseeable in the future. Under these circumstances, increasing water productivity in agriculture holds the greatest potential for improving food security and preventing environmental degradation.

The exact extent of salt-affected soils in Iran is not known. Based on a recent estimate, 34 Mha or nearly 20 percent of the surface area of the country is salt-affected. This includes 25.5 Mha of slightly to moderately and 8.5 Mha of severely salt-affected soils. Salt-affected soils are mainly distributed in the central plateau, southern coastal plain, Khuzestan plain and inter-mountain valleys. The salinization of land and water resources has been the consequence of both naturally-occurring phenomena and anthropogenic activities. Secondary salinization has been the main cause of the spread of salinity and, as a result, the waterlogged area has increased from 160 000 ha in 1977 to 0.7 Mha today, only in areas under the command of new dams.

In spite of the extent of soil salinity in the country, there are few studies which have been carried out to assess and monitor soil salinity. A number of these studies have been concerned with the evaluation of soil salinity in irrigated fields as a means of evaluating the appropriateness of on-going management practices. In these studies, soil salinity has been assessed in terms of laboratory measurement of electrical conductivity. The results from one such study in the south-west of the country show that wheat fields irrigated with saline water of electrical conductivity ranging from 8–12 dS m⁻¹ produced a relatively good yield of 4–6.5 tonnes ha⁻¹. Monitoring of root zone salinity throughout the growing season for three years showed that mean root salinity in almost all the fields was nearly equal to the electrical conductivity of irrigation water, suggesting a high leaching fraction as a result of poor on-farm water management.

The use of advanced technology for assessment and monitoring of soil salinity based on bulk soil electrical conductivity is still in its early stages. Although instrumentation such as EM38, four electrode probe and time-domain-reflectometry (TDR) are now available, their actual field use has been very limited. In a case study an EM38 device was used for soil salinity survey in a pistachio orchard in central part of the country. Results indicated that the device could be conveniently used as a rapid measurement tool for soil salinity appraisal both in surface and depth.

On the other hand, remotely sensed data, especially satellite images, have been used more often for assessing and monitoring soil salinity in different parts of the country. Most of these studies utilized images obtained by Landsat TM and ETM sensors combined with other auxiliary data such as field measurements, DEM, slope and aspect maps. Methodologies used included visual interpretation of FCC map, image classification (both supervised and unsupervised) and selection of effective band(s) or indices.

Advances in the assessment and monitoring of soil salinization for managing salt-affected habitats in Egypt

M. H. Gomaa

Head of Land Reclamation Sector, Ministry of Agriculture and Land Reclamation, Egypt.
E-mail: gomaa_1999@yahoo.com

Agricultural development in Egypt is facing human-induced degradation, mainly due to salinity and waterlogging. The causes include seepage from irrigation canals, inadequate drainage systems, poor irrigation practices, use of slightly saline (drainage or mixed) water for irrigation without proper management and inappropriate agronomic practices. The presence of excess salts in the soil directly or indirectly affects plant growth. Salinity can affect the survival of normal field crops leading to completely barren soils which need to be reclaimed. As a result, soil salinity has become one of the most frequently used parameters to characterize field variability for application to sustainable agriculture.

The value of spatial measurements of soil electrical conductivity (EC) to sustainable agriculture is widely acknowledged, but soil EC is still often misunderstood and misinterpreted. The following areas are discussed with particular emphasis on spatial EC measurements: a brief introduction to soil degradation, soil salinity (worldwide) and the extent and causes of salt-affected soils in Egypt. This paper also presents an overview of advances in technology used for assessing and monitoring soil salinity such as remote sensing (RS), geographic information systems (GIS), mobile instrumental systems and computer assisted expert systems. These advances in technology can contribute to the planning and implementation of measures to combat salinization. Selecting the method of detection depends on several factors such as the accuracy required, cost of the project, available funds and the area to be surveyed. The future of sustainable agriculture rests on the reliability, reproducibility and understanding of these technologies.

Recent evolution of soil salinization and its driving processes in China

J. Yang

Institute of Soil Science, Chinese Academy of Sciences, 71 East Beijing Road, Nanjing 210008, China.
E-mail: jsyang@issas.ac.cn

Salt-affected soils are spread extensively in China, covering a wide area from tropical to temperate zones, from the coast to the inland and from the semi-arid to desert regions. Currently the total area of salt-affected soils is about 36 Mha, which occupies 4.88 percent of usable land in the whole country. About 9 Mha of arable land is salt-affected, accounting for 6.62 percent of the total land in the country.

Salt-affected soils are mainly distributed in Northwest China, North China, Northeast China and coastal regions. All these areas have hot spots of soil salinization. In the Songnen Plain and the Great Bend region of the Yellow River, the extent and degree of salinization are increasing. The area of salt-affected soils reaches 3.5 Mha in the Songnen plain, with an annual rate of salinization increase as high as 1–1.4 percent. About 45 percent of salt-affected land in this area has been degraded to severe saline land and abandoned. Salinity in the Great Bend region of Inner Mongolia has increased 1–3 percent annually in the last three decades due to inappropriate irrigation. Similarly, secondary salinization has occurred in the West Corridor of Gansu.

Irrigation with brackish water has been an important cause of salinization in the north part of China. According to monitoring data from Guyuan region in Ningxia, soil salt content increased to 2.3 g kg⁻¹ after 5 years of brackish water irrigation and up to 8.3 g kg⁻¹ after 14 years. Shallow ground in most river irrigation districts of Ningxia, Inner Mongolia and Xinjiang has been observed and salinization occurred as a result. The salt-water regime has changed under drip irrigation practices in some regions of Xinjiang, causing salt accumulation. Approximately 35 percent of irrigated land in Gansu, Xinjiang and Ningxia and 50 percent irrigated land in Inner Mongolia, are currently risking salinization. There are also imbalances in salt-water movement in coastal regions, such as the Yellow River delta.

The main driving factors of evolution of salinization in China are inappropriate water resource and land management as well as climate change. Inefficient water use is the dominant cause of soil salinization in irrigation districts, due to large seepage of irrigation canals, low irrigation efficiency, unnecessary high irrigation norms, poor drainage systems, blockage of drainage ditches and the rise of groundwater table from the construction of plain reservoirs. Micro irrigation practices such as drip irrigation in arid regions can potentially change the salt-water regime and therefore may cause salt accumulation. Local salinization has already been found in some areas with drip irrigation and such phenomena tend to expand.

Inappropriate land management in semi-arid and arid regions is also a driving factor in accelerating salinization, worsening soil physical and chemical properties (due to poor fertility management), increasing the evaporation surface and impairing soil drainage by removing vegetation cover from over-grazing, deforestation.

Climate change is considered as another significant driving factor on salinization. Currently climate in North China and Northeast China has been warming and precipitation has decreased. Such changes have promoted the evolution of salinization in the region. Climate change may also cause sea level rise. Salt leaching from soils in coastal regions, including the Yellow River and Changjiang River deltas, is therefore subject to a disadvantage owing to blockage of salt outflow and seawater intrusion. Consequently, local salinization expansion is observed.

Assessment and management of salt-affected soils in Sudan

A. A. El-Mobarak

Associate Professor, Land Evaluation Research Section, Land and Water Research Centre, P.O. Box 388, Wad Medani, Sudan.

E-mail: melmobarak@yahoo.com

Sudan, the largest country in Africa, has an area of approximately 2.5 million km². Its climate is diverse, ranging from tropical humid in the south to desert in the north. This paper highlights the changes in climatic conditions and the properties of salt-affected soils in relation to salinity and sodicity. The distribution and area covered in every region is shown. The system adopted in Sudan for the assessment of a soil as being salt-affected is derived from the USDA system with amendments to fit the Sudan conditions (based primarily on the results of research trials). The research activity in reclamation of salt-affected soils is also covered and some of the results of the trials are shown.

The studies on climatic changes showed that there has been a considerable decrease in the annual rainfall. It reaches 19 percent in the arid climatic zone and 1 percent in the semi-arid climatic zone compared to the averages for 1941–1970 and 1971–1999. This, combined with human misuse of natural resources, water scarcity and removal of vegetation cover, necessarily implies a deterioration in soil properties. The increase in population and the limited extent of good agricultural lands in the Khartoum area and along the river banks in the northern regions have led to expansion into the marginal lands of the higher terraces of the Nile, where soils are affected by varying degrees of salinity and/or sodicity.

In many areas of the world the presence of salts (including sodium) in arid and semi-arid zones is a widespread phenomenon and Sudan is no exception. The salinity or sodicity may result as a consequence of physical weathering under low rainfall conditions coupled with low humidity and high temperatures by evaporation or by capillarity. All these conditions exist or have previously existed in Sudan. The reclamation of salt-affected soils was very limited until the 1970s, and was confined to some experiments in the Gezira, Soba and Hudeiba research stations. In the 1970s, Sudan witnessed some environmental, political, scientific and demographic changes that accelerated the need for reclamation of salt-affected soils, including:

- the decrease in good agricultural lands due to desertification, sand encroachment and river bank erosion that led to thinking about expansion into new lands to cover the food gap and support exports;
- the adoption of a local government system, so the different states tried to develop and exploit their resources, e.g. the northern states – as wheat-producing areas – expanded their areas into the high terrace salt-affected soils;
- the foundation of the Land and Water Research Centre, to classify, evaluate and reclaim salt-affected soils.

Salt-affected soils in Sudan fall under three soil orders: Vertisols, Aridisols and Entisols. They extend along vast areas at latitudes 14–22° N, including the White Nile, North Gezira, Khartoum state, along the river Nile, crossing the river Nile and the northern states. The land suitability classification system adopted in Sudan by the Land and Water Research Centre is derived from the FAO system, but some amendments were made to fit the Sudan conditions, e.g. exchangeable sodium percentage (ESP) values of up to 35 in the topsoil and 50 in the subsoil were incorporated as acceptable limits for the potential rating of the soil as marginally suitable in Gezira. These values were generalized over other soils in Sudan. Very good efforts were made for reclamation by

researchers from agricultural research stations in Gezira, Hudeiba and Soba and the farm of the University of Khartoum. The Land and Water Research Centre and other experts reported that about 268 636 ha were found to be affected by salinity and/or sodicity.

From research results on the reclamation of salt-affected soils in Sudan, it was concluded that gypsum is a good amendment for leaching salts from the topsoil, while organic amendments like farmyard manure, chicken manure and dry sewage increased crop yields. The strategy for research should focus on a well-defined soil series, for ease of transferring the technology to other similar soils. Encouraging breeding programmes for salt and heat tolerance is of paramount importance, as is the promotion of extension programmes to transfer the technological packages to farmers. Finally, the Land and Water Research Centre of the Agricultural Research Corporation is carrying out a very important programme to have a soil database of all the soils in Sudan with special reference to salt-affected soils. We now have an up-to-date geo-referenced soil information database with digital maps.

SESSION 3: MODELLING FOR SALINITY/SODICITY DEVELOPMENT

Overview of salinity modelling approaches at different spatial-temporal scales

J. Batlle Sales

Departamento de Biología Vegetal, Universidad de Valencia, Spain.

E-mail: Jorge.Batlle@uv.es

Salinity is a natural property of many natural environments, arising from the geomorphic and geochemical processes creating landforms and coastal areas. During recent decades, the environmental role of natural salt-affected ecosystems has been recognized and the conservation and restoration of such habitats has become a priority policy worldwide. On the other hand, salinization is a progressive, human-driven, soil and water degradation process and an increasing environmental concern that affects the most productive agro-ecosystems under irrigated agriculture in arid and semi-arid regions.

Modelling is a valuable mathematical tool for studying salinization, dealing with simplified representations of systems; revealing complex interrelations of properties of the system under study; and creating scenarios for investigating “what if?” questions. Modelling of natural saline environments, as well as agricultural environments subject to salinization, has been addressed in several ways, from deterministic soil-plant-water-atmosphere approaches to systems dynamics model development, including social, economic or cultural factors, among others.

Each modelling exercise tries to provide an answer to a particular question formulated; hence the input information required to run the models ranges in complexity, as do the data acquisition requirements. Some models are used only by researchers whereas others are affordable by farmers. Many models are dynamic in concept, while others reflect a steady-state condition. The scale of application, geometry of the system, biological, chemical and physical processes represented, as well as the capability of representing an evolving system, are the main differences between models.

Some aspects that are important under normal agricultural practices are not well reproduced by some codes nowadays. Management practices, geometry of irrigation and evaporation, sinks of solutes (plant uptake) and interaction of fertilizers with soil components are incorporated in an uneven way in the available codes, and should be developed further.

Although it is desirable that the models become more and more deterministic and mechanistic, this requires a very intensive effort in research and computing. There are uncertainties associated with the values of the input parameters, with the computation procedure or with the inaccurate description of the system. The parameter estimation, analysis of sensitivity and validation procedures are refinements applicable to most models.

This paper gives an overview of different modelling approaches; compares the most used models based on their implementation, advantages and limitations; and discusses the possibilities of nested approaches and the advantage of clearly establishing an exit strategy in the management plans.

Keywords: Soil salinity, salinity modelling.

SMSS: a soil-plant model describing the impact of irrigation on soil and runoff water salinity

M. Lahlou and M. Badraoui

Institut Agronomique et Vétérinaire Hassan II, Po Box 6202, Rabat-Instituts 10101, Morocco.

E-mail: m.lahlou@iav.ac.ma

In arid and semi-arid regions, water is the main limiting factor for agricultural production. In fact, both the quantity and the quality of irrigation water fundamentally affect soil quality and crop production. The presence of salts in irrigation water and the evaporation potential in the irrigated areas usually lead to the salinization and alkalization of soils, particularly in arid and semi-arid zones. The monitoring of both soil and water qualities in irrigated areas is necessary to measure the sustainability of the production system. Thus, modelling the movements of salts in irrigated soils is a means to predict their evolution. For this purpose, two models (SMSS2 and SMSS3) were developed in order to predict the impact of irrigation on soil and water qualities.

The first model (SMSS2), based on the Laudelout model (1994), is used to analyze the movements of salts in saturated conditions with perfect drainage. It is a simple model essentially allowing the prediction of salinity/sodicity of soil after a long period of irrigation. It is controlled by two types of parameters: the exchange selectivity coefficients and the volume and quality of irrigation water.

The second model (SMSS3), improved from LEACHM (Hutson *et al.*, 1992) and UNSATCHEM (Šimůnek *et al.*, 1993) models, is used to simulate water and solutes transport in partially saturated soil profiles. Over a short period of time it allows the follow-up of the soil salinity and sodicity, which are controlled by different parameters such as: (i) simulation period, (ii) the conditions at the boundaries, (iii) soil physical properties, (iv) soil chemical properties (exchange selectivity and ionic activity products), (v) the crop, (vi) the temporal distribution of applied chemicals, tillage, rain and irrigation water and (vii) weekly distribution of the crop coefficients and potential evapotranspiration.

The validation of the SMSS2 model with the data of the lower Euphrates valley in Syria showed that the simulated data agreed with those measured in the field. The mean differences found between the simulated and the measured data in the field varied between 0.5 and 1 dS m⁻¹.

The validity of SMSS3 model, applied to the Doukkala region in Morocco, allowed predicting with acceptable precision the evolution of soil salinity and sodicity in the studied area. The obtained mean differences between the measured data in the field and those simulated vary between -0.25 and 0.13 dS m⁻¹. A sensitivity study of this model showed that the use of ionic activity products (IAP) of soluble minerals is preferable to their solubility products. For example, the use of the IAP of calcite produced a gain of approximately 18 percent in precision of the electrical conductivity in comparison with the result obtained using its solubility product.

Keywords: salinity, alkalinity, irrigation, modelling, Euphrates, Syria, Doukkala, Morocco.

SESSION 4: MAPPING AND INTERPRETATION OF SPATIAL DATA

Emerging challenges addressing the characterization and mapping of salt-induced land degradation

M. Qadir^{1,2} and E. De Pauw¹

¹International Center for Agricultural Research in the Dry Areas (ICARDA), Aleppo, Syria.

E-mail: m.qadir@cgiar.org

²International Water Management Institute (IWMI), P.O. Box 2075, Colombo, Sri Lanka.

Salt-induced soil degradation is an inherent and natural feature of the arid and semi-arid regions. Over the last few decades, it has increased steadily in several major irrigation schemes throughout the world. This has triggered imbalances between the goods and services supplied by the natural resources (land and water) and the demands of societies. Since soil degradation occurs both 'on-site' and 'off-site', it affects the livelihoods within and outside the farming communities. Therefore, salt-affected soils are usually considered as a major environmental and agricultural productivity constraint. However, these soils are a valuable resource that cannot be neglected, especially in areas where significant investments have already been made in irrigation infrastructure. From the perspective of managing salt-affected soils for economically feasible productivity enhancement, it is imperative to characterize different types of salt-affected soils for suitable technical interventions. Several efforts have already been done in the past to characterize salt-affected soils through different approaches. Owing to the complexity of different processes leading to the development of salt-affected soils and the use of a variety of assessment criteria, there is no single system that fits well with the characterization and classification of these soils.

Understanding the processes that lead to natural or anthropogenic soil salinization is as important as the characterization of salt-affected soils. From an agroecological perspective a distinction needs to be made between primary and secondary salinization. Primary salinization refers to the build-up of salts as a result of lithological inheritance or topographical position and is a natural process in arid zones. Secondary salinization is not linked to natural factors, but involves anthropogenic activities related to inappropriate irrigation management using both freshwater and saline water without drainage provision, groundwater depletion, and/or seawater intrusion.

The current status of salinity assessment in the dry areas is unsatisfactory. There is a lack of good information on the actual location, extent and severity of the salinity problems in most countries. This has severe implications on the ability of the respective governments to formulate appropriate policies and target investment towards control or reclamation projects. Although soil salinity is easy to detect, most soil maps, particularly at small scales, show primary salinity because of its association with geological or geomorphological features, which are easy to map at these scales. The mapping of secondary salinization is more complicated because of high spatial and temporal variability. Therefore, reliable figures are however hard to obtain. Secondary salinity is spotty both in horizontal and vertical distribution and can affect any soil type, which makes it difficult to obtain representative sampling points from which to extrapolate its true extent. Secondary salinity responds more rapidly to management practices than primary salinity and is therefore more difficult to capture in classical soil surveys, which provide one-time shots of a highly dynamic picture.

Mapping salinity is expensive and in order to achieve a rapid and effective assessment of its extent and severity, a multi-scale strategy is essential. The key principle is to build up an understanding of the spatial variability at different scales using indicators appropriate for each scale. These indicators can include spectral signatures from

different remote sensing platforms, secondary data such as existing thematic maps, direct field measurements using appropriate sampling schemes and interpolation methods and participatory assessments with farmers along transects. Used in a practical and complementary manner, these approaches will help to zoom rapidly into the salinity hotspots at different resolutions and be far less costly than the classical field surveys, even where assisted by geostatistical tools.

Given the highly dynamic nature of salinity, assessments of extent and intensity need to be undertaken at regular intervals for trend monitoring and require an institutional setting. The most effective way might be by adapting the mission and mandates of the national institutions with the capacity of undertaking soil surveys and land resource studies to the new information requirements of development projects and environmental monitoring agencies. The United Nations Convention for Combating Desertification (UNCCD) could be a useful framework for capacity building in long-term salinity monitoring.

Monitoring, predicting and quantifying soil salinity, sodicity and alkalinity in Hungary at different scales: Past experiences, current achievements and outlook with special regard to European Union initiatives

T. Tóth

Research Institute for Soil Science and Agricultural Chemistry of the Hungarian Academy of Sciences, Budapest.

E-mail: tiber@rissac.hu

Approximately 13 percent of Hungary is considered to be salt-affected and with this large extent it is unique in Europe. There are large areas of naturally saline and sodic soils, but secondary salinization is also known to occur.

Due to the geological and hydrological conditions, the country demonstrates the most characteristic features of natural continental (not marine) salinization, sodification and alkalization. Since the most important direct source of soil salinization is the shallow groundwater level below the lowland surface, there is a chance of irrigation-related salinization in two dominant situations: when the abundant use of river waters causes waterlogging and rise of saline groundwater (salinization from below); and when typically saline tubewell-waters are used for irrigation (salinization from above).

The spatial assessment of salt-affected areas began with the systematic mapping of salt-affected areas. There is a series of ten maps describing different aspects (salt-affected soil types, vegetation types, salt-efflorescences) of the salinity-status nationwide from 1897 onward, with the latest survey finished in 2006.

Besides the national scale of 1:500 000, soil salinity is also mapped at the scale of 1:100 000 on the AGROTOPO map sheets and 1:25 000 in the Kreybig Soil Information System (spatial vector data for maps and database for profiles and test boring). In spite of the two systems being digitally available, the information collected at the scale of 1:10 000 is available only for two-thirds of the country and is not digitised.

Very early maps at field scale, later at regional scale showed numerical salinity/sodicity values. At present field scale numerical maps are analysed in order to optimise salinity mapping in space and time.

Systematic monitoring of soil salinization in irrigated areas dates back to 1989 in the irrigation district of Tiszafüred in the county of Jász-Nagykun-Szolnok. A nationwide Soil Information and Monitoring System was initiated from 1991. In this system, from the 1 236 soil profiles, 69 profiles classified as salt-affected are sampled and analysed yearly for the indicators of salinity and alkalinity.

When the large-scale irrigation projects were planned in the second half of the last century, prediction of soil salinization was based on the concept of the critical depth of saline groundwater. As an alternative, a numerical rule-based algorithm was developed for the prediction of the risk of soil salinization in irrigated fields. Numerical process-based modelling with LEACHM and UNSATCHEM simulation programs was tested, but with limited success so far for Hungarian areas.

Based on environmental correlation, there is a long history of predicting the occurrences of salt-affected areas. A unique physical modelling system using compensation lysimeters are used for testing the effect of saline groundwater at different depths and with drainage management in a dominantly sodic area.

Reclamation, including installation drainage and afforestation of salt-affected soils, has lost its momentum in the country due to a decrease in financial profitability. On the

other hand, over the last few years a clear picture has been drawn of how efficient the different reclamation techniques are and how afforestation changes the salt distribution of soils and underlying strata.

Hungarian soil scientists play a leading role in the development of good practice for the management of salt-affected areas in Europe. Based on the initiatives of the Soil Framework Directive of the European Community, salinization is considered as a serious “soil threat” to agriculture. There is a well-developed concept of salinization and a set of common criteria for delineating areas threatened by salinization, are agreed on. A currently running EU-financed research project (ENVASSO) focuses on the assessment methodologies for quantifying soil threats. Another project (RAMSOIL) catalogues and evaluates the available risk assessment methodologies of each EU member country. To facilitate the spatial assessment of soil salinity/sodicity/alkalinity according to the EU-wide legal framework, the EU provides grants for promising research proposals to establish sensor-based digital field soil mapping methodologies. Such projects are based on the rich history of salinity sensing with field devices, such as the *Geophilus Electricus* system for multiple depth assessment of salinity.

Salinization and sodification of irrigated soils in Eastern Kenya

S. N. Wanjogu¹ and P. T. Gicheru²

¹Kenya Soil Survey, P.O. Box 14733 – 00800, Nairobi.

E-mail: kss@iconnect.co.ke

²Centre Director, National Agricultural Research Laboratories, P.O. Box 14733 – 00800, Nairobi.

E-mail: cdnarl@iconnect.co.ke

The Hola irrigation scheme covers 1 700 ha of land of which 872 ha are under irrigation. The scheme is in agro-climatic zone VI where the ratio of rainfall to evaporation is 15–25 percent. This zone covers 12.4 Mha of land in the country of which 2.3 Mha have soils that are slightly saline, 1.4 Mha are moderately saline and 1.9 Mha are severely saline. The major land uses in the research zone are semi-nomadism and nomadism. The soils in the study area are classified as Mollic and Haplic Solonetz, sodic or salic phase; Gleyic, Vertic and Calcaric Cambisols, sodic or salic-sodic phase and are developed on marine sediments of Pleistocene to Recent age and old alluvial deposits. Irrigation in the area started more than fifty years ago but increased salinization and sodification of the soils has led to the abandonment of irrigated farming in many fields since the mid-1980s.

The objective of this study was to assess irrigation induced changes in soil characteristics in the period 1987–2002 with a view to giving recommendations on possible management remedial measures in addition to assessing the possible impacts on the soil characteristics by the introduced shrub *Prosopis juliflora* in the study area.

Though the quality of irrigation water had been assessed to be satisfactory, sodicity and salinity of the soils have been increasing, while porosity has been decreasing in the topsoils, during the fifteen years of monitoring. Mean electrical conductivity (EC) values initially increased from 1.4 dS m⁻¹ to 2.1 dS m⁻¹ in the first nine years, but it has decreased to 1.1 dS m⁻¹ during the last six years. This may be an indication of increased leaching after the introduction of the shrub *Prosopis juliflora*, which has improved soil physical characteristics and thus enhanced flushing of the dissolved salts. Mean exchangeable sodium percentage (ESP) values have more than doubled from 10 percent to 24 percent, while porosity has reduced from 53 percent to 48 percent. Increase in EC and ESP indicate that processes of salinization and sodification have been taking place possibly due to inefficient use of irrigation water. A decrease in porosity indicates deteriorating soil physical characteristics.

Nature and distribution of salts in the Upper Wami-Ruvu Plains, Morogoro, Tanzania: Implications for land management methods

M. Kilasara¹ and E. L. Kaboni²

¹Sokoine University of Agriculture, P.O. Box 3003 Morogoro, Tanzania.

E-mail: mmkilasara@yahoo.com

²Ministry of Agriculture and Food Security, ARDI Maruku, P.O. Box 127 Bukoba, Tanzania.

E-mail: rykasuga@yahoo.com

Tanzania has about 7 Mha of potential irrigated agricultural land, which is located in 9 major river basins. The area under irrigated agriculture almost doubled to reach 3 percent of the total potential irrigated land over the past 10 years. While the government of Tanzania has ambitious plans to rapidly expand the area under irrigation as one of the strategies to address the Millennium Development Goals, there is evidence of decline in crop yield in farms that have been under irrigation for some time. The decline is associated with salinization, which is attributed to poor land management. Salinization is considered one of the key factors accounting for low returns in small-scale irrigation schemes in Tanzania. This study was conducted within the upper part of the Wami-Ruvu river basin in Tanzania in order to characterize and map salt-affected soils for the purpose of developing an appropriate land management strategy suitable for small-scale irrigated and rainfed rice production.

A 250 ha area of farmland located along the Ngerengere River (located between 6° 49' 00" S and 37° 35' 10" E) representative of the upper part of the Wami-Ruvu basin was chosen for the study. Soil auger observations were carried out at 50 m grid points to the depth of 100 cm in order to characterize the distribution of soils in the area. Soil samples from representative soil profiles were analysed for selected chemical, physical and mineralogical characteristics. The soils were classified according to FAO system. Rock, soil and surface and groundwater samples were collected and analysed in the laboratory so as to determine the source of salinity in the studied area. Supplementary 10 m grid soil observations were carried out at a depth of 0–50 cm in a 2 ha area. The area had patches that showed clear features of salt accumulation at the soil surface coupled with pockets in which the paddy rice crop performed poorly or the soil surface was bare. Soil samples from these points were analysed in the laboratory for pH and electrical conductivity (EC). Salinity and alkalinity classes were developed using established criteria. The performance of paddy rice crop was evaluated at the vegetative stage prior to flowering on the basis of crop vigour.

Three salt-affected and four taxonomic types of soils exist in the study: saline soil (Stagnic Solonchak – Chloridic, Clayic), normal soil (Stagnic Cutanic Luvisol – Hypereutric, Profondic, Clayic), sodic soil (Luvic Stagnosol – Eutric, Clayic) and Stagnic, Salic Solonetz (Abruptic, Clayic). Both the Solonchak and Stagnosol contained swelling type of clays. Basic gneiss, amphibolite and hornblendite were among the dominant rocks in the study area. All of them were rich in bases, which can be considered to be an important source of salinity. The EC of the water in the Ngerengere River remained low throughout the year while the content of sodium increased from 0.3–0.4 mg l⁻¹ during the rainy season to 29.9 mg l⁻¹ by the end of the dry season. Its use for irrigation is limited due to its sodicity. Most of the groundwater samples showed high EC and, as with the surface waters, chlorides, carbonates and bicarbonates of sodium were dominant. These are taken to be the direct source of salinity and possibly alkalinity in the studied soils.

Drastic spatial variations in EC and pH occurred over short distance in the saline-sodic soil. They had a pattern similar to that of the performance of paddy rice during the late vegetative growth stage. The mean values for EC in the root zone were 0.5, 4.0 and 20.2 dS m⁻¹ in the patches with healthy, wilted and withered crop or bare ground surface, while the corresponding mean pH values were 7.0, 8.1 and 8.9, respectively.

On the basis of the linkage between the performance of the paddy rice crop and the corresponding pH and EC values, the studied soils could be put into three broad salinity-alkalinity categories: very slightly acid to moderately alkaline and none saline to slightly saline soils, moderately alkaline to very strongly alkaline and moderately saline to strongly alkaline soils. These could serve as the basis for land management in the area. It is recommended that knowledge of the salt patch distribution, their characteristics and the chemistry of both ground and surface waters over the entire growing season should be investigated prior to any attempt to solve the problem of salinity and alkalinity in the study area. In view of the inconveniences that small-scale farmers are likely to face while linking with specialized laboratories to have soil and water samples analysed for EC and pH, efforts should be made to facilitate field-based characterisation and interpretation.

Salt-affected soils in Romania

M. Dumitru, C. Simota, D. Raducu, E. Dumitru, I. Calciu and N. Florea

National Research and Development Institute for Soil Science, Agricultural Chemistry and Environment (ICPA), Bulevard Marasti Nr. 61, Bucharest, Romania.

E-mail: office@icpa.ro, elisa_dumitru@yahoo.com

This paper presents information related to: (A) the assessment of salt-affected soil based on specific indicators and on (B) diagnostic criteria for their classification; (C) a distribution of salt-affected soils in Romania including (D) a map of salt-affected soils and vulnerable or susceptible soils to salinization; (E) a simple example in pilot area; and (F) the research concerning reclamation and management practices applied in Romania for salt-affected soils.

The assessment of the quality of salt-affected soils has been carried out using classical indicators of soil salinity. Soil sodicity is characterized by total salt content (TSC) in mg 10^{-2} g soil, exchangeable sodium percentage (ESP) in percentage of cation exchange capacity (%CEC), sodium adsorption ratio (SAR), in mg me l^{-1} and pH (soil:water = 1:2.5). Saline soils are characterized by their electrical conductivity (EC), measured in mMho cm^{-1} . From the taxonomical point of view salt-affected soils are assessed using the main specific diagnostic horizons: Salic Horizon (sa), Hyposalic Horizon (sc), Natric Horizon (na), Solonetzic Horizon (Bt_{na}), Hyponatric horizon (ac), Sulphuric Horizon. According to these criteria, the main soil types affected by salts in Romania are Solonchaks, Solonetz and Salinez soils, a category of non-saline soils with high risk or potential for salinity under certain conditions.

The areas affected – Romanian Plain (200 600 ha), Western Plain (175 000 ha), Moldavia (114 000 ha), Transylvania (20 400 ha) and Dobrogea (104 000 ha) – total 614 000 ha. To this area we can add the surface area at risk of salinity, which is 1.221 Mha. The Map of Salt-affected Soils and Susceptible Soils to be Affected by Salt in Romania shows the distribution and degree to which soils are affected by, or susceptible to, salts, as well as classes and sub-classes of soils and the dominant salinization type. Other details include relief units and groundwater depth and salinity.

Research concerning the reclamation and management practices for salt-affected soils has had a long tradition in Romania. Various institutions have been investigated techniques for management of reclaimed land affected by salinization, such as the National Research and Development Institute for Soil Science, Agricultural Chemistry and Environment (ICPA), Bucharest and the Research and Development Station for Improvement of Salt-affected Soils, Braila. They have tested traditional agricultural management methods. According to the National Soil Quality Monitoring System, these soils have been a particular problem since 1977. The amount of land cultivated for rice, one of the most profitable farming systems, doubled between 1980 and 1990 in the Maxineni-Corbu area, from 20 000 ha to 49 000 ha. However, from 1990 onwards, this area decreased dramatically. Other solutions for controlling soil salinization and improving soil status and crop yields, which were tested in experimental fields, involve drainage, land levelling and modelling, leaching, gypsum amendment, chiselling, soil improvement fertilization, halotolerant crops and crop rotation.

This paper presents a simple example of a study concerning salt-affected soils in a pilot area in the west of Romania.

SESSION 7: SUSTAINABLE BIOSALINE AGRICULTURAL SYSTEMS

Biosaline agriculture: Prospects and potential within global and regional context

S. Ismail and F. Taha

International Center for Biosaline Agriculture (ICBA), Dubai, United Arab Emirates.

E-mail: s.ismail@biosaline.org.ae

During the last few decades, net agricultural production has suffered a significant drop, although productivity per unit area has increased. Among the many reasons, salinity and associated factors, like waterlogging and/or drought, have contributed significantly. The increase in saline areas has been directly attributed to both water and soil salinity problems. An increase in groundwater pumping has resulted in the intrusion of seawater, in both coastal and inland areas. Furthermore, fresh water aquifers have been exhausted and, hence overlying saline water layers mix with fresh water, resulting in the increase of salinity in the groundwater. The most common reasons for the increase in salt-affected lands are the mis-management of irrigated areas. Inadequate or absence of drainage systems and high rates of irrigation water application have resulted in the movement of salts in the soil profile, especially in the dry areas, where these salts are brought to the surface as a result of high evapo-transpiration rates.

A number of approaches have been evaluated and implemented to combat the salinity problems, based on specific types of site, regional and global problems. These include soil reclamation, water management and plant-based approaches. The first approach of soil reclamation in large areas requires not only high financial investment, but also continuous maintenance to make it technically feasible. The other two approaches are management strategies using appropriate irrigation and drainage systems and selection of appropriate agriculture production systems, suitable for specific edaphic and climatic conditions. These approaches, when applied in an integrated form, constitute the backbone of biosaline agriculture.

Plant-based approaches to using salt-affected lands in an economically and environmentally safe manner are based on the salinity ranges of soil, water (including groundwater) and other associated factors. The selected production system(s) not only helps in halting further deterioration of marginal lands, but also has direct commercial uses such as food, forage/fodder, livestock industry, medicinal uses, wood, etc. In addition, the use of these marginal resources also provides many secondary and indirect products, including bio-fuel and bio-energy, carbon sequestration, phyto-remediation, etc.

Strategies for plant-based solutions to the use of salt-affected areas include the production of new genotypic material through conventional breeding or biotechnological methods (for glycophytes) or selection and adaptation of existing salt-tolerant germplasms (both glycophytes and halophytes). Both strategies are equally important and feasible based on the nature of the salinity conditions.

This paper will cover the latter approach of identifying, evaluating, screening and optimizing the productivity of existing salt-tolerant germplasms. Different management strategies to increase productivity will also be discussed. Case studies will be presented from different regions of the world (especially from third world countries) to use marginal and saline resources in different types of agricultural production systems. The paper will also describe the economic and environmental benefits of the case studies.

Greywater use for irrigation of home gardens in peri-urban areas of Jordan

M. Bino

The Inter-Islamic Network on Water Resources Development and Management, Amman, Jordan.
E-mail: muradinw@nic.net.jo

Marginal water includes saline water, waste water and runoff water. These water sources differ in location with regard to the application site, quality, quantity and stability of supply. Marginal water is usually deficient in one or more qualities that make it fit for open irrigation or other beneficial uses.

In a water-scarce region like the Middle East, it is not surprising that marginal water is considered to be an important water resource. Reclaimed waste water is mainly used in agriculture and its proportion in comparison to ground and surface water will be on the rise as the volume of municipal water flow increases and the respective collection and treatment systems are expanded and enhanced.

Projects implemented in Jordan show that implementation of greywater re-use systems could lead to an increased efficiency in the use of water, as well as a decrease in water demand by 15 percent and water bills by 27 percent on average. This provides an opportunity that should not be missed for improving the life quality of urban poor people as well as rural households. Greywater re-use for urban agriculture is especially relevant given that, by 2015, it is expected that 85 percent of Jordan's population will be living in urban areas.

The Inter-Islamic Network on Water Resources Development and Management (INWRDAM), Amman, Jordan, has implemented more than 1 200 greywater use installations since 2000 in different parts of Jordan. Experience has shown that the potential of greywater use at household level is not fully utilized and that there are no social or religious barriers to greywater use for irrigation of home garden crops under restricted irrigation practices.

This paper will address technical and socio-economic issues of greywater use at household level in peri-urban areas of Jordan and how greywater use is gaining increasing popularity in nearby countries such as in Lebanon, Yemen, the West Bank and Gaza and other countries in the region. The paper will address issues related to greywater treatment options and quality that it is possible to achieve by low-cost treatment methods, types of crops to grow, environmental impacts of greywater on soil and plants, benefit-cost analysis of re-use, acceptance by the local community and recommended re-use guidelines that it is possible to apply to regulate greywater re-use. The paper will also discuss and recommend policy options for increased use of greywater as an untapped marginal water resource.

Saline agriculture: Pakistan scenario

R. H. Qureshi¹ and J. Akhtar²

¹Higher Education Commission Sector H-9, Islamabad, Pakistan.

E-mail: rqureshi@hec.gov.pk

²Saline Agriculture Research Centre, University of Agriculture, Faisalabad, Pakistan.

Pakistan, having a land area of 800 000 km², is highly dependent on irrigated agriculture to feed over 155 million people. Irrigation has been one of the most important causes of salinization in the cultivated areas. Of the approximately 6.3 Mha of salt-affected land, about half is situated within the canal commands, thus having serious social and economic consequences for Pakistan.

These salt-affected soils vary not only in terms of physical and chemical characteristics, depth of water table, availability of water for leaching of salts as well as crop cultivation, but also their distribution pattern is highly variable. Therefore, there are different options to reclaim these lands and for their profitable agricultural use. Various types of salt-affected lands in Pakistan have been defined.

A summary of the wealth of knowledge accumulated through detailed experimentation involving screening programs for salt tolerance of crop species, studies in laboratory and agronomic studies in field have been described in this paper. Some indicators of economic returns of saline agriculture have also been discussed. The history of saline agriculture in Pakistan and the success story of the studies carried out in different parts of the country are included in the paper.

Extent of salt-affected land in Central Asia: Biosaline agriculture and the utilization of salt-affected resources

K. Toderich¹, S. Ismail², I. Massino³, M. Wilhelm⁴, S. Yusupov⁵, T. Kuliev⁶ and S. Ruziev⁷

¹International Center for Biosaline Agriculture (ICBA), Central Asia and Caucasus sub-office, Tashkent, Uzbekistan

E-mail: ktoderich@cgiar.org

²International Center for Biosaline Agriculture (ICBA HQ), Dubai, UAE

E-mail: s.ismail@biosaline.org.ae

³Scientific Production Center for Maize Production, Tashkent region, Uzbekistan

⁴Uzbek Research Institute of Karakul Sheep Breeding and Desert Ecology Research, Samarkand, Uzbekistan

E-mail: ecokar@rol.uz

⁵Institute of Agriculture and Agroecology in Pre-Aralie, Kyzylorda, Kazakhstan

E-mail: pniiacsx@rambler.ru

⁶Department of Botany, Gulistan State University, Gulistan Uzbekistan

E-mail: kuliev@mail.ru

⁷Ministry of Agriculture of the Republic of Turkmenistan, Ashgabad, Turkmenistan

E-mail: rozyews@online.tm

The economy of Central Asia countries (Uzbekistan, Kazakhstan, Turkmenistan, Tajikistan and Kyrgyzstan) is based primarily on agriculture and agricultural processing, with cotton and wheat being the major export crops. This dependence of the economy on agrarian-based activities is unlikely to change in the foreseeable future as most of the countries mentioned have embarked on a staged process of market liberalization. With the collapse of the former Soviet Union, economies in several of these countries have contracted with associated socio-economic, environmental and food security issues attaining greater prominence.

The large expansion of the irrigated area in the Aral Sea Basin has exacted a substantial toll on land and water resources in the Basin. Elevated water tables associated with poor irrigation management and non-functional drainage infrastructure, due to the lack of financial resources, have resulted in significant salinization of crop lands that invariably results in abandonment of land due to declining wheat and cotton yields. Approximately 600 000 ha of irrigated cropland in Central Asia have become derelict over the last decade due to waterlogging and salinization. It is estimated that approximately 20 000 ha of irrigated land of the Mirzachuli Steppe (marginal transboundary areas of Uzbekistan, northern Tajikistan and the southern part of Kazakhstan) are lost to salinity and invariably abandoned every year. The proportion of irrigated land that is salinized to some extent has risen from 48 percent in 1990 to approximately 64 percent in 2003. In some downstream provinces of Uzbekistan (Navoiy, Bukhara, Surkhandarya, Khorezm and Karakalpakstan), Turkmenistan (Dashauz) and around the Syrdarya delta from the Kazakh side, 86–96 percent of irrigated lands are salinized. Saline areas are generally found in poorer areas of the region with per capita incomes 30 percent lower than the national average and where unemployment levels are 40 percent higher. Soil salinity and waterlogging are a heavy burden for resource-poor farmers who are located in arid/hyper-arid degraded zones. This phenomenon has had a major impact on the livelihoods of rural communities who are dependent on land and water resources for goods and services. The conventional approach to rehabilitating these salinized areas requires major technical expertise and investments that are beyond the means of national budgets, as well as farmers' investment capacity, in these emerging/

transition economies.

Degradation of natural desert pastures, throughout all the Central Asia states has reached an alarming degree, requiring prompt action for the fragile ecosystems. The average annual rainfall here varies from 80 to 120 mm. Soils are sandy loam to loamy in texture, with poor vegetation cover that is highly saline with a low fertility. *Artemisia*-ephemeral and psammophytic rangelands available for grazing of animals have been replaced by halophytic plant communities with less palatable fodder plants. In addition, these *Artemisia* pastures tend to disappear under excessive and permanent grazing and also because they are currently being heavily uprooted for fuel. The anthropogenic transformation of pasture vegetation is evident near wells or watering places, saline lakes, settlements, sheep folds and along roads.

The poor natural drainage system of marginal cropping irrigated lands and sandy deserts has also caused an increase in the salt contents of surface soils and groundwater, which has induced secondary salinization. This has resulted in the migration of the local population (mostly Kazakh and Uzbeks) to neighbouring cities or countries. A consequence of pasture degradation is a big decline in the livestock system and the livelihoods of the communities in the region.

Nowadays, a unique source for development of agropastoral livestock-feed systems in the remote desert/semi-desert zones of all Central Asia countries is the reclamation of saline pastures (currently occupying 3.2 Mha), where halophytes that have the ability to grow adequately under prevailing edaphic conditions play a valuable role as feed for livestock and for medicinal purposes. Until recently, no serious research efforts have been made on the cultivation of wild halophytes, so they have had no market value as yet.

The role of plants in the remediation of saline and sodic soils is an emerging low-cost approach in the reclamation of abandoned irrigated lands. In this respect, the creation of highly productive fodder systems through the establishment of palatable halophytes has been shown to remediate saline/sodic soils as well as providing an income to resource-poor farmers. These genetic resources play a very important role in (i) the rehabilitation of degraded lands; (ii) controlling high water tables; (iii) utilization of non-conventional water resources; and (iv) landscaping. An innovative programme and experimental fields on domestication and utilization of *Glychyrryza glabra*, *Hippophae ramnoides*, *Elaeagnus angustifolia*, *Artemisia diffusa* and *Alhagi pseudoalhagi* with suitable modern agro-technologies were initiated in 2006 in the Central Kyzylkum Desert. Incorporation of these plants into a biosaline farming system represents the only source of income for many poor rural families, who live far away from markets. These plants, due to their ability to be propagated by both reproductive (seeds) and vegetative (suckers) means, are the target fodder species for rehabilitation of degraded pastures; sand-fixing; water-table and soil erosion control; haymaking and silage; better stock feeds (feed blocks) for animals in the late autumn/winter; bee-keeping and honey production; and the volatile oil of *Artemisia* for traditional medicine. The fruit of *Hippophae* and *Elaeagnus*, as they are rich in sugar, falconoids and various vitamins, can be used by local people for jam and wine production and dyes. Glychyrric acid extracted from *Glychyrryza glabra* roots is used as flavouring in food, as well as in tobacco, alcohol, candies and cosmetics.

Options for improving the livelihoods of the rural population in some Central Asia regions by using artesian thermal and groundwater are evident. The artesian waters could be used for development of arid fodder production systems, as well as for recreation, vegetable production, and other purposes through appropriate management practices. The establishment of highly-productive fodder systems will ensure the safety of the natural habitat and increase the income levels of the poor farmers. However, since the whole issue involves using the saline artesian water for long-term sustainable production, care needs to be taken of management and environmental issues.

Transferring new technology or methodology of the International Centre for Biosaline Agriculture (ICBA) in the planting of both perennial and annual valuable halophytes (based on worldwide dataset from similar sites and conditions) is an alternative approach that is already being tested in some Central Asia regions. Preliminary results are very encouraging and demonstrate the potential of salt-tolerant plants and halophytes for food, animal feed, fuel wood, bioenergy and other products. These species offer an alternative to the traditional cotton growing systems, the yield of which has been significantly reduced due to an increase in salt-affected areas.

Many farmers who had abandoned their cotton farms have returned to the production system and are gaining economic returns. Sorghum and pearl millet, among conventional forage, and *Atriplex* spp. and *Acacia ampliceps*, among the non-conventional forage crops, have gained a good deal of interest in the region. Such integrated research is both novel and timely for bringing these countries into the modern world.

Extent and utilization of salt-affected lands: Biosaline agriculture and marginal resources in Tajikistan

S. Sanginboy

Tajikistan Academy of Agricultural Sciences (TAAS), Dushanbe, Tajikistan.

E-mail: sanginov@yahoo.com

The agricultural sector is an important part of the national economy due to its contribution to the GDP, employment and exports. Considering that three-quarters of the population lives in rural areas, the agriculture sector will play a crucial role in poverty alleviation. While arable land is scarce, experience in recent years, based on the ongoing restructuring of the large state-owned cooperative farms, has shown that the transition to private farms can raise yields substantially.

Food security is a big issue for Tajikistan. It is a landlocked country, and the Government has emphasized that self-sufficiency in wheat production is a national priority, flour being the single staple food. Agricultural land in Tajikistan, especially arable land, is a very scarce resource. Total agricultural land is 4.57 Mha, of which arable land counts for 0.7 Mha, or 0.11 ha per capita, including pasture lands. The major problem is the access of peasants to land and its fair distribution. Efficient use of, and fair access to, land and water resources is one of the sector's priorities in fighting poverty. A key issue for agricultural development in irrigated zones of Tajikistan are the pumping and maintenance costs of groundwater and irrigation facilities. High expenditure is required every year to keep them in working condition. With the transition of the country to a market system, a fee on water consumption was introduced, starting in 1996. However, farmers are only able to afford a small part of the maintenance costs. The main concern still remaining is the state of pumping stations covering almost 300 000 ha, or 40 percent of irrigated land. A rural population of about 2 million resides and earns a living in this area.

Low efficiency of water consumption in agriculture is another big problem. Lack of inputs and technical resources and destruction of their distribution systems, destruction of irrigation system infrastructures, among others, has caused soil degradation by salinization and waterlogging. This has resulted in decreasing productivity of crops by as much as 50–60 percent, poor crop quality and more energy for pumping water. As a result, at present, Tajikistan has more than 116 000 hectares of saline and 30 000 hectares of waterlogged land. The irrigation and drainage water in valleys has raised the groundwater level (1–2 m to soil surface) in the lower-lying lands and territories. Many of the cotton-producing areas of the country are now suffering from marked salinity.

Development of irrigation in the republic has been accompanied by an increase in water use. For 80 years, it was 5.5 km³ year⁻¹. Currently it is 11.2 km³ year⁻¹ with water consumption above 20 000 m³ ha⁻¹ year⁻¹. As a result, 30 000 hectares of irrigated land are experiencing secondary salinization and waterlogging.

Biosaline agriculture provides a big opportunity to use these marginal resources in wastelands and turn these into production systems. The use of physical, chemical and biological management practices could help in flushing the salts down in the soil profile. As a result of plant succession using different salt-tolerant plant species, economic returns can be achieved and thus help to alleviate poverty.

The salinity problem in the Sultanate of Oman: Past, present and future perspectives

S. B. A. R. Al-Rasbi

Ministry of Agriculture and Fisheries, Sultanate of Oman.

E-mail: salrasbi@yahoo.com, agricop@omantel.net.om

The Sultanate of Oman is one of the countries located in dryland areas. Its average annual rainfall is less than 100 mm. In the previous three decades the salinity problem has increased dramatically. The main reason besides lower rainfall was the over pumping of fresh water. The Ministry of Agriculture and Fisheries (currently the Ministry of Agriculture) recently became aware the problem. Through a field survey it was found that at least 50 percent of groundwater in the agricultural coastal region of Batinah was saline. Some control measures were taken, including legislation preventing the digging of new wells. Detailed surveys were conducted to study the soil and water salinity problem and attempts were made to use the saline water. Programmes were established and still being conducted to select salt-tolerant crops. Several experiments were conducted on field crops, forest trees, shrubs and vegetables. A scheme of transferring plants that consume high quantities of water away from the salinity affected areas is the current focus. The ministry cooperated with international institutes, such as the International Center of Biosaline Agriculture (ICBA) and the Food and Agriculture Organization of the United Nations (FAO), as well as local agencies like the University of Sultan Qaboos and the Ministry of Regional Municipalities and Water Resources.

A programme for the development of salt-tolerant cultivars and varieties was started by ICBA in 2003. Under this programme, different lines of varieties of pearl millet, sorghum, barley and canola (*Brassica sp*) were studied under moderately and highly saline irrigation water. Shrub species *Atriplex lentiformis*, *Atriplex halimus* and *Atriplex nummularia* and tree species *Acacia ampliceps* were experimented at higher water salinity levels. In addition, demonstration plots of grass species *Leptochloa fusca*, *Paspalum vaginatum*, *Distichlis spicata*, *Sporobolus virginicus* and a nursery of grass species *Leptochloa fusca*, *Paspalum vaginatum*, *Distichlis spicata*, *Sporobolus virginicus* were also established at higher water salinity level. The most tolerant lines of the tested varieties were then replanted to select the most suitable to distribute to farmers.

Moreover, the Ministry recently got involved in a programme with the University of Sultan Qaboos. This programme has the following objectives: monitoring and preparation of salinity maps; identification of management practices to tackle soil and water salinity; cultivation of salt-tolerant fodders/grasses; selection of sowing techniques; fish culture with saline water; and socio-economic analysis.

To identify the quantity and quality of the brackish water in whole of the Sultanate, the Ministry of Municipalities, Water Resources and Environment conducted a field survey. In this survey remote sensing tools were used. Areas of aquifers were identified and results were presented in tables, graphs and maps. An area of about 107.8 km² of brackish groundwater was found in the inland areas of Oman. Options were suggested for using brackish water, including using the water directly, economic model projects, allocation to large-scale farming, small aquaculture farms and desalinization with reverse osmosis to produce potable water.

The salinity problem in the sultanate could be solved through continuing to apply the legislation of minimizing groundwater pumping; developing new salt-tolerant plants through screening and breeding techniques; and developing planting and irrigation techniques which suit saline agriculture.

Water shortage in western US and saline recycled water use for urban irrigation

K. B. Marcum

Dept. of Applied Biological Sciences, Wanner Hall, Arizona State University, East Campus, 7001 E. Williams Field Rd., Mesa, AZ 85212, U.S.A.

E-mail: Kennth.Marcum@asu.edu, <http://ctas.poly.asu.edu/marcum>

Critical water shortages are occurring throughout the western U.S. due to population pressures resulting from urban influx and prolonged drought due to climate change. Land and water used for irrigation is rapidly shifting from agriculture to urban use and most (50–70 percent) of urban water use is for landscape irrigation. Rapid urban development along coastlines has resulted in salt water intrusion due to over-pumping, causing salinizing of fresh water aquifers. For these reasons, governmental policies have been adopted in western states requiring use of recycled sewage water, brackish groundwater and other secondary, saline water sources for urban landscape irrigation.

Landscape managers face unique challenges due to high traffic and the inability to cultivate soil, exacerbating sodicity issues due to compaction pressures. Special management is often required, such as irrigation scheduling, irrigation acid injection, modified root zones and subsurface drainage, in addition to the use of salt-tolerant species and cultivars. Relative salinity tolerance and associated physiological responses of turf grasses for urban landscapes were investigated.

Tolerance ranged from salt-sensitive to halophytic. Species were ranked for salt tolerance, and recommendations made. Salinity tolerance was associated with increased root/shoot ratios, shoot saline ion exclusion, minimal shoot osmotic adjustment, salt gland activity and compatible solute accumulation.

Part 3

Findings and reports from working groups and field visit

FINDINGS

One day of the SPUSH Consultation was devoted to group discussions. The participants were divided into three groups and discussed the following issues:

- The nature of the problem, including the extent, causes, distribution and degree of salinity development, the criteria, classification and diagnosis of salt-affected soils applied in participating countries, the availability of information and the importance of geo-referenced data to classify a soil as salt-affected.
- Practical methodologies, which are applied for assessing and monitoring of salt-affected soils, as well as lessons learned at field, basin and national levels, including mapping of salt-affected soils and availability of satellite imagery for mapping salt-affected soils in participating countries.
- Predicting and modelling for salinity/sodicity development. Identifying weaknesses, strengths and opportunities for creating models closer to field conditions.
- Policies and strategies for assessment and monitoring of salt-affected soils
- Main solutions and proven technologies applied in the participating countries to combat salinization/sodication and salinity problems.

Half a day was devoted to participatory group discussions on the way forward for SPUSH. The following issues were discussed:

- Determination of priorities for future activities, research and cooperation between member countries.
- Communication strategies.
- The next Network Expert Consultation.

The following are the findings integrating presentations and discussions from the different discussion groups.

Causes of salinity development in some participating countries

- Climate (mainly low rainfall and high evaporation).
- Improper land use and poor agronomic practices.
- Mineralization of groundwater.
- Natural origin.
- Rise in water table.
- Introduction of large-scale irrigation without proper management.
- Seawater intrusion.
- Use of poor quality water for irrigation without proper soil/water and nutrient management.
- Land tenure systems.
- Poor drainage systems.
- Over-pumping of groundwater for irrigation with low recharge.
- Less coordination between institutions concerned.

- Water-saving strategies.
- Clearance of mangroves, shrimp farming, backflow of seawater during the dry season.
- Seawater spray resulting from breaking of wave crests during typhoon surges.

Assessment and monitoring methods used in some participating countries

- Many countries do not have any programmes for assessing or monitoring salt-affected soils. Some have very old data and no national salinity maps.
- Few countries conducted soil surveys, soil sampling and diagnostics at different scales for the whole country.
- Remote sensing and GIS for assessing, monitoring and quantifying salt-affected soils were used in few countries.
- Satellite imagery can be used successfully where salt-affected soils occur in the topsoil, and it can be cost-effective if good ground truthing methods are available.
- In some countries monitoring programmes only cover monitoring of well water, but not necessarily for the whole country. Very few countries carry out monitoring programmes in the whole country on an annual basis, such as in Hungary, Romania and Tajikistan. In such cases, fixed points (benchmarks) for long-term monitoring should be identified. Proper site selection for effective monitoring is a must.
- In most countries, the soil electrical conductivity of the soil saturated extract is used as an indicator of soil salinity, while the ESP or SAR of the saturation extract is the indicator for sodification. For a large-scale assessment of soil salinity, mobile electromagnetic induction EM38 and EM34 are used for delineating areas of different levels of salinity. However, calibration techniques should be well established.
- Salinity sensors and tensiometers for simulation of salt and water dynamics are used in few countries (e.g. China).

Modelling of salinity and sodicity development

- Models could be of limited use if they are not well designed. Some models can be very vulnerable to particular parameters if not properly developed.
- Comparison of two models (SMSS2 and SMSS3) with reference to irrigation-induced soil and water quality parameters was presented by Morocco. The statistical results from the model outputs supported the reliable use of models. The character of soil physics should be studied for reliable prediction models. The SWAP model has been used efficiently and could be shared with SPUSH member countries.
- Limitation of the use of modelling under saline conditions is due to the dynamic nature of salinity problems which should be clearly understood by model users.

Mapping and interpretation of spatial data

- The spatial assessment of salt-affected soils starts with the systematic mapping of salt-affected areas.
- Some participating countries have standardized mapping protocols that could eventually be shared with other member countries.
- Scales of maps (if available), classification systems and technologies used vary in the different member countries. Presenting salinity in different units and scales in participating countries is a problem for comparison purposes. However, standardizing different criteria and methodologies for salinity mapping is a challenge. This could be a future activity for the SPUSH Network.
- Standards and sampling protocols are needed for different mapping scales. This would be also useful for the production of cost-effective monitoring systems.

Main technological solutions

- Improved irrigation to prevent rise in water table, preventive and effective drainage systems and appropriate land use.
- In general, for sodic soil, chemical amelioration amendments accompanied by suitable crop husbandry constitute the main technological intervention.
- For saline, waterlogged soils, leaching and drainage accompanied by suitable agronomic practices are effective for land reclamation.
- The use of salt-tolerant crops, grasses, trees and halophytes is promising, as demonstrated in some INBA countries.
- Construction of protective embankments with flow-regulating gates.
- Rainwater harvesting and use of water efficient technologies.
- Appropriate management of brackish water for irrigation use.
- Afforestation.
- Biosaline agriculture, including breeding salt-tolerant species can contribute to the improvement of farmers' income.

REPORTS FROM WORKING GROUPS

The following are the reports produced by the rapporteurs and drafting group. They are organized according to the topics discussed.

Assessment and monitoring of salt-affected soils

A regular monitoring system for salinity assessment at country, regional and global level is essential to convince policy-makers of the threat/impact of salinity, and the opportunities available to tackle problems. It was highlighted that:

- In most of the member countries there was no strategy or policy at national level for assessing, monitoring, and mapping of salt-affected soils. Various countries lack a systematic national system to monitor secondary salinity. Studies were carried out on an *ad hoc* basis and reliable data to establish baseline conditions are also lacking. However, some data were available from sources such as Terrastat, Aquastat, Agenda 21 and State of Environment reports.
- Financial and human resources to monitor salt-affected soils are not easily available in most countries. More efforts to increase human resources and capacity building should be made to strengthen research, technology development and transfer.
- SPUSH member countries should identify hot spots and use them as a benchmark for monitoring. They are also encouraged to adopt early warning systems to prevent more salinization and develop farmer-friendly salinity assessment and monitoring methodologies.
- Countries like India, Tanzania and Romania have up-to-date databases related

to salt-affected soils and poor quality water up to 2006/2007. Other SPUSH countries need to update their databases. This will not only benefit the country but will also facilitate the preparation of a more accurate map of salt-affected soils at global level by FAO, to raise the awareness of the situation and request funds for rehabilitation programmes.

Raising awareness and strengthening coordination at national level

- Member countries are encouraged to publish quality papers on various aspects of soil salinity. This will strengthen the Network already established for this purpose.
- Some countries have standardized assessment, modelling and mapping methodologies and procedures which could be shared with other SPUSH partners to update the knowledge on the extent, distribution, nature and classification of salt-affected soils in member countries.
- Member countries may jointly develop brochures or other material on assessment and monitoring techniques, technologies developed for the reclamation and management of salt-affected soils, and judicious use of poor quality/wastewaters to be shared with other countries.
- As more than one authority or institute in each member country is concerned and has related activities, data, maps, reports, research and development programmes, special institutional arrangements should be made to allow effective cooperation and avoid overlapping of activities at national level.
- The education of the younger generations is important and universities should be encouraged to include salinity issues as part of their education programme on soils.

Standardization of methods

Participants thought there was a need for standard analytical methods as well as terminology, without compromising existing national and international standards. Scientists should recommend how to standardize these methods and terminology, after which correlations with national methods and terminology can be developed. Participants recommended that international institutions coordinate this activity.

With regard to standardization, almost all participants agreed that:

- Soil electrical conductivity of the saturated extract (EC_{sat}) should continue to be used as the indicator of soil salinity, and the exchangeable sodium percentage (ESP) as the indicator of sodication.
- For a large-scale assessment of soil salinity, a methodology based on the EM38 sensor can be used, and maps can be created delineating areas with different levels of salinity.
- Data must be standardized, fit for purpose and stored properly in a format that is easy to access and process. Data that are incorrect, misleading, incomplete and outdated must be cleaned.
- A uniform risk assessment methodology may be developed to rank areas according to the relative probability of being affected by a given level of salinity, sodicity and alkalinity, and use the data to determine whether farming practices increase soil salinity, soil sodicity and soil alkalinity. The data may be converted and expressed in common units/scales.

Modelling for salinity/sodicity development

- Modelling soil-water-salt-plant relationships is important for developing, using and transferring technologies as well as developing decision support systems. Scientists need to consider how models can better represent field conditions, since several factors play significant roles in the behaviour of biological systems

such as agricultural practices. Scientists should consider issues like climate change and its potential impacts in salt-water dynamics under actual conditions.

- Models require very intensive efforts in research and computing which is problematic for most countries. Physically-based models simulating water and solute transport represent an essential tool for predicting soil salinity and/or sodicity. These models allow for comparing different options to develop strategies for sustainable irrigation in the short- and long-term. However, calibration and validation of these models against soil and crop field data are needed to check the accuracy of the predicted values before these models can be used to develop reliable management scenarios.
- The following processes need to be included in models: transient water transport, transient salt transport, ion exchange, precipitation/dissolution index of salts, changing groundwater level, waterlogging, preferential flow (where applicable), freezing/thawing (where applicable), changing irrigation water composition, plant uptake of ions, CO₂ partial pressure and its effects, heat transfer, depth distribution of roots in relation to water uptake from soil and shallow groundwater and solution chemistry, including pH.
- Assessment and monitoring of associated salts/metals like boron, iron, aluminium, manganese, arsenic, selenium, nitrates, etc. should also be considered.

Mapping and interpretation of spatial data

Assessment methods and management techniques to deal with salt-affected soils must take into account economic and social factors. Geomorphology must be the first information source for sampling design. Satellite imagery can be used successfully where salinity occurs in the topsoil, and it can be cost- and time-effective, with good verification prospects. Map coordinates are used for the site location and fixed points (benchmarks) are required for long-term monitoring. The ratio of desired number of samples to associated cost of sampling can be optimized through adequate sampling design.

An expert group needs to be constituted within the SPUSH Network to address issues of:

- Perfection of GIS and remote sensing methodologies to delineate waterlogged saline soils from waterlogged soils. However, from a crop production/soil salinization point of view, waterlogged soils are those with water table within 2 m from ground surface.
- Dependence of appearance of salt crust on accumulation of hygroscopic versus non-hygroscopic salts that may affect the remote sensing data.
- Presence of salts in sub-surface layers and presence of associated toxic salts such as aluminium, selenium, etc.

Saline agriculture

- In many areas, there is a need to adapt to salinity. Several countries under the SPUSH and INBA umbrella have identified promising germplasm of trees, bushes/ grasses/crops, as well as their varieties, and fish, which need to be shared with other partner countries.
- Developing joint network projects/programmes addressing the issues of utilization of salt-affected soils and saline water with the objective of sharing information and regional coordination and of monitoring and assessment could be effective to improve efficiency and share costs.
- Reclamation of salt-affected land is largely attempted through high cost input technology. However, in certain situations, growing salt-tolerant plants (crops/pasture/trees) may be more cost-effective for rehabilitation of such lands. Therefore, programmes aimed at screening and developing plant varieties with

high salt-tolerance should be put in place. Any alternative varieties selected for inclusion in a diversification programme should have satisfactory salt-tolerance, and be economically worthwhile.

- A strong extension programme would also be required. Exchange of salt-tolerant germplasm between member countries could have immediate impact. A seed bank in FAO may have a role in facilitating the exchange.

Management strategies

- To facilitate technology-sharing and ensure that it reaches farmers, farmer associations and field technician services, research and policy making should be strengthened in the member countries.
- Strengthening the capacity of field technicians is critical if the technologies are to be disseminated to the target groups. There are various ways of strengthening technology sharing, such as specialised training, farmer field schools, volunteer soil doctor, model farms and information campaigns.
- Farmers should be active participants in the development of the appropriate management systems for salt-affected soils, for which appropriate training is needed.
- The Network should develop guidelines and a database on resource surveys and management practices and make this available to those involved in land use planning for irrigated agriculture, extension officers, other advisors to farmers, and farmers. This database must be maintained and updated regularly.
- Although the definition of salt-affected soils in relation to nature and quantity of salts in soils and water is clear, guidelines need to be developed by Network members to choose management strategies that reflect local conditions, such as:
 - agro-climatic zone
 - soil texture and drainage characteristics
 - rainfall patterns
 - distribution of salts in the soil profile
 - other associated salts/metals
 - expected land use
 - level of management/cultivation practices
 - quality of irrigation water
- Promoting water use efficiency, affecting control over water tables and preventing salinization are major objectives of the management strategies. While much can be achieved through improved management, special attention should be given to communal drainage networks, on-farm drainage and preventing leakage from irrigation channels.
- Issues of property rights should be addressed, in collaboration with the World Intellectual Property Organization (WIPO), so that the member countries can share knowledge while developing new technologies to address salt-affected soils.
- Various technologies were presented during the Expert Consultation. It is recommended that member countries share technologies developed for reclamation, management, assessment, monitoring and mapping of salt-affected soils and judicious use of poor quality/waste waters. Human resources development/capacity-building initiatives should be encouraged to strengthen research development and technology transfer systems between participating countries.

FIELD VISIT

On 29 November 2007, three hours were devoted to visiting ICBA facilities and learning about their programmes and activities.

ICBA is a scientific research institution with a unique focus on developing and promoting systems that facilitate agricultural production in areas characterized by saline water or soils. It has modern facilities dedicated solely to the development of saline agriculture. The Center was established in September 1999 with its headquarters in Dubai, UAE, and funded by the Islamic Development Bank, the OPEC Fund, the Arab Fund for Economic and Social Development and the Government of the United Arab Emirates. When the Center became operational, it focused on research that would directly benefit agriculture in the six Gulf Cooperation Council (GCC) countries. The Center extended its work to the members of the Islamic Development Bank and gradually broadened its activities, including regional projects, to cover other countries where farmers cultivate their crops in saline soils or irrigate them with saline water.

Its mission is to demonstrate the value of saline water resources for the production of environmentally and economically useful plants. It actively ensures that the results of the research are of interest to those countries using salt-affected soils and saline water for agriculture and wishing to preserve fresh water resources by growing salt-tolerant forage species in marginal areas.

In order to become acquainted with the field activities undertaken by ICBA and to see the facilities of the Center, the participants, accompanied by technical staff, visited its experimental farm. It has a 100 ha research farm, of which 35 ha is developed (14 blocks of 2.5 ha each) and already under research and experimental programmes and the other 65 ha is under construction, rehabilitation and development. The farm has two sources of water for irrigation: groundwater of 30 dS m⁻¹ and Ein Municipal water of only 3 dS m⁻¹ to be blended to obtain three levels of saline water (5–15 dS m⁻¹ for crops and forages, 15–20 dS m⁻¹ for halophytes and >20 dS m⁻¹ for halophyte and mangroves) for irrigation, mainly using sprinkler or drip irrigation systems. It also has a large tank for seawater of 40 dS m⁻¹ as a third source of irrigation water. The farm is under a tile drainage system to draining into two evaporation ponds. This water is also used for irrigation and growing mangroves. The Center has four air-conditioned greenhouses and a large shadehouse providing climate-controlled conditions for a wide range of experiments. These facilities also ensure a secure environment for the production of seeds adapted to saline environments. In addition, the Center has a computer controlled irrigation system to allow precise control of the amount and salinity of irrigation water. It has a well-equipped laboratory for each facet of scientific investigation and a weather station and also 75 lysimeters of 70 cm depth which are used for growing *Leptochloa fusca*, *Conocarpus lancifolius* and *Atriplex canescens*.

The work of ICBA falls into four main areas:

- plant genetic resources – collection and characterization of germplasm; maintenance of a gene bank for salt-tolerant crops; and exchange of genetic resources;
- sustainable production and management systems – identification of effective irrigation systems and methods; evaluation and selection of field and forage crops and halophytes that can flourish under saline conditions;
- information management and networking – development of collaborative research networks, dissemination of technology and information on saline irrigated agriculture, creation of biosaline agriculture information networks;
- extension and training – publication of bulletins, brochures and newsletters; holding workshops, seminars and conferences; provision of opportunities for the on-the-job training and convening field days and open days.

During the visit, participants held comprehensive discussions and shared views on the possible future activities of the SPUSH Network and exchanged experience on the available methodologies and technologies on assessment, monitoring and modelling of salt-affected soils.

Possible ways of collaboration between the International Center for Biosaline Agriculture and its Inter-Islamic Network on Biosaline Agriculture (INBA) and the FAO Global Network (SPUSH) were discussed.

ANNEX I

Information on the networks

THE GLOBAL NETWORK ON SALINIZATION PREVENTION AND PRODUCTIVE USE OF SALT-AFFECTED HABITATS (SPUSH)

The SPUSH Network is a neutral forum to connect research institutions, land users and policy makers. The Network aims at disseminating and exchanging information; facilitating the application of technology; contributing to the design of relevant policies and promoting focused scientific research. The main topics covered by the Network include:

- assessment and monitoring tools,
- extent and types of salt-affected soils,
- impact of salinization, sodication and waterlogging on agricultural production, ecosystems and ecosystem services,
- effects of salinization, sodication and waterlogging on other factors related to agricultural production (e.g. nutrient uptake, diseases, pests),
- management practices to prevent salinization, sodication and waterlogging,
- management practices to increase the productivity of salt-affected habitats, including biosaline agriculture,
- practices for the rehabilitation and reclamation of salt-affected habitats,
- linkages between climate change and salinization, sodication and waterlogging.

These topics consider primary (of natural origin) and secondary (human-induced) salinization, sodication and waterlogging; salt-affected habitats in rainfed, irrigated and coastal areas; and technical and socio-economic aspects linked to management of salt-affected habitats.

THE INTER-ISLAMIC NETWORK ON BIOSALINE AGRICULTURE (INBA)

The Inter-Islamic Network on Biosaline Agriculture (INBA) is a non-political, non-profit, independent and autonomous body promoting biosaline agriculture under the auspices of the Organization of the Islamic Conference (OIC) Ministerial Committee on Scientific and Technological Cooperation (COMSTECH). It was established in 2002 at the 10th General Assembly meeting of COMSTECH.

The Inter-Islamic Network on Biosaline Agriculture aims to provide a forum for collaborative action and networking activities that stimulate and strengthen national and international institutions and aid agencies. Members include the Ministries of Agriculture and Water Resources, universities, national, regional and international agricultural research and development agencies, extension services, and end-users, including farmer groups and non-governmental organizations (NGOs).

ANNEX II

SPUSH expert consultation and INBA meeting programme

Monday, 26 November 2007

- 08:00–09:00 Registration
- 09:00–10:00 Inaugural Session
- Welcome address by Dr. Shawki Barghouti, Director General ICBA
- Address by **Mr. Fawzi AL-Sultan**, Chairman ICBA BoD
- Address by **Mr. Kayan Jaff**, FAO representative in the UAE
- Address by **Dr. Amin Mohamed Mashali**, FAO
- Address by **Mr. Mohammad Tourie**, IDB
- Address by **Dr. Anwar Nasim**, COMSTECH
- Address by **Prof. Dr. Faisal Taha**, Director Technical Programme ICBA
- 10:00–10:20 Tea / Coffee Break
- 10:20–10:50 Introduction to the SPUSH Network and the Expert Consultation
Dr. Amin Mohamed Mashali, FAO
- Session 1 Assessment and monitoring of salt-affected soils (at field, landscape and irrigation district levels)
- Chairperson: **Dr. Jorge Batlle-Sales**, Spain
- Rapporteur: **Dr. Abdullah Dakheel**, ICBA
- 10:50–11:20 Assessing and monitoring the risk of salinization in a Sicilian vineyard using the Geonics EM38
Dr. Giuseppina Crescimanno, UNESCO and **Dr. Kenneth B. Marcum**, USA (Keynote presentation)
- 11:20–11:40 Use of an above-ground electromagnetic induction meter for assessing salinity changes in natural landscapes and agricultural fields
Dr. Janette Arriola-Morales, Mexico and **Dr. Jorge Batlle-Sales**, Spain

- 11:40–12:00 Primary soil salinity, sodicity and alkalinity status of different water management areas in South Africa
Dr. J.P. Nell, South Africa
- 12:00–12:40 Discussions
- 12:40–13:40 Prayer and Lunch break
- Session 2 Assessment and monitoring of salt-affected soils at national and regional levels

Chairperson: **Dr. Giuseppina Crescimanno, UNESCO**

Rapporteur: **Dr. Shoaib Ismail, ICBA**
- 13:40–14:00 Advances in assessment of salt-affected soils for mapping, monitoring and management strategies in India
Dr. Gurbachan Singh, India
- 14:00–14:20 Salt-affected soils in Thailand: Assessment and monitoring of salinization
Dr. Rungsun Im-Erb, Thailand
- 14:20–14:40 An overview of salinity problems in Iran: Assessment and monitoring technology
Mr. Y. Hasheminejad, Iran
- 14:40–15:00 Advances in assessment and monitoring of soil salinization for managing salt-affected habitats in Egypt
Dr. Mohammed H. Gomaa, Egypt
- 15:00–15:20 Tea/Coffee Break
- Session 2 Assessment and monitoring of salt-affected soils at national and regional levels (continuation)

Chairperson: **Dr. M. Qadir, ICARDA/IWMI**

Rapporteur: **Dr. Mahmoud Abdelfattah, ICBA**
- 15:20–15:40 Recent evolution of soil salinization in China and its driving processes
Dr. Jingsong Yang, China
- 15:40–16:00 Assessment and management of salt-affected soils of Sudan
Dr. Abdelmagid Ali El-Mobarak, Sudan
- 16:00–17:00 Discussions

Tuesday, 27 November 2007

- Session 3 Modelling for Salinity/Sodicity Development.
- Chairperson: **Dr. Anwar Nasim**, COMSTECH
- Rapporteur: **Dr. Nurul Akhand**, ICBA
- 08:30–09:00 Overview on salinity modelling approaches at different spatial-temporal scales
Dr. Jorge Batlle-Sales, Spain (Keynote presentation)
- 09:00–09:20 SMSS a soil-plant model describing impact of irrigation on salinity of soil and run-off water.
Dr. Mouanis Lahlou, Morocco
- 09:20–10:00 Discussions
- 10:00–10:20 Tea / Coffee Break
- Session 4 Mapping and interpretation of spatial data
- Chairperson: **Dr. Kenneth B. Marcum**, USA
- Rapporteur: **Dr. Shabbir Shahid**, ICBA
- 10:20–10:50 Emerging challenges addressing the characterization and mapping of salt-induced land degradation
Dr. M. Qadir, ICARDA/ IWMI (Keynote presentation)
- 10:50–11:10 Monitoring, predicting and quantifying soil salinity, sodicity and alkalinity in Hungary at different scales. Past experience, current achievements and outlook with special regard to European Union initiatives
Dr. Tibor Toth, Hungary
- 11:10–11:30 Salinization and sodication on irrigated soils in Eastern Kenya
Dr. Patrick Gicheru, Kenya
- 11:30–11:50 Nature and distribution of salts in the upper Ruvu-Wami plains, Morogoro, Tanzania: implications for land management
Dr. Method Kilasara, Tanzania
- 11:50–12:10 Salt-affected soils in Romania
Dr. Elisabeta Dumitru, Romania
- 12:10–12:30 Extent and utilization of salt-affected lands: Biosaline agriculture and marginal resources in Tajikistan
Dr. Sanginov Sanginboy, Tajikistan
- 12:30–13:00 Discussions

- 13:00–14:00 Prayer and Lunch break
- Session 5 Expert rounds on the use of assessment and monitoring tools for management of salt-affected areas
- 14:00–17:30 Participants will be divided into three working groups to provide guidelines and recommendations on the following topics: Assessment and monitoring of salt-affected soils; modelling for salinity/sodicity development; mapping and interpretation of spatial data.

Wednesday, 28 November 2007

Session 5 Expert rounds on the use of assessment and monitoring tools for management of salt-affected areas (continuation)

08:30–11:30 Continuation of expert rounds discussions - Session 5

11:30–12:00 Tea / Coffee Break

12:00–13:00 Presentation of Reports, discussions and recommendations - Session 5

Chairperson: **Dr. Amin Mohamed Mashali**, FAO

13:00–14:00 Prayer and Lunch break

14:00–15:00 Presentation of Reports, discussions and recommendations - Session 5 (continuation)

15:00–15:20 Tea / Coffee Break

Session 6 The way forward for SPUSH

Chairperson: **Dr. Amin Mohamed Mashali**, FAO

Rapporteur: **Dr. Nanduri Rao**, ICBA

15:20–18:00 Determination of priorities for future activities, research and cooperation between member countries.

Communication strategies.

Next network expert consultation.

Conclusions and recommendations.

Thursday, 29 November 2007**Meeting on 'Status and Progress of Biosaline Agriculture'**

- 08:30–10:00 Visit to ICBA: On-going programmes, activities and facilities
- 10:00–10:30 Tea / Coffee Break
- Session 7 Sustainable Biosaline Agricultural Systems
- Chairperson: **Prof. Dr. Faisal Taha**, ICBA
- Rapporteur: **Mrs. Carla Mellor**, ICBA
- 10:30–11:00 Biosaline Agriculture: Prospects and potential within global and regional context
Dr. Shoaib Ismail and Prof. Dr. Faisal Taha, ICBA (Keynote presentation)
- 11:00–11:20 Greywater use for irrigation of home gardens in peri-urban areas of Jordan
Dr. Murad Bino, Jordan
- 11:20–11:40 Saline agriculture: Pakistan scenario
Dr. Riaz H. Qureshi, Pakistan
- 11:40–12:00 Extent of salt-affected land in Central Asia: Biosaline agriculture and utilization of salt-affected resources
Dr. Kristina Toderich, Uzbekistan and **Dr. Shoaib Ismail**, ICBA
- 12:00–12:15 Short break
- 12:15–12:35 Necessity of Biosaline agriculture and irrigation management for sustainable agriculture
Dr. Ali A. Aljalod, Kingdom of Saudi Arabia
- 12:35–12:55 Salinity problem in the Sultanate of Oman: Past, present and future perspectives
Mr. Salim Ben Abdullah Rashid Al-Rasbi, Sultanate of Oman
- 12:55–13:15 Water shortage in Western U.S. and saline recycled water use for urban irrigation
Dr. Kenneth Marcum, USA
- 13:15–13:35 Discussions
- 13:35–14:35 Prayer and Lunch break

Session 8 Future Strategies for Sustainable Biosaline Agricultural Systems

Chairperson: **Dr. Riaz H. Qureshi**, Pakistan

Rapporteur: **Dr. Shoaib Ismail**, ICBA

14:35–16:30 Discussions

Recommendations

16:30–16:45 Tea / Coffee Break

16:45–17:15 Closing Session

Concluding Remarks by IDB/COMSTECH

Concluding Remarks by INBA/ICBA

Concluding Remarks by FAO

ANNEX III

List of participants

Dr. Mahmoud ABDELFAH
Soil Scientist on Secondment, ICBA
P.O. Box 14660, Dubai, UAE.
Tel: +971 4 3361100
Fax: +971 4 3361155
Mobile: +971 50 693 8412
E-mail: m.abdelfattah@biosaline.org.ae
www.biosaline.org

Dr. Ali A. AL-JALOOD
Soil Science and Fertilizer Use Specialist,
King Abdulaziz City for Science
& Technology, Natural Resources &
Environment Research Institute, P.O. Box:
28322, Riyadh, Kingdom of Saudi Arabia.
Tel: +966 4883555/4813610
Fax: +966 4813611
Mobile: +966 5549044
E-mail: aljaloud@kacst.edu.sa

Dr. Ahmed AL MASOUM
Deputy Director General, ICBA,
P.O. Box 14660, Dubai, UAE.
Tel: +971 4 3361100
Fax: +971 4 3361155
Mobile: +971 50 643 6568
E-mail: a.masoum@biosaline.org.ae,
www.biosaline.org

Mr. Salim Ben Abdullah Rashid AL-RASBI
Head of Biosaline Research Laboratory,
Ministry of Agriculture and Fisheries
PO Box 467 – Postal Code: 113, Muscat,
Sultanate of Oman.
Tel: +968 4696280 / 24696367
Fax: +968 24605304
E-mail: agricop@omantel.net.om,
salrasbi@yahoo.com, www.maf.gov.om

Dr. Fawzi AL-SULTAN
Chair of the Board, ICBA
P.O. Box 14660, Dubai, UAE.
Tel: +971 4 3361100
Fax: +971 4 3361155
E-mail: fawzi@fandnconsultancy.com,
www.biosaline.org

Dr. Shawki BARGHOUTI
Director General, ICBA, P.O. Box 14660,
Dubai, UAE.
Tel: +971 4 3361100
Fax: +971 4 3361155
Mobile: +971 50 655 6022
E-mail: s.barghouti@biosaline.org.ae,
www.biosaline.org

Dr. Jorge BATLLE-SALES
Departamento de Biología Vegetal, Facultad
de Farmacia, Universitat de Valencia, Estudi
General, Avda. Vicent Andres Estelles, 46100
Burjasot, Valencia, Spain.
Tel: +34 96 3544289
Fax: +34 963864289
Mobile: +34 653411584
E-mail: Jorge.Batlle@uv.es

Dr. Murad Jabay BINO
Executive Director, Inter-Islamic Network
on Water Resources Development &
Management (INWRDAM), P.O. Box 1460,
PC 11941 Jubieha, Amman, Jordan.
Tel: +962 6 5332993
Fax: +962 6 5332969
Mobile: 079 5545311
E-mail: inwrdam@nic.net.jo,
muradinw@nic.net.jo

Dr. Giuseppina CRESCIMANNO
Head of Science, UNESCO-BRESCIA in
Venice and Professor at Palermo University,
Dipartimento d'Ingegneria e Tecnologie
Agro-Forestali ITAF,
Viale delle Scienze, 09128 Palermo, Italy.
Tel: +39 091 7028109
Fax: +39 091 484035
E-mail: gcrescim@mbox.unipa.it,
g.crescimanno@unesco.org, gcrs@hotmail.
com, gcrescim@unipa.it, <http://portal.unesco.org/en/>

Dr. Abdullah DAKHEEL
Field and Forage Crop Scientists, ICBA
P.O. Box 14660, Dubai, UAE.
Tel: +971 4 3361100
Fax: +971 4 3361155
Mobile: +971 50 623 6580
E-mail: a.dakheel@biosaline.org.ae,
www.biosaline.org

Dr. Elisabeta DUMITRU
National Research and Development
Institute for Soil Science, Agricultural
Chemistry and Environment (ICPA)
Bucharest Bulevard Marasti Nr. 61;
Romania.
Tel: +40 21 318 4349 / 4458
Fax: +40 21 3184348
E-mail: office@icpa.ro, elisa_dumitruu@
yahoo.com

Dr. Abdelmagid Ali EL-MOBARAK
Researcher, Agriculture Research
Corporation, Land and Water Research
Center, P.O. Box 388, Wadi Medani, Sudan.
Tel: +249 51 143035
E-mail: melmobarak@yahoo.com,
melmobarak@maktoob.com

Dr. Patrick GICHERU
Director, National Agricultural Research
Laboratories, Box 14733, Nairobi, Kenya.
E-mail: cdnarl@iconnect.co.ke

Dr. Mohammed H. GOMAA
Tel: +2 02 3372441
Fax: +2 02 3373048
Home: +2 02 6708394
Mobile: +2 0102891442
E-mail: gomaa_1999@yahoo.com

Mr. Y. HASHEMINEJHAD
Chairman Sector Land Reclamation,
Ministry of Agriculture and Land
Reclamation, Nady Al-saaed street, Dokki,
Giza. National Salinity Research Center,
Iran.
Tel:+98 351 7212311
Fax:+98 3517212312
E-Mail: cheraghi@insrc.org

Dr. Rungsun IM-ERB
Chief Soil Salinity Section, Land
Development Department, Ministry of
Agriculture and Cooperative, Phahonyothin
Road, Chatuchak, Bangkok 10900, Thailand.
Tel: +66 2 579 5546/0111 ext.1217
Fax: +66-2-5620312
Home: +66 2 5891172
Mobile: +66 011736371
E-mail: rungsuni@hotmail.com,
rungsun@ladd.go.th

Dr. Shoaib ISMAIL
Halophyte Agronomist, International
Center for Biosaline Agriculture, ICBA,
P.O. Box 14660, Dubai, UAE.
Tel: +971 4 3361100
Fax: +971 4 3361155
Mobile: +971 50 5385952
E-mail: s.ismail@biosaline.org.ae,
www.biosaline.org

Mr. Kayan JAFF
Programme Officer/FAO Representative,
UNDP UAE
Box 3490 United Arab Emirates, UN
House, Baniyas Street 18/2 Plot 86 Abu
Dhabi, UAE.
Tel: +971 2 4493248
Fax: +971 2 4495112
E-mail: FAO-AE@fao.org

Dr. Method KILASARA
Department of Soil Science, Sokoine
University of Agriculture, P.O. Box 3008
Morogoro, Tanzania.
Tel: +255 732930160
Mobile: +255 754 493668
E-mail: mmkilasara@yahoo.com

Dr. Mouanis LAHLOU
IAV Hassan II, P.O. BOX 6202, Rabat-
Institute, Rabat, Morocco.
Tel: +212 37 7717 58/59/45
Fax: +212 37 778135
E-mail: m.lahlou@iav.ac.ma,
mouanis.lahlou@gmail.com

Dr. Clemencia LICONA MANZUR
Soil Reclamation and Development
Officer, Agriculture Department, Food and
Agriculture Organization of the UN, Viale
delle Terme di Caracalla, 00153 Rome, Italy
Tel: +39 06 570 55044
Fax: +39 06 570 56275
E-mail: Clemencia.LiconManzur@fao.org,
<http://www.fao.org>

Dr. Kenneth B. MARCUM
Dept. of Applied Biological Sciences,
Wanner Hall, Arizona State University, East
Campus, 7001 E. Williams Field Rd., Mesa,
AZ 85212, U.S.A.
Tel: +1 480 727 1213
E-mail: Kenneth.Marcum@asu.edu,
<http://ctas.poly.asu.edu/marcum>

Dr. Amin Mohamed MASHALI
FAO Consultant
Viale delle Terme Di Caracalla-00153 Rome,
Italy.
Tel: +39 06 57053806
Fax: +39 06 57056275
Home: +39 06 51957805
E-mail: amin.mashali@fao.org

Mrs. Carla MELLOR
Librarian, ICBA,
P.O. Box 14660, Dubai, UAE.
Tel: +971 4 3361100
Fax: +971 4 3361155
Mobile: +971 50 256 1018
E-mail: c.mellor@biosaline.org.ae,
www.biosaline.org

Dr. Anwar NASIM
Advisor Science, COMSTECH
COMSTECH Secretariat, 3-Constitution
Avenue,
G-5/2, Islamabad 44000, Pakistan.
Tel: +92 51 9220681/3
Fax: +92 51 9211115/9220265
E-mail: anwar_nasim@yahoo.com,
comstech@isb.comsats.net.pk,
www.comstech.org.pk

Dr. J.P. NELL
ARC-Institute for Soil, Climate and Water,
Private Bag X79, Pretoria 0001, South
Africa.
Tel: +27 12 310 2600
Fax: + 27 12 323 1157
Mobile: 082 772 6914
E-mail: pnell@arc.agric.za

Dr. Manzoor QADIR
IWMI-ICARDA Joint Appointment
Marginal -quality Water Resources
Management and Salt-prone Soils,
International Center for Agricultural
Research in the Dry Areas (ICARDA), P.O.
Box 5466, Aleppo, Syria.
Tel: +963 21 2213433
Fax: +963 21 2213490
E-mail: m.qadir@cgiar.org,
<http://www.icarda.org>

Dr. Riaz H. QURESHI
Advisor (QA & LI), Higher Education
Commission, Sector H9, Islamabad,
Pakistan.
Tel: +92 51 4444676
Fax: 051-9258744
Mobile: +92 300 5071063
E-mail: rqureshi@hec.gov.pk,
www.hec.gov.pk

Dr. Nanduri RAO
Plant Genetic Resource Scientist, ICBA
P.O. Box 14660, Dubai, UAE.
Tel: +971 4 3361100
Fax: +971 4 3361155
Mobile: +971 436 3929
E-mail: n.rao@biosaline.org.ae,
www.biosaline.org

Dr. Sanginov SANGINBOY
Tajikistan Academy of Agricultural Sciences,
21a Rudaki Prospect, Dushanbe 734025,
Tajikistan.
Tel: +992 372 271979
Fax: +992 372 213207
Mobile: +992 93 5700710
E-mail: soil@tajik.net

Dr. Shabbir SHAHID
Salinity Management Scientist, ICBA,
P.O. Box 14660, Dubai, UAE.
Tel: +971 4 3361100
Fax: +971 4 3361155
Mobile: +971 50 413 4149
E-mail: s.shahid@biosaline.org.ae,
www.biosaline.org

Dr. Gurbachan SINGH
Director, Central Soil Salinity Research
Institute (CSSRI), Indian Council of
Agricultural Research. Zarifa Farm,
Kachhwa Road, Karnal – 132001, India.
Tel: +91 184 280801
Fax: +91 184 280480
E-mail: director@cssri.ernet.in

Dr. Faisal TAHA
Director Technical Programme ICBA
P.O. Box 14660, Dubai, UAE
Tel: +971 4 3361100
Fax: +971 4 3361155
E-mail: f.taha@biosaline.org.ae
www.biosaline.org

Dr. Kristina TODERICH
Plant Scientist, ICBA. 6 Murtazaeva Street,
PO Box 4564, 700000 Tashkent, Uzbekistan.
Tel: +998 71 1372130
Fax: +998 71 1207125
E-mail: k.toderich@cgiar.com
ktoderich@yahoo.com

Dr. Tibor TOTH
RISSAC (Research Institute for Soil
Science and Agricultural Chemistry of the
Hungarian Academy of Sciences, MTA
TAKI, Herman Otto ut 15, Budapest, H-
1022 Hungary.
Tel/Fax: +36 1 1558839
Telex: 227223 AGROK-H
E-mail: h3700tot@ella.hu

Eng. Mohamed Ali TOURE
Acting Director & Senior S&T Expert
Science & Technology Office, Islamic
Development Bank, P.O. 5925, Jeddah
21432, Saudi Arabia.
Tel. +966 2 6466935
Fax: +966 2 6467828
E-mail: MToure@isdb.org,
<http://www.isdb.org>

Dr. Jingsong YANG
Head of Soil Physics and Soil Salinity
Division, The Institute of Soil Science,
Chinese Academy of Sciences, 71 East
Beijing Road, P.O. Box 821, 210008 Nanjing,
Jiangsu, China.
Phone: +86 25 86881222
Fax: +86 25 86881000
E-mail: jsyang@issas.ac.cn

Advances in the assessment and monitoring of salinization and status of biosaline agriculture

Report of an expert consultation held in Dubai, United Arab Emirates, 26–29 November 2007

The causes of salinity and sodicity, which vary between countries and regions, need to be identified, assessed and monitored carefully so that they can be managed and controlled. There is a need for practical and cost-effective methodologies for assessing, monitoring and mapping the extent and distribution of salt-affected soils; for identifying the causes and sources of the problem; and for choosing management options and evaluating the effectiveness of those options. The objective of the Expert Consultation on Advances in Assessment and Monitoring of Salinization for Managing Salt-affected Habitats was to exchange experiences with data collection and analysis for the assessment and monitoring of salinity and sodicity, with particular emphasis on practical applications at local, national, regional and global levels. The Meeting on the Status and Progress of Biosaline Agriculture of the Inter-Islamic Network on Biosaline Agriculture was an opportunity to present the work of the hosting institution and to exchange information between the two Networks.

ISBN 978-92-5-106439-9 ISSN 0532-0488



9 789251 064399

11220E/1/10.09/700



Università degli Studi
di Palermo



ASU
ARABIAN SCIENCE UNIVERSITY

