

Cultch and oyster density standards  
for reef restoration,  
sustainable harvest  
and  
fishing at  
maximum sustained yield

Tom Soniat, Nathan Cooper  
University of New Orleans

Eric Powell, Laura Solinger, Kathleen Hemeon, Sara Pace  
University of Southern Mississippi

GULF STATES MARINE FISHERIES COMMISSION

10/12/2021

### Academic Partners:

University of New Orleans--Tom Soniat, Nathan Cooper

University of Southern Mississippi--Eric Powell, Sara Pace, Kathleen Hemeon,  
Laura Solinger

### State Agency Partners:

Texas Parks and Wildlife Department--Christine Jensen

Louisiana Department of Wildlife and Fisheries--Carolina Bourque

Mississippi Department of Marine Resources-- Eric Broussard, Jason Rider

Alabama Department of Conservation and Natural Resources--Jason Herrmann

Florida Fish and Wildlife Conservation Commission--Ryan Gandy

### Industry Partners:

Hillman Shrimp and Oyster Company (Texas)--Clifford Hillman

AmeriPure Oysters (Louisiana)--John Tesvich

Crystal Seas Oysters (Mississippi)--Jennifer Jenkins

Bon Secour Fisheries (Alabama)--Chris Nelson

Water Street Seafood (Florida)--Steven Nash

### Federal Partner:

NOAA Saltonstall-Kennedy Program

# Introduction to the model


- Initially developed for Louisiana oyster fishery
- Annual stock assessment provides: oyster density, oyster numbers by size, cultch density and reef area
- These are inputs to the model which simulates growth, natural and fishing mortality, and cultch loss or gain
- Shells of dead oysters are added to the reef and shells of fished oysters are debited
- Growth and mortality are size and time dependent
- Fishing can occur for seed and/or sack oysters
- Fishing rate is time dependent
- Initial sustainability goal: no-net cultch loss (NNL)

# Incorporation of cultch and oyster density standards

- New sustainability goal: oyster density/cultch density standard (OCS) for fishing at Maximum Sustained Yield (*msy*)
- How was the OCS derived?  
(Solinger et al., submitted to *Canadian Journal of Fisheries and Aquatic Sciences*)

# How many oysters can be harvested?


- Oyster population at equilibrium assumed (unchanging live population and reef structure)
- Dermo mortality = 10% chosen as common
- Maximum catch occurs between 20 and 25% F, yielding 2.9 million oysters
- Thus, at a level of 10% Dermo mortality, an F of about 20% is recommended
- Total mortality (Natural + Dermo + Fishing) is then 40% per year



Increasing Catch

# What is the required oyster density?


About 33 market-sized oysters are required to support an oyster population with a Dermo mortality of 10% and fishing mortality of 20%





# What is the required cultch density?

- As is expected, increasing fishing mortality (F) reduces volume of cultch per square meter
- Note that cultch is relatively consistent with Dermo, because Dermo continues to contribute cultch to the shell bed
- At  $D = 0.10$  and  $F = 0.20$ , 2.2 kg of cultch is needed to sustain the population




# Application of oyster and cultch density standard (OCS) to the Gulf of Mexico

OCS: 30 oysters/m<sup>2</sup> (> 75 mm)  
2000 g/m<sup>2</sup> cultch (surficial)



# Sample Locations



# Mean monthly water temperatures

Month	Temp. (°C)
January	11
February	12
March	16
April	20
May	24
June	28
July	29
August	29
September	28
October	23
November	17
December	13

Data from Eugene Island, central coastal Louisiana:

<https://www.currentresults.com/Oceans/Temperature/louisiana-alabama-average-water-temperature.php#c>



# Low, moderate and high salinity year assumptions

Month	Low Salinity	Moderate Salinity	High Salinity
J	9.8	18.3	18.8
F	9.0	15.0	21.8
M	7.3	15.3	20.5
A	6.5	13.8	22.3
M	7.3	12.8	21.8
J	9.0	12.8	17.8
J	5.5	11.3	16.3
A	7.3	11.0	18.5
S	9.0	14.0	19.3
O	11.8	16.0	23.0
N	12.3	16.3	23.8
D	11.5	16.0	25.0
<b>Annual Mean</b>	<b>8.8</b>	<b>14.4</b>	<b>20.7</b>


# Gulf of Mexico simulated harvests

	Salinity	TX		LA		AL		FL	
<b>OCS</b>		Sack	Seed	Sack	Seed	Sack	Seed	Sack	Seed
	Low	0	N/A	0	0	0	N/A	26,732	N/A
	Moderate	0	N/A	0	0	0	N/A	5,228	N/A
	High	0	N/A	0	0	0	N/A	0	N/A
<b>NNL</b>									
	Low	1,130,391	N/A	478,229	25,203	31,343	N/A	62,070	N/A
	Moderate	941,158	N/A	458,929	6,741	230	N/A	14,690	N/A
	High	439,469	N/A	267,104	8,043	9,706	N/A	246	N/A

Soniat et al., submitted to JSR

- Sack and seed harvest is in Louisiana sacks (52.85 Liters)
- Salinity is the monthly salinity scenario
- Low salinity scenario yields largest harvests
- Seed fishing harvest in LA only
- Sustainable fishing by OCS standard only in FL
- Sack harvest allowable under NNL in all States
- Seed harvest allowable under NNL standard in LA
- Texas has the highest allowable harvest under NNL standard

# Oyster reef management: a general theory of oyster reef functionality




# Oyster reef management: cultch

- Shell is a temporary resource
  - Shell half-lives measured *in situ* confirm that shell half lives are about 5 years in the Gulf of Mexico (Pace et al. 2020. JSR 39:245–256)
- Clean shell (ESA) diminishes very quickly
  - The decay rate of the “clean factor” associated with newly planted shell has been measured: Half life = 0.81 years (Ashton-Alcox 2021. JSR 40:191-211). Implies annual planting under NNL.



# Oyster reef management: cultch vs. recruitment



65 year  
time series

Delaware Bay


Moderate  
mortality reefs

Hemond et al.  
2020. JSR  
39:633-645,

- Differential recruitment among substrate types and regions in Delaware Bay were used to estimate ESA (clean cultch)
- Larvae preferentially set on live oysters and boxes vs. other cultch
- Habitat quality is a surrogate for broodstock.  
**Total ESA defines recruitment potential!**
- The relationship follows a Beverton-Holt process (asymptomatic)  
**Excessive ESA/cultch does not always lead to increased recruitment!**
- There is a non-zero level of ESA/cultch at which recruitment ceases  
**Reefs can be *effectively* extinct with ESA/cultch still present!**

# Oyster reef management: cultch vs. live surface area

- Haskin Rule: minimum ratio 40% live animals to cultch, used in Delaware Bay. Fishing ends with < 40% ratio
- Live surface area to other cultch assessed here
- Model results converge on Haskin “rule of thumb”
- Live ratio/Haskin Rule could serve as a “quick and dirty” check of overfishing



Greater Proportion of Live Surface Area

# Oyster reef management: NNL vs OCS

- NNL
  - Doesn't include oyster density/cultch density standard
  - Greater harvests vs OCS (presently)
  - Not self-sustaining (requires cultch planting)
  - Provides no benchmarks for reef restoration
- OCS
  - Includes oyster density/cultch density standard
  - Restricted harvests vs NNL (presently)
  - Self sustaining (no cultch planting required)
  - Provides benchmarks for reef restoration
  - Identifies harvest at maximum sustained yield
- OCS/NNL Hybrid
  - Work toward the cultch standard but fish as NNL
  - Be cognizant of the oyster density standard
  - Don't allow cultch density of  $< 1000 \text{ g/m}^2$

# Oyster reef management: summary

Requirements for successful management of oyster reefs and sustainable fishing are available

- Oyster density and cultch standards serve as restoration targets
- Simulations using OCS provide harvest estimates at maximum sustained yield

# Future Activities

- Transfer code (Ruby) to Agencies  
(make request to Nathan Cooper, [ncooper@my.uno.edu](mailto:ncooper@my.uno.edu))
- Hold R modeling session for Agency personnel  
Jan./Feb. ? 2022  
([Laura.Solinger @usm.edu](mailto:Laura.Solinger@usm.edu))

Thanks to all!