

**MATH 651
 HOMEWORK II
 PARTIAL SOLUTIONS
 SPRING 2013**

Problem 6. A free loop in a space X is a map $S^1 \rightarrow X$ with no condition on the basepoint. Since $\pi_1(X, x_0) \cong [S^1, (X, x_0)]_*$, there is a natural map

$$\Lambda : \pi_1(X, x_0) \rightarrow [S^1, X].$$


- (1) Show that Λ is surjective if and only if X is path-connected.
- (2) Show that $\Lambda([\gamma]) = \Lambda([\delta])$ if and only if the loops γ and δ are conjugate in $\pi_1(X, x_0)$.

Solution. Recall that $S^1 \cong [0, 1]/\{0, 1\}$. It will be convenient to go back and forth between maps $S^1 \rightarrow X$ and maps $[0, 1] \rightarrow X$ which agree at the endpoints. We can use this for homotopies as well: a homotopy $h : S^1 \times I \rightarrow X$ is equivalent to a homotopy $h' : I \times I \rightarrow X$ such that $h'(0, t) = h'(1, t)$ for every t .


- (1) (\Rightarrow) Suppose that Λ is surjective. Let x_1 be a point in X . Then there is a loop γ at x_0 such that $\Lambda([\gamma]) = [c_{x_1}]$. In other words, there is a homotopy $h : \gamma \simeq c_{x_1}$. Then $h(0, t)$ defines a path in X from x_0 to x_1 .

(\Leftarrow) Assume that X is path connected, and let $f : S^1 \rightarrow X$ be any map. We write δ for the resulting path $[0, 1] \rightarrow S^1 \rightarrow X$ and x_1 for the basepoint of δ . Let φ be any path from x_0 to x_1 . Then the path-composite $\varphi * (\delta * \varphi^{-1})$ is a loop at x_0 . Write γ for this loop. It remains to show that $\Lambda([\gamma]) = [\delta]$. In other words, we must show that $\gamma \simeq \delta$ as maps $S^1 \rightarrow X$. A homotopy is given by

$$h(s, t) = \begin{cases} \varphi(t + 2s) & 0 \leq 2s \leq 1 - t \\ \delta\left(\frac{4s - 2 + 2t}{1 + 3t}\right) & 2 - 2t \leq 4s \leq 3 + t \\ \varphi(4(1 - s) + t) & 3 + t \leq 4s \leq 4. \end{cases}$$



- (2) (\Rightarrow) Suppose that $\Lambda([\gamma]) = \Lambda([\delta])$. Then there is a homotopy $h : \gamma \simeq \delta$ as maps $S^1 \rightarrow X$. Note that $h(0, t)$ defines a loop φ at x_0 . Since h is a homotopy through maps out of the circle, $h(1, t)$ is the same loop φ . We want to show that $[\gamma] = [\varphi * \delta * \varphi^{-1}]$. A path-homotopy is illustrated in the figure to the right.



Said differently, let $i_0 : I \rightarrow I \times I$ be the inclusion as the bottom edge and let $i_1 : I \rightarrow I \times I$ be a path that first travels up the vertical left edge, then across the top edge, then down the right edge. By the definition of h , the path γ is $h \circ i_0$ and the path $\varphi * \delta * \varphi^{-1}$ is $h \circ i_1$. But the two maps i_0 and i_1 are certainly path-homotopic, as $I \times I$ is contractible (even convex). It follows that $\gamma \simeq_p \varphi * \delta * \varphi^{-1}$.

(\Leftarrow) Assume $h : \gamma \simeq_p \varphi * \delta * \varphi^{-1}$ is a path-homotopy for some loop φ . Since h is a path-homotopy, it must be constant on the vertical edges of the square $I \times I$. Collapsing each of these edges produces a disc D^2 , and we get an induced map $\bar{h} : D^2 \rightarrow X$. The desired free homotopy $H : \gamma \simeq \delta$ is given by the following composition, where the first map is any homeomorphism identifying the boundary as indicated.

