

WFW2  
TJB  
GA

NOAA Technical Memorandum NWS WR-176



---

APPROXIMATIONS TO THE PEAK SURFACE WIND GUSTS  
FROM DESERT THUNDERSTORMS

Salt Lake City, Utah  
June 1982



NOAA TECHNICAL MEMORANDA  
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P. O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U. S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22151. Prices vary for all paper copy; \$3.50 microfiche. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operational Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidli, April 1968 (revised February 1981).
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB-178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. Capt. Thomas D. Murphy, U.S.A.F., May 1968. (AD-673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB-179084)
- 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB-179289)
- 35 Joint ESSA/FAA ARTC Radar Weather Surveillance Program. Herbert P. Benner and DeVon B. Smith, December 1968 (revised June 1970). AD-681857)
- 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (PB-183055)
- 37 Disposal of Logging Residues without Damage to Air Quality. Owen P. Cramer, March 1969. (PB-183057)
- 39 Upper-Air Lows over Northwestern United States. A. L. Jacobson, April 1969. (PB-184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970's. L. W. Snellman, August 1969. (PB-185068)
- 42 Analysis of the Southern California Santa Ana of January 15-17, 1966. Barry B. Aronovitch, August 1969. (PB-185670)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB-185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB-187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB-190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J. C. Burnash, December 1969. (PB-188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB-190157)
- 49 Predicting Precipitation Type. Robert J. C. Burnash and Floyd E. Hug, March 1970. (PB-190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB-191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB-193102)
- 52 Sacramento Weather Radar Climatology. R. G. Pappas and C. M. Veliquette, July 1970. (PB-193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB-194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB-194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB-194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM-71-00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM-71-00016)

NOAA Technical Memoranda (NWS WR)

- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon. Arthur W. Fritz, October 1970. (COM-71-00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM-71-00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM-71-00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM-71-00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM-72-10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM-72-10554)
- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM-72-10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM-72-11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM-72-11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM-72-11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM-72-10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM-72-10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM-73-10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (revised March 1978). (COM-73-10251)
- 83 A Comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM-73-10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM-73-11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y. G. Lee, June 1973. (COM-73-11276)
- 89 Objective Forecast Precipitation over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, Sept. 1973. (COM-73-11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM-73-10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM-74-11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM-74-11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM-74-11555/AS)
- 96 Map Type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM-75-10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB-250-711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM-75-10719/AS)
- 99 A Study of Flash Flood Susceptibility--A Basin in Southern Arizona. Gerald Williams, August 1975. (COM-75-11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB-246-902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB-253-053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB-252-866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB-254-650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB-254-649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB-259-594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB-260-437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB-273-677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB-264-655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB-265-941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB-273-694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB-273-676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB-268-847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R. F. Quiring, April 1977. (PB-272-831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB-268-740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB-271-290/AS)

NOAA Technical Memorandum NWS WR-176

APPROXIMATIONS TO THE PEAK SURFACE WIND GUSTS  
FROM DESERT THUNDERSTORMS

Darryl Randerson

National Weather Service Nuclear Support Office  
Las Vegas, Nevada  
June 1982

UNITED STATES  
DEPARTMENT OF COMMERCE  
Malcolm Baldrige, Secretary

National Oceanic and  
Atmospheric Administration  
John V. Byrne, Administrator

National Weather  
Service  
Richard E. Hailgren, Director



This publication has been reviewed  
and is approved for publication by  
Scientific Services Division,  
Western Region.

A handwritten signature in cursive script, reading "L. W. Snellman". The signature is written in black ink and is positioned above the typed name and title.

L. W. Snellman, Chief  
Scientific Services Division  
Western Region Headquarters  
Salt Lake City, Utah

## CONTENTS

	<u>Page</u>
Figures .....	iv
Tables .....	iv
Abstract .....	1
I. Introduction .....	1
II. Procedure .....	2
III. Data .....	3
IV. Analysis .....	6
V. Application .....	13
VI. Summary .....	14
VII. Acknowledgements .....	14
VIII. References .....	16

FIGURES

	<u>Page</u>
Figure 1. Plot of observed $V_m$ and $\Delta T$ data for 49 thunderstorms that occurred over southern Nevada. The solid line is the line of best fit to these data and is given by Eq. 1. The 90 percent confidence limits are plotted as dashed lines. ....	7
Figure 2. Exponential plot of the 49 cases on log-linear paper. The solution to Eq. 3 is plotted as the solid line. ....	9
Figure 3. Exponential plot for 10 cases where the difference between observed and predicted $\Delta T$ is $\leq 5^\circ\text{F}$ . Solid line is the line of best fit as given by Eq. 6. ....	12
Figure 4. Reproduction of Fig. 1 with only the plotted data. ....	15

TABLES

Table 1. Tabulation of data used to develop Figures 1, 2, and 3. Stations (STN) used are Yucca Flat (UCC), Las Vegas (LAS), and Desert Rock (DRA). Observed maximum temperatures ( $T_m$ ) and observed thunderstorm-associated minimum temperatures ( $T_{\min}$ ) are in $^\circ\text{F}$ and come from MF1-10B forms. Observed minimum temperatures are assumed to be those occurring with the thunderstorm downdraft and are not the minimums observed near sunrise. The observed temperature difference ( $\Delta T$ ) is in $^\circ\text{F}$ . The peak wind gust ( $V_m$ ) is in mph and comes from MF1-10B forms. Total precipitation for the day (R) is in inches. The minimum downdraft temperature derived from the sounding data from UCC (and DRA) are listed under the $T_e$ column in $^\circ\text{F}$ . Calculated values of $\Delta T$ are listed under the $\Delta T_c$ column and represent $T_m - T_e$ . ....	4
Table 2. Data listing of maximum wind gust ( $V_m$ ) by wind gust temperature ( $\Delta T$ ) category. The median $V_m$ value in each $\Delta T$ category is underlined in the $V_m$ column. The average maximum surface wind gust ( $\bar{V}_m$ ) is calculated for each $\Delta T$ category. ....	5

Table 3.	Tabulation of the estimates of $V_m$ according to the regression equations developed in this report. ....	11
----------	---	----

# APPROXIMATIONS TO THE PEAK SURFACE WIND GUSTS FROM DESERT THUNDERSTORMS

Darryl Randerson  
Nuclear Support Office  
National Weather Service

ABSTRACT. Simple procedures for estimating the peak surface wind gusts from desert thunderstorms are proposed. The fundamental assumption is that the peak surface wind gust is related to the difference ( $\Delta T$ ) between the maximum ambient air temperature attained prior to the thunderstorm and the minimum temperature of the cooler air generated by the thunderstorm downdraft. A total of 49 independent cases are analyzed statistically. Six different regression equations are developed. The simplest and perhaps the most useful expression suggests that the range of the peak gust can be calculated as  $(15 + \Delta T) \pm 12$  mph with 90 percent confidence, provided  $\Delta T$  is forecast perfectly.

## I. INTRODUCTION

One of the important forecast problems associated with the prediction of thunderstorms is to ascertain the maximum speed of the surface wind gust accompanying the storm. Simple procedures for estimating this gust are addressed in this report. The proposed procedures are simple mathematical expressions derived from regression analysis of a rather homogeneous set of thunderstorm data.

Pioneering work on estimating the peak surface wind gust from thunderstorms was completed by Brancato (1942) and by Jordan (1945). Based on these works, Fawbush and Miller (1954) developed a forecasting scheme for predicting the peak surface wind gust. Their scheme was adopted and modified slightly by the Air Weather Service (1956). This procedure and another outlined by Miller (1972) are used by the Air Weather Service (Crisp, 1979) for predicting surface wind gusts accompanying severe thunderstorms in the eastern half of the United States. The basis for the Fawbush-Miller technique is that the peak surface wind gust is related to the difference between the ambient surface temperature observed just prior to the thunderstorm and the temperature associated with the thunderstorm downdraft. Specifically, the Fawbush-Miller technique relates the peak wind gust to the difference between the prethunderstorm surface temperature and the surface temperature of the saturation adiabat passing through the wet-bulb temperature at the freezing level. This technique has not functioned satisfactorily with desert thunderstorms. Observed minimum temperatures accompanying thunderstorm downdrafts are usually 10 to 15° F warmer than those predicted by the Fawbush-Miller scheme. In addition, the bases of desert thunderstorms tend to be higher than those associated with midwestern thunderstorms (MacDonald,



1976). Higher cloud bases generally result from the small mixing ratios occurring over the desert southwest. In fact, for many desert thunderstorms, moist adiabatic descent to the ground may be terminated with complete evaporation of the precipitation before the downdraft reaches the ground. Dry adiabatic descent will then ensue, as the downdraft continues to accelerate downward. Upon reaching the ground, the downdraft air spreads out under the thunderstorm, creating the often observed haboob (Idso et al., 1972) over the desert regions of the world.

Analyses of atmospheric soundings taken during desert thunderstorm situations in southern Nevada tend to show that when precipitation reaches the ground the maximum surface wind gust is part of a moist adiabatic downdraft that descends to the ground and spreads out under the thunderstorm. The saturation adiabat followed during this process is the one passing through the convective condensation level (Saucier, 1959, p. 71) and intersecting the ground at the observed surface pressure. The temperature difference between the observed surface temperature theoretically identified by this adiabat and the observed maximum ambient air temperature just prior to the thunderstorm is highly correlated with the observed peak surface wind gust ( $V_m$ ).

## II. PROCEDURE

A survey of the MF1-10A and 10B forms for Yucca Flat (UCC), Desert Rock (DRA), and Las Vegas, Nevada (LAS) revealed that local thunderstorms form during a variety of weather conditions and at almost any time of day (Quiring, 1972). Moreover, the data show great variability in the measured peak surface wind gust for individual thunderstorms. Such variety is common in meteorology. One approach to simplifying such a complex physical situation is to attempt to homogenize the data set. Homogenization is justified on the basis of finding physical phenomena with uniform dynamics and constant forcing functions. The following set of criteria were used to select a homogeneous set of thunderstorm events for this study:

1. To help restrict the thunderstorm activity to that modulated by summertime surface heating, consider only those thunderstorms that occurred in the period June through September between the hours of 1000 LST and 2200 LST daily.
2. To obtain similar amounts of surface heating for each case, no ceilings were permitted below 20,000 feet AGL although high broken to overcast conditions were acceptable.
3. No precipitation was permitted before 1000 LST.
4. To help assure that the observed surface wind gusts were associated with moist adiabatic descent, the thunderstorms must have been observed to occur at a weather station and have been recorded in column 5 of MF1-10A (formerly WBAN-10A).
5. To help assure that descending air followed the moist adiabatic process to the ground, at least 0.01" of precipitation had to be measured during the thunderstorm.

6. The peak wind gust must have been observed to occur during the thunderstorm.
7. Estimated values of the peak wind gust (e.g. July 19, 1969, at UCC) were not allowed; the peak wind gust must have been read from a recorder by an observer on duty at a weather station.
8. For days with more than one thunderstorm occurrence, only the peak surface wind gust from the first storm was used.

All the above criteria had to be met for a peak surface wind gust to be entered into the developmental data base.

### III. DATA

All the data used were collected from three stations located in southern Nevada. Data from UCC, DRA, and LAS were used because they were easily accessible. Both UCC and DRA are located on the Nevada Test Site (NTS). UCC is near the center of the NTS on the western edge of a normally dry lake bed at an elevation of nearly 1,200 m above mean sea level (MSL). This station is surrounded by mountains with the highest terrain to the northwest rising nearly 1,000 m above the valley floor. DRA, located 40 km south of UCC, is situated in the southern part of the NTS at an elevation of nearly 1,000 m above MSL. The terrain slopes gradually upward to the north. The northern end of the Spring Mountains is to the south of DRA. Located approximately 120 km southeast of the NTS and 10 km south of downtown Las Vegas, at McCarran International Airport, LAS is nearly 660 m above MSL. This WSO station is in a broad valley surrounded by mountains ranging from near 600 m to 3,000 m above the valley floor (near 550 m above MSL). The tallest mountains are west and north of the city.

The developmental data set consists of 49 thunderstorm-generated, peak surface wind gusts during the period June through September for 1963 through 1980. Included in this data sample are 28 cases from UCC for 1963 through 1977, 18 cases from LAS for 1971 through 1980, and 3 cases from DRA from 1978 through 1980. All the data used are tabulated in Table 1 and categorized in Table 2.

Peak surface wind gust data were extracted from columns 71, 72, and 73 of Form MF1-10B (formerly WBAN 10B). To confirm that the peak wind gust was from the observed thunderstorm, the MF-10B data were compared with the hourly (and special) observations entered on Form MF1-10A (formerly WBAN 10A). On Form MF1-10A, special attention was given to present weather conditions, to reported surface winds, and to the remarks column where "PK WND" reports are listed.

Observations of the maximum temperature prior to the thunderstorm and of the minimum temperature during the storm were obtained from columns 47 and 48 of Form MF1-10B, respectively. These observations were compared with the hourly observations and with columns 82, 83, and 84 of MF1-10B to confirm that

Table 1. Tabulation of data used to develop Figures 1, 2, and 3. Stations (STN) used are Yucca Flat (UCC), Las Vegas (LAS), and Desert Rock (DRA). Observed maximum temperatures ( $T_m$ ) and observed thunderstorm-associated minimum temperatures ( $T_{min}$ ) are in  $^{\circ}F$  and come from MF1-10B forms. Observed minimum temperatures are assumed to be those occurring with the thunderstorm downdraft and are not the minimums observed near sunrise. The observed temperature difference ( $\Delta T$ ) is in  $^{\circ}F$ . The peak wind gust ( $V_m$ ) is in mph and comes from MF1-10B forms. Total precipitation for the day ( $R$ ) is in inches. The minimum downdraft temperature derived from the sounding data from UCC (and DRA) are listed under the  $T_e$  column in  $^{\circ}F$ . Calculated values of  $\Delta T$  are listed under the  $\Delta T_c$  column and represent  $T_m - T_e$ .

DATE	STN	Observed					Calculated	
		$T_m$	$T_{min}$	$\Delta T$	$V_m$	$R(in)$	$T_e$	$\Delta T_c$
081563	UCC	97	76	21	38	.05	65	32
081265	UCC	92	71	21	30	.18	66	26
081365	UCC	94	68	26	37	.12	64	30
072366	UCC	97	78	19	42	.56	66	31
072966	UCC	100	64	36	48	.36	60	40
081866	UCC	97	79	18	37	.13	61	36
080967	UCC	94	76	18	35	.05	65	29
081467	UCC	98	85	13	25	.01	63	35
081967	UCC	98	72	26	35	.16	68	30
090667	UCC	87	66	21	22	.39	59	28
070668	UCC	97	65	32	41	.05	61	36
070768	UCC	93	65	28	44	.21	M	M
080668	UCC	95	64	31	47	.25	65	30
071369	UCC	96	72	24	36	.10	64	32
071869	UCC	97	62	33	54	.77	66	31
091669	UCC	83	69	14	31	2.13	57	26
081670	UCC	94	74	20	30	.03	66	28
073071	LAS	110	93	17	40	.05	65	45
080871	LAS	104	73	31	58	.35	65	39
060472	UCC	90	63	27	51	.01	57	33
060472	LAS	96	84	12	28	.01	57	39
071672	LAS	107	72	35	54	.13	67	40
081472	LAS	89	81	8	21	.12	62	27
073074	UCC	102	72	30	31	.57	62	40
080174	LAS	102	93	9	26	.01	63	39
080274	UCC	95	67	28	41	.03	64	31
090574	LAS	103	85	18	28	.16	64	39
070475	LAS	95	86	9	31	.01	63	32
072675	UCC	104	90	14	34	.01	62	42
072775	UCC	100	70	30	37	.02	61	39
091675	LAS	94	71	23	24	.92	59	35
091875	UCC	90	86	4	19	.08	64	26
072476	LAS	100	75	25	43	.20	64	36

DATE	STN	Observed					Calculated	
		T <sub>m</sub>	T <sub>min</sub>	ΔT	V <sub>m</sub>	R(in)	T <sub>e</sub>	ΔT <sub>c</sub>
072576	LAS	97	73	24	38	1.25	65	32
072676	UCC	93	63	30	40	1.10	65	28
072876	LAS	103	88	15	44	.04	66	37
072976	UCC	94	76	18	39	.08	64	30
072976	LAS	102	69	33	70	.35	64	38
090476	LAS	104	95	9	20	.16	61	43
090776	UCC	83	69	14	29	.28	60	23
092576	UCC	74	61	13	22	.05	60	14
092576	LAS	82	78	4	12	.15	60	22
081277	UCC	99	75	24	39	.05	64	35
072878	LAS	102	78	24	45	.19	65	37
080378	LAS	112	75	37	58	.04	68	44
080678	LAS	114	80	34	55	.46	65	49
090478	DRA	96	74	22	35	.06	64	32
072380	DRA	107	74	33	63	.29	67	40
073080	DRA	<u>102</u>	<u>75</u>	<u>27</u>	<u>37</u>	<u>.01</u>	<u>67</u>	<u>35</u>
Mean	-	97.0	74.9	22.1	37.6	0.26	63.3	33.4
S.D.	-	7.6	8.6	8.8	12.1	0.39	2.8	6.6

Table 2. Data listing of maximum surface wind gust (V<sub>m</sub>) by wind-gust temperature (ΔT) category. The median V<sub>m</sub> value in each ΔT category is underlined in the V<sub>m</sub> column. The average maximum surface wind gust ( $\bar{V}_m$ ) is calculated for each ΔT category.

ΔT(°F) Category	$\bar{\Delta T}$ (°F)	N	V <sub>m</sub> (mph)	$\bar{V}_m$
8	5	2	12, 19	15.5
8-12	10	5	20, 21, <u>26</u> , 28, 31	25.2
13-17	15	7	22, 25, 29, <u>31</u> , 34, 40, 44	32.1
18-22	20	10	22, 28, 30, 30, <u>35</u> , 35, 37, 38, 39, 42	33.6
23-27	25	10	24, 35, 36, 37, <u>37</u> , 38, 39, 43, 45, 51	38.5
28-32	30	8	31, 37, 40, <u>41</u> , 41, 44, 47, 58	42.4
33-37	35	7	48, 54, 54, <u>55</u> , 58, 63, 70	57.4
37	40	0		

they occurred before and during the thunderstorm event, respectively. The moist adiabatic downdraft temperature is assumed to produce the minimum temperature ( $T_{\min}$ ) observed during the thunderstorm. The maximum temperature prior to the thunderstorm was assumed to represent the maximum possible temperature ( $T_m$ ) achieved before the arrival of the downdraft. The difference between these two temperatures ( $T_m - T_{\min}$ ) is referred to as the wind-gust temperature,  $\Delta T$ .

In Table 2, the  $V_m$  data are classified according to  $\Delta T$  categories. This table shows that  $V_m$  values are spread across a wide range of  $\Delta T$ 's. The mean  $V_m$  is 37.6 mph (16.8 m/s), the median  $V_m$  is 37 mph (16.5 m/s), and the mode is 37 mph. In addition, the  $V_m$  for each  $\Delta T$  class increases as  $\Delta T$  gets larger. Tables 1 and 2 also help establish the bounds on  $\Delta T$ . In general, we can expect  $\Delta T$  to range between 0° and 40° F (or 0° to 22°C).

#### IV. ANALYSIS

To determine if  $V_m$  is related to  $\Delta T$ , the data tabulated in Table 1 were plotted on linear graph paper. The independent variable  $\Delta T$  was plotted along the abscissa and the dependent variable  $V_m$  along the ordinate as in Figure 1. The resulting scatter diagram (Figure 1) shows that a relationship does exist between  $V_m$  and  $\Delta T$ .

Also plotted in Figure 1 are the threshold wind speeds for severe thunderstorms and for special weather advisories for the NTS. The plotted data show that 8 percent of the wind gusts were in the severe thunderstorm category and that 22 percent were equal to or greater than the threshold for an NTS wind advisory.

A linear relationship appears to exist for the data plotted in Figure 1. Consequently, the data were analyzed using a simple linear regression program. A Hewlett-Packard (HP) statistical package for use with an HP-65 calculator contains such a program (STAT 1-22A). This program was used to determine the line of best fit to the data. This line is plotted in Figure 1 according to the derived expression,

$$V_m = 13.24 + 1.1\Delta T \quad (1)$$

where  $V_m$  is in mph and  $\Delta T$  is in °F. In Eq. 1, 64 percent of the variance of  $V_m$  is accounted for by  $\Delta T$ . The residual mean square or the standard error of the estimate of  $V_m$  on  $\Delta T$  is 7.3 mph, the standard error of the  $V_m$  intercept (13.24 mph) is 2.85 mph, and the standard error of the slope of the line (1.1 mph/°F) is 0.12 mph/°F. Without much loss in precision, Eq. 1 can be simplified to

$$V_m = 15 + \Delta T \quad (2)$$

where the 13.24 has been rounded up to compensate for the 10 percent loss in the coefficient for  $\Delta T$ . Equation 2 provides a very simple method for estimating the average maximum surface wind gust as a function of  $\Delta T$ . Equation 2 is suggested as the first approximation to the peak surface wind gust (in mph)




Figure 1. Plot of observed  $V_m$  and  $\Delta T$  data for 49 thunderstorms that occurred over southern Nevada. The solid line is the line of best fit to these data and is given by Eq. 1. The 90 percent confidence limits are plotted as dashed lines.

from thunderstorms occurring over southern Nevada. To be valid, all the criteria in Section II must be satisfied and  $\Delta T$  must be forecast perfectly. In other words, the proposed model is a "perfect prog" model.

Point estimates of  $V_m$  are not very meaningful unless some measure of the possible error in the estimate is given. An estimate of the peak surface wind gust should be accompanied by some sort of number interval together with a measure of assurance that the true  $V_m$  lies within the interval. Confidence or prediction intervals are a useful means of providing the necessary limits on estimates of  $V_m$ . Ostle (1963, pp. 170-174) differentiates between confidence intervals and prediction intervals. Ostle adopts the concept of a confidence interval to specify limits of acceptability when predictions by the dependent variable are used to estimate the mean of a population. A prediction interval is calculated when the dependent variable is used to predict an individual value rather than the mean. Both schemes can be included under the generic concept of confidence intervals and this terminology is used here only in the generic sense because the prediction interval is actually calculated and plotted in Figure 1.

A procedure for calculating the prediction interval is described by Ostle. In general, this interval is a function of the estimated variance of predicted individual  $V_m$  values for given  $\Delta T$  values. Use is also made of the t-distribution to account for different levels of confidence. Based on the procedure outlined by Ostle, the 90 percent confidence interval is bounded by the dashed lines drawn in Figure 1. This interval tells us that we can be 90 percent confident that the observed value of an individual  $V_m$  will lie within the plotted interval. For example, if  $\Delta T = 30^\circ\text{F}$ , we can be 90 percent confident that the observed  $V_m$  will lie between 33 and 59 mph. In other words,  $V_m = 46 \pm 13$  mph with 90 percent confidence. Or, in a practical sense, we could advise that the peak surface wind gusts expected with thunderstorms would range from 35 to 60 mph.

Near the mean wind-gust temperature ( $\overline{\Delta T} = 22^\circ\text{F}$  for the data in Table 1) the prediction interval narrows. For  $\Delta T = 22^\circ\text{F}$ , the prediction interval for  $V_m$  is  $\pm 12$  mph. This result means that any prediction of an individual  $V_m$  associated with a given  $\Delta T$  will be more meaningful for those values of  $\Delta T$  near  $\overline{\Delta T}$ .

Errors in the application of Eq. 2 (and 1) can enter through imprecise estimates of the maximum temperature and from inaccurate determination of the mean mixing ratio near the ground. Equations 1 and 2 show that a  $1^\circ\text{F}$  error in  $\Delta T$  will result in an error rate of 1 mph/ $^\circ\text{F}$  in estimates of  $V_m$ .

Other regression fits to the  $V_m$  and  $\Delta T$  data were applied to determine if the unexplained variance could be reduced further. An exponential curve fit to the data yields an  $r^2$  of 0.66 for

$$V_m = 17.68e^{0.032 \Delta T} \quad (3)$$

where  $V_m$  is in mph and  $\Delta T$  is in  $^\circ\text{F}$ . This regression line is plotted in Figure 2. The main difference between Eq. 3 and Eq. 1 (or 2) is that Eq. 3




Figure 2. Exponential plot of the 49 cases on log-linear paper. The solution to Eq. 3 is plotted as the solid line.



gives larger estimates of  $V_m$  for  $\Delta T > 30^\circ\text{F}$  (see Table 3). This difference increases to 6 mph for  $\Delta T = 40^\circ\text{F}$ . Without much loss in precision, Eq. 3 can be estimated by

$$V_m = 18e^{0.032 \Delta T} \quad (4)$$

This equation is proposed as an alternative approximation to  $V_m$ .

The same sources of error are found in Eqs. 3 and 4 as in Eqs. 1 and 2; however, the error rate,  $\epsilon$ , varies exponentially in Eqs. 3 and 4. For Eq. 3,  $\epsilon = 0.56e^a$  where  $a = 0.032\Delta T$ . Consequently, for  $\Delta T < 20^\circ\text{F}$ ,  $\epsilon < 1 \text{ mph}/^\circ\text{F}$ . As  $\Delta T$  approaches  $40^\circ\text{F}$ ,  $\epsilon$  approaches  $2 \text{ mph}/^\circ\text{F}$ .

A power-curve was also fit to the  $V_m$  and  $\Delta T$  data. The resulting expression is

$$V_m = 7.1 \Delta T^{0.54} \quad (5)$$

where  $V_m$  is in mph and  $\Delta T$  is in  $^\circ\text{F}$ . Equation 5 has an  $r^2$  of 0.67 so that 67 percent of the variance of  $V_m$  is accounted for by  $\Delta T$ . This equation yields values of  $V_m$  smaller than those of Eqs. 1 through 4 for  $\Delta T > 25^\circ\text{F}$  (see Table 3).

A special analysis was conducted to help confirm the theory that the maximum surface wind gust from desert thunderstorms can be closely approximated from the difference  $T_m - T_{\min}$ . Consequently, the only cases considered were those in which the calculated difference ( $T_m - T_e = \Delta T_c$ ) was within  $\pm 5^\circ\text{F}$  of the observed  $\Delta T$ . In addition, only those cases close to the upper-air sounding site (UCC) were used. Ten cases in Table 1 satisfy these conditions. An exponential curve fit to these 10 points explains slightly more variance ( $r^2 = 0.92$ ) than linear regression ( $r^2 = 0.86$ ).<sup>\*</sup> The derived exponential expression is,

$$V_m = 13.8e^{0.037 \Delta T} \quad (6)$$

where  $V_m$  is in mph and  $\Delta T$  is in  $^\circ\text{F}$ .<sup>\*\*</sup> The estimates of  $V_m$  from this equation are similar to those from Eqs. 1 and 3 (see Table 3). Equation 6 and the related data are plotted in Figure 3. These 10 cases are probably as close to an ideal sample as possible. This exercise confirms that there is a strong connection between observed  $\Delta T$  and  $V_m$  and suggests it is possible to derive an estimate of  $T_e$  from the vertical temperature/humidity profile. Furthermore, the analysis emphasizes that the reliability of the prediction of  $V_m$  is closely tied to the accuracy with which  $\Delta T$  can be estimated.

---

<sup>\*</sup>Linear regression equation for  $N = 10$  is  $V_m = 4.0 + 1.3 \Delta T$ .

<sup>\*\*</sup>There are two other cases that satisfy the temperature criteria ( $\leq 5^\circ\text{F}$ ). These two cases are for LAS. If added to the data base,  $N = 12$ ,  $V_m = 12.6e^{0.042 \Delta T}$  and  $r^2 = 0.81$ .

Insight into the behavior of derived  $T_e$  values relative to observed  $T_{min}$  and to "observed"  $\Delta T$  values can be obtained from Table 1. In general, the tabulated data demonstrate that values of  $T_e$  derived from the UCC (or DRA) soundings are usually colder than the observed thunderstorm-related minimum temperatures ( $T_{min}$ ). Consequently, for a fixed  $T_m$ , the calculated  $\Delta T$  (listed in Table 1 under  $\Delta T_c$ ) normally will be larger than the observed  $\Delta T$ . In fact, Table 1 shows that  $\Delta T_c < \Delta T$  (or  $T_e > T_{min}$ ) for only three cases. Therefore, the proposed models (Eqs. 1, 3, and 6) may tend to predict wind speeds that are too fast. Or from a different perspective, the models may be considered to yield conservative estimates of the peak surface wind gusts from desert thunderstorms.

Table 3 summarizes values of  $V_m$  as functions of  $\Delta T$  according to the equations developed in this report. The table demonstrates that the greatest difference in the  $V_m$  predictions is 11 mph for  $\Delta T = 40^\circ F$ . In general, the  $V_m$  values differ only by 5 to 7 mph for a given  $\Delta T$  so that the prediction from one equation is probably as valid as that from any of the others.

Table 3. Tabulation of the estimates of  $V_m$  according to the regression equations developed in this report.\*

$\Delta T(^{\circ}F)$	$V_m(\text{mph})$					
	Eq. 1	Eq. 2	Eq. 3	Eq. 4	Eq. 5	Eq. 6
5	19	20	21	21	17	17
10	24	25	24	25	25	20
15	30	30	28	30	31	24
20	35	35	33	34	36	29
25	41	40	39	40	40	35
30	46	45	46	47	45	42
35	52	50	54	55	48	50
40	57	55	63	65	52	61

\*To convert from mph to m/s, multiply mph by 0.447.  
 To convert from mph to kt, multiply mph by 0.868.  
 To convert from mph to km/h, multiply mph by 1.61.




Figure 3. Exponential plot for 10 cases where the difference between observed and predicted  $\Delta T$  is  $\leq 5^{\circ}\text{F}$ . Solid line is the line of best fit as given by Eq. 6.

## V. APPLICATION

To use the equations and diagrams in this report, the wind-gust temperature ( $\Delta T$ ) must be calculated. This temperature is found by subtracting the moist adiabatic downdraft temperature ( $T_e$ ) from the predicted maximum temperature ( $T_m$ ) occurring before the thunderstorm. There are numerous methods for predicting  $T_m$  and they will not be described here except to mention that persistence is a powerful predictor in the summertime. In fact, for those days when the selected thunderstorms occurred only on the NTS,

$$T_m (\text{today}) = 6.7 + 0.92 T_m (\text{yesterday})$$

with  $r^2 = 0.90$ ,  $N = 31$  cases.

To find  $T_e$  on a thermodynamic chart, the following steps are suggested for southern Nevada with the use of the 1200 GMT sounding for DRA.

1. Estimate  $T_m$  in  $^{\circ}\text{F}$ .
2. Determine the mean mixing ratio from the surface to the 850-mb level (approximately a depth of 50 mb).
3. For use in adiabatic ascent, reduce the predicted  $T_m$  by  $3^{\circ}\text{C}$  to account for super-adiabatic conditions near the ground. Using the parcel method, lift the parcel with temperature ( $T = T_m - 3^{\circ}$ ) dry adiabatically to the CCL.
4. Read the moist-adiabatic temperature at the CCL and follow this process line to the ground. Read  $T_e$  in  $^{\circ}\text{F}$ .
5. Calculate  $T_m - T_e = \Delta T$  in  $^{\circ}\text{F}$ . Here  $T_e$  is assumed to be an accurate estimate of  $T_{\text{min}}$ .
6. Enter the abscissa in, say, Figure 1 with  $\Delta T$  and find  $V_m$  or calculate  $V_m$  using the desired equation.

As an example, predict  $T_m$  to be  $92^{\circ}\text{F}$  ( $33^{\circ}\text{C}$ ,  $P_{\text{sfc}} = 900$  mb) and let the mean mixing ratio be 10 g/kg giving a dew-point temperature of  $54^{\circ}\text{F}$ . Subtract  $3^{\circ}\text{C}$  from  $T_m$  and find the CCL to be near 700 mb so that the CCL lies on the  $22^{\circ}\text{C}$  saturation adiabat. Follow this saturation adiabat to the ground ( $P_{\text{sfc}} = 900$  mb) and find  $T_e = 65^{\circ}\text{F}$ . The value for  $T_m$  is predicted to be  $92^{\circ}\text{F}$  so that  $\Delta T = 27^{\circ}\text{F}$ . For this temperature difference, Figure 1 yields a maximum gust of approximately 43 mph. Within the confidence interval we could say the peak gusts will range from approximately 30 to 55 mph. We would be 90 percent confident that the observed  $V_m$  would lie within this speed range when  $\Delta T$  is forecast perfectly.

## VI. SUMMARY

In using the proposed schemes it is important to emphasize that Figure 1 (and 2 and 3) will only estimate the peak surface wind gust to be expected from a desert thunderstorm in the summertime. In using all three figures, the thunderstorm must pass over the forecast site, measurable precipitation must occur, and the other criterion listed in Section II must take place to attain the estimated peak speeds. If all these conditions are not met, the proposed scheme may tend to overestimate the peak surface wind gusts. It is not known if this scheme is applicable to other sites in the desert southwest or for desert sites elsewhere in the world. Perhaps the most useful aspect of the derived expressions is that they give the user an objective basis for providing forecast guidance on expected peak surface wind gusts from desert thunderstorms. The schemes may be especially useful in alerting forecasters to the potential for locally severe thunderstorms.

The reader should understand that the strength of surface outflow winds generated by thunderstorm downdrafts varies greatly. For example, Fritsch and Rodgers (1981) note that it depends upon such physical parameters as the vertical shear of the horizontal wind, midlevel intrusions of dry air, and cloud microphysical processes. This report has endeavored to draw together a physically homogeneous set of thunderstorm-generated, wind-gust data. Differences in cloud microphysics, macroscale dynamics, distance from the initial ground contact by the downdraft, and other factors will all contribute to the variable strength of observed outflow surface wind gusts. Such variability is portrayed by the scatter of the data plotted in Figures 1, 2, and 3.

A plot of only the  $\Delta T$  and  $V_m$  data used in this study is presented in Figure 4. This figure is made available for individuals who might want to test and compare the proposed schemes for their areas of concern.

## VII. ACKNOWLEDGEMENTS

My sincere thanks to H. G. Booth, N. C. Kennedy, and R. F. Quiring for their helpful comments and useful recommendations, and to the other staff members at WSNSO who supported this study. A special thanks to the meteorological technicians who helped draw together and check some of the observations.




Figure 4. Reproduction of Fig. 1 with only the plotted data.

## REFERENCES

- Air Weather Service, 1956: Severe weather forecasting. AWSM 105-37, Headquarters, Air Weather Service, MATS, U.S. Air Force, Washington, D.C., 147 pp.
- Brancato, G. N., 1942: The Meteorological Behavior and Characteristics of Thunderstorms, USWB, April 1942.
- Crisp, C. A., 1979: Training Guide for Severe Weather Forecasters, Air Force Global Weather Central, Offutt AFB, NE, 74 pp.
- Fawbush, E. J. and R. C. Miller, 1954: A basis for forecasting peak wind gusts in nonfrontal thunderstorms. Bull. Am. Meteorol. Soc., 35:14-19.
- Fritsch, J. M. and D. M. Rodgers, 1981, The Ft. Collins Hailstorm, Bull. Am. Meteorol. Soc., 62:1560-1569.
- Idso, S. B., R. S. Ingram, and J. M. Pritchard, 1972: An American Haboob, Bull. Am. Meteorol. Soc., 53:930-935.
- Jordan, H., 1945: Gust Forecast, AAF Weather Service Bulletin, 3:22-23.
- MacDonald, A., 1976: Gusty surface winds and high level thunderstorms, Western Regional Technical Attachment No. 76-14, NWS, Western Region Headquarters, Salt Lake City, Utah, 5 pp.
- Miller, R. C., 1972: Notes on analysis and severe-storm forecasting procedures of the Air Force Global Weather Central, AWSTR 200 (Rev), Air Force Global Weather Central, Offutt AFB, Nebraska.
- Ostle, B., 1963: Statistics in Research. Iowa State University Press, Ames, Iowa, 585 pp.
- Quiring, R. F., 1972: Frequency of occurrence and duration of thunderstorms and associated phenomena at Yucca Flat, Nevada. Weather Service Nuclear Support Office, Las Vegas, Nevada, 8 pp + 14 tables.
- Saucier, W. J., 1959, Principals of Meteorological Analysis, University of Chicago Press, Chicago, Illinois, 438 pp.

- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R. F. Quiring, June 1977. (PB-271-704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB-271-742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB-272-661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB-273-155/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB-281-387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB-283-080/AS)
- 129 Fire Whirls. David W. Goens, May 1978. (PB-283-866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB-286-014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB-289-916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R. G. Pappas, R. Y. Lee, B. W. Finke, October 1978. (PB289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., Jan. 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather District. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 145 On the Use of Solar Radiation and Temperature Models to Estimate the Snap Bean Maturity Date in the Willamette Valley. Earl M. Bates, August 1979. (PB80-160971)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80-155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80-160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80-174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80-220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80-196033)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins, June 1980. (PB80-225493) (Out of print.)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80-225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB81-108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. TenHarkel, July 1980. (PB81-106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81-102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81-108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81-134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81-133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81-205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81-224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J. C. Burnash and R. Larry Ferral, April 1981. (PB81-224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. Lt. Colonel Keith R. Cooley, April 1981. (PB81-225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81-23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82-230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82-127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981. (revised September 1981). (PB82-196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82-127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981.
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82-153883)
- 171 Verification of 72-Hour 500-mb Map-Type Predictions. R. F. Quiring, November 1981. (PB82-158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82-177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981.
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82-193103)
- 175 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1981. Emil B. Gunther and Staff, June 1982.



## NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

*The National Oceanic and Atmospheric Administration* was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

**PROFESSIONAL PAPERS** — Important definitive research results, major techniques, and special investigations.

**CONTRACT AND GRANT REPORTS** — Reports prepared by contractors or grantees under NOAA sponsorship.

**ATLAS** — Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

**TECHNICAL SERVICE PUBLICATIONS** — Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

**TECHNICAL REPORTS** — Journal quality with extensive details, mathematical developments, or data listings.

**TECHNICAL MEMORANDUMS** — Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



*Information on availability of NOAA publications can be obtained from:*

**ENVIRONMENTAL SCIENCE INFORMATION CENTER (D822)  
ENVIRONMENTAL DATA AND INFORMATION SERVICE  
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION  
U.S. DEPARTMENT OF COMMERCE**

**6009 Executive Boulevard  
Rockville, MD 20852**