

NOAA Technical Memorandum NWS WR-184



COLLECTION AND USE OF LIGHTNING STRIKE DATA IN THE WESTERN U.S.
DURING SUMMER 1983

Salt Lake City, Utah
February 1984

**U.S. DEPARTMENT OF
COMMERCE**

/ National Oceanic and
Atmospheric Administration

/ National Weather
Service



NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22; except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P. O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U. S. Department of Commerce, Sillis Building, 5285 Port Royal Road, Springfield, Virginia 22151. Prices vary for all paper copy; \$3.50 microfiche. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operational Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidli, April 1968 (Revised December 1983)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB-178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. Capt. Thomas D. Murphy, U.S.A.F., May 1968. (AD-673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belasky, July 1968. (PB-179084)
- 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB-179289)
- 35 Joint ESSA/FAA ARTC Radar Weather Surveillance Program. Herbert P. Benner and Devon B. Smith, December 1968 (revised June 1970). AD-681857)
- 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (PB-183055)
- 37 Disposal of Logging Residues without Damage to Air Quality. Owen P. Cramer, March 1969. (PB-183057)
- 39 Upper-Air Lows over Northwestern United States. A. L. Jacobson, April 1969. (PB-184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970's. L. W. Snellman, August 1969. (PB-185068)
- 42 Analysis of the Southern California Santa Ana of January 15-17, 1966. Barry B. Aronovitch, August 1969. (PB-185670)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB-185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB-187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB-190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J. C. Burnash, December 1969. (PB-188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB-190157)
- 49 Predicting Precipitation Type. Robert J. C. Burnash and Floyd E. Hug, March 1970. (PB-190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB-191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB-193102)
- 52 Sacramento Weather Radar Climatology. R. G. Pappas and C. M. Veliquette, July 1970. (PB-193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB-194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB-194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB-194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM-71-00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM-71-00016)

NOAA Technical Memoranda (NWS WR)

- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon. Arthur W. Fritz, October 1970. (COM-71-00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM-71-00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM-71-00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM-71-00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM-72-10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM-72-10554)
- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM-72-10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM-72-11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM-72-11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM-72-11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM-72-10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM-72-10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM-73-10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (revised March 1978). (COM-73-10251)
- 83 A Comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM-73-10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM-73-11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y. G. Lee, June 1973. (COM-73-11276)
- 89 Objective Forecast Precipitation over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, Sept. 1973. (COM-73-11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM-73-10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM-74-11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM-74-11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM-74-11555/AS)
- 96 Map Type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM-75-10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB-250-711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM-75-10719/AS)
- 99 A Study of Flash Flood Susceptibility--A Basin in Southern Arizona. Gerald Williams, August 1975. (COM-75-11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB-246-902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB-253-053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB-252-866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB-254-650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB-254-649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB-259-594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB-260-437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB-273-677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB-264-655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB-265-941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB-273-594/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB-273-676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB-268-847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R. F. Quiring, April 1977. (PB-272-831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB-268-740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB-271-290/AS)

NOAA Technical Memorandum NWS WR-184

COLLECTION AND USE OF LIGHTNING STRIKE DATA IN THE WESTERN U.S.
DURING SUMMER 1983

Glenn E. Rasch and Mark Mathewson

Scientific Services Division
National Weather Service Western Region
Salt Lake City, Utah
February 1984

UNITED STATES
DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary

National Oceanic and
Atmospheric Administration
John V. Byrne, Administrator

National Weather
Service
Richard E. Hallgren, Director



This publication has been reviewed
and is approved for publication by
Scientific Services Division,
Western Region.



Glenn E. Rasch, Chief
Scientific Services Division
Western Region Headquarters
Salt Lake City, Utah

TABLE OF CONTENTS

	<u>Page</u>
I. Introduction	1
II. ALDS System Operation	3
III. National Weather Service ALDS Processor -- 1983	5
IV. 1983 Case Studies	7
V. Concluding Remarks and Recommendations	11
VI. References	13
VII. Appendices	14

I. INTRODUCTION

Since 1977, the Bureau of Land Management (BLM) has operated an Automatic Lightning Detection system (ALDS) in the western United States. The primary use of the system has been by the BLM and other land management agencies for the detection of lightning caused wildfires. The system is designed to detect cloud-to-ground strikes only. A secondary use has been for thunderstorm forecasting by National Weather Service (NWS) forecasters. The references document some early NWS efforts to use the ALDS data. However, these efforts were hampered by the fact that the lightning strike information, in general, has not been made available to forecasters in the form of real-time displays that could be compared to other meteorological data.

During the summer of 1982, the NWS's Western Region conducted an experiment to determine (1) if the lightning strike data could be rapidly and economically distributed and displayed at NWS offices, and (2) whether or not the data were useful as diagnostic and forecasting tools. The experiment was confined to a small portion of the overall BLM system. It consisted of two of the twelve separate lightning networks operated by the BLM. These two were chosen because they were readily available to Western Region Headquarters from the BLM in Salt Lake City, Utah. Areal coverage consisted of most of Nevada, Utah, and Colorado, and parts of Idaho, Wyoming, Arizona, New Mexico, and California. The strikes were processed in real-time by a Nova 4/X minicomputer. Every half hour a one-half hour accumulation graphic was created and sent to the Western Region Automation of Field Operations and Services (AFOS) computer. The graphics were not transmitted on the AFOS regional distribution circuit and, therefore, were routinely available only to forecasters at WSFO Salt Lake City.

The graphics were found to be timely, reliable, and a useful supplement to satellite and radar observations for detecting and forecasting thunderstorms. They were especially helpful in areas where radar coverage is poor. Many such areas exist in the western United States due to wide spacing between radar sites and blockage of radar signals by mountain ranges. In such a limited experiment, it was not possible to relate significant rainfall rates to the concentration of lightning activity in a quantitative way. However, the lightning graphics did permit forecasters to determine accurately where the most active thunderstorm cells were located. They also proved more timely than either radar maps or satellite pictures. The lightning graphics were typically available to Salt Lake City forecasters within two minutes after the end of each half hour accumulation period.

Two Western Region technical attachments, demonstrating the usefulness of the 1982 lightning data, are included as attachments to this report (Appendices 1 and 2). They also serve to illustrate the display format used in 1982.

Due to the encouraging results of 1982, the experiment was expanded to include the entire BLM network for the summer of 1983. Again it was decided to use AFOS as the primary distribution and display device. Additionally, alphanumeric products, consisting of digits plotted in grid boxes to represent lightning activity level, were made available to dial-in users. This enabled non-AFOS sites to access lightning data using a variety of terminal devices.

The next section of this report describes the BLM ALDS system. The remainder of the report focuses on the 1983 experiment -- Section III describes how the data were processed and delivered to NWS forecasters; Section IV illustrates several examples of data collected. Finally, concluding remarks and recommendations for future use and improvements of the lightning strike network are made in Section V.

II. ALDS SYSTEM OPERATION

The automated Lightning Detection System is manufactured by Lightning, Location, and Protection, Inc. The ALDS system consists of two major components: direction finders and position analyzers. A typical installation has four direction finders bordering the area of interest and one position analyzer. The direction finders consist of a flat plate antenna to detect signal strength and polarity and a loop antenna to detect direction. The direction finders are best located away from all external electrical and magnetic noise sources such as in an open field. Unfortunately, the majority of sensors must be placed in a secure area such as on the top of a building. This placement can induce errors in the data.

When lightning occurs, the direction finders within a 400 kilometer radius will usually detect it. On flat land, the effective lightning detectability is 80% at 400 kilometers. In mountainous country, it appears that mountain shadows can occur. The direction finder records the direction of the stroke as well as the magnitude of its electromagnetic signal. The direction finder performs waveform matching to determine if the stroke was a cloud-to-cloud or cloud-to-ground stroke. The electrical waveforms created by the lightning stroke have varying characteristics that differ between the two types of lightning. For example, if the rise time of the waveform is slow, the odds are that the stroke was a cloud-to-cloud. If the rise time of the waveform is fast, the stroke was probably a cloud-to-ground. Secondary flash magnitudes are also measured and used to determine lightning type. If the magnitude of the second peak is less than 60% of the first, the lightning is classified as cloud-to-ground. The reliability of detecting the type of lightning decreases with distance since the waveform is smoothed by the atmosphere. For example, a cloud-to-ground stroke at 50 km, may have a 95% probability that it is detected as a cloud-to-ground stroke. The same stroke detected at 300 km, may only have an 85% probability that it is detected as a cloud-to-ground stroke.

The direction finder by itself cannot locate the lightning stroke since it can only detect direction. There are usually four direction finders per position analyzer. A lightning stroke location is computed by the received direction and magnitude from two direction finders, using triangulation methods. The accuracy of the direction finder directions is about one degree. Hence, the accuracy of the located lightning is within several kilometers if the angles are ideal. If the perceived angles are within six degrees from the baseline (direct line between the two direction finders), the lightning is located on the baseline using the ratio of the stroke magnitudes to determine distance along the baseline. If the direction finders detect

except between the hours of 15 and 20Z. These hours were avoided due to the already high volume of communication traffic. Four hour accumulations were transmitted five times daily, and a 24 hour accumulation was transmitted once a day. Examples of 30 minute and 24 hour accumulation AFOS products are presented in Figure 4. Legends on the AFOS products are interpreted as follows:

#S = Total number of strikes detected

Gs = Grid size in degrees latitude/longitude

Mc = Minimum activity level for which a contour is drawn

Ci = Contour interval

Dt = Minimum activity level for which a "+" is plotted

Mx = Minimum activity level for which a maximum value will be labeled. The maximum value is plotted with an "x" representing the location of the maximum value.

The Western Region lightning processing and dissemination program was implemented on the Nova minicomputer system on June 16, 1983. The system functioned with very few problems until the lightning network was de-activated for the winter on September 26, 1983.

Most of the problems encountered with the ALDS system had nothing to do with the Nova or its software. They were AFOS communications problems or problems with the BLM network. For example, if the AFOS system at WSFO Salt Lake City was down or went down during product transmission from the Nova, the product was missed by all AFOS stations on the loop. This occurred several times during the summer. Another problem was that occasionally a DF or PA would fail. When such a failure occurred, there was no automatic way to alert forecasters of the problem. Operators of the ALDS system at Boise were very helpful in identifying outages and notifying forecasters of such problems. However, an automated method for identifying DF and PA failures is needed.

As in 1982, no quantitative attempts were made in 1983 to correlate lightning activity levels with heavy rainfall amounts or with severe convective storms. Comments received from forecasters indicated that on several occasions the lightning data were helpful in the timely issuance of flash flood and severe thunderstorm warnings. More study is needed in this area.

IV. 1983 CASE STUDIES

The following three examples serve to illustrate the usefulness of lightning strike graphics. They also demonstrate some problems encountered with the data. The lightning graphics used in the first two examples are single strike graphics generated specifically for this study. Each plotted dot represents a strike. The graphics in these two examples are not the same as those created for AFOS transmission. The third example uses the actual operational products as they were generated for AFOS transmission.

A. August 24, 1983 - The first example illustrates a problem noted in the northern Utah area. Figure 5 shows satellite pictures for 2015 and 2045Z, August 24, 1983. The 2015Z picture is a visual image while the one for 2045Z is a combination visual and enhanced infrared. Areas where cloud temperatures are below -30 degrees C appear enhanced. These areas also depict where thunderstorms are likely to be found. Figure 6 is a plot of all lightning strikes detected by the BLM network between 2015 and 2045Z. Each strike is represented by a dot. Three hundred strikes were detected during the 30 minute period, including out-of-range strikes (i.e., strikes having less than two DFs within 400 kms). In most areas there is good correspondence between the enhanced areas on satellite imagery and where lightning strikes were detected. For example, good correspondence exists in western Colorado, western New Mexico, and eastern Arizona. The same is true in the three corners area of Idaho, Oregon, and Nevada. In these areas the lightning strike data determined which clouds on the satellite photos were thunderstorm clouds and which were not. This distinction is important for pinpointing areas of strong convection that might lead to flash flooding and severe weather, especially in areas of poor radar detection like western Colorado and eastern Utah. The lightning strike data also help detect thunderstorms embedded in other clouds, which is important to the safe operation of aircraft.

Obviously, a problem exists in northern Utah. Several strikes are plotted in western Utah and near the southeastern shore of the Great Salt Lake where the satellite photos show the atmosphere to be cloud free. The errors in northern Utah may be due in part to the fact that none of the 30 direction finders are located in or very close to northern Utah (see Figure 1). One way to improve the accuracy of strike location in northern Utah would be to add another DF in that area.

To determine more precisely what factors may have contributed to the location errors in northern Utah, the lightning data from each of the position analyzers were plotted separately in Figure 7. Solid lines on the chart indicate the effective range of the PA, assumed to be that area within 400 kms of at least two DFs. Strikes outside the lines would have been eliminated

from the operational products. The PAs that contributed erroneous strikes were in Colorado and Wyoming. the Rock Springs direction finder is common to these two PA systems. Late in the summer the Rock Springs DF was found to have a bad analog-to-digital converter that could have caused these large position errors. The problem was not corrected until September 1, 1983. Example C below illustrates a case in which lightning strikes occurred in northern Utah after the analog-to-digital converter was replaced. The remaining PAs, southern Nevada, southern Idaho, and northern Nevada, appear to be locating strikes quite accurately.

B. August 10, 1983 - The 0835Z radar chart (Figure 8) shows that the early morning hours of August 10 were convectively quite active. The 0815Z enhanced satellite picture (Figure 9) indicates the main areas of enhanced cold clouds (temperature less than -30 degrees C) to be in southern British Columbia, northeast Oregon, central Idaho, east central Utah, and northern Wyoming. The ALDS recorded 1422 cloud-to-ground strikes during the one hour period from 08 to 09Z (Figure 10). This total includes duplicate strikes and strikes outside the assumed 400 km range of the DFs.

In order to examine this situation more carefully, lightning strikes from several individual PAs were plotted separately in Figures 11-20. The heavy solid lines on these charts indicate the effective range of the PA. Strikes outside this line were eliminated from the operational products and should be ignored.

First of all, let's examine the activity in central and southern Arizona. Only one strike was observed in this area by the southern Nevada network (Figure 11) and none by the New Mexico PA (figure 12). Most of the activity in this area appears weak on radar (Figure 8) and may not have been producing much lightning. However, note also that coverage is poor in southern Arizona. This portion of the state is outside the effective range of both PAs. More DFs are needed in this area to adequately observe cloud-to-ground lightning.

In the north there are several areas of interest, and all active thunderstorm cells appear to be depicted accurately by the PAs in Montana, northern and southern Idaho, Washington, Oregon, and Wyoming Figures 13-18. The strong cell in central Idaho, with a cloud top temperature less than -55 degrees C, has a heavy concentration of strikes detected by the PAs in Wyoming, northern and southern Idaho, and Washington. The convection in northeast Oregon is nicely depicted by the strikes from the Oregon and Washington PAs. The Washington PA is also quite accurate in depicting lightning strikes in southern British Columbia, even though they were on the fringe of the coverage area. The Montana PA even shows strikes that line up remarkably

well with radar echoes in southwest Montana. The Montana PA also recorded many strikes in northeast Wyoming where the enhanced satellite picture shows a large complex of thunderstorm cells.

However, as in the first example, problems are evident in the lightning data over northern Utah and also in southwest Wyoming. The PAs responsible for the erroneous strikes are again Colorado and Wyoming (Figures 18 and 19). And, as mentioned in the previous case, the one DF common to both PAs is the one at Rock Springs, Wyoming.

Figure 20 depicts an enlargement of the graphic sent on AFOS covering the 30 minute period 0810 - 0840Z. One grid box in southwest Wyoming had 30 strikes. Several areas in northern Utah and southwest Wyoming show activity levels exceeding three strikes per grid box. (Minimum contouring level is three strikes). Based on satellite imagery and radar, these strikes are badly misplaced, and the evidence points to problems with the Rock Springs DF. As mentioned in the previous example, this DF had a bad analog-to-digital converter that was replaced September 1, 1983.

C. September 3, 1983 - This case was chosen because it occurred after the hardware problem with the Rock Springs DF was corrected and because thunderstorms were in evidence over northern Utah. A sequence of satellite pictures, lightning graphics, and radar charts are shown in Figures 21-24. The satellite photograph, lightning chart, and radar summary in each figure are at approximately the same time. The satellite pictures and radar charts are valid at 45 and 35 minutes after the hour, respectively, while the lightning charts are 30 minute accumulations. Beginning time of the sequence is about 1400Z and ending time about 1800Z.

On the lightning graphics one or two strikes per grid box are represented by a plus sign. The first contouring level occurs where three or more strikes are recorded in a box. The contouring interval is 20, so the second contouring level occurs where ever 23 or more strikes per grid box are detected. Maximum values are noted by an "X" if they are 10 or greater.

The satellite pictures are a combination of visible at low levels and enhanced infrared at high levels. The warmest contoured temperature is -30 degrees C.

In this example, there is obviously a much better relationship between radar echoes, cold cloud temperatures on satellite pictures, and lightning strike locations in northern Utah than in the two previous examples.

For example, in Figure 21 along the Utah/Wyoming border north of Evanston, the radar echoes and enhanced cloud tops

and shadow producing cumulonimbus on the satellite image are in areas where lightning strikes were detected. As the thunderstorms developed in northeast Nevada and northwest Utah, lightning strikes appeared in those areas (Figures 23 and 24). Similarly, lightning strikes were detected along and south of the Uinta Mountains of northeast Utah as convection increased and moved or developed eastward with time in that area (Figures 23-24).

However, there are also questionable areas of lightning strike data. For example, in Figure 22(a) an enhanced area of cloud tops colder than -30 degrees C appears in northwest Utah. No strikes were detected. Nothing conclusive can be said about this area, since it is possible no cloud-to-ground strikes occurred. The strikes plotted near the southeast shore of the Great Salt Lake in Figures 21(c) and 22(c) are also suspect. The airport observing station is located there, and the observer reported no cloud-to-ground strikes during the period 1410-1540Z. The observer could see CBs to the northeast and south through southeast, which is where most of the strikes were recorded. But strikes were not seen at the station.

In Figure 23(c) the southern-most strikes plotted in central Utah appear mislocated. Neither the satellite picture nor the radar chart support thunderstorm activity so far south.

This example shows that the severe mislocation errors that occurred on a regular basis prior to September 1 were no longer evident. However, it also indicates that significant location errors were still likely. The reason is not known for sure, but it is possible that the errors were due in part to the large spacing between DFs.

V. CONCLUDING REMARKS AND RECOMMENDATIONS

The experiment conducted during the summer of 1983 demonstrated that cloud-to-ground lightning strike data from the BLM network could be collected and processed effectively and efficiently at a central location and then distributed to NWS users in near real time. No noticeable negative impact was evident on the AFOS system, which was used as the primary distribution and display system. Non-AFOS sites were also able to receive data on an alphanumeric display by dialing the central computer in Salt Lake City as needed.

The lightning data proved useful to forecasters in a number of ways. They helped fill gaps in the radar network. Gaps are a significant problem in the West due to wide spacing of radars and mountain blockage of radar signals. The data enabled forecasters to distinguish thunderstorm clouds from non-thunderstorm clouds. They also enabled forecasters to identify embedded thunderstorms that may have otherwise gone undetected. They also helped forecasters to pinpoint the location and movement of very active (in terms of number of strikes) thunderstorm cells. Although a relationship was not established between the level of lightning activity and rainfall rate, several cells with high frequencies of lightning activity did produce strong radar echoes and heavy precipitation amounts.

Several problems were encountered with the data. First of all, due to the large number of overlapping networks, numerous strikes were detected by two or more networks simultaneously. The majority of these duplicate strikes were identified and eliminated, but some duplicates no doubt went undetected. Secondly, it was impossible to automatically monitor the status of the various PAs and DFs. One or more PAs and/or DFs were sometimes inoperable without the user's knowledge. Third, several areas have poor or totally inadequate coverage. Northern Utah and western Nevada appear to have poor coverage. Although strikes are detected in these areas, location errors are probably large. Southern Arizona is an area with inadequate coverage. A large portion of southern Arizona does not have the necessary two DFs within 400 kms. A fourth problem is large positioning errors when a strike occurs on or near the baseline connecting two DFs.

All of the above problems could be overcome with some modifications to the existing network and system software. One position analyzer with 30 or more direction finders rather than 12 PAs each with three or four DFs would eliminate duplicate strikes. The software could be written such that only the two DFs detecting the strongest signal would be used in the triangulation process. If the strike occurred near the baseline connecting these two DFs, a third DF could be used to more accurately locate the


strike. The software could also be modified to automatically report the operational status of the PAs and DFs. Finally, more DFs could be added to the network to improve the coverage. One is badly needed somewhere in northern Utah and another is needed in southern Arizona.

The results of the 1983 experiment are very encouraging. With improvements as described above, the lightning data could be even more useful. Also studies relating lightning strikes to heavy rainfall and severe convection should be conducted to see if the data can be used to improve NWS watch and warning programs.

VI. REFERENCES

Rea, J. A. and C. E. Fontana, 1980: "An Automatic Lightning Detection system in Northern California". NOAA Technical Memorandum NWS WR-153, 15 pp.

Rea, J. A. and T. Getchell, 1981: "Status of the Western U.S. ALDS". Western Region Technical Attachment No. 81-32.



WESTERN REGION TECHNICAL ATTACHMENT
No. 82-32
July 27, 1982

EXPERIMENTAL LIGHTNING DETECTION CHARTS HELP
LOCATE THUNDERSTORMS

The Bureau of Land Management (BLM) has been operating a network of lightning direction-finding (DF) stations in the western United States for several years (Figure 1). The DF stations are designed to detect only cloud-to-ground strokes. It is estimated that the system detects between 80 and 90% of all cloud-to-ground flashes, and that only 1-2% of all triggers are caused by cloud discharges [1]. The data from several DF stations are transmitted to a central position-analyzing computer where the location of individualized flashes are computed automatically by triangulation. The automatic lightning detection system (ALDS) is used by BLM and other agencies to aid in detection of lightning caused fires.

Last summer Western Region TA No. 81-32 gave a status report of the ALDS and discussed an example of how the system is used in finding lightning caused fires [2]. However, output from ALDS has not been routinely available to NWS weather forecasters.

This summer, as an experiment, data from two of the many analyzers are being transmitted to the Western Region Headquarters over telephone line from the BLM in Salt Lake City. One analyzer is located in Las Vegas, Nevada and the other in Grand Junction, Colorado. The two analyzers are capable of detecting cloud-to-ground lightning strikes over most of Nevada, Utah, and Colorado, as well as parts of Idaho, Wyoming, Arizona, New Mexico, and California. Each lightning strike is plotted as a dot on a U.S. map background in AFOS. A map is produced every half hour and stored as a test graphic in the Western Region AFOS system (NMCGPHT21). Each map shows the lightning strikes that occurred during the half hour previous to the time shown on the map. A half hour strike total is presented in the upper left-hand corner. Figure 2 shows an example of such a map using a 1:1 zoom ratio.

The purpose of the experiment is to determine the utility of the lightning strike data to forecasters and to determine the best way(s) to display this information. As the following example illustrates, the data looks very useful as a supplement to radar and satellite data in detecting thunderstorms.

During the afternoon of Wednesday July 14, thunderstorms developed in several different places over the Great Basin within range of the DF stations. Figures 3 through 7 show a sequence of five ALDS charts, one mile visible satellite pictures, and Western Region radar maps, each an hour apart. Although the various displays are not valid at exactly the same time (the ALDS data is actually a half hour time composite), they are close enough in time to make some interesting comparisons.

First of all, note how the convection early in the afternoon (Figure 2) is orographically induced by the mountains in central Colorado, and northern New Mexico, and in northeast Utah by the Uinta Mountains and Book Cliffs. The ALDS system detected lightning strikes in many of these areas early in the day (not shown), and in some areas, before echoes appeared on radar. A strong correlation is apparent between lightning strikes and cloudy areas on satellite pictures during early and mid-afternoon in Figures 3 and 4.

Some cloudy areas, such as the Book Cliffs in northeast Utah, do not show lightning strikes. Perhaps convection had not reached a stage where thunderstorms were occurring.

The correlation is not as good between lightning strikes and radar echoes, particularly in western Colorado (Figures 3, 4, and 5). Even the national map (Figure 8), which includes stations not available on the Western Region map, cannot account for the strikes in many locations in western Colorado. This is not surprising since the probability of detection is low in that area (Figure 9).

Thus the lightning detection data proved useful in supplementing radar by providing additional information in poor detection areas. It also supplemented satellite pictures by differentiating thunderstorm clouds from other cloud types.

Later in the afternoon convection began developing along a weak frontal boundary located in a northeast to southwest line across northwest Utah into eastern Nevada (Figures 5, 6, and 7). The first cloud-to-ground stroke was detected between 2130 and 2200Z and was plotted on a map and available in AFOS by 2202Z (Figure 5A). This confirmed that the clouds seen on satellite (Figure 5B) were thunderstorms. Echoes were not seen by radar until 2335Z, and this chart (Figure 7C) was not available to forecasters until about 00Z, nearly two hours after the lightning data was available in AFOS.

We are currently experimenting in SSD with various display formats for lightning data and will be monitoring the utility and accuracy of this data. It appears to be very useful as a supplement to radar and satellite in detecting thunderstorms. We hope this data can be made routinely available to Western Region forecasters in the near future.

References:

- [1] Krider, E.P., R. C. Nogle, A.E. Pifer, and D. L. Vance: "Lightning Direction-Finding Systems for Forest Fire Detection", Bulletin of the American Meteorological Society, Vol. 61, No. 9, September 1980, pp 980-986.
- [2] Rea, James, and Jamie Getchell: "Status of the Western U.S. ALDS", Western Region Technical Attachment No. 81-32, August 25, 1981.




FIGURE 1. THE LOCATIONS OF LIGHTNING DF STATIONS IN THE WESTERN UNITED STATES. LARGE CIRCLES SHOW STATIONS THAT ALSO HAVE POSITION-ANALYZING COMPUTERS.















FIGURE 2. LIGHTNING STRIKE PLOT FOR 30 MINUTE INTERVAL FROM 2130-2200Z, JULY 19, 1982






FIGURE 7A. LIGHTNING STRIKE PLOT, 2230-0000Z JULY 14, 1982




FIGURE 7B. ONE MILE VISIBLE SATELLITE PICTURE, 2315Z JULY 14, 1982





FIGURE 7C. RADAR MAP, 2335Z JULY 14, 1982



98R-RADAR FIGURE 8. NATIONAL RADAR COMPOSITE, 2135Z JULY 14, 1982




FIGURE 9. PROBABILITY OF DETECTION OF RAIN OR RAIN SHOWERS BY WESTERN U.S. ARTC RADAR



WESTERN REGION TECHNICAL ATTACHMENT
No. 82-37
August 10, 1982

USING THE LIGHTNING DETECTION CHART IN REAL TIME . . . TWO CASES

Steve Todd and Ken Labas
Salt Lake City, Utah WSFO

During a recent convective outbreak, the new lightning detection chart (T22) became an important real-time tool for the forecasters at WSFO Salt Lake City (see Western Region TA #82-32 for background). In one instance, potential flash flood producing cells were monitored closely with respect to movement and development. The data correlated well with satellite imagery and spotter reports. In the second, satellite imagery and radar were deficient in displaying convective development while T22 was most revealing.

On July 27, 1982, Utah was in the midst of one of the most significant general flash flood situations in several years. Precipitable water values ranged in excess of 200% of normal and a dynamic trigger was approaching from Nevada. Figure 1A is the 700-mb analysis as of 27/12Z. A Flash Flood Watch was issued statewide at 1730Z for that afternoon and evening. By 1900Z, Figure 1B, lightning was detected over many areas of central Utah. The most active cells were apparently between Beaver (BVR) and Fillmore (FIL) as noted in satellite imagery, Figure 1C. Figures 2A through 2D chronicle the rapid expansion of convection in this area during the next hour or two. Lightning strikes indicate quite well where the most active cells are located and their subsequent propagation northward towards Nephi (NPH). At 2010Z a report from Greenville, 3 miles west of BVR, indicated rainfall of 1" in 20 minutes and 2.25" in one hour. Note the correlation between this and strike locations in Figure 1B and 2A and B. Other problems developed about 2100Z between Delta (U24) and Manti (MTI), when a state highway was washed out. Compare Figure 2C.

While concentration of cloud-to-ground lightning strikes does not necessarily indicate significant rainfall, it was a good indicator of exactly where the most active cells were. In this particular case, they produced excessive rainfall. Also note that potentially dangerous rainfall rates seem well matched with cloud top temperatures which reached the -54° to -60° C threshold (see Figures 1C and 2D).

The next day, July 28, the scene shifted to the Wasatch Front during the predawn hours. Surface observation data is very sparse this time of day and the forecaster has to rely on satellite imagery, radar, and now T22 to follow convective developments. Very moist, tropical air covered northern Utah and an area of thunderstorms was occurring from the northeast shore of the Great Salt Lake into southeast Idaho. Figure 3 shows the situation a couple of hours after midnight. A well defined cyclonic circulation is shown approaching from northern Nevada (Figure 4).

The cells in northeast Utah would appear to be about stationary (in the vicinity of the col) although in actuality the flow across the area was in the process of changing from east or southeast to westerly. Figures 5-9 (from 0930Z to 1100Z) show the convection building southward along the Wasatch Front until it reached the Salt Lake airport about 0952Z. Satellite imagery does show development southward, however, the coldest tops remain well north of SLC and cells never appear to be as far south as the thunderstorms actually occurred. In a real-time environment, at 2 or 3 a.m. with no surface obser-

vations available except at HIF to north of SLC and those from the Highway Patrol, the true developmental tendency of these cells was in question and very critical to the early morning forecast.

The real monitor of this situation was T22. Figures 5 to 9 vividly show the rather explosive southward expansion of these cells towards SLC and a little beyond. The forecaster, by 0900Z, was able to determine that a threat was materializing rapidly to the north of his local forecast area and coming right at him!

Also, note the 1035Z radar plot at the height of the storm (Figure 10). The most important rain producing cells are barely evident. This was in part due to the fact that the SLC radar had its STC on and the echoes depicted are from the Rock Springs, Wyoming radar. The final result in the bucket was nothing short of record breaking -- 1.11 inches of rain fell at the Salt Lake airport in little over an hour. A new daily record was established and we were put on the threshold of setting a new all-time monthly precipitation record for July.






FIGURE 1A 700mb 12Z JULY 27, 1962

FIGURE 1B 27/1962Z

POLAR STEREOGRAPHIC PROJECTION, TRUE AT LATITUDE 90°

1946 27JL62 36E-1MB 02297 24623 SA1






FIGURE 2A 27/1930Z





FIGURE 2B 27/2000Z



27/2100Z

3 22962 SB6 0815 28 JUL 82







FIGURE 3



700mb 00Z JULY 28, 1982
SCALE 1:10,000,000

FIGURE 4



700mb 12Z JULY 28, 1982
SCALE 1:10,000,000




FIGURE 1. Bureau of Land Management ALDS network. Dots represent direction finder (DF) locations. Area inside solid line is where at least two DFs are within 400 kms.




FIGURE 2. Coverage areas for ALDS alphanumeric products.







FIGURE 4. Sample AFOS depictions of lightning strikes. (a) 30 minute accumulation on a standard U.S. background for September 4, 1983, 0440-0510Z. (b) Same as (a) except display is on a 9:1 zoom. (c) 24 hour accumulation on a standard U.S. background for June 25, 1983, 0000-2359Z. (d) Same as (c) except display is on a 9:1 zoom.

2015 24AU83 38A-1 02284 24901 SA1



2045 24AU83 38A-1C4 02283 24893 SA1



FIGURE 5. Half-mile satellite pictures. (a) Visible image, 2015Z, August 24, 1983. (b) Combination visible and enhanced infrared image, 2045Z, August 24, 1983.









FIGURE 7. Same as Figure 6 except plots are for individual position analyzers.

0815 10AUG83 38E-2HF 00921 22771 SB6




FIGURE 8. RADAR SUMMARY
0835Z August 10, 1983

0835Z 10AUG1983

RADAR SUMMARY











FIGURE 9. One-mile enhanced infrared satellite picture, 0815Z, August 10, 1983.




1645 03SE83 38A-1C4 02297 24914 SA1




(a) Same as Figure 21(a) except time is 1645Z.

1745 03SE83 38A-1C4 02282 24903 SA1



(a) Same as Figure 21(a) except time is 1745Z.


FIGURE 23.



(b) Same as Figure 21(b) except time is 1635Z.

(c) Same as Figure 21(c) except time is 1640-1710Z.

FIGURE 24



(b) Same as Figure 21(b) except time is 1735Z.

(c) Same as Figure 21(c) except time is 1710-1740Z.

- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R. F. Quiring, June 1977. (PB-271-704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB-271-742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB-272-651)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB-273-155/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB-281-387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB-283-080/AS)
- 129 Fire Whirls. David W. Goens, May 1978. (PB-283-866/AS)
- 130 Flash-Flood Procedures. Ralph C. Hatch and Gerald Williams, May 1978. (PB-286-014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB-289-916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R. G. Pappas, R. Y. Lee, B. W. Finke, October 1978. (PB289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., Jan. 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather District. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 145 On the Use of Solar Radiation and Temperature Models to Estimate the Snap Bean Maturity Date in the Willamette Valley. Earl M. Bates, August 1979. (PB80-160971)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80-155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80-160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80-174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80-220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80-196033)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins, June 1980. (PB80-225493) (Out of print.)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80-225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB81-108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. TenHarkel, July 1980. (PB81-106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81-102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81-108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81-134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81-133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81-205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81-224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J. C. Burnash and R. Larry Ferral, April 1981. (PB81-224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. Lt. Colonel Keith R. Cooley, April 1981. (PB81-225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81-23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82-230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82-127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981. (revised September 1981). (PB82-196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82-127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981.
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82-153883)
- 171 Verification of 72-Hour 500-mb Map-Type Predictions. R. F. Quiring, November 1981. (PB82-158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82-177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82-196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82-193103)
- 175 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1981. Emil B. Gunther and Staff, June 1982. (PB82-252420)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82-253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidli, April 1969 (revised March 1983).
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E. B. Gunther, June 1983.
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83-24904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteries, Vernon C. Bissel, David G. Bennett, August, 1983.
- 181 Quantitative and Spatial Distribution of Winter Precipitation Along Utah's Wasatch Front. Lawrence B. Dunn, August, 1983.
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December, 1983.
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January, 1984.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS — Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS — Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS — Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS — Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS — Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS — Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

**ENVIRONMENTAL SCIENCE INFORMATION CENTER (D822)
ENVIRONMENTAL DATA AND INFORMATION SERVICE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE**

**6009 Executive Boulevard
Rockville, MD 20852**