



NOAA Technical Memorandum NWS WR-219

**A CASE STUDY OF THE OPERATIONAL
USEFULNESS OF THE SHARP WORKSTATION
IN FORECASTING A MESOCYCLONE-INDUCED
COLD SECTOR TORNADO EVENT IN CALIFORNIA**

**John P. Monteverdi
Department of Geosciences
San Francisco State University
San Francisco, California**

March 1993

**U.S. DEPARTMENT OF
COMMERCE**

National Oceanic and
Atmospheric Administration

National Weather
Service



NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P.O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidli, April 1968 (Revised March 1986). (PB86 177672/AS). (Revised October 1991 - PB92-115062/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB 179084)
- 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. (PB 184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 1969. (PB 185066)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)
- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliuette, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 65 Climate of Sacramento, California. Tony Martini, April 1990. (Fifth Revision) (PB89 207781/AS)
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)
- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251)
- 83 A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11655/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1967. (PB87 143160/AS)
- 96 Map type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 126 Climate of San Francisco. E. Jan Null, February 1978. Revised by George T. Pericht, April 1988. (PB88 208624/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)

NOAA Technical Memorandum NWS WR-219

**A CASE STUDY OF THE OPERATIONAL
USEFULNESS OF THE SHARP WORKSTATION
IN FORECASTING A MESOCYCLONE-INDUCED
COLD SECTOR TORNADO EVENT IN CALIFORNIA**


John P. Monteverdi
Department of Geosciences
San Francisco State University
San Francisco, California

March 1993

*UNITED STATES
DEPARTMENT OF COMMERCE
Ronald H. Brown, Secretary*

*National Oceanic and
Atmospheric Administration
John A. Knauss, Under Secretary
and Administrator*

*National Weather Service
Elbert W. Friday, Jr., Assistant
Administrator for Weather Services*



This publication has been reviewed
and is approved for publication by
Scientific Services Division,
Western Region



Kenneth B. Mielke, Chief
Scientific Services Division
Salt Lake City, Utah

TABLE OF CONTENTS

I.	INTRODUCTION	1
II.	FORECASTER AWARENESS OF ROTATIONAL POTENTIAL OF CALIFORNIA THUNDERSTORMS	3
III.	CASE STUDY: SOUTHERN SACRAMENTO VALLEY TORNADOES OF DECEMBER 17, 1992	7
IV.	DISCUSSION AND CONCLUSIONS	19
V.	REFERENCES	21

TABLE OF FIGURES

Figure 1.	Station location map	2
Figure 2.	NGM analysis of 500 mb heights (dam) and absolute vorticity for 1200 UTC 17 December 1992	2
Figure 3.	Infrared satellite image for 2201 UTC (1401 LST) 17 December 1992	8
Figure 4.	Schematic diagram for synoptic "type" associated with intense "cold sector" thunderstorms in California	9
Figure 5.	Oakland sounding for:	10
	(a) 1200 UTC (0400 LST) 17 December 1992	
	(b) 0000 UTC (1600 LST) 18 December 1992	
Figure 6.	Opening display from SHARP Workstation	12
Figure 7.	1200 UTC 17 December 1992 Oakland hodograph display from SHARP Workstation	12
Figure 8.	Scatter diagram for relationship between buoyant energy (CAPE or B+) and 0-2 km AGL positive wind shear	14
Figure 9.	Subsynoptic analysis of altimeter settings for 2200 UTC 17 December 1992	14
Figure 10.	Radar summary for 2335 UTC 17 December 1992	15
Figure 11.	Infrared satellite image for 0101 UTC 18 December 1992	17
Figure 12.	Surface moisture flux convergence field for 2100 UTC 17 December 1992	18
Figure 13.	Bogus 2200 UTC sounding for Marysville area produced by SHARP Workstation	18
Figure 14.	Bogus 2200 UTC hodograph for MYV area produced by SHARP Workstation	20
Figure 15.	Opening display from SHARP Workstation after analyses of bogus 2200 UTC sounding and hodograph data for Marysville area	20

A Case Study of the Operational Usefulness of the SHARP Workstation in Forecasting a Mesocyclone-induced Cold Sector Tornado Event in California

John P. Monteverdi

*Department of Geosciences, San Francisco State University
San Francisco, California*

ABSTRACT

An illustration of the operational usefulness of the SHARP Workstation in providing supplementary guidance to forecasters in a situation in which two tornadoes occurred in California's Sacramento Valley is presented. Use of the SHARP Workstation in analyzing the initial hodograph and in producing a bogus afternoon sounding and hodograph for the Sacramento Valley indicated that buoyancy and shear were in the correct range for moderate to strong mesocyclone-induced tornadoes. Conventional wisdom would have suggested that weak funnel clouds and small hail were the chief threats in this weather pattern. However, forecasters, aware of the role of shear in inducing storm rotation and of the potential for this weather pattern to be associated with favorable buoyancy and shear parameters in certain regions of California, would have been alert to the possibility of damaging and potentially life-threatening tornadoes.

I. INTRODUCTION

Between 2200 and 2345 UTC 17 December 1992, two tornadoes occurred in the southern Sacramento Valley near Oroville (Butte County) and another at the town of Loma Rica near Marysville (Yuba County), California (see Fig. 1 for locations). Damage surveys have established that both tornadoes were of moderate intensity (F1) (personal communication, Mr. John Quadros, Warning Preparedness Meteorologist, National Weather Service Forecast Office (WSFO), Redwood City). Although the

Marysville tornado occurred in a relatively unpopulated area, the Oroville tornado passed near the center of town, causing significant property damage (personal communication, Mr. Chris Fontana, Meteorologist-in-Charge (MIC), Weather Service Office (WSO) Redding).

An earlier tornado, rated as F0, in the Oakland-San Leandro area (1725 UTC) also had been verified. Other funnel clouds were observed in the Sacramento Valley and in the San Francisco Bay




Figure 1. Station location map. Contour labeling: 1 corresponds to 1000 ft (315 m) and 5 corresponds to 5000 ft (1575 m). MYV is designator for Marysville.




Figure 2. NGM analysis of 500 mb heights (dam) and absolute vorticity ($10^{-5}s^{-1}$) for 1200 UTC 17 December 1992..

region on this day. These tornadoes and funnel clouds occurred in the midst of a month-long period in which at least twelve tornadoes took place in northern and central California (personal communication, Mr. Jack Hales, Lead Forecaster, National Severe Storms Forecast Center (NSSFC) Kansas City, Missouri).

The primary purpose of this report is to illustrate the extent to which subsidiary guidance, available to operational forecasters, would have provided invaluable "forewarning" of the tornadic activity in a weather pattern in which conventional wisdom dictated that weak funnel clouds and small hail were the chief "severe" weather threats. "Subsidiary guidance" is defined here as personal computer or AFOS-resident programs which augment the usual synoptic-scale guidance and thermodynamic diagnoses available to operational meteorologists. The reader is referred to Doswell (1987) for a thorough overview of the synoptic and thermodynamic controls on convective events and Doswell (1992) for a review of the role of interactive workstations in the modernized National Weather Service.

The weather pattern which occurred on 17 December 1992 had been identified in two previous published studies (Monteverdi et al. 1988; Braun and Monteverdi, 1991) as one associated with strong to severe thunderstorms producing funnel clouds and occasional tornadoes in northern and central California. Those studies illustrated the usefulness of quasi-geostrophic diagnostics (e.g., those produced by the "ua-program" developed by Foster (1988)) in highlighting that portion of California most likely to experience

dynamics contributing to strong convection.

The importance of destabilization and wind shear in the development of California storms with supercellular characteristics was discussed by Braun and Monteverdi (1991). While the pattern recognition alluded to above would have already directed the attention of the forecaster to the probability of strong thunderstorms over the general area, the present preliminary study indicates that analyses of the 1200 UTC Oakland (OAK) sounding, produced interactively by the **Skew-T/Hodograph Analysis and Research Programs (SHARP) Workstation** (Hart and Korotky, 1991) and the forecaster, would have provided "quantitative" evidence that the wind shear was favorable for strong storm rotation. In this case, the usefulness of the SHARP Workstation as a "nowcasting" tool is verified.

This report provides a preliminary assessment of the meteorological controls on the Butte/Yuba County events. A more complete case study of these and other northern and central California tornadoes in December 1992 will be submitted as a possible separate Technical Attachment or Memorandum.

II. FORECASTER AWARENESS OF ROTATIONAL POTENTIAL OF CALIFORNIA THUNDERSTORMS

A. Measures of Buoyancy

Operational forecasters have been utilizing various indices evaluated from sounding data to judge thunderstorm potential. As observed by Doswell (1985) and alluded to by Braun and Monteverdi (1991), unquestioning acceptance of the thunderstorm probability and severity

thresholds defined on the bases of these indices without the application of sound meteorological reasoning can lead the forecaster to bad judgments. This is especially true in the case of California cold sector convective situations which are characterized by quite different thermodynamic characteristics than those observed in the Great Plains or Midwest severe thunderstorm patterns for which such thresholds were defined.

One such index, often misused in California, is the 500 mb Lifted Index (LI), which is essentially a measure of the buoyancy as a function of the temperature deficit (excess) of the environment at 500 mb compared to that of a parcel lifted to 500 mb. In convective situations east of California, the 500 mb LI (or, even, that at 300 mb) often qualitatively "describes" the buoyancy at the level at which the convectively-driven vertical acceleration is maximum. As pointed out in Monteverdi et al. (1988) and Braun and Monteverdi (1991), a LI based upon 700 mb temperature information is more appropriate for convective situations typical of the cool season in California. Such indices may be strongly negative (unstable) while those at 500 mb are positive, possibly leading a forecaster to underestimate convective potential.

The 500 mb LI, as a measure of potential instability in California convective situations, is a very poor quantitative estimator of total buoyancy, since the index computed for other levels often has very little relation to that computed at 500 mb. On the other hand, the Convective Available Potential Energy (CAPE), or Positive Buoyancy (B+), provides a measure of instability which is vertically integrated and not biased by the arbitrary

level at which the calculation is performed. CAPE can be qualitatively estimated from an examination of the "positive" area on a sounding plotted on a Skew-T/Log P diagram. Moreover, graphics available on AFOS and output from the SHARP Workstation programs can provide the forecaster with a quantitative estimate of CAPE.

Finally, a common pitfall for forecasters is to estimate rotational potential based upon any of the measures of sounding instability mentioned above. As pointed out by Weisman and Klemp (1982), Lazarus and Droegemeier (1990), and many others, rotational potential for a storm depends upon a favorable shear profile superimposed on an unstable environment. Marginal instabilities, as estimated by CAPE, may be associated with strong to severe tornadoes if winds veer with height and, more importantly, if the 0-3 km shear vector veers with height. Thus, as observed in the documentation for the SHARP Workstation, tornadoes have been observed for B+ values as low as 200 J kg⁻¹ with appropriate shear values. Thus, while estimations of the potential for large hail may be based solely upon some sort of LI or B+, **it is not possible to judge the rotational potential of a storm without consideration of the storm-relative shear environment.**

B. Measures of Favorable Shear

The question arises of how an operational meteorologist is to judge the degree to which the shear environment is "favorable" for the development of rotating thunderstorms. It is not something quickly or easily calculated from a glance at the mandatory and significant level winds plotted on the right

of the sounding on AFOS. However, with practice, forecasters can become adept at estimating "rotation potential" from a visual examination of the observed or modified hodograph.

Many studies have shown that rotational potential of thunderstorms is a function of the degree to which horizontal vorticity may be tilted into the vertical by the updraft of the convective cell. For example, a low-level wind at nearly right angles to the wind aloft will be characterized by strong shear vorticity, which may be converted to vertical vorticity as this horizontal air stream is tilted upward in the updraft core. This vorticity (which you cannot "see") may be converted to mesoscale circulation (which you can "see") if the thunderstorm is moving in a manner favorable for the generation of rotation (discussed below).

Weisman and Klemm (1982), and many others, have shown that a low-level wind veering and increasing with height is a necessary but not a sufficient characteristic of the hodograph in situations in which rotating thunderstorms develop. The low-level wind, particularly in the layer between 0 and 2 to 3 km, must vary in such a manner that the hodograph appears "curved" in a clockwise sense. Only for situations characterized by curved hodographs is the production of horizontal vorticity maximized. Once generated, such vorticity may be converted to storm-relative rotation in the manner conceptually described above. A quantitative measure of this storm-relative rotation is the so-called storm-relative helicity (s-r helicity).

Davies-Jones et al. (1990) have shown that s-r helicity values of around $150 \text{ m}^2\text{s}^{-2}$ are often found for thunderstorms with

forming mesocyclones. (Such storms, then, are classified as "rotating thunderstorms"; i.e., supercells.) Thus, while ratios of buoyancy to shear (e.g., the Bulk Richardson Number (BRN)) might suggest rotation in thunderstorms, **storms will show precursor rotation to tornado development only in those cases in which this ratio is in the proper range (discussed below) AND s-r helicity indicates that the shear is favorable for rotation.** (It is important to note s-r helicity values only give an indication of tornado potential. Once a storm begins to rotate (i.e., becomes supercellular), the forward and rear flank downdrafts then interact to contribute to tornado formation.)

C. Judging Buoyancy and Shear in the Central Valley from Oakland Sounding

It is well known to operational forecasters that the convective risk in the Central Valley of California is not often directly indicated by an evaluation of the OAK sounding. The same thing may be true for estimating thunderstorm risk for other large valleys in the Coast Range, particularly those which have a topographic configuration similar to that of the Sacramento Valley (e.g., the so-called "Santa Rosa Plain"). The SHARP Workstation programs allow the forecaster to quickly modify the 1200 UTC sounding for observed low-level conditions in the valley or to alter temperature and wind information to estimate changes which might occur synoptically and subsynoptically by the time of peak convective risk. Since the Sacramento Valley (and, to a lesser extent, the San Joaquin Valley) have been identified as regions particularly prone to tornado occurrence (e.g., Hales,

1985 and Braun and Monteverdi, 1991), it is especially important for forecasters to be cognizant of those mesoscale conditions which might enhance the threat for severe weather in the Central Valley when pattern recognition has already indicated a likelihood of strong convection region-wide.

Monteverdi et al. (1988) and Braun and Monteverdi (1991) document two cases in which a weather pattern, similar to the one which occurred on 17 December 1992 (discussed below), was associated with a number of tornadoes (including an F2) and funnel clouds in the Sacramento and San Joaquin Valleys. This pattern is associated with an enhanced risk of small-to-moderate hail and cold sector funnels across northern and central California. In the case of the 24 September 1986 tornadoes near Redding and Chico (documented in Braun and Monteverdi, 1991), manual alterations of the OAK sounding and hodograph produced buoyancy and wind shear profiles for the valley well within the ranges expected for right-moving, rotating (supercell) thunderstorms and with mesocyclone-related tornadoes. The SHARP Workstation programs allow the forecaster familiar with the operation of the software to perform these alterations in approximately five to ten minutes.

D. The Role of Pattern Recognition in Estimating Rotation Potential

It is important for the forecaster to recognize those synoptic patterns associated with severe weather in the Central Valley. There are dangers, however, in reliance on mere pattern recognition to direct the forecaster's attention to the severe weather threat. Rotating thunderstorms can occur in any

situation in which buoyancy and shear are in favorable ranges. While certain patterns (as is the case of the one described in this study) are most prone to be characterized by the proper buoyancy to shear ratios, California tornadoes and severe thunderstorms have been observed in other synoptic configurations (e.g., Reed and Blier, 1986 and Hales, 1985). **Forecasters who understand the cooperative role of buoyancy and shear will be able to anticipate tornado potential when any pattern produces favorable wind and instability profiles.**

Moreover, mere reliance on pattern recognition cannot provide the forecaster with the guidance necessary to judge where the mesoscale focus of severe convection is likely to occur. Outmoded notions in California, that once this pattern occurs, tornadic convection "is random" and cannot be anticipated, must be discarded. There is ample evidence in the literature that this is as much NOT the case in California as it is NOT the case in portions of the country where operational meteorologists are accustomed to evaluating severe weather potential.

There is a pervasive attitude among operational meteorologists that since F0 and F1 tornadoes are classed as "weak", they do not compare with the type of tornadoes that forecasters must anticipate east of the Rockies. Such reasoning often leads to a minimization of the risk associated with such tornadoes. The fact is that most tornado events in Kansas, Oklahoma, and Texas are in the F0 to F1 range, with most Tornado Warnings issued for F0, F1, and F2 events. Moderate (F1) and even strong (F2) tornadoes occur with some frequency in California (e.g., Hales, 1985

and Braun and Monteverdi, 1991). Furthermore, even a weak (F0) tornado can cause significant damage and hazard to life. **The forecaster must be aware that despite the lower frequencies of tornado events in California (compared to those observed in the Great Plains), similar risks are associated with their occurrence.**

III. CASE STUDY: SOUTHERN SACRAMENTO VALLEY TORNADOES OF DECEMBER 17, 1992

A. Pattern Recognition

1. Synoptic-scale Controls

The synoptic pattern which occurred on December 17, 1992 is illustrated in Fig. 2, the NGM analysis of 500 mb heights/absolute vorticity for 1200 UTC. This pattern was similar to that described in Monteverdi et al. (1988), associated with funnel clouds and large hail in the San Joaquin Valley on 21 March 1987, and in Braun and Monteverdi (1991), associated with a number of tornadoes, including an F2, in the Sacramento Valley on 24 September 1986. In each of these three cases, cyclonic vorticity advection associated with an advancing jet streak seemed to play an important role in diagnosing a vertical motion field which enhanced thunderstorm development and contributed to destabilization. For the present case, the jet streak was evident (Fig. 2) by the vorticity dipole centered at 43°N, 130°W.

Satellite imagery (Fig. 3) indicated open cellular cumulus with greatest enhancement under the left front quadrant of the advancing jet streak. This pattern was quite similar to the

schematic "type" associated with strong to severe convection in northern and central California, as illustrated by Fig. 4. Armed with this knowledge, a forecaster would certainly anticipate thunderstorm activity in California which might approach severe limits and which probably would be associated with funnel clouds.

2. Sounding

The 1200 UTC 17 December 1992 OAK sounding is given in Fig. 5a. A glance at the sounding indicates fairly stable initial conditions at OAK. In fact, the initial LI was +6, indicative of a stable atmosphere. Moreover, since heating effects are minimized during the winter season, it could be expected that very little destabilization due to diurnal effects would occur and that the low-level characteristics of the sounding during the day, even in the Sacramento Valley, would remain unaltered.

The fact that strong cold advection in the middle and lower troposphere, in association with the trough (Fig. 2) advancing southeastward, would destabilize the sounding profoundly can be considered part of the "pattern recognition" portion of the forecasting decision. Figure 5b, the 0000 UTC 18 December 1992 OAK sounding, indicates that such destabilization did occur and was associated with between a 5°C to a 10°C cooling for every level from the top of the mixed layer at (920 mb) to about the 550 mb. It is interesting to note that a similar initial sounding configuration and destabilization occurred in association with cold advection during the 24 September 1986 tornado case documented for the same general area (Braun and Monteverdi, 1991).

2201 17DE92 19E-22A 00864 17072 CB3




Figure 3. Infrared satellite image for 2201 UTC (1401 LST) 17 December 1992. "O" indicates enhanced open cellular cumulus. Letters "A" and "B" refer to cloud features discussed in the text.




Figure 4. Schematic diagram showing location of features for synoptic "type" often associated with intense "cold sector" thunderstorms in California (after Monteverdi et. al. 1988).





Figure 5. OAK sounding and plotted mandatory and significant level winds.
 (a) 1200 UTC (0400 LST) 17 December 1992.
 (b) 0000 UTC (1600 LST) 18 December 1992.

3. Wind Shear Profile

The mandatory and significant level winds, plotted on the right side of Fig. 5a, show a wind veering and increasing in strength with height. As described above, such a wind profile QUALITATIVELY may indicate the proper shear for thunderstorm rotation, if the hodograph of this wind information is curved in a clockwise sense.

The forecaster would first need to decide if the favorable wind profile was likely to persist until the cold advection and vertical motion fields described above would destabilize the atmosphere enough for thunderstorms to form. A consideration of the probable destabilization of 1200 UTC sounding and the favorable wind shear combined with recognition of the synoptic pattern would have already directed the forecaster's attention to the risk of funnel clouds and strong thunderstorms in the forecast area.

Pattern recognition might also lead the forecaster to seek a focus for the threat in the Sacramento Valley. As described in Braun and Monteverdi (1991), this type of synoptic pattern is often associated with mesoscale, perhaps leeside, troughs situated in the Central Valley. In such a pattern, low-level winds at OAK will shift to northwesterly but will remain southeasterly in the Central Valley. When this occurs, the OAK hodograph will indicate unfavorable shear profile for rotation (see winds plotted on right of Fig. 5b) even when the shear profile in the Central Valley continues to be favorable for rotating thunderstorms. When such a shear profile in the Central Valley is combined with other destabilizing influences, possible

development of supercell thunderstorms should be anticipated.

B. Interactive Analysis of Sounding and Hodograph by SHARP Workstation and Forecaster

1. Unaltered Sounding and Hodograph

The SHARP Workstation can provide additional insight to the forecaster even without alteration of the original sounding and hodograph. For this particular case, the indicators of rotation potential suggested by the SHARP Workstation analyses will be stressed. The forecaster should realize that the programs may also be used to provide information on hail and strong (downburst) wind forecasts.

Figure 6 is an example of the first screen that greets the forecaster on the SHARP Workstation analysis of a sounding and hodograph. On the 1200 UTC OAK sounding, the LI of +6 is consistent with the lack of positive buoyancy ($B+ = 0$). Other indices, including the Showalter Index, the K-Index, and the Totals-Totals index, also suggest no risk of thunderstorm development. However, each of these indices has been developed to assess the risk of thunderstorms in portions of the country where the roots of the convection are in deep layers of moisture. This is often not the case in California, and forecasters are urged not to use these indices to produce a "cookbook" thunderstorm forecast.

Since positive buoyancy ($B+$) is zero, the Bulk Richardson Number (BRN) is not defined for the 1200 UTC sounding.

However, the forecaster should keep in mind that small BRNs are associated with supercell (rotating) thunderstorms, whereas larger values are associated with multi-cellular convection. BRNs less than about 2 indicate the shear is too great for the buoyancy and thunderstorms tend to "shear out". The higher the BRN the more "upright" the convection. However, too high of a BRN would imply a thunderstorm in which the downdraft would "overwhelm" the updraft; such a thunderstorm could not develop the self-sustaining nature so characteristic of supercells. **Supercells have been observed for BRNs between 2 and 45.** However, the forecaster must keep in mind that the BRN is a "bulk" measure (i.e., based upon absolute value of shear and not whether or not it has the right characteristic). A BRN in the correct range of values for rotating thunderstorms is a necessary but not a sufficient condition for supercell development.

The low-level portion of the 1200 UTC hodograph (Fig. 7), plotted by the SHARP Workstation, is curved in the clockwise sense, indicative of a wind shear conducive to storm rotation. The degree to which the low-level shear profile is favorable for storm rotation can be estimated from the 0-3 km sr-helicity, the 0-2 km positive shear, and the energy/helicity index. The 0-3 km s-r helicity is calculated by the SHARP Workstation assuming storms develop with a storm motion (shown by arrow on the hodographs) dictated by the mean wind. **Davies-Jones et al. (1990) advise that 0-3 km s-r helicities approaching 150 (m/s)^2 support mesocyclone development, $151\text{-}299 \text{ (m/s)}^2$ weak tornadoes, $300\text{-}449 \text{ (m/s)}^2$ strong tornadoes, and greater than 450 (m/s)^2 violent tornadoes.** It is

important that the forecaster note that such values should only be used to evaluate rotation potential and never should be used in isolation from other considerations which determine whether or not a storm will become tornadic. In the present case, the 0-3 km s-r helicity of 446 (m/s)^2 stands out as an indicator of strong rotation potential IF thunderstorms develop and IF buoyancy achieves values in the proper range (discussed below).

Johns et al. (1990) have shown that low-level shear associated with a wind veering and increasing with height (called positive shear) in the 0-2 km layer is a parameter most highly correlated with tornado occurrence with **values between $6 \times 10^{-3} \text{ sec}^{-1}$ to $25 \times 10^{-3} \text{ sec}^{-1}$ encompassing all of the tornado events documented in the study.** The value of $12 \times 10^{-3} \text{ sec}^{-1}$ for the 1200 UTC sounding also suggests potential for rotating thunderstorms if the shear profile persisted during the time of storm development. In fact, Fig. 8 indicates that if such a shear were associated with a B+ of around 500 J kg^{-1} , strong or violent mesocyclone-induced tornadoes may result.

Finally, the Energy/Helicity Index (EHI) is undefined for the 1200 UTC sounding because it is another measure of the ratio between buoyancy and shear. Rather than being a bulk measure (as is the BRN), it uses the 0-2 km s-r helicity rather than the absolute value of the shear. This index is still undergoing operational testing; however, **values of the EHI of around 1 indicate a tendency for rotation to support strong (F2 and F3) tornadoes.**




Figure 8.

Scatter diagram (from Johns et. al. 1990) showing the relationship between buoyant energy (CAPE or B+) and 0-2 km AGL positive wind shear for 242 strong and violent tornado cases. Solid curved line is their suggested lower bound for the buoyant energy/shear combination associated with strong or violent tornadoes.




Figure 9.

Subsynoptic analysis of altimeter settings for 2200 UTC 17 December 1992. Broad solid line is main cold front, light dashed line is post-frontal trough. Approximate location of tornadic thunderstorm at this time indicated by "T".

Radar Summary with cell movement




Figure 10. Radar summary for 2335 UTC 17 December 1992. "A" indicates front and "B" the tornadic thunderstorm.

2. Forecaster-altered Sounding and Hodograph

Pattern recognition on the part of the operational forecaster would have already established the regional risk of thunderstorm activity for this case. An evaluation of the initial sounding and hodograph and consideration of the initial s-r helicity obtained from the SHARP Workstation would definitively establish the threat of tornadoes in any portion of the forecasting area in which the hodograph and buoyancy would be favorable. Since the low-level winds often remain southeasterly in the Sacramento Valley for the duration of such a pattern and that diurnal heating effects are often much greater there than along the coast, a first "guess" should establish an enhanced risk of tornadoes for the valley. The forecaster should not dismiss the threat for other interior valleys in the forecast area, however.

Figure 9 gives the 2200 UTC 17 December subsynoptic analysis for northern and central California. Note that up-valley, southerly flow occurred ahead of a subsynoptic scale trough located in the southern Sacramento Valley even though northwesterly winds characterized the low-level flow in the San Francisco Bay region (and in the 0000 UTC OAK hodograph). The main cold front (wide solid line) had already passed through northern California and was evidenced by a rain area on radar (indicated by the letter A on Fig. 10) and a large cloud band on satellite imagery (indicated by the letter A on Fig. 3 and Fig. 11). A large thunderstorm (indicated by the letter B on Figs. 3, 10, and 11) developed in the area of surface moisture flux convergence (Fig. 12) ahead of the trough line and north of the main front. The southeastward motion of the cell is

evident from a comparison of Figs. 3, 10, and 11.

Note the prestorm moisture flux convergence field over California was characterized by two maxima, one associated with the frontal system over central California and another associated with the trough line. Maxima of moisture flux convergence indicate a mesoscale focus for destabilization and/or wind convergence (Doswell, 1985 and many others). **Moisture flux convergence occurring in association with post frontal troughs was also found to be an important feature in the pre-storm environment with the 24 September 1986 tornadoes in the Sacramento Valley** (Braun and Monteverdi, 1991). The equivalent of the field of moisture flux convergence given in Fig. 12 may be obtained from the MESOS or ADAP programs resident on AFOS.

Figure 13 gives a "hindcasted" bogus 2200 UTC sounding and Fig. 14 a bogus hodograph for Marysville (MYV) constructed on the SHARP Workstation. The bogus sounding was created by inserting the surface data for MYV at 2200 UTC and altering the temperatures of all levels above the mixed layer by the amount suggested by the 1200 UTC advection patterns (for simplicity in this case study, determined from a comparison of the 1200 UTC and 0000 UTC OAK soundings). The bogus hodograph was created from the 1200 UTC OAK hodograph by substitution of the 2200 UTC surface wind at MYV and by insertion of the true storm motion obtained from Sacramento (SAC) weather radar (indicated by arrow). Figure 15 displays the opening screen from the SHARP Workstation when analyzing the sounding and hodograph information.





Figure 11. Infrared satellite image for 0101 UTC 18 December 1992. "A" indicates front and "B" the tornadic thunderstorm.

Surface Moisture convergence ($10^{-7}/s$)



INTERVAL: 0.5

Figure 12. Surface moisture flux convergence field ($10^{-7} s^{-1}$) for 2100 UTC 17 December 1992.




Figure 13. Bogus 2200 UTC sounding for MYV area produced by SHARP Workstation. Dotted line indicates temperature profile of lifted surface parcel.

The sounding was not modified for the greater cooling which would have occurred in the middle and lower middle troposphere over the Sacramento Valley. Greatest cold advection at these levels occurred over northern California, as is indicated by the brighter aspect of the cloudiness, which moved over that portion of California in Figs. 3 and 11. The temperature changes above the mixed layer evident in Fig. 13 and Fig. 5b (the unaltered 0000 UTC OAK sounding) from 1200 UTC (Fig. 5a) probably underestimate the degree to which destabilization occurred over the storm development region.

The dotted line (Fig. 13) represents the temperature of a surface lifted parcel in the Marysville area at 2200 UTC. Note that the equilibrium level lies just below 500 mb, resulting in an LI of +3 (Fig. 15). Radar echo tops (in this case, 27,000 feet), were much lower than the types of tops found with tornadic storms in the eastern portion of the United States. **It is important for the forecaster not to bias judgment based upon preconceived notions that severe thunderstorms have radar tops over an arbitrary elevation, such as 40,000 or 50,000 feet.**

The sounding analyses indicate that the atmosphere was buoyant beneath the 500 mb level. The 700 mb LI of -3 is more indicative of the fact that positive buoyancy is quite evident in this sounding. The B+ value of 552 J kg^{-1} is small, but when combined with the nature of the shear, this value may be associated with strong tornadic thunderstorms according to Fig. 8.

The modified hodograph shows the sort of low-level curvature indicative of high rotational potential. As would be

expected, s-r helicity of 454 (m/s)^2 , positive shear of $13 \times 10^{-3} \text{ sec}^{-1}$, and EHI of 1.17 would suggest that storms in the southern Sacramento Valley could be associated with tornadoes and that such tornadoes could be strong (F2 or F3).

IV. DISCUSSION AND CONCLUSIONS

This study illustrates the advantages and insights that the interactive use of the SHARP Workstation provided on the severe weather threat in California on 17 December 1992. Many of the AFOS products used by forecasters are useful in establishing the synoptic controls on a regional convective event but provide little insight to those factors which might "focus" the threat subsynoptically. An evaluation of such products in this case would lead to the recognition of the pattern as one often associated with severe weather phenomena in the state. Subtle aspects of the pattern include a post frontal subsynoptic or mesoscale low in the Central Valley which remains there until the main upper trough passes. Thus, while low-level winds may veer to northwesterly in the OAK hodograph, southeasterly flow can persist in that portion of the Central Valley east of the surface post frontal trough axis.

Previous studies have shown that such southeasterly flow is subjected to lifting as it moves northward in the Sacramento Valley and is associated with moisture advection and moisture flux convergence. In the present case, cold advection in the middle and lower middle troposphere could have been expected to lower temperatures at those levels over all of northern and central California, with the greatest changes over the northern

portions of the state. All of these factors would have acted to destabilize the sounding in the Sacramento Valley by the afternoon of the 17th and transform the sounding from one in which buoyancy was absent into one in which thunderstorms could develop.

Once the likelihood of thunderstorm development in the valley is recognized, monitoring of those rotation-generation factors which might be present must occur. The operational meteorologist would need to become familiar with those "indices" which estimate buoyancy and shear, and become adept at utilizing these indicators in assessing the rotational characteristics of developing thunderstorms. Such indicators include the Bulk Richardson Number, the Positive Shear, the Energy Helicity Index, and, most importantly, the storm-relative helicity.

The present case illustrates that the storm-relative helicity evident in the SHARP Workstation analyses of the 1200 UTC OAK sounding and hodograph, modified for conditions in the valley, was favorable for rotating thunderstorms. This storm-relative helicity was dependent upon low-level winds veering from southeasterly at the surface to west-southwesterly at the 3 km level. The keys to establishing the mesoscale focus for the tornado threat in northern and central California in this case were: (a) understanding of the role of storm-relative helicity in storm rotation; and (b) isolating the portion of the region which would retain low-level southeasterly flow (and, thus, favorable storm-relative helicity) until the main upper trough passed and all aspects of the hodograph became unfavorable.

Finally, California forecasters must remember that tornadoes are an important feature of California climatology in certain weather patterns. Once the operational meteorologist recognizes the risk for such storms, the same techniques used by forecasters in more tornado-prone portions of the country must be applied to establish a focus for the threat.

Acknowledgments. The author gratefully acknowledges the support and encouragement from Area Manager Norman Hoffmann and Deputy Meteorologist-in-Charge Roger Williams and all the forecasters of the WSFO San Francisco (Redwood City) in the joint interactions between San Francisco State University and the WSFO. Particular thanks are extended to Lead Forecasters E. Jan Null and John Plankinton and Warning Preparedness Meteorologist John Quadros who, with the author under the auspices of the COMET program, are pursuing studies of various severe and unusual weather events in California and of the integration of subsidiary guidance into the forecasting routine. The subsynoptic plots included in this report were produced by the SDM/WXP programs obtained by the Department of Geosciences from UNIDATA and meticulously installed, customized, and documented by the Department's site manager, Dr. David Dempsey. This research was partially sponsored by COMET grants S9209 and S9236.

V. REFERENCES

Braun, S.A. and J.P. Monteverdi, 1991: An analysis of a mesocyclone-induced tornado occurrence in northern California. *Wea. Forecasting*, **6**, 13-31.

Davies-Jones, R. D. Burgess, and M. Foster, 1990: Test of helicity as a tornado forecast parameter. *16th AMS Conference on Severe Local Storms*, Kananaskis Park, Alberta, 588-592.

Doswell, C.A. III, 1985: The operational meteorology of convective weather--Volume II: Storm scale analysis. 240 pp. NOAA Tech. Memo. ERL ESG-15. [Available from the National Severe Storms Lab, Norman, Oklahoma].

_____, 1987: The distinction between large-scale and mesoscale contribution to severe convection: A case study example. *Wea. Forecasting*, **2**, 17-31.

_____, 1992: Some issues on forecaster workstation design. *Wea. Forecasting*, **7**, 47-56.

Foster, M., 1988: Upper air diagnostics program for IBM-PC. NWSFO-Norman, Oklahoma. [Available from the NWS Western Region, P.O. Box 11188 Federal Building, Salt Lake City, Utah].

Hales, J., 1985: Synoptic features associated with Los Angeles tornado occurrences. *Bull. Amer. Meteor. Soc.*, **66**, 657-662.

Hart, J.A. and J. Korotky, 1991: The SHARP Workstation--A Skew-T/Hodograph analysis and research program. NOAA/NWS NWSFO Charleston. [Available from the National Weather Service Forecast Office, Charleston, West Virginia].

Johns, R.H., J.M. Davies, and P.W. Leftwich, 1990: An examination of the relationship of 0-2 km AGL "positive" wind shear to potential buoyant energy in

strong and violent tornado situations. *16th AMS Conference on Severe Local Storms*, Kananaskis Park, Alberta, 593-598.

Lazarus, S.M. and K.K. Droegemeier, 1990: The influence of helicity on the stability and morphology of numerically simulated storms. *16th AMS Conference on Severe Local Storms*, Kananaskis Park, Alberta, 269-274.

Monteverdi, J.P., S.A. Braun, and T.C. Trimble, 1988: Funnel clouds in the San Joaquin Valley, California. *Mon. Wea. Rev.*, **104**, 1289-1296.

Reed, R. and W. Blier, 1986: A case study of comma cloud development in the Eastern Pacific. *Mon. Wea. Rev.*, **114**, 1681-1695.

Weisman, M.L. and J.B. Klemp, 1982: The dependence of numerically simulated convective storms on vertical shear and buoyancy. *Mon. Wea. Rev.*, **110**, 504-520.

- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins (Retired) and Alexander R. Smith. Fifth Revision, July 1992. (PB92 220177)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82 158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82 177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 196103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidl, April 1969 (Revised December 1986). (PB87 142063/AS)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteris, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85 110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85 132694)
- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 187887AS)
- 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lussky, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lussky, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Radid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
- 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
- 199 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Lussky, April 1987. (PB87 185229/AS)
- 200 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke, September 1987. (PB88 110895/AS)
- 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson and George Clark, September 1987. (PB88 114038/AS)
- 202 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88 101935/AS)
- 203 An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern California. John P. Monteverdi and Scott A. Braun, October 1988. (PB89 121297/AS)
- 204 Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Carven Scott, November 1988. (PB89 128649/AS)
- 205 Forecast Guidelines for Fire Weather and Forecasters -- How Nighttime Humidity Affects Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS)
- 206 A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarters, Scientific Services Division, August 1989. (PB89 230633/AS)
- 207 The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, June 1990. (PB90-240268)
- 208 Meteorological Factors Contributing to the Canyon Creek Fire Blowup. September 6 and 7, 1988. David W. Goens, June 1990. (PB90-245085)
- 209 Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow Whitlatch, December 1990. (PB91-129239)
- 210 Hydrotools. Tom Egger, January 1991. (PB91-151787/AS)
- 211 A Northern Utah Soaker. Mark E. Struthwolf, February 1991. (PB91-168716)
- 212 Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 1991. (PB91-208439)
- 213 Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Mollner, July 1991. (PB91-227405/AS)
- 214 Emergency Operational Meteorological Considerations During an Accidental Release of Hazardous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424)
- 215 WeatherTools. Tom Egger, October 1991.
- 216 Creating MOS Equations for RAWS Stations Using Digital Model Data. Dennis D. Gettman, December 1991. (PB92-131473/AS)
- 217 Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 1992. (PB92-196104)
- 218 NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 1992. (PB93-146785)

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.


CONTRACT AND GRANT REPORTS---Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS--Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161