



NOAA Technical Memorandum NWS WR-222

**CONVECTIVE AND ROTATIONAL PARAMETERS
ASSOCIATED WITH THREE TORNADO EPISODES
IN NORTHERN AND CENTRAL CALIFORNIA**

**John P. Monteverdi
Department of Geosciences,
San Francisco State University,
San Francisco, California**

**John Quadros
National Weather Service Forecast Office
San Francisco, California**

September 1993

**U.S. DEPARTMENT OF
COMMERCE**

/ National Oceanic and
Atmospheric Administration

/ National Weather
Service



NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). In beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P.O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidli, April 1968 (Revised March 1986). (PB86 177672/AS). (Revised October 1991 - PB92-115062/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB 179084)
- 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. PB 184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 1969. (PB 185068)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)
- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 65 Climate of Sacramento, California. Tony Martini, April 1990. (Fifth Revision) (PB89 207781/AS)
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)

- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251)
- 83 A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS)
- 96 Map type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 126 Climate of San Francisco. E. Jan Null, February 1978. Revised by George T. Perich, April 1988. (PB88 208624/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudgel, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Hellick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)

NOAA Technical Memorandum NWS WR-222

**CONVECTIVE AND ROTATIONAL PARAMETERS
ASSOCIATED WITH THREE TORNADO EPISODES
IN NORTHERN AND CENTRAL CALIFORNIA**

John P. Monteverdi
Department of Geosciences,
San Francisco State University,
San Francisco, California

John Quadros
National Weather Service Forecast Office
San Francisco, California

September 1993

*UNITED STATES
DEPARTMENT OF COMMERCE
Ronald H. Brown, Secretary*

*National Oceanic and
Atmospheric Administration
(Vacant), Under Secretary
and Administrator*

*National Weather Service
Elbert W. Friday, Jr., Assistant
Administrator for Weather Services*



This publication has been reviewed
and is approved for publication by
Scientific Services Division,
Western Region

A handwritten signature in black ink that reads "Ken Mielke". The signature is written in a cursive, flowing style.

Kenneth B. Mielke, Chief
Scientific Services Division
Salt Lake City, Utah

LIST OF FIGURES

Figure 1	Locations of stations used in subsynoptic analyses and of 1992 tornadoes discussed in text. STS (Santa Rosa), MRY (Monterey), and Oroville-Marysville (MYV) indicate the closest observations to three tornado episodes	2
Figure 2	Schematic diagram showing location of features for synoptic "type" often associated with intense "cold sector" thunderstorms in California (after Monteverdi et al., 1988)	2
Figure 3	Layer temperature change ($^{\circ}\text{C}$) at RDD for the 12 hours ending 0000 UTC 25 September 1986	6
Figure 4	(a) Bogus RDD sounding, 0000 UTC 24 September 1986 Dashed line shows surface lifted parcel. Surface conditions indicated by arrow	6
	(b) Bogus RDD hodograph, 0000 UTC 24 September 1986 Arrow indicates true storm motion as determined from SAC weather radar.	6
Figure 5	Photograph of first Sebastopol tornado taken from Santa Rosa, approximately 8 km away, looking west-southwest. Note two vortices. Photo courtesy of Kent Porter, Santa Rosa Press Democrat	8
Figure 6	NGM analysis of 500 mb heights (dam) and absolute vorticity (10^{-5} s^{-1}) for 0000 UTC 3 December 1992	10
Figure 7	Subsynoptic analysis of altimeter settings for 0000 UTC 3 December 1992. Broad solid line is main cold front, light dashed line is post-frontal trough. Approximate location of tornadic thunderstorm at this time indicated by "T"	10

Figure 8	Advanced Very High Resolution Radiometer (AVHRR) visible image for 2225 UTC 2 December 1992.	
	Note suggestion of mesoscale vortex in the cloud mass (indicated by "T"). First tornado occurred under the southernmost portion of the vortex	11
Figure 9	Layer temperature change (°C) at STS for the 12 hours ending 0000 UTC 3 December 1992	13
Figure 10	(a) Bogus STS sounding, 0000 UTC 3 December 1992. Dashed line shows surface lifted parcel. Surface conditions indicated by arrow	13
	(b) Bogus STS hodograph, 0000 UTC 3 December 1992. Arrow indicates true storm motion as determined from tornado tracks	13
Figure 11	Analysis of 500 mb heights (dam) and temperatures (°C) for 0000 UTC 7 December 1992	13
Figure 12	AVHRR infrared image for 2317 UTC 6 December 1992. Thunderstorm cells southwest of Monterey peninsula produced waterspouts and tornadoes about 45 minutes after image time	14
Figure 13	Layer temperature change (°C) at MRY for the 12 hours ending 0000 UTC 7 December 1992	16
Figure 14	(a) Bogus MRY sounding, 0000 UTC 7 December 1992. Dashed line shows surface lifted parcel. Surface conditions indicated by arrow	16
	(b) Bogus MRY hodograph, 0000 UTC 7 December 1992. Arrow indicates true storm motion as determined from SAC weather radar	16
Figure 15	NGM analysis of 500 mb heights (dam) and absolute vorticity (10^{-5} s^{-1}) for 1200 UTC 17 December 1992	16

Figure 16	Subsynoptic analysis of altimeter settings for 2200 UTC 17 December 1992. Broad solid line is main cold front, light dashed line is post-frontal trough. Approximate location of tornadic thunderstorm at this time indicated by "T"	18
Figure 17	Surface moisture flux convergence field (10^{-7} s^{-1}) for 2100 UTC 17 December 1992. Positive values indicate moisture flux convergence	18
Figure 18	AVHRR infrared image for 2245 UTC 17 December 1992. Tornadic thunderstorm indicated by letter "T"	19
Figure 19	Layer temperature change ($^{\circ}\text{C}$) at MYV for the 12 hours ending 2200 UTC 17 December 1992	21
Figure 20	(a) Bogus MYV sounding, 2200 UTC 17 December 1992. Dashed line shows surface lifted parcel. Surface conditions indicated by arrow	21
	(b) Bogus MYV hodograph, 2200 UTC 17 December 1992. Arrow indicates true storm motion as determined from SAC weather radar	21

LIST OF TABLES

Table 1 Summary of severe weather reports associated with
 tornado cases discussed in text 3

Table 2 Convective and rotational parameters obtained from
 analyses of bogus soundings for four tornado occurrences
 in northern and central California 3

TABLE OF CONTENTS

I. INTRODUCTION	1
II. NORTHERN AND CENTRAL CALIFORNIA TORNADO "PROTOTYPE"	4
III. SELECTED TORNADO EVENTS OF DECEMBER 1992 IN NORTHERN AND CENTRAL CALIFORNIA	7
IV. DISCUSSION AND CONCLUSIONS	20
V. REFERENCES	22

Convective and Rotational Parameters Associated With Three Tornado Episodes in Northern and Central California

John P. Monteverdi

Department of Geosciences, San Francisco State University, San Francisco, California

John Quadros

National Weather Service Forecast Office, San Francisco, California

ABSTRACT

An overview of the synoptic and subsynoptic controls on three tornado episodes (seven tornadoes) in northern and central California during December 1992 is presented and compared to the "prototype" documented for the 24 September 1986 mesocyclone-induced F2 event in the Sacramento Valley. Convective and rotational parameters calculated interactively on the SHARP Workstation verified anecdotal evidence that two of the three December episodes were mesocyclone-induced. The study indicates that careful consideration of subsynoptic analyses and buoyancy and shear parameters can indicate a mesoscale focus for supercellular development in California "cold sector" thunderstorm environments.

I. INTRODUCTION

Twelve verified tornado events occurred in northern and central California during December 1992 (personal communication, Mr. Jack Hales, Lead Forecaster, National Severe Storms Forecast Center (NSSFC)). National Weather Service (NWS) field damage surveys, undertaken for the tornadoes of 2 December in the Santa Rosa area, of 6 December in the Monterey area, and of 17 December in the Oroville-Marysville areas, indicated that each event was characterized by multiple touchdowns of F1 (moderate) tornadoes (Table 1). Other reports of funnel clouds, large hail, and unconfirmed tornadoes or waterspouts also occurred on these days. The locations of these and other associated severe weather events discussed in the text are shown in Fig. 1.

Most California tornadoes occur in a cold sector environment which, until recently, had been

thought to be characterized only by non-rotating thunderstorms (see, e.g., Cooley, 1978 and Halvorson, 1971). Hales (1985) first suggested that the interaction of topographic factors in the Los Angeles Basin with flow patterns in certain cold sector weather types might create an environment favorable for supercellular convection. Braun and Monteverdi (1991) documented a mesocyclone-induced F2 tornado in the Sacramento Valley which occurred in a cold sector environment in which a favorable shear profile was created by topographic channelling of the low-level flow.

It is clear that the foci for the "typical" cold sector funnel cloud and very weak (F0) tornado events may be difficult to isolate operationally. It is equally clear that many, perhaps most, of the stronger (F1 and F2) events in California are mesocyclone-induced and associated with synoptic and subsynoptic focusing mechanisms, which may be resolved in an operational setting.




Figure 1 Locations of stations used in subsynoptic analyses and of 1992 tornadoes discussed in text. map. STS (Santa Rosa), MRY (Monterey), and Oroville-Marysville (MYV) indicate the closest observation sites to three tornado episodes.




Figure 2 Schematic diagram showing location of features for synoptic "type" often associated with intense "cold sector" thunderstorms in California (after Monteverdi et al., 1988).

Location of Tornado	Date of Tornado	Tornado Intensity	Path Length (miles)	Other Unconfirmed Tornado, Funnel or Waterspout in Area	Hail and/or Wind Reports	Photo or Observation of Tornado
Vina	9/24/86	F2	15	2 Tornadoes, Many Funnel	Golf ball, 60 mph	Yes
Sebastopol 1	12/2/92	F1	7.5	Wall cloud	1/2" hail	Yes
Sebastopol 2	12/2/92	F1	3	None	None	No
Windsor	12/2/92	F1	1	Wall cloud	None	No
Carmel	12/6/92	F1	7	Waterspout	None	Yes
				Funnel clouds	None	
Monterey	12/6/92	F1	1	None	None	No
Oroville	12/17/92	F1	4	Funnel Clouds	Golf ball	No
				F0 Tornado OAK		Yes
Loma Rica	12/17/92	F1	5	Funnel Clouds		Yes

Table 1. Summary of severe weather reports associated with tornado cases discussed in text.

(Sources: USDC Storm Data; John Quadros, Warning Preparedness Meteorologist, WSFO, San Francisco; Chris Fontana, WSO, Redding)

Location of Bogus Sounding	Date of Tornado	Location of Tornado(es)	Tornado Intensity	Meso-cyclone Induced?	500 mb LI (C)	700 mb LI (C)	B+ (J/kg)	Positive Shear (0-2 km) (X 10 ⁻³ s ⁻¹)	Storm Relative Helicity (m/s) ²	Energy Helicity Index	Bulk Richardson Number
Redding	9/24/86	Vina, Chico	F2	Yes	-3	-5	1806	9.7	342	3.2	15
Santa Rosa	12/2/92	Sebastopol, Windsor	F1,F1,F1	Yes	-2	-3	546	9.4	284	0.98	7
Monterey	12/6/92	Carmel, Monterey	F1,F1	No	-1	-3	446	8.6	254	0.6	5
Marysville	12/17/92	Oroville, Loma Rica	F1,F1	Yes	3	-3.5	552	12.5	454	1.17	3

Table 2. Convective and Rotational Parameters Obtained from Analyses of Bogus Soundings for Four Tornado Occurrences in North-Central California

The same general techniques in use by severe weather forecasters in other parts of the country may be utilized in California to establish the threat of strong to severe thunderstorms and tornadoes, and to determine the factors which would localize the threat subsynoptically.

Supercell thunderstorms have been shown to be associated with the majority of moderate, strong, and severe tornado events in the United States (Davies-Jones, 1986 and many others). Many studies have shown that the synoptic and mesoscale factors creating a favorable buoyancy and shear environment for supercellular convection can be diagnosed operationally (Doswell, 1985; Doswell, 1987; Johns and Doswell, 1992; and many others). The key element in anticipation of such tornado-producing thunderstorms is forecaster awareness of the role of shear in inducing storm rotation and of the potential for certain weather patterns to be associated with favorable buoyancy and shear parameters. The recent spate of tornadoes in California underscores that tornado forecasting is also an important part of the operational problem in certain California weather patterns.

A previous report (Monteverdi, 1993) described the operational usefulness of the Skew T/Hodograph Analysis and Research Programs (SHARP) Workstation (Hart and Korotky, 1991) in assessing the thermodynamic and wind shear conditions in the Sacramento Valley conducive to supercellular-type convection for the 17 December 1992 event in the Oroville area. The study indicated that a focus for the tornadic activity could have been judged by forecasters: (i) alert to the severe weather potential of the synoptic pattern on that day; and, (ii) able to evaluate information available from interactive sounding and hodograph analyses performed with the SHARP Workstation program.

The purpose for this report is to provide general documentation for those December tornado events in north-central California which occurred in a region best represented by the Oakland (OAK) radiosonde site. Convective and rotational parameters calculated for the December events will be compared to those

summarized for the 24 September 1986 F2-tornado-producing thunderstorm (Braun and Monteverdi, 1991) in the Sacramento Valley. Finally, a brief discussion of the subsynoptic or local factors which focused the threat will be discussed for each case.

II. NORTHERN AND CENTRAL CALIFORNIA TORNADO "PROTOTYPE"

Two "weather types" associated with "cold sector" tornadoes in California have been documented. Reed and Blier (1986) and Hales (1985) have discussed cases in which tornadoes in central and southern California occurred in association with cutoff mid- and upper-tropospheric troughs. Northern California tornadoes can occur in similar patterns, but most frequently occur in progressive situations similar that shown schematically in Fig. 2.

In a typical sequence, a moderate to strong surface disturbance passes through northern and central California. This disturbance is typically associated with a mid- and upper-tropospheric short-wave trough moving southeastward along the upstream side of a long-wave trough. The short-wave trough is often negatively-tilted and associated with moderate to strong mid-tropospheric cyclonic vorticity advection (CVA) and strong mid-tropospheric cold advection. As pointed out by Doswell (1987), mid-tropospheric CVA (often termed "dynamics") diagnoses upper-tropospheric divergence and an upward vertical motion field which encourages convection. Such synoptic-scale lifting of a layer destabilizes the atmosphere and changes environmental lapse rates such that the Level of Free Convection (LFC) is lowered, effectively increasing the positive buoyancy of lifted parcels.

It is important to remember, however, that differential vorticity advection approximated by CVA patterns comprises only a portion of the quasi-geostrophic forcing for vertical motions. The shape and sign of the temperature advection field also contributes to vertical motion. As a first approximation, operational forecasters can assess the combined effects of both terms by

examining CVA by the thermal wind as determined by an overlay of the 700 mb vorticity field on the 1000-500 mb thickness pattern (to determine the sign and relative magnitude of the vertical velocity at the 700 mb level). More accurate assessment of the quasi-geostrophic forcing for vertical motion can be obtained by examining the Q-vector divergence field, as computed and displayed by the PCGRIDS (Petersen, 1992) or by the "UA" programs (Foster, 1988) resident on PC-workstations at most Weather Service Forecast Offices (WSFOs).

California operational forecasters know that the pattern depicted in Fig. 2 is often associated with moderate to strong cold advection in the lower and middle troposphere. The cold advection near the surface is mitigated by sensible heating of southeastward moving air streams by the Pacific and by diurnal heating over the continent. The net effect of these processes is to destabilize the air mass over California in the lowest two-thirds of the troposphere. Because of this, the 700 mb Lifted Index (LI) is a better "indicator" of buoyancy than the 500 mb LI. In addition, cold sector thunderstorms are often associated with relatively low tropopauses and equilibrium levels even though the lower atmosphere may be quite unstable. The very strong mid-tropospheric cooling (Fig. 3), which occurred in the hours preceding the Vina tornado (Braun and Monteverdi, 1991), is representative of the marked destabilization which occurs in these patterns.

Another important feature common to many of the recent California tornado and funnel cloud episodes (e.g., Monteverdi et al., 1988; Braun and Monteverdi, 1991; and Monteverdi, 1993) is the presence of a jet streak on the southwestern periphery of the advancing short-wave trough. The upward mid-tropospheric vertical motion found east of synoptic-scale troughs tends to be augmented when the divergent front left quadrant of an advancing upper-tropospheric jet streak approaches and passes to the east of the trough axis (as documented by Uccellini and Kocin, 1988; Meier, 1993; and others). Rapid destabilization is often evident in those portions

of California which lay north of the jet axis and east of the main mid-tropospheric trough axis in patterns similar to that shown in Fig. 2.

Tornadic convection occurs most often in the "cold" air north or northwest of the main cold front and, occasionally, along the cold front itself. This cold air is often marked by "open cellular" cumulus on satellite images in the Pacific before making landfall. Recent case studies (Braun and Monteverdi, 1991; and Monteverdi, 1993) have shown that post-frontal mesoscale troughs, or low pressure areas, often develop in the Central Valley of California behind the surface cold front under the area of synoptically-forced vertical motion associated with the main mid- and upper-tropospheric trough. The eastern portions of such surface features are characterized by southerly or southeasterly upvalley flow and can be a focus for significant moisture flux convergence. It is not certain that jet streak-induced circulations have a role in the production of such surface troughs. However, the intersection of the left front quadrant of the jet streak with the surface trough line was the site for thunderstorm initiation in the 1986 F2 Vina Tornado case (Braun and Monteverdi, 1991).

The prestorm sounding (Fig. 4a) and hodograph (Fig. 4b) for the mesocyclone-induced tornado of 24 September 1986 at Vina can be considered as a "prototype" sounding and hodograph for the purposes of this study. The sounding was constructed on the SHARP Workstation by insertion of the Redding (RDD) surface temperature and wind information into the OAK 0000 UTC [approximate time of thunderstorm initiation as explained in Braun and Monteverdi (1991) radiosonde data and by assuming the wind veered smoothly between the surface and the top of the coastal mountains at 1500 m (5000 ft)].

Important buoyancy and rotational parameters for the sounding/hodograph (Figs. 4a and 4b) are summarized in Table 2. Similar information for the December 1992 events considered in this study is also provided and will be discussed in this Technical Memorandum. The reader is referred to Weisman and Klemp (1982) for a




Figure 3 Layer temperature change (°C) at RDD for the 12 hours ending 0000 UTC 25 September 1986.




Figure 4 (a) Bogus RDD sounding, 0000 UTC 24 September 1986. Dashed line shows surface lifted parcel. Surface conditions indicated by arrow.
 (b) Bogus RDD hodograph, 0000 UTC 24 September 1986. Arrow indicates true storm motion as determined from SAC weather radar.

discussion of the Bulk Richardson Number (BRN), Davies-Jones et al. (1990) for an overview of the significance of the storm-relative helicity (s-r helicity) in the development of thunderstorm rotation, Johns et al. (1990) for discussion of the relationship of the curvature of the low-level hodograph to storm rotation (positive shear), and Hart and Korotky (1991) for a brief discussion of the significance of all of the various parameters (including the Energy/Helicity Index -- EHI) displayed in the SHARP Workstation output.

To facilitate interpretation of the Table 2 and the soundings and hodographs provided in this study, the reader is reminded that supercell thunderstorms have been observed for BRNs between 2 and 45. However, the forecaster must keep in mind that the BRN is a "bulk" measure (i.e., based upon absolute value of shear and not whether it has the right characteristic). A BRN in the correct range of values for rotating thunderstorms and associated with favorable s-r helicities is a necessary, but not sufficient, condition for supercell development. Davies-Jones et al. (1990) advise that 0-3 km s-r helicities approaching $150 \text{ m}^2 \text{ s}^{-2}$ support mesocyclone development, 151-299 $\text{m}^2 \text{ s}^{-2}$ support weak tornadoes, 300-449 $\text{m}^2 \text{ s}^{-2}$ support strong tornadoes, and greater than 450 $\text{m}^2 \text{ s}^{-2}$ support violent tornadoes. In the case of the information for the Vina tornado given in Table 2, the BRN of 15 combined with the s-r helicity of $342 \text{ m}^2 \text{ s}^{-2}$ certainly would suggest a threat for strong (F2 or F3) tornadoes.

Johns et al. (1990) have shown that low-level shear, associated with a wind veering and increasing with height (positive shear), in the 0-2 km layer is a parameter most highly correlated with tornado occurrence. Particularly their results indicate values between $6 \times 10^{-3} \text{ s}^{-1}$ to $25 \times 10^{-3} \text{ s}^{-1}$ encompassed all of the tornado events. Finally, the EHI is another measure of the ratio of the buoyancy and shear, using the 0-2 km s-r helicity rather than the absolute value of the shear (as does the BRN). This index is still undergoing operational testing, however, values of the EHI of around 1 indicate a tendency for rotation to support strong (F2 and F3) tornadoes.

The positive shear value of $9.7 \times 10^{-3} \text{ s}^{-1}$ for the hodograph strongly suggests potential for rotating thunderstorms when combined with the buoyancy (B+) of 1806 J kg^{-1} . In fact, Johns et al. (1990) show that such shear can be associated with F2 and F3 tornadoes with a B+ of only around 500 J kg^{-1} . The EHI of 3.2 is also suggestive of high risk for strong and violent tornadoes.

III. SELECTED TORNADO EVENTS OF DECEMBER 1992 IN NORTHERN AND CENTRAL CALIFORNIA

A. General Overview

The tornado events of 2 December, 6 December, and 17 December 1992 in northern and central California occurred "synoptically near" the OAK radiosonde site. "Synoptically near" is defined here as less than half the distance to the neighboring Medford and Winnemucca radiosonde observations. The authors make the assumption that the OAK radiosonde data, modified for the low-level temperature and wind conditions at the stations nearest the tornado occurrences, are representative.

A summary of the severe weather reports for each of the tornadoes observed in the events, and for the 1986 Vina tornado, is given in Table 1. Although damage survey teams have investigated tornado sites in California before, the information for the Santa Rosa and Monterey tornadoes was obtained from the first two intensive ground surveys ever undertaken from the San Francisco WSFO (personal communication, Mr. Roger Williams, Deputy Meteorologist in Charge, WSFO, San Francisco). The Oroville tornado was investigated by a team from the Redding Weather Service Office (WSO) and the Loma Rica (near Marysville--MYV) tornado by a team from the Sacramento WSO.

The authors believe that heightened awareness to the risk of severe weather in California on the part of the WSFOs and WSOs will lead to more complete damage surveys. For example, it is interesting to note that NSSFC data for the period 1950-1988 indicate the mean path length




Figure 5

Photograph of first Sebastopol tornado taken from Santa Rosa, approximately 8 km away, looking west-southwest. Note two vortices. Photo courtesy of Kent Porter, Santa Rosa Press Democrat.

for tornadoes in California during that period to have been about 1 mile. The mean path length for the tornadoes considered in this study was about 4 miles with two verified tracks of 7 miles or greater. This suggests that California tornadoes have path lengths which are on average about the same as those observed in other parts of the country.

Post-event examinations of the data suggest that each event was associated with multiple touchdowns of tornadoes from the same thunderstorm. In addition, although all of the tornadoes were classified as F1, the authors concluded that the damage associated with the first Sebastopol tornado suggested winds approaching (but not quite the same as) those observed with F2 tornadoes. These observations contradict the conventional wisdom that cold sector funnels most often only touch down briefly, singly, and are very weak.

B. Sebastopol/Windsor Tornadoes of 2 December 1992

i. Description

On 2 December 1992, a number of tornadoes were reported in the Santa Rosa (STS) area (see Fig. 1 and Table 1). Witness reports of a quasi-stationary lowered base were verified by a number of photographs and video images, which substantiated the lowering was a wall-cloud that remained stationary for around one-half hour. It is interesting to note that other thunderstorms in northern California moved northeastward at 20 to 25 knots at the time of the tornado sightings and quasi-stationary wall-cloud observations in the Santa Rosa area.

At approximately 2300 UTC, a large cone-shaped tornado descended from the lowered base. The photograph (Fig. 5) was taken from Santa Rosa looking southwest when the forming tornado was around 5 miles distant. There were no other thunderstorms in the vicinity, and the region southwest of the descending tornado was in sunlight.

The tornado (Fig. 5) was the first of three in the area and was documented with a path length of 7.5 miles. As it moved slowly north-northeastward, it dissipated to be replaced by a second tornado with a track slightly east of the first. This tornado had a path length of 3 miles before dissipating. At this time, video images of the wall cloud were captured by television cameramen. It slowly moved northeastward towards Windsor, at which time the third and final tornado in this sequence descended.

Damage surveys and reports indicated that the parent thunderstorm moved northeastward at 9 knots while other thunderstorms in the region were moving 20 to 25 knots. This slow movement is consistent with the quasi-stationary and quasi-steady nature of supercell thunderstorms.

ii. Synoptic and Subsynchronous-scale Controls

The 500 mb height and absolute vorticity patterns for 0000 UTC 3 December (Fig. 6) show that the mid-tropospheric trough affecting northern California was closed, suggesting that the pattern was similar to that documented by Hales (1985) for central and southern California. The history of the short-wave trough (not given) indicated that it had progressed around the north side of the long-wave ridge in the Gulf of Alaska and approached California from the northwest, as also occurred for the prototype discussed above. At the time of Fig. 6, moderate to strong mid-tropospheric CVA in southwesterly flow characterized the middle third of California.

A series of subsynoptic surface analyses indicated that a post-frontal trough had developed in the region of California northwest of the major cold front. A wave had developed on this surface trough by 0000 UTC (Fig. 7) with a meso-low in the area between Ukiah and STS. Unlike the prototype discussed above, winds at OAK had not yet veered to the northwest, undoubtedly because the mid- and upper-tropospheric short-wave was located farther off the coast in this case. Thus, all of the coastal valleys, including those in the San Francisco Bay region and the Central Valley, were situated to






Figure 6 NGM analysis of 500 mb heights (dam) and absolute vorticity ($10^{-5}s^{-1}$) for 0000 UTC 3 December 1992.



Plot of Surface Altimeter setting (in Hg) for 0Z 3 DEC 92

Figure 7 Subsynoptic analysis of altimeter settings for 0000 UTC 3 December 1992. Broad solid line is main cold front, light dashed line is post-frontal trough. Approximate location of tornadic thunderstorm at this time indicated by "T".



2 DEC. 1992 2225 Z
AVHRR CH.1 (VIS .62 microns)
1.1 KM RESOLUTION

Figure 8 Advanced Very High Resolution Radiometer (AVHRR) visible image for 2225 UTC 2 December 1992. Note suggestion of mesoscale vortex in the cloud mass (indicated by "T"). First tornado occurred under the southernmost portion of the vortex.

channel low-level flow to a southeasterly direction. The meso-low acted as a local focus in the vicinity of STS, which at the time of Fig. 7 was reporting an east wind. The "T" on Fig. 7 gives the approximate position of the second tornado at the time of the chart.

The enhanced Advanced Very High Resolution Radiometer (AVHRR) visible satellite image for 2225 UTC (Fig. 8), about the time of the first tornado touchdown, shows numerous showers and thundershowers arrayed around the mid- and upper-tropospheric circulation center west of Point Arena. The initial development in the Santa Rosa area was the southern-most in a complex of thunderstorms which extended northward and then northwestward across Cape Mendocino. A mesoscale circulation, possibly related to the meso-low near STS, is indicated by the tight, comma shaped spiral indicated by the letter "T".

iii. Buoyancy and Wind Shear Parameters

As in the case of the prototype, strong, cold advection produced profound changes in the sounding from 1200 UTC to 0000 UTC. The layer temperature changes in the mid-troposphere (Fig. 9) are even greater than those for the prototype (Fig. 3).

The bogus STS sounding (Fig. 10a) was obtained on the SHARP Workstation by insertion of the 2300 UTC STS temperature and wind information into the OAK 0000 UTC sounding. 2300 UTC was chosen because this was the time closest to initial tornado development. The bogus hodograph (Fig. 10b) was constructed by insertion of the surface wind for STS at 2300 UTC and by assuming that the narrow valley in which STS lays would prevent any directional shear until the crest at 5000 m (1500 ft). Storm motion was determined from tornado track information as summarized above.

The parameters given for the modified sounding and hodograph in Table 2 verify ground observations of storm rotation. Buoyancy and shear were in the correct range for the development of supercells and mesocyclone-

induced tornadoes. The EHI value of slightly less than 1 combined with a s-r helicity of 284 J kg^{-1} were consistent with the development of tornadoes in the F1 to F2 range.

C. Monterey Tornadoes of 6 December 1992

i. Description

Two F1 tornadoes occurred in the Monterey-Carmel area between 2300 UTC 6 December and 0000 UTC 7 December (see Fig. 1 and Table 1). Damage consisted mainly of many uprooted trees and shingle loss to roofs. There were several reports of waterspouts and funnel clouds off the Carmel coast. Observations of the Carmel tornado suggested that it occurred from the flat base of the parent thunderstorm and no observations of wall-clouds or lowered bases reported. Damage surveys indicated that the path length of the Carmel tornado was 7 miles and that of the second tornado in Monterey was 1 mile. These surveys also indicated that the tornadoes moved along with the thunderstorm motion, as observed on SAC weather radar.

ii. Synoptic and Sub-synoptic Controls

The 500 mb analysis for 0000 UTC 7 December 1992 (Fig. 11) shows a negatively-tilted trough extended from the Gulf of Alaska southeastward over California. The trough was associated with strong mid-tropospheric CVA (not shown). The axis of the upper-tropospheric jet stream intersected the California coast near Vandenberg Air Force Base. The mid-tropospheric expression of a jet streak is evident by the height contour packing on the southern periphery of the trough (Fig. 11).

At the time of the tornado reports, a cold front was in the process of advancing through the north-central portions of California. AVHRR infrared imagery for 2317 UTC (Fig. 12), about 45 minutes before the first tornado report, indicated that considerable enhancement of the frontal cloud band had occurred east of the main mid-tropospheric trough axis in the vicinity of the left front quadrant of the advancing jet streak. Strong thunderstorm development was




Figure 9 Layer temperature change (°C) at STS for the 12 hours ending 0000 UTC 3 December 1992.





Figure 10 (a) Bogus STS sounding, 0000 UTC 3 December 1992. Dashed line shows surface lifted parcel. Surface conditions indicated by arrow.
 (b) Bogus STS hodograph, 0000 UTC 3 December 1992. Arrow indicates true storm motion as determined from tornado tracks.




Figure 11 Analysis of 500 mb heights (dam) and temperatures (°C) for 0000 UTC 7 December 1992.




Figure 12

AVHRR infrared image for 2317 UTC 6 December 1992. Thunderstorm cells southwest of Monterey peninsula produced waterspouts and tornadoes about 45 minutes after image time.

evident from just east of San Francisco southwestward across the coastline near Monterey.

Since there are few observations in the Monterey area and the front lay over the data-sparse Pacific, subsynoptic analyses could not be used to provide insights to the focus for this case. However, since the surface winds in the Monterey area, the San Francisco Bay region, the southern Sacramento Valley, and the northern San Joaquin Valley were all southeasterly at this time, it appears that no evidence for a mesoscale focus would be discerned from subsynoptic analyses. In addition, since the funnel clouds and waterspouts which preceded the tornadoes were observed over the Pacific, west of Carmel, it is probable that any such focusing mechanisms would have been active west of the coastline and away from any observation sites.

iii. Buoyancy and Wind Shear Parameters

Mid-tropospheric cold advection and low-level diurnal heating produced 12 h temperature changes (Fig. 13) in the Monterey area in the same sense (but of a smaller magnitude) than those observed for the other cases considered. The destabilization for this case was not as marked as that which occurred for the other cases. In fact, the B+ (Table 2) obtained from a bogus 0000 UTC MRY sounding (Fig. 14a), constructed on the SHARP Workstation, is the smallest of the cases discussed in this study.

The bogus hodograph for MRY (Fig. 14b) was constructed by insertion of the 0000 UTC wind observation for MRY and the true storm motion as determined from SAC weather radar. The 0-2 km positive shear and 0-3 km s-r helicity, as estimated from the SHARP Workstation analysis of the bogus hodograph, were also somewhat smaller than those observed for the other cases (Table 2). The curvature of the hodograph between the surface and 2000 m was clockwise and, with the storm motion indicated, produced a s-r helicity of $254 \text{ m}^2 \text{ s}^{-2}$, a "rotational potential" which can support weak tornadoes, according to Davies-Jones et al. (1990).

Although the BRN obtained from the sounding (Fig. 14a) was in a range which suggested developing thunderstorms might have supercellular characteristics, the buoyancy and shear evident in Table 2 for this case were likely too small for mesocyclone development. This is substantiated by the EHI of 0.6, by far the smallest of such values for the cases considered in this study.

The parameters summarized in Table 2 for this case suggest that the potential for cold sector funnels and weak tornadoes region-wide was high in this case. However, no discernible mesoscale or subsynoptic focus was evident which would have aided forecasters in localizing the threat to the Monterey area. This lack of focus is probably consistent with the fact that both anecdotal evidence and buoyancy/shear parameters indicate that these tornadoes were not mesocyclone-induced.

D. Oroville-Marysville (Loma Rica) Tornadoes of 17 December 1992

i. Description

Two tornadoes occurred in the southern Sacramento Valley between 2125 and 2330 UTC 17 December 1992 (see Fig. 1 and Table 1). The first tornado passed through the town of Oroville at 2125 UTC and produced substantial damage (personal communication, Mr. Chris Fontana, Meteorologist in Charge, WSO Redding) along its path length of 3/4 mile. A previous report (Monteverdi, 1993) described the usefulness of the SHARP Workstation in providing guidance to forecasters in anticipating this event.

The second tornado occurred at 2330 UTC and passed through an unpopulated area near Loma Rica, which is a small village around 15 miles northeast of Marysville. Videotapes and eyewitness reports verified both the presence of a wall-cloud and the subsequent touchdown of the tornado outside of Loma Rica. Other reports indicate that the tornado was "cone-shaped" as it traversed its approximately 5 mile path length.




Figure 13 Layer temperature change ($^{\circ}\text{C}$) at MRY for the 12 hours ending 0000 UTC 7 December 1992.





Figure 14 (a) Bogus MRY sounding, 0000 UTC 7 December 1992. Dashed line shows surface lifted parcel. Surface conditions indicated by arrow.
 (b) Bogus MRY hodograph, 0000 UTC 7 December 1992. Arrow indicates true storm motion as determined from SAC weather r2




Figure 15 NGM analysis of 500 mb heights (dam) and absolute vorticity (10^{-5}s^{-1}) for 1200 UTC 17 December 1992.

Analyses of radar and satellite imagery confirm that both tornadoes were associated with a thunderstorm which moved southeastward through the Sacramento Valley. There were unconfirmed reports of golf-ball size hail and other funnel clouds when the storm was between Oroville and Marysville.

ii. Synoptic and Subsynoptic-scale Controls

The synoptic pattern which occurred on 17 December 1992 (Fig. 15) was the most similar of the other December 1992 cases to the prototype described earlier. CVA associated with an advancing jet streak seemed to play an important role in diagnosing a vertical motion field which enhanced thunderstorm development and contributed to destabilization. The jet streak was evident (Fig. 15) by the vorticity dipole centered at 43°N, 130°W.

Satellite imagery during the morning of 17 December showed open cellular cumulus west of the coastline in the hours before the initiation of convection in the Sacramento Valley, with greatest enhancement under the left front quadrant of the advancing jet streak. This pattern was quite similar to the schematic "type" associated with strong to severe convection in northern and central California (Fig. 2).

Figure 16 gives the 2200 UTC 17 December subsynoptic analysis for northern and central California. Note that upvalley, southerly flow was occurring ahead of a subsynoptic scale trough located in the southern Sacramento Valley even though northwesterly winds characterized the low-level flow in the San Francisco Bay region. Mesoscale or subsynoptic lee-side troughs are common in patterns which closely resemble the prototype surface and mid-tropospheric patterns (Fig. 2), discussed in section 2.

The prestorm moisture flux convergence field over California (Fig. 17) was characterized by two maxima, one associated with the frontal system over central California and another associated with the trough line. Maxima of moisture flux convergence indicate a mesoscale


focus for destabilization and/or wind convergence (Doswell, 1985 and many others). The tornadic thunderstorm (indicated by the letter T on Figs. 16 and 18) initially developed in the northern-most area of surface moisture flux convergence ahead of the trough line and north of the main front. Moisture flux convergence occurring in association with post-frontal troughs was also found to be an important feature in the prestorm environment with the 24 September 1986 tornadoes in the Sacramento Valley (Braun and Monteverdi, 1991).

iii. Buoyancy and Wind Shear Characteristics

Strong cold advection in the mid- and lower-troposphere, in association with the trough (Fig. 15) advancing southeastward, caused pronounced destabilization over northern and central California. The cooling in the 900-400 mb layer was very marked, as is evident in Fig. 19, which shows the 12 h layer temperature changes for the period from 1200 UTC 17 December and 0000 UTC 18 December. Figure 19 was obtained by substitution of the surface data for MYV into the respective OAK soundings. It is interesting to note that the development of favorable buoyancy for the three mesocyclone-induced tornado cases considered in this study was associated with similar 12 h layer temperature changes, as is evident from a comparison of Figs. 3, 9, and 20.


The bogus 2200 UTC MYV sounding (Fig. 20a) was constructed on the SHARP Workstation. The sounding information (Table 2) substantiates the fact that weak positive buoyancy characterized the lower-troposphere even though the 500 mb Lifted Index (LI) indicated negative buoyancy at that level. As pointed out in Braun and Monteverdi (1991) and Monteverdi (1993), the 700 mb LI provides a more accurate indicator of instability in cold sector California events, particularly during the cool season (-3.5 in this case).

The bogus MYV hodograph (Fig. 20b) was created from the 1200 UTC OAK hodograph by substitution of the 2200 UTC surface wind at MYV and by insertion of the true storm motion



Plot of Surface Altimeter setting (in Hg) for 22Z 17 DEC 92

Figure 16 Subsynoptic analysis of altimeter settings for 2200 UTC 17 December 1992. Broad solid line is main cold front, light dashed line is post-frontal trough. Approximate location of tornadic thunderstorm at this time indicated by "T".



INTERVAL: 0.5

Figure 17 Surface moisture flux convergence field (10^{-7} s^{-1}) for 2100 UTC 17 December 1992. Positive values indicate moisture flux convergence.




Figure 18 AVHRR infrared image for 2245 UTC 17 December 1992. Tornadic thunderstorm indicated by letter "T".

obtained from SAC weather radar (indicated by arrow). The modified hodograph shows the type of low-level curvature indicative of high rotational potential, verified by the s-r helicity of $454 \text{ m}^2 \text{ s}^{-2}$. The weak buoyancy of 552 J kg^{-1} is in the correct range to support strong and violent tornadoes if the positive shear exceeded $10 \times 10^{-3} \text{ s}^{-1}$ (Johns et al., 1990). The positive shear of $13 \times 10^{-3} \text{ s}^{-1}$ combined with the s-r helicity gives an EHI of 1.17, suggesting that storms in the southern Sacramento Valley could be associated with strong (F2 or F3) tornadoes.

IV. DISCUSSION AND CONCLUSIONS

Previous studies (Braun and Monteverdi, 1991; and Monteverdi, 1993) have pointed out that shear profiles which support s-r helicities favorable for storm rotation can be created or augmented locally by most valleys in northern and central California. Since both the Coast Range and the Sierra Nevada trend northwest-to-southeast, most ridges and valleys are oriented in the "proper" direction to channel low-level southwesterly and southerly flow to southeasterly. For the cases considered in this study, this low-level channelling combined with the west-southwest flow in the mid-troposphere, created a situation in which moderate to strong streamwise vorticity was generated in the low-level flow. As a result, in all of the cases considered in this study, s-r helicities achieved values which could support moderate to strong tornadoes.

The southern Sacramento tornado events of 17 December 1992 occurred in a synoptic and subsynoptic setting very similar to that associated with the "prototype" 24 September 1986 Vina tornado. Of the soundings and hodographs considered here, those for 17 December 1992 (Figs. 20a and 20b) most closely resembled those for the Vina tornado (Figs. 4a and 4b). In the Vina event (Braun and Monteverdi, 1991), a post-frontal subsynoptic trough focused the topographically channelled low-level flow. It appears that, in the case of the 17 December 1992 tornadoes, proper phasing of synoptic-scale features also supported the development of a subsynoptic, leeside trough in the Sacramento

Valley which, in turn, augmented the channelled flow in the same manner. The s-r helicities for these two events were the greatest of those calculated for all events considered here and actually were in a range which could support strong (24 September 1986) and strong to violent (17 December 1992) tornadoes (Table 2).

Strong and violent tornadoes have been observed for a wide range of buoyancy and shear ratios (Johns et al., 1990). While moderate buoyancy was associated with the Vina event (Table 2), only weak instability occurred for the December events. This was partially due to the fact that although all of the events considered here were "cold sector", the Vina tornado occurred in a warm season environment characterized by much higher temperatures and dew points than those that occurred with the December 1992 tornadoes. This study substantiates the conclusions of Johns et al. (1990), and many others, that weak instability can be associated with at least moderate mesocyclone-induced tornadoes if shear values are in favorable ranges. In the case of the events considered here, only the Monterey tornadoes occurred in an environment in which the buoyancy to shear ratios failed to support strong thunderstorm rotation.

Each of the three December events occurred in a synoptic pattern which corresponded to one of the two already discussed in the literature as being associated with cold-sector tornadoes in California (Hales, 1985; Braun and Monteverdi, 1991). The Monterey and southern Sacramento Valley tornadoes were associated with a progressive mid-tropospheric pattern similar to that of the "prototype" discussed earlier in this study. The Santa Rosa/Sebastopol tornadoes occurred in a similar pattern except that the mid-tropospheric trough became closed off west of the coastline. This apparently kept the greatest upper-tropospheric divergence and ascent along the coast.

This study indicates that careful subsynoptic analyses and thoughtful consideration of buoyancy and shear information, obtained interactively on the SHARP Workstation (or any similar analysis system) by the forecaster, can




Figure 19 Layer temperature change (°C) at MYV for the 12 hours ending 2200 UTC 17 December 1992.





Figure 20 (a) Bogus MYV sounding, 2200 UTC 17 December 1992. Dashed line shows surface lifted parcel. Surface conditions indicated by arrow.
 (b) Bogus MYV hodograph, 2200 UTC 17 December 1992. Arrow indicates true storm motion as determined from SAC weather radar.

indicate a mesoscale or subsynoptic focus for those tornado events in California which are mesocyclone-induced. On the other hand, determination of such a local focus may not be possible for thunderstorms which do not have supercellular characteristics and those supercellular events which are initiated over the data-sparse Pacific.

This study also suggests that buoyancy and shear parameters may yield operationally useful guidance in distinguishing between the threat for funnel clouds, or weak, moderate, and strong tornadoes in California cold sector thunderstorm events. Research is continuing at San Francisco State University (SFSU) to determine buoyancy and shear information for all of the tornado and funnel cloud events in northern and central California since 1950.

Acknowledgments.

AVHRR imagery was supplied courtesy of Mr. Steve Hipskind, Earth System Science Division, NASA Ames Research Center and was processed by Ms. Kathy Pagan, Research Technician, SFSU. The subsynoptic plots included in this report were produced by the SDM/WXP programs obtained by the Department of Geosciences from UNIDATA and meticulously installed, customized, and documented by the site manager, Dr. David Dempsey. This research was partially sponsored by COMET grants S9209 and S9236. The authors gratefully acknowledge the support and encouragement from Area Manager Norman Hoffmann, Deputy Meteorologist in Charge Roger Williams, and all the forecasters of the WSFO San Francisco (Redwood City) in the joint interactions with SFSU and the WSFO.

V. REFERENCES

Braun, S.A. and J.P. Monteverdi, 1991: An analysis of a mesocyclone-induced tornado occurrence in Northern California. *Wea. Forecasting*, **6**, 13-31.

Cooley, J., 1978: Cold air funnel clouds. *Mon. Wea. Rev.*, **106**, 1368-1372.

Davies-Jones, R.D., 1986: Tornado dynamics. *Thunderstorm Morphology and Dynamics*. E. Kessler, Ed., University of Oklahoma Press, 197-236.

Davies-Jones, R. D., Burgess and M. Foster, 1990: Test of helicity as a tornado forecast parameter. *16th AMS Conference on Severe Local Storms*, Kananaskis Park, Alberta, 588-592.

Doswell, C.A. III, 1985: The operational meteorology of convective weather--Volume II: Storm-scale analysis. 240 pp. NOAA Tech. Memo. ERL ESG-15. [Available from the National Severe Storms Lab, Norman, Oklahoma].

_____, 1987: The distinction between large-scale and mesoscale contribution to severe convection: A case study example. *Wea. Forecasting*, **2**, 17-31.

Foster, M., 1988: Upper air diagnostics program for IBM-PC. NWSFO-Norman, Oklahoma. [Available from the NWS Western Region, P.O. Box 11188 Federal Building, Salt Lake City, Utah 84147].

Hales, J., 1985: Synoptic features associated with Los Angeles tornado occurrences. *Bull. Amer. Meteor. Soc.*, **66**, 657-662.

Halvorson, D.A., 1971: Tornado and funnel clouds in San Diego County. *Western Region Technical Attachment No. 71-33*. NWS Western Region, P.O. Box 11188, Salt Lake City, Utah 84147.

Hart, J.A. and J. Korotky, 1991: The SHARP Workstation--A Skew T/Hodograph analysis and research program. NOAA/NWS WSFO Charleston. [Available from the National Weather Service Forecast Office, Charleston, West Virginia].

Johns, R.H., J.M. Davies, and P.W. Leftwich, 1990: An examination of the relationship of 0-2 km AGL "positive" wind shear to potential buoyant energy in strong and violent tornado situations. *16th AMS Conference on Severe Local Storms*, Kananaskis Park, Alberta, 593-598.

Johns, R.H., and C.A. Doswell III, 1992: Severe local storms forecasting. *Wea. Forecasting*, **7**, 588-612.

Meier, Keith, 1993: Jet streak dynamics: Effects of curvature. *Western Region Technical Attachment No. 93-03*. [Available from the NWS Western Region, P.O. Box 11188, Salt Lake City, Utah 84147].

Monteverdi, J.P., 1993: A case study of the operational usefulness of the SHARP Workstation in forecasting a mesocyclone-induced cold sector tornado event in California. *NOAA Technical Memorandum. NWS WR-219*. 22 pp.

Monteverdi, J.P., S.A. Braun, and T.C. Trimble, 1988: Funnel clouds in the San Joaquin Valley, California. *Mon. Wea. Rev.*, **104**, 1289-1296.

Reed, R., and W. Blier, 1986: A case study of comma cloud development in the Eastern Pacific. *Mon. Wea. Rev.*, **114**, 1681-1695.

Uccellini, L.W., and P.J. Kocin, 1987: The interaction of jet streak circulations during heavy snow events along the East Coast of the United States. *Wea. Forecasting*, **2**, 289-308.

Weisman, M.L., and J.B. Klemp, 1982: The dependence of numerically simulated convective storms on vertical shear and buoyancy. *Mon. Wea. Rev.*, **110**, 504-520.

- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins (Retired) and Alexander R. Smith. Fifth Revision, July 1992. (PB92 220177)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82 158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82 177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidli, April 1969 (Revised December 1986). (PB87 142063/AS)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteris, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85 110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85 132694)
- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 1878887AS)
- 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lussky, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lussky, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Radid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
- 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
- 199 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Lussky, April 1987. (PB87 185229/AS)
- 200 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke, September 1987. (PB88 110895/AS)
- 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson and George Clark, September 1987. (PB88 114038/AS)
- 202 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88 101935/AS)
- 203 An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern California. John P. Monteverdi and Scott A. Braun, October 1988. (PB89 121297/AS)
- 204 Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Carven Scott, November 1988. (PB89 128649/AS)
- 205 Forecast Guidelines for Fire Weather and Forecasters -- How Nighttime Humidity Affects Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS)
- 206 A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarters, Scientific Services Division, August 1989. (PB89 230833/AS)
- 207 The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, June 1990. (PB90-240268)
- 208 Meteorological Factors Contributing to the Canyon Creek Fire Blowup, September 6 and 7, 1988. David W. Goens, June 1990. (PB90-245085)
- 209 Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow Whitlatch, December 1990. (PB91-129239)
- 210 Hydrotools. Tom Egger, January 1991. (PB91-151787/AS)
- 211 A Northern Utah Soaker. Mark E. Struthwolf, February 1991. (PB91-168716)
- 212 Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 1991. (PB91-208439)
- 213 Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Mollner, July 1991. (PB91-227405/AS)
- 214 Emergency Operational Meteorological Considerations During an Accidental Release of Hazardous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424)
- 215 WeatherTools. Tom Egger, October 1991. (PB93-184950)
- 216 Creating MOS Equations for RAWs Stations Using Digital Model Data. Dennis D. Gettman, December 1991. (PB92-131473/AS)
- 217 Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 1992. (PB92-196104)
- 218 NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 1992. (PB93-146785)
- 219 A Case Study of the Operational Usefulness of the Sharp Workstation in Forecasting a Mesocyclone-Induced Cold Sector Tornado Event in California. John P. Monteverdi, March 1993. (PB93-178697)
- 220 Climate of Pendleton, Oregon. Claudia Bell, August 1993. (PB93-227536)
- 221 Utilization of the Bulk Richardson Number, Helicity and Sounding Modification in the Assessment of the Severe Convective Storms of 3 August 1992. Eric C. Evenson, September 1993.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS--Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS--Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161