



NOAA Technical Memorandum NWS WR-228

**FORECASTING MINIMUM TEMPERATURES IN
THE SANTA MARIA AGRICULTURAL DISTRICT**

**Wilfred Pi
National Weather Service Forecast Office
Monterey, California**

**Peter Felsch
National Weather Service Office
Missoula, Montana**

December 1994

**U.S. DEPARTMENT OF
COMMERCE**

/ National Oceanic and
Atmospheric Administration

/ National Weather
Service



NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P.O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 6285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidli, April 1968 (Revised March 1986). (PB86 177672/AS). (Revised October 1991 - PB92-115062/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB 179084)
- 32 Probability Forecasting—A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento—Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. (PB 184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 1969. (PB 185068)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185782)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)
- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquettes, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon. Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 65 Climate of Sacramento, California. Tony Martini, April 1990. (Fifth Revision) (PB89 207781/AS)
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)

- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251)
- 83 A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS)
- 96 Map Type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 694)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Card, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 126 Climate of San Francisco. E. Jan Null, February 1978. Revised by George T. Pericht, April 1988. (PB88 208624/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudgel, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298618/AS)

NOAA Technical Memorandum NWS WR-228

**FORECASTING MINIMUM TEMPERATURES IN
THE SANTA MARIA AGRICULTURAL DISTRICT**

Wilfred Pi
National Weather Service Forecast Office
Monterey, California


Peter Felsch
National Weather Service Office
Missoula, Montana

December 1994

UNITED STATES
DEPARTMENT OF COMMERCE
Ronald H. Brown, Secretary

National Oceanic and
Atmospheric Administration
D. James Baker, Under Secretary
and Administrator

National Weather Service
Elbert W. Friday, Jr., Assistant
Administrator for Weather Services



**This publication has been reviewed
and is approved for publication by
Scientific Services Division,
Western Region**



**Kenneth B. Mielke, Chief
Scientific Services Division
Salt Lake City, Utah**

TABLE OF CONTENTS

I.	INTRODUCTION	1
II.	KEY TEMPERATURE STATION DESCRIPTION/CLIMATOLOGY AND MINIMUM TEMPERATURE FORECASTING SCHEME	3
	A. WINTER KEY TEMPERATURE STATIONS	4
	B. SPRING KEY TEMPERATURE STATIONS	8
III.	CONCLUSION AND RECOMMENDATIONS	13
IV.	REFERENCES	13

FIGURES AND TABLES

FIGURE 1	LOCATION OF KEY AGRICULTURAL STATIONS	15
FIGURE 2	PRINTOUT OF ANALOGOUS DAYS PROGRAM	16
TABLE 1	CORRECTION CONSTANTS FOR DEW POINT AND HUMIDITY	17
TABLE 2	KEY STATION STATISTICS	18

FORECASTING MINIMUM TEMPERATURES IN THE SANTA MARIA AGRICULTURAL DISTRICT

Wilfred Pi
National Weather Service Forecast Office
Monterey, California

Peter Felsch
National Weather Service Office
Missoula, Montana

ABSTRACT

The National Weather Service Office in Santa Maria has developed a method of forecasting minimum temperatures for each of the 14 key agricultural temperature stations in San Luis Obispo and northern Santa Barbara Counties. Data were collected from the past 15 winter and spring frost seasons to develop a PC program called "Analogous Days". For further assistance in forecasting minimum temperatures, data from the past four seasons were used to develop a minimum temperature forecasting scheme, using surface pressure differences, 850 mb winds and temperatures, and winds at 1000 and 2000 feet. The purpose of this project is to develop a better method of accurately forecasting minimum temperatures for each of the key agricultural temperature stations. Growers in the Agricultural District need to know the expected minimum temperatures in order to take precautions for their frost sensitive crops and plan field activities. This project will also benefit the National Weather Service after Modernization and Reorganization (MAR) occurs.

I. INTRODUCTION

The Agricultural District of San Luis Obispo County and northern Santa Barbara County has extremely complex terrain consisting of small mountain ranges traversing both counties in different directions to form several valleys and canyons. These mountain ranges may funnel cold or warm air into a valley or canyon during the nighttime hours, dependent on the wind conditions. Because each mountain range is oriented in a different direction, each of the valleys and canyons have a different nighttime temperature structure based on the direction and speed of the wind.

The key agricultural temperature stations are located in different valleys and/or canyons, and thus, have their own distinct

nighttime temperature structure. There are two major topographical features in this Agricultural District influencing this temperature structure: the Pacific Ocean and the Santa Lucia Mountains. The Pacific Ocean, on the west side of the Agricultural District, affects those nearby valleys. The Santa Maria Valley and the Lopez Lake Canyon are examples of coastal valleys. Temperatures at key agricultural temperature stations (Oceano, Upper Arroyo Grande, Santa Maria, Bonita School) in these two valleys are typically warmer, especially in a westerly wind flow, because the stations are close to the ocean. Other key temperature stations such as Los Alamos, Zaca, Santa Ynez, Sisquoc, Nipomo Mesa, and Nipomo Foothill are normally cooler, due to sheltering by surrounding local mountains. Two other key temperature stations, Morro Grange

and Price Canyon, are in narrow coastal valleys, and temperatures are strongly dependent on wind conditions. The Santa Lucia Mountains in San Luis Obispo County, which are oriented northwest to southeast about 15 miles from the ocean, isolate the ocean effects to the Salinas and Cuyama Valleys. If the winds are light, key temperature stations in these valleys (Radike Vineyard, Estrella River, Shandon) will have strong radiational cooling and generally have the lowest temperatures among the key temperature stations. Figure 1 shows the location of the key agricultural temperature stations.

Vegetables, citrus, and avocados are the principle crops grown near the key temperature stations Santa Maria, Bonita School, Nipomo Mesa, Nipomo Foothill, Morro Grange, Oceano, and Upper Arroyo Grande. The other key temperature stations are located in grape vineyards and apple orchards. The winter growing season is confined to the aforementioned key temperature stations during the period when the grapes and apples are dormant. The spring frost season is extended to include all the other key temperature stations during the period when buds form on the grapevines and apple trees. Accurately forecasting minimum temperatures on expected cold nights is extremely important to the growers, as all of the crops are sensitive to temperatures below freezing.

Forecasting minimum temperatures for all the key temperature stations are dependent on the predicted minimum temperature at Santa Maria. A useful tool to help forecast the minimum temperature at Santa Maria is the FOUS POP/Ceiling/Wind Guidance product. Another useful tool is the Analogous Days PC program, developed at the National Weather Service Office in Santa Maria by

Jeff Smith, currently Hydrologist at the River Forecast Center in Salt Lake City. This program was improved by Dave Rosenberg, Met Intern at WSFO San Francisco, using the Paradox spreadsheet. The Analogous Days Program has been developed using data from the past 15 winter and spring frost seasons and will continue to be updated after every frost season.

The first part of the program calculates minimum temperatures for Santa Maria during the winter and spring seasons, and for Paso Robles during the spring season based only on the following hygrometric equation:

$$\begin{aligned} \text{SMX MIN} &= Dp - ((RH-30)/3) + VD + VH \\ \text{PRB MIN} &= Dp - ((RH-18)/3) + VD + VH \end{aligned}$$

where Dp is the dew point

VD is the correction constant for dew point
VH is the correction constant for humidity
(correction constants are shown in Table 1)

This hygrometric equation was originally developed by Ellison (1928) and has been used in the past to forecast minimum temperatures in other Agricultural Districts (Fritz, 1970; Tuft, 1975; Hill, 1976; Oard, 1977; Rogers and Rohli, 1991). The numerical values of the equation and correction constants were adjusted for Santa Maria and Paso Robles by Wilbur Shigehara, currently Meteorologist in Charge at NWS San Diego, using raw data. The temperature and dew point from 1600 local time on that afternoon are entered into the program by the forecaster. The relative humidity, correction constants, and minimum temperature forecast are then calculated by the program.

The second part of the program searches through an archive file of the past 15 years of data to find analogous days based on the

Santa Maria and Paso Robles forecast minimum temperatures and other meteorological parameters as described below.

1. The Santa Maria to Bakersfield surface pressure difference gives an indication of the east-west surface pressure gradient. If the pressure at Santa Maria is higher than at Bakersfield, the pressure gradient force is directed toward lower pressure and suggests a westerly flow.
2. The Salinas to Santa Barbara surface pressure difference gives a general indication of whether strong winds can be expected. If the surface pressure at Salinas is about 3.5 mb higher than surface pressure at Santa Barbara, then it is suggested that there will be strong northwest winds. Northwest winds play a role in modifying minimum temperatures at some of the key stations.
3. The closest upper-air sounding is taken at Vandenberg Air Force Base, 10 miles southeast of Santa Maria, which provides important data consisting of the 850 mb and 500 mb temperatures and winds, as well as the winds at 1000 and 2000 feet.

The forecaster enters the 850 mb and 500 mb temperatures, winds, and the magnitude of the surface pressure differences from Santa Maria to Bakersfield and from Salinas to Santa Barbara from 0000 UTC into the Analogous Days Program. The criteria for finding the analogous days for the minimum temperature the next morning from the entered data are:

850 mb temperature: $\pm 4^{\circ}\text{C}$
Pressure differences: $\pm 1.5\text{ mb}$

Dew point: $\pm 2^{\circ}\text{F}$
Formula minimum(s): $\pm 4^{\circ}\text{F}$

The analogous days that meet all the above requirements are listed, displaying the minimum temperatures for all the key temperature stations for that day. Archived weather maps from those analogous days can be compared to the current weather maps. The analogous day(s), with a weather pattern that most resembles the current weather pattern, are selected and the forecast minimum temperature for Santa Maria is determined. A sample printout of the Analogous Days Program is shown in Fig. 2.

Once a forecast minimum temperature for Santa Maria is obtained, forecast minimum temperatures for the other key stations can be obtained by using the developed minimum temperature forecasting schemes, which use the Analogous Day Program as guidance.

II. KEY TEMPERATURE STATION DESCRIPTION/CLIMATOLOGY AND MINIMUM TEMPERATURE FORECASTING SCHEME

The climatology of each key agricultural temperature station is based mainly upon the Santa Maria-Bakersfield surface pressure difference, the Salinas-Santa Barbara surface pressure difference, and the upper-air data as discussed earlier. Another meteorological parameter used is the surface pressure difference between Santa Maria and Paso Robles, which gives an indication of the north-south pressure gradient. If the pressure of Santa Maria is higher than at Paso Robles, the pressure gradient force is positive and the flow is suggested to have a southerly component. The minimum temperature forecasting

scheme was developed after the 1990-1991 winter and spring frost seasons using the key agricultural temperature station climatology as a guide, and then was fine-tuned after verification from the last two frost seasons. This scheme is to be used along with the Analogous Days Program and given equal weight. Frequently, the Analogous Days Program will not have any days to compare. In this case, the minimum temperature formulas are used along with this scheme. The following is a brief description of the climate of each key agricultural temperature station and its minimum temperature forecasting scheme.

SMX = Santa Maria

PRB = Paso Robles

SMX-BFL = Santa Maria - Bakersfield surface pressure difference.

SMX-PRB = Santa Maria - Paso Robles surface pressure difference.

SNS-SBA = Salinas - Santa Barbara surface pressure difference.

A. WINTER KEY TEMPERATURE STATIONS

i) **SANTA MARIA** When the Santa Maria-Bakersfield surface pressure difference is ≤ -2.0 mb and the Santa Maria-Paso Robles difference is ≤ -1.0 mb, consideration of northeast or east surface wind continuing through the night may moderate minimum temperatures. However, surface winds are more likely with a surface pressure difference of -3 or -4 mb. Likewise, in springtime if the pressure difference from Salinas-Santa Barbara is ≥ 6 mb, there is a good chance of nighttime wind from the north or northwest. This is especially true if you have pressure difference ≥ 7 mb and an 850 mb wind of ≥ 35 kts (Note: This particular rule holds for most of the key temperature stations, winter or spring). However, keep in mind that a very cold

airmass, i.e. an 850 mb temperature of $\leq -3^{\circ}\text{C}$, can stabilize the airmass, diminishing the wind beneath an inversion and producing a cold night. This occurs when high winds aloft are decoupled from the surface winds by the temperature inversion.

Santa Maria can be one of the colder winter key temperature stations under clear skies and light wind. A situation often encountered during mid-winter occurs when a cold front moves through the district late in the afternoon or early in the evening and has a cold airmass behind it (i.e. 850 mb temp $< 1^{\circ}\text{C}$). During these situations, the forecast often will call for showers or partial clearing, but strong downward vertical velocity behind the front creates a nearly clear night with light wind. This leads to an unexpected early morning frost. Satellite pictures and upper-air data are the greatest aid in detecting these potentially troublesome situations. These situations can take place at any of the winter key temperature stations, not just Santa Maria.

The coldest situations (2/15/90, 1/16/87, 1/6/78) over the entire district occur when: (1) the 850 mb temperature is $\leq -3^{\circ}\text{C}$, (2) the surface dewpoint is in the lower 20s $^{\circ}\text{F}$ or 10s $^{\circ}\text{F}$, and (3) the daytime surface temperature is generally $\leq 55^{\circ}\text{F}$. Surface pressure differences in these situations are usually only light to moderate offshore, but stronger surface winds may prevent very cold temperatures.

SANTA MARIA SCHEME (Winter Season)

1. a) If 850 mb winds are 190° to 280° and ≥ 20 kts, onshore flow will allow low- to mid-level clouds to be advected into the coastal plain: thus, forecast clouds for SMX.

- b) If SMX-PRB $\geq +1.5$ mb, and SNS-SBA $\leq +3.0$ mb, with 850 mb winds ≤ 25 kts, a southerly surface flow is more dominant than northwest flow: thus, forecast clouds for SMX.
2. If no clouds are forecast, and 500 mb heights are ≤ 560 dam, with 850 mb temperature of $\leq 0^\circ\text{C}$, forecast temperatures $\leq 32^\circ\text{F}$. A cold airmass will produce cold minimum temperatures regardless of winds.
 3. If no clouds are expected and 850 mb winds are ≤ 15 kts, forecast closer to the predicted formula temperature from the Analogous Days Program than the FOUS guidance. Otherwise, forecast closer to the FOUS guidance. Verification of comparisons between FOUS and the formula minimum temperature has shown that the formula forecast is better in this situation.

In the spring, the combination of longer days and a greater influence of an ocean breeze cause Santa Maria and the other key temperature stations near the coast and coastal valleys except Morro Grange, to warm by 1 or 2°F at night.

SANTA MARIA SCHEME (Spring Season)

1. a) If 850 mb winds are 190° to 280° and ≥ 20 kts, forecast clouds for SMX.
b) If SMX-PRB $\geq +1.5$ mb, and SNS-SBA $\leq +3.0$ mb with 850 mb winds ≥ 25 kts, forecast clouds.
2. Use the Analogous Days formula from 0 to $+2^\circ\text{F}$, if you are expecting no clouds and no wind modification.

a) If clouds are expected, add $\geq 4^\circ\text{F}$ to the Analogous Days formula.

b) If SNS-SBA $\geq +4.0$ mb, then wind will modify temperatures $\geq 4^\circ\text{F}$.

3. If 500 mb heights are ≤ 543 dam, forecast minimum temperatures near 32°F .

ii) BONITA SCHOOL The same rules used at Santa Maria apply to Bonita School, although Bonita School is usually about 1 to 2°F warmer than Santa Maria. Its proximity to the ocean or a bit more wind at night causes the difference.

BONITA SCHOOL SCHEME (Winter and Spring Season)

1. a) If SMX-BFL and SMX-PRB $\geq +0.1$ mb, forecast 0 to 1°F warmer than SMX. If SNS-SBA $> +3.5$ mb, forecast $\geq 2^\circ\text{F}$ warmer than SMX. The combination of an onshore flow and wind modification (increased moisture at low levels) causes the warming.
b) If SMX-BFL and SMX-PRB < -0.1 mb, forecast 0 to 2°F colder than SMX. If SNS-SBA $< +1.5$ mb and 850 mb winds are 20° to 160° , forecast 2 to 3°F colder than SMX. A light offshore flow allows for cool air drainage and good radiational cooling.
2. If no clouds are forecast, and 500 mb heights are ≤ 560 dam, and 850 mb temperature is $\leq 0^\circ\text{C}$, forecast temperatures $\leq 32^\circ\text{F}$. When the airmass is cold, the station gets cold.

iii) OCEANO This station is usually 1- 3°F warmer than Santa Maria, but can get slightly cooler if Santa Maria's temperature gets moderated by wind. A light onshore

breeze can help modify minimum temperatures if dewpoints are in the middle 40s°F near the ocean. The coldest mornings are in light offshore flow conditions. This station is susceptible to moderate air drainage down Arroyo Grande Creek Canyon.

OCEANO SCHEME (Winter and Spring Season)

1. a) If SMX-BFL and SMX-PRB $\geq +0.1$ mb, forecast 0 to 2°F warmer than SMX. If SNS-SBA $> +3.5$ mb, forecast $\geq 2^\circ\text{F}$ warmer than SMX. See Bonita School reasoning.
 - b) If SMX-BFL and SMX-PRB is between $+0.1$ and -1.0 mb, forecast 0 to 2°F colder than SMX. If SNS-SBA $< +3.0$ mb and 850 mb winds are 20° to 160° , forecast 2 to 3°F colder than SMX. See Bonita School reasoning.
 - c) If SMX-BFL and SMX-PRB < -1.0 mb, forecast $\geq 2^\circ\text{F}$ higher than SMX. A north to northeast drainage and offshore flow will modify minimum temperatures.
2. If no clouds expected, and 500 mb heights are ≤ 560 dam and 850 mb temperature is $\leq 0^\circ\text{C}$, forecast temperatures $\leq 32^\circ\text{F}$. If the air mass is cold, the station will be cold.

iv) NIPOMO FOOTHILL This is the one of toughest winter key temperature station for which to forecast. On the coldest mornings there is a light onshore flow when the Santa Maria-Bakersfield surface pressure difference is between 2.0 mb and -1.0 mb and the Santa Maria-Paso Robles pressure difference is between 2.0 mb and -1.5 mb. This seems to weaken the normal drainage wind and allows good radiational cooling. In this situation, this station will

be 1 to 3°F colder than Santa Maria and about as cold as Nipomo Mesa. This station is susceptible to moderate air drainage off the hills making the station 3 to 7°F warmer than Santa Maria. If surface offshore pressure differences are moderate to strong (Santa Maria-Bakersfield and Santa Maria-Paso Robles pressure differences < -2.0 mb), there is a good chance of nighttime wind modification. If the winds at 1000 or 2000 feet are west, northwest, or northeast and ≥ 15 knots, then wind modification of the nighttime temperature can be expected. Temperature modification is not likely if the wind direction is north or southeast, regardless of wind strength due to the orientation of the surrounding coastal hills.

NIPOMO FOOTHILL SCHEME (Winter and Spring Season)

1. a) If SMX-BFL < -2.0 mb and SMX-PRB < -1.5 mb, forecast 0 to 3°F warmer than SMX. If SNS-SBA $> +3.0$ mb, forecast $\geq 3^\circ\text{F}$ warmer than SMX. North to northeast drainage and offshore winds will modify temperatures.
- b) If SMX-BFL and SMX-PRB $> +2.0$ mb, forecast 1 or 2°F warmer than SMX. A good onshore flow will advect moisture from the ocean and modify temperatures.
- c) If SMX-BFL is between $+2.0$ and -2.0 mb; and if SMX-PRB is between $+2.0$ and -1.5 mb, forecast 0 to 3°F colder than SMX. Wind will not be much of a factor and there will be good radiational cooling. If SNS-SBA $> +3.0$ mb, forecast the about the same minimum temperature as SMX. Northwest winds will also help to modify temperatures if this criteria is met.

2. If no clouds are expected, 500 mb heights are ≤ 560 dam, and the 850 mb temperature $\leq 0^{\circ}\text{C}$, forecast minimum temperatures $\leq 32^{\circ}\text{F}$.

Note: Generally if the winds at 1000 feet are ≥ 15 knots, minimum temperatures will be modified. If light winds prevail at 1000

feet, then forecast colder minimum temperatures.

v) NIPOMO MESA NE About the only time this site is not the coldest of the stations is with a light onshore flow. When that happens, Nipomo Foothill and Morro Grange can get slightly colder than Nipomo Mesa. Also, when strong offshore flow is occurring (Santa Maria-Bakersfield pressure difference < -3.5 mb, plus strong northeasterly winds at 2000 feet), this station will be a little warmer than Santa Maria.

NIPOMO MESA SCHEME (Winter Season)

1. a) If SMX-BFL < -2.5 mb and SMX-PRB < -2.0 mb, forecast $\pm 1^{\circ}\text{F}$ from the SMX forecast minimum temperature. A strong north to northeast flow will modify temperatures.

b) If no clouds are expected, and if SMX-BFL is between $+2.0$ and -2.5 mb, and if SMX-PRB is between $+2.0$ and -2.0 mb, forecast 1 to 4°F colder than SMX. If SNS-SBA $< +2.0$ mb forecast 3 to 5°F colder than SMX. A light offshore flow draining in cold air is the coldest scenario for Nipomo Mesa. A light onshore flow starts to bring in some moisture from the ocean, modifying temperatures.

c) If SMX-BFL and SMX-PRB $> +2.0$

mb, forecast 0 to 2°F colder than SMX.

NIPOMO MESA SCHEME (Spring Season)

1. a) If no clouds and SMX-BFL and SMX-PRB $> +2.0$ mb, forecast 0 to 2°F colder than SMX. See winter season.

b) If SMX-BFL < -2.5 mb and SMX-PRB < -2.0 mb, forecast $\pm 1^{\circ}\text{F}$ from SMX. See winter season.

c) If SMX-BFL > -2.5 mb and SMX-PRB > -2.0 mb, forecast 1 to 4°F colder than SMX. See winter season.
2. If rule 1.c above is true, and the 850 mb temperature is $\leq +1^{\circ}\text{C}$, then forecast $\geq 4^{\circ}\text{F}$ colder than SMX unless SNS-SBA $> +5.0$ mb, modify to 1 to 3°F colder than SMX.
3. If no clouds are expected, forecast 1°F colder than SMX Analogous Days formula.

vi) UPPER ARROYO GRANDE This is the warmest of the winter key temperature stations. This station almost always has a nighttime drainage wind, as it is situated in an east-west valley that drains from the mountains to its east. This station will fall to near or below freezing only in our coldest situations, and will seldom drop below $27-29^{\circ}\text{F}$ even when Santa Maria has dropped to 22 or 23°F .

UPPER ARROYO GRANDE SCHEME (Winter and Spring Season)

If SMX-BFL and SMX-PRB $> +0.5$ mb and 850 mb temperature is $\leq 0^{\circ}\text{C}$, forecast minimum temperatures 0 to 3°F colder than SMX. The combination of onshore

flow and a cold airmass will suppress the drainage wind from the mountains. Otherwise forecast $\geq 2^{\circ}\text{F}$ higher than SMX.

vii) MORRO GRANGE This station is also a difficult location to forecast. This station is typically one of the warmest, especially with any kind of offshore flow, but gets cold with a light onshore flow when the nighttime wind is from a northwest direction. This can be the coldest key temperature station during the winter on the first night after a cold front has passed with northwest flow (i.e. Santa Maria-Bakersfield pressure difference is ≥ -1.5 mb and Santa Maria-Paso Robles pressure difference is ≥ -1.0 mb). The reason for this is that the surrounding hills block a north to northwest flow creating nearly calm conditions at Morro Grange.

MORRO GRANGE SCHEME (Winter Season)

1. If SMX-BFL and SMX-PRB ≥ -1.0 mb, forecast 3 to 5°F colder than SMX. If SNS-SBA $< +2.5$ mb, forecast 6°F colder than SMX. The flow is onshore and blocked by the surrounding hills.
2. If SMX-BFL and SMX-PRB is between -1.0 and -2.0 mb, and SNS-SBA $< +2.0$ mb, forecast between 2°F colder and 1°F warmer than SMX. If SNS-SBA $> +3.0$ mb, forecast $\geq 2^{\circ}\text{F}$ warmer. A drainage and offshore flow becomes a factor to begin modifying temperatures. A strong northwest flow will also help modify temperatures.
3. If SMX-BFL and SMX-PRB < -2.0 mb, forecast $\geq 2^{\circ}\text{F}$ warmer than SMX. A moderate to strong offshore flow will modify temperatures considerably.

4. If high temperature from the previous day $\geq 70^{\circ}\text{F}$ and the high temperature on the next day is expected to be the same, forecast minimum temperatures of $\geq 30^{\circ}\text{F}$ on expected cold nights. Verification has shown that the diurnal variation on warm days does not exceed 40°F . Therefore, enough warm air from the day should prevent very cold temperatures from occurring.

In the spring, the surrounding hills prevent the ocean breeze from modifying minimums at this station like the other winter key temperature stations. As a result, the temperature departure from Santa Maria is larger in the spring season than in the winter season.

MORRO GRANGE SCHEME (Spring Season)

1. If no clouds are expected, 500 mb heights are ≤ 570 dam, and 850 mb temperature is $\leq +3^{\circ}\text{C}$, forecast minimum temperatures $\leq 32^{\circ}\text{F}$. A cool airmass along with the typical spring onshore flow brings cold nights.
2. If no clouds are expected, forecast 3°F colder than SMX Analogous Days formula.
3. If SMX-BFL and SMX-PRB ≥ -1.0 mb, forecast 4 to 6°F colder than SMX. Otherwise, forecast at or above SMX Analogous Days formula. If SNS-SBA $> +2.5$ mb, add 2 to 3°F . A strong onshore flow will help to modify temperatures a few degrees.

B. SPRING KEY TEMPERATURE STATIONS

i) SHANDON This station, at 1100 feet elevation, is the coldest key temperature

station in the district. At times, the newer Radike Vineyard station can be just as cold. This station's minimum temperatures are significantly lower than all other stations when the 850 mb wind is northwesterly and greater than 40 knots, because all other stations will remain above 32°F due to wind. Shandon is protected from the northwest wind, especially at night, even if there is a northwest wind during the day. This station has dropped to 27°F when all other key temperature stations were in the upper 30s°F and 40s°F. Generally, Shandon will be around 7°F colder than Santa Maria and 3°F or so colder than Paso Robles. Occasionally irrigation in the alfalfa field across the road can be a source of modification, but it has not been quantified.

When the surface pressure difference between Paso Robles and Fresno becomes ≤ -1 mb, the possibility exists for nighttime wind at Shandon from the east as the site opens out into the Central Valley through a series of canyons. Under these conditions the Radike Vineyard station will usually be colder than Shandon.

SHANDON SCHEME

1. If there are clouds and it rained, use #2 only.
2. Add 2 to 3°F to the PRB Analogous Days formula, if one of the following is true:
 - a) there are clouds and/or it has rained,
 - b) the SNS-SBA $> +6.0$ mb,
 - c) 850 mb winds are N to NE greater than 40 kts.

3. If no clouds/rain are forecast, and
 - a) If SMX-BFL and SMX-PRB is between -0.5 and -2.6 mb, forecast 10 to 14°F colder than SMX. Light to moderate offshore flow is the coldest scenario.
 - b) If SMX-BFL and SMX-PRB is between $+0.5$ and $+2.0$ mb, forecast 5 to 9°F colder than SMX. An onshore flow begins to modify temperatures because there is more moisture in the air.
 - c) If SMX-BFL and SMX-PRB $> +2.0$ mb, forecast ≤ 4 °F colder than SMX. The onshore flow is stronger.

ii) RADIKE VINEYARD This station is located at 1000 feet elevation and is surrounded by small hills. Any winds ≥ 25 kts at 850 mb will cause minimum temperatures to be moderated. (Note: The winds at 1000 and 2000 feet from the Vandenberg upper-air sounding are not applicable to Shandon and Radike Vineyard because of the Santa Lucia Mountains.) Normally, this station will be about 1 or 2°F higher than Shandon. On nights when you do not suspect wind to be a major factor, forecast about the same minimum temperature or a degree higher than Shandon.

RADIKE VINEYARD SCHEME

1. If no clouds/rain are forecast, forecast 0 to 3°F colder than PRB Analogous Days formula.
2. Forecast 2 to 3°F warmer than the PRB Analogous Days formula if either,
 - a) the 850 mb winds are ≥ 30 kts, or
 - b) if you expect clouds or rain.

3. If no clouds/rain are forecast and
 - a) If SMX-BFL and SMX-PRB is between -2.0 and -1.0 mb, forecast 2 to 6°F colder than SMX. A light to moderate offshore flow begins to modify temperatures.
 - b) If SMX-BFL and SMX-PRB is between -1.0 and -0.5 mb, forecast 8°F colder than SMX.
 - c) If SMX-BFL and SMX-PRB is between -0.5 and +1.5 mb, forecast 9 to 12°F colder than SMX. A light onshore flow is the coldest scenario as the surrounding mountains block north to northwest winds.
 - d) If SMX-BFL and SMX-PRB > +1.5 mb, forecast 4 to 7°F colder than SMX. A stronger onshore flow begins to modify temperatures.

iii) ESTRELLA RIVER The minimum temperatures at Estrella River averages 1°F lower than Paso Robles and 2 to 3°F higher than Shandon. Both Paso Robles and Estrella River are susceptible to northwest nighttime wind after a frontal passage. If calm conditions exist, the Analogous Days Program works fairly well. This station can be moderated by morning fog after a precipitation event.

ESTRELLA RIVER SCHEME

1. If there are clouds and it has rained, use #2 only.
2. Add 2 to 3°F to the PRB formula if one of the following is true:
 - a) there are clouds, it has rained,
 - b) if SNS-SBA > +6.0 mb,

c) if 850 mb winds are N to NE, ≥ 40 kts.

3. If no clouds/rain are forecast and
 - a) If SMX-BFL and SMX-PRB is between +0.5 and -2.6 mb, forecast 7 to 12°F colder than SMX. See Shandon reasoning.
 - b) If SMX-BFL and SMX-PRB is between +0.5 and +2.0 mb, forecast 3 to 7°F colder than SMX. See Shandon reasoning.
 - c) If SMX-BFL and SMX-PRB > +2.0 mb, forecast $\leq 2^\circ\text{F}$ colder than SMX. See Shandon reasoning.

iv) PRICE CANYON This station is usually about 2 or 3°F colder than Santa Maria except when you expect it to be windy at San Luis Obispo Airport. Strong north to northwest wind can be expected at this station all night, as well as at San Luis Obispo Airport when the surface pressure difference between Paso Robles and San Luis Obispo exceeds 2 or 3 mb and when the 850 mb wind is in excess of 25 kts.

PRICE CANYON SCHEME

1. If winds up to 850 mb are N to NE at ≥ 25 kts and/or SMX-PRB ≤ -3.0 mb, do not forecast freezing temperatures and go to #3. A strong N to NE wind modifies temperatures.
2. a) If SMX-BFL and SMX-PRB is between -1.5 and +2.5 mb, and if SNS-SBA $\leq +4.0$ mb, forecast 2 to 5°F colder than SMX. A light offshore flow or a light to moderate onshore flow are the coldest scenarios as the coastal hills protect the station.

b) If SMX-BFL and SMX-PRB ≤ -2.0 mb, forecast $\geq 2^\circ\text{F}$ warmer than SMX. A moderate to strong offshore flow moderates temperatures.

c) If SMX-BFL and SMX-PRB $> +2.5$ mb, forecast 0 to 2°F warmer than SMX. Strong onshore flow will modify temperatures.

3. If no clouds are forecast, use SMX Analogous Days formula.

v) SISQUOC This station runs about 2 to 3°F colder than Santa Maria but can be affected by both drainage and strong nighttime offshore wind.

SISQUOC SCHEME

1. If clouds are forecast and/or SNS-SBA $> +6.0$ mb, use #2 only.

2. a) If SMX-BFL and SMX-PRB $\geq +2.0$ mb, forecast 3 to 4°F colder than SMX. The station is protected from an onshore flow by hills to the west, which is the coldest scenario.

b) If SMX-BFL and SMX-PRB is between $+1.5$ to -1.5 mb, forecast $\leq 2^\circ\text{F}$ colder than SMX. A weak flow will allow some drainage wind from the east.

3. a) If 850 mb temperature is $\leq +1^\circ\text{C}$, forecast 3 to 4°F colder than SMX. When the airmass is cold, this station will be cold regardless of pressure differences.

b) If 850 mb temperature is $\geq +2^\circ\text{C}$, forecast $\leq 2^\circ\text{F}$ colder than SMX.

4. If no clouds and wind are forecast, then use SMX Analogous Days formula.

vi) LOS ALAMOS This station runs 2 to 3°F colder than Santa Maria, but can occasionally run 4 or 5°F colder early in the spring during still nights when little wind is expected. This station has always been a bit confusing since it's closer to the ocean than the other nearby stations and yet it is colder. Perhaps the drainage of cold air from the surrounding hills combined with pooling of the cold air in this area of the valley is responsible for this. Nighttime northwest wind can be a moderating factor but nighttime drainage wind is not.

LOS ALAMOS SCHEME

1. If clouds are expected and SNS-SBA $> +6.0$ mb, forecast above freezing. A strong onshore flow will modify temperatures.

2. If no clouds and wind as #1 above, and 850 mb temperature is:

a) $\leq -1^\circ\text{C}$, forecast 6°F colder than SMX.

b) 0 to $+1^\circ\text{C}$, forecast 3 to 5°F colder than SMX.

c) $> +2^\circ\text{C}$, forecast $\leq 2^\circ\text{F}$ colder than SMX.

Like Sisquoc, with a cold airmass, the station will be cold. The warmer the airmass, the warmer the station.

3. If no clouds and SNS-SBA $\geq +6.0$ mb, forecast 1 degree colder than SMX. If SNS-SBA $\leq +6.0$ mb, forecast 2 to 3°F colder.

vii) ZACA This is about the warmest of the "southern four" key temperature stations (Sisquoc, Zaca, Santa Ynez and Los Alamos). It regularly gets a drainage wind

from the east or gets mixing from nighttime northwesterly winds. When wind is not a factor, this station will be just slightly warmer than the other three stations and about 1 to 2°F colder than Santa Maria.

ZACA SCHEME

1. If clouds are expected and/or SNS-SBA $\geq +7.0$ mb, forecast above freezing. A strong onshore flow will modify temperatures.
2. If no clouds are forecast and wind as in #1, then:
 - a) If SMX-BFL and SMX-PRB is between +1.5 to +2.5 mb, forecast 5 to 7°F colder than SMX. A light to moderate onshore flow is the coldest scenario for minimum temperatures.
 - b) If SMX-BFL and SMX-PRB is between +1.5 and -1.5 mb, forecast 2 to 5°F colder than SMX. A weak flow will allow for some drainage wind.
 - c) If SMX-BFL and SMX-PRB < -1.5 mb, forecast 1°F colder to 2°F warmer than SMX. With an offshore flow, a drainage wind will modify temperatures.
3. If no clouds and wind are forecast, then use SMX Analogous Days formula.

viii) SANTA YNEZ This station is interesting since it is one of the coldest of the "southern four" late in the winter through about mid-March (due to cold air draining into the area around it). As the days warm up, it becomes one of the warmest of the "southern four" late in spring, as nighttime cold air is offset by daytime warming. Therefore, in late

winter minimum temperatures at this station can be as much as 6°F lower than Santa Maria, then only 1 or 2°F lower by late spring. Temperatures at this station can be moderated by fog after rain.

SANTA YNEZ SCHEME

1. Use the Santa Maria Analogous Days formula and forecast 0 to 2°F colder if no clouds and wind.
 - a) If clouds are expected, add 3°F or more to the Analogous Days formula.
 - b) If SNS-SBA $> +5.0$ mb, wind will modify minimum temperatures up about 5°F.
2.
 - a) If SMX-BFL and SMX-PRB are between -2.5 and +0.5 mb, forecast 6 to 9°F colder than SMX. A light onshore and a light to moderate offshore flow are the coldest scenarios for minimum temperatures.
 - b) If SMX-BFL and SMX-PRB are between +0.5 to +2.0 mb, forecast 2 to 5°F colder.
 - c) If SMX-BFL and SMX-PRB are +2.0 mb or higher, forecast $\geq 2^\circ\text{F}$ warmer. An onshore flow brings in moisture from the west to modify temperatures.
3. If clouds are expected, use rule 1.a only.
4. After March, add 2 to 3°F to all of the above conditions.

Table 2 shows the average departure of the minimum temperatures of all the key agricultural temperature stations from the average minimum temperature at Santa Maria. Nights with high temperature ceilings are those preceded by a day with a

maximum temperature of $\leq 60^{\circ}\text{F}$, implying cold air advection with weak or no inversion, resulting in widespread freezing temperatures. Conversely, nights with low temperature ceilings are those preceded by a day with a maximum temperature of 60°F or higher, implying radiational cooling and a moderate to strong inversion, resulting in only isolated areas of freezing temperatures.

III. CONCLUSION AND RECOMMENDATIONS

Map types have been developed for different weather situations that cause cold temperatures to occur over the Agricultural District (Hamilton and Lussky, 1985; Crossan, 1981). A few examples are at 500 mb: (1) a ridge just off the coast of California, (2) a ridge directly over California, and (3) a cutoff low east of California. Using map types and examining the 36 to 60 hour prognostic charts from the NGM and AVN models, potentially cold periods within the district can be anticipated. Within 12 hours, the Analogous Days Program is used to predict the minimum temperature for Santa Maria and Paso Robles. Then, the minimum temperatures for each key agricultural temperature station are determined using the forecasting schemes, with guidance from the Analogous Days Program.

This method of forecasting is not only useful now, but will be even more useful after MAR comes into place. Under the current plan of MAR, the National Weather Service Office in Santa Maria is scheduled to be closed in the late 1990s. The forecasting for this Agricultural District may have to be handled by the Weather Forecast Offices (WFOs) in Oxnard and/or Monterey with guidance from the National Weather Service Office

in Riverside. This project will give the WFOs a good idea of the climatology of each key agricultural temperature station and help them with their agriculture forecasting responsibilities for this district.

The National Weather Service Office in Santa Maria will continue the development of the Analogous Days Program and the forecasting scheme as more data becomes available in the coming frost seasons and possibly combine the two into one PC forecasting program.

IV. REFERENCES

Crossan, Thomas R., 1981: Central San Joaquin Map Types, NOAA Technical Memorandum, NWS WR-173, Salt Lake City, Utah.

Ellison, Eckley S., 1928: A Critique on the Construction and Use of Minimum-Temperature Formulas, *Mon. Wea. Rev.*, **50**, 485-495.

Fritz, Arthur W., 1970: An Aid in Forecasting the Minimum Temperature at Medford, Oregon. NOAA Technical Memorandum NWS-WR-60, NWS Western Region, Salt Lake City, Utah.

Hamilton, Ronald S. and Glenn R. Lussky, 1985: Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest, NOAA Technical Memorandum NWS WR-191, Salt Lake City, Utah.

Hill, Christopher D., 1976: Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada During the Summer Months, NOAA Technical Memorandum NWS WR-104, NWS Western Region, Salt Lake City, Utah.

Oard, Michael J., 1977: A Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression, NOAA Technical Memorandum NWS WR-113, NWS Western Region, Salt Lake City, Utah.

Rogers, Jeffrey C. and Robert V. Rohli, 1991: Florida Citrus Freezes and Polar Anticyclones in the Great Plains, *J. Climate*, 4, 1103-1113.

Tuft, Wesley L., 1975: A Set of Rules for Forecasting Minimum Temperatures in Napa and Sonoma Counties, NOAA Technical Memorandum NWS WR-102, Salt Lake City, Utah.




Figure 1 - Locations of key agricultural stations

MINIMUM TEMPERATURE / ANALAGOUS DAYS PROGRAM

04/13/93

INITIAL DATA SET

SMX 00Z TEMP	61	DEG F	PRB 00Z TEMP	75	DEG F
SMX 00Z DWPNT	48	DEG F	PRB 00Z DWPNT	36	DEG F
SMX 00Z RH	63	%	PRB 00Z RH	24	%

CORRESPONDING NEXT MORNING'S FORMULAE MINIMUMS

SMX FCST MIN	39	DEG F	VD.. 3	VH..-1
SMX NAPA MIN	30	DEG F	VD..-4	VH.. 1
PRB FCST MIN	31	DEG F	VD.. 0	VH..-5

ADDITIONAL ANALAGOUS DAYS DATA

VBG 00Z 850 TEMP	10	DEG C	VBG 00Z 850 MB WIND	33015	
VBG 00Z 500 MB TEMP	-18	DEG C	VBG 00Z 500 MB WIND	34030	
SNS-SBA PRES DIFF	004.2	MB	SMX-BFL PRES DIFF	000.4	MB

SPRING SEASON

MORNING MINIMUMS

ANALAGOUS DATA

DATE	MORNING MINIMUMS											ANALAGOUS DATA				
	SM	NM	MG	SH	ER	RD	PC	SQ	LA	ZA	SY	PR	GRADIENTS sns sba	smx bf1	VBG 00Z 850 mb t wind	VBG 00Z 500 mb t wind
3/26/77	37	34	XX	31	XX	XX	XX	36	33	XX	35	34	2.8	-.6	6 50 21	-15 10 60
3/28/84	42	34	XX	31	37	XX	XX	39	34	XX	35	43	5.6	-.2	11 20 21	-18 350 28
5/09/79	38	37	XX	32	38	XX	XX	35	35	XX	35	37	XXXX	1.3	6 10 19	-12 340 87

Figure 2 - Printout of Analogous Days Program

FORMULAE

Santa Maria Airport Formula for Minimum Temperature Using 00Z Data

Paso Robles Formula for Minimum Temperature (00Z)

Formula: Min=

Min=

$$\text{DEW POINT} - \frac{\text{HUMIDITY}-30}{3} + V_d + V_h$$

$$\text{DEW POINT} - \frac{\text{HUMIDITY}-25}{3} + V_d + V_h$$

Constants

DEW POINT	V _d	HUMIDITY	V _h	DEW POINT	V _d	HUMIDITY	V _h
7-15.....	.20	13-14.....	-12	10-12.....	.9	10.....	-8
16.....	.19	15.....	-11	13-16.....	.8	11-16.....	-7
17.....	.18	16-17.....	-10	17-19.....	.7	17-19.....	-6
18.....	.17	18.....	-9	20-22.....	.6	20-24.....	-5
19.....	.14	19.....	-8	23-24.....	.5	25-29.....	-4
20.....	.11	20.....	-7	25-27.....	.4	30-34.....	-3
21-22.....	.10	21.....	-6	28-29.....	.3	35-39.....	-2
23.....	.9	22-23.....	-5	30-33.....	.2	40-44.....	-1
24.....	.8	24.....	-4	34.....	.1	45-49.....	0
25.....	.6	25.....	-3	35-37.....	.0	50-54.....	2
26.....	.5	26.....	-2	38-39.....	-.1	55-59.....	3
27.....	.4	27.....	-1	40-42.....	-.2	60-64.....	4
28.....	.3	28-32.....	0	43.....	-.3	65-69.....	5
29-32.....	.2	33-47.....	-1	44-48.....	-.4	70-74.....	6
33-37.....	.1	48-63.....	-2	49.....	-.5	75-79.....	7
38-45.....	.2	64-78.....	-3	50.....	-.6	80-84.....	8
46-48.....	.3	88.....	-4			85-89.....	9
49.....	.4					90-94.....	10
						95-98.....	11
						99-100.....	12

Table 1 - Correction constants for dew point and humidity

KEY STATION STATISTICS

Departure from average minimum temperature at Santa Maria through
11 winter and spring seasons. (Winter, 1982 - Spring, 1993)

WINTER

	LOW CEILING	HIGH CEILING	ALL COLD NIGHTS
BONITA SCHOOL	+0.7	+1.3	+0.9
NIPOMO MESA	-3.8	-2.3	-3.4
NIPOMO FOOTHILL	+0.3	+1.8	+0.7
UPPER ARROYO GRANDE	+5.1	+3.0	+4.3
OCEANO	+0.6	+1.8	+0.9
MORRO GRANGE	+1.9	+0.2	+1.3 (4 YRS)

SPRING

	LOW CEILING	HIGH CEILING	ALL COLD NIGHTS
NIPOMO FOOTHILL	-0.3	-0.6	-0.3
NIPOMO MESA	-4.4	-4.0	-4.3 (9 YRS)
UPPER ARROYO GRANDE	+1.5	+1.0	+1.4
OCEANO	+1.1	+1.0	+1.1
BONITA SCHOOL	+0.7	+0.1	+0.6
SHANDON	-7.3	-7.6	-7.4
ESTRELLA RIVER	-3.3	-2.4	-3.1
RADIKE VINEYARD	-6.6	-5.7	-6.4 (8 YRS)
PRICE CANYON	-2.2	-2.6	-2.3 (9 YRS)
SISQUOC	-2.4	-1.5	-2.2
LOS ALAMOS	-2.4	-2.2	-2.3
ZACA	-0.7	-0.9	-0.7
SANTA YNEZ	-2.2	-1.5	-2.1
MORRO GRANGE	-3.4	-4.0	-3.6 (4 YRS)
PASO ROBLES	-4.0	-2.5	-3.7

- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The EART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196053)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins (Retired) and Alexander R. Smith. Fifth Revision, July 1992. (PB92 220177)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gal M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82 158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82 177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidli, April 1969 (Revised December 1986). (PB87 142063/AS)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteria, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109636)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85 110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85 132694)
- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 187887AS)
- 190 Great Salt Lake Effect Snowfall. Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Lusky, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Lusky, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Radid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
- 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
- 199 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Lusky, April 1987. (PB87 185229/AS)
- 200 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth B. Mielke, September 1987. (PB88 110895/AS)
- 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson and George Clark, September 1987. (PB88 114038/AS)
- 202 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88 101935/AS)
- 203 An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern California. John P. Monteverdi and Scott A. Braun, October 1988. (PB89 121297/AS)
- 204 Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Carven Scott, November 1988. (PB89 128649/AS)
- 205 Forecast Guidelines For Fire Weather and Forecasters - How Nighttime Humidity Affects Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS)
- 206 A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarters, Scientific Services Division, August 1989. (PB89 230833/AS)
- 207 The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, June 1990. (PB90-240268)
- 208 Meteorological Factors Contributing to the Canyon Creek Fire Blowup, September 6 and 7, 1988. David W. Goens, June 1990. (PB90-245085)
- 209 Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow Whitlatch, December 1990. (PB91-129239)
- 210 Hydrotools. Tom Egger, January 1991. (PB91-151787/AS)
- 211 A Northern Utah Soaker. Mark E. Struthwolf, February 1991. (PB91-168716)
- 212 Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 1991. (PB91-208439)
- 213 Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Mollner, July 1991. (PB91-227405/AS)
- 214 Emergency Operational Meteorological Considerations During an Accidental Release of Hazardous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424)
- 215 WeatherTools. Tom Egger, October 1991. (PB93-184950)
- 216 Creating MOS Equations for RAWS Stations Using Digital Model Data. Dennis D. Gettman, December 1991. (PB92-131473/AS)
- 217 Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 1992. (PB92-195104)
- 218 NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 1992. (PB93-146785)
- 219 A Case Study of the Operational Usefulness of the Sharp Workstation in Forecasting a Mesocyclone-Induced Cold Sector Tornado Event in California. John P. Monteverdi, March 1993. (PB93-178697)
- 220 Climate of Pendleton, Oregon. Claudia Bell, August 1993. (PB93-227536)
- 221 Utilization of the Bulk Richardson Number, Helicity and Sounding Modification in the Assessment of the Severe Convective Storms of 3 August 1992. Eric C. Evenson, September 1993. (PB94-131943)
- 222 Convective and Rotational Parameters Associated with Three Tornado Episodes in Northern and Central California. John P. Monteverdi and John Quadros, September 1993. (PB94-131943)
- 223 Climate of San Luis Obispo, California. Gary Ryan, February 1994. (PB94-162062)
- 224 Climate of Wenatchee, Washington. Michael W. McFarland, Roger G. Buckman, and Gregory E. Matzen, March 1994. (PB94-164308)
- 225 Climate of Santa Barbara, California. Gary Ryan, December 1994.
- 226 Climate of Yakima, Washington. Greg DeVoir, David Hogan, and Jay Neher, December 1994.
- 227 Climate of Kalispell, Montana. Chris Maier, December 1994.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS--Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS--Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161