



**NOAA TECHNICAL MEMORANDUM
NWS WR-251**

**A COMPOSITE STUDY EXAMINING FIVE HEAVY
SNOWFALL PATTERNS FOR SOUTH-CENTRAL
MONTANA**

**Jonathan D. Van Ausdall
Thomas W. Humphrey
National Weather Service Office
Billings, MT**

February 1998

**U.S. DEPARTMENT
OF COMMERCE**

National Oceanic and
Atmospheric Administration

National Weather
Service



NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.


Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, 125 South State Street - Rm 1210, Salt Lake City, Utah 84138-1102. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; microfiche are \$3.50. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (Revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (Revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidli, April 1968 (Revised March 1986). (PB86 177672/AS). (Revised October 1991 - PB92-115062/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB 179084)
- 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. (PB 184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 1969. (PB 185068)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)
- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394)
- 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)
- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 65 Climate of Sacramento, California. Richard Honton and Tony Martini (Retired), August 1996. (Fifth Revision) (PB89 207781/AS)
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)
- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)
- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological Charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (Revised March 1978). (COM 73 10251)
- 83 A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 1277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS)
- 96 Map type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christo, Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAN/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 126 Climate of San Francisco. E. Jan Null, February 1978. (Revised by George T. Pericht, April 1988 and January 1995). (PB88 208624/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudgel, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David W. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Moliner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Helfick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric



**NOAA TECHNICAL MEMORANDUM
NWS WR-251**

**A COMPOSITE STUDY EXAMINING FIVE HEAVY
SNOWFALL PATTERNS FOR SOUTH-CENTRAL
MONTANA**


**Jonathan D. Van Ausdall
Thomas W. Humphrey
National Weather Service Office
Billings, MT**

February 1998

UNITED STATES
DEPARTMENT OF COMMERCE
William M. Daley, Secretary

National Oceanic and
Atmospheric Administration
D. James Baker, Under
Secretary and Administrator

National Weather Service
Robert W. Winokur, Acting Assistant
Administrator for Weather Services



**This publication has been reviewed
and is approved for publication by
Scientific Services Division,
Western Region**



**Delain A. Edman, Chief
Scientific Services Division
Salt Lake City, Utah**

TABLE OF CONTENTS

LIST OF FIGURES AND TABLES	iv
ABSTRACT	1
I. INTRODUCTION	1
II. PROCEDURE	2
III. ANALYSIS AND OBSERVATIONS	2
A. Southwest Flow	2
1. All Southwest Flow Events	3
2. "Big" Southwest Flow Events	3
3. "Super" Southwest Flow	3
B. Amplified Southwest Flow	3
1. All Amplified Southwest Flow	3
2. "Big" Amplified Southwest Flow Events	4
3. "Super" Amplified Southwest Flow Events	4
C. West Flow	4
1. All West Flow Events	4
2. "Big" West Flow Events	5
3. "Super" West Flow Events	5
D. Northwest Flow	5
1. All Northwest Flow Events	5
2. "Big" Northwest Flow Events	6
3. "Super" Northwest Flow Events	6
E. Closed Lows	7
1. All Closed Low Events	7
2. "Big" Closed Low Events	7
3. "Super" Closed Low Events	7
F. Snowfall Distribution Statistics	8
1. Amount and Duration of Events	8
2. Snowfall Event Distribution By Month	8
IV. SUMMARY AND CONCLUSIONS	9
ACKNOWLEDGMENTS	10
APPENDIX A - LIST OF EVENTS WITH SNOWFALL AMOUNTS	40

LIST OF FIGURES AND TABLES

FIGURES

- Figure 1: a) 500 mb heights (dm), b) Mean Sea-Level Pressure (MSLP) in mb, and c) 250 mb winds (knots) for composite Southwest Flow cases.
- Figure 2: Topography for Montana and northern Wyoming. The location of Billings is shown. The contour interval is 500 feet.
- Figure 3: a) Standard deviation (STD) of 500 mb heights (m), b) STD of MSLP (mb), c) 250 mb wind STD (knots) for Southwest Flow cases.
- Figure 4: Same as Figure 1, except for "Big" Southwest Flow cases.
- Figure 5: Same as Figure 1, except for a "Super" Southwest Flow case.
- Figure 6: a) 500 mb heights (dm), b) MSLP (mb), and c) 250 mb winds (knots) for Amplified Southwest Flow cases.
- Figure 7: a) 500 mb height STD (m), b) MSLP STD (mb), and c) 250 mb wind STD (knots) for Amplified Southwest Flow cases.
- Figure 8: Same as Figure 6, except for "Super" Amplified Southwest Flow (Case 1). No 250 mb wind data available for this case.
- Figure 9: a) 500 mb heights (dm), b) MSLP (mb), and c) 250 mb winds (knots) for "Super" Amplified Southwest Flow (Case 2).
- Figure 10: a) 500 mb heights (dm), b) MSLP (mb), and c) 250 mb winds (knots) for West Flow cases.
- Figure 11: a) 500 mb height STD (m), b) MSLP STD (mb), and c) 250 mb wind STD (knots) for West Flow cases.
- Figure 12: a) 500 mb heights (dm), b) MSLP (mb), and c) 250 mb winds (knots) for Northwest Flow cases.
- Figure 13: a) 500 mb height STD (m), b) MSLP STD (mb), and c) 250 mb wind STD (knots) for Northwest Flow cases.
- Figure 14: Same as Figure 12, except for "Big" Northwest Flow cases.

- Figure 15: Same as Figure 12, except for "Super" Northwest Flow (Case 1). No 250 mb wind data available for this case.
- Figure 16: Same as Figure 15, except for Case 2.
- Figure 17: a) 500 mb heights (dm), b) MSLP (mb), and c) 250 mb winds (knots) for "Super" Northwest Flow (Case 3).
- Figure 18: Same as Figure 17, except for Case 4.
- Figure 19: a) 500 mb heights (dm), b) MSLP (mb), and c) 250 mb winds (knots) for Closed Low cases.
- Figure 20: a) 500 mb height STD (m), b) MSLP STD (mb), and c) 250 mb wind STD (knots) for Closed Low cases.
- Figure 21: a) 500 mb heights (dm), b) MSLP (mb) for "Super" Closed Low (Case 1). No 250 mb data was available for this case.
- Figure 22: a) 500 mb heights (dm), b) MSLP (mb), and c) 250 mb winds (knots) for "Super" Closed Low (Case 2).
- Figure 23: Snowfall Event Distribution By Month - All Flow Patterns.
- Figure 24a: Same as Figure 23, except for Southwest Flow.
- Figure 24b: Same as Figure 23, except for Amplified Southwest Flow.
- Figure 24c: Same as Figure 23, except for West Flow.
- Figure 24d: Same as Figure 23, except for Northwest Flow.
- Figure 25e: Same as Figure 23, except for Closed Low.

TABLES

- Table 1: Snowfall Distribution statistics for each flow pattern showing number and percentage of "Big" and "Super" events.
- Table 2: Snowfall Distribution statistics for each flow pattern showing the average snowfall (in inches), average length of each event (in days), percentage of one-day events, two-day events, and events longer than two days.

A COMPOSITE STUDY EXAMINING FIVE HEAVY SNOWFALL PATTERNS FOR SOUTH-CENTRAL MONTANA

*Jonathan D. Van Ausdall and Thomas W. Humphrey
National Weather Service - Billings, Montana*

Abstract

This study identifies heavy snow patterns for south-central Montana. A snow amount of 6 inches or greater for each event was considered heavy snow for this study. Data for the years 1948-1994 was compiled using the NCEP Version III Grid Point Data Set. Patterns were examined during the time of heaviest snow, and composites made of 5 distinct flow patterns. The five flow patterns composited were: Southwest Flow, Amplified Southwest Flow, West Flow, Northwest Flow, and Closed Low. Composites were obtained by using the GEMCD program. Fields chosen for the composites were: 500 mb heights, Mean Sea-Level Pressure (MSLP), and 250 mb winds. Data for 250 mb winds was only available since 1965. To isolate the heavier snowfalls, separate composites were performed for single day totals of 5-9 inches. Single day snows of 10+ inches were rare, so they were examined individually.

I. INTRODUCTION

Heavy snows affect large segments of the population in southern Montana. The general public, as well as special interest groups such as ranchers, emergency managers, and transportation officials, can benefit greatly from an accurate snowfall forecast. Prior to this study, no documentation was available to aid the forecaster with identifying heavy snow patterns for southern Montana. This study attempted to diagnose surface and upper-level patterns most commonly linked to heavy snowfall. For this study, heavy snow was defined as 6 inches of snow falling in an event. An event was defined as six inches or more snowfall on a single day, or at least 2 inches of

snowfall on consecutive days totaling 6 inches or more. Data for a 47-year period, 1948-1994, was used to determine heavy snowfall events. The 500 mb heights during the period of heaviest snow were used as a "snapshot" of each event's upper-level flow. Based upon these snapshots, the events were then divided into five distinct upper-level flow patterns. Composites of Mean Sea-Level Pressure (MSLP), 500 mb heights, and 250 mb winds were performed for each of the five upper-level flow patterns. These composites were the focus of this study.

II. PROCEDURE

To determine heavy snowfall events, daily climate data for Billings, Montana was examined for the period 1948-1994. These data were obtained from the World Weatherdisc. Composites were generated using the GEMCD program, developed by Dan St. Jean (NWSFO Boise), which used data from the National Center for Environmental Prediction (NCEP) Grid Point Data Set: Version III CD-ROM. Surface and 500 mb data were available for the period: June 1948 through December 1994. The 250 mb wind data were available from 1965 through 1994.

Before compositing, 500 mb heights for each case were extracted from the NCEP CD. The 500 mb heights were then regrouped into similar flow patterns. Five distinct patterns were identified: Southwest Flow, Amplified Southwest Flow, West Flow, Northwest Flow, and Closed Lows. For each pattern, MSLP, 500 mb heights, and 250 mb winds were composited. Standard deviations were computed to get a measure of the spread of each field for a given flow pattern. To see if events within each flow pattern had a tendency to be "quick-hitters" or long-lived, statistics on the length of events were calculated. Separate composites were made for "Big" events, which were categorized as 5-9 inches on a single day. "Super" events, 10 inches or more on a single day, were rare, so they were not composited.

III. ANALYSIS AND OBSERVATIONS

A. Southwest Flow

1. All Southwest Flow Events (18 Cases)

The Southwest Flow cases were characterized by a split-flow regime over western North America (Fig. 1a). The southern stream trough axis was located near the West Coast, which resulted in broad southwest flow over southern Montana. The northern stream flowed south in western Canada and merged with the southern stream in southern Saskatchewan. The ridge in the northern stream over western Canada provided a source of arctic air for the northern Plains, as evidenced by the 1024+ mb surface high in northwest Canada (Fig. 1b). The location of the arctic high provided northeast winds at the surface over south-central Montana. The Rocky Mountains in Montana have a northwest-southeast orientation through the western half of the state (Fig. 2). Thus, northeast winds provide good upslope for southern Montana. Moist, southwest flow aloft originating off the West Coast resulted in a good overrunning situation (i.e. isentropic lift) over the area. At 250 mb (Fig. 1c), the core of the jet stream (~85 knots) extended from northern Utah into the Dakotas. This configuration would place the left-front quadrant of approaching jet-streaks over southern Montana, enhancing vertical motion. The upslope flow over south-central Montana was further enhanced by surface waves moving through central Wyoming, which

can be inferred from the Region of lower pressure extending from near Salt Lake City to the Nebraska panhandle (Fig. 1b).

Standard deviations at 500 mb (Fig. 3a) are largest in Canada and the West Coast of the U.S., and smallest from eastern Montana southeast into Colorado. The large spread in Canada reflects the varying strength and position of the northern stream ridge, while the differences in the western U.S. stem from the position and amplitude of the southern stream trough. Deviations in MSLP, (Fig. 3b) greatest in Canada, are related to the strength of the arctic high.

At 250 mb (Fig. 3c), the greatest variations, located from the eastern Pacific into the Dakotas, are due to the location and strength of the jet.

2. "Big" Southwest Flow Events (10 Cases)

The composite plots of "Big" Southwest flow cases (Fig. 4a-c), showed some important, though subtle, differences from the composite plots of all Southwest Flow cases. At 500 mb, the northern stream was farther south, which resulted in a slightly stronger arctic high (~2 mb). The southern stream at 500 mb had a tighter gradient, and not surprisingly, a stronger jet at 250 mb. The configuration of the 250 mb jet (Fig. 4c) put southern Montana in an area of divergence aloft, which helped enhance vertical motions.

Standard deviations from the "Big" Southwest Flow cases were similar to the composite of all Southwest Flow cases, in terms of both magnitude and location of minimum and maximum values.

3. "Super" Southwest Flow Events (1 Case)

The one "Super" Southwest flow case was vastly different than the composites of the "Big" cases and the composite of all Southwest Flow cases. In this case, all three fields (Fig. 5a-c) were much stronger than the composites in Sections 1 and 2. At 500 mb (Fig. 5a), there was a very tight gradient in the southern stream, with a 145 knot west-southwest jet (Fig. 5c) at 250 mb. This jet supported a 996 mb surface low which propagated from southwest Montana across northern Wyoming. The arctic high in northwest Canada was not very intense, but the pressure gradient of 20 mb from northeastern to southwestern Montana was more than twice as strong as either of the composites in Sections 1 and 2, which resulted in enhanced upslope flow.

B. Amplified Southwest Flow

1. All Amplified Southwest Flow Events (23 Cases)

The Amplified Southwest Flow cases (Fig. 6a-c) had a sharp trough at 500 mb extending from central Canada to the western Great Basin. This trough was sharper and farther east than the Southwest Flow cases. Also, the distinct split-flow in the previous flow pattern was not as apparent in the Amplified Southwest Flow cases, although there was a hint of a southern stream in the western Great Basin (Fig. 6a). The configuration of the 250 mb jet (Fig. 6c) put southern Montana in an area of divergence aloft. At the surface (Fig. 6b), the arctic high was centered in central Alberta, with the lowest pressures in

eastern Utah and western Colorado. So while the upper-air patterns for the Southwest Flow and Amplified Southwest Flow events were quite different, the surface patterns in the two flow regimes were similar.

Standard deviations at 500 and 250 mb (Figs. 7a&c) were largest near the trough axis, which were related to the position and amplitude of the trough. The largest deviations of MSLP (Fig. 7b) were in Canada, associated with the strength and location of the arctic high.

2. "Big" Amplified Southwest Flow Events (14 Cases)

The "Big" Amplified Southwest Flow composites (not shown) looked nearly identical to the composites done with all cases. The similarity was due in large part to the large number of cases (14), that were included in both sets of composites. Moisture availability was likely higher in the "Big" cases, but there was no means of quantifying moisture in this study. The standard deviations were also similar, with maxima and minima in the same locations. However, the magnitudes were somewhat smaller with the "Big" cases.

3. "Super" Amplified Southwest Flow Events (2 Cases)

The first "Super" case (Fig. 8a-b) had a split-flow pattern at 500 mb. In the southern stream, a sharp 500 mb trough axis was located in central Nevada, with southwest flow from the eastern Great Basin into the Plains. The northern stream had a ridge in the Gulf of Alaska,

with northerly flow over Alberta. The ridge over the northeast Pacific Ocean supported a 1040+ mb surface high in northwest Canada. A low-pressure center was located in central Utah. There were no 250 mb winds available for this case, but based upon the flow at 500 mb, there was likely a jet from Utah into the Dakotas.

Case 2 (Fig. 9a-c) had a split-flow at 500 mb also, but it was weaker than Case 1. The trough axis in the southern stream was located from western Wyoming into western Colorado. Southwest flow extended from Colorado into the eastern Dakotas. The placement of the northern stream ridge and trough were the same as Case 1. At 250 mb (Fig. 9c), there was a 130+ knot jet from Nebraska into eastern North Dakota. At the surface, an arctic high was located in northwest Canada, with a low-pressure center in the Nebraska panhandle. Implied surface winds in this case were northwest in southern Montana, but strong dynamics compensated for the lack of upslope flow.

A stronger jet was associated with both "Super" cases, which meant stronger dynamical forcing. This would account for the heavier snowfall in the "Super" cases.

C. West Flow

1. All West Flow Events (13 Cases)

West Flow cases (Fig. 10a-c) had a split-flow at 500 mb (Fig. 10a). The southern branch was nearly zonal, and confined mostly to the U.S. The northern branch had a ridge in the northeast Pacific, with

northwest flow across Alberta and Saskatchewan. The flow in the northern branch became more zonal in central Canada where the two streams combined. At the surface (Fig. 10b), an arctic high was centered in northwest Canada, with an area of low pressure centered over southern Idaho. The jet (Fig. 10c) at 250 mb extended from northeast Oregon into western North Dakota, and was 85-90 knots. The surface pattern and the location of the jet indicated frontal waves likely helped enhance vertical motion over southern Montana.

Standard deviations (Fig. 11a-c) at 500 mb (Fig. 11a) were largest in Canada, associated with the strength of the northern stream. MSLP standard deviations (Fig. 11b) were fairly uniform, but were largest in Canada and the northern Plains, due to the intensity of the arctic high. The 250 mb standard deviations were on the order of 20 knots, with largest values in the upper Midwest, related to the strength of the confluence of the northern and southern streams.

2. "Big" West Flow Events (7 Cases)

Composites of "Big" West Flow events (not shown) were not significantly different from the composites of all West Flow events. Over half of West Flow events were "Big", which contributed to the similarity of the two composites. The jet was about 10 knots stronger in the "Big" cases, but moisture availability was likely the determining factor in these scenarios.

Standard deviations (not shown) were similar to those in Section 1, in terms of the magnitude and location of minimum and maximum values.

3. "Super" West Flow Events (None)

There were no West Flow events which had more than 10 inches of snow in a single day.

D. Northwest Flow

1. All Northwest Flow Events (19 Cases)

Northwest Flow cases (Fig. 12a-c) were characterized by a Hudson Bay Low at 500 mb (Fig. 12a), with northwest flow over western Canada and the U.S.. A large surface high (1032+ mb) was located in northwest Canada (Fig. 12b). Ridging extended into northeast Montana, which resulted in upslope flow in southern Montana. The jet at 250 mb (Fig. 12c) extended from British Columbia into the central Plains, with a core of 100 knots over central Montana. The southerly ageostrophic flow associated with the exit Region of the jet produced isentropic lift over the arctic boundary which rested along the Rocky Mountains.

Standard deviations (Fig. 13a-c) at 500 mb (Fig. 13a) were smallest along the Rocky Mountains. Largest values were located in central Canada, associated with the strength and location of the Hudson Bay Low, and off the Pacific Northwest coast, due to the amplitude of the ridge. Surface standard deviations

(Fig. 13b) were largest on the cold side of the arctic boundary, attributable to the strength and eastward extent of the arctic high. At 250 mb (Fig. 13c), deviations were greatest over British Columbia and the southern Rockies, and can be related to the strength and orientation of the jet.

2. "Big" Northwest Flow Events (6 Cases)

The "Big" cases showed noticeable differences from the previous events. At 500 mb (Fig. 14a), the Hudson Bay Low was located farther east. The flow into the west coast of North America was more westerly, which would allow greater moisture influx. At the surface (Fig. 14b), pressures were weaker over Montana (~4 mb), with a hint of a surface wave in central Wyoming. The shallower layer of arctic air over southern Montana would result in higher moisture capacity at low- and mid-levels. The low-pressure wave rippling along the arctic front would concentrate the available moisture and enhance snowfall rates. At 250 mb (Fig. 14c), the jet was slightly weaker with more of a westerly orientation.

Standard deviations (not shown) were similar to those in Section 1, in terms of location and magnitude.

3. "Super" Northwest Flow Events (4 Cases)

The four "Super" cases were different enough from each other to prohibit compositing, but did share common elements: a source of cold air and a jet near Montana. Case 1 (Fig. 15a-b) had a strong Hudson Bay Low at 500 mb (Fig.

15a) with northwest flow over Montana, and west flow associated with the southern branch of the jet over Oregon and California. At the surface (Fig. 15b), an area of high pressure was located over northern British Columbia, with a wave of low pressure in western Nebraska. Ridging in northern Montana produced upslope flow in southern Montana. Case 2 (Fig. 16a-b) had a polar vortex over Hudson Bay at 500 mb (Fig. 16a) with northwest flow over Montana and no southern stream. At the surface (Fig. 16b), a large arctic high was over northern Alberta, with the arctic boundary lying against the Rockies. Pressure gradients across Montana were weak, but there was upslope flow in southern Montana. Case 3 (Fig. 17a-c) had low heights and northwest flow over Montana at 500 mb (Fig. 17a). Surface features (Fig. 17b) were similar to Case 1, with a wave of low pressure near the Montana/Wyoming/South Dakota border. Again, ridging into northeast Montana provided upslope flow. Aloft, a 120+ kt jet at 250 mb (Fig. 17c) was entering western Montana, with the favorable left-front quadrant of the jet over southern Montana. The 500 mb flow in Case 4 (Fig. 18a) was similar to Case 3, but the gradient across Montana was stronger. This case had downslope flow (southwest winds) at the surface (Fig. 18b), but a 115 knot jet moving into southern Montana at 250 mb (Fig. 18c), likely minimized these effects.

The surface patterns in each of the "Super" cases were distinctly different, but each case did have a polar vortex in the vicinity of Hudson Bay, which supplied the cold air to southern

Montana. Also, a strong jet at 250 mb was near southern Montana in each case (implied from the 500 mb flow in Cases 1 and 2), which helped produce the necessary lift needed for heavy snow. Upslope flow at the surface was present in three of four cases, which probably served to enhance snowfall, but was not a requirement for generating the snow.

E. Closed Lows

1. All Closed Low Events (21 Cases)

The composite of all Closed Low events (Fig. 19a-c) had a closed circulation at 500 mb (Fig. 19a) near the Idaho/Utah/Wyoming border with southeast flow over southern Montana. The dominant surface feature was a low-pressure area over southeast Colorado (Fig. 19b), with high pressure located in central Canada. Upslope surface flow was not a factor in these cases, which were dynamically driven for the most part. The 250 mb winds (Fig. 19c) were strongest across New Mexico and Arizona, with the nose of the jet entering southeast Colorado, supporting the low-pressure area there. Over southern Montana, the flow was diffluent, and likely divergent as well, enhancing vertical motion.

Standard deviations at 500 mb (Fig. 20a) were uniform over the western U.S. and Canada. Strength of the closed circulation accounted for most of the variability. Deviations at the surface (Fig. 20b) were less than 10 mb across the West. This fairly small variability was the result of the consistent strength and

position of the Canadian high and Colorado low. At 250 mb (Fig. 20c), values were largest in the western and southern U.S., associated with the strength of the jet, and smallest in the deformation zone over central Montana.

2. "Big" Closed Low Events (13 Cases)

Due to the large number of "Big" cases (13, or 62 percent of the total number of cases), the composites and standard deviations of the "Big" cases (not shown) were almost identical to those in Section 1.

3. "Super" Closed Low Events (2 Cases)

Case 1 of the "Super" Closed Lows (Fig. 21a-b) was the biggest snowfall on record in Billings, where 42 inches of snow fell in two days! At 500 mb (Fig. 21a), the center of the closed circulation was located over southeast Wyoming, with diffluent east flow over southern Montana. A surface low-pressure center was over western Nebraska, with an inverted trough extending into southwest Saskatchewan (Fig. 21b). An arctic high was centered over northwest Canada. The flow around the 500 mb low-pressure center allowed for Gulf of Mexico moisture to be advected over the arctic air, which is an extremely rare occurrence for southern Montana. Case 2 (Fig. 22a-c) was a more typical event, with the 500 mb low (Fig. 22a) over southwest Montana. Weak high pressure (~1022 mb) was centered over southern Alberta (Fig. 22b), with the low-pressure center in southeast Colorado, and an inverted

trough extending into eastern Montana. The jet at 250 mb (Fig. 22c) was diving south from the Pacific Northwest into southern Utah, which is farther north and west than the composite in Section 1. A deformation zone was located over southern Montana, enhancing vertical motion.

The 500 mb closed circulations in both "Super" cases were farther north than the composite in Section 1. The northward displacement places the deformation zone over southern Montana, and allows for large amounts of moisture to overrun the arctic air. Though less frequent than in northwest flow, closed lows have a higher snowfall potential than the other four cases due to possible entrainment of Gulf of Mexico moisture, and slow movement of the system. Case 1 shows the result of both these parameters acting together.

F. Snowfall Distribution Statistics

1. Amount and Duration of Events

The total number of events for each flow pattern was distributed fairly evenly (Table 1). Each pattern other than West Flow represented about 20 percent of the total. The percentage of "Big" events was over 50 percent for four of five flow patterns. The exception was Northwest Flow, but that was compensated by a higher percentage of "Super" events. The most significant numbers were the percentage of "Super" events for Northwest and West Flow. No 10-inch one-day snowfalls were observed in West Flow patterns, whereas over 20 percent of Northwest Flow cases were "Super"

events. A possible explanation for the absence of "Super" events in West Flow was that moisture had to travel over several mountain ranges before reaching southern Montana while moisture in Northwest Flow only had to pass over the Canadian Rockies. Also, the powerful jet in Northwest Flow cases meant stronger dynamical forcing, which would lead to greater snowfall rates.

Each flow pattern averaged over 7 inches of snow per event (Table 2), with Northwest Flow and Closed Lows each having average snowfalls over 10 inches. The average length of each event was smallest for Amplified Southwest Flow, and largest for Closed Low. The average snowfall showed a direct relationship with the average length of each event. Taking the ratio of the average snowfall and average length for each event yielded similar values. This showed that the average snow total was determined more by the length of the event, rather than snowfall intensity. Breaking down the length of each event, Closed Low events were the longest lasting, with almost 30 percent of the events more than two days long. Amplified Southwest Flow events were the shortest, with over 65 percent of the cases lasting only one day.

2. Snowfall Event Distribution By Month

The seasonal distribution for all events (Fig. 23), showed a peak in early Spring, with nearly 20 events in March. A minimum in the number of events occurred in February. The "Big" and "Super" events were spread out evenly between October and April.

Southwest Flow (Fig. 24a) events peaked in January, with secondary peaks in March and April. No events were recorded in February throughout the 47-year data set, which is climatologically the driest month in Billings. "Big" events were evenly distributed from October through March, with the only "Super" event occurring in January.

Amplified Southwest Flow (Fig. 24b) events peaked in fall and spring, with only four events during the winter months. The majority of the "Big" and "Super" snows were also in the fall and spring.

West Flow (Fig. 24c) cases were most common in November, December, and March, with none occurring in January. "Big" cases were distributed in a similar manner. The lack of cases in January could be due to a southward displacement of the jet stream.

The vast majority of Northwest Flow (Fig. 24d) cases (17 of 19) occurred during the winter months, with no spring events. This can be explained by the seasonal strength of the polar vortex, which is at its maximum strength during the winter. The distribution of the "Big" and "Super" cases followed a similar pattern.

Closed Low cases (Fig. 24e) showed an opposite signature when compared to Northwest Flow. The peak was in the spring, with a few cases in October. No Closed Low events were recorded from late fall through winter. This correlated well with the climatological occurrence of closed lows. The two "Super" cases occurred in mid to late spring, which could be due to the higher moisture availability during this time.

IV. SUMMARY AND CONCLUSIONS

Heavy snow events in south-central Montana were broken down into five different flow patterns. The identified patterns were: Southwest, Amplified Southwest, West, Northwest and Closed Lows.

In all patterns except Closed Low, upslope flow, jet energy, and overrunning moisture were key ingredients to the heavy snowfall. In the Closed Low events, a slow-moving deformation zone coupled with abundant moisture were the key factors.


Southwest Flow, Amplified Southwest Flow, and West Flow events were fairly evenly distributed from September through May, whereas Northwest Flow and Closed Low cases were mainly winter and spring events, respectively.

Closed Low and Northwest Flow patterns were the biggest snow producers, which was directly related to the length of those events.


Hopefully, these composites will help alert forecasters to potential developing heavy snow patterns, and lead to longer lead times for watches and warnings.

ACKNOWLEDGMENTS


The authors would like to thank Keith Meier (BIL) and Dave Bernhardt (GTF) for their reviews and helpful suggestions regarding this manuscript. Thanks also goes out to Dan St. Jean (BOI) for providing the GEMCD program, which allowed us to easily access and display the data for this study. Finally, we would like to thank Sally Springer (BIL) for her help in producing the tables.



A) 500MB Heights (dm)



B) MSLP (mb)



C) 250MB Wind (knots)

Figure 1 - Southwest Flow Cases







Figure 2 - Topography of Montana and northern Wyoming
Contour Interval = 500 feet



A) 500MB Heights (m)




B) MSLP (mb)




C) 250MB Winds (knots)


Figure 3 - Southwest Flow Standard Deviation



A) 500MB Heights (dm)




B) MSLP (mb)




C) 250MB Winds (knots)


Figure 4 - "Big" Southwest Flow Cases



A) 500MB Heights (dm)




B) MSLP (mb)




C) 250MB Winds (knots)


Figure 5 - "Super" Southwest Flow Case



A) 500MB Heights (dm)




B) MSLP (mb)




C) 250MB Wind (knots)


Figure 6 - Amplified Southwest Flow Cases



A) 500MB Heights (m)




B) MSLP (mb)




C) 250MB Winds (knots)

Figure 7 - Amplified Southwest Flow Standard Deviation




A) 500MB Heights (dm)




B) MSLP (mb)


Figure 8 - "Super" Amplified Southwest Flow (Case 1)



A) 500MB Heights (dm)




B) MSLP (mb)




C) 250MB Winds (knots)


Figure 9 - "Super" Amplified Southwest Flow (Case 2)



A) 500MB Heights (dm)




B) MSLP (mb)




C) 250MB Wind (knots)


Figure 10 – West Flow Cases



A) 500MB Heights (m)




B) MSLP (mb)




C) 250MB Winds (knots)


Figure 11 - West Flow Standard Deviation



A) 500MB Heights (dm)




B) MSLP (mb)




C) 250MB Winds (knots)


Figure 12 - Northwest Flow Cases



A) 500MB Heights (m)




B) MSLP (mb)




C) 250MB Winds (knots)


Figure 13 - Northwest Flow Standard Deviation



A) 500MB Heights (dm)




B) MSLP (mb)




C) 250MB Winds (knots)

Figure 14 - "Big" Northwest Flow Cases




A) 500MB Heights (dm)




B) MSLP (mb)

Figure 15 - "Super" Northwest Flow (Case 1)




A) 500MB Heights (dm)




B) MSLP (mb)


Figure 16 - "Super" Northwest Flow (Case 2)



A) 500MB Heights (dm)




B) MSLP (mb)





C) 250MB Winds (knots)

Figure 17 - "Super" Northwest Flow (Case 3)




A) 500MB Heights (dm)






A) 500MB Heights (dm)




B) MSLP (mb)




C) 250MB Winds (knots)


Figure 19 - Closed Low Cases



A) 500MB Heights (m)




B) MSLP (mb)




C) 250MB Winds (knots)

Figure 20 - Closed Low Standard Deviation




A) 500MB Heights (dm)




B) MSLP (mb)


Figure 21 - "Super" Closed Low (Case 1)



A) 500MB Heights (dm)



B) MSLP (mb)



C) 250MB Winds (knots)

Figure 22 - "Super" Closed Low (Case 2)

Snowfall Distribution -- Table 1					
Flow Pattern	Number of Events	“Big” Events	% “Big” Events	“Super” Events	% “Super” Events
Southwest	18	10	55.6	1	5.6
Amplified SW	23	14	60.9	2	8.7
West	13	7	53.8	0	0
Northwest	19	6	31.6	4	21.1
Closed Low	21	13	61.9	2	9.5

Snowfall Distribution -- Table 2					
Flow Pattern	Average Snowfall (inches)	Avg. Length of Event (days)	Percentage of 1-Day Events	Percentage of 2-Day Events	Percentage of Events >2 Days
Southwest	8.5	1.8	44.4	38.9	16.7
Amplified SW	7.4	1.4	65.2	30.4	4.4
West	7.9	1.8	30.8	61.5	7.7
Northwest	10.7	2.0	15.8	73.7	10.5
Closed Low	12.7	2.2	14.3	57.1	28.6

Snowfall Event Distribution By Month All Flow Patterns




Figure 23 - Snowfall Distribution - All Flow Patterns

Snowfall Event Distribution By Month Southwest Flow




Figure 24a - Snowfall Distribution - Southwest Flow

Snowfall Event Distribution By Month Amplified Southwest Flow




Figure 24b - Snowfall Distribution - Amplified Southwest Flow

Snowfall Distribution By Month West Flow




Figure 24c - Snowfall Distribution - West Flow

Snowfall Event Distribution By Month Northwest Flow




Figure 24d - Snowfall Distribution - Northwest Flow

Snowfall Event Distribution By Month Closed Low




Figure 24e - Snowfall Distribution - Closed Low

APPENDIX A - LIST OF EVENTS AND SNOWFALL AMOUNTS

Southwest Flow Cases			
Date(s) of Event	Snowfall (inches)	"Big" Event	"Super" Event
December 3-4, 1948	8.2	✓	
January 7-8, 1949	8.0	✓	
January 13-16, 1954	13.2	✓	
December 23-24, 1955	7.4	✓	
March 12-13, 1957	8.4	✓	
April, 22-23, 1958	6.1		
April 23-25, 1961	9.7		
April 9-10, 1963	6.5		
March 12-13, 1967	7.6		
January 26-28, 1968	6.8		
October 15-16, 1971	7.4	✓	
January 2-3, 1973	6.1		
January 25-26, 1975	14.0		✓
November 18-19, 1977	11.0	✓	
November 17-19, 1978	10.8	✓	
April 2-4, 1982	6.6		
January 22-23, 1989	10.2	✓	
March 22-23, 1994	6.7	✓	

Amplified Southwest Flow Cases			
Date(s) of Event	Snowfall (inches)	"Big" Event	"Super" Event
February 26-27, 1951	7.2	✓	
March 24, 1954	6.9	✓	
March 2-4, 1955	8.7		
April 1-2, 1956	6.7		
November 1-2, 1957	6.0	✓	
November 15-17, 1958	7.8	✓	
March 9-10, 1962	6.3		
September 7-8, 1962	6.3		
March 22, 1964	10.5		✓
December 6-7, 1971	7.1	✓	
April 17-18, 1972	7.2	✓	
November 26-28, 1975	7.4		
November 19, 1979	6.0	✓	
January 17-18, 1980	7.6	✓	
September 18-19, 1983	7.5		
September 23-24, 1984	6.9	✓	
October 7-8, 1985	6.0	✓	
April 12-13, 1986	7.3		
March 7-9, 1987	6.1	✓	
April 25, 1988	6.6	✓	
October 27-28, 1991	9.6	✓	
January 11-12, 1993	7.1	✓	
November 17, 1994	12.4		✓

West Flow Cases			
Date(s) of Event	Snowfall (inches)	"Big" Event	"Super" Event
February 8-9, 1955	9	✓	
May 2-3, 1956	6.7	✓	
March 4-5, 1957	7.7		
November 14-15, 1961	8.5		
November 22, 1962	6.3	✓	
March 3, 1972	7.5	✓	
December 2-3, 1972	7.6		
December 10-13, 1975	10.7		
November 17-19, 1978	10.8	✓	
March 3-5, 1980	7.5		
March 1-2, 1989	7.5		
March 22-23, 1990	7	✓	
April 28-29, 1990	6.2	✓	

Northwest Flow Cases			
Date(s) of Event	Snowfall (inches)	"Big" Event	"Super" Event
December 16-17, 1955	21.3		✓
February 13-16, 1958	8.1		
Dec 4-7, 1958	11.3		
February 16-18, 1959	9	✓	
November 10-12, 1959	18.8		✓
January 6-8, 1962	10.6	✓	
January 21-22, 1963	6.5		

Northwest Flow Cases			
Date(s) of Event	Snowfall (inches)	"Big" Event	"Super" Event
December 16-17, 1963	13	✓	
February 27-28, 1970	6.1		
December 31, 1971 - January 2, 1972	17.6		✓
October 31 - November 4, 1973	12.8	✓	
December 27-29, 1973	9.3		
February 24-25, 1978	8.4	✓	
December 4, 1978	13.7		✓
December 23, 1984	7.9	✓	
February 3-5, 1988	6.6		
December 19-21, 1989	7.1		
January 16-17, 1994	5.9		

Closed Low Cases			
Date(s) of Event	Snowfall (inches)	"Big" Event	"Super" Event
October 17-20, 1949	8	✓	
March 11-12, 1954	7		
October 23-25, 1954	8.6		
April 2-5, 1955	42.3*		✓
March 1-3, 1966	14.2	✓	
April 29-30, 1967	9	✓	
April 26-28, 1970	19.2	✓	
April 19-20, 1973	9.6		

Closed Low Cases			
Date(s) of Event	Snowfall (inches)	"Big" Event	"Super" Event
March 25-27, 1975	6.8		
April 7-8, 1975	8.7	✓	
March 28-29, 1977	6.7	✓	
October 15-16, 1980	14.7	✓	
March 17-18, 1981	9.3	✓	
May 11-12, 1981	15.6		✓
March 19-20, 1982	11.8	✓	
May 11-12, 1983	11.9	✓	
May 8-9, 1986	8.3	✓	
April 27-28, 1989	6		
April 10-13, 1991	22.7	✓	
October 7-8, 1993	7.8	✓	

* - Largest Billings snowfall on record

- Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 148 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 149 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033)
- 152 Climate of Salt Lake City, Utah. William J. Alder, Sean T. Buchanan, William Cope (Retired), James A. Cisco, Craig C. Schmidt, Alexander R. Smith (Retired), Wilbur E. Figgins (Retired), April 1996 - Sixth Revision (PB96 175583)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, Revised September 1981. (PB82 196783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82-158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82-177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidli and Austin Jamison, April 1969 (Revised July 1996). (PB96-191614)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasternis, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85-110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85-132694)
- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 1878887AS)
- 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Luskky, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Luskky, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Radiol Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)
- 198 Preliminary Lightning Climatology Studies for Idaho. Christopher D. Hill, Carl J. Gorski, and Michael C. Conger, April 1987. (PB87 180196/AS)
- 199 Heavy Rains and Flooding in Montana: A Case for Slantwise Convection. Glenn R. Luskky, April 1987. (PB87 185229/AS)
- 200 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1986. Roger L. Cross and Kenneth-B. Mielke, September 1987. (PB88 110895/AS)
- 201 An Inexpensive Solution for the Mass Distribution of Satellite Images. Glen W. Sampson and George Clark, September 1987. (PB88 114038/AS)
- 202 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1987. Roger L. Cross and Kenneth B. Mielke, September 1988. (PB88-101935/AS)
- 203 An Investigation of the 24 September 1986 "Cold Sector" Tornado Outbreak in Northern California. John P. Monteverdi and Scott A. Braun, October 1988. (PB89 121297/AS)
- 204 Preliminary Analysis of Cloud-To-Ground Lightning in the Vicinity of the Nevada Test Site. Carven Scott, November 1988. (PB89 128649/AS)
- 205 Forecast Guidelines for Fire Weather and Forecasters - How Nighttime Humidity Affects Wildland Fuels. David W. Goens, February 1989. (PB89 162549/AS)
- 206 A Collection of Papers Related to Heavy Precipitation Forecasting. Western Region Headquarters, Scientific Services Division, August 1989. (PB89 230833/AS)
- 207 The Las Vegas McCarran International Airport Microburst of August 8, 1989. Carven A. Scott, June 1990. (PB90-240268)
- 208 Meteorological Factors Contributing to the Canyon Creek Fire Blowup, September 6 and 7, 1988. David W. Goens, June 1990. (PB90-245085)
- 209 Stratus Surge Prediction Along the Central California Coast. Peter Felsch and Woodrow Whitlatch, December 1990. (PB91-129239)
- 210 Hydrotools. Tom Egger, January 1991. (PB91-151787/AS)
- 211 A Northern Utah Soaker. Mark E. Struthwolf, February 1991. (PB91-168716)
- 212 Preliminary Analysis of the San Francisco Rainfall Record: 1849-1990. Jan Null, May 1991. (PB91-208439)
- 213 Idaho Zone Preformat, Temperature Guidance, and Verification. Mark A. Molner, July 1991. (PB91-227405/AS)
- 214 Emergency Operational Meteorological Considerations During an Accidental Release of Hazardous Chemicals. Peter Mueller and Jerry Galt, August 1991. (PB91-235424)
- 215 WeatherTools. Tom Egger, October 1991. (PB93-184950)
- 216 Creating MOS Equations for RAWs Stations Using Digital Model Data. Dennis D. Gettman, December 1991. (PB92-131473/AS)
- 217 Forecasting Heavy Snow Events in Missoula, Montana. Mike Richmond, May 1992. (PB92-196104)
- 218 NWS Winter Weather Workshop in Portland, Oregon. Various Authors, December 1992. (PB93-146785)
- 219 A Case Study of the Operational Usefulness of the Sharp Workstation in Forecasting a Mesocyclone-Induced Cold Sector Tornado Event in California. John P. Monteverdi, March 1993. (PB93-178697)
- 220 Climate of Pendleton, Oregon. Claudia Bell, August 1993. (PB93-227536)
- 221 Utilization of the Bulk Richardson Number, Helicity and Sounding Modification in the Assessment of the Severe Convective Storms of 3 August 1992. Eric C. Evenson, September 1993. (PB94-131943)
- 222 Convective and Rotational Parameters Associated with Three Tornado Episodes in Northern and Central California. John P. Monteverdi and John Quadros, September 1993. (PB94-131943)
- 223 Climate of San Luis Obispo, California. Gary Ryan, February 1994. (PB94-162062)
- 224 Climate of Wenatchee, Washington. Michael W. McFarland, Roger G. Buckman, and Gregory E. Matzen, March 1994. (PB94-164308)
- 225 Climate of Santa Barbara, California. Gary Ryan, December 1994. (PB95-173720)
- 226 Climate of Yakima, Washington. Greg DeVoir, David Hogan, and Jay Neher, December 1994. (PB95-173688)
- 227 Climate of Kalispell, Montana. Chris Maier, December 1994. (PB95-169488)
- 228 Forecasting Minimum Temperatures in the Santa Maria Agricultural District. Wilfred Pi and Peter Felsch, December 1994. (PB95-171088)
- 229 The 10 February 1994 Oroville Tornado--A Case Study. Mike Staudenmaier, Jr., April 1995. (PB95-241873)
- 230 Santa Ana Winds and the Fire Outbreak of Fall 1993. Ivory Small, June 1995. (PB95-241865)
- 231 Washington State Tornadoes. Tresté Huse, July 1995. (PB96-107024)
- 232 Fog Climatology at Spokane, Washington. Paul Frisbie, July 1995. (PB96-106604)
- 233 Storm Relative Isentropic Motion Associated with Cold Fronts in Northern Utah. Kevin B. Baker, Kathleen A. Hadley, and Lawrence B. Dunn, July 1995. (PB96-106596)
- 234 Some Climatological and Synoptic Aspects of Severe Weather Development in the Northwestern United States. Eric C. Evenson and Robert H. Johns, October 1995. (PB96-112958)
- 235 Climate of Las Vegas, Nevada. Paul H. Skrbac and Scott Cordero, December 1995. (PB96-135553)
- 236 Climate of Astoria, Oregon. Mark A. McInerney, January 1996.
- 237 The 6 July 1995 Severe Weather Events in the Northwestern United States: Recent Examples of SSWEs. Eric C. Evenson, April 1996.
- 238 Significant Weather Patterns Affecting West Central Montana. Joe Lester, May 1996. (PB96-178751)
- 239 Climate of Portland, Oregon. Clinton C. D. Rockey, May 1996. (PB96-17603)
- 240 Downslope Winds of Santa Barbara, CA. Gary Ryan, July 1996. (PB96-191697)
- 241 Operational Applications of the Real-time National Lightning Detection Network Data at the NWSO Tucson, AZ. Darren McCollum, David Bright, Jim Meyer, and John Glueck, September 1996. (PB97-108450)
- 242 Climate of Pocatello, Idaho. Joe Heim, October 1996. (PB97-114540)
- 243 Climate of Great Falls, Montana. Matt Jackson and D. C. Williamson, December 1996. (PB97-126684)
- 244 WSR-88D VAD Wind Profile Data Influenced by Bird Migration over the Southwest United States. Jesus A. Haro, January 1997. (PB97-135263)
- 245 Climatology of Cape for Eastern Montana and Northern Wyoming. Heath Hockenberry and Keith Meier, January 1997. (PB97-133425)
- 246 A Western Region Guide to the Eta-29 Model. Mike Staudenmaier, Jr., March 1997. (PB97-144075)
- 247 The Northeast Nevada Climate Book. Edwin C. Clark, March 1997. (First Revision - January 1998 - Andrew S. Gorelow and Edwin C. Clark) (PB97-152789)
- 248 Climate of Eugene, Oregon. Clinton C. D. Rockey, April 1997. (PB97-155303)
- 249 Climate of Tucson, Arizona. John R. Glueck, October 1997.
- 250 Northwest Oregon Daily Extremes and Normals. Clinton C. D. Rockey, October 1997.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications.

PROFESSIONAL PAPERS--Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS--- Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS--Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS -- Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS--Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS--Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

NATIONAL TECHNICAL INFORMATION SERVICE

U. S. DEPARTMENT OF COMMERCE

5285 PORT ROYAL ROAD

SPRINGFIELD, VA 22161