

only copy

NOAA TM NWS WR 72

A UNITED STATES
DEPARTMENT OF
COMMERCE
PUBLICATION



NOAA Technical Memorandum NWS WR72

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Weather Service

A Paradox Principle in the Prediction of Precipitation Type

THOMAS J. WEITZ

*Regional Office Library,
U. S. Weather Bureau
Salt Lake City, Utah*

Western Region

SALT LAKE CITY,
UTAH

February 1972

*M(25.5)
WB-WR
1972-F*

NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 23 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS.

Papers 1 to 23, except for 5 (revised edition) and 10, are available from the National Weather Service Western Region, Scientific Services Division, P. O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84111. Papers 5 (revised edition), 10, and all others beginning with 24 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Bldg., 5285 Port Royal Road, Springfield, Va. 22151. Price: \$3.00 paper copy; \$0.95 microfiche. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda

- WRTM 1 Some Notes on Probability Forecasting. Edward D. Diemer, September 1965. (Out of print.)
WRTM 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
WRTM 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965 to February 20, 1966. Edward D. Diemer, March 1966.
WRTM 4 Use of Meteorological Satellite Data. March 1966.
WRTM 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (revised November 1967, October 1969). (PB-178000)
WRTM 6 Improvement of Forecast Wording and Format. C. L. Glenn, May 1966.
WRTM 7 Final Report on Precipitation Probability Test Programs. Edward D. Diemer, May 1966.
WRTM 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (revised January 1967). (Out of print.)
WRTM 9 A Collection of Papers Related to the 1966 NMC Primitive-Equation Model. June 1966.
WRTM 10 Sonic Boom. Loren Crow (6th Weather Wing, USAF, Pamphlet), June 1966. (Out of print.) (AD-479366)
WRTM 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
WRTM 12 A Comparison of Fog Incidence at Missoula, Montana, with Surrounding Locations. Richard A. Dightman, August 1966. (Out of print.)
WRTM 13 A Collection of Technical Attachments on the 1966 NMC Primitive-Equation Model. Leonard W. Snellman, August 1966. (Out of print.)
WRTM 14 Application of Net Radiometer Measurements to Short-Range Fog and Stratus Forecasting at Los Angeles. Frederick Thomas, September 1966.
WRTM 15 The Use of the Mean as an Estimate of "Normal" Precipitation in an Arid Region. Paul C. Kangieser, November 1966.
WRTM 16 Some Notes on Acclimatization in Man. Edited by Leonard W. Snellman, November 1966.
WRTM 17 A Digitalized Summary of Radar Echoes Within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
WRTM 18 Limitations of Selected Meteorological Data. December 1966.
WRTM 19 A Grid Method for Estimating Precipitation Amounts by Using the WSR-57 Radar. R. Granger, December 1966. (Out of print.)
WRTM 20 Transmitting Radar Echo Locations to Local Fire Control Agencies for Lightning Fire Detection. Robert R. Peterson, March 1967. (Out of print.)
WRTM 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparans, April 1967.
WRTM 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.
WRTM 23 "K" Chart Applications to Thunderstorm Forecasts Over the Western United States. Richard E. Hambidge, May 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- WBTM 24 Historical and Climatological Study of Grinnell Glacier, Montana. Richard A. Dightman, July 1967. (PB-178071)
WBTM 25 Verification of Operational Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
WBTM 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
WBTM 27 Objective Minimum Temperature Forecasting for Helena, Montana. D. E. Olsen, February 1968. (PB-177827)
WBTM 28 Weather Extremes. R. J. Schmidli, April 1968 (revised July 1968). (PB-178928)
WBTM 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB-178425)
WBTM 30 Numerical Weather Prediction and Synoptic Meteorology. Capt. Thomas D. Murphy, U.S.A.F., May 1968. (AD-673365)
WBTM 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB-179084)
WBTM 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB-179289)
WBTM 33 Objective Forecasting. Philip Williams, Jr., August 1968. (AD-680425)
WBTM 34 The WSR-57 Radar Program at Missoula, Montana. R. Granger, October 1968. (PB-180292)
WBTM 35 Joint ESSA/FAA ARTC Radar Weather Surveillance Program. Herbert P. Benner and DeVon B. Smith, December 1968 (revised June 1970). (AD-681857)
WBTM 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (Out of print.) (PB-183055)
WBTM 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB-183057)
WBTM 38 Climate of Phoenix, Arizona. R. J. Schmidli, P. C. Kangieser, and R. S. Ingram, April 1969. (Out of print.) (PB-184295)
WBTM 39 Upper-Air Lows Over Northwestern United States. A. L. Jacobson, April 1969. (PB-184296)
WBTM 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L. W. Snellman, August 1969. (PB-185068)
WBTM 41 High Resolution Radiosonde Observations. W. S. Johnson, August 1969. (PB-185673)
WBTM 42 Analysis of the Southern California Santa Ana of January 15-17, 1966. Barry B. Aronovitch, August 1969. (PB-185670)
WBTM 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB-185762)
WBTM 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB-187763)
WBTM 45/1 Precipitation Probabilities in the Western Region Associated with Winter 500-mb Map Types. Richard A. Augulis, December 1969. (PB-188248)

U. S. DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL WEATHER SERVICE

NOAA Technical Memorandum NWSTM WR-72

A PARADOX PRINCIPLE IN THE PREDICTION OF PRECIPITATION TYPE

Thomas J. Weitz
Weather Service Office
Walla Walla, Washington



WESTERN REGION
TECHNICAL MEMORANDUM NO. 72

SALT LAKE CITY, UTAH
JANUARY 1972

TABLE OF CONTENTS

	<u>Page</u>
Abstract	1
I. Introduction	1-2
II. The Basis of the "Paradox Principle"	2
III. Data Investigated	2-3
IV. Evolution of a Type Index	3-4
V. Comparison of Predictors	4
VI. Testing on Independent Data	4
VII. Application in Forecasting	4-5
VIII. Summary	5
IX. Acknowledgment	5-6
X. References	6

LIST OF FIGURES AND TABLES

	<u>Page</u>
Figure 1. Illustration of the "Paradox Principle"	7
Figure 2. Relation of Rain Cases and Snow Cases to Thickness and Temperature	8
Figure 3. A Sample of Graph of the Y-Index	9
Table 1. Summary of Developmental Data	10
Table 2. Testing of Independent Data	10
Table 3. Correction to be Applied to 18,000-Foot Temperature to Approximate 500-mb Temperature	11

A PARADOX PRINCIPLE IN THE PREDICTION OF PRECIPITATION TYPE

ABSTRACT

Utility of 1000 - 500-mb thickness in forecasting precipitation type can be enhanced significantly by also considering the mean lapse rate in that layer. This can be done readily by relating thickness to 500-mb temperature. The 500-mb temperature is a relatively easy parameter to forecast in comparison to the uncertainty of temperature forecasts for lower levels. The paradox is developed that inclusion of 500-mb temperature produces a diametrically opposite effect to that which might be expected.

I. INTRODUCTION

Precipitation forecasting is more than a question of occurrence or nonoccurrence as has been the object of most studies. Type of precipitation has received much less attention. This may be attributable to precipitation forecast verification systems which are concerned only with amount and totally ignore type. Precipitation type is of great practical importance. One inch of rainfall may be little more than a wet inconvenience; the same amount of precipitation falling as snow may paralyze a community with a blanket of a foot or more. The need for further investigation on this problem has been pointed out by the National Weather Service's Western Region Scientific Services Division. In discussing uses for thickness forecasts by teletype head "FOUS" the comment is made, "Examination of relationships between thickness and rain versus snow occurrence is needed" [1].

The type of precipitation occurring in a given air mass is closely related to the thickness, or mean temperature, of a shallow layer near the surface such as the 1000 - 850-mb layer. It is less closely related to the thickness of the deeper 1000 - 500-mb layer. The thickness of a shallow layer is, however, much more difficult to forecast than that of a deep layer. Also there are very few objective forecasts available for such shallow layers, and what little is available or can be derived does not have the necessary precision for successful precipitation type forecasting.

In contrast, there is an abundance of facsimile prognostic material for the 500-mb level available, as well as computer produced 1000 - 500-mb thickness values available on teletype on "FOUS" transmissions. This deeper layer is by far the most practical one for use in forecasting precipitation type. This thickness can be a valuable forecasting tool in the important rain versus snow decision, especially if the mean lapse rate of the layer is also considered.

Thickness, or mean temperature, of the 1000 - 500-mb layer has been the subject of numerous investigations, ably summarized by Penn [2]. One of the more comprehensive of these studies was made by Wagner [3]. Wagner's charts indicate an equal probability of rain or snow, for a 1000 - 500-mb thickness of approximately 17,400 feet (5304 meters) for Walla Walla, Washington. Wagner's figure is in close agreement with this investigation which determined a critical value of 5323 meters for thickness considered alone.

II. THE BASIS OF THE "PARADOX PRINCIPLE"

The concept of warmer air at 500-mb favoring the production of snow and colder air favoring rain may appear preposterous without further examination. Nevertheless, the logic of the concept can be readily demonstrated as illustrated in Figure 1.

Shown are two sample vertical temperature soundings between 1000 and 500 mb. A constant lapse rate is assumed in each case for simplicity. The solid line, representing the snow case, S,S', has a small lapse rate with -20C at 500 mb and 0C at 1000 mb. The dashed line, representing the rain case, R,R', has a steeper lapse rate with -25C at 500 mb and +5C at 1000 mb.

The mean temperature through the layer is the same for each sounding, that is, the 1000 - 500-mb thicknesses are the same in each case. Note that the snow case has a higher temperature at 500 mb than does the rain case.

The principle is therefore proposed that, for a given thickness value, the warmer the air at 500 mb the greater the probability of precipitation occurring in the form of snow rather than rain, and vice versa. This is the "paradox principle".

With a major part of the forecaster's attention being focused on the 500-mb chart and its temperature pattern, this may be somewhat surprising.

III. DATA INVESTIGATED

Data required for investigations of this hypothesis were the 1000 - 500-mb thickness, the 500-mb temperature, and associated precipitation type. In this study 500-mb heights and temperatures for 1200 GCT were extracted from printed "Daily Weather Maps" published by the National Weather Service. One thousand-millibar heights were derived from Walla Walla station pressures. Since Walla Walla's barometer elevation of 976 feet msl averages quite close to the 1000-mb level, a straight-line conversion was used based on the U. S. Standard Atmosphere as given in Smithsonian Meteorological Tables [4].

Developmental data were taken from the months of December, January, and February for the winters 1968-69 and 1969-70. Cases included

were all those which had a measurable amount of precipitation for the 12-hour period centered on 1200 GCT, that is, from 0600 GCT to 1800 GCT. Cases of both rain and snow, or of sleet, were not included. Freezing rain was classified as "rain". A total of 61 cases was obtained consisting of 25 cases of snow and 36 cases of rain.

IV. EVOLUTION OF A TYPE INDEX

A pictorial representation of the joint relationship among the 1000 - 500-mb thickness, 500-mb temperature, and the type of precipitation (rain or snow) is shown in Figure 2. Symbols are conventional with dots for rain cases and asterisks for snow. A line of "best separation" between rain and snow might easily be drawn on this chart by "eye", but it was felt that a more stable solution would result by using statistical technique to determine the line of "best separation"; in this case, a linear discriminant analysis was made. Such an analysis [5], resulted in the following discriminant function:

$$L = 30.7514 + .0567T - .0055K$$

where T = 500-mb temperature in degrees Celsius

k = 1000 - 500-mb thickness in meters.

For convenience of calculation, this function was converted to the following by dividing through by .0055, changing signs and setting the result equal to Y:

$$Y = - \frac{L}{.0055} = K - 10.3T - 5591.$$

Zero values of the discriminant function or precipitation type index, Y, are represented by the solid slanting line in Figure 2. Y will have positive values at points above the line in the "rain" area, and negative values below the line in the "snow" area. Note that the slope of the Y = 0 line clearly substantiates the "paradox principle".

Similar functions were computed for each of the variables alone which resulted in the following discriminant values:

$$T = - 26.02 \text{ } ^\circ\text{C}$$

$$K = 5323 \text{ m.}$$

These values are represented in Figure 2 by the vertical and horizontal dashed lines.

The general equation for the type index, Y, would be:

$$Y = K - aT - C.$$

This equation appears useful at other locations, provided the local values of the constants a and C are determined by empirical investigation.

V. COMPARISON OF PREDICTORS

The relative accuracy of the three predictors, Y (type index as described above), K (thickness alone), and T (temperature alone), were compared using the critical value of each which produced the optimum relationship to precipitation type. Considering each indication as a "forecast", results are shown in Table 1.

The 1000 - 500-mb thickness is a fair discriminator between rain and snow, and is certainly better than the 500-mb temperature. Due to the "paradox principle", however, combining 500-mb temperature with thickness results in an index which shows significant improvement over thickness alone.

VI. TESTING ON INDEPENDENT DATA

Critical values of Y, K, and T as determined from the two developmental winters were tested on independent data from the winter 1970-71. Data were extracted in the same manner as for the developmental winters. Results are shown in Table 2.

This 1970-71 winter was abnormally dry and mild, affording only 17 cases for testing. Although the predictor Y showed a gratifying high accuracy (only one erroneous forecast), this was undoubtedly higher than could normally be expected. Still, it showed results superior to the other two predictors. The overall record for the predictor Y for all three winters, two of developmental data and one of independent testing, shows an accuracy of 88.4%, which may be considered a more representative evaluation.

VII. APPLICATION IN FORECASTING

The Y value may be computed for each 12-hour interval for which radiosonde data are available and plotted on a running graph. For locations not near a radiosonde station, the 500-mb height and temperature may be extracted from facsimile charts. The 1000-mb height may be computed from station pressure at locations suitably near that level, or it may be taken from sea-level pressure using the formula:

$$H = 8(P - 1000)$$

where H = 1000-mb height in meters and

P = sea-level pressure in millibars.

(The precise factor according to Smithsonian Tables is 8.0008 geopotential meters per millibar change in pressure, at 0 C and 1000 mb) [4].

With thickness forecasts available twice daily by teletype (headed "FOUS"), as well as 18,000-foot temperatures on the FD forecasts, the Y value may be extrapolated. Using 24-hour thickness forecasts on FOUS and the third section of the FD forecasts, the Y values may be projected for 24 hours. The nighttime transmissions will produce Y projections for 0000 GCT the following afternoon, and the midday transmissions the Y value for 1200 GCT the following morning.

Stations not located at grid points for which FOUS and FD forecasts are made may effectively employ interpolation methods because of the usual smoothness of values at the 500-mb level. Temperature at the 500-mb level may not, of course, be identical with that at 18,000 feet (5486 meters) due to variations in the 500-mb surface. Compensation for this can be made by noting the prognostic 500-mb height and applying an appropriate temperature correction. This correction involves raising the 18,000-foot temperature when the 500-mb height is less than 18,000 feet (5486 meters) or lowering it when the 500-mb height is above that level. The standard tropospheric lapse rate according to Smithsonian Meteorological Tables [4] is 0.0065 degrees Celsius per meter, or one degree Celsius for each 154 meters (to whole meters). Correction values are given in Table 3.

Figure 3 illustrates a sample Y chart with values determined each 12 hours. Twenty-four hour projections based on the FOUS and FD forecasts are plotted as small Xs.

Projections of Y for more extended periods based on facsimile prognostic charts have thus far met with only limited success. Although positions of major features are usually rather good on these charts, absolute heights are frequently inaccurate. This may be due to sparseness of data over the ocean as most inaccuracies are most marked over the western United States.

VIII. SUMMARY

The Y value based on the "Paradox Principle" appears to be a substantial improvement over use of thickness alone in the important distinction between rain and snow. It is readily computed from parameters for which computer forecasts are now available. It appears suitable for use at locations other than Walla Walla, Washington, provided local values of the constants are determined by empirical investigation.

IX. ACKNOWLEDGMENT

The author is indebted to Mr. Woodrow W. Dickey, Scientific Services Division, National Weather Service Western Region, for his review,

helpful suggestions, and analyses, and to Mr. Leonard W. Snellman, Chief Scientific Services Division, National Weather Service Western Region, for his interest and encouragement. Their assistance is gratefully acknowledged.

X. REFERENCES

- [1] National Weather Service, Western Region Technical Attachment No. 69-45, November 10, 1969.
- [2] Penn, Samuel, 1957: The Prediction of Snow vs. Rain. Forecasting Guide No. 2, U. S. Department of Commerce, Weather Bureau.
- [3] Wagner, A. James, 1957: Mean Temperature from 1000 Millibars to 500 Millibars as a Predictor of Precipitation Type. Bulletin of the American Meteorological Society No. 38.
- [4] Smithsonian Meteorological Tables, Sixth Revised Edition, Smithsonian Institution, 1951.
- [5] Panofsky, H. A., and Brier, C. W. : Some Applications of Statistics to Meteorology. The Pennsylvania State University, 1958.




FIGURE 1. ILLUSTRATION OF THE "PARADOX PRINCIPLE".




FIGURE 2. RELATIONSHIP OF RAIN (•) AND SNOW (*) OCCURRENCES TO 1000 - 500-MB THICKNESS (K) AND 500-MB TEMPERATURE (T).




FIGURE 3. A SAMPLE GRAPH OF THE Y-INDEX. THE SOLID LINE IS A RUNNING GRAPH OF ACTUAL VALUES AT 12-HOUR INTERVALS; "Xs" ARE 24-HOUR PROJECTIONS.

TABLE 1

SUMMARY OF DEVELOPMENTAL DATA

CATEGORICAL "BEST FIT" FORECASTS	Y	K	T
Rain forecast, rain occurred	33	28	26
Rain forecast, snow occurred	5	6	10
Snow forecast, rain occurred	3	8	10
Snow forecast, snow occurred	20	19	15
Total cases correct	53	47	41
Total cases incorrect	8	14	20
Percent correct	86.9	77.0	67.2

TABLE 2

TESTING ON INDEPENDENT DATA

CATEGORICAL "BEST FIT" FORECASTS	Y	K	T
Rain forecast, rain occurred	11	9	8
Rain forecast, snow occurred	0	0	2
Snow forecast, rain occurred	1	3	4
Snow forecast, snow occurred	5	5	3
Total cases correct	16	14	11
Total cases incorrect	1	3	6
Percent correct	94.1	82.3	64.7

TABLE 3

CORRECTION TO BE APPLIED TO THE TEMPERATURE AT
18,000 FEET TO APPROXIMATE THE CORRESPONDING TEMPERATURE AT 500 MB

<u>500-MB HEIGHT IN DECAMETERS</u>	<u>CORRECTION DEGREES C.</u>
495 to 510	+3
511 to 525	+2
526 to 540	+1
541 to 556	0
557 to 571	-1
572 to 587	-2
588 to 602	-3

Western Region Technical Memoranda: (Continued)

- No. 45/2 Precipitation Probabilities in the Western Region Associated with Spring 500-mb Map Types. Richard P. Augulis. January 1970. (PB-189434)
- No. 45/3 Precipitation Probabilities in the Western Region Associated with Summer 500-mb Map Types. Richard P. Augulis. January 1970. (PB-189414)
- No. 45/4 Precipitation Probabilities in the Western Region Associated with Fall 500-mb Map Types. Richard P. Augulis. January 1970. (PB-189435)
- No. 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates. December 1969. (PB-190476)
- No. 47 Statistical Analysis as a Flood Routing Tool. Robert J. C. Burnash. December 1969. (PB-188744)
- No. 48 Tsunami. Richard A. Augulis. February 1970. (PB-190157)
- No. 49 Predicting Precipitation Type. Robert J. C. Burnash and Floyd E. Hug. March 1970. (PB-190962)
- No. 50 Statistical Report of Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona 1969. Wayne S. Johnson. April 1970. (PB-191743)
- No. 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell. July 1970. (PB-193102)
- No. 52 Sacramento Weather Radar Climatology. R. G. Pappas and C. M. Veliquette. July 1970. (PB-193347)
- No. 53 Experimental Air Quality Forecasts in the Sacramento Valley. Norman S. Benes. August 1970. (PB-194128)
- No. 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch. August 1970.
- No. 55 Application of the SSARR Model to a Basin Without Discharge Record. Vail Schermerhorn and Donald W. Kuehl. August 1970. (PB-194394).
- No. 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck. September 1970. (PB-194389)
- No. 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote. September 1970. (PB-194710)
- No. 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson. October 1970. (COM-71-00017)
- No. 59 Application of P.E. Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman. October 1970. (COM-71-00016)

NOAA Technical Memoranda NWS

- No. 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon. Arthur W. Fritz, October 1970. (COM-71-00120)
- No. 61 Relationship of Wind Velocity and Stability to SO₂ Concentrations at Salt Lake City, Utah. Werner J. Heck, January 1971. (COM-71-00232)
- No. 62 Forecasting the Catalina Eddy. Arthur L. Eichelberger, February 1971. (COM-71-00223)
- No. 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner. February 1971. (COM-71-00349)
- No. 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- No. 65 Climate of Sacramento, California. Wilbur E. Figgins, June 1971. (COM-71-00764)
- No. 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM-71-00829)
- No. 67 Precipitation Detection Probabilities by Los Angeles ARTC Radars. Dennis E. Ronne, July 1971.
- No. 68 A Survey of Marine Weather Requirements. Herbert P. Benner, July 1971. (COM-71-00889)
- No. 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM-71-00956)
- No. 70 Predicting Inversion Depths and Temperature Influences in the Helena Valley. David E. Olsen, October 1971.
- No. 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972.