

only copy

A UNITED STATES
DEPARTMENT OF
COMMERCE
PUBLICATION



NOAA Technical Memorandum NWS WR 73

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Weather Service

A Synoptic Climatology for Snow- storms in Northwestern Nevada

BERT L. NELSON, PAUL M. FRANSIOLI AND
CLARENCE M. SAKAMOTO

Western Region

SALT LAKE CITY,
UTAH

February 1972

NOAA TECHNICAL MEMORANDA
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 23 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS.

Papers 1 to 23, except for 5 (revised edition) and 10, are available from the National Weather Service Western Region, Scientific Services Division, P. O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84111. Papers 5 (revised edition), 10, and all others beginning with 24 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Bldg., 5285 Port Royal Road, Springfield, Va. 22151. Price: \$3.00 paper copy; \$0.95 microfiche. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda

- WRTM 1 Some Notes on Probability Forecasting. Edward D. Diemer, September 1965. (Out of print.)
WRTM 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
WRTM 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965 to February 20, 1966. Edward D. Diemer, March 1966.
WRTM 4 Use of Meteorological Satellite Data. March 1966.
WRTM 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (revised November 1967, October 1969). (PB-178000)
WRTM 6 Improvement of Forecast Wording and Format. C. L. Glenn, May 1966.
WRTM 7 Final Report on Precipitation Probability Test Programs. Edward D. Diemer, May 1966.
WRTM 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (revised January 1967). (Out of print.)
WRTM 9 A Collection of Papers Related to the 1966 NMC Primitive-Equation Model. June 1966.
WRTM 10 Sonic Boom. Loren Crow (6th Weather Wing, USAF, Pamphlet), June 1966. (Out of print.) (AD-479366)
WRTM 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
WRTM 12 A Comparison of Fog Incidence at Missoula, Montana, with Surrounding Locations. Richard A. Dightman, August 1966. (Out of print.)
WRTM 13 A Collection of Technical Attachments on the 1966 NMC Primitive-Equation Model. Leonard W. Snellman, August 1966. (Out of print.)
WRTM 14 Application of Net Radiometer Measurements to Short-Range Fog and Stratus Forecasting at Los Angeles. Frederick Thomas, September 1966.
WRTM 15 The Use of the Mean as an Estimate of "Normal" Precipitation in an Arid Region. Paul C. Kangieser, November 1966.
WRTM 16 Some Notes on Acclimatization in Man. Edited by Leonard W. Snellman, November 1966.
WRTM 17 A Digitalized Summary of Radar Echoes Within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
WRTM 18 Limitations of Selected Meteorological Data. December 1966.
WRTM 19 A Grid Method for Estimating Precipitation Amounts by Using the WSR-57 Radar. R. Granger, December 1966. (Out of print.)
WRTM 20 Transmitting Radar Echo Locations to Local Fire Control Agencies for Lightning Fire Detection. Robert R. Peterson, March 1967. (Out of print.)
WRTM 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparans, April 1967.
WRTM 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.
WRTM 23 "K" Chart Applications to Thunderstorm Forecasts Over the Western United States. Richard E. Hambidge, May 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- WBTM 24 Historical and Climatological Study of Grinnell Glacier, Montana. Richard A. Dightman, July 1967. (PB-178071)
WBTM 25 Verification of Operational Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
WBTM 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
WBTM 27 Objective Minimum Temperature Forecasting for Helena, Montana. D. E. Olsen, February 1968. (PB-177827)
WBTM 28 Weather Extremes. R. J. Schmidli, April 1968 (revised July 1968). (PB-178928)
WBTM 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB-178425)
WBTM 30 Numerical Weather Prediction and Synoptic Meteorology. Capt. Thomas D. Murphy, U.S.A.F., May 1968. (AD-673365)
WBTM 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB-179084)
WBTM 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB-179289)
WBTM 33 Objective Forecasting. Philip Williams, Jr., August 1968. (AD-680425)
WBTM 34 The WSR-57 Radar Program at Missoula, Montana. R. Granger, October 1968. (PB-180292)
WBTM 35 Joint ESSA/FAA ARTC Radar Weather Surveillance Program. Herbert P. Benner and DeVon B. Smith, December 1968 (revised June 1970). (AD-681857)
WBTM 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (Out of print.) (PB-183055)
WBTM 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB-183057)
WBTM 38 Climate of Phoenix, Arizona. R. J. Schmidli, P. C. Kangieser, and R. S. Ingram, April 1969. (Out of print.) (PB-184295)
WBTM 39 Upper-Air Lows Over Northwestern United States. A. L. Jacobson, April 1969. (PB-184296)
WBTM 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L. W. Snellman, August 1969. (PB-185068)
WBTM 41 High Resolution Radiosonde Observations. W. S. Johnson, August 1969. (PB-185673)
WBTM 42 Analysis of the Southern California Santa Ana of January 15-17, 1966. Barry B. Aronovitch, August 1969. (PB-185670)
WBTM 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB-185762)
WBTM 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB-187763)
WBTM 45/1 Precipitation Probabilities in the Western Region Associated with Winter 500-mb Map Types. Richard A. Augulis, December 1969. (PB-188248)

U. S. DEPARTMENT OF COMMERCE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
NATIONAL WEATHER SERVICE

NOAA Technical Memorandum NWSTM WR-73

A SYNOPTIC CLIMATOLOGY FOR SNOWSTORMS
IN NORTHWESTERN NEVADA

Bert L. Nelson
Weather Service Office
Reno, Nevada

Paul M. Fransioli
Desert Research Institute
University of Nevada System
Reno, Nevada

Clarence M. Sakamoto
Weather Service Office
Reno, Nevada



WESTERN REGION
TECHNICAL MEMORANDUM NO. 73

SALT LAKE CITY, UTAH
FEBRUARY 1972

TABLE OF CONTENTS

	<u>Page</u>
Abstract	i
I. Introduction	i
II. Data	i
III. Results	2-4
IV. Snowstorm-Thunderstorm of May 20 - 21, 1971	4-5
V. Conclusions	5
VI. Acknowledgment	5
VII. References	6
VIII. Bibliography	6

LIST OF FIGURES AND TABLES

	<u>Page</u>
Figure 1. Type I. Surface and 500-mb Charts for 1200Z December 24, 1968	7
Figure 2. Type I. Surface and 500-mb Charts for 1200Z December 25, 1968	8
Figure 3. Type II. Surface and 500-mb Charts for 0600Z and 0000Z respectively, March 14, 1963	9
Figure 4. Type II. Surface and 500-mb Charts for 0600Z and 0000Z respectively, March 15, 1963	10
Figure 5. Type III. Surface and 500-mb Charts for 1200Z January 9, 1964	11
Figure 6. Type III. Surface and 500-mb Charts for 1200Z January 10, 1964	12
Figure 7. Type III. Surface and 500-mb Charts for 1200Z April 12, 1970	13
Figure 8. Type III. Surface and 500-mb Charts for 1200Z April 13, 1970	14
Figure 9. Type III. Surface and 500-mb Charts for 1200Z April 14, 1970	15
Figure 10. Type IV. Surface and 500-mb Charts for 1200Z February 18, 1969	16
Figure 11. Type IV. Surface and 500-mb Charts for 1200Z February 19, 1969	17
Figure 12A. Western Region Type 4 Winter Mean Map	18
Figure 12B. Western Region Type 6 Winter Mean Map	18
Figure 13A. Western Region Type 3 Spring Mean Map	19
Figure 13B. Western Region Type 7 Spring Mean Map	19
Figure 14. Surface and 500-mb Charts for 1200Z May 19, 1971	20
Figure 15. Surface and 500-mb Charts for 1200Z May 20, 1971	21
Figure 16. Surface and 500-mb Charts for 1200Z May 21, 1971	22

	<u>Page</u>
Table 1. Distribution of Snowstorms at Reno, Lovelock, and Winnemucca, Nevada	23
Table 2. Frequency of the Five Types of Snowstorms of 1 Inch or More and 4 Inches or More in Northwestern Nevada	24

A SYNOPTIC CLIMATOLOGY FOR SNOWSTORMS IN NORTHWESTERN NEVADA

ABSTRACT

The purpose of this study is to provide a climatological aid for forecasting snow in northwestern Nevada. A total of 112 snowstorms affecting Reno, Lovelock, or Winnemucca were analyzed to determine if these storms could be categorized into separate types. Five separate categories were defined and are discussed. Examples of four of these types are provided. A separate discussion of the unusual thunderstorm-snowstorm occurrence on May 20 - 21, 1971, is also provided.

I. INTRODUCTION

Northwestern Nevada experiences several heavy snowstorms each year. These storms affect economic activity and at times bring transportation to a standstill. Snowstorms often hamper airline schedules, closing fields because of hazardous ice, snow and/or reduced visibility. Strong winds often accompany these storms. Commuters have difficulty getting to and from work and transportation of goods on Interstate 80 is affected. Analysis of these storms may aid those involved from the standpoint of timely anticipation and prediction by meteorologists. Therefore, a study was initiated to develop a climatological aid for forecasting snow for northwestern Nevada. The objective of this study was to categorize those storms which are similar in synoptic patterns and movements. For the purpose of this analysis, a snowstorm was defined as one which resulted in a snowfall of one inch or more during a 24-hour period. Furthermore, if a storm lasted more than one day, e.g., 3 days, then the total snowfall for the three days was used as the storm total. Subjective analysis of maps was used to identify storm duration. A heavy snowstorm was defined as a snowfall of 4 inches or more.

II. DATA

Reno, Lovelock, and Winnemucca were chosen as locations for determining areal distribution and frequency of snow storms. All three cities are located on Interstate 80, the main artery across northern Nevada. The data base of 10 years from 1961 to 1970 was selected, as it was readily available locally. Furthermore, the 10-year period was considered sufficiently long to obtain a reasonable sample size for analysis. Climatological Data for Nevada [1] was used to verify storm totals. Daily Weather Maps [5] were utilized to determine storm tracks, associated upper-air patterns, synoptic situations and frequency of each type of snowstorm.

III. RESULTS

I. Areal Distribution.

A total of 112 snowstorms of one inch or more at Reno, Lovelock, or Winnemucca was recorded for the 10-year period. Of these 112, Lovelock experienced only 23 snowfalls of at least one inch while Reno had 65 and Winnemucca 68. Table I shows the distribution of these storms at each location. Some of these storms were common to all three locations; most of them were not.

Analysis of each snowfall indicated that a snowstorm occurred only twice in Lovelock without occurring at Reno or Winnemucca. A snowstorm was common to both Reno and Winnemucca 23 times. Thus, 42 (65 minus 23) of Reno's snowfalls did not produce one inch or more of snow in Winnemucca and 45 (68 minus 23) of Winnemucca's snowfalls did not produce one inch or more of snow in Reno. This breakdown does not consider storms where it snowed in Winnemucca and rained in Reno or vice versa. There was a total of 29 cases of heavy snowfalls. Only 8 of these occurred at both Reno and Winnemucca. The average heavy snowfall was 7.5 inches for Reno, 6 inches for Winnemucca, and 4 inches (2 cases) for Lovelock. During the period 1961 - 1970, the heaviest snowfalls were 16.7 inches in Reno, 8.4 inches in Winnemucca, and 4.0 inches in Lovelock for one snowstorm. Heavier amounts, however, have fallen in a 24-hour period at Lovelock and Winnemucca prior to 1961.

Topography is a major factor affecting the distribution of snowfall in northwestern Nevada. For example, Reno is located in Washoe Valley with the Sierra Nevada to the west and smaller mountain ranges to the east and south. Elevation differences within the valley are also a factor in the quantity of snowfall. The airport elevation is 4,411 feet. Urban housing surrounding Reno extends up to 5,000 feet in elevation. Sometimes it may be snowing at the higher elevation area and raining at the airport. Reno is also susceptible to easterly flow that funnels through the Truckee River Valley. This easterly flow is normally associated with cyclogenesis and wave development in central Nevada. When this occurs, snowfall is usually heavy and steady, and stations on the east side of the Sierra receive greater amounts of snow than those on the normally wetter west slope.

Winnemucca (elevation 4,303 feet) is located in Paradise Valley and is not as well protected by mountains as Reno or as Lovelock. The Sonoma Range is southeast of Winnemucca and provides protection from southeasterly winds. Other low-lying hills provide some protection, but Winnemucca is particularly exposed to Pacific cold fronts approaching from the northwest.

Lovelock is located in the Humboldt River Valley which is oriented north-south with the Trinity Range to the west and the Humboldt Range to the east. Lovelock's elevation of 3,903 feet is slightly lower than Reno and Winnemucca. Lovelock is in a warm area that extends from the lee of Sierra into portions of Humboldt, Pershing, and Churchill counties. Lovelock's average yearly snowfall is only 8.7 inches compared with 24.1 inches at Reno and 30.3 inches at Winnemucca. Thus, the combination of low elevation and protected location contributes to the semiarid climate at Lovelock.

2. Storm Types and Frequencies.

One hundred twelve storms were typed using mean storm tracks for the United States after Bowie and Weightman [2]. The North Pacific Storm Track was interpreted to be the path of cyclones moving inland from northern California, Washington, and Oregon and was broken down into three types: Types I, II, and III. These were defined as follows:

Type I: A north Pacific cyclone tracking across Washington or Oregon with no wave development along the front. This type is generally associated with a 500-mb trough along the Pacific Coast with weak or no cold air advection into Nevada. (See Figures 1 and 2.)

Type II: A north Pacific cyclone tracking southeasterly across Nevada. This type is associated with a 500-mb trough along the Pacific Coast that is deepening quite rapidly and has strong cold air advection (Figures 3 and 4). (The cold advection would be more apparent on 700-mb and 850-mb charts.)

Type III: A north Pacific cyclone similar to Type I but with wave development along the frontal system. This type is associated with a 500-mb trough along the Pacific Coast moving inland with a moderate amount of cold air advection (Figures 5 through 9).

Cyclones that move inland from southern California were categorized into another type.

Type IV: A South Pacific Storm Track moving inland into California and tracking across southern Nevada. The 500-mb trough is very deep and extends over Baja, California. At times the 500-mb map shows a closed low over Baja, California (Figures 10 and 11).

Type V: Snowfall that does not conform to one of the other four storm tracks.

Table 2 shows snowstorm frequency distribution by type of storm. The North Pacific Storm Track (Types I, II, and III) had 90 cases, which is 80 percent of the total number of storms. Heavy snowfalls (4 inches or more) were quite evenly distributed among Types I, II, and III. In spring, Type III storms associated with wave formation result in precipitation more often in the form of rain than snow, at the lower elevations.

The South Pacific Storm Track (Type IV) was associated with only 7 snowfalls of an inch or more. This is because low centers from the south pick up warm, moist air off the southern California coast and generally bring rain rather than snow into northwestern Nevada. Therefore, temperature is critical in this type of storm.

Comparison of snowfall cases with 500-mb maps analyzed by Augulis [3,4] showed that many of the storms conform to his specific 500-mb patterns. Twelve cases correspond to Augulis' Winter Type 4 (Figure 12A) where the trough line lies between 130 - 140°W and 18 cases correspond to Winter Type 6 (Figure 12B) with the trough line lying just inside the West Coast. Eighteen cases correspond to Spring Type 3 (Figure 13A) which has a trough line that lies between 110 - 120°W. A total of 9 cases fit Spring Type 7 (Figure 13B), which shows a negative tilt trough between 125 - 130°W.

IV. SNOWSTORM-THUNDERSTORM OF MAY 20 - 21, 1971

This storm, although resembling a Type III storm, warrants further discussion. The storm formed and moved with such rapidity that it produced thunderstorms together with snow and a near blizzard condition which reduced visibility to zero. Temperatures preceding the storm reached a high of 72 degrees in the afternoon of May 20 and dropped to 33 degrees 9 hours later. A total of 6.3 inches of snow fell at Reno during the evening of the 20th and the morning of the 21st. Similarly, 4.3 inches fell at Winnemucca while Lovelock recorded only a trace. The snow was relatively dense as the rapid influx of cold air was preceded by relatively warm, moist layer over Nevada.

Figures 14 through 16 show the synoptic patterns beginning with May 19, 1971. A weak cold front moved through Nevada on the 19th (Figure 14) and by 1200Z May 20 had passed Las Vegas with a suggestion of a wave formation in southern Nevada (Figure 15). The 500-mb map on the 19th (Figure 14) showed a very deep trough off Alaska, extending southeastward through Washington. The following day, a closed low formed over Washington with a weak trough extending southward through northern California (Figure 15). At the same time, the

ridge over western Canada was building. By 1200Z on the 21st, a closed surface low had formed over Nevada with a cold low aloft over northern California (Figure 16). Analysis of 6-hourly surface maps showed a definite deepening of the surface low on the 1800Z map and by 0000Z on the 22nd, further deepening was indicated.

The rapidity with which the storm formed and the dense snowfall accompanying it caused heavy property damage, primarily from broken branches and trees, power outages, and damage to agricultural crops.

V. CONCLUSIONS

Analysis of 112 snowstorms during a 10-year period shows the following:

1. Snowstorms moving through northwestern Nevada are much more likely to affect Reno and Winnemucca than Lovelock.
2. Local differences have a large effect on the amount of snowfall observed in any storm.
3. Over 80 percent of the snowstorms were associated with the North Pacific Storm Track, (Types I, II, and III).
4. The South Pacific Storm Track (Type IV) and Cyclogenesis resulted in very few snowstorms in northwestern Nevada.
5. Twenty-five of the 29 heavy (4 inches or more) snowstorms were associated with the North Pacific Storm Track (Types I, II, and III).

VI. ACKNOWLEDGMENT

This study was initiated by a need expressed in a graduate course in Climatology in which Clarence M. Sakamoto participated at the University of Nevada. Joseph Ganser, Meteorologist in Charge, Weather Service Office, Reno, and members of his staff were helpful with their review and suggestions. The authors also acknowledge comments and constructive criticisms by the Scientific Services Division, Western Region, National Weather Service.

VII. REFERENCES

- [1] *Climatological Data for Nevada, 1961 - 1970*. U. S. Department of Commerce, Environmental Data Service.
- [2] BOWIE, H., and R. H. WEIGHTMAN. *Types of Storms of the United States and Their Average Movements*. Monthly Weather Review, Supplement 1.
- [3] AUGULIS, RICHARD P. *Precipitation Probabilities in the Western Region Associated with Winter 500-mb Map Types*. U. S. Department of Commerce, Technical Memorandum WBTM WR45-1, 1969.
- [4] AUGULIS, RICHARD P. *Precipitation Probabilities in the Western Region Associated with Spring 500-mb Map Types*. U. S. Department of Commerce, Technical Memorandum WBTM WR45-1, 1970.
- [5] *Daily Weather Maps, 1961 - 1970*. U. S. Department of Commerce, Weather Bureau.

VIII. BIBLIOGRAPHY

1. HOVEY, WILLIAM B., and MARK D. SHULMAN. "A Synoptic Climatology of Heavy Snowstorms for New Jersey," *The Bulletin, New Jersey Academy of Science*, 10, No. 2, 1965.
2. KLEIN, W. H., D. L. JORGENSON and A. F. KORTE. "Relation Between Upper Air Lows and Winter Precipitation in the Western Plateau States," *Monthly Weather Review*, 96, pp. 162 - 168, 1968.
3. KLEIN, W. H., D. L. JORGENSON and A. F. KORTE. "A Synoptic Climatology of Winter Precipitation for 700-mb Lows for Intermountain Areas of the West," *Journal of Applied Meteorology*, 6, pp. 782 - 790, 1967.
4. ROGERS, C. FREDERICK. "A Mountain Precipitation Study," *DRI Thesis 431*, Library, University of Nevada, Reno, Nevada, 1970.
5. WEAVER, DONALD. "Meteorology of Hydrologically Critical Storms in California," *Hydrometeorological Report #37*, U. S. Department of Commerce, December 1962.





FIGURE 1





FIGURE 2





FIGURE 3





FIGURE 4





FIGURE 5





FIGURE 6





FIGURE 7





FIGURE 8





FIGURE 9





FIGURE 10









FIGURE 11








FIGURE 14





FIGURE 15




FIGURE 16

TABLE 1

DISTRIBUTION OF SNOWSTORMS AT RENO, LOVELOCK, AND WINNEMUCCA, NEVADA (1961 - 1970)

SNOW AMOUNT	LOCATION		
	Reno	Lovelock	Winnemucca
At least 1 inch	65	23	68
At least 2 inches	42	15	39
4 inches or more	19	2	18

TABLE 2

FREQUENCY OF THE FIVE TYPES OF SNOWSTORMS OF 1 INCH OR MORE AND 4 INCHES OR MORE IN NORTHWESTERN NEVADA (1961 - 1970)

STORM TRACK Type	NUMBER OF SNOWSTORMS	
	1 inch or more	4 inches or more
Type I	35	6
Type II	27	9
Type III	28	10
Type IV	7	1
Type V	15	3

Western Region Technical Memoranda: (Continued)

- No. 45/2 Precipitation Probabilities in the Western Region Associated with Spring 500-mb Map Types. Richard P. Augulis. January 1970. (PB-189434)
- No. 45/3 Precipitation Probabilities in the Western Region Associated with Summer 500-mb Map Types. Richard P. Augulis. January 1970. (PB-189414)
- No. 45/4 Precipitation Probabilities in the Western Region Associated with Fall 500-mb Map Types. Richard P. Augulis. January 1970. (PB-189435)
- No. 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates. December 1969. (PB-190476)
- No. 47 Statistical Analysis as a Flood Routing Tool. Robert J. C. Burnash. December 1969. (PB-188744)
- No. 48 Tsunami. Richard A. Augulis. February 1970. (PB-190157)
- No. 49 Predicting Precipitation Type. Robert J. C. Burnash and Floyd E. Hug. March 1970. (PB-190962)
- No. 50 Statistical Report of Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona 1969. Wayne S. Johnson. April 1970. (PB-191743)
- No. 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell. July 1970. (PB-193102)
- No. 52 Sacramento Weather Radar Climatology. R. G. Pappas and C. M. Veliquette. July 1970. (PB-193347)
- No. 53 Experimental Air Quality Forecasts in the Sacramento Valley. Norman S. Benes. August 1970. (PB-194128)
- No. 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch. August 1970.
- No. 55 Application of the SSARR Model to a Basin Without Discharge Record. Vail Schermerhorn and Donald W. Kuehl. August 1970. (PB-194394).
- No. 56 Areal Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck. September 1970. (PB-194389)
- No. 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote. September 1970. (PB-194710)
- No. 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson. October 1970. (COM-71-00017)
- No. 59 Application of P.E. Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman. October 1970. (COM-71-00016)

NOAA Technical Memoranda NWS

- No. 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon. Arthur W. Fritz, October 1970. (COM-71-00120)
- No. 61 Relationship of Wind Velocity and Stability to SO₂ Concentrations at Salt Lake City, Utah. Werner J. Heck, January 1971. (COM-71-00232)
- No. 62 Forecasting the Catalina Eddy. Arthur L. Eichelberger, February 1971. (COM-71-00223)
- No. 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Woerner. February 1971. (COM-71-00349)
- No. 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- No. 65 Climate of Sacramento, California. Wilbur E. Figgins, June 1971. (COM-71-00764)
- No. 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM-71-00829)
- No. 67 Precipitation Detection Probabilities by Los Angeles ARTC Radars. Dennis E. Ronne, July 1971.
- No. 68 A Survey of Marine Weather Requirements. Herbert P. Benner, July 1971. (COM-71-00889)
- No. 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM-71-00956)
- No. 70 Predicting Inversion Depths and Temperature Influences in the Helena Valley. David E. Olsen, October 1971.
- No. 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972.
- No. 72 A Paradox Principle in the Prediction of Precipitation Type. Thomas J. Weitz, February 1972.