



STATE OF CALIFORNIA  
MARINE RESEARCH COMMITTEE



**CALIFORNIA COOPERATIVE OCEANIC  
FISHERIES INVESTIGATIONS**

ATLAS No. 20



CALIFORNIA  
COOPERATIVE  
OCEANIC  
FISHERIES  
INVESTIGATIONS

*Atlas No. 20*

STATE OF CALIFORNIA  
MARINE RESEARCH COMMITTEE

*Cooperating Agencies:*

CALIFORNIA ACADEMY OF SCIENCES  
CALIFORNIA DEPARTMENT OF FISH AND GAME  
STANFORD UNIVERSITY, HOPKINS MARINE STATION  
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION, NATIONAL MARINE FISHERIES SERVICE  
UNIVERSITY OF CALIFORNIA, SCRIPPS INSTITUTION OF OCEANOGRAPHY

June, 1974

## THE CALCOFI ATLAS SERIES

This is the twentieth in a series of atlases containing data on the hydrography and plankton from the region of the California Current. The field work was carried out by the California Cooperative Oceanic Fisheries Investigations,<sup>1</sup> a program sponsored by the State of California under the direction of the State's Marine Research Committee. The cooperating agencies in the program are:

California Academy of Sciences  
California Department of Fish and Game  
Stanford University, Hopkins Marine Station  
National Oceanic and Atmospheric Administration, National Marine Fisheries Service<sup>2</sup>  
University of California, Scripps Institution of Oceanography

CalCOFI atlases<sup>3</sup> are issued as individual units as they become available. They provide processed physical, chemical and biological measurements of the California Current region. Each number may contain one or more contributions. A general description of the CalCOFI program with its objectives appears in the preface of Atlas No. 2.

This atlas was prepared by the Data Collection and Processing Group of the Marine Life Research Program, Scripps Institution of Oceanography.

CalCOFI Atlas Editorial Staff:

Abraham Fleminger and John G. Wyllie, Editors

CalCOFI atlases in this series, through June 1974, are:

- No. 1. Anonymous, 1963. CalCOFI atlas of 10-meter temperatures and salinities 1949 through 1959.
- No. 2. Fleminger, A., 1964. Distributional atlas of calanoid copepods in the California Current region, Part I.
- No. 3. Alvarino, A., 1965. Distributional atlas of Chaetognatha in the California Current region.
- No. 4. Wyllie, J.G., 1966. Geostrophic flow of the California Current at the surface and at 200 meters.
- No. 5. Brinton, E., 1967. Distributional atlas of Euphausiacea (Crustacea) in the California Current region, Part I.
- No. 6. McGowan, J.A., 1967. Distributional atlas of pelagic molluscs in the California Current region.
- No. 7. Fleminger, A., 1967. Distributional atlas of calanoid copepods in the California Current region, Part II.
- No. 8. Berner, L.D., 1967. Distributional atlas of Thaliacea in the California Current region.
- No. 9. Kramer, D., and E. H. Ahlstrom, 1968. Distributional atlas of fish larvae in the California Current region: Northern Anchovy, *Engraulis mordax* (Girard). 1951 through 1965.
- No. 10. Isaacs, J. D., A. Fleminger and J. K. Miller, 1969. Distributional atlas of zooplankton biomass in the California Current region: Spring and Fall 1955-1959.
- No. 11. Ahlstrom, E. H., 1969. Distributional atlas of fish larvae in the California Current region: jack mackerel, *Trachurus symmetricus*, and Pacific hake, *Merluccius productus*, 1951 through 1966.
- No. 12. Kramer, D., 1970. Distributional atlas of fish eggs and larvae in the California Current region: Pacific sardine, *Sardinops caerulea* (Girard). 1951 through 1966.
- No. 13. Smith, P. E., 1971. Distributional atlas of zooplankton volume in the California Current region, 1951 through 1966.
- No. 14. Isaacs, J. D., A. Fleminger and J. K. Miller, 1971. Distributional atlas of zooplankton biomass in the California Current region: Winter 1955-1959.
- No. 15. Wyllie, J. G., and R. J. Lynn, 1971. Distribution of temperature and salinity at 10 meters, 1960-1969 and mean temperature, salinity and oxygen at 150 meters, 1950-1968 in the California Current.
- No. 16. Crowe, F. J. and R. A. Schwartzlose, 1972. Release and recovery records of drift bottles in the California Current region, 1955 through 1971.
- No. 17. Ahlstrom, E. H., 1972. Distributional atlas of fish larvae in the California Current region: six common mesopelagic fishes—*Vinciguerria lucetia*, *Triphoturus mexicanus*, *Stenobranchius leucopsarus*, *Leuroglossus stilbius*, *Bathylagus wesethi* and *Bathylagus ochotensis*. 1955 through 1960.
- No. 18. Brinton, E., 1973. Distributional atlas of Euphausiacea (Crustacea) in the California Current region, Part II.


- No. 19. Bowman, T. E. and M. W. Johnson, 1973. Distributional atlas of calanoid copepods in the California Current region, 1949 and 1950.
- No. 20. Thomas, W. H. and D. L. R. Seibert, 1974. Distribution of nitrate, nitrite, phosphate and silicate in the California Current region, 1969.
- Owen, R. W., Jr., 1974. Distribution of primary production, plant pigments and Secchi depth in the California Current region, 1969.
- Smith, P. E., 1974. Distribution of zooplankton volumes in the California Current region, 1969.

<sup>1</sup> Usually abbreviated CalCOFI, sometimes CALCOFI or CCOFI.

<sup>2</sup> Formerly called U.S. Fish and Wildlife Service, Bureau of Commercial Fisheries.

<sup>3</sup> For citation this issue in the series should be referred to as CalCOFI Atlas No. 20.

Library of Congress Catalog Card Number 67-4238.



CONTENTS

W. H. Thomas and D. L. R. Seibert

Distribution of Nitrate, Nitrite, Phosphate and Silicate  
in the California Current region, 1969 ..... vii  
Charts 2-97

R. W. Owen Jr.

Distribution of Primary Production, Plant Pigments and  
Secchi Depth in the California Current region, 1969..... xi  
Charts 98-117

P. E. Smith

Distribution of Zooplankton Volumes in the  
California Current region, 1969 ..... xv  
Charts 118-125

CalCOFI Atlas No. 20  
A. Fleminger and J. G. Wyllie, Editors  
Data Collection and Processing Group  
Marine Life Research Program  
Scripps Institution of Oceanography  
La Jolla, California  
June, 1974

# DISTRIBUTION OF NITRATE, NITRITE, PHOSPHATE AND SILICATE IN THE CALIFORNIA CURRENT REGION, 1969

W. H. Thomas and D. L. R. Seibert<sup>1</sup>

## Text

Introduction .....	vii
Methods .....	ix
10 meter nutrients .....	ix
Nutrients integrated over 0 to 50 meters	ix
100 meter nutrients .....	ix

## Charts

10 meter Nitrate - N distribution. ....	2-9
10 meter Nitrite - N distribution. ....	10-17
10 meter Phosphate - P distribution. ....	18-25
10 meter Silicate - Si distribution. ....	26-33
Nitrate - N integrated over 0-50 meters. . .	34-41
Nitrite - N integrated over 0-50 meters. . .	42-49
Phosphate - P integrated over 0-50 meters. .	50-57
Silicate - Si integrated over 0-50 meters. . .	58-65
100 meter Nitrate - N distribution. ....	66-73
100 meter Nitrite - N distribution. ....	74-81
100 meter Phosphate - P distribution. ....	82-89
100 meter Silicate - Si distribution. ....	90-97

## Introduction

Nutrient levels are important in controlling the productivity of phytoplankton in the sea and in turn the other biota that are dependent on phytoplankton as a food source. Coastal regions where nutrient concentrations are high are areas of major fishing activity. An example is the Peru Current where nutrient replenishment due to widespread upwelling supports the worlds largest fishery.

In the California Current region thousands of nutrient determinations have been made since the CalCOFI program commenced in 1949. Sources of data on nutrient concentrations off California prior to 1969 are summarized in Table 1.

Phosphate data are abundant for the years 1949 to 1951 and moderately so for 1955. However, for nearly all years from 1951 to 1960, no nutrient data were collected by CalCOFI cruises. In 1961, the collecting of chemical data increased with the advent of newer, more reliable methods of analysis, and other nutrients in addition to phosphate were measured. In 1964 to 1966, five nutrients were analyzed during an extensive series of cruises carried out off southern Baja California by the Scripps Tuna Oceanography Research (STOR) project. Chemical nutrients were not measured by the CalCOFI program in 1966 and 1967, but the Food Chain Research Group (FCRG) of the Institute of Marine Resources carried out an extensive time series of analyses at three nearshore stations off La Jolla. Other nearshore data are available for 1959 and 1960.

In 1969 nutrient measurements off California was greatly expanded. A more complete suite of nutrients was analyzed and measurements were obtained in almost every month of the year. It is especially useful that one of the principal nitrogen sources, nitrate, was measured since it has been shown to limit phytoplankton production off Baja California and southern California (Thomas, 1969; Thomas et al, 1974).

In this atlas we present charts of the horizontal distribution of major phytoplankton nutrients, nitrate, nitrite, phosphate and silicate in the California Current region during 1969.

---

1. Scripps Tuna Oceanography Research Program  
Institute of Marine Resources  
Scripps Institution of Oceanography, La Jolla,  
California.



TABLE 1. SOURCES OF NUTRIENT DATA OFF CALIFORNIA AND BAJA CALIFORNIA, 1949-1968

Date	Cruise	Nutrients Analyzed	Remarks:	References
1949	CCOFI	PO <sub>4</sub> -P	Nutrient data available from many stations. Seasonal coverage.	Rakestraw et al. (1957)
1950	CCOFI	PO <sub>4</sub> -P	Nutrient data available from many stations. Seasonal coverage.	Rakestraw et al. (1960)
1951	CCOFI	PO <sub>4</sub> -P	Nutrient data available from many stations. Seasonal coverage.	Reid et al. (1963 a)
1953	Transpac	PO <sub>4</sub> -P SiO <sub>3</sub> -Si	1 station within California Current Area	Reid et al. (1965 a)
1955	NORPAC	PO <sub>4</sub> -P	Moderate number of stations	Rakestraw et al. Norpac (1960)
1956	SCOPE	SiO <sub>3</sub> -Si PO <sub>4</sub> -P	1 station off southern Baja California	Reid et al. (1963 b)
1959	STOR TO59-1	NO <sub>2</sub> -N PO <sub>4</sub> -P	3 stations off southern Baja California	Reid et al. (1965 b)
1959	Allan Hancock Foundation	PO <sub>4</sub> -P SiO <sub>3</sub> -Si	Nearshore surface stations in Southern California Bight, October-November	Chambers (1965)
1960	Allan Hancock Foundation	PO <sub>4</sub> -P SiO <sub>3</sub> -Si	Nearshore surface stations in Southern California Bight, April	Chambers (1965)
1961	CCOFI 6107-8	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N	Nutrient data available from many stations in July and August	Anon., SIO Ref. 62-16 (1962)
1962	CCOFI 6210-11	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N	Nutrient data available from many stations in October and November	Anon., SIO Ref. 63-25 (1963 a)
1963	CCOFI 6301-2, 6304, 6311	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N	Nutrient data available from many stations in January, February, April, May, November, December (latter 2 months with only a few nearshore stations off southern Baja California)	Anon., SIO Ref. 64-2 (1963 b) Anon., SIO Ref. 64-13 (1964 a) Anon., SIO Ref. 65-1 (1964 b)
1964	CCOFI 6401, 6404, 6407	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N	Nutrient data available from many stations in January, February, March, April, July	Anon., SIO Ref. 65-7 (1965 a) Anon., SIO Ref. 66-20 (1966)
	STOR TO-64-1 TO-64-2	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N NO <sub>3</sub> -N NH <sub>4</sub> -N	Nutrient data available from many stations off southern Baja California in June and August	Anon., SIO Ref. 69-4 (1969) IMR Ref. 69-8
1965	CCOFI 6501, 6504, 6507	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N	Nutrient data available for many stations in January, February, March, April, June, July, August	Anon., SIO Ref. 66-4 (1965 b) Anon., SIO Ref. 67-16 (1967 a) Anon., SIO Ref. 67-17 (1967 b)
	STOR TO-65-1	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N NO <sub>3</sub> -N NH <sub>4</sub> -N	Many stations off southern Baja California in September	Thomas (Unpublished data)
1966	STOR TO-66-1	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N NO <sub>3</sub> -N NH <sub>4</sub> -N	Many stations off southern Baja California in November	Thomas (Unpublished data)
1967	FCRG	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N NO <sub>3</sub> -N	3 stations off La Jolla occupied weekly for 5-month period April-September	Strickland (1968)
1968	CCOFI 6804, 6806	PO <sub>4</sub> -P SiO <sub>3</sub> -Si NO <sub>2</sub> -N NO <sub>3</sub> -N	Nutrient data available for many stations in April, May, June	Anon., SIO-Ref. 71-3 (1971)

## Methods

Water samples were collected by Nansen bottles or with a Niskin sampler. Hydrographic casts to 1000 meters were made at selected stations. The data from these stations were used for estimating the abundance of nutrients in the upper 50 meters, by integration, and for the distribution at 100 meters. In addition to the hydrographic stations intermediate positions were sampled at 10-meter depths.

Most of the samples were analyzed immediately after collection by means of the autoanalyzer technique (Strickland and Parsons, 1968). The limits of detection of nitrate, nitrite, phosphate and silicate using these procedures are 0.1, 0.01, 0.03 and 0.1  $\mu\text{g-atoms/liter}$  respectively. In the lattermost cruises of 1969 samples were frozen and processed ashore using the autoanalyzer.

Integration of the data from 0 to 50 meters and the plotting of the 10 meter, 100 meter and integrated values was carried out with the aid of a CDC 3600 computer; contours were fitted by eye.

## 10 Meter Nutrients

To illustrate near-surface mixed layer nutrient distributions charts are presented showing contoured values from a depth of 10 meters. Especially high

nutrient values were found during periods of seasonal upwelling along the central California coast, Point Conception, Punta Eugenia and Punta Abreojos. Generally lower values were found farther offshore.

## Nutrients Integrated 0 - 50 Meters

These data provide a general idea of the nutrients available to phytoplankton in the euphotic zone. Their distributions are similar to those at 10 meters.

## 100 Meter Nutrients

These charts show the distribution below the mixed layer. They may vary more due to physical than to biological processes. The distributions show the nutrient levels that might enter the euphotic zone as a result of upwelling or vertical mixing.


## Acknowledgements

Technicians of the Data Collecting and Processing Group, Marine Life Research Program, Scripps Institution of Oceanography were responsible for collecting and analyzing the samples and for computing the results. The preparation of these charts was partially supported by the Oceanography Program, National Science Foundation, Grant GA32529X.

## REFERENCES CITED

- Anonymous, Scripps Inst. of Oceanogr., 1962. Data Rept. Phys. and Chem. Data CCOFI Cruise 6107-08. SIO Ref. No. 62-16.
- \_\_\_\_\_, 1963 a. Data Rept. Phys. and Chem. Data CCOFI Cruises 6210-11 and 6212, SIO Ref. No. 63-25.
- \_\_\_\_\_, 1963 b. Data Rept. Phys. and Chem. Data CCOFI Cruise 6301-2. SIO Ref. No. 64-2.
- \_\_\_\_\_, 1964 a. Data Rept. Phys. and Chem. Data CCOFI Cruises 6304 and 6306. SIO Ref. No. 64-13.
- \_\_\_\_\_, 1964 b. Data Rept. Phys. and Chem. Data CCOFI Cruises 6310 and 6311. SIO Ref. No. 65-1.
- \_\_\_\_\_, 1965 a. Data Rept. Phys. and Chem. Data CCOFI Cruise 6401. SIO Ref. No. 65-7.
- \_\_\_\_\_, 1965 b. Data Rept. Phys. and Chem. Data CCOFI Cruise 6501. SIO Ref. No. 66-4.
- \_\_\_\_\_, 1966. Data Rept. Phys. and Chem. Data CCOFI Cruises 6404 and 6407. SIO Ref. No. 66-20.
- \_\_\_\_\_, 1967 a. Data Rept. Phys. and Chem. Data CCOFI Cruise 6504. SIO No. 67-16.
- \_\_\_\_\_, 1967 b. Data Rept. Phys. and Chem. Data CCOFI Cruises 6507 and 6509. SIO Ref. No. 67-17.

- \_\_\_\_\_, 1969. Data Rept. Phys. and Chem. and Biol. Data. Cruises TO-64-1 and TO-64-2. SIO Ref. No. 69-4, IMR Ref. No. 69-8.
- \_\_\_\_\_, 1971. Data Rept. Phys. and Chem. Data CCOFI Cruises 6801, 6804 and 6806. SIO Ref. No. 71-3.
- Chambers, L. A. 1965. An oceanographic and biological survey of the southern California mainland shelf. Calif. State Water Qual. Cont. Bd. Publ. No. 27: 1-231.
- Rakestraw, N. W., P. L. Horner, and W. S. Wooster (Eds.). 1957. Oceanic observations of the Pacific: 1949. Univ. Calif. Press, Berkeley, 363 p.
- Rakestraw, N. W., J. L. Reid, Jr., P. Rudnick and W. S. Wooster (Eds.). 1960. Oceanic observations of the Pacific: 1950. Univ. Calif. Press, Berkeley, 508 p.
- \_\_\_\_\_, 1960. Oceanic observations of the Pacific: 1955. The NORPAC data. Univ. Calif. Press, Berkeley, 532 p.
- Reid, J. L., Jr., R. S. Arthur, and E. B. Bennett (Eds.). 1963 a. Oceanic observations of the Pacific: 1951. Univ. Calif. Press, Berkeley, 598 p.
- \_\_\_\_\_, 1963 b. Oceanic observations of the Pacific: 1956. Univ. Calif. Press, Berkeley, 458 p.
- \_\_\_\_\_, 1965 a. Oceanic observations of the Pacific: 1953. Univ. Calif. Press, Berkeley, 576 p.
- \_\_\_\_\_, 1965 b. Oceanic observations of the Pacific: 1959. Univ. Calif. Press, Berkeley, 901 p.
- Strickland, J. D. H. 1968. Research on the marine food chain; progress report, Jan. 1967-June 1968. Part III. Data record, coastal plankton survey, 1967. Section I. Chemistry and productivity. IMR Ref. No. 68-8.
- Strickland, J. D. H., and T. R. Parsons, 1968. A practical handbook of seawater analysis. Fish. Res. Bd. Canada Bull., 167: 1-311.
- Thomas, W. H. 1969. Phytoplankton nutrient enrichment experiments off Baja California and in the Eastern equatorial Pacific Ocean. J. Fish. Res. Bd. Canada, 26: 1133-1145.
- Thomas, W. H., D. L. R. Seibert, and A. N. Dodson (in press). Phytoplankton enrichment experiments and bioassays in natural coastal seawater and in-sewage outfall receiving waters off Southern California. Estuarine and Coastal Marine Science.



### CALCOFI CRUISE 6901


6 - 30 JANUARY 1969

### 10 METER NITRATE-N

CONTOUR INTERVAL  $2 \mu\text{g-at/liter}$   
( $0.1 \mu\text{g-at/liter}$  contour dashed)


10m NITRATE-N

6901




10m NITRATE-N

6902




10m NITRATE-N

6904



10m NITRATE-N


6905-6



10 m NITRATE-N


6907





10m NITRATE-N

6908-9



CALCOFI CRUISE 6910


9 OCTOBER - 8 NOVEMBER 1969

10 METER NITRATE-N

CONTOUR INTERVAL  $2 \mu\text{g-at/liter}$   
( $0.1 \mu\text{g-at/liter}$  contour dashed)


10m NITRATE-N

6910



10m NITRATE-N


6912



10m NITRITE-N


6901






10m NITRITE-N

6904



10m NITRITE-N


6905-6



10m NITRITE-N


6907






10m NITRITE-N

6908-9




10m NITRITE-N

6910



10m NITRITE-N

6912



CALCOFI CRUISE 6901


6 - 30 JANUARY 1969

10 METER PHOSPHATE - P

CONTOUR INTERVAL 0.25  $\mu\text{g-at/liter}$


10m PHOSPHATE-P

6901




10m PHOSPHATE-P

6902




10m PHOSPHATE-P

6904



10m PHOSPHATE-P


6905-6



10m PHOSPHATE - P


6907






10m PHOSPHATE-P

6908-9




10m PHOSPHATE - P

6910



10m PHOSPHATE-P

6912



CALCOFI CRUISE 6901


6 - 30 JANUARY 1969

10 METER SILICATE - Si

CONTOUR INTERVAL 5  $\mu\text{g-at/liter}$   
(2  $\mu\text{g-at/liter}$  contour dashed)


10m SILICATE-Si

6901




10m SILICATE-Si

6902




10m SILICATE-Si

6904



10m SILICATE-Si

6905-6



# CALCOFI CRUISE 6907

10 - 29 JULY 1969


## 10 METER SILICATE - Si

CONTOUR INTERVAL  $5 \mu\text{g-at/liter}$   
( $2 \mu\text{g-at/liter}$  contour dashed)

10m SILICATE-Si


6907






10m SILICATE-Si

6908-9




10 m SILICATE - Si

6910



10m SILICATE-Si


6912



INTEGRATED NITRATE-N


6901






INTEGRATED NITRATE-N

6904



INTEGRATED NITRATE-N


6905-6



INTEGRATED NITRATE-N


6907






INTEGRATED NITRATE-N

6908-9




INTEGRATED NITRATE-N

6910




INTEGRATED NITRATE-N

6912




INTEGRATED NITRITE-N

6901




INTEGRATED NITRITE-N

6902




INTEGRATED NITRITE-N

6904



INTEGRATED NITRITE-N


6905-6



INTEGRATED NITRITE-N


6907






INTEGRATED NITRITE-N

6908-9




INTEGRATED NITRITE-N

6910




INTEGRATED NITRITE-N

6912




INTEGRATED PHOSPHATE-P

6901




INTEGRATED PHOSPHATE-P

6902




INTEGRATED PHOSPHATE-P

6904



INTEGRATED PHOSPHATE-P


6905-6



INTEGRATED PHOSPHATE-P


6907






INTEGRATED PHOSPHATE-P

6908-9




INTEGRATED PHOSPHATE-P

6910




INTEGRATED PHOSPHATE-P

6912




INTEGRATED SILICATE-Si

6901




INTEGRATED SILICATE-Si

6902




INTEGRATED SILICATE-Si

6904



INTEGRATED SILICATE-Si


6905-6



INTEGRATED SILICATE-Si


6907






INTEGRATED SILICATE-Si

6908-9




INTEGRATED SILICATE-Si

6910




INTEGRATED SILICATE-Si

6912



100m NITRATE-N

6901



### CALCOFI CRUISE 6902


13 FEBRUARY - 11 MARCH 1969

### 100 METER NITRATE-N

CONTOUR INTERVAL 5  $\mu\text{g-at/liter}$

100m NITRATE-N

6902



CALCOFI CRUISE 6904


2 - 26 APRIL 1969

100 METER NITRATE-N

CONTOUR INTERVAL  $5\mu\text{g-at/liter}$


100m NITRATE-N

6904



100m NITRATE-N

6905-6




CALCOFI CRUISE 6907  
10 - 29 JULY 1969  
100 METER NITRATE-N  
CONTOUR INTERVAL  $5 \mu\text{g-at/liter}$

100m NITRATE-N


6907






100m NITRATE-N

6908-9




100m NITRATE-N

6910




100m NITRATE-N

6912



100m NITRITE-N


6901



100m NITRITE-N

6902





### CALCOFI CRUISE 6905-6


5-29 MAY and 9-28 JUNE 1969

### 100 METER NITRITE-N

CONTOUR INTERVAL 0.05 $\mu\text{g-at/liter}$   
(0.01 $\mu\text{g-at/liter}$  contour dashed)

100m NITRITE-N


6905-6



100m NITRITE-N


6907





100m NITRITE-N


6908-9



CALCOFI CRUISE 6910  
9 OCTOBER - 8 NOVEMBER 1969  
100 METER NITRITE - N  
CONTOUR INTERVAL 0.05  $\mu\text{g-at/liter}$   
(0.01  $\mu\text{g-at/liter}$  contour dashed)


100m NITRITE-N

6910




100m NITRITE-N

6912



100m PHOSPHATE-P


6901



CALCOFI CRUISE 6902  
13 FEBRUARY - 11 MARCH 1969  
100 METER PHOSPHATE - P  
CONTOUR INTERVAL 0.5  $\mu\text{g-at/liter}$

100m PHOSPHATE-P

6902



CALCOFI CRUISE 6904


2 - 26 APRIL 1969

100 METER PHOSPHATE - P

CONTOUR INTERVAL 0.5 μg-at/liter


100m PHOSPHATE-P

6904



100m PHOSPHATE-P

6905-6




CALCOFI CRUISE 6907  
10-29 JULY 1969  
100 METER PHOSPHATE - P  
CONTOUR INTERVAL 0.5 μg-at/liter

100 m PHOSPHATE - P

6907






CALCOFI CRUISE 6908-9  
6 AUG. - 8 SEPT. and 11 SEPT. - 7 OCT. 1969  
100 METER PHOSPHATE - P  
CONTOUR INTERVAL 0.5  $\mu\text{g-at/liter}$

100 m PHOSPHATE-P


6908-9



CALCOFI CRUISE 6910  
9 OCTOBER - 8 NOVEMBER 1969  
100 METER PHOSPHATE - P  
CONTOUR INTERVAL 0.5  $\mu\text{g-at/liter}$


100m PHOSPHATE - P

6910




100m PHOSPHATE-P

6912




100m SILICATE-Si

6901




100m SILICATE-Si

6902




100m SILICATE-Si

6904



100m SILICATE-Si


6905-6



100m SILICATE-Si


6907






100m SILICATE-Si

6908-9



100m SILICATE-Si

6910



100m SILICATE-Si

6912

# DISTRIBUTION OF PRIMARY PRODUCTION, PLANT PIGMENTS, AND SECCHI DEPTH IN THE CALIFORNIA CURRENT REGION, 1969

R. W. Owen, Jr.<sup>1</sup>

## Text

Introduction .....	xi
Phytoplankton pigments .....	xi
Phytoplankton production - simulated in situ method .....	xii
Secchi depth .....	xiii
Chart preparation .....	xiii

## Charts

Primary Production .....	98-101
Chlorophyll <i>a</i> at the sea surface .....	102-105
Chlorophyll <i>a</i> integrated 0-150 meters .....	106-109
Phaeo pigments integrated 0-150 meters ...	110-113
Secchi depth .....	114-117

## Introduction

The charts presented in the phytoplankton section of this atlas summarize measurements obtained systematically in the California Current region during 1969. Together with subsequent measurements, the data assume special significance since they are the only extant source for estimating seasonal cycles, annual changes, and spatial variations in the region. As phytoplankton supports the natural marine food resources available to man, the postulated sensitivity of potential fish yield to phytoplankton variations and the sensitivity of these variations to more easily measured and perhaps predictable environmental changes may be confirmed through these measurements.

## Phytoplankton Pigments

Pigment concentrations were determined by fluorometry using water samples from prescribed regular and special CalCOFI stations. More than 10 levels were usually sampled between the surface and 150m depth in 1969. A secchi disc measurement was

usually used to determine appropriate sampling depths at each station. Where no Secchi reading was possible, standardized depths were sampled. Methods of pigment analyses were used consistent with procedures recommended by the Biological Methods Panel Committee on Oceanography, National Academic of Science (1969).

Samples in 270-ml plastic bottles were filtered under low vacuum (not exceeding -7 inches of mercury) through Whatman GF/C glass fiber filters (2.4-cm diameter) having a thin magnesium carbonate coating. Filtration was usually completed within a half hour after sample collection. In some instances, however, samples were chilled and stored in darkness up to 12 hours before processing.

Filters were placed in 90%-v/v-spectral-quality acetone, ground for 30 seconds in a motor-driven tissue grinder and stored dark for 10 minutes to permit full extraction of the pigments. Filters and extract were separated by centrifuging for 10 minutes. Extract volume was brought up to 10.0 ml; about 5 ml was placed in a cuvette and submitted to a Turner fluorometer, Model 111, for fluorometric analysis by the method of Holm-Hansen, Lorenzen, Holmes and Strickland (1965). A blue-pass Corning 5-60 primary filter and a red-pass Corning 2-64 secondary filter were used in conjunction with the light source and the detector, a red-sensitive multiplier phototube (Hamamatsu R 136).

On occasion, a fluorometer failure occurred and sample filters were folded, stored dark in a freezer,

---

1. National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Southwest Fisheries Center  
La Jolla, California 92037

and analyzed later aboard ship or returned frozen to La Jolla for analysis. Values so obtained were multiplied by 1.1 to account for the consistent 10% loss that occurs when samples are stored for more than a day (Author's data: Lorenzen, pers. comm.).

All phytoplankton data were card punched, verified, and submitted to the CDC 3600 computer for calculation of basic phytoplankton parameters. Computed and listed by cruise, station and depth were chlorophyll *a* concentration, phaeo-pigment concentration<sup>1</sup>, total pigment concentration, phytoplankton production per unit volume and several parameters for use in editing and interpreting these basic values. Also printed were interpolations and vertical profiles of pigment and production values.

Pigment concentrations were calculated from the equations given by Lorenzen (1966):

$$\text{chlorophyll } a \text{ (mgm}^{-3}\text{)} = \frac{\frac{Fo/Fa_{\max}}{(Fo/Fa_{\max}) - 1} (k_x) (Fo - Fa)}{\text{liters filtered}}$$

$$\text{phaeophytin } a \text{ (mgm}^{-3}\text{)} = \frac{\frac{Fo/Fa_{\max}}{(Fo/Fa_{\max}) - 1} (k_x) [Fo/Fa_{\max} (Fa) - Fo]}{\text{liters filtered}}$$

where:  $Fo$  = fluorescence before acidification  
 $Fa$  = fluorescence after acidification  
 $Fo/Fa_{\max}$  = maximum acid factor which can be expected in the absence of phaeophytin  
 $k_x$  = calibration constant for a specific sensitivity scale

The  $k_x$  factors were determined by submitting known concentrations of chlorophyll *a* to each fluorometer and observing fluorescence units. Linearity of each instrument was checked by serial dilution of pigment extracts. The maximum acid factor was determined by observing fluorescence before and after acidification of extracts of diatom and dinoflagellate cultures sampled in logarithmic phase of growth when phaeo-pigments are absent.

Field measurements were examined station-by-station. Constancy of the shape of each vertical profile was considered and in the few cases where a second deep maximum occurred that was defined by a single observation, a scale reading error was pre-

sumed to have occurred and the value was rejected. In addition, the vertical profile of the acid factor (the proportional change of pigment fluorescence upon acidification of the extract) was examined. This factor normally decreased monotonically from surface layer values near 2 to deep values near 1. Where single significant departures from this pattern occurred, sample values either were justified from the original data, or, if no simple change of values could be supported in terms of analyst error, the observation was rejected. Both apparent errors and rejections were rare, never exceeding 1% on most cruises.

Plant pigments, both chlorophyll *a* and phaeo-pigment concentrations, were integrated to 150 m to express the total pigment under a square meter of sea surface. Only minute quantities of plant pigment occur below 150 m except in the extreme, offshore portion of the survey tracks. The integration programs were based on Simpson's Rule for irregular areas and therefore use a linear interpolation between observations. Where water depth was too shallow to permit sampling to 150 m depth, integration was performed to the deepest sample. Integration values at such stations probably underestimate water column pigment totals since the deepest sample may not be close to the bottom.

#### Phytoplankton production-simulated in situ method

Production measurements were made by a modification of the <sup>14</sup>C method of Steemann-Nielsen (1952). Sampling depths were chosen to span the range of light intensities over which phytoplankton production exceeds respiration. Samples drawn from biologically inert Lexan plastic samplers were placed in 270 ml glass bottles and inoculated with 10 to 20  $\mu$ Ci of sterile NaH<sup>14</sup>CO<sub>2</sub> solution buffered at pH 9.0. Directly after inoculation the samples were loaded into two cylindrical incubators. One incubator was completely opaque and the other was screened to transmit intensity of incident solar radiation present at each of the seven sampled depths. Once loaded, the incubators were placed under full exposure to sun and sky radiation. The time for collecting samples was arranged so that incubation could be started at local apparent noon. Incubation continued until local sunset, when the samples were recovered for filtration. Spectral quality of the light was not controlled. Surface sea water was pumped through the incubators to control temperature.

<sup>1</sup>. Phaeo-pigments are degradation products of chlorophyll *a* and represent detrital pigment.

Filtration was conducted in subdued light at a vacuum not ordinarily exceeding -7 inches of mercury using Millipore membrane filters (2.5 cm diameter) with 0.45- $\mu$ m pore diameters. The filters were then glued to identified copper planchets, placed in perforated pill boxes and stored in desiccators. Upon their return to shore for radioassay, the filters were fumed over concentrated HCl for 10 minutes, sorted by station and returned to their desiccators to await beta-assay. As soon as possible after fuming, samples were turned over to the General Atomic Corporation, La Jolla, California, where they were counted on Beckman wide-beta gas flow counters. Detector end windows were of 80- $\mu$ g/cm<sup>2</sup> Mylar with 5.72-cm diameters. Samples were counted for 3 minutes. Background counts, which usually did not exceed three per minute, were determined and subtracted from each sample count. Counter efficiencies were 27.0 and 34.0% during the entire period of analyses, as determined from point-source standards calibrated against those maintained by the National Bureau of Standards.

Production rate of the phytoplankton was calculated from the equation:

Production (mg C/m<sup>3</sup>/day) =

$$(L - D) \frac{W}{A \times E} \times 1.05 \times 2$$

where: L = counting rate (counts per minute) of the phytoplankton in the light bottle.  
 D = counting rate (counts per minute) of the phytoplankton in the dark bottle.  
 W = weight of carbonate carbon in the water assumed to be 25,000 mg/m<sup>3</sup>.  
 A = discharges per minute by the radiocarbon added to each sample; usually 4.44 X 10<sup>7</sup> dpm  
 E = counter efficiency, determined experimentally for each counter.  
 1.05 = factor to allow for difference in uptake rate of <sup>14</sup>C as compared with that of <sup>12</sup>C  
 2 = correction from the half day incubation period to full day.

Criteria for rejection of production values, while applied over the whole period of the field program, were reworked frequently to ensure consistency. Averaged dark counts were submitted for a single high dark value when the latter occurred at a station, if the corresponding light count was consistent with known profile characteristics. Sets of high dark counts were accepted if they were not erratic (indicating contamination) and if they were well

below the corresponding light counts. Values for entire stations were rejected if the vertical profile was so erratic as to indicate light leaks in incubators, inadvertent exposure to ambient light during filtration, or contamination of the <sup>14</sup>C stock.

Production values for the water column were obtained by computer integration to the euphotic depth. As with plant pigment integrations, Simpson's Rule was used.

### Secchi Depth

Diffuse attenuation of light in the water was estimated from the Secchi depth, determined by lowering a flat 30-cm white disc until it disappeared from sight, then raising the disc until it reappeared and averaging the corresponding depths. Euphotic depth, taken to be at 1% of surface light intensity, was assumed for experimental purposes to occur at 3 X the Secchi depth. (Author's data; R.W. Eppley and J.D.H. Strickland, pers. comm.)

### Chart Preparation

The spatial density of phytoplankton measurements obtained in any single cruise was considered too sparse to warrant contouring the resulting values. A number of stations were sampled repeatedly, however, and sufficient internal agreement between plant pigment and Secchi depth values was obtained from one cruise period to the next to permit pooling of the data into 3-month intervals. The coherence of the contoured patterns that arise from pooling help to confirm the validity of pooling. The particular 3-month intervals presented were chosen to span chronologically the year of observations rather than to represent extremes in the variations encountered. Inspection of the time variations over the region revealed that these pooling intervals were at least as representative of the variations as any other set of 3-month intervals, although more optimal periods might be chosen to represent seasonal change in some sub-areas of the region.

For contouring the charts of plant pigment and Secchi depth distributions, values at stations occupied more than once in the pooling interval were averaged. Contour intervals were selected simply for clarity of representing the distributions: a quasi-geometric progression was necessary for the plant pigment distributions, whereas constant intervals most clearly represented the distribution of water clarity as measured by Secchi depth. In situ production observations were so sparse in time and space as to preclude meaningful contouring.

Computed values by station and cruise of chlorophyll a, phaeo-pigments, production and Secchi depths, from which these charts were prepared, are

reported by Owen and Sanchez (1974) together with insolation and mixed-layer depth.

#### REFERENCES CITED


Biological Methods Panel, NRC Committee on Oceanography, 1969. Recommended procedures for measuring the productivity of plankton standing stock and related ocean properties. Nat. Acad. Sci. Std. book No. 309-01760-2. 59 pp.

Holm-Hansen, O., C.J. Lorenzen, R.W. Holmes and J.D.H. Strickland, 1965. Fluorometric determination of chlorophyll. J. Cons. 30: 3-15.

Lorenzen, C.J., 1966. A method for the continuous measurement of in vivo chlorophyll concentration. Deep-Sea Res. 13: 223-227.


Owen, R.W., Jr. and C.K. Sanchez, 1974. Phytoplankton pigment and production measurements in the California Current region. 1969-72. National Marine Fisheries Service (U.S.) Data Report No. 91.

Steemann-Nielsen, E., 1952. The use of radioactive carbon for measuring organic production in the sea. J. Cons. 18: 117-140.




PRIMARY PRODUCTION  
FEBRUARY - MARCH 1969






PRIMARY PRODUCTION


MAY - JUNE 1969



PRIMARY PRODUCTION  
AUGUST - SEPTEMBER 1969



PRIMARY PRODUCTION  
NOVEMBER - DECEMBER 1969




CHLOROPHYLL *a*

AT THE SEA SURFACE, mg/m<sup>3</sup>

JAN. - FEB. - MAR. 1969

CONTOUR INTERVALS

0.1 mg/m<sup>3</sup>


0.5 mg/m<sup>3</sup> 

(0.05 mg/m<sup>3</sup> contour dashed)


CHLOROPHYLL *a*

SURFACE

JAN. - FEB. - MAR. 1969




CHLOROPHYLL a  
SURFACE  
APR. - MAY - JUN. 1969




CHLOROPHYLL  $a$   
SURFACE  
JUL. - AUG. - SEP. 1969






CHLOROPHYLL a  
SURFACE  
OCT. - NOV. - DEC. 1969




CHLOROPHYLL *a*  
INTEGRATED  
JAN. - FEB. - MAR. 1969







CHLOROPHYLL a  
INTEGRATED  
APR. - MAY - JUN. 1969




CHLOROPHYLL  $a$   
INTEGRATED  
JUL. - AUG. - SEP. 1969




CHLOROPHYLL *a*  
INTEGRATED  
OCT. - NOV. - DEC. 1969




PHAEO PIGMENTS  
JAN. - FEB. - MAR. 1969



PHAEO PIGMENTS  
APR. - MAY - JUN. 1969




PHAEO PIGMENTS  
JUL. - AUG. - SEP. 1969




PHAEO PIGMENTS

OCT. - NOV. - DEC. 1969




SECCHI DEPTH  
JAN. - FEB. - MAR. 1969






SECCHI DEPTH

APR. - MAY - JUN. 1969



SECCHI DEPTH

JUL. - AUG. - SEP. 1969



SECCHI DEPTH  
OCT. - NOV. - DEC. 1969  
CONTOUR INTERVAL 5 METERS

SECCHI DEPTH  
OCT. - NOV. - DEC. 1969

# DISTRIBUTION OF ZOOPLANKTON VOLUMES IN THE CALIFORNIA CURRENT REGION, 1969

Paul E. Smith<sup>1</sup>

Text	
Introduction .....	xv
Charts	
Zooplankton Volumes, 1969 .....	118-125

## Introduction

This report presents contoured charts summarizing displacement volumes of zooplankton samples that were collected systematically during regular CalCOFI cruises carried out in 1969.

The sampling in 1969 differs in two details from that carried out in the years 1951 to 1966 described by Smith (1971): 1. The depth of the standard CalCOFI oblique tow was increased to approximately 210 meters by paying out and retrieving 300 meters of cable at the usual rates of release and return (50 meters per minute out and 20 meters per minute in). 2. The standard CalCOFI net was constructed of a single grade of nylon fabric with mesh apertures of 0.505 mm. Previous nets were constructed of two grades of bolting silk, the anterior section having apertures of 0.55 mm and the posterior section having apertures of 0.25 mm.

A sampling study considering the effects of the two types of nets and the two depths of tow on the

displacement volume was carried out at CalCOFI station 93.30 which lies ten miles west of La Jolla (latitude 32° 50.5' N, longitude 117° 31.0' W, bottom depth 805 meters). The object was to facilitate comparison of the systematic measurements of zooplankton volumes obtained in 1969 with those from previous CalCOFI cruises. Sets of replicate samples were collected at 10, 6 and 2 hours before and after local apparent noon by oblique tows with silk and nylon nets deployed in randomized order from 140 m and 210 m depth. A total of 42 tows were made between 2100 PST June 28 and 0300 PST July 1, 1968 (Table 1). All other field and laboratory procedures used to obtain and process the test samples and the samples from the 1969 cruises follow those described by Kramer et al (1972). Omitting the abnormally large catch by the nylon net taken at 1020PST (838 cc/1000m<sup>3</sup>) the general results (Table 2) indicate that in the test area zooplankton concentrates above 140 meters are much greater proportionately than between 140 m and 210 m depth, in both day and night sampling periods. The day to night ratio of average volume of zooplankton remains virtually the same between the 140 meter silk net tows (D/N=0.797) and the 210 meter nylon tows (D/N = 0.803). However, in terms of cc/1000m<sup>3</sup> the volumes from the 1969 type tows tend to be 25 - 30% less than those from the previous years (Smith 1971), a decrease that will affect the contoured abundance to a limited degree.

1. National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Southwest Fisheries Center  
La Jolla, California 92037

TABLE 1

## A COMPARISON OF PLANKTON VOLUMES BY PRE-1969 AND POST-1969 SAMPLING TECHNIQUES

TOW TYPE	NIGHT					DAY				
	TIME	VOLUME FILTERED	ESTIMATED DEPTH	DISPLACEMENT VOLUME		TIME	VOLUME FILTERED	ESTIMATED DEPTH	DISPLACEMENT VOLUME	
	PST	m <sup>3</sup>	m	cc/1000m <sup>3</sup>	ml m <sup>-2</sup>	PST	m <sup>3</sup>	m	cc/1000m <sup>3</sup>	ml m <sup>-2</sup>
CalCOFI Silk	2153	444	138	347	47.89	1018	436	139	250	34.75
Net to 140 m	0153	438	138	215	29.67	1313	455	135	217	29.30
As 1951-68	0548	429	138	182	25.12	1743	447	136	204	27.74
Type A	2138	441	140	295	41.30	0908	456	140	261	36.54
	0143	433	141	296	41.74	1343	449	141	276	38.92
	0623	432	138	331	45.68	1738	431	138	304	41.95
	2203	454	138	432	59.62					
	0108	422	140	434	60.76					
Mean		436.6	138.9	316.5	43.97		445.7	138.2	252.0	34.87
Standard Deviation		9.9	1.2	90.68	12.63		10.2	2.3	37.13	5.50
Mult. Error Factor				1.36	1.36				1.16	1.18
CalCOFI Silk	2120	617	210	170	35.70	0950	630	213	130	27.69
Net to 210 m	0225	604	208	200	41.60	1345	660	204	129	26.32
Type B	0515	645	207	155	32.09	1815	630	209	255	53.30
(Test)	2110	667	210	336	70.56	0950	643	210	218	45.78
	0215	631	211	212	44.73	1415	641	211	192	40.51
	0515	619	208	205	42.64	1710	656	206	189	38.93
	2140	637	212	308	65.30					
	0140	606	211	271	57.18					
Mean		628.3	209.6	232.1	48.73		643.3	208.8	185.5	38.76
Standard Deviation		21.3	1.8	65.51	14.01		12.6	3.3	49.42	10.40
Mult. Error Factor				1.32	1.33				1.32	1.32
CalCOFI Nylon	2225	695	209	204	42.64	0915	743	201	137	27.54
Net to 210 m	0115	644	207	236	48.85	1420	696	206	124	25.54
Type C	0620	670	212	131	27.77	1715	674	205	172	35.26
As 1969-ff	2205	679	209	196	40.96	1020*	686	209	838*	175.14*
	0110	663	210	244	51.24	1305	679	208	212	44.10
	0550	656	208	243	50.54	1810	665	208	262	54.50
	2105	643	209	274	57.27					
	0205	618	211	280	59.08		690.5	206.2	290.83	60.35
Mean		658.5	209.4	226.0	47.29		27.8	2.9	272.78	57.26
Standard Deviation		23.9	1.6	48.32	10.07				2.01	2.03
Mult. Error Factor				1.28	1.27					
*Omitted from Mean									181.4	37.39
Standard Deviation									56.34	12.04
Mult. Error Factor									1.36	1.37

TABLE 2.


<u>Type of Net and Tow</u>	VOLUME FILTERED	
	<u>Average Volume Strained m<sup>3</sup></u>	
140 meter Silk	440	
210 meter Silk	635	increase 44.3% over 140 m Silk
210 meter Nylon	671	increase 52.5% over 140 m Silk increase 5.7% over 210 m Silk
	CONCENTRATION	
	<u>Average Concentration cc/1000m<sup>3</sup></u>	
140 meter Silk	289	
210 meter Silk	212	decrease 26.6% from 140 m Silk
210 meter Nylon	209*	decrease 27.7% from 140 m Silk decrease 1.4% from 210 m Silk


\*Omitting large 1020 PST day value for 210 meter Nylon net.

## REFERENCES CITED


Smith, P.E., 1971. Distributional atlas of zooplankton volume in the California Current region, 1951 through 1966. CalCOFI Atlas No. 13.


Kramer, D., Mary J. Kalin, Elizabeth G. Stevens, James R. Thraikill and James R. Zweifel, 1972. Collecting and Processing Data on Fish Eggs and Larvae in the California Current Region. National Oceanic and Atmospheric Administration Tech. Rpt. National Marine Fisheries Service. Circ. - 370.























Zooplankton Displacement Volume





These maps are designed to show essential details of the area most intensively studied by the California Cooperative Oceanic Fisheries Investigations. This is approximately the same area as is shown in color on the front cover. Geographical place names are those most commonly used in the various publications emerging from the research. The cardinal station lines extending southwestward from the coast are shown. They are 120 miles apart. Additional lines are utilized as needed and can be as closely spaced as 12 miles apart and still have individual numbers. The stations along the lines are numbered with respect to the station 60 line, the numbers increasing to the west and decreasing to the east. Most of them are 40 miles apart, and are numbered in groups of 10. This permits adding stations as close as 4 miles apart as needed. An example of the usual identification is 120.65. This station is on line 120, 20 nautical miles southwest of station 60.

The projection of the front cover is Lambert's Azimuthal Equal Area Projection. The detail maps are a Mercator projection.

## CONTENTS

W. H. Thomas and D. L. R. Seibert

Distribution of Nitrate, Nitrite, Phosphate and Silicate  
in the California Current region, 1969 ..... vii

Charts 2-97

R. W. Owen Jr.

Distribution of Primary Production, Plant Pigments and  
Secchi Depth in the California Current region, 1969..... xi

Charts 98-117

P. E. Smith

Distribution of Zooplankton Volumes in the  
California Current region, 1969 ..... xv

Charts 118-125