

## About Thermal and Non-thermal Ignition of Nuclear Fusion Reactions

HORA Heinrich<sup>(a)</sup>, KRASA Josef<sup>(b)</sup>, HOFFMANN Dieter H.H.<sup>(c)</sup>, ELIEZER Shalom<sup>(d)</sup>, NISSIM Noaz<sup>(e)</sup>, KIRCHHOFF Jan<sup>(f)</sup>, McKENZIE Warren<sup>(g)</sup>, KIRCHHOFF Götz<sup>(h)</sup>, LALOUSIS Paraskevas<sup>(i)</sup>, MILEY G.H.<sup>(j)</sup>, MARGARONE Daniele<sup>(k)</sup>, PICCIOTTO Antonino<sup>(l)</sup>, GIUFFRIDA Lorenzo<sup>(m)</sup>, CIRRONE Pablo G.A.<sup>(n)</sup>, KORN Georg<sup>(o)</sup>, LIU Bing<sup>(p)</sup>, HO Mark<sup>(q)</sup>,

- (a) Department of Theoretical Physics, University of NSW, Sydney, Australia and HB11 Energy Pty.Ltd. Freshwater, Sydney 2096, Australia, [h.hora@unsw.edu.au](mailto:h.hora@unsw.edu.au)
- (b) Institute of Physics, Czech Academy of Science, Czech Republic [krasa@fzu.cz](mailto:krasa@fzu.cz)
- (c) Department of Nuclear Physics, Tech. University Darmstadt, and Xian Joao Tong University Xian, China [Hoffmann@physik.tu-darmstadt.de](mailto:Hoffmann@physik.tu-darmstadt.de)
- (d) SOREQ Rsearch Center, Yavne, Israel, [Shalom.Eliezer@gmail.com](mailto:Shalom.Eliezer@gmail.com)
- (e) SOREQ Rsearch Center, Yavne, Israel, [noaznissim@gmail.com](mailto:noaznissim@gmail.com)
- (f) HB11 Energy . Freshwater, Sydney 2096, Australia, [jan.kirchhoff@hb11.energy](mailto:jan.kirchhoff@hb11.energy)
- (g) HB11 Energy Freshwater, Sydney 2096, Australia Warren, [McKenzie@HB11.Energy](mailto:McKenzie@HB11.Energy)
- (h) HB11 Energy Freshwater, Sydney 2096, Australia, [goetz.kirchhoff@hb11.energy](mailto:goetz.kirchhoff@hb11.energy)
- (i) Inst. Electronic Structure and lasers FORTH Heraklion, Greece [lalousis@iesl.forth.gr](mailto:lalousis@iesl.forth.gr)
- (j) Dept. Nucl. Plasma & Radiol. Engineering University of Illinois, [gmiley@illinois.edu](mailto:gmiley@illinois.edu)
- (k) Institute of Physics, Czech Acad. Sc., Czech Rep. [Daniele.Margarone@eli-beams.eu](mailto:Daniele.Margarone@eli-beams.eu)
- (l) Fondazione Bruno Kessler, Trento, Italy [picciotto@fbk.eu](mailto:picciotto@fbk.eu)
- (m) Institute of Physics, Czech Acad. Sc., Czech Rep [Lorenzo.Giuffrida@eli-beams.eu](mailto:Lorenzo.Giuffrida@eli-beams.eu)
- (n) INFN Istituto Nazionale de Fisica Nucleare, [pablo.cirrone@lns.infn.it](mailto:pablo.cirrone@lns.infn.it)
- (o) Institute of Physics, Czech Acad. Sc., Czech Rep. [Georg.Korn@eli-beams.eu](mailto:Georg.Korn@eli-beams.eu)
- (p) Hebei Key Laboratory of Compact Fusion, Langfang 065001, China ENN Science and Technology Development Co., Ltd., Langfang 065001, China, [liubing@enn.ch](mailto:liubing@enn.ch)
- (q) ANSTO, Kirrawee 2223, Australia, [mho@ansto.gov.au](mailto:mho@ansto.gov.au)

Dr. Ursula von der Leyen, President of the European Commission, mentioned in her inauguration speech 1st DEC 2019 in Portugal that one of her two priorities is the decarbonisation of energy generation. This is not only to prevent a catastrophe but is a problem of *existential survival*. The attempt to use renewables is acknowledged but for reaching low carbon, this is nearly an impossibly monstrous task. Apart from the expected doubling of energy demand within the next 30 years it is of such a gigantic volume that this can never be fully reached by present measures. One way is thanks to Lord Rutherford and Otto Hahn, to focus on nuclear energy for generators of electricity based on ten million times higher energy per reaction than chemical energy as from carbon. Fission of uranium under extreme control produces now more than 10% of all electricity but the problem of dangerous nuclear radiation or catastrophic accidents as Chernobyl cannot be excluded by 100%.


What remains is nuclear fusion from the energy by joining together very light nuclei to heaviers. This is the energy source of myriades of the stars. The more than million times higher nuclear reaction energy than from burning carbon needs then ignition temperatures of many million degrees Celsius. The sun burns hydrogen into helium at 15 million degrees. To repeat this in power stations is tried since 60 years but

„...Fusion always seems to be 30 years away“, see J.M. Windridge in [1].

A radical change of this situation is now neverless open thanks to getting rid of the temperatures above of the million degrees Celsius by using the just reached ultra-extreme CPA (Chirped Pulse Amplification) laser pulses for LASER BORON FUSION without the problems of dangerous nuclear radiation. The equation of motion for the force density  $\mathbf{f}$  in high-temperature plasmas is given by hydrodynamics for nuclear fusion

$$\mathbf{f} = -\nabla p + \mathbf{f}_{NL} \quad (1)$$

consists of a first term with the gas dynamic pressure  $p$  given by the density and the thermal equilibrium temperature  $T$ . This is in the range below eV (electronvolt = 8135 °C) as chemical energy when burning coal, but is at about ten million times higher for nuclear energy. For ignition, this range is of many dozens of million degree Celsius °C and was well reached with magnetic plasma confinement in tokamaks or stellarators or in aimed billion degrees in reverse field tri-alpha-energy configurations or in laser driven spherical compression with lasers of nanosecond pulses, but for too short times than needed for building a reactor for generating electricity.



**Fig. 1.** Side-on framing camera picture of a plasma produced from an aluminum sphere of  $80 \mu\text{m}$  radius at the time indicated after irradiation from the left hand side by a  $30 \text{ ns}$  ruby laser pulse focused to  $0.4 \text{ mm}$  diameter at the mentioned times after irradiation. The second frame shows a rapidly expanding half-moon-like plasma and a inner spherical part [5].

Apart from the first term with the thermal pressure  $p$  in Eq.(1), the presence of electric and magnetic fields and plasma-optical responses can generate the non-thermal electromagnetic nonlinear force  $f_{\text{NL}}$ . This could directly be seen in the experiments by Sucov et al. [2] at laser interaction in Figure 1 resulting first in the non-transient equation of motion Eq. (10) with Maxwell's plasma stress tensor needing additional nonlinear terms in Schlüter's earlier formulation seen from momentum conservation [3]. Finally for the general transient case clarifying six controversial solutions, the Lorentz and gauge invariant Eq. (6b) [4] was derived.

Evaluating about 500 pictures like Fig. 1 [5] for different laser pulse energies, pulse durations and sphere diameters showed the central plasma thermally expanding with temperatures around  $20 \text{ eV}$  fully following the expected thermal plasma property of the heated sphere taking the varying energy depending on the cases of the interacting laser energy of the cross section of values of about 90%.

The maximum energy of the ions in the half-moons were in the range up to  $5 \text{ keV}$  showing a non-thermal behaviour [5]. This definitely could not be from bremsstrahlung in the corresponding range of  $20 \text{ MeV}$  temperature. Very detailed hydrodynamic computation of plane geometry condition of deuterium plasma at  $10^{17} \text{ W/cm}^2$  laser intensity up to the critical density during  $1.5 \text{ ps}$  laser irradiation showed a plasma block acceleration of  $10^{20} \text{ cm/s}^2$  [6]. This ultrahigh acceleration against the direction of the laser beam was measured from blue Doppler-shifted spectral lines by Sauerbrey [7] using CPA laser

pulses [8] for which discovery the 2018 Physics Nobel Prize was awarded to Donna Strickland and Gerard Mourou. The nonlinear force driven acceleration [7] was about 100,000 times higher than thermal acceleration by the world's largest NIF laser.

One highlight of non-thermal energy density by nonlinear force conditions against the thermal energy density with plasma temperatures of hundred Million  $^\circ\text{C}$  is the measurement by Sven Steinke where  $18 \text{ nm}$  thin diamond films absorbed 99% energy of laser pulses and produces an energy density of  $6.55 \times 10^{12} \text{ J/cm}^3$  [9][10]. This is much higher than the hundred million degrees of thermal equilibrium ignition especially for the very low gains of hydrogen H with the boron isotope 11 (HB11 reaction) that is primarily environmentally clean without radioactivity. This supports the design (Fig. 16 of [11]) for an absolutely clean, safe, low-cost and abundant electricity generator.

The new result is not only that a way was found to a drastic reduction of the carbon energy by remembering the ten million times higher nuclear reaction energy. The HB11 Energy has on top the advantage to be free from radioactivity with keeping the very minor secondary reactions sufficiently easily under control. The main new achievement is that the difficulty of thermal equilibrium of hundred million degrees temperatures for fusion is now overcome by nonlinear forces from CPA laser pulses for using non-thermal ignition [12].

- [1] Miley G.H. and H. Hora. Fusion Science and Technology. 75 (2019) 575
- [2] Sucov EW, Pack JL, Phelps AV and Engelhardt AG. Physics of Fluids 10, (1967) 2035
- [3] Hora, H., (1969) Physics of Fluids 12, (1969) 182
- [4] Hora H., The transient electrodynamic forces at laser-plasma-interaction. Physics of Fluids. 28 (1985) 3705
- [5] Hora H., Experimental results of free targets. Laser Interaction and Related Plasma Phenomena, Plenum Press, New York NY Vol.1, 1971, p. 273, see [6] Fig. 1.2.
- [6] Hora H., Physics of Laser Driven Plasmas, Wiley Press, New York 1981, Fig 10.18 a&b
- [7] Sauerbrey R. Physics of Plasmas 3, (1996) 4712-4716.
- [8] Strickland D. & Mourou G. (1985) Opt.Comm., 56, 2019
- [9] Steinke S., Hening A., Schnürer M., Sokolnik T., Nickles P.V., Jung D., Kiefer D., Hörlein R., Schreiber J., Tajima T., Yan XQ, Hegelich M., Meyer-ter-Vehn J., Sandner W., and Habs D. Laser and Particle Beams 28 (2010) 215
- [10] Hora H., G.H. Miley, S. Eliezer and H. Nissim. High Energy Density Physics 35 (2020) 100739
- [11] Hora Heinrich, Georg Korn, Lorenzo Giuffrida, Daniele Margarone, A. Picciotto, Josef Krása, Karel Jungwirth, Jiri Ullschmied, Paraskevas Lalouis, Shalom Eliezer, George H. Miley, Stavros Moustazis and Gérard Mourou. Laser and Particle Beams 33 (2015) 607
- [12] Giuffrida L., Belloni F., Margarone D., Petringa G., Milluzzo, Scuderi V., Velyhan V., Rosinsky M., Picciotto A., Kucharik M., Dostal M., Dudzak R., Krása, J., Istokskaja V., Catalano R., Tudisco R., Verona, C., Jungwirth, K., Bellutti P., Korn G., Cirrone G.A.P. Physical Review E 101 (2020) 013204