
Convergence analysis for Lasserre’s measure–based hierarchy of upper bounds for polynomial optimization

Etienne de Klerk · Monique Laurent · Zhao Sun

Received: date / Accepted: date

Abstract We consider the problem of minimizing a continuous function f over a compact set \mathbf{K} . We analyze a hierarchy of upper bounds proposed by Lasserre in [*SIAM J. Optim.* 21(3) (2011), pp. 864 – 885], obtained by searching for an optimal probability density function h on \mathbf{K} which is a sum of squares of polynomials, so that the expectation $\int_{\mathbf{K}} f(x)h(x)dx$ is minimized. We show that the rate of convergence is $O(1/\sqrt{r})$, where $2r$ is the degree bound on the density function. This analysis applies to the case when f is Lipschitz continuous and \mathbf{K} is a full-dimensional compact set satisfying some boundary condition (which is satisfied, e.g., for polytopes and the Euclidean ball). The r th upper bound in the hierarchy may be computed using semidefinite programming if f is a polynomial of degree d , and if all moments of order up to $2r + d$ of the Lebesgue measure on \mathbf{K} are known, which holds for example if \mathbf{K} is a simplex, hypercube, or a Euclidean ball.

Keywords Polynomial optimization · Semidefinite optimization · Lasserre hierarchy

Mathematics Subject Classification (2000) 90C22 · 90C26 · 90C30

Etienne de Klerk
Tilburg University
PO Box 90153, 5000 LE Tilburg, The Netherlands
E-mail: E.deKlerk@uvt.nl

Monique Laurent
Centrum Wiskunde & Informatica (CWI), Amsterdam and Tilburg University
CWI, Postbus 94079, 1090 GB Amsterdam, The Netherlands
E-mail: monique@cwi.nl

Zhao Sun
Tilburg University
PO Box 90153, 5000 LE Tilburg, The Netherlands
E-mail: z.sun@uvt.nl

1 Introduction and Preliminaries

1.1 Background

We consider the problem of minimizing a continuous function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ over a compact set $\mathbf{K} \subseteq \mathbb{R}^n$. That is, we consider the problem of computing the parameter:

$$f_{\min, \mathbf{K}} := \min_{x \in \mathbf{K}} f(x).$$

Our main interest will be in the case where f is a polynomial, and \mathbf{K} is defined by polynomial inequalities and equations. For such problems, active research has been done in recent years to construct tractable hierarchies of (upper and lower) bounds for $f_{\min, \mathbf{K}}$, based on using sums of squares of polynomials and semidefinite programming (SDP). The starting point is to reformulate $f_{\min, \mathbf{K}}$ as the problem of finding the largest scalar λ for which the polynomial $f - \lambda$ is nonnegative over K and then to replace the hard positivity condition by a suitable sum of squares decomposition. Alternatively, one may reformulate $f_{\min, \mathbf{K}}$ as the problem of finding a probability measure μ on K minimizing the integral $\int_K f d\mu$. These two dual points of view form the basis of the approach developed by Lasserre [16] for building hierarchies of semidefinite programming based lower bounds for $f_{\min, \mathbf{K}}$ (see also [17, 20] for an overview). Asymptotic convergence to $f_{\min, \mathbf{K}}$ holds (under some mild conditions on the set \mathbf{K}). Moreover, error estimates have been shown in [25, 24] when K is a general basic closed semi-algebraic set, and in [4, 5, 6, 7, 9, 11, 26] for simpler sets like the standard simplex, the hypercube and the unit sphere. In particular, [25] shows that the rate of convergence of the hierarchy of lower bounds based on Schmüdgen's Positivstellensatz is in the order $O(1/\sqrt[r]{2r})$, while [24] shows a convergence rate in $O(1/\sqrt[r]{\log(2r/c)})$ for the (weaker) hierarchy of bounds based on Putinar's Positivstellensatz. Here, c, c' are constants depending only on \mathbf{K} and $2r$ is the selected degree bound. For the case of the hypercube, [4] shows a convergence rate in $O(1/r)$ using Bernstein approximations.

On the other hand, by selecting suitable probability measures on \mathbf{K} , one obtains upper bounds for $f_{\min, \mathbf{K}}$. This approach has been investigated, in particular, for minimization over the standard simplex and when selecting some discrete distributions over the grid points in the simplex. The multinomial distribution is used in [23, 6] to show convergence in $O(1/r)$ and the multivariate hypergeometric distribution is used in [7] to show convergence in $O(1/r^2)$ for quadratic minimization over the simplex (and in the general case assuming a rational minimizer exists).

Additionally, Lasserre [18] shows that, if we fix any measure μ on \mathbf{K} , then it suffices to search for a polynomial density function h which is a sum of squares and minimizes the integral $\int_K f h d\mu$ in order to compute the minimum $f_{\min, \mathbf{K}}$ over \mathbf{K} (see Theorem 1 below). By adding degree constraints on the polynomial density h we get a hierarchy of upper bounds for $f_{\min, \mathbf{K}}$ and our main objective in this paper is to analyze the quality of this hierarchy of upper bounds for $f_{\min, \mathbf{K}}$. Next we will recall this result of Lasserre [18] and then we describe our main results.

1.2 Lasserre's hierarchy of upper bounds

Throughout, $\mathbb{R}[x] = \mathbb{R}[x_1, \dots, x_n]$ is the set of polynomials in n variables with real coefficients, and $\mathbb{R}[x]_r$ is the set of polynomials with degree at most r . $\Sigma[x]$ is the set of sums of squares of polynomials, and $\Sigma[x]_r = \Sigma[x] \cap \mathbb{R}[x]_{2r}$ consists of all sums of squares of polynomials with degree at

most $2r$. We now recall the result of Lasserre [18], which is based on the following characterization for nonnegative continuous functions on a compact set \mathbf{K} .

Theorem 1 [18, Theorem 3.2] *Let $\mathbf{K} \subseteq \mathbb{R}^n$ be compact, let μ be an arbitrary, fixed, finite Borel measure supported by \mathbf{K} , and let f be a continuous function on \mathbb{R}^n . Then, f is nonnegative on \mathbf{K} if and only if*

$$\int_{\mathbf{K}} g^2 f d\mu \geq 0 \quad \forall g \in \mathbb{R}[x].$$

Therefore, the minimum of f over K can be expressed as

$$f_{\min, \mathbf{K}} = \inf_{h \in \Sigma[x]} \int_{\mathbf{K}} h f d\mu \quad \text{s.t.} \quad \int_{\mathbf{K}} h d\mu = 1. \quad (1)$$

Note that formula (1) does not appear explicitly in [18, Theorem 3.2], but one can derive it easily from it. Indeed, one can write $f_{\min, \mathbf{K}} = \sup \{ \lambda : f(x) - \lambda \geq 0 \text{ over } \mathbf{K} \}$. Then, by the first part of Theorem 1, we have $f_{\min, \mathbf{K}} = \sup \{ \lambda : \int_{\mathbf{K}} h(f - \lambda) d\mu \geq 0 \forall h \in \Sigma[x] \}$. As $\int_{\mathbf{K}} h(f - \lambda) d\mu = \int_{\mathbf{K}} h f d\mu - \lambda \int_{\mathbf{K}} h d\mu$, after normalizing $\int_{\mathbf{K}} h d\mu = 1$, we can conclude (1).

If we select the measure μ to be the Lebesgue measure in Theorem 1, then we obtain the following reformulation for $f_{\min, \mathbf{K}}$, which we will consider in this paper:

$$f_{\min, \mathbf{K}} = \inf_{h \in \Sigma[x]} \int_{\mathbf{K}} h(x) f(x) dx \quad \text{s.t.} \quad \int_{\mathbf{K}} h(x) dx = 1.$$

By bounding the degree of the polynomial $h \in \Sigma[x]$ by $2r$, we can define the parameter:

$$\underline{f}_{\mathbf{K}}^{(r)} := \inf_{h \in \Sigma[x]_r} \int_{\mathbf{K}} h(x) f(x) dx \quad \text{s.t.} \quad \int_{\mathbf{K}} h(x) dx = 1. \quad (2)$$

Clearly, the inequality $f_{\min, \mathbf{K}} \leq \underline{f}_{\mathbf{K}}^{(r)}$ holds for any $r \in \mathbb{N}$. Lasserre [18] gives conditions under which the infimum is attained in the program (2).

Theorem 2 [18, Theorems 4.1 and 4.2] *Assume $\mathbf{K} \subseteq \mathbb{R}^n$ is compact and has nonempty interior and let f be a polynomial. Then, the program (2) has an optimal solution for every $r \in \mathbb{N}$ and*

$$\lim_{r \rightarrow \infty} \underline{f}_{\mathbf{K}}^{(r)} = f_{\min, \mathbf{K}}.$$

We now recall how to compute the parameter $\underline{f}_{\mathbf{K}}^{(r)}$ in terms of the moments $m_{\alpha}(\mathbf{K})$ of the Lebesgue measure on \mathbf{K} , where

$$m_{\alpha}(\mathbf{K}) := \int_{\mathbf{K}} x^{\alpha} dx \quad \text{for } \alpha \in \mathbb{N}^n,$$

and $x^{\alpha} := \prod_{i=1}^n x_i^{\alpha_i}$.

Let $N(n, r) := \{\alpha \in \mathbb{N}^n : \sum_{i=1}^n \alpha_i \leq r\}$, and suppose $f(x) = \sum_{\beta \in N(n, d)} f_\beta x^\beta$ has degree d . If we write $h \in \Sigma[x]_r$ as $h(x) = \sum_{\alpha \in N(n, 2r)} h_\alpha x^\alpha$, then the parameter $\underline{f}_{\mathbf{K}}^{(r)}$ from (2) can be reformulated as follows:

$$\begin{aligned} \underline{f}_{\mathbf{K}}^{(r)} &= \min \sum_{\beta \in N(n, d)} f_\beta \sum_{\alpha \in N(n, 2r)} h_\alpha m_{\alpha+\beta}(\mathbf{K}) \\ \text{s.t.} \quad &\sum_{\alpha \in N(n, 2r)} h_\alpha m_\alpha(\mathbf{K}) = 1, \\ &\sum_{\alpha \in N(n, 2r)} h_\alpha x^\alpha \in \Sigma[x]_r. \end{aligned} \quad (3)$$

Hence, if we know the moments $m_\alpha(\mathbf{K})$ for any $\alpha \in \mathbb{N}^n$ with $|\alpha| := \sum_{i=1}^n \alpha_i \leq d + 2r$, then we can compute the parameter $\underline{f}_{\mathbf{K}}^{(r)}$ by solving the semidefinite program (3) which involves a LMI of size $\binom{n+2r}{2r}$.

When \mathbf{K} is the standard simplex $\Delta_n = \{x \in \mathbb{R}_+^n : \sum_{i=1}^n x_i \leq 1\}$, the unit hypercube $\mathbf{Q}_n = [0, 1]^n$, or the unit ball $B_1(0) = \{x \in \mathbb{R}^n : \|x\| \leq 1\}$, there exist explicit formulas for the moments $m_\alpha(\mathbf{K})$. Namely, for the standard simplex, we have

$$m_\alpha(\Delta_n) = \frac{\prod_{i=1}^n \alpha_i!}{(|\alpha| + n)!}, \quad (4)$$

see e.g., [15, equation (2.4)] or [14, equation (2.2)]. From this one can easily calculate the moments for the hypercube \mathbf{Q}_n :

$$m_\alpha(\mathbf{Q}_n) = \int_{\mathbf{Q}_n} x^\alpha dx = \prod_{i=1}^n \int_0^1 x_i^{\alpha_i} dx_i = \prod_{i=1}^n \frac{1}{\alpha_i + 1}.$$

To state the moments for the unit Euclidean ball, we will use the notation $[n] := \{1, \dots, n\}$, the Euler gamma function $\Gamma(\cdot)$, and the notation for the double factorial of an integer k :

$$k!! = \begin{cases} k \cdot (k-2) \cdots 3 \cdot 1, & \text{if } k > 0 \text{ is odd,} \\ k \cdot (k-2) \cdots 4 \cdot 2, & \text{if } k > 0 \text{ is even,} \\ 1 & \text{if } k = 0 \text{ or } k = -1. \end{cases}$$

In terms of this notation, the moments for the unit Euclidean ball are given by:

$$m_\alpha(B_1(0)) = \begin{cases} \frac{\pi^{n/2} \prod_{i=1}^n (\alpha_i - 1)!!}{\Gamma(1 + \frac{n+|\alpha|}{2}) 2^{|\alpha|/2}} = \frac{\pi^{(n-1)/2} 2^{(n+1)/2} \prod_{i=1}^n (\alpha_i - 1)!!}{(n+|\alpha|)!!} & \text{if } \alpha_i \text{ is even for all } i \in [n], \\ 0 & \text{otherwise.} \end{cases} \quad (5)$$

One may prove relation (5) using

$$\int_{B_1(0)} x^\alpha dx = \frac{1}{\Gamma(1 + (n + |\alpha|)/2)} \int_{\mathbb{R}^n} x^\alpha \exp(-\|x\|^2) dx$$

(see e.g. [19, Theorem 2.1]), together with the fact (see e.g., page 872 in [18]) that

$$\int_{-\infty}^{+\infty} t^p \exp(-t^2/2) dt = \begin{cases} \sqrt{2\pi}(p-1)!! & \text{if } p \text{ is even,} \\ 0 & \text{if } p \text{ is odd,} \end{cases}$$

and the identity $\Gamma(1 + \frac{k}{2}) = \frac{k!!}{2^{(k+1)/2}} \sqrt{\pi}$ for any integer $k \in \mathbb{N}$ (see e.g., [1, Section 6.1.12]).

For a general polytope $\mathbf{K} \subseteq \mathbb{R}^n$, it is a hard problem to compute the moments $m_\alpha(\mathbf{K})$. In fact, the problem of computing the volume of polytopes of varying dimensions is already #P-hard [10]. On the other hand, any polytope $\mathbf{K} \subseteq \mathbb{R}^n$ can be triangulated into finitely many simplices (see e.g., [8]) so that one could use (4) to obtain the moments $m_\alpha(\mathbf{K})$ of \mathbf{K} . The complexity of this method depends on the number of simplices in the triangulation. However, this number can be exponentially large (e.g., for the hypercube) and the problem of finding the smallest possible triangulation of a polytope is NP-hard, even in fixed dimension $n = 3$ (see e.g., [8]).

Example

Consider the minimization of the Motzkin polynomial $f(x_1, x_2) = x_1^4 x_2^2 + x_1^2 x_2^4 - 3x_1^2 x_2^2 + 1$ over the hypercube $\mathbf{K} = [-2, 2]^2$, which has four global minimizers at the points $(\pm 1, \pm 1)$, and $f_{\min, \mathbf{K}} = 0$. Figure 1 shows the computed optimal sum of squares density function h^* , for $r = 12$, corresponding to $\underline{f}_{\mathbf{K}}^{(12)} = 0.406076$. We observe that the optimal density h^* shows four peaks at the four global minimizers and thus, it appears to approximate the density of a convex combination of the Dirac measures at the four minimizers.




Fig. 1 Graph and contour plot of $h^*(x)$ on $[-2, 2]^2$ ($r = 12$ and $\deg(h^*) = 24$) for the Motzkin polynomial.

We will present several more numerical examples in Section 4.

1.3 Our main results

In this paper we analyze the quality of the upper bounds $\underline{f}_{\mathbf{K}}^{(r)}$ from (2) for the minimum $f_{\min, \mathbf{K}}$ of f over K . Our main result is an upper bound for the range $\underline{f}_{\mathbf{K}}^{(r)} - f_{\min, \mathbf{K}}$, which applies to the case when f is Lipschitz continuous on \mathbf{K} and when \mathbf{K} is a full-dimensional compact set satisfying the additional condition from Assumption 1, see Theorem 3 below. We will use throughout the following notation about the set \mathbf{K} .

We let $D(\mathbf{K}) = \max_{x, y \in \mathbf{K}} \|x - y\|^2$ denote the diameter of the set \mathbf{K} , where $\|x\| = \sqrt{\sum_{i=1}^n x_i^2}$ is the ℓ_2 -norm. Moreover, $w_{\min}(\mathbf{K})$ is the minimal width of \mathbf{K} , which is the minimum distance between

two distinct parallel supporting hyperplanes of \mathbf{K} . Throughout, $B_\epsilon(a) := \{x \in \mathbb{R}^n : \|x - a\| \leq \epsilon\}$ denotes the Euclidean ball centered at $a \in \mathbb{R}^n$ and with radius $\epsilon > 0$. With γ_n denoting the volume of the n -dimensional unit ball, the volume of the ball $B_\epsilon(a)$ is given by $\text{vol}B_\epsilon(a) = \epsilon^n \gamma_n$.

Assumption 1 *There exist constants $\eta_{\mathbf{K}} > 0$ and $\epsilon_{\mathbf{K}} > 0$ such that, for any point $a \in \mathbf{K}$,*

$$\text{vol}(B_\epsilon(a) \cap \mathbf{K}) \geq \eta_{\mathbf{K}} \text{vol}B_\epsilon(a) = \eta_{\mathbf{K}} \epsilon^n \gamma_n, \quad \text{for all } 0 < \epsilon \leq \epsilon_{\mathbf{K}}. \quad (6)$$

For instance, full-dimensional polytopes and the Euclidean balls satisfy Assumption 1, see Section 5.1 for details. Moreover, for any compact set $\mathbf{K} \subseteq \mathbb{R}^n$ satisfying Assumption 1, define

$$r_{\mathbf{K}} := \max \left\{ \frac{D(\mathbf{K})e}{2\epsilon_{\mathbf{K}}^3}, n \right\} \quad \text{if } \epsilon_{\mathbf{K}} \leq 1 \quad \text{and} \quad r_{\mathbf{K}} := \frac{D(\mathbf{K})e}{2} \quad \text{if } \epsilon_{\mathbf{K}} \geq 1. \quad (7)$$

We can now present our main result.

Theorem 3 *Assume that $\mathbf{K} \subseteq \mathbb{R}^n$ is compact and satisfies Assumption 1. Then there exists a constant $\zeta(\mathbf{K})$ (depending only on \mathbf{K}) such that, for any Lipschitz continuous function f with Lipschitz constant M_f on \mathbf{K} , the following inequality holds:*

$$\underline{f}_{\mathbf{K}}^{(r)} - f_{\min, \mathbf{K}} \leq \frac{\zeta(\mathbf{K})M_f}{\sqrt{r}} \quad \text{for any } r \geq r_{\mathbf{K}}. \quad (8)$$

Moreover, if f is a polynomial of degree d and \mathbf{K} is a convex body, then

$$\underline{f}_{\mathbf{K}}^{(r)} - f_{\min, \mathbf{K}} \leq \frac{2d^2 \zeta(\mathbf{K}) \sup_{x \in \mathbf{K}} |f(x)|}{w_{\min}(\mathbf{K}) \sqrt{r}} \quad \text{for any } r \geq r_{\mathbf{K}}. \quad (9)$$

The key idea to show this result is to select suitable sums of squares densities which we are able to analyse. For this, we will select a global minimizer a of f over K and consider the Gaussian distribution with mean a and, as sums of squares densities, we will select the polynomials $H_{r,a}$ obtained by truncating the Taylor series expansion of the Gaussian distribution, see relation (14).

1.4 Contents of the paper

Our paper is organized as follows. In Section 2, we give a constructive proof for our main result Theorem 3. In Section 3 we show how to obtain feasible points in \mathbf{K} that correspond to the bounds $\underline{f}_{\mathbf{K}}^{(r)}$ through sampling. This is followed by a section with numerical examples (Section 4). Finally, in the concluding remarks (Section 5), we revisit Assumption 1, and discuss computational perspectives of the approach studied here.

2 Proof of our main result in Theorem 3

In this section we prove our main result in Theorem 3. Our analysis will hold for Lipschitz continuous f , so we will start by reviewing some relevant properties in Section 2.1. In the next step we indicate in Section 2.2 how to select the polynomial density function h as a special sum of squares that we will be able to analyze. Namely, we let a denote a global minimizer of the function f over the set $\mathbf{K} \subseteq \mathbb{R}^n$. Then we consider the density function G_a in (12) of the Gaussian distribution with mean a and the polynomial $H_{r,a}$ in (14), which is obtained from the truncation at degree $2r$ of the Taylor series expansion of the Gaussian density function G_a . The final step will be to analyze the quality of the bound obtained by selecting the polynomial $H_{r,a}$ and this will be the most technical part of the proof, carried out in Section 2.3.

2.1 Lipschitz continuous functions

A function f is said to be Lipschitz continuous on \mathbf{K} , with Lipschitz constant M_f , if it satisfies:

$$|f(y) - f(x)| \leq M_f \|y - x\| \quad \text{for all } x, y \in \mathbf{K}.$$

If f is continuous and differentiable on \mathbf{K} , then f is Lipschitz continuous on \mathbf{K} with respect to the constant

$$M_f = \max_{x \in \mathbf{K}} \|\nabla f(x)\|. \quad (10)$$

Furthermore, if f is an n -variate polynomial with degree d , then the Markov inequality for f on a convex body \mathbf{K} reads as

$$\max_{x \in \mathbf{K}} \|\nabla f(x)\| \leq \frac{2d^2}{w_{\min}(\mathbf{K})} \sup_{x \in \mathbf{K}} |f(x)|,$$

see e.g., [3, relation (8)]. Thus, together with (10), we have that f is Lipschitz continuous on \mathbf{K} with respect to the constant

$$M_f \leq \frac{2d^2}{w_{\min}(\mathbf{K})} \sup_{x \in \mathbf{K}} |f(x)|. \quad (11)$$

2.2 Choosing the polynomial density function $H_{r,a}$

Consider the function

$$G_a(x) := \frac{1}{(2\pi\sigma^2)^{n/2}} \exp\left(-\frac{\|x-a\|^2}{2\sigma^2}\right), \quad (12)$$

which is the probability density function of the Gaussian distribution with mean a and standard variance σ (whose value will be defined later). Let the constant $C_{\mathbf{K},a}$ be defined by

$$\int_{\mathbf{K}} C_{\mathbf{K},a} G_a(x) dx = 1. \quad (13)$$

Observe that $G_a(x)$ is equal to the function $\frac{1}{(2\pi\sigma^2)^{n/2}} e^{-t}$ evaluated at the point $t = \frac{\|x-a\|^2}{2\sigma^2}$.

Denote by $H_{r,a}$ the Taylor series expansion of G_a truncated at the order $2r$. That is,

$$H_{r,a}(x) = \frac{1}{(2\pi\sigma^2)^{n/2}} \sum_{k=0}^{2r} \frac{1}{k!} \left(-\frac{\|x-a\|^2}{2\sigma^2} \right)^k. \quad (14)$$

Moreover consider the constant $c_{\mathbf{K},a}^r$, defined by

$$\int_{\mathbf{K}} c_{\mathbf{K},a}^r H_{r,a}(x) dx = 1. \quad (15)$$

The next step is to show that $H_{r,a}$ is a sum of squares of polynomials and thus $H_{r,a} \in \Sigma[x]_{2r}$. This follows from the next lemma.

Lemma 1 *Let $\phi_{2r}(t)$ denote the (univariate) polynomial of degree $2r$ obtained by truncating the Taylor series expansion of e^{-t} at the order $2r$. That is,*

$$\phi_{2r}(t) := \sum_{k=0}^{2r} \frac{(-t)^k}{k!}.$$

Then ϕ_{2r} is a sum of squares of polynomials. Moreover, we have

$$0 \leq \phi_{2r}(t) - e^{-t} \leq \frac{t^{2r+1}}{(2r+1)!} \quad \text{for all } t \geq 0. \quad (16)$$

Proof First, we show that ϕ_{2r} is a sum of squares. As ϕ_{2r} is a univariate polynomial, by Hilbert's Theorem (see e.g., [20, Theorem 3.4]), it suffices to show that $\phi_{2r}(t) \geq 0$ for any $t \in \mathbb{R}$. As $\phi_{2r}(-\infty) = \phi_{2r}(+\infty) = +\infty$, it suffices to show that $\phi_{2r}(t) \geq 0$ at all the stationary points t where $\phi'_{2r}(t) = 0$. For this, observe that $\phi'_{2r}(t) = \sum_{k=1}^{2r} (-1)^k \frac{t^{k-1}}{(k-1)!}$, so that it can be written as $\phi'_{2r}(t) = -\phi_{2r}(t) + \frac{t^{2r}}{(2r)!}$. Hence, for any t with $\phi'_{2r}(t) = 0$, we have $\phi_{2r}(t) = \frac{t^{2r}}{(2r)!} \geq 0$.

Next, we show that $\phi_{2r}(t) \geq e^{-t}$ for all $t \geq 0$. Fix $t \geq 0$. Then, by Taylor Theorem (see e.g., [30]), one has $e^{-t} = \phi_{2r}(t) + \frac{\phi^{(2r+1)}(\xi)t^{2r+1}}{(2r+1)!}$ for some $\xi \in [0, t]$. As $\phi^{(2r+1)}(\xi) = -e^{-\xi}$, one can conclude that $e^{-t} - \phi_{2r}(t) = -\frac{e^{-\xi}t^{2r+1}}{(2r+1)!} \leq 0$ and $e^{-t} - \phi_{2r}(t) \geq -\frac{t^{2r+1}}{(2r+1)!}$. \square

We now consider the parameter $f_{\mathbf{K},a}^{(r)}$ defined as

$$f_{\mathbf{K},a}^{(r)} := \int_{\mathbf{K}} f(x) c_{\mathbf{K},a}^r H_{r,a}(x) dx. \quad (17)$$

Our main technical result is the following upper bound for the range $f_{\mathbf{K},a}^{(r)} - f_{\min,\mathbf{K}}$, whose proof is given in Section 2.3 below. Theorem 3 follows then as a direct application of Theorem 4.

Theorem 4 *Assume $\mathbf{K} \subseteq \mathbb{R}^n$ is compact and satisfies Assumption 1, and consider the parameter $r_{\mathbf{K}}$ from (7). Then there exists a constant $\zeta(\mathbf{K})$ (depending only on \mathbf{K}) such that, for any Lipschitz continuous function f with Lipschitz constant M_f on \mathbf{K} , the following inequality holds:*

$$f_{\mathbf{K},a}^{(r)} - f_{\min,\mathbf{K}} \leq \frac{\zeta(\mathbf{K})M_f}{\sqrt{2r+1}}, \quad \text{for any } r \geq \frac{r_{\mathbf{K}}}{2}. \quad (18)$$

Moreover, if f is a polynomial of degree d and \mathbf{K} is a convex body, then

$$f_{\mathbf{K},a}^{(r)} - f_{\min,\mathbf{K}} \leq \frac{2d^2 \zeta(\mathbf{K}) \sup_{x \in \mathbf{K}} |f(x)|}{w_{\min}(\mathbf{K}) \sqrt{2r+1}}, \quad \text{for any } r \geq \frac{r_{\mathbf{K}}}{2}. \quad (19)$$

Proof (of Theorem 3) Assume f is Lipschitz continuous with Lipschitz constant M_f on K and a is a minimizer of f over the set \mathbf{K} . Using the definitions (2) and (17) of the parameters and the fact that $H_{r,a}$ is a sum of squares with degree $4r$, it follows that

$$\underline{f}_{\mathbf{K}}^{(2r+1)} \leq \underline{f}_{\mathbf{K}}^{(2r)} \leq f_{\mathbf{K},a}^{(r)}, \quad \text{for any } r \in \mathbb{N}.$$

Then, from inequality (18) in Theorem 4, one obtains

$$\underline{f}_{\mathbf{K}}^{(2r+1)} - f_{\min,\mathbf{K}} \leq \underline{f}_{\mathbf{K}}^{(2r)} - f_{\min,\mathbf{K}} \leq f_{\mathbf{K},a}^{(r)} - f_{\min,\mathbf{K}} \leq \frac{\zeta(\mathbf{K})M_f}{\sqrt{2r+1}} \quad \text{for any } r \geq \frac{r_{\mathbf{K}}}{2}.$$

Hence, for any $r \geq r_{\mathbf{K}}$,

$$\begin{aligned} \underline{f}_{\mathbf{K}}^{(r)} - f_{\min,\mathbf{K}} &\leq \frac{\zeta(\mathbf{K})M_f}{\sqrt{r+1}} \leq \frac{\zeta(\mathbf{K})M_f}{\sqrt{r}} \quad \text{for even } r, \\ \underline{f}_{\mathbf{K}}^{(r)} - f_{\min,\mathbf{K}} &\leq \frac{\zeta(\mathbf{K})M_f}{\sqrt{r}} \quad \text{for odd } r. \end{aligned}$$

This concludes the proof for relation (8), and relation (9) follows from (19) in an analogous way. This finishes the proof of Theorem 3. \square

2.3 Analyzing the polynomial density function $H_{r,a}$

In this section we prove the result of Theorem 4. Recall that a is a global minimizer of f over \mathbf{K} . For the proof, we will need the following four technical lemmas.

Lemma 2 *Assume $\mathbf{K} \subseteq \mathbb{R}^n$ is compact and satisfies Assumption 1. Then, for any $0 < \epsilon \leq \epsilon_{\mathbf{K}}$ and $r \in \mathbb{N}$, we have:*

$$c_{\mathbf{K},a}^r \leq C_{\mathbf{K},a} \leq \frac{(2\pi\sigma^2)^{n/2} \exp\left(\frac{\epsilon^2}{2\sigma^2}\right)}{\eta_{\mathbf{K}} \epsilon^n \gamma_n}. \quad (20)$$

Proof By Lemma 1, $\phi_{2r}(t) \geq e^{-t}$ for all $t \geq 0$, which implies $H_{r,a}(x) \geq G_a(x)$ for all $x \in \mathbb{R}^n$. Together with the relations (13) and (15) defining the constants $C_{\mathbf{K},a}$ and $c_{\mathbf{K},a}^r$, we deduce that $c_{\mathbf{K},a}^r \leq C_{\mathbf{K},a}$. Moreover, by the definition (13) of the constant $C_{\mathbf{K},a}$, one has

$$\begin{aligned} \frac{1}{C_{\mathbf{K},a}} &= \int_{\mathbf{K}} G_a(x) dx = \int_{\mathbf{K}} \frac{1}{(2\pi\sigma^2)^{n/2}} \exp\left(-\frac{\|x-a\|^2}{2\sigma^2}\right) dx \\ &\geq \int_{\mathbf{K} \cap B_{\epsilon}(a)} \frac{1}{(2\pi\sigma^2)^{n/2}} \exp\left(-\frac{\|x-a\|^2}{2\sigma^2}\right) dx \\ &\geq \frac{1}{(2\pi\sigma^2)^{n/2}} \exp\left(-\frac{\epsilon^2}{2\sigma^2}\right) \text{vol}(\mathbf{K} \cap B_{\epsilon}(a)). \end{aligned}$$

We now use relation (6) from Assumption 1 in order to conclude that $\text{vol}(\mathbf{K} \cap B_\epsilon(a)) \geq \eta_{\mathbf{K}} \epsilon^n \gamma_n$, which gives the desired upper bound on $C_{K,a}$. \square

Lemma 3 *Given $\tilde{x} \in \mathbb{R}^n$ and a function $F : \mathbb{R}_+ \rightarrow \mathbb{R}$, define the function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ by $f(x) = F(\|x - \tilde{x}\|)$ for any $x \in \mathbb{R}^n$. Then, for any $\rho_2 \geq \rho_1 \geq 0$, one has*

$$\int_{B_{\rho_2}(\tilde{x}) \setminus B_{\rho_1}(\tilde{x})} f(x) dx = n \gamma_n \int_{\rho_1}^{\rho_2} z^{n-1} F(z) dz,$$

where $\gamma_n = \frac{\pi^{(n-1)/2} 2^{(n+1)/2}}{n!!}$ is the volume of the unit Euclidean ball in \mathbb{R}^n .

Proof Apply a change of variables using spherical coordinates as explained, e.g., in [2]. \square

Lemma 4 *For any positive integers r and n , one has $\left(\frac{1}{2r+1}\right)^{-\frac{n}{4(2r+1)+2n}} < 6n$.*

Proof Let $n \in \mathbb{N}$ be given. Denote

$$g(r) := \left(\frac{1}{2r+1}\right)^{-\frac{n}{4(2r+1)+2n}} = (2r+1)^{\frac{n}{4(2r+1)+2n}} \quad (r \geq 0).$$

Observe that, $g(0) = 1$, $g(r) > 0$ for all $r \geq 0$, $\ln(g(r)) = \frac{n}{8r+4+2n} \ln(2r+1)$, and thus $\lim_{r \rightarrow \infty} g(r) = 1$. It suffices to show $g(r^*) < 6n$ for any stationary point r^* . Since

$$\frac{d \ln(g(r))}{dr} = \frac{-8n \ln(2r+1)}{(8r+4+2n)^2} + \frac{2n}{(2r+1)(8r+4+2n)},$$

and $g'(r) = \frac{1}{g(r)} \frac{d \ln(g(r))}{dr}$, any stationary point r^* satisfies

$$\frac{d \ln(g(r^*))}{dr} = 0 \iff (2r^*+1) [\ln(2r^*+1) - 1] = \frac{n}{2}.$$

Since

$$(2r^*+1)(\ln(3) - 1) \leq (2r^*+1) [\ln(2r^*+1) - 1] = \frac{n}{2},$$

one has $2r^*+1 \leq \frac{n}{2(\ln(3)-1)} < 6n$. Since $g(r) \leq 2r+1$ for all $r \geq 0$, one has $g(r^*) \leq 2r^*+1 < 6n$. \square

Lemma 5 *Assume $\mathbf{K} \subseteq \mathbb{R}^n$ is compact and satisfies Assumption 1. Then, for any $0 < \epsilon \leq \epsilon_{\mathbf{K}}$, one has*

$$\int_{\mathbf{K}} C_{\mathbf{K},a} \|x - a\| G_a(x) dx \leq \epsilon + \frac{n \sigma^{n+1} p(n)}{\epsilon^n \eta_{\mathbf{K}}} e^{\frac{\epsilon^2}{2\sigma^2}},$$

where $p(n) := \int_0^{+\infty} t^n e^{-t^2/2} dt$ is a constant depending on n , given by

$$p(n) = \begin{cases} 1 & \text{if } n = 1, \\ \sqrt{\frac{\pi}{2}} \prod_{j=1}^k (2j-1) & \text{if } n = 2k \text{ and } k \geq 1, \\ \prod_{j=1}^k (2j) & \text{if } n = 2k+1 \text{ and } k \geq 1. \end{cases} \quad (21)$$

Proof Let $\varphi := \int_{\mathbf{K}} C_{\mathbf{K},a} \|x - a\| G_a(x) dx$ denote the integral that we need to upper bound. We split the integral φ as $\varphi = \varphi_1 + \varphi_2$, depending on whether x lies in the ball $B_\epsilon(a)$ or not.

First, we upper bound the term φ_1 as

$$\varphi_1 := \int_{\mathbf{K} \cap B_\epsilon(a)} \|x - a\| C_{\mathbf{K},a} G_a(x) dx \leq \epsilon \int_{\mathbf{K} \cap B_\epsilon(a)} C_{\mathbf{K},a} G_a(x) dx \leq \epsilon \int_{\mathbf{K}} C_{\mathbf{K},a} G_a(x) dx = \epsilon.$$

Second, we bound the integral

$$\varphi_2 := C_{\mathbf{K},a} \int_{\mathbf{K} \setminus B_\epsilon(a)} \|x - a\| G_a(x) dx.$$

Since $\mathbf{K} \subseteq B_{\sqrt{D(\mathbf{K})}}(a)$, one has

$$\varphi_2 \leq C_{\mathbf{K},a} \int_{B_{\sqrt{D(\mathbf{K})}}(a) \setminus B_\epsilon(a)} \|x - a\| G_a(x) dx,$$

where the right hand side, by Lemma 3, is equal to

$$\frac{C_{\mathbf{K},a} n \gamma_n}{(2\pi\sigma^2)^{n/2}} \int_{\epsilon}^{\sqrt{D(\mathbf{K})}} z^n \exp\left(-\frac{z^2}{2\sigma^2}\right) dz.$$

By a change of variable $t = \frac{z}{\sigma}$, one obtains

$$\varphi_2 \leq \frac{C_{\mathbf{K},a} n \gamma_n \sigma}{(2\pi)^{n/2}} \int_{\epsilon/\sigma}^{\sqrt{D(\mathbf{K})}/\sigma} t^n \exp\left(-\frac{t^2}{2}\right) dt,$$

and thus

$$\varphi_2 \leq \frac{C_{\mathbf{K},a} n \gamma_n \sigma}{(2\pi)^{n/2}} \int_0^{+\infty} t^n \exp\left(-\frac{t^2}{2}\right) dt = \frac{C_{\mathbf{K},a} n \gamma_n \sigma}{(2\pi)^{n/2}} p(n).$$

Here we have set $p(n) := \int_0^{+\infty} t^n e^{-\frac{t^2}{2}} dt$ which can be checked to be given by (21) (e.g., using induction on n). Now, combining with the upper bound for $C_{\mathbf{K},a}$ from (20), we obtain

$$\varphi_2 \leq \frac{n\sigma^{n+1} p(n)}{\epsilon^n \eta_{\mathbf{K}}} e^{-\frac{\epsilon^2}{2\sigma^2}}.$$

Therefore, we have shown:

$$\varphi = \varphi_1 + \varphi_2 \leq \epsilon + \frac{n\sigma^{n+1} p(n)}{\epsilon^n \eta_{\mathbf{K}}} e^{-\frac{\epsilon^2}{2\sigma^2}},$$

which shows the lemma. \square

We are now ready to prove Theorem 4.

Proof (of Theorem 4) Observe that, if f is a polynomial, then we can use the upper bound (11) for its Lipschitz constant and thus the inequality (19) follows as a direct consequence of the inequality (18). Therefore, it suffices to show the relation (18).

Recall that a is a minimizer of f over \mathbf{K} . As f is Lipschitz continuous with Lipschitz constant M_f on K , we have

$$f(x) - f(a) \leq M_f \|x - a\| \quad \forall x \in \mathbf{K}.$$

This implies

$$f_{\mathbf{K},a}^{(r)} - f_{\min,\mathbf{K}} = \int_{\mathbf{K}} c_{\mathbf{K},a}^r H_{r,a}(x) (f(x) - f(a)) dx \leq M_f \int_{\mathbf{K}} \|x - a\| c_{\mathbf{K},a}^r H_{r,a}(x) dx.$$

Our objective is now to show the existence of a constant $\zeta(\mathbf{K})$ such that

$$\psi := \int_{\mathbf{K}} c_{\mathbf{K},a}^r \|x - a\| H_{r,a}(x) dx \leq \frac{\zeta(\mathbf{K})}{\sqrt{2r+1}}, \quad \text{for any } r \geq r_{\mathbf{K}}, \text{ (see (7))}$$

by which we can then conclude the proof for (18).

For this, we split the integral ψ as the sum of two terms:

$$\psi = \underbrace{\int_{\mathbf{K}} c_{\mathbf{K},a}^r \|x - a\| G_a(x) dx}_{=: \psi_1} + \underbrace{\int_{\mathbf{K}} c_{\mathbf{K},a}^r \|x - a\| (H_{r,a}(x) - G_a(x)) dx}_{=: \psi_2}.$$

First, we upper bound the term ψ_1 . As $c_{\mathbf{K},a}^r \leq C_{\mathbf{K},a}$ (by (20)), we can use Lemma 5 to conclude that, for any $0 < \epsilon \leq \epsilon_{\mathbf{K}}$,

$$\psi_1 \leq \int_{\mathbf{K}} C_{\mathbf{K},a} \|x - a\| G_a(x) dx \leq \epsilon + \frac{n\sigma^{n+1}p(n)}{\epsilon^n \eta_{\mathbf{K}}} e^{\frac{\epsilon^2}{2\sigma^2}} = \epsilon \underbrace{\left[1 + \frac{n\sigma^{n+1}p(n)}{\epsilon^{n+1}\eta_{\mathbf{K}}} e^{\frac{\epsilon^2}{2\sigma^2}} \right]}_{=: \mu_1} = \epsilon \mu_1. \quad (22)$$

Second we bound the integral

$$\psi_2 = \int_{\mathbf{K}} c_{\mathbf{K},a}^r \|x - a\| (H_{r,a}(x) - G_a(x)) dx.$$

We can upper bound the function $H_{r,a}(x) - G_a(x)$ using the estimate from (16) and we get

$$H_{r,a}(x) - G_a(x) \leq \frac{1}{(2\pi\sigma^2)^{n/2}} \frac{\|x - a\|^{4r+2}}{(2\sigma^2)^{2r+1} (2r+1)!}.$$

Then we have

$$\psi_2 \leq \frac{1}{(2\pi\sigma^2)^{n/2}} \int_{\mathbf{K}} c_{\mathbf{K},a}^r \frac{\|x - a\|^{4r+3}}{(2\sigma^2)^{2r+1} (2r+1)!} dx = \frac{1}{(2\pi\sigma^2)^{n/2}} \frac{c_{\mathbf{K},a}^r}{(2\sigma^2)^{2r+1} (2r+1)!} \int_{\mathbf{K}} \|x - a\|^{4r+3} dx.$$

Now we upper bound the integral $\int_{\mathbf{K}} \|x - a\|^{4r+3} dx$. Since $\mathbf{K} \subseteq B_{\sqrt{D(\mathbf{K})}}(a)$, one has

$$\int_{\mathbf{K}} \|x - a\|^{4r+3} dx \leq \int_{B_{\sqrt{D(\mathbf{K})}}(a)} \|x - a\|^{4r+3} dx,$$

where the right hand side, by Lemma 3, is equal to

$$n\gamma_n \int_0^{\sqrt{D(\mathbf{K})}} z^{4r+n+2} dz = \frac{n\gamma_n D(\mathbf{K})^{\frac{4r+n+3}{2}}}{4r+n+3} \leq n\gamma_n D(\mathbf{K})^{\frac{4r+n+3}{2}}.$$

Thus, we obtain

$$\psi_2 \leq \frac{1}{(2\pi\sigma^2)^{n/2}} \frac{c_{\mathbf{K},a}^r}{(2\sigma^2)^{2r+1}(2r+1)!} n\gamma_n D(\mathbf{K})^{\frac{4r+n+3}{2}}.$$

We now use the upper bound for $c_{\mathbf{K},a}^r$ from (20):

$$c_{\mathbf{K},a}^r \leq \frac{(2\pi\sigma^2)^{n/2} \exp\left(\frac{\epsilon^2}{2\sigma^2}\right)}{\eta_{\mathbf{K}} \epsilon^n \gamma_n}$$

and we obtain

$$\psi_2 \leq \frac{n \exp\left(\frac{\epsilon^2}{2\sigma^2}\right) D(\mathbf{K})^{\frac{4r+n+3}{2}}}{\eta_{\mathbf{K}} \epsilon^n (2r+1)! (2\sigma^2)^{2r+1}}.$$

Finally we use the Stirling's inequality:

$$(2r+1)! \geq \sqrt{2\pi(2r+1)} \left(\frac{2r+1}{e}\right)^{2r+1},$$

and obtain

$$\begin{aligned} \psi_2 &\leq \underbrace{\frac{n \exp\left(\frac{\epsilon^2}{2\sigma^2}\right) D(\mathbf{K})^{\frac{n+1}{2}}}{\eta_{\mathbf{K}}}}_{=: \mu_2} \left(\frac{D(\mathbf{K})e}{2\sigma^2 \epsilon^{n/(2r+1)} (2r+1)}\right)^{2r+1} \frac{1}{\sqrt{2\pi(2r+1)}} \\ &= \frac{\mu_2}{\sqrt{2\pi(2r+1)}} \left(\frac{D(\mathbf{K})e}{2\sigma^2 \epsilon^{n/(2r+1)} (2r+1)}\right)^{2r+1}. \end{aligned} \quad (23)$$

We can now upper bound the quantity $\psi = \psi_1 + \psi_2$, by combining the upper bound for ψ_1 in (22) with the above upper bound (23) for ψ_2 . That is,

$$\psi \leq \epsilon\mu_1 + \frac{\mu_2}{\sqrt{2\pi(2r+1)}} \left(\frac{D(\mathbf{K})e}{2\sigma^2 \epsilon^{n/(2r+1)} (2r+1)}\right)^{2r+1}.$$

We now indicate how to select the parameters ϵ and σ .

First we select $\sigma = \epsilon$, so that both parameters μ_1 and μ_2 appearing in (22) and (23) are constants depending on n and \mathbf{K} , namely

$$\mu_1 = 1 + \frac{np(n)e^{1/2}}{\eta_{\mathbf{K}}} \quad \text{and} \quad \mu_2 = \frac{ne^{1/2}D(\mathbf{K})^{\frac{n+1}{2}}}{\eta_{\mathbf{K}}}.$$

Next we select ϵ so that $\frac{D(\mathbf{K})e}{2\epsilon^{2+n/(2r+1)}(2r+1)} = 1$, i.e.,

$$\epsilon = \left(\frac{D(\mathbf{K})e}{2(2r+1)} \right)^{\frac{2r+1}{2(2r+1)+n}} = \left(\frac{D(\mathbf{K})e}{2} \right)^{\frac{2r+1}{2(2r+1)+n}} \left(\frac{1}{2r+1} \right)^{\frac{1}{2} - \frac{n}{4(2r+1)+2n}}.$$

Summarizing, we have shown that

$$\begin{aligned} \psi &\leq \left(\frac{1}{2r+1} \right)^{\frac{1}{2} - \frac{n}{4(2r+1)+2n}} \left[\left(\frac{D(\mathbf{K})e}{2} \right)^{\frac{2r+1}{2(2r+1)+n}} \mu_1 + \frac{\mu_2}{\sqrt{2\pi}} \left(\frac{1}{2r+1} \right)^{\frac{n}{4(2r+1)+2n}} \right] \\ &\leq \left(\frac{1}{2r+1} \right)^{\frac{1}{2}} 6n \left(\mu_1 \max \left\{ 1, \sqrt{\frac{D(\mathbf{K})e}{2}} \right\} + \frac{\mu_2}{\sqrt{2\pi}} \right). \end{aligned} \quad (24)$$

To obtain the last inequality (24), we use the inequality $\left(\frac{1}{2r+1} \right)^{-\frac{n}{4(2r+1)+2n}} < 6n$ (recall Lemma 4), together with the two inequalities $\left(\frac{D(\mathbf{K})e}{2} \right)^{\frac{2r+1}{2(2r+1)+n}} \leq \max \left\{ 1, \sqrt{\frac{D(\mathbf{K})e}{2}} \right\}$ and $\left(\frac{1}{2r+1} \right)^{\frac{n}{4(2r+1)+2n}} \leq 1$.

Since we have assumed $\epsilon \leq \epsilon_{\mathbf{K}}$ (recall Lemma 2), this implies the condition $r \geq \frac{D(\mathbf{K})e}{4} \epsilon_{\mathbf{K}}^{-(2+\frac{n}{2r+1})} - \frac{1}{2}$, i.e., the inequality (24) holds for any $r \geq \frac{D(\mathbf{K})e}{4} \epsilon_{\mathbf{K}}^{-(2+\frac{n}{2r+1})} - \frac{1}{2}$. If $\epsilon_{\mathbf{K}} \leq 1$ and $r \geq n/2$, then we have $\epsilon_{\mathbf{K}}^{-(2+\frac{n}{2r+1})} \leq \epsilon_{\mathbf{K}}^{-3}$ and thus the inequality (24) holds for any $r \geq \max \left\{ \frac{D(\mathbf{K})e}{4\epsilon_{\mathbf{K}}^3}, \frac{n}{2} \right\}$. If $\epsilon_{\mathbf{K}} \geq 1$ then $\epsilon_{\mathbf{K}}^{-(2+\frac{n}{2r+1})} \leq 1$ and thus (24) holds for any integer $r \geq \frac{D(\mathbf{K})e}{4}$. Hence, the inequality (24) holds for any $r \geq r_{\mathbf{K}}/2$, where $r_{\mathbf{K}}$ is as defined in (7).

Finally, by defining the constant

$$\zeta(\mathbf{K}) := 6n \left(\mu_1 \max \left\{ 1, \sqrt{\frac{D(\mathbf{K})e}{2}} \right\} + \frac{\mu_2}{\sqrt{2\pi}} \right),$$

which indeed depends only on \mathbf{K} and its dimension n , we can conclude the proof for (18). \square

Remark 1 Note that in the proof of Theorem 4, we use Assumption 1 only for the selected minimizer $a \in \mathbf{K}$ (and we use it only in the proof of Lemma 2). Hence, if the selected point a lies in the interior of \mathbf{K} , i.e., if there exists $\delta > 0$ such that $B_\delta(a) \subseteq \mathbf{K}$, then the result of Theorem 4 (and thus Theorem 3) holds when selecting $\eta_{\mathbf{K}} = 1$ and $\epsilon_{\mathbf{K}} = \delta$.

Our results extend also to unconstrained global minimization:

$$f^* := \min_{x \in \mathbb{R}^n} f(x),$$

if we know that f has a global minimizer a and we know a ball $B_\delta(0)$ containing a . We can then indeed minimize f over a compact set K , which can be chosen to be the ball $B_\delta(0)$ or a suitable hypercube containing a .

3 Obtaining feasible solutions through sampling

In this section we indicate how to sample feasible points in the set \mathbf{K} from the optimal density function obtained by solving the semidefinite program (2).

Let $f \in \mathbb{R}[x]$ be a polynomial. Suppose $h^*(x) \in \Sigma[x]_r$ is an optimal solution of the program (2), i.e., $\underline{f}_{\mathbf{K}}^{(r)} = \int_{\mathbf{K}} f(x)h^*(x)dx$ and $\int_{\mathbf{K}} h^*(x)dx = 1$. Then h^* can be seen as the probability density function of a probability distribution on \mathbf{K} , denoted as $\mathcal{T}_{\mathbf{K}}$ and, for any random vector $X = (X_1, \dots, X_n) \sim \mathcal{T}_{\mathbf{K}}$, the expectation of $f(X)$ is given by:

$$\mathbb{E}[f(X)] = \int_{\mathbf{K}} f(x)h^*(x)dx = \underline{f}_{\mathbf{K}}^{(r)}. \quad (25)$$

As we now recall one can generate random samples $x \in \mathbf{K}$ from the distribution $\mathcal{T}_{\mathbf{K}}$ using the well known *method of conditional distributions* (see e.g., [21, Section 8.5.1]). Then we will observe that with high probability one of these sample points satisfies (roughly) the inequality $f(x) \leq \underline{f}_{\mathbf{K}}^{(r)}$ (see Theorem 5 for details).

In order to sample a random vector $X = (X_1, \dots, X_n) \sim \mathcal{T}_{\mathbf{K}}$, we assume that, for each $i = 2, \dots, n$, we know the cumulative conditional distribution of X_i given that $X_j = x_j$ for $j = 1, \dots, i-1$, defined in terms of probabilities as

$$F_i(x_i | x_1, \dots, x_{i-1}) := \Pr[X_i \leq x_i \mid X_1 = x_1, \dots, X_{i-1} = x_{i-1}].$$

Additionally, we assume that we know the cumulative marginal distribution function of X_i , defined as:

$$F_i(x_i) := \Pr[X_i \leq x_i].$$

Then one can generate a random sample $x = (x_1, \dots, x_n) \in \mathbf{K}$ from the distribution $\mathcal{T}_{\mathbf{K}}$ by the following algorithm:

- Generate x_1 with cumulative distribution function $F_1(\cdot)$.
- Generate x_2 with cumulative distribution function $F_2(\cdot|x_1)$.
- \vdots
- Generate x_n with cumulative distribution function $F_n(\cdot|x_1, \dots, x_{n-1})$.

Then return $x = (x_1, x_2, \dots, x_n)^T$.

There remains to explain how to generate a (univariate) sample point x with a given cumulative distribution function $F(\cdot)$, since this operation is carried out at each of the n steps of the above algorithm. For this one can use the classical *inverse-transform method* (see e.g., [21, Section 8.2.1]), which reduces to sampling from the uniform distribution on $[0, 1]$ and can be described as follows:

- Generate a sample u from the uniform distribution over $[0, 1]$.
- Return $x = F^{-1}(u)$ (if F is strictly monotone increasing, or $x = \min\{y : F(y) \geq u\}$ otherwise).

As an illustration, we now indicate how to compute the cumulative marginal and conditional distributions $F_i(\cdot)$ and $F_i(\cdot | x_1 \dots x_{i-1})$ for the case of the hypercube $\mathbf{Q}_n = [0, 1]^n$. We will then apply this method to several examples of polynomial minimization over the hypercube \mathbf{Q}_n in the

next section. As before we are given a sum of squares density function $h^*(x)$ on $K = [0, 1]^n$. For $i = 1, \dots, n$, define the function $f_{1\dots i} \in \mathbb{R}[x_1, \dots, x_i]$ by

$$f_{1\dots i}(x_1, \dots, x_i) = \int_0^1 \cdots \int_0^1 h^*(x_1, \dots, x_n) dx_{i+1} \cdots dx_n.$$

Then the cumulative marginal distribution function $F_1(\cdot)$ is given by

$$F_1(x_1) = \int_0^{x_1} f_1(y) dy$$

and, for $i = 2, \dots, n$, the cumulative conditional distribution function $F_i(\cdot \mid x_1 \dots x_{i-1})$ is given by

$$F_i(x_i \mid x_1 \dots x_{i-1}) = \frac{\int_0^{x_i} f_{1\dots i}(x_1, \dots, x_{i-1}, y) dy}{f_{1\dots(i-1)}(x_1, \dots, x_{i-1})}.$$

We now observe that if we generate sufficiently many samples from the distribution $\mathcal{T}_{\mathbf{K}}$ then, with high probability, one of these samples is a point $x \in \mathbf{K}$ satisfying (roughly) $f(x) \leq \underline{f}_{\mathbf{K}}^{(r)}$.

Theorem 5 *Let $X \sim \mathcal{T}_{\mathbf{K}}$. For any $\epsilon > 0$,*

$$\Pr \left[f(X) > \underline{f}_{\mathbf{K}}^{(r)} + \epsilon \left(\underline{f}_{\mathbf{K}}^{(r)} - f_{\min, \mathbf{K}} \right) \right] < \frac{1}{1 + \epsilon}.$$

Proof Let $X \sim \mathcal{T}_{\mathbf{K}}$ so that $\mathbb{E}[f(X)] = \underline{f}_{\mathbf{K}}^{(r)}$. Define the nonnegative random variable

$$Y := f(X) - f_{\min, \mathbf{K}}.$$

Then, one has $\mathbb{E}[Y] = \underline{f}_{\mathbf{K}}^{(r)} - f_{\min, \mathbf{K}}$. Given $\epsilon > 0$, the Markov Inequality (see e.g., [22, Theorem 3.2]) implies

$$\Pr [Y \geq (1 + \epsilon)\mathbb{E}[Y]] \leq \frac{1}{1 + \epsilon}.$$

This completes the proof. □

For given $\epsilon > 0$, if one samples N times independently from $\mathcal{T}_{\mathbf{K}}$, one therefore obtains an $x \in \mathbf{K}$ such that

$$f(x) \leq \underline{f}_{\mathbf{K}}^{(r)} + \epsilon \left(\underline{f}_{\mathbf{K}}^{(r)} - f_{\min, \mathbf{K}} \right)$$

with probability at least $1 - \left(\frac{1}{1 + \epsilon} \right)^N$. For example, if $N \geq 1 + \frac{1}{\epsilon}$ then this probability is at least $1 - 1/e$.

Table 1 Test functions

Name	Formula	Minimum ($f_{\min, \mathbf{K}}$)	Search domain (\mathbf{K})
Booth Function	$f = (x_1 + 2x_2 - 7)^2 + (2x_1 + x_2 - 5)^2$	$f(1, 3) = 0$	$[-10, 10]^2$
Matyas Function	$f = 0.26(x_1^2 + x_2^2) - 0.48x_1x_2$	$f(0, 0) = 0$	$[-10, 10]^2$
Three-Hump Camel Function	$f = 2x_1^2 - 1.05x_1^4 + \frac{1}{6}x_1^6 + x_1x_2 + x_2^2$	$f(0, 0) = 0$	$[-5, 5]^2$
Motzkin Polynomial	$f = x_1^4x_2^2 + x_1^2x_2^4 - 3x_1^2x_2^2 + 1$	$f(\pm 1, \pm 1) = 0$	$[-2, 2]^2$
Styblinski-Tang Function (with $n = 2, 3, 4$)	$f = \sum_{i=1}^n \frac{1}{2}x_i^4 - 8x_i^2 + \frac{5}{2}x_i$	$f(-2.093534, \dots, -2.093534) = -39.16599n$	$[-5, 5]^n$
Rosenbrock Function (with $n = 2, 3, 4$)	$f = \sum_{i=1}^{n-1} 100(x_{i+1} - x_i^2)^2 + (x_i - 1)^2$	$f(1, \dots, 1) = 0$	$[-2.048, 2.048]^n$

Table 2 $f_{\mathbf{K}}^{(r)}$ for Booth, Matyas, Three-Hump Camel and Motzkin Functions

r	Booth Function		Matyas Function		Three-Hump Camel Function		Motzkin Polynomial	
	Value	Time (sec.)	Value	Time (sec.)	Value	Time (sec.)	Value	Time (sec.)
1	244.680	0.30	8.26667	0.26	265.774	0.44	4.2	0.17
2	162.486	0.34	5.32223	0.34	29.0005	0.38	1.06147	0.28
3	118.383	0.41	4.28172	0.27	29.0005	0.31	1.06147	0.08
4	97.6473	0.39	3.89427	0.41	9.58064	0.39	0.829415	0.13
5	69.8174	0.55	3.68942	0.47	9.58064	0.55	0.801069	0.06
6	63.5454	0.59	2.99563	0.69	4.43983	0.55	0.801069	0.13
7	47.0467	0.64	2.54698	0.72	4.43983	0.59	0.708889	0.13
8	41.6727	0.70	2.04307	0.76	2.55032	0.67	0.565553	0.16
9	34.2140	0.83	1.83356	0.81	2.55032	0.70	0.565553	0.16
10	28.7248	0.94	1.47840	0.87	1.71275	0.84	0.507829	0.22
11	25.6050	1.03	1.37644	0.94	1.71275	0.84	0.406076	0.31
12	21.1869	1.48	1.11785	1.25	1.27749	1.11	0.406076	0.27

Table 3 $f_{\mathbf{K}}^{(r)}$ for Styblinski-Tang and Rosenbrock Functions (with $n = 2, 3$)

r	Sty.-Tang ($n = 2$)		Rosenb. ($n = 2$)		Sty.-Tang ($n = 3$)		Rosenb. ($n = 3$)	
	Value	Time (sec.)	Value	Time (sec.)	Value	Time (sec.)	Value	Time (sec.)
1	-12.9249	0.41	214.648	0.34	-18.8832	0.34	629.086	0.37
2	-25.7727	0.31	152.310	0.34	-36.0339	0.38	394.187	0.34
3	-34.4030	0.39	104.889	0.35	-44.9525	0.65	295.811	0.44
4	-41.4436	0.36	75.6010	0.33	-54.4424	0.98	206.903	0.53
5	-45.1032	0.41	51.5037	0.50	-60.5823	0.66	168.135	0.66
6	-51.0509	0.50	41.7878	0.45	-67.6027	0.98	121.558	1.05
7	-56.4050	0.52	30.1392	0.41	-74.5791	1.33	101.953	1.23
8	-58.6004	0.58	25.8329	0.42	-79.1261	2.28	77.4797	1.92
9	-60.7908	0.67	19.4972	0.55	-82.9581	3.53	66.6954	3.08
10	-64.0147	0.83	17.3999	0.61	-87.6127	7.82	53.0369	4.44
11	-65.7111	0.86	13.6289	0.76	-91.0233	10.53	46.5871	7.89
12	-66.5532	1.23	12.5024	0.94	-93.2038	19.47	38.4281	13.99

Table 4 $f_{\mathbf{K}}^{(r)}$ for Styblinski–Tang and Rosenbrock Functions (with $n = 4$)

r	Sty.-Tang ($n = 4$)		Rosenb. ($n = 4$)	
	Value	Time (sec.)	Value	Time (sec.)
1	-24.6541	0.25	1048.19	0.34
2	-45.5192	0.34	690.332	0.42
3	-55.0577	0.61	536.367	0.48
4	-66.8202	0.78	382.729	0.72
5	-74.7215	1.37	314.758	1.39
6	-82.8699	3.09	236.709	3.09
7	-90.8863	9.98	202.674	6.61
8	-97.1192	28.64	156.295	19.62
9	-102.387	83.01	137.015	60.59

4 Numerical examples

In this section, we consider several well-known polynomial test functions from global optimization that are listed in Table 1.

For these functions, we calculate $f_{\mathbf{K}}^{(r)}$ by solving the SDP (3) for increasing r .

We performed the computation on a PC with AMD Phenom(tm) 9600B Quad-Core CPU (2.30 GHz) and with 4 GB RAM. Moreover, we use CVX [12, 13] in MATLAB, selecting SDPT3 [27, 28] as the SDP solver.

We record the values $f_{\mathbf{K}}^{(r)}$ as well as the CPU times (needed to solve the SDP) in Tables 2, 3 and 4.

Furthermore, for each order r , we use the method described in Section 3 to generate samples that are feasible solutions of (2), for the bivariate Rosenbrock and the Three–Hump Camel function in Table 1. For each order, the sample sizes 20 and 1000 are used. We also generate samples uniformly from the feasible set, for comparison. We give the results in Tables 5 and 6, where we record the mean, variance and the minimum value of these samples together with $f_{\mathbf{K}}^{(r)}$ (which equals the sample mean by (25)).

Note that the average of the sample function values approximate $f_{\mathbf{K}}^{(r)}$ reasonably well for sample size 1000, but poorly for sample size 20. Moreover, the average sample function value for uniform sampling from \mathbf{K} is much higher than $f_{\mathbf{K}}^{(r)}$. Also, the minimum function value for sampling from $\mathcal{T}_{\mathbf{K}}$ is significantly lower than the minimum function value obtained by uniform sampling for most values of r . In terms of generating “good” feasible solutions, sampling from $\mathcal{T}_{\mathbf{K}}$ therefore outperforms uniform sampling from \mathbf{K} for these examples, as one would expect.

5 Concluding remarks

We conclude with some additional remarks on Assumption 1, and some discussion on the computation perspectives of the approach studied here for global optimization.

Table 5 Sampling results for the Rosenbrock Function ($n = 2$)

r	$f_{\mathbf{K}}^{(r)}$	Mean	Variance	Minimum	Sample Size
1	214.648	121.125	14005.5	0.00451826	20
		209.9	80699.0	0.0008754	1000
2	152.310	184.496	58423.9	4.94265	20
		149.6	54455.0	0.02805	1000
3	104.889	146.618	64611.2	0.0113339	20
		110.1	26022.0	0.0665	1000
4	75.6010	62.4961	5803.21	0.0542813	20
		75.65	45777.0	0.007285	1000
5	51.5037	58.4032	4397.0	0.668679	20
		50.64	6285.0	0.01382	1000
6	41.7878	35.4183	2936.24	1.16154	20
		37.64	3097.0	0.06188	1000
7	30.1392	29.6545	1022.2	1.05813	20
		27.11	1332.0	0.02044	1000
8	25.8329	19.5392	301.334	0.505628	20
		34.32	4106.0	0.074	1000
9	19.4972	20.8982	328.475	0.564992	20
		18.65	593.6	0.07951	1000
10	17.3999	9.37959	146.496	0.562473	20
		15.33	685.7	0.1448	1000
11	13.6289	8.74923	52.1436	0.75774	20
		15.7	7498.0	0.1719	1000
12	12.5024	5.43151	66.561	0.438172	20
		12.7	764.7	0.0945	1000
Uniform Sample		489.722	433549.0	9.0754	20
		465.729	361150.0	0.0771463	1000

5.1 Revisiting Assumption 1

In this section we consider in more detail our Assumption 1. First we recall another condition, known as the *interior cone condition*, which is classically used in approximation theory (see, e.g., Wendland [29]).

Definition 1 [29, Definition 3.1] A set $\mathbf{K} \subseteq \mathbb{R}^n$ is said to satisfy an interior cone condition if there exist an angle $\theta \in (0, \pi/2)$ and a radius $\rho > 0$ such that, for every $x \in \mathbf{K}$, a unit vector $\xi(x)$ exists such that the set

$$C(x, \xi(x), \theta, \rho) := \{x + \lambda y : y \in \mathbb{R}^n, \|y\| = 1, y^T \xi(x) \geq \cos \theta, \lambda \in [0, \rho]\} \quad (26)$$

is contained in \mathbf{K} .

In fact, one can show that any set satisfying an interior cone condition also satisfies Assumption 1.

Lemma 6 *If a set $\mathbf{K} \subseteq \mathbb{R}^n$ satisfies the interior cone condition (26) then \mathbf{K} also satisfies Assumption 1, where we set*

$$\eta_{\mathbf{K}} = \left[\frac{\sin \theta}{1 + \sin \theta} \right]^n \quad \text{and} \quad \epsilon_{\mathbf{K}} = \rho.$$

Table 6 Sampling results for the Three–Hump Camel Function

r	$f_{\mathbf{K}}^{(r)}$	Mean	Variance	Minimum	Sample Size
1	265.774	216.773	177142.0	0.106854	20
		261.23	193466.0	0.11705	1000
2	29.0005	28.0344	2964.85	1.1718	20
		27.712	6712.8	0.014255	1000
3	29.0005	14.9951	523.904	0.452655	20
		32.363	16681.0	0.0088426	1000
4	9.58064	2.99756	14.1201	0.175016	20
		10.364	1944.0	0.010013	1000
5	9.58064	4.41907	14.1358	0.419394	20
		9.1658	643.88	0.0015924	1000
6	4.43983	7.98481	245.089	0.126147	20
		4.5791	493.12	0.0035581	1000
7	4.43983	3.96711	20.3193	0.260331	20
		3.7911	57.847	0.0076111	1000
8	2.55032	2.18925	3.87943	0.0310113	20
		2.2302	8.3767	0.0028817	1000
9	2.55032	1.38102	2.27433	0.138641	20
		3.2217	812.18	0.00014805	1000
10	1.71275	1.03179	0.992636	0.0645815	20
		1.5069	3.9581	0.0014225	1000
11	1.71275	1.30757	1.90985	0.0320489	20
		1.6379	7.2518	0.0021144	1000
12	1.27749	0.841194	0.914514	0.0369565	20
		1.2105	2.3	0.0005154	1000
Uniform Sample		304.032	163021.0	1.65885	20
		243.216	183724.0	0.00975034	1000

Proof Assume that \mathbf{K} satisfies the interior cone condition (26). Then, using [29, Lemma 3.7], we know that, for every $x \in \mathbf{K}$ and $h \leq \rho/(1 + \sin \theta)$, the closed ball $B_{h \sin \theta}(x + h\xi(x))$ is contained in $C(x, \xi(x), \theta, \rho)$ and thus in \mathbf{K} . Then, for any $x_0 \in \mathbf{K}$ and $\epsilon \in (0, \rho]$, after setting $h = \epsilon/(1 + \sin \theta)$, one can obtain

$$\frac{\text{vol}(B_\epsilon(x_0) \cap \mathbf{K})}{\text{vol}B_\epsilon(x_0)} \geq \frac{\text{vol}C(x_0, \xi(x_0), \theta, \epsilon)}{\text{vol}B_\epsilon(x_0)} \geq \frac{\text{vol}B_{h \sin \theta}(x_0 + h\xi(x_0))}{\text{vol}B_\epsilon(x_0)} = \left[\frac{\sin \theta}{1 + \sin \theta} \right]^n.$$

Thus, Assumption 1 holds after setting $\eta_{\mathbf{K}} = \left[\frac{\sin \theta}{1 + \sin \theta} \right]^n$ and $\epsilon_{\mathbf{K}} = \rho$. \square

For instance, every Euclidean ball with radius $\epsilon > 0$ satisfies an interior cone condition, with radius ϵ and angle $\theta = \pi/3$, see e.g., [29, Lemma 3.10]. Moreover we now show that any full-dimensional polytope satisfies an interior cone condition.

Theorem 6 *Any full-dimensional polytope satisfies an interior cone condition.*

Proof Let $\mathbf{K} \subseteq \mathbb{R}^n$ be a full-dimensional polytope with set of vertices $\{u_1, \dots, u_N\}$. Since \mathbf{K} is full-dimensional then, for any vertex u_i ($i \in [N]$), there exist a unit vector ξ_i , an angle θ_i and a radius ρ_i such that $C(u_i, \xi_i, \theta_i, \rho_i) \subseteq \mathbf{K}$. Set $\theta := \min_{i \in [N]} \theta_i$ and $\rho := \min_{i \in [N]} \rho_i$. Then, for any vertex u_i ($i \in [N]$), one has $C(u_i, \xi_i, \theta, \rho) \subseteq \mathbf{K}$.

We now claim that, for any $x \in \mathbf{K}$, a unit vector $\xi(x)$ exists such that $C(x, \xi(x), \theta, \frac{\rho}{n+1}) \subseteq \mathbf{K}$.

We may assume w.l.o.g. that $x = \sum_{i=1}^{n+1} \alpha_i u_i$ with $\alpha_i \geq 0$ for any $i \in [n+1]$ and $\sum_{i=1}^{n+1} \alpha_i = 1$. One can easily see that there exists $j \in [n+1]$ such that $\alpha_j \geq \frac{1}{n+1}$, and we can assume w.l.o.g. that $j = 1$, that is, $\alpha_1 \geq \frac{1}{n+1}$.

From the fact that $C(u_1, \xi_1, \theta, \rho) \subseteq \mathbf{K}$, we can obtain that, for any unit vector $y \in \mathbb{R}^n$ with $y^T \xi_1 \geq \cos \theta$ and for any $0 \leq r \leq \rho$, then $u_1 + ry \in \mathbf{K}$ holds.

We now consider $x + \lambda y$, where $0 \leq \lambda \leq \frac{\rho}{n+1}$. We have

$$x + \lambda y = \alpha_1 u_1 + \lambda y + \sum_{i=2}^{n+1} \alpha_i u_i = \alpha_1 \left(u_1 + \frac{\lambda}{\alpha_1} y \right) + \sum_{i=2}^{n+1} \alpha_i u_i.$$

As $\alpha_1 \geq \frac{1}{n+1}$ and $0 \leq \lambda \leq \frac{\rho}{n+1}$, we deduce that $0 \leq \frac{\lambda}{\alpha_1} \leq \rho$ and thus $u_1 + \frac{\lambda}{\alpha_1} y \in \mathbf{K}$.

Hence, $x + \lambda y \in \mathbf{K}$ and thus $C(x, \xi_1, \theta, \frac{\rho}{n+1}) \subseteq \mathbf{K}$. □

Combining with Lemma 6, we get the following corollary.

Corollary 1 *Full-dimensional polytopes and Euclidean balls satisfy Assumption 1.*

For example, as the hypercube \mathbf{Q}_n is a full-dimensional polytope, it satisfies an interior cone condition and thus Assumption 1. This can also be seen directly. Set

$$\theta = \arcsin \frac{1}{\sqrt{n}}, \quad \rho = 1/2, \quad \text{and} \quad \xi(x) = -\frac{s(x - \frac{1}{2}e)}{\|x - \frac{1}{2}e\|},$$

where $s(x)$ denotes the sign vector of x for any $x \in \mathbb{R}^n$. Then, one can check that

$$C(x, \xi(x), \theta, \rho) \subseteq \mathbf{Q}_n \quad \text{for any } x \in \mathbf{Q}_n.$$

Furthermore, one can also easily check that \mathbf{Q}_n satisfies Assumption 1 with the constants

$$\eta_{\mathbf{Q}_n} = \frac{1}{2^n} \quad \text{and} \quad \epsilon_{\mathbf{Q}_n} = \frac{1}{2}.$$

5.2 Computational perspectives for global optimization

Recall that the computation of the upper bound $f_{\mathbf{K}}^{(r)}$ by solving the semidefinite programs (3) involve matrix variables of order $\binom{n+2r}{2r}$. Thus one is limited to relatively small values of n and r , when using interior point SDP solvers.

Having said that, the sampling approach of Section 3 often provides good feasible solutions for the examples in Section 4, even for small values of r . One may therefore explore using the sampling technique (for small r) as a way of generating starting points for multi-start global optimization algorithms.

Another possibility to enhance computation would be to investigate more general sufficient conditions for nonnegativity of h on \mathbf{K} , than the sum-of-squares condition studied here. This may result in a faster rate of convergence than for $f_{\mathbf{K}}^{(r)}$.

Acknowledgements

We thank Jean Bernard Lasserre for bringing our attention to his work [18] and for several valuable suggestions, and Dorota Kurowicka for valuable discussions on multivariate sampling techniques.

References

1. Abramowitz, M., Stegun, I.A.: Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables. National Bureau of Standards Applied Mathematics Series 55 (1972)
2. Blumenson, L.E.: A Derivation of n -Dimensional Spherical Coordinates. *The American Mathematical Monthly* 67(1), 63–66 (1960)
3. De Klerk, E., Den Hertog, D., Elabwabi, G.: On the complexity of optimization over the standard simplex. *European journal of operational research*, 191, 773–785 (2008)
4. De Klerk, E., Laurent, M.: Error bounds for some semidefinite programming approaches to polynomial minimization on the hypercube. *SIAM Journal on Optimization*, 20(6), 3104–3120 (2010)
5. De Klerk, E., Laurent, M., Parrilo, P.: A PTAS for the minimization of polynomials of fixed degree over the simplex. *Theory of Computer Science* 361(2-3), 210–225 (2006)
6. De Klerk, E., Laurent, M., Sun, Z.: An alternative proof of a PTAS for fixed-degree polynomial optimization over the simplex. *Mathematical Programming*, DOI: 10.1007/s10107-014-0825-6 (2014)
7. De Klerk, E., Laurent, M., Sun, Z.: An error analysis for polynomial optimization over the simplex based on the multivariate hypergeometric distribution. arXiv:1407.2108 (2014)
8. De Loera, J., Rambau, J., Santos, F.: Triangulations: Structures and algorithms, Book manuscript (2008)
9. Doherty, A.C., Wehner, S.: Convergence of SDP hierarchies for polynomial optimization on the hypersphere. arXiv:1210.5048v2 (2013)
10. Dyer, M.E., Frieze, A.M.: On the Complexity of Computing the Volume of a Polyhedron. *SIAM J. Comput.*, 17(5), 967–974 (1988)
11. Faybusovich, L.: Global optimization of homogeneous polynomials on the simplex and on the sphere. In C. Floudas and P. Pardalos, editors, *Frontiers in Global Optimization*. Kluwer Academic Publishers (2003)
12. Grant, M., Boyd, S.: Graph implementations for nonsmooth convex programs, Recent Advances in Learning and Control (a tribute to M. Vidyasagar), V. Blondel, S. Boyd, and H. Kimura, editors, pages 95–110, *Lecture Notes in Control and Information Sciences*, Springer, http://stanford.edu/~boyd/graph_dcp.html (2008)
13. Grant, M., Boyd, S.: CVX: Matlab software for disciplined convex programming, version 2.0 beta. <http://cvxr.com/cvx> (2013).
14. Grundmann, A., Moeller, H.M.: Invariant integration formulas for the n -simplex by combinatorial methods. *SIAM J. Numer. Anal.* 15, 282–290 (1978)
15. Lasserre, J.B., Zeron, E.S.: Solving a class of multivariate integration problems via Laplace techniques. *Applicaciones Mathematicae*, 28(4), 391–405 (2001)
16. Lasserre, J.B.: Global optimization with polynomials and the problem of moments. *SIAM J. Optim.* 11, 796–817 (2001)
17. Lasserre, J.B.: *Moments, Positive Polynomials and Their Applications*. Imperial College Press (2009)
18. Lasserre, J.B.: A new look at nonnegativity on closed sets and polynomial optimization. *SIAM J. Optim.* 21(3), 864–885 (2011)
19. Lasserre, J.B.: Unit balls of constant volume: which one has optimal representation? Preprint at arXiv: 1408.1324 (2014)
20. Laurent, M.: Sums of squares, moment matrices and optimization over polynomials. In *Emerging Applications of Algebraic Geometry*, Vol. 149 of IMA Volumes in Mathematics and its Applications, M. Putinar and S. Sullivant (eds.), Springer, pages 157–270 (2009)
21. Law, A.M.: *Simulation Modeling and Analysis* (4th edition). Mc Graw-Hill (2007)
22. Motwani, R., Raghavan, P.: *Randomized Algorithms*. Cambridge University Press (1995)
23. Nesterov, Y.: Random walk in a simplex and quadratic optimization over convex polytopes. CORE Discussion Paper 2003/71, CORE-UCL, Louvain-La-Neuve (2003)
24. Nie, J., Schweighofer, M.: On the complexity of Putinar’s Positivstellensatz. *Journal of Complexity*, 23(1), 135–150 (2007)
25. Schweighofer, M.: On the complexity of Schmüdgen’s Positivstellensatz. *Journal of Complexity*, 20, 529–543 (2004)

-
26. Sun, Z.: A refined error analysis for fixed-degree polynomial optimization over the simplex. *Journal of the Operations Research Society of China*, 2(3), 379–393 (2014)
 27. Toh, K.C., Todd, M.J., Tutuncu, R.H.: SDPT3—a Matlab software package for semidefinite programming. *Optimization Methods and Software*, 11, 545–581 (1999)
 28. Tutuncu, R.H., Toh, K.C., Todd, M.J.: Solving semidefinite-quadratic-linear programs using SDPT3. *Mathematical Programming Ser. B*, 95, 189–217 (2003)
 29. Wendland, H.: Scattered Data Approximation. Cambridge University Press (2005)
 30. Whittaker, E.T., Watson, G.W.: A course of modern analysis (4ed). Cambridge University Press, New York (1996)