


Post-Exposure Prophylaxis (PEP) After Sexual Assault to Prevent an HIV Infection



Texas Association Against Sexual Assault
Strengthening Community Responses for Adult Survivors

Start here



Risk Tolerance

Low Risk sexual assault

- Barrier protection/dry sexual contact
- oral
- vaginal
- all the way up to receptive anal sex

High Risk


- forced receptive anal sex with multiple attackers
- attackers who threaten you with their HIV status
- attackers who share needles or have receptive anal sex with other males
- attackers who have sex for money or pay for sex

- ➔ While some sexual assaults might be considered low risk, or you might hear this term used it does not equal no risk. If this is something you are concerned about you have the right to request this medication.
- ➔ Ask your SANE nurse and talk to the Emergency Room Doctor about your options.
- ➔ Every patient should be offered this medication.

Risk vs. Benefit

- ➔ The Emergency Room doctor will discuss risk versus benefits based on your current health status and the medications or drugs that you take daily.
- ➔ This therapy is tolerated well. It is what is given to healthcare providers when they have a contaminated needle stick at work.

Most Common Side Effects:



fatigue
(tired feeling)



nausea
(feeling sick to your stomach)



vomiting or
diarrhea

If you have any of these symptoms your healthcare provider can recommend the best medications to help you tolerate these medications.



Practicalities



28 day therapy of anti-viral medication



Isentress



Isentress and Truvada



Return immediately to the emergency room for vomiting, belly pain, yellow skin or any concerns on how you are tolerating these medications.



You will need blood work done at the time these medications are started and again in about 10 days to see how your body is tolerating them.



Financial Resources

If you cannot afford these medication and do not have insurance the drug is available from the manufacturer on an Emergency Application. You will need the doctors signature on these forms so ask them to print them now and get them signed while you are in the Emergency Room.

➔ Manufacturer-based Programs:

Gilead's "Advancing Access®": 1-800-727-5400

Merck's SUPPORT™: 1-800-850-3430

➔ Non Manufacturer-based Programs:

Patient Advocate Foundation Co-Pay Relief Program: Hotline: 866-512-3861 #1
aidsetc.org | PEpline: 1-888-448-4911



References

Kimberly A. Workowski, MD & Gail A. Bolan, MD. Sexually Transmitted Diseases Treatment Guidelines, 2015. *MMWR Recomm Rep* 2015;64. Pages 2-17, 21-24, 104-109.

A National Protocol for Sexual Assault Medical Forensic Examinations Adults/Adolescents Second Edition. *U.S. Department of Justice. Office on Violence Against Women.* April 2013. NCJ 228119. <https://www.ojp.gov/pdffiles1/ovw/241903.pdf>

Authored by
Rebecca Fears RN, CEN, SANE-A, CA-SANE, CP-SANE
Forensic Nurse Coordinator
UT Health Tyler

Reviewed by
Meghan Richardson DNP, MSN, APRN, FNP-C, CA-SANE, CP-SANE

Robert G. Creath MD, FACEP, FAAEM
Chief of Emergency Services UT Health East Texas at Tyler
Chief of Emergency Services UT Health South Broadway-Tyler
Medical Director Section of Emergency Medicine-UTHT

The information on this brochure is not intended or implied to be a substitute for professional medical advice, diagnosis or treatment. All content, including text, graphics, images and information, contained on or available through this web site is for general information purposes only. Reference in this brochure to any specific commercial product, process, or service is for the information and convenience of the public, and does not constitute endorsement or recommendation by TAASA.