

The Use of Ambient Monitoring to Estimate the Atmospheric Loading of Persistent Toxic Substances to the Great Lakes

**(Phase II of Transportation and Deposition of
Persistent Toxic Substances to the Great Lakes Basin:
Deposition Monitoring Element)**

**Draft
May 8, 1997**

**prepared for the International Joint Commission's
International Air Quality Advisory Board**

by

Dr. Mark Cohen

and

Dr. Paul Cooney

**Center for the Biology of Natural Systems (CBNS)
Queens College, City University of New York
718-670-4180 (phone); 718-670-4189 (fax)**

Preface

This report, on the use of ambient monitoring to estimate the atmospheric loading of persistent toxic substances to the Great Lakes, was commissioned by the IJC International Air Quality Advisory Board, the "IJC Air Board".

It is the last in a series of four closely related reports prepared for the IJC Air Board. The first three reports deal with (1) the capability of specific persistent toxic substances to be subjected to long range atmospheric transport; (2) the status and capabilities of associated emissions inventories; and (3) modeling the atmospheric transport and deposition of persistent toxic substances to the great lakes. A summary of the four components has also been prepared.

These reports were prepared as background documents for the IJC-sponsored Joint International Air Quality Board and Great Lakes Water Quality Board Workshop on Significant Sources, Pathways and Reduction/Elimination of Persistent Toxic Substances, to be held May 21-22, in Romulus Michigan. It is expected that the discussion at the Workshop will serve to elaborate upon and extend the analysis presented in this background report.

Acknowledgments

The assistance of the following people is gratefully acknowledged:

Barry Commoner, of CBNS, for useful discussions, guidance and assistance with the project; and Sharon Peyser, of CBNS, for providing administrative support throughout.

John McDonald and the members of the International Joint Commission International Air Quality Advisory Board for providing assistance, guidance and information.

The following people, who provided information about ambient monitoring of air and/or water in the Great Lakes region:

Mark Allen k	Phil Cook d	Carrie Monosmith i
George Apgar j	Tom Dann c	Dan Orr e
Joel Baker m	Dave De Vault d	Bruce Rodger k
Angela Bandemehr d	Gary Evans d	Frank Schaedlich t
Thomas O. Barnwell, Jr. d	Ray Hoff c	Peter Steer e
Illora Basu v	Alan Hoffman d	Malcolm Still c
Terry Bidleman c	Keri Hornbuckle s	Deborah Swackhamer r
John Bourbon d	Gerald Keeler u	Clyde Sweet h
Ken Brice c	Serge L'Italien c	Stephen Vermette s
C.H. Chan c	Maris Lusia c	Glen Warren d

c. Environment Canada	m. University of Maryland
d. U.S. Environmental Protection Agency	r. University of Minnesota
e. Ontario Ministry of Environment and Energy	s. State University of New York at Buffalo
h. Illinois State Water Survey	t. Tekran, Inc.
i. Michigan Dept. of Envr. Quality	u. University of Michigan
j. Vermont Agency of Natural Resources	v. Indiana University
k. Wisconsin Dept. of Natural Resources	

Table of Contents

Preface	i
Acknowledgments	ii
Table of Contents	iii
List of Figures	v
List of Tables	v
List of Abbreviations	vii
1. INTRODUCTION	1
A. Overall Scope of this Analysis	1
B. Compounds being Considered	3
2. OVERVIEW of the USE of AMBIENT MONITORING of PERSISTENT TOXIC SUBSTANCES TO ESTIMATE NET ATMOSPHERIC LOADINGS to the GREAT LAKES	5
A. Wet Deposition	5
B. Dry Deposition: Particle-Phase Material	5
C. Dry Deposition: Vapor-Phase Material (“Gas Exchange”) . . .	6
D. Droplet Resuspension	8
E. Fog Deposition	8
F. Indirect Atmospheric Loading	9
G. Overall Atmospheric Loading	9
3. MONITORING of AMBIENT AIR and PRECIPITATION for PERSISTENT TOXIC SUBSTANCES in the GREAT LAKES BASIN	13

4.	AMBIENT MONITORING of PERSISTENT TOXIC SUBSTANCES in the WATER of the GREAT LAKES	42
A.	Overview of Monitoring Programs	42
i.	Metal/Organometallics	46
ii.	Organochlorine Biocides	46
iii.	Industrial/Miscellaneous	47
iv.	Chlorobenzenes	48
v.	Polychlorinated Dioxins and Furans	48
vi.	Polychlorinated Biphenyls (PCBs)	49
vii.	Polycyclic Aromatic Hydrocarbons (PAHs) ...	50
viii.	Suspended Sediment Properties	50
B.	Summary of Monitoring of Persistent Toxic Substances in the Water of the Great Lakes	51
5.	SPATIAL REPRESENTATIVENESS of AMBIENT MONITORING DATA for the ESTIMATION of the NET LOADING of PERSISTENT TOXIC SUBSTANCES to the GREAT LAKES	57
A.	Lake Michigan Urban Air Toxics Study	57
B.	Lake Michigan Mass Balance Study	62
C.	Ambient Air Monitoring of Persistent Toxic Substances Conducted by the Michigan Department of Environmental Quality	65
6.	SUMMARY AND CONCLUSIONS	67
7.	BIBLIOGRAPHY and REFERENCES	75

List of Figures

Figure 1.	Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Lake Superior	25
Figure 2.	Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Lake Michigan	29
Figure 3.	Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Huron	33
Figure 4.	Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Lake Erie	37
Figure 5.	Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Lake Ontario	41
Figure 6.	Spatial Variations in Air Concentrations of Biocides and PCB's (Lake Michigan Urban Air Toxics Study)	60
Figure 7.	Spatial Variations in Air Concentrations of PAH's (Lake Michigan Urban Air Toxics Study)	61

List of Tables

Table 1.	Compounds and Compound Groups Targeted in the Binational Virtual Elimination Strategy (BVES) for Persistent Toxic Substances in the Great Lakes Basin	4
Table 2.	Parameters Typically Used to Estimate the Net Atmospheric Deposition to a Given Lake or Lake Area	12
Table 3.	Air and Precipitation Monitoring Programs in the Great Lakes Region Identified in this Analysis	15
Table 4.	Air and Precipitation Monitoring Sites in the Great Lakes Region Identified in this Analysis	16-17
Table 5.	Monitoring of BVES Compounds in Ambient Air and Precipitation in the Great Lakes Region: Canadian Programs	18-19

Table 6.	Monitoring of BVES Compounds in Ambient Air and Precipitation in the Great Lakes Region: U.S. Programs	21-21
Table 7.	Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Superior	22-23
Table 8.	Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Michigan	26-27
Table 9.	Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Huron	30-31
Table 10.	Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Erie	34-35
Table 11.	Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Ontario	38-39
Table 12.	Water Measurement Programs in the Great Lakes: (1986 - 1996)	43
Table 13.	Compounds Measured in Water Monitoring Programs in the Great Lakes	44-45
Table 14.	Summary of Water Monitoring of BVES Persistent Toxic Substances in the Great Lakes (1986 - 1996)	52-56
Table 15.	BVES Compounds for which air measurements were made in the Lake Michigan Urban Air Toxics Study	58
Table 16.	Summary of Air Monitoring Data from a Cruise On Lake Michigan During Summer 1994	62
Table 17.	Summary of Air and Precipitation Monitoring Data at Nine Sites in the Vicinity of Lake Michigan	64
Table 18.	Summary of Air and Water Monitoring Programs for Persistent Toxic Substances in the Great Lakes Region	70-73
Table 19.	Summary of Air and Water Monitoring Issues Identified for BVES Compounds in the Great Lakes Region	74

List of Abbreviations

AEOLOS	..	Atmospheric Exchange Over Lakes and Oceans research project
AES	Atmospheric Environment Service, Environment Canada
ANR	Agency of Natural Resources, Vermont
BVES	Binational Virtual Elimination Strategy
CN	CAMNet (Canadian Mercury Network), various sponsors
COA	Canada Ontario Agreement
DNR	Department of Natural Resources
EC	Environment Canada
EHD	Ecosystem Health Division, Environment Canada
EPA	United States Environmental Protection Agency
HACMP	..	Hazardous Air Contaminants Monitoring Program, Vermont ANR
HgR	Mercury monitoring program, 1995-96 (info. from Gary Evans, EPA)
IADN	Integrated Atmospheric Deposition Network
LCS	Lake Champlain Study
LMMBS	..	Lake Michigan Mass Balance Study
LMUATS	..	Lake Michigan Urban Air Toxics Study
LVL	Level
NAPS	National Air Pollution Surveillance Network, Environment Canada
NWRI	National Water Research Institute
OMOEE	..	Ontario Ministry of Environment and Energy
ox	oxychlordanes
PAH	polycyclic aromatic hydrocarbons
part.	particulate phase of atmosphere
PCDD/F	..	polychlorinated dibenzo- <i>p</i> -dioxins and dibenzofurans
pre	precipitation
precip	precipitation
tn	<i>trans</i> -nonachlor
VOC	volatile organic compounds

1. INTRODUCTION

Atmospheric deposition is an important pathway for the entry of many pollutants to the Great Lakes. This report describes attempts to estimate the net atmospheric loading to the Great Lakes through the use of ambient pollutant measurements. Many examples of these estimates will be discussed in this report. Some of the central examples of these types of estimates include the analyses by Eisenreich *et al.* (1981), Strachan and Eisenreich (1988), Eisenreich and Strachan (1992), and Hoff *et al.* (1996).

The methodology used to estimate atmospheric loading to one or more of the Great Lakes from ambient measurements can be briefly summarized in the following way. First, atmospheric deposition is considered to occur by both wet and dry pathways, i.e., in both the presence and absence of precipitation. Loading from *wet deposition pathway* is estimated from the precipitation rate and the concentration of pollutant in the precipitation. The estimation of loading by the *dry deposition pathway* is somewhat more complicated. In essence, the estimation is based on the concentration of pollutant in the air above the lake, and, as discussed below, for gaseous pollutants (as opposed to pollutant which exists on particles in the atmosphere), the loading estimate also depends on the concentration of pollutant in the water near the surface of the lake. The rate of dry deposition also depends on the meteorological conditions above the lake, and in some cases, the hydrodynamic characteristics of the water near the lake's surface. Reviews of some or all of these phenomena include those by Arimoto (1989), Bidleman and McConnell (1995), Hoff (1994) and Hoff *et al.* (1996), and Slinn *et al.* (1978).

The approach used to estimate loadings from ambient measurements is discussed in Section 2 of this report.

A. Overall Scope of this Analysis

This analysis is primarily limited to U.S. and Canadian government efforts which have monitored one or more of the compounds or groups of compounds listed in Table 1, below, in the Great Lakes region, and/or have used monitoring data to estimate the net atmospheric loading of particular pollutants to the Great Lakes.

In a few cases, particularly relevant non-governmental monitoring/loading efforts and/or efforts for other compounds have been included, but, comprehensive coverage of these other types of analyses was beyond the scope of this analysis.

The analysis began by attempting to assemble the universe of air, precipitation, and water monitoring programs in the Great Lakes region, with basic details about each program (i.e., locations, sampling periods, compounds measured). It turned out that

this effort required a substantial amount of resources. The information presented in this report regarding the universe of monitoring programs relative to the BVES compounds represents the bulk of the work that could be performed in this limited analysis.

Unfortunately, in essentially all cases, it was not possible to obtain and/or analyze monitoring data in the course of this analysis, due to lack of time and/or lack of data availability.

As a general observation, the establishment and maintenance of a clearinghouse for information about and data from monitoring programs in the Great Lakes region would certainly be a useful tool for the public, the research community and the regulatory community.

In the course of this work, it was discovered that there appear to be several parallel efforts underway to assemble the universe of monitoring programs in Canada and the U.S. Details of these efforts were not available for this study, but, it is hoped that they may be available in the future.

Finally, data from two recent research efforts undertaken in the Great Lakes region under the auspices of the U.S. EPA — the AEOLOS project and the Lake Michigan Mass Balance Study — were not generally available for this analysis. These data are being assembled and analyzed by others and will be presented by them elsewhere. It must be noted that these projects would appear to be very relevant to the subject of this report. The public release of the results of these studies is eagerly anticipated.

In addition, two important documents regarding the subject of this report are expected to be released soon but are not currently available. These are:

- Proceedings of the Conference on Atmospheric Deposition to the Great Waters, sponsored by the Air and Waste Management Association, the Great Lakes Center for Environmental Research & Education at Buffalo State College, and the U.S. EPA Great Lakes National Program Office, held October 28-30, 1996, in Niagara Falls, NY.
- Atmospheric Deposition to the Great Lakes and Coastal Waters, edited by Joel Baker, University of Maryland, to be published by the Society of Environmental Toxicology and Chemistry.

Both of these documents are expected to contain very relevant information relative to the subject of this study.

B. Compounds Being Considered

The IJC International Air Quality Advisory Board selected a target list of 27 chemicals or chemical groups to be considered in this analysis, including twelve Level I substances or groups and fifteen Level II substances or groups:

- Level I substances are the 11 Critical Pollutants identified by the IJC's Great Lakes Water Quality Board, plus two additional Critical Pollutant identified by the Lake Superior LaMP and the Lake Ontario Toxics Management Plan (octachlorostyrene and chlordane).
- Level II Substances are those substances identified by the Canada-Ontario Agreement respecting the Great Lakes Basin Ecosystem (COA) as "Tier II" chemicals, plus additional substances of concern identified by LaMP and RAP processes and the Great Lakes Water Quality Guidance in the U.S.

A list of the compounds or groups included is given in Table 1, with the Level indicated in parentheses following the name of the compound.

Table 1. Compounds and Compound Groups Targeted in the Binational Virtual Elimination Strategy (BVES) for Persistent Toxic Substances in the Great Lakes Basin
(Envr. Canada and U.S. EPA, 1996) (Level indicated in parentheses)

METALS / ORGANOMETALLICS

Alkylated Lead (I)

including, but not necessarily limited to:
tetra-, tri- and di-ethyl lead,
tetra-, tri- and di-methyl lead

Cadmium and Cadmium Compounds (II)

including, but not necessarily limited to:
cadmium, cadmium oxide,
cadmium dichloride, cadmium sulfide

Mercury and Mercury Compounds (I)

including, but not necessarily limited to:
elemental mercury, mercury dichloride,
mercury oxide, monomethyl mercury, and
particulate mercury

Tributyltin Compounds (II)

ORGANOCHLORINE BIOCIDES

Aldrin / Dieldrin (I)
Chlordane (I)
DDT / DDD / DDE (I)
Endrin (II)
Heptachlor / Heptachlor Epoxide (II)
Hexachlorocyclohexanes ($\alpha, \beta, \delta,$ and γ) (II)
Methoxychlor (II)
Mirex (I)
Pentachlorophenol (II)
Toxaphene (I)

INDUSTRIAL / MISCELLANEOUS

4-Bromophenyl Phenyl Ether (II)
3,3'-Dichlorobenzidene (II)
Hexachloro-1,3-Butadiene (II)
4,4'-Methylene bis (2-Chloroaniline) (II)
Octachlorostyrene (I)

CHLOROBENZENES

1,4-dichlorobenzene (II)
Tetrachlorobenzenes (several congeners) (II)
Pentachlorobenzene (II)
Hexachlorobenzene (I)

POLYCHLORINATED DIBENZO-P-DIOXINS and DIBENZOFURANS

2,3,7,8-TCDD and 2,3,7,8-TCDF (I)
1,2,3,7,8-PeCDD (I)
1,2,3,4,7,8-HxCDD (I)
1,2,3,6,7,8-HxCDD (I)
1,2,3,7,8,9-HxCDD (I)
1,2,3,4,6,7,8-HpCDD (I)
OCDD (I)
1,2,3,7,8-PeCDF (I)
2,3,4,7,8-PeCDF (I)
1,2,3,4,7,8-HxCDF (I)
1,2,3,6,7,8-HxCDF (I)
1,2,3,7,8,9-HxCDF (I)
2,3,4,6,7,8-HxCDF (I)
1,2,3,4,6,7,8-HpCDF (I)
1,2,3,4,7,8,9-HpCDF (I)
OCDF (I)

POLYCHLORINATED BIPHENYLS (PCB'S)

PCB's (I) [there are 209 PCB congeners]

POLYCYCLIC AROMATIC HYDROCARBONS

Benzo[a]Pyrene (I)
Dinitropyrenes (several congeners) (II)

plus PAH's as a group (II)
including but not limited to:
Phenanthrene, Anthracene
Benz[a]Anthracene, Perylene
Benzo[g,h,i]Perylene

To form a group of PAH's for this analysis,
the following additional PAH's were added,
consisting of the remaining compounds in the
EPA's 16-PAH list & the ATSDR 17-PAH list:

Naphthalene, Acenaphthene
Acenaphthylene, Fluorene, Pyrene
Fluoranthene, Chrysene,
Benzo[b]Fluoranthene, Benzo[j]Fluoranthene
Benzo[k]Fluoranthene, Benzo[e]Pyrene
Dibenz[a,h]Anthracene,
Indeno[1,2,3-c,d]Pyrene

2. OVERVIEW of the USE of AMBIENT MONITORING of PERSISTENT TOXIC SUBSTANCES to ESTIMATE NET ATMOSPHERIC LOADINGS to the GREAT LAKES

As discussed above, briefly, in the introduction, in the use of ambient monitoring data to estimate loadings of pollutants to the Great Lakes through the atmospheric pathway, both wet and dry deposition phenomena are considered.

A. Wet Deposition

To estimate wet deposition flux, pollutant concentrations measured in sampled precipitation are multiplied by precipitation rates, e.g. (with one possible set of a consistent units)¹:

$$(1a) \quad \begin{aligned} & [“L_{\text{wet}}”, \text{ wet deposition loading } (g/\text{year})] = \\ & \quad [“C_p”, \text{ concentration of pollutant in precipitation } (g/m^3)] \\ & \quad \times [“R_p”, \text{ precipitation rate } (m/\text{year})] \\ & \quad \times [“A”, \text{ area of lake } (m^2)]. \end{aligned}$$

Using the symbols defined in the above relation, the loading can be expressed as:

$$(1b) \quad L_{\text{wet}} = C_p R_p A$$

Examples of wet deposition flux estimates to the Great Lakes are included in analyses by Eisenreich *et al.* (1981), Strachan and Eisenreich (1988), Gatz *et al.* (1989), Eisenreich and Strachan (1992), Voldner and Alvo (1993), Chan *et al.* (1994), and Hoff *et al.* (1996).

B. Dry Deposition: Particle-Phase Material

To estimate dry deposition flux from ambient measurements, vapor and particle-associated fractions of a given pollutant are typically considered separately. First, the fraction of a given pollutant that is expected to exist in the vapor phase in the atmosphere, and the fraction that is expected to exist in the particle phase are estimated, either from direct measurement or theoretical considerations.

¹. For the reader's convenience, terms used by Hoff *et al.* (1996) are given the same symbol in this analysis. For terms not used by Hoff *et al.* (1996), a closely related or new symbol is used, as appropriate.

The particle-phase fraction of a given pollutant is generally considered to be deposited at a rate defined by an assumed deposition velocity, and the deposition rate is estimated using ambient measurements, e.g., (with one possible set of consistent units):

$$\begin{aligned}
 (2a) \quad & [L_{\text{dry,part}}, \text{ particle-associated dry deposition loading (g/year)}] = \\
 & [C_a, \text{ total concentration of pollutant in atmosphere (g/m}^3\text{)}] \\
 \times & [\phi_a, \text{ fraction of atmospheric pollutant associated with atmospheric} \\
 & \text{ particles (dimensionless)}] \\
 \times & [v_d, \text{ deposition velocity (m/year) (generally an assumed value)}] \\
 \times & [A, \text{ area of lake, (m}^2\text{)}]
 \end{aligned}$$

Using the symbol defined in the above relation, the loading can be written as:

$$(2b) \quad L_{\text{dry,part}} = C_a \phi_a v_d A$$

Examples of approaches for estimating the dry deposition flux of particle-associated pollutants include those by Hicks and Williams (1980), Slinn and Slinn (1980), Williams (1982) and Schmidt (1982). Discussions of the appropriate value to use for the average deposition velocity for particle phase material are given by Eisenreich *et al.* (1981), Strachan and Eisenreich (1988), Eisenreich and Strachan (1992), and Hoff *et al.* (1996).

As is frequently noted, this approach is regarded as somewhat uncertain, as the exact “average” value of the dry deposition velocity for particulate phase material is not known. Further, as the dry deposition velocity will obviously depend on the particle size distribution in the atmosphere, and the size distribution for the particle-associated material of different pollutants will generally vary from pollutant to pollutant, the use of a single average deposition velocity for all particulate pollutants is obviously somewhat of an oversimplification.

In most of the ambient air monitoring that takes place in the Great Lakes region, particle size distributions are not routinely measured. Measurements of pollutant-specific particle size distributions are even less common.

C. Dry Deposition: Vapor-Phase Material (“Gas Exchange”)

The estimation of the deposition rate (or volatilization rate) of the vapor-phase fraction of a given pollutant to a lake is generally estimated in the following way. The net direction and driving force for the flux of pollutant is considered to result from the degree of departure from thermodynamic equilibrium between the air and water near the lake surface. This equilibrium is generally assumed to be governed by Henry’s Law, a commonly used convention with trace, volatile species in water-air systems. For a given pollutant, the air and water phases are considered to be at equilibrium if the

concentrations satisfy the following condition (again, with one possible set of consistent units):

$$(3a) \quad \begin{aligned} & [“C_{w,dissolved}”, \text{ truly dissolved water phase concentration of pollutant in the} \\ & \quad \text{water below the surface (g/m}^3\text{)}] = \\ & \quad [“C_{a,vapor}”, \text{ vapor phase concentration of pollutant in air above the} \\ & \quad \text{water surface (g/m}^3\text{)}] \\ & \times \quad [“R”, \text{ gas constant (Pa-m}^3 / \text{mol -}^\circ\text{K)}] \\ & \times \quad [“T”, \text{ temperature (}^\circ\text{K)}] \\ & \div \quad [“H”, \text{ Henry’s Law coefficient for the pollutant (Pa-m}^3\text{/mole)}] \end{aligned}$$

Similar to the situation in the atmosphere, a pollutant can exist associated with particles in the water (i.e., suspended sediment) or in a truly dissolved state. It is the truly dissolved pollutant that is relevant to the above thermodynamic equilibrium condition. Using the symbols defined in the above relation, the equilibrium condition above can be defined expressed as:

$$(3b) \quad C_{w,dissolved} = C_{a,vapor} RT / H$$

When the actual pollutant concentrations in the air and water do not satisfy the above equation, the system is said to depart from equilibrium, and the degree of departure, or the *thermodynamic driving force* due to the concentration imbalance, “ Δ_c ”, is expressed by the difference between the two sides of “equation” 3b above:

$$(4) \quad \Delta_c = C_{a,vapor} (RT / H) - C_{w,dissolved}$$

As is typically done in many mass transfer situations, the net rate of flux in a non-equilibrium situation is estimated as the product of the thermodynamic driving force and a mass transfer coefficient,² e.g.,

$$(5) \quad L_{dry,vapor} = \Delta_c K_{OL} A = [C_{a,vapor} (RT/H) - C_{w,dissolved}] K_{OL} A$$

where K_{OL} , the mass transfer coefficient has units of m/yr (to be consistent with the above discussion). K_{OL} depends on the degree of mixing and diffusion on both sides of the surface, i.e., in both the water phase and the air, and is often parameterized as a function of the wind speed at a particular height above the water surface.

It can be seen from equation (5) above that the net flux of dry, vapor “deposition” can be positive (from the air to the water) or negative (from the water to the air),

². An analogous approach is taken in heat transfer situations, where a heat transfer coefficient is used.

depending on the relative concentrations of a given pollutant in the vapor and truly dissolved in the liquid phase.

There have been many studies that have attempted to estimate the direction and rate of gas exchange of different pollutants with one or more of the Great Lakes.

Examples include:

- Hoff *et al.* (1996): a range of compounds measured in the IADN program;
- Hoff *et al.* (1993): toxaphene;
- Achman *et al.* (1993), Jeremiason *et al.* (1994), Hornbuckle *et al.* (1995), and Honrath *et al.* (1997): PCB's;
- Baker and Eisenreich (1990): PAH's and PCB's;
- McConnell *et al.* (1993) and Ridal *et al.* (1996): α -HCH and γ -HCH.

Bidleman and McConnell (1995) have recently reviewed the gas-exchange phenomenon.

D. Droplet Resuspension

While it is somewhat poorly understood at the present, it is also possible for water droplets to be "ejected" from the lake. While some of these droplets would so large that they would quickly fall back into the lake, some are small enough to be carried aloft into the atmosphere above the lake. The water in these droplets will strive to reach thermodynamic equilibrium with the water vapor in the ambient air (as characterized, for example, by the relative humidity) and most or all of the water in the droplets will evaporate relatively quickly. Much of the inorganic and organic material contained in the droplets when they were first formed will remain in the new aerosol particles, even after the water has partially or completely evaporated. These new particles, then, can be seen as a way in which pollutants in the lake can be resuspended into the atmosphere, in a particle-related fashion. A qualitative "term" is added to the loading equation to represent this phenomena.

Most estimates of atmospheric loading do not currently attempt to make quantitative estimates of this phenomenon.

E. Fog Deposition

Another rather poorly characterized deposition phenomenon is that related to fog. The depositional behavior of fog droplets will be different from particle-phase deposition under non-fog conditions. It is unlikely that fog deposition is reliably

measured in conventional precipitation sampling programs. A qualitative “term” is added to the loading equation to represent this phenomena.

Most estimates of atmospheric loading do not currently attempt to make quantitative estimates of this phenomenon.

F. Indirect Atmospheric Loading

In addition to the direct atmospheric loading to a given lake, it is recognized that atmospheric deposition to land areas in a lake’s watershed can contribute, indirectly, to the lake’s pollutant loading. Pollutants deposited in the watershed can be washed by precipitation runoff directly to a lake or to a tributary which empties into a lake. It is believed that these indirect processes are probably less important than the direct deposition phenomena described above. However, much less is known about indirect atmospheric loading, and it is difficult to even make semi-quantitative estimates at this time.

Most estimates of atmospheric loading do not currently attempt to quantitatively include this phenomenon.

G. Overall Atmospheric Loading

The total, direct, net atmospheric loading to a given lake or lake subsection is the sum of the wet and (net) dry deposition amounts estimated above, i.e.:

$$(6a) \quad L_{\text{total,direct}} = L_{\text{wet}} + L_{\text{dry,part}} + L_{\text{dry,vapor}} + [\text{fog deposition}] - [\text{droplet resuspension}]$$

$$(6b) \quad L_{\text{total,direct}} = A \{ C_p [R_p] + C_a [\phi_a v_d + (1-\phi_a)(K_{OL}RT/H)] - C_{w,dissolved} [K_{OL}] \} + [\text{fog deposition}] - [\text{droplet resuspension}]$$

A particle-associated fraction ϕ_w in the water phase can be defined as in the air phase, above, and C_w can be defined as the total water concentration, analogous to the total air concentration C_a . Using these definitions, equation 6b can be rewritten as:

$$(6c) \quad L_{\text{total,direct}} = A \{ C_p [R_p] + C_a [\phi_a v_d + (1-\phi_a)(K_{OL}RT/H)] - C_w (1-\phi_w)[K_{OL}] \} + [\text{fog deposition}] - [\text{droplet resuspension}]$$

Adding a “conceptual” term to represent indirect loadings and slightly rearranging equation 6, above, the following expression is obtained, giving the total atmospheric loading to a lake:

$$\begin{aligned}
(7) \quad L_{\text{total}} = & \quad A R_p C_p && \text{(wet deposition)} \\
& + \quad A \phi_a v_d C_a && \text{(dry deposition of particle-phase material)} \\
& + \quad \text{“fog deposition”} \\
& - \quad \text{“pollutant losses from the lake due to droplet resuspension”} \\
& + \quad A K_{\text{OL}} \{ [(1-\phi_a) (RT/H) C_a] - [(1-\phi_w) C_w] \} && \text{(net deposition of vapor-phase material)} \\
& + \quad \text{“indirect atmospheric loadings”}
\end{aligned}$$

Except for the gas constant (R), and the area of the Lake (A)³, all of the parameters in the above equations will be time- and location-dependent. That is, at any given time, the following parameters will vary from place to place on a given lake, and, at any given location, each will vary over time:

- (a) the concentration of the pollutant in precipitation (C_p), air (C_a), and water (C_w);
- (b) the partitioning behavior of the pollutant in the air and water phases, expressed in the above equations as ϕ_a and ϕ_w , the particle-associated fractions in the air and water phases, respectively.
- (c) meteorological variables, such as precipitation rate, temperature (which appears directly in the equations, and, which influences H), and wind speed (which influences K_{OL} and v_d and droplet resuspension phenomena);

Ambient measurements cannot obviously be made at every location in the air and water near the surface of a given lake. Thus, one issue that arises in applying the above methodology is the extent to which a given set of measurements “captures” enough of the spatial variations to allow an accurate estimate for a given lake or lake portion. For example, if measurements at only one location are made and used to estimate the net atmospheric deposition to a given lake, the question obviously arises

³. Obviously, the area of the lake or lake subsection being considered can change, but, the *magnitude* and *rate* of these changes are relatively small compared to the changes in essentially every other parameter involved.

as to how representative of the “average” the measurements are. The same questions arise even when multiple measurement locations are used.

Moreover, while meteorological measurements at a given site can be made more or less continuously, measurements of chemical concentrations at a given site tend to be made only periodically. Thus, when measurements at only specific times are used, an analogous question arises regarding the extent to which the measurements of any parameter are representative enough to construct accurate time-averages.

Thus, the degree of accuracy of the above methodology will depend in detail on the representativeness of the measurements.

In Table 2, below, the parameters which are typically used to estimate the atmospheric loading to the Great Lakes (or any lake, for that matter) are presented, along with a note about how the parameters are obtained.

This report will discuss various aspects of the loading equations above and the parameters summarized below, including:

- the extent to which ongoing measurement programs provide data for loading estimates;
- strengths and weaknesses (i.e., challenges) in making estimates of atmospheric loadings; and
- estimated atmospheric loadings to the Great Lakes.

**Table 2. Parameters Typically Used to Estimate the
Net Atmospheric Deposition to a Given Lake or Lake Area
(all the parameters below will vary in time and space; thus, averages are used)**

	Parameter	How Obtained (in typical situation)
Wet Deposition of Gas and Particle Phase Pollutant	Concentration of the Pollutant in Precipitation	Measured
	Precipitation Rate	Measured
Dry Deposition of Particle-Phase Pollutant	Concentration of the Pollutant in the Air Near the Lake Surface	Measured
	Vapor/Particle Partitioning Characteristics	Sometimes measured, sometimes estimated. Estimates depend on physical/chemical properties of pollutant, temperature, the nature of the atmospheric aerosol, and the degree to which vapor/particle equilibrium is achieved.
	Dry Deposition Velocity of Particle-Associated Pollutant	Typically estimated; often a constant value is assumed
Net Dry Deposition Flux of Vapor Phase Pollutant	Concentration of the Pollutant in the Air Near the Lake Surface	Measured
	Vapor/Particle Partitioning Characteristics	Sometimes measured, sometimes estimated. Estimates depend on physical/chemical properties of pollutant, temperature, the nature of the atmospheric aerosol, and the degree to which vapor/particle equilibrium is achieved.
	Concentration of Pollutant Truly Dissolved in the Water Near the Lake Surface	Measured, or estimated from the total concentration of the pollutant measured in the water
	Henry's Law Constant	Based on existing laboratory measurements; temperature dependent
	Temperature	Measured
	Air-Water Mass Transfer Coefficient	Estimated, using correlation-based semi-empirical theories derived from experimental measurements. Correlations are often based on the wind speed, measured at a given height above the surface.

3. MONITORING of AMBIENT AIR and PRECIPITATION for PERSISTENT TOXIC SUBSTANCES in the GREAT LAKES BASIN

There are a number of air and/or precipitation monitoring programs for persistent toxic substances in the Great Lakes region.

The most notable is the Integrated Atmospheric Deposition Network (IADN). IADN was established by the 1987 revision to the Great Lakes Water Quality Agreement, and involves coordination among a number of Federal and State/Provincial agencies in the United States and Canada. A summary of this program is given in the IADN Quality Assurance Program Plan (QAPP) (Environment Canada *et al.*, 1994), and by Sweet *et al.* (1995) and Hoff *et al.* (1996). The program will be briefly summarized here.

Air and precipitation samples are collected at three U.S. and two Canadian master stations. There are two satellite stations in the U.S., at which air and precipitation samples are collected. There are twelve satellite stations operated by one or more Canadian federal or provincial agencies (1 of these is in Minnesota; the rest are in Canada). At the four of the Canadian-managed stations operated by OMOEE, air and precipitation samples are collected. At the seven Canadian-managed satellite stations operated by Environment Canada's Ecosystem Health Division (EHD), precipitation samples are collected. At the satellite station at Egbert operated by Environment Canada's Centre for Atmospheric Research Experiments (CARE), air samples are collected. Different compounds are monitored in various portions of the IADN program.

Several other monitoring programs in the Great Lakes region were identified in this analysis. The IADN program and these other programs are summarized in Tables 3-11, below.

Table 3 gives a list of the monitoring programs identified, with the number of sites in different proximity ranges to the Great Lakes. The one-letter code denoting each program is used in Tables 4-11.

Table 4 shows the universe of monitoring sites identified in the programs shown in Table 3. The 2-3 character codes denoting the locations are used in Tables 5-11. Also, the codes describing "site character", e.g., R = Rural, are used in tables 7-11.

In assembling the universe of monitoring sites, all sites identified in any State or Province adjoining a Great Lake were included, with Quebec, Manitoba, and Vermont added. Obviously, all things being equal, sites closer to a given Lake will probably be more representative than a site further away. However, the Great Lakes region is so

large, sites on the order of 100-200 km away from a given Lake cannot obviously be said to be irrelevant.

Tables 5 and 6 give a summary of the coverage of BVES compounds in the monitoring programs operated by Canadian and U.S. agencies, respectively.

Tables 7 through 11 give a summary for each Great Lake of the coverage of BVES compounds in monitoring locations at the monitoring sites in the vicinity of the Lake.

Table 3. Air and Precipitation Monitoring Programs in the Great Lakes Region Identified in this Analysis								
c o d e	Program	Sponsor or Coordinator	Phase(s) Sampled	BVES Compounds Included	# of sites (*)			
					0-10 km	10-100 km	> 100 km	
U.S. IADN Program								
u	U.S. IADN Sites	EPA (IN Univ)	vapor, part, precip	many	5	0	0	
Canadian IADN Program								
c	Air Monitoring at Master Sites	EC AES	vapor, part	many	2	0	0	
o	OMOEE IADN Satellite Sites	OMOEE	vapor, part, precip	many	3	1	0	
e	IADN Satellite site at Egbert	EC AES	vapor, part	many	0	1	0	
g	Great Lakes Precip Network	EC EHD	precip	many	8	2	0	
n	IADN precip at Pt. Petre	EC NWRI	precip	many	1	0	0	
Toxaphene Monitoring in the Great Lakes Region								
b	Research at Pt. Petre	EC	vapor, part	Toxaphene	1	0	0	
h	Research at Eagle Harbor	Indiana Univ.	vapor, part	Toxaphene	1	0	0	
National Air Pollution Surveillance Network, Environment Canada								
t	sites measuring PAH's	EC	vapor + part	PAH's	4	1	4	
t	sites measuring COA substances	EC	vapor + part	OCS, DNP, HCB,PCP	4	1	0	
t	sites measuring VOC's	EC	vapor	VOC's	7	4	11	
t	sites measuring PCDD/F's	EC	vapor + part	PCDD/F	4	1	4	
t	sites measuring metals	EC	particulate	total Cd, Pb, Hg	3	2	5	
Air Toxics Monitoring Program, Ontario Ministry of Environment and Energy								
a	sites measuring PAH's	OMOEE	vapor + part	PAH's	5	2	1	
y	sites measuring VOC's	OMOEE	vapor	VOC's	10	4	1	
p	sites measuring PCDD/F's	OMOEE	vapor + part	PCDD/F	1	1	0	
p	sites measuring PCDD/F's	OMOEE	precipitation (Dorset)	PCDD/F	0	1	0	
Mercury Monitoring Programs								
d	Mercury Deposition Network	various	precipitation	total mercury	1	5	6	
k	Mercury monitoring (CAMNet)	EC	vapor	total mercury	1	1	2	
s	Mercury: Dorset	OMOEE	vapor, precip	total Hg	0	1	0	
q	Univ Mich Air Quality Laboratory	Univ. Michigan	(?) vapor, part, precip	total mercury (?)	0	1	0	
f	Univ Mich Air Quality Laboratory	"trust fund"	(?) vapor, part, precip	total mercury (?)	1	0	0	
l	Lake Champlain study	(?)	(?) vapor, part, precip	total mercury (?)	0	0	1	
r	Mercury research (1995-1996)	EPA	vapor, part	total mercury	5	1	4	
Additional Monitoring Programs in the Great Lakes Region								
m	Air toxics monitoring	Michigan DNR	vapor + particulate	ongoing (?)	1	3	0	
v	Haz. Air Contam. Monitoring	Vermont ANR	vapor + particulate	many	0	0	5	
w	Green Bay Urban Air Toxics	Wisconsin DNR	vapor + particulate	many	1	0	0	
* If information was available, the number of monitoring sites within 0-10 km of a Great Lake, the number of sites between 10-100 km of a Great Lake, and the number of sites greater than 100 km from a Great Lake are listed (in States/Provinces adjoining the Great Lakes, with Manitoba, Quebec, and Vermont added).								

Table 4. Air and Precipitation Monitoring Sites in the Great Lakes Region Identified in this Analysis (a)								
Site Code (b)	Site Name	State or Province	Program(s) at Site (c)	Lat (d)	Long (d)	site type (e)	distance from Lake (km) (f)	orientation to Lake (g)
Sites in the Vicinity of Lake Superior								
EH	Eagle Harbor	MI	u h f r	47.46	88.15	R	0.1	s
BR	Brule River	WI	u d	46.75	91.61	R	0.4	s
Sb	Sibley	ON	g	48.50	88.68	(R)	~ 1	n
WR	Wolf Ridge	MN	o	47.41	91.26	(R)	~ 1	n
TB	Thunder Bay	ON	y	48.40	89.25	U	~ 1	n
SSM	(2 sites) Sault Ste. Marie	ON	a y	46.65	84.35	U	~ 1	e
TL	Turkey Lakes	ON	g	47.03	84.38	(R)	~ 25	e
Fb	Fernberg	MN	d	47.95	91.50	(R)	70	w
TrL	Trout Lake	WI	d	46.05	89.65	(R)	70	s
PR	Popple River	WI	d	45.80	88.40	(R)	100	s
MF	Marcell Expt Forest	MN	d	47.53	93.47	(R)	150	w
CC	Cedar Creek	MN	r	45.40	93.30	(R)	180	sw
CR	Camp Ripley	MN	d	46.25	94.50	(R)	200	sw
Lt	Lamberton	MN	d	44.24	95.30	(R)	380	sw
Wp1	(65 Ellen St.) Winnipeg	MB	t	49.90	97.15	UC	530	nw
Wp2	(301 Weston St.) Winnipeg	MB	t	49.80	97.10	UC	530	nw
Sites in the Vicinity of Lake Michigan								
SBD	Sleeping Bear Dunes	MI	u r	44.76	86.06	R	1	e
Ch	IIT - Chicago	IL	u	41.83	87.62	U	1.5	w
GBy	Green Bay	WI	w	44.50	88.00	U	~ 1	w
SH	South Haven	MI	m	42.40	86.30	(R)	~ 1	e
Pe	Pellston	MI	m	45.55	84.80	(R)	20	e
LG	Lake Geneva	WI	d	42.58	88.50	(R)	60	w
Bo	Bondville	IL	d r	40.05	88.37	(R)	200	sw
WM	Wildcat Mountain	WI	r	43.70	90.57	(R)	230	w
Sites in the Vicinity of Lake Huron								
BI	Burnt Island	ON	c g	45.83	82.95	R	~ 1	island
GB	Grand Bend	ON	o g	43.33	81.75	(R)	~ 1	e
Sa	Sarnia	ON	t	42.98	82.40	UR	~ 1	s
Dv	Deckerville	MI	m	43.50	82.70	(R)	20	w
Eg	Egbert	ON	e k (t?)	44.26	79.79	(R)	40	se
Do	Dorset	ON	o s p y	45.20	78.85	R	90	e
Sites in the Vicinity of Lake Erie								
SP	Sturgeon Point	NY	u r	42.69	79.06	(R)	0.1	s
PS	Port Stanley	ON	o	42.67	81.17	(R)	~ 1	n
PI	Pelee Island	ON	g	41.97	82.52	R	~ 1	island
RP	Rock Point	ON	g	42.85	79.55	(R)	~ 1	w
Sm	Simcoe	ON	t	42.88	80.29	R	~ 1	n
Na	(2 sites) Nanticoke	ON	a y	42.80	80.08	U	~ 1	n
We	Welland	ON	a	42.98	79.25	U	10	n
Lw	Longwoods Conservation Area	ON	t	42.90	81.49	R	30	n
Wd1	(University Ave) Windsor	ON	p a y	42.32	83.04	UC	30	n
Wd2	(College & South St.) Windsor	ON	t	42.29	83.08	U	30	n
Wd3	Windsor	ON	a y	42.30	83.10	U	30	n
SC	St. Clair	ON	g	42.38	82.40	(R)	30	n
Dx	Dexter	MI	q m r	42.35	83.90	(R)	80	w
SF	Salt Fork Lake	OH	r	40.10	81.50	(R)	150	s
HC	Hill Creek St. Park	PA	d	41.83	77.17	(R)	170	se
AP	Allegheny Portage Nat'l Hist. Site	PA	d	40.33	78.50	(R)	240	s

Table 4. Air and Precipitation Monitoring Sites in the Great Lakes Region Identified in this Analysis (a)								
Site Code (b)	Site Name	State or Province	Program(s) at Site (c)	Lat (d)	Long (d)	site type (e)	distance from Lake (km) (f)	orientation to Lake (g)
Sites in the Vicinity of Lake Ontario								
PP	Pt. Petre	ON	c n g b k t r	43.83	77.15	R	~ 0.1	n
Bu	Burlington	ON	g	43.38	79.85	U	~ 1	w
To1	(Evans and Arnold Ave) Toronto	ON	t	43.62	79.53	UI	~ 1	n
To2	(Junction Triangle) Toronto	ON	t	43.67	79.45	UC	~ 1	n
To3	(Downtown) Toronto	ON	t	43.66	79.39	UC	~ 1	n
MZ	(Metro Zoo) Toronto	ON	g	43.87	79.19	(U)	~ 1	n
Ha1	Hamilton	ON	p y	43.26	79.86	U	~ 1	w
Ha2-5	(4 additional sites) Hamilton	ON	y	43.26	79.86	U	~ 1	w
Ha6	Hamilton	ON	t	43.26	79.86	UC	~ 1	w
St	Stouffville	ON	t	43.98	79.27	UR	30	n
Lo	London	ON	y	43.00	81.25	U	40	n
Co	Cornwall	ON	a y	45.05	74.70	U	150	ne
Ot1	(Rideau & Wurtenburg) Ottawa	ON	t	45.43	75.68	UC	150	ne
Ot2	(88 Slater) Ottawa	ON	t	45.42	75.70	UC	150	ne
SA	Saint-Anicet	QU	t k	45.14	74.31	R	200	ne
BV	Burlington	VT	v	44.47	73.21	U	250	ne
Wn	Winooski	VT	v	44.49	73.12	U	250	ne
Rt	Rutland	VT	v	43.61	72.98	U	260	ne
Un	Underhill	VT	v l	44.50	72.90	(R)	275	ne
Mo1	(Pte. aux Trembles) Montréal	QU	t	45.64	73.50	UR	280	ne
Mo2	(1125 Ontario Est) Montréal	QU	t	45.52	73.56	UC	280	ne
Mo3	(Duncan/Decarie) Montréal	QU	t	45.50	73.66	UC	280	ne
Mo4	(1001 Boul Maison. O.) Montréal	QU	t	45.50	73.58	UC	280	ne
Mo5	(Parc Oceanie, Brossard) Montréal	QU	t	45.44	73.47	UR	280	ne
Mo6	(7650 Charteauneuf, Anjou) Montréal	QU	t	45.60	73.56	UR	280	ne
As	l'Assomption	QU	t	45.82	73.43	R	300	ne
MS	Mt. Sutton	QU	t	45.08	72.68	R	300	ne
Bb	Brattleboro	VT	v	42.84	72.56	U	300	ne
Fr	St. Francoise	QU	t	46.02	71.93	R	400	ne
QC	Québec City	QU	t	46.82	71.22	UI	500	ne
Jo	Jonquière	QU	t	48.44	71.20	UI	620	ne
Notes								
a	Sites Identified in the "Great Lakes Region"	Includes all identified sites in any State or Province adjoining one or more Great Lakes, with sites in Manitoba, Quebec, and Vermont added. This table does not include any additional sites used in L. Mich. Mass Balance Study or the AEOLOS project; information about these was not available for this study.						
b	Site Code	These codes are used in other tables, as well						
c	Programs at Site	See Table 3 for program code descriptions. With only a few exceptions, are sites/programs are "active" as of April 1997. Former sites/programs are generally not included.						
d	Latitude and Longitude	In some cases, these are somewhat approximate.						
e	Site Type	Approximate Character of Site (all subject to confirmation and checking): Codes in parentheses are guess-timates and should be checked. R = Rural; U = Urban; UI = Urban Industrial; UC = Urban Commercial; UR = Urban Residential						
f	Distance from Lake (km)	Many of the sites are also in somewhat close proximity to more than one Great Lake. The distance given is the approximate distance to the closest Great Lake. An "~" indicates that precise distance from Lake was not determined in this study, but, that an approximate value is given.						
g	Orientation to Lake	Approximate direction from Lake to Site, i.e., "sw" = site is southwest of stated Lake						

Table 5. Monitoring of BVES Compounds in Ambient Air and Precipitation in the Great Lakes Region: Canadian Programs (a,b)

Site Category -->	Master Sites			Satellite Sites			National Air Pollution Surveillance Network (NAPS) (c)					OMOEE Toxics Monitoring		
program -->	IADN	IADN	IADN	IADN			VOC	PAH	PCDD/F	COA	PART	PAH	VOC	PCDD/F
sponsor / coordinator	EC AES	EC EHD	EC NWRI	EC AES	OMOEE	EC EHD	EC					OMOEE	OMOEE	OMOEE
sites -->	BI, PP	BI, PP	PP	Eg	Do, WR, PS, GB	Sb, TL, GB, SC, PI, RP, Bu	PP, To2-3, St, Eg, Ha6, Sm, Lw, Sa, W d2	PP, To2, Ha6, Sm, W d2	PP, To2, Ha6, Sm, W d2	PP, To2, Ha6, Sm, W d2	PP, To1, To3, W d2	SSM(2), W d1&3, Na(2), We, Co	SSM(2), TB W d1&3, Lo Na(2), Do Ha1-5, Co	Ha1, W d1 [+ precip at Do]
phase(s) -->	vapor part.	precip	precip	vapor part.	vap. part. precip	precip	vapor	combined vapor+part	combined vapor+part	combined vapor+part	part	combined vap+part	vapor	combined vapor+part
Compound or Group	Lvl	Notes: (a) An "x" = compound is measured in given program. (b) see Tables 3, 4 and List of Abbreviations. (c) The NAPS program also includes sites in Quebec and Manitoba												
MERCURY and MERCURY COMPOUNDS														
Elemental Mercury (Hg ⁰)	I													
Divalent Hg (e.g., HgCl ₂)	I													
Monomethyl Mercury	I													
Total Gaseous Mercury	I	k: PP			k	s: Do								
Particulate Mercury	I										x			
Total Mercury	I					s: Do (pre)								
OTHER METALS / ORGANOMETALLICS														
Alkylated Lead	I													
Total Cadmium (Cd)	II	x	x	x	x	x	x				x			
Individual Cd Species	II													
Tributyltin Compounds	II													
ORGANOCHLORINE BIOCIDES														
Aldrin	I	x	x	x	x	x	x							
Dieldrin	I	x	x	x	x	x	x							
Chlordane	I	α,γ,tn,ox	α,γ	α,γ,tn	α,γ,tn,ox	α,γ,tn,ox	α,γ							
DDT / DDD / DDE	I	x	x	x	x	x	x							
Endrin	II	x	x	x	x		x							
Heptachlor	II	x	x	x	x	x	x							
Heptachlor Epoxide	II	x	x	x	x	x	x							
Hexachlorocyclohexanes	II	α,β,δ,γ	α,γ	α,γ	α,β,δ,γ	α,β,γ	α,γ							
Methoxychlor	II	x	x	x	x	x	x							
Mirex	I	x		x	x	x								
Pentachlorophenol	II									x				
Toxaphene	I	b: PP												

INDUSTRIAL / MISCELLANEOUS															
4-Bromophenyl Phenyl Ether	II														
3,3'-Dichlorobenzidene	II														
Hexachloro-1,3-Butadiene	II							x							
4,4'-Methylene bis(2-Chloroaniline)	II														
Octachlorostyrene	I					x						x			
CHLOROENZENES															
1,4-dichlorobenzene	II		x	x			x		x					x	
Tetrachlorobenzenes	II		x	x		x	x								
Pentachlorobenzene	II		x	x		x	x								
Hexachlorobenzene	I	x	x	x		x	x	x				x			
POLYCHLORINATED DIBENZO-P-DIOXINS & DIBENZOFURANS (PCDD/F'S) and POLYCHLORINATED BIPHENYLS (PCB'S)															
PCDD/F's	I						p:Do(pre)					x			x
PCB's	I	x		x		x	x								
POLYCYCLIC AROMATIC HYDROCARBONS															
Acenaphthene	II-a	x	x	x		x		x			x			x	
Acenaphthylene	II-a	x	x	x		x		x			x			x	
Anthracene	II	x				x					x			x	
Benz (a) Anthracene	II	x		x		x	x				x			x	
Benzo (b) Fluoranthene	II-a	x		x		x	x				x			x	
Benzo (j) Fluoranthene	II-a														
Benzo (k) Fluoranthene	II-a	x				x	x				x			x	
Benzo (a) Pyrene	I	x		x		x	x				x			x	
Benzo (e) Pyrene	II-a	x				x					x			x	
Benzo (g,h,i) Perylene	II	x				x	x				x			x	
Chrysene	II-a	x		x		x	x				x			x	
Dibenz(a,h)Anthracene	II-a	x				x	x				x			x	
Dinitropyrenes	II											x			
Fluoranthene	II-a	x	x	x		x	x	x			x			x	
Fluorene	II-a	x	x	x		x		x			x			x	
Indeno(1,2,3-cd)Pyrene	II-a	x				x	x				x			x	
Naphthalene	II-a									x				x	
Phenanthrene	II	x		x		x	x				x			x	
Perylene	II										x			x	
Pyrene	II-a	x	x	x		x	x	x			x			x	

Table 6. Monitoring of BVES Compounds in Ambient Air and Precipitation in the Great Lakes Region: U.S. Programs (a,b,c)

Site Category-->	Master Sites	Satellite Sites	Michigan DNR	Wisconsin DNR	Mercury	Mercury	Vermont
program-->	IADN	IADN	air toxics	air toxics	MDN	HgR	HACMP
sponsor / coordinator	EPA	EPA	Mich DNR	Wisc DNR	various	EPA (?)	Vermont ANR
sites -->	EH, SBD, SP	BR, Ch	SH, Pe, Dx, Dx (ongoing?)	GBy (moved to Wisconsin Rapids in June 1997)	MF, Fb, CR, Lt, AP, HC, PR, TrL, LG	EH, SBD, SP, BI, PP, Bo, WM, SF, Dx, CC (1995-1996 only)	BV, Wn, Rt, Un, Bb (sites 250-300 km from Lk. Ontario)
phase(s) -->	vapor part. precip	vapor part. precip	combined (?) vapor+part	combined vapor+part	precip	vapor part	vapor+part
Compound or Group	Lvl	Notes: (a) "x" = compound is measured in given program (b) see Tables 3, 4 & List of Abbrev. (c) Table does not include info. regarding L. Mich. Mass Balance Study or AEOLOS project					
MERCURY and MERCURY COMPOUNDS							
Elemental Mercury (Hg ⁰)	I						
Divalent Hg (e.g., HgCl ₂)	I						
Monomethyl Mercury	I						
Total Gaseous Mercury	I	r, f, EH		q: Dx		x	Un: LCS
Particulate Mercury	I	r, f, EH				x	Un: LCS
Total Mercury	I		d: BR			x	
OTHER METALS / ORGANOMETALLICS							
Alkylated Lead	I						
Total Cadmium (Cd)	II	x	x	x	x		x?
Individual Cd Species	II						
Tributyltin Compounds	II						
ORGANOCHLORINE BIOCIDES							
Aldrin	I						
Dieldrin	I	x	x	x	x		
Chlordane	I	α,γ,tn	α,γ,tn		x		
DDT / DDD / DDE	I	x	x	x	DDT		
Endrin	II						
Heptachlor	II						
Heptachlor Epoxide	II						
Hexachlorocyclohexanes	II	α,γ	α,γ	α,γ	γ		
Methoxychlor	II						
Mirex	I						
Pentachlorophenol	II						
Toxaphene	I	h: research					

INDUSTRIAL / MISCELLANEOUS							
4-Bromophenyl Phenyl Ether	II						
3,3'-Dichlorobenzidene	II						
Hexachloro-1,3-Butadiene	II				x		x
4,4'-Methylene bis(2-Chloroaniline)	II						
Octachlorostyrene	I						
CHLOROBENZENES							
1,4-dichlorobenzene	II				x		x
Tetrachlorobenzenes	II						
Pentachlorobenzene	II						
Hexachlorobenzene	I	x	x	x			
POLYCHLORINATED DIBENZO-P-DIOXINS & DIBENZOFURANS (PCDD/F'S) and POLYCHLORINATED BIPHENYLS (PCB'S)							
PCDD/F's	I						x
PCB's	I	x	x	x	"total pcb's"		x
POLYCYCLIC AROMATIC HYDROCARBONS							
Acenaphthene	II-a	x	x				x
Acenaphthylene	II-a	x	x	x			x
Anthracene	II	x	x	x			x
Benz (a) Anthracene	II	x	x		x		x
Benzo (b) Fluoranthene	II-a	x	x				x
Benzo (j) Fluoranthene	II-a						
Benzo (k) Fluoranthene	II-a	x	x				x
Benzo (a) Pyrene	I	x	x	x	x		x
Benzo (e) Pyrene	II-a	x	x				
Benzo (g,h,i) Perylene	II	x	x				x
Chrysene	II-a	x	x		x		x
Di benz(a,h)Anthracene	II-a	x	x				x
Dinitropyrenes	II						
Fluoranthene	II-a	x	x	x	x		x
Fluorene	II-a	x	x	x			x
Indeno(1,2,3-cd)Pyrene	II-a	x	x				x
Naphthalene	II-a			x	x		x
Phenanthrene	II	x	x	x	x		x
Perylene	II						
Pyrene	II-a	x	x	x	x		x

Table 7. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Superior

(for descriptions of codes in columns, see tables 3 and 4)

Compound or Compound Group ↓	Level	Air Monitoring					Precipitation Monitoring						
		Eagle Harbor	Brule River	Wolf Ridge	Thunder Bay	Sault Ste. Marie (2 sites)	Additional Sites	Eagle Harbor	Brule River	Wolf Ridge	Stibley	Turkey Lakes	Additional Sites
distance from Lake (km) -->		0.1	0.4	~ 1	~ 1	~ 1	CC = 180 Wp1,2 = 530	0.1	0.4	~ 1	~ 1	~ 25	70 - 380 km
character of site-->		R	R	(R)	U	U		R	R	(R)	(R)	(R)	
MERCURY and MERCURY COMPOUNDS													
Elemental Mercury (Hg ⁰)	I												
Divalent Hg (e.g., HgCl ₂)	I												
Monomethyl Mercury	I												
Total Gaseous Mercury	I	fr					r: CC						
Particulate Mercury	I	fr					r:CC; t:Wp1						
Total Mercury	I							d					d: Fb, TrL, PR, MF, CR, Lt; r: CC
OTHER METALS / ORGANOMETALLICS													
Alkylated Lead	I												
Total Cadmium (Cd)	II	u	u	o			t: Wp1	u	u	o	g	g	
Individual Cd Species	II												
Tributyltin Compounds	II												
ORGANOCHLORINE BIOCIDES													
Aldrin	I			o						o	g	g	
Dieldrin	I	u	u	o				u	u	o	g	g	
Chlordane	I	u	u	o				u	u	o	g	g	
DDT / DDD / DDE	I	u	u	o				u	u	o	g	g	
Endrin	II										g	g	
Heptachlor	II			o						o	g	g	
Heptachlor Epoxide	II			o						o	g	g	
Hexachlorocyclohexanes	II	u	u	o				u	u	o	g	g	
Methoxychlor	II			o						o	g	g	
Mirex	I			o						o			
Pentachlorophenol	II												
Toxaphene	I	h											
INDUSTRIAL / MISCELLANEOUS													
4-Bromophenyl Phenyl Ether	II												
3,3'-Dichlorobenzidine	II												
Hexachloro-1,3-Butadiene	II						t: Wp1						
4,4'-Methylene bis(2-Chloroaniline)	II												
Octachlorostyrene	I			o						o			


Table 7. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Superior

(for descriptions of codes in columns, see tables 3 and 4)

Compound or Compound Group ↓	Level	Air Monitoring					Precipitation Monitoring						
		Eagle Harbor	Brule River	WolfRidge	Thunder Bay	Sault Ste. Marie (2 sites)	Additional Sites	Eagle Harbor	Brule River	WolfRidge	Sibley	Turkey Lakes	Additional Sites
distance from Lake (km) -->		0.1	0.4	~ 1	~ 1	~ 1	CC = 180 Wp1,2 = 530	0.1	0.4	~ 1	~ 1	~ 25	70 - 380 km
character of site-->		R	R	(R)	U	U		R	R	(R)	(R)	(R)	
CHLOROBENZENES													
1,4-dichlorobenzene	II				y	y	t Wp1				g	g	
Tetrachlorobenzenes	II			o						o	g	g	
Pentachlorobenzene	II			o						o	g	g	
Hexachlorobenzene	I	u	u	o				u	u	o	g	g	
POLYCHLORINATED DIBENZO-P-DIOXINS & DIBENZOFURANS (PCDD/F'S) and POLYCHLORINATED BIPHENYLS (PCB'S)													
PCDD/F's	I						t Wp1,Wp2						
PCB's	I	u	u	o				u	u	o			
POLYCYCLIC AROMATIC HYDROCARBONS													
Acenaphthene	II-a	u	u			a	t Wp1,Wp2	u	u		g	g	
Acenaphthylene	II-a	u	u			a	t Wp1,Wp2	u	u		g	g	
Anthracene	II	u	u	o		a	t Wp1,Wp2	u	u	o			
Benz (a) Anthracene	II	u	u	o		a	t Wp1,Wp2	u	u	o			
Benzo (b) Fluoranthene	II-a	u	u	o		a	t Wp1,Wp2	u	u	o			
Benzo (j) Fluoranthene	II-a												
Benzo (k) Fluoranthene	II-a	u	u	o		a	t Wp1,Wp2	u	u	o			
Benzo (a) Pyrene	I	u	u	o		a	t Wp1,Wp2	u	u	o			
Benzo (e) Pyrene	II-a	u	u			a	t Wp1,Wp2	u	u				
Benzo (g,h,i) Perylene	II	u	u	o		a	t Wp1,Wp2	u	u	o			
Chrysene	II-a			o		a	t Wp1,Wp2			o			
Di benz(a,h)Anthracene	II-a	u	u	o		a	t Wp1,Wp2	u	u	o			
Dinitropyrenes	II												
Fluoranthene	II-a	u	u	o		a	t Wp1,Wp2	u	u	o	g	g	
Fluorene	II-a	u	u			a	t Wp1,Wp2	u	u		g	g	
Indeno(1,2,3-cd)Pyrene	II-a	u	u	o		a	t Wp1,Wp2	u	u	o			
Naphthalene	II-a				y	y	t Wp1						
Phenanthrene	II	u	u	o		a	t Wp1,Wp2	u	u	o			
Perylene	II					a	t Wp1,Wp2						
Pyrene	II-a	u	u	o		a	t Wp1,Wp2	u	u	o	g	g	

(this page intentionally left blank)

Figure 1. Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Lake Superior



symbol	program code (see Tbl 3)	description
★	u	IADN Master Site (air and precipitation) The site in this figure, Eagle Harbor, is a U.S. Site; Toxaphene is being monitored at this site in program "h" Mercury is/was monitored at this site in program "p" and "r", respectively
◆	u	U.S. IADN Satellite Site (air and precipitation) Brule River, is also part of the Mercury Deposition Network, program "d"
●	g	IADN Satellite Site, Environment Canada EHD (precipitation)
■	o	OMOEE IADN Satellite Site (air and precipitation)
□	t	Nat'l Air Pollutant Surveillance Network, Envr. Canada (air) both sites in Winnipeg measure PCDD/F's and PAH's one of the sites in Winnipeg measures VOC's and particulate metals
+	y	OMOEE Air Toxics Monitoring, VOC's
*	a,y	OMOEE Air Toxics Monitoring, PAH's and VOC's
△	d	Mercury Deposition Network (precipitation only)
▽	r	EPA Mercury Research Project (1995-96) (air and precip.)

Table 8. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Michigan
(for descriptions of codes in columns, see tables 3 and 4)

Compound or Compound Group	Level	air monitoring					precipitation monitoring			
		Sleeping Bear Dunes	ITT - Chicago	Green Bay (ctb)	South Haven	Pellston	Additional Sites	Sleeping Bear Dunes	ITT - Chicago	Additional Sites
distance from Lake (km) -->		1	1.5	~ 1	~ 1	20				
character of site-->		R	U	U	(R)	(R)				
MERCURY and MERCURY COMPOUNDS										
Elemental Mercury (Hg ⁰)	I									
Divalent Hg (e.g., HgCl ₂)	I									
Monomethyl Mercury	I									
Total Gaseous Mercury	I	r					r: WM,Bo			
Particulate Mercury	I	r					r: WM,Bo			
Total Mercury	I									d: LG, Bo
OTHER METALS / ORGANOMETALLICS										
Alkylated Lead	I									
Total Cadmium (Cd)	II	u	u	w	m	m		u	u	
Individual Cd Species	II									
Tributyltin Compounds	II									
ORGANOCHLORINE BIOCIDES										
Aldrin	I									
Dieldrin	I	u	u	w	m	m		u	u	
Chlordane	I	u	u	w				u	u	
DDT / DDD / DDE	I	u	u	w	m	m		u	u	
Endrin	II									
Heptachlor	II									
Heptachlor Epoxide	II									
Hexachlorocyclohexanes	II	u	u	w	m	m		u	u	
Methoxychlor	II									
Mirex	I									
Pentachlorophenol	II									
Toxaphene	I									
INDUSTRIAL / MISCELLANEOUS										
4-Bromophenyl Phenyl Ether	II									
3,3'-Dichlorobenzidene	II									
Hexachloro-1,3-Butadiene	II			w						
4,4'-Methylene bis(2-Chloroaniline)	II									
Octachlorostyrene	I									

Table 8. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Michigan
(for descriptions of codes in columns, see tables 3 and 4)

Compound or Compound Group	Level	air monitoring					precipitation monitoring			
		Sleeping Bear Dunes	ILT - Chicago	Green Bay (ch)	South Haven	Pellston	Additional Sites	Sleeping Bear Dunes	ILT - Chicago	Additional Sites
CHLOROBENZENES										
1,4-dichlorobenzene	II									
Tetrachlorobenzenes	II			w						
Pentachlorobenzene	II									
Hexachlorobenzene	I	u	u		m	m		u	u	
POLYCHLORINATED DIBENZO-P-DIOXINS & DIBENZOFURANS (PCDD/F'S) and POLYCHLORINATED BIPHENYLS (PCB'S)										
PCDD/F's	I									
PCB's	I	u	u	w	m	m		u	u	
POLYCYCLIC AROMATIC HYDROCARBONS										
Acenaphthene	II-a	u	u					u	u	
Acenaphthylene	II-a	u	u		m	m		u	u	
Anthracene	II	u	u		m	m		u	u	
Benz (a) Anthracene	II	u	u	w				u	u	
Benzo (b) Fluoranthene	II-a	u	u					u	u	
Benzo (j) Fluoranthene	II-a									
Benzo (k) Fluoranthene	II-a	u	u					u	u	
Benzo (a) Pyrene	I	u	u	w	m	m		u	u	
Benzo (e) Pyrene	II-a	u	u					u	u	
Benzo (g,h,i) Perylene	II	u	u					u	u	
Chrysene	II-a			w						
Di benz(a,h)Anthracene	II-a	u	u					u	u	
Di nitro pyrenes	II									
Fluoranthene	II-a	u	u	w	m	m		u	u	
Fluorene	II-a	u	u		m	m		u	u	
Indeno(1,2,3-cd)Pyrene	II-a	u	u					u	u	
Naphthalene	II-a			w	m	m				
Phenanthrene	II	u	u	w	m	m		u	u	
Perylene	II									
Pyrene	II-a	u	u	w	m	m		u	u	

(this page intentionally left blank)

Figure 2. Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Lake Michigan



<u>symbol</u>	<u>program code</u> (see <u>Tbl 3</u>)	<u>description</u>
★	u	IADN Master Site (air and precipitation) The site in this figure, Sleeping Bear Dunes, is a U.S. Site; Mercury was monitored at this site in program "r"
◆	u	U.S. IADN Satellite Site (air and precipitation)
☆	w	Wisconsin Department of Natural Resources
○	m	Michigan Department of Natural Resources (air)
△	d	Mercury Deposition Network (precipitation only) Mercury was monitored at Bondville in program "r"
▽	r	EPA Mercury Research Project (1995-96) (air and precip.)

(Note: this map does not include additional sampling locations utilized in the Lake Michigan Mass Balance Study or the AEOLUS project; accounts of these research projects are given by others elsewhere.)

Table 9. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Huron
(for descriptions of codes in columns, see Tables 3 and 4)


Compound or Compound Group	Level	Air Monitoring						Precipitation Monitoring		
		Burnt Island	Grand Bend	Sarnia	Dorset	Deckerille	Egbert	Burnt Island	Grand Bend	Dorset
distance from Lake (km) -->		~ 1	~ 1	~ 1	90	20	40	~ 1	~ 1	90
character of site-->		R	(R)	UR	(R)	(R)	(R)	R	(R)	(R)
MERCURY and MERCURY COMPOUNDS										
Elemental Mercury (Hg ⁰)	I									
Divalent Hg (e.g., HgCl ₂)	I									
Monomethyl Mercury	I									
Total Gaseous Mercury	I	r k summer 97			s		k			
Particulate Mercury	I	r					t			
Total Mercury	I									s
OTHER METALS / ORGANOMETALLICS										
Alkylated Lead	I									
Total Cadmium (Cd)	II	c	o		o	m	e t	g	g	o
Individual Cd Species	II									
Tributyltin Compounds	II									
ORGANOCHLORINE BIOCIDES										
Aldrin	I	c	o		o		e	g	g	o
Dieldrin	I	c	o		o	m	e	g	g	o
Chlordane	I	c	o		o		e	g	g	o
DDT / DDD / DDE	I	c	o		o	m	e	g	g	o
Endrin	II	c					e	g	g	
Heptachlor	II	c	o		o		e	g	g	o
Heptachlor Epoxide	II	c	o		o		e	g	g	o
Hexachlorocyclohexanes	II	c	o		o	m	e	g	g	o
Methoxychlor	II	c	o		o		e	g	g	o
Mirex	I	c	o		o		e			o
Pentachlorophenol	II									
Toxaphene	I									
INDUSTRIAL / MISCELLANEOUS										
4-Bromophenyl Phenyl Ether	II									
3,3'-Dichlorobenzidene	II									
Hexachloro-1,3-Butadiene	II			t			t			
4,4'-Methylene bis(2-Chloroaniline)	II									
Octachlorostyrene	I		o		o					o

Table 9. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Huron
(for descriptions of codes in columns, see Tables 3 and 4)

Compound or Compound Group	Level	Air Monitoring						Precipitation Monitoring		
		Burnt Island	Grand Bend	Sarnia	Dorset	Deception	Egbert	Burnt Island	Grand Bend	Dorset
distance from Lake (km) -->		~ 1	~ 1	~ 1	90	20	40	~ 1	~ 1	90
character of site -->		R	(R)	UR	(R)	(R)	(R)	R	(R)	(R)
CHLOROBENZENES										
1,4-dichlorobenzene	II			t	y		t	g	g	
Tetrachlorobenzenes	II		o		o			g	g	o
Pentachlorobenzene	II		o		o			g	g	o
Hexachlorobenzene	I	c	o		o	m	e	g	g	o
POLYCHLORINATED DIBENZO-P-D IOXINS & DIBENZOFURANS (PCDD/F'S) and POLYCHLORINATED BIPHENYLS (PCB'S)										
PCDD/F's	I									p
PCB's	I	c	o		o		e			o
POLYCYCLIC AROMATIC HYDROCARBONS										
Acenaphthene	II-a	c					e	g	g	
Acenaphthylene	II-a	c				m	e	g	g	
Anthracene	II	c	o		o	m	e			o
Benz (a) Anthracene	II	c	o		o		e			o
Benzo (b) Fluoranthene	II-a	c	o		o		e			o
Benzo (j) Fluoranthene	II-a									
Benzo (k) Fluoranthene	II-a	c	o		o		e			o
Benzo (a) Pyrene	I	c	o		o	m	e			o
Benzo (e) Pyrene	II-a	c					e			
Benzo (g,h,i) Perylene	II	c	o		o		e			o
Chrysene	II-a	c	o		o		e			o
Di benz(a,h)Anthracene	II-a	c	o		o		e			o
Dinitropyrenes	II									
Fluoranthene	II-a	c	o		o	m	e	g	g	o
Fluorene	II-a	c				m	e	g	g	
Indeno(1,2,3-cd)Pyrene	II-a	c	o		o		e			o
Naphthalene	II-a			t	y	m	t			
Phenanthrene	II	c	o		o	m	e			o
Perylene	II									
Pyrene	II-a	c	o		o	m	e	g	g	o

(this page intentionally left blank)

Figure 3. Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Lake Huron



<u>symbol</u>	<u>program code (see Tbl 3)</u>	<u>description</u>
★	c	IADN Master Site (air and precipitation) The site in this figure, Burnt Island, is a Canadian site; Air monitoring conducted by Atmospheric Environment Service, Envr. Canada Precipitation monitoring by Envr. Canada EHD (program "g") Mercury was monitored at this site in program "r" Monitoring of total gaseous mercury may begin in summer 1997 (program "k")
▲	e	IADN Satellite Site at Egbert, Ontario, Environment Canada (air) total gaseous mercury is measured here also, program "k" this is also reportedly a NAPS site ("t") where VOC's and particulate metals are measured
■	o	OMOEE IADN Satellite Site (air and precipitation) At Dorset, total mercury in precipitation and total gaseous mercury are measured (program "s") Also at Dorset, PCDD/F's in precipitation and VOC's in air are measured by OMOEE (programs "p" and "y") At Grand Bend, precipitation monitoring by EHD is also performed (program "g")
□	t	Nat'l Air Pollutant Surveillance Network, Envr. Canada (air) at the NAPS site at Sarnia, VOC's are measured
○	m	Michigan Department of Natural Resources (air)

Table 10. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Erie

(for descriptions of codes in columns, see Tables 3 and 4)

Compound or C Compound Group	Level	Air Monitoring									Precipitation Monitoring					
		Sturgeon Point	Port Stanley	Simcoe	Nanticoke (2 sites)	Welland	Windsor (*)	Longwoods	Dexter	Additional Sites	Sturgeon Point	Port Stanley	Pelee Island	Rock Point	St. Clair	Additional Sites
		0.1	~ 1	~ 1	~ 1	10	30	30	80		0.1	~ 1	~ 1	~ 1	30	
character of site-->	(R)	(R)	R	U	U	U	R	(R)		(R)	(R)	R	(R)	(R)		
MERCURY and MERCURY COMPOUNDS																
Elemental Mercury (Hg ⁰)	I															
Divalent Hg (e.g., HgCl ₂)	I															
Monomethyl Mercury	I															
Total Gaseous Mercury	I	r						q r		r, SF						
Particulate Mercury	I	r					t	q r		r, SF						
Total Mercury	I									r						q:Dx r:Dx,SF d:HC,AP
OTHER METALS / ORGANOMETALLICS																
Alkylated Lead	I															
Total Cadmium (Cd)	II	u	o				t		m		u	o	g	g	g	
Individual Cd Species	II															
Tributyltin Compounds	II															
ORGANOCHLORINE BIOCIDES																
Aldrin	I		o								u	o	g	g	g	
Dieldrin	I	u	o						m		u	o	g	g	g	
Chlordane	I	u	o								u	o	g	g	g	
DDT / DDD / DDE	I	u	o						m		u	o	g	g	g	
Endrin	II												g	g	g	
Heptachlor	II		o									o	g	g	g	
Heptachlor Epoxide	II		o									o	g	g	g	
Hexachlorocyclohexanes	II	u	o						m		u	o	g	g	g	
Methoxychlor	II		o									o	g	g	g	
Mirex	I		o									o				
Pentachlorophenol	II			t			t									
Toxaphene	I															
INDUSTRIAL / MISCELLANEOUS																
4-Bromophenyl Phenyl Ether	II															
3,3'-Dichlorobenzidene	II															
Hexachloro-1,3-Butadiene	II			t			t	t								
4,4'-Methylene bis(2-Chloroaniline)	II															
Octachlorostyrene	I		o	t			t					o				

Table 10. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Erie


(for descriptions of codes in columns, see Tables 3 and 4)

Compound or Compound Group	Level	Air Monitoring									Precipitation Monitoring					
		Sturgeon Point	Port Stanley	Simcoe	Nanticoke (2 sites)	Welland	Windsor (*)	Longwoods	Dexter	Additional Sites	Sturgeon Point	Port Stanley	Pelée Island	Rock Point	St. Clair	Additional Sites
		0.1	~ 1	~ 1	~ 1	10	30	30	80		0.1	~ 1	~ 1	~ 1	30	
character of site-->	(R)	(R)	R	U	U	U	R	(R)		(R)	(R)	R	(R)	(R)		
CHLOROBENZENES																
1,4-dichlorobenzene	II			t	y		ty	t					g	g	g	
Tetrachlorobenzenes	II		o									o	g	g	g	
Pentachlorobenzene	II		o									o	g	g	g	
Hexachlorobenzene	I	u	o	t			t		m		u	o	g	g	g	
POLYCHLORINATED DIBENZO-P-DIOXINS & DIBENZOFURANS (PCDD/F'S) and POLYCHLORINATED BIPHENYLS (PCB'S)																
PCDD/F's	I			t			tp									
PCB's	I	u	o						m		u	o				
POLYCYCLIC AROMATIC HYDROCARBONS																
Acenaphthene	II-a	u		t	a	a	ta				u		g	g	g	
Acenaphthylene	II-a	u		t	a	a	ta		m		u		g	g	g	
Anthracene	II	u	o	t	a	a	ta		m		u	o				
Benz (a) Anthracene	II	u	o	t	a	a	ta				u	o				
Benzo (b) Fluoranthene	II-a	u	o	t	a	a	ta				u	o				
Benzo (j) Fluoranthene	II-a															
Benzo (k) Fluoranthene	II-a	u	o	t	a	a	ta				u	o				
Benzo (a) Pyrene	I	u	o	t	a	a	ta		m		u	o				
Benzo (e) Pyrene	II-a	u		t	a	a	ta				u					
Benzo (g,h,i) Perylene	II	u	o	t	a	a	ta				u	o				
Chrysene	II-a		o	t	a	a	ta					o				
Dibenz(a,h)Anthracene	II-a	u	o	t	a	a	ta				u	o				
Dinitropyrenes	II															
Fluoranthene	II-a	u	o	t	a	a	ta		m		u	o	g	g	g	
Fluorene	II-a	u		t	a	a	ta		m		u		g	g	g	
Indeno(1,2,3-cd)Pyrene	II-a	u	o	t	a	a	ta				u	o				
Naphthalene	II-a			t	y		ty	t	m							
Phenanthrene	II	u	o	t	a	a	ta		m		u	o				
Perylene	II			t	a	a	ta									
Pyrene	II-a	u	o	t	a	a	ta		m		u	o	g	g	g	

(*) NAPS operates one site in Windsor, measuring VOC's, PAH's, PCDD/F's, selected COA substances, and particulate metals. OMOEE operates two sites in Windsor. At both sites, PAH's and VOC's are measured; at the University Avenue site, PCDD/F's are measured in air.

(this page intentionally left blank)

Figure 4. Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Lake Erie



<u>symbol</u>	<u>program code (see Tbl 3)</u>	<u>description</u>
★	u	IADN Master Site (air and precipitation) The site in this figure, Sturgeon Point, is a U.S. site; Mercury was monitored at this site in program "r" Monitoring of mercury in precipitation may start again soon (program "d")
■	o	OMOEE IADN Satellite Site (air and precipitation) At Dorset, total mercury in precipitation and total gaseous mercury are measured (program "s") At Grand Bend, precipitation monitoring by EHD is also performed (program "g")
□	t	Nat'l Air Pollutant Surveillance Network, Envr. Canada (air) At the Windsor NAPS site, VOC's, PCDD/F's, PAH's, particulate metals, and selected COA substances are measured; OMOEE also monitors PCDD/F's at a site in Windsor (program "p") At the Longwoods NAPS site, VOC's are measured At the Simcoe NAPS site, VOC's, PCDD/F's, PAH's, & selected COA substances are measured;
x	a	OMOEE Air Toxics Monitoring, PAH's (site on this map: Welland, Ontario)
+	y	OMOEE Air Toxics Monitoring, VOC's (site on this map: London, Ontario)
*	a,y	OMOEE Air Toxics Monitoring, PAH's and VOC's (sites on this map: Windsor (2 sites) and Nanticoke (2 sites), Ontario)
○	m	Michigan Department of Natural Resources (air) Mercury is/was being monitored at Dexter in programs "q" and "r", respectively
△	d	Mercury Deposition Network (precipitation only)
▽	r	EPA Mercury Research Project (1995-96) (air and precip.)

Table 11. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Ontario

(for description of codes in columns, see Tables 3 and 4)

Compound or Compound Group	Level	Air Monitoring						Precipitation Monitoring				
		Pt. Peire	Toronto (*)	Hamilton (**)	Stouffville	London	Cornwall	Additional Sites	Pt. Peire	Toronto	Burlington	Additional Sites
distance from Lake (km) -->		0.1	~ 1	~ 1	30	40	150	all ≥ 150 km	0.1	~ 1	~ 1	all ≥ 200 km
character of site-->		R	U	U	UR	U	U		R	U	U	
MERCURY and MERCURY COMPOUNDS												
Elemental Mercury (Hg ⁰)	I											
Divalent Hg (e.g., HgCl ₂)	I											
Monomethyl Mercury	I											
Total Gaseous Mercury	I	kr						k SA; l: Un				
Particulate Mercury	I	tr	t					t: O12, Mo2, Mo6, QC; l: Un				
Total Mercury	I								r			k SA; l: Un
OTHER METALS / ORGANOMETALLICS												
Alkylated Lead	I											
Total Cadmium (Cd)	II	ct	t					t: O12, Mo2, Mo6, QC; v: BV, Wn, Rt, Un, Bb	ng	g	g	
Individual Cd Species	II											
Tributyltin Compounds	II											
ORGANOCHLORINE BIOCIDES												
Aldrin	I	c							ng	g	g	
Dieldrin	I	c							ng	g	g	
Chlordane	I	c							ng	g	g	
DDT / DDD / DDE	I	c							ng	g	g	
Endrin	II	c							ng	g	g	
Heptachlor	II	c							ng	g	g	
Heptachlor Epoxide	II	c							ng	g	g	
Hexachlorocyclohexanes	II	c							ng	g	g	
Methoxychlor	II	c							ng	g	g	
Mirex	I	c							n			
Pentachlorophenol	II	t	t	t								
Toxaphene	I	b										
INDUSTRIAL / MISCELLANEOUS												
4-Bromophenyl Phenyl Ether	II											
3,3'-Dichlorobenzidene	II											
Hexachloro-1,3-Butadiene	II	t	t	t	t			t: O1,2,SA, Mo1,2,4,5,As,MS,Fr; v: BV, Wn, Rt, Un, Bb				
4,4'-Methylene bis(2-Chloroaniline)	II											
Octachlorostyrene	I	t	t	t								

Table 11. Summary of Ambient Air and Precipitation Monitoring of BVES Substances in the Vicinity of Lake Ontario

(for description of codes in columns, see Tables 3 and 4)


Compound or Compound Group	Level	Air Monitoring							Precipitation Monitoring			
		Pt. Peire	Toronto (*)	Hamilton (**)	Stouffville	London	Cornwall	Additional Sites	Pt. Peire	Toronto	Burlington	Additional Sites
distance from Lake (km) -->		0.1	~ 1	~ 1	30	40	150	all ≥ 150 km	0.1	~ 1	~ 1	all ≥ 200 km
character of site-->		R	U	U	UR	U	U		R	U	U	
CHLOROBENZENES												
1,4-dichlorobenzene	II	t	t	ty	t	y	y	t:Ot1,2,SA, Mo1,2,4,5,As,MS,Fr; v: BV, Wn, Rt, Un, Bb	ng	g	g	
Tetrachlorobenzenes	II								ng	g	g	
Pentachlorobenzene	II								ng	g	g	
Hexachlorobenzene	I	ct	t	t					ng	g	g	
POLYCHLORINATED DIBENZO-P-DIOXINS & DIBENZOFURANS (PCDD/F'S) and POLYCHLORINATED BIPHENYLS (PCB'S)												
PCDD/F's	I	t	t	tp				t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb				
PCB's	I	c						v: BV, Wn, Rt, Un, Bb	n			
POLYCYCLIC AROMATIC HYDROCARBONS												
Acenaphthene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	ng	g	g	
Acenaphthylene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	ng	g	g	
Anthracene	II	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb				
Benz (a) Anthracene	II	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	n			
Benzo (b) Fluoranthene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	n			
Benzo (j) Fluoranthene	II-a											
Benzo (k) Fluoranthene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb				
Benzo (a) Pyrene	I	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	n			
Benzo (e) Pyrene	II-a	ct	t	t			a	t: Mo2,Jo				
Benzo (g,h,i) Perylene	II	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb				
Chrysene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	n			
Di benz(a,h)Anthracene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb				
Di nitropyrenes	II											
Fluoranthene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	ng	g	g	
Fluorene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	ng	g	g	
Indeno(1,2,3-cd)Pyrene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb				
Naphthalene	II-a	t	t	ty	t	y	y	t:Ot1,2,SA, Mo1,2,4,5,As,MS,Fr; v: BV, Wn, Rt, Un, Bb				
Phenanthrene	II	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	n			
Perylene	II	t	t	t			a	t: Mo2,Jo				
Pyrene	II-a	ct	t	t			a	t: Mo2,Jo; v: BV, Wn, Rt, Un, Bb	ng	g	g	

(*) Three NAPS sites in Toronto; not all compounds are measured at all sites (VOC's at all 3; PAH, PCDD/F and COA at 1 site only; particulate metals at 2 sites)

(**) One NAPS site in Hamilton; Five OMOEE air toxics monitoring sites in Hamilton: VOC's measured at all five, PCDD/F in air measured at one site (downtown)

(this page intentionally left blank)

Figure 5. Air and Precipitation Sampling Sites for One or More BVES Compounds in the Vicinity of Lake Ontario



symbol	program code (see Tbl 3)	description
★	c	IADN Master Site (air and precipitation) The site in this figure, Point Petre, is a Canadian site; Air monitoring is conducted by Atmospheric Environment Service, Envr. Canada ("c") Precipitation monitoring by National Water Research Institute, Envr. Canada ("n") Precipitation monitoring also by Envr. Canada EHD (program "g") Total gaseous mercury monitored by Environment Canada (program "k") Toxaphene monitored by Environment Canada (program "b") Mercury was monitored at this site in program "r"
●	g	IADN Satellite Site, Environment Canada EHD (precipitation)
□	t	Nat'l Air Pollutant Surveillance Network, Envr. Canada (air) Toronto (3 sites): VOC's (3 sites); particulate metals (2 sites); PAH's, PCDD/F's and selected COA substances (1 site); Stouffville, St. Anicet, l'Assomption, St. Francoise, Mt. Sutton: VOC's only; Quebec City: particulate metals are measured; Ottawa (2 sites): VOC's measured at both, particulate metals at 1 site Simcoe & Hamilton: VOC's, PCDD/F's, PAH's, & selected COA substances measured; At one site in Hamilton, OMOEE also measures PCDD/F's (program "p") Jonquiere: PAH's and PCDD/F's measured Montreal (six sites): VOC's (4 sites); PAH's and PCDD/F's (1 site); particulate metals (2 sites)
+	y	OMOEE Air Toxics Monitoring, VOC's (sites on this map: Hamilton, Ontario (5 sites))
*	a,y	OMOEE Air Toxics Monitoring, PAH's and VOC's (site on this map: Cornwall, Ontario)
◇	v	Vermont Agency of Natural Resources (air)

4. AMBIENT MONITORING of PERSISTENT TOXIC SUBSTANCES in the WATER of the GREAT LAKES

A. Overview of Monitoring Programs

There have been many measurements of toxic compounds in the water of the Great Lakes. These measurements are typically made on cruises of research vessels in which samples of lake water are collected at different locations (“stations”) in a given Lake.

Table 12 gives a list of the different research cruises that carried out water measurements for each of the Great Lakes during the period 1986-1996. In some cases, information on 1997 cruises was included. For each cruise, the sponsoring agency is listed followed by the principal investigator, where known, in parentheses. Then, the seasons in which the cruises took place (Sp = Spring, Su = Summer, etc.) are listed, with the number of sampling stations in parentheses. Lastly, there is a reference to a “suite number”, which is an abbreviation for the suite of chemical compounds measured in the particulate program. Each suite number corresponds to a column in Table 13 where the compounds measured for that particular cruise are indicated (compounds that were measured are marked with an “x”).

The rows in Table 13 list the BVES compounds as well as several others that have been measured in the water of the Great Lakes. Each of the columns correspond to the specific research projects or cruises that have carried out water measurements in one or more of the Great Lakes between 1986 and 1996. The table is organized by toxic compound groupings: Metals/Organometallics, Organochlorine Biocides, Industrial/Miscellaneous, Chlorobenzenes, Dioxins, PCB’s, and PAH’s.

In the following, a narrative summary of monitoring efforts will be presented, considering the research done in accordance with the above chemical groupings.

While a substantial effort was made to assemble a comprehensive list of cruises and compounds measured, it is acknowledged that it is likely that one or more efforts may have been missed. It is hoped that in the further review and discussion of this document, any errors of omission or content can be corrected.

As discussed in Section 2, the phase of a given pollutant in the water column may be important for estimating the net direction and rate of gas exchange. When it could be obtained, information regarding the phase of the measurements made — dissolved or associated with suspended sediment — will be presented.

Table 12. Water Measurement Programs of One or More BVES Compounds in the Great Lakes: 1986-1997

Year	Lake Superior	Lake Huron	Lake Michigan	Lake Erie	Lake Ontario
1997	<ul style="list-style-type: none"> EC(L'Italien): Sp & Su, (14 stations), Suite 4; EPA: Sp (6 stations) Suite 6; DS: Sp(?) (5-9 sta.) Suite 11 	<ul style="list-style-type: none"> EPA: Sp (6 stations) Suite 6 	<ul style="list-style-type: none"> EPA: Sp (6 stations) Suite 6; DS: Sp? (5-9 sta.) Suite 11 	<ul style="list-style-type: none"> EPA: Sp (6 stations), Suite 6 	<ul style="list-style-type: none"> EPA: Sp (6 stations) Suite 6
1996	<ul style="list-style-type: none"> EC(L'Italien): Sp & Su, (14 stations), Suite 4; EPA: Sp (6 stations) Suite 6; DS: Sp(?) (5-9 sta.) Suite 11 	<ul style="list-style-type: none"> EPA: Sp (6 stations) Suite 6 	<ul style="list-style-type: none"> EPA: Sp (6 stations) Suite 6 	<ul style="list-style-type: none"> EPA: Sp (6 stations), Suite 6 	<ul style="list-style-type: none"> EPA: Sp (6 stations) Suite 6
1995			<ul style="list-style-type: none"> EPA, LMMBS: 8 cruises (41 stations), Suite 5; DS: Sp? (5-9 sta.) Suite 11 	<ul style="list-style-type: none"> EC(L'Italien): Sp,Su,Fa (9 stations), Suite 3 	
1994			<ul style="list-style-type: none"> EPA, LMMBS: 8 cruises (41 stations), Suite 5; DS: Sp? (5-9 sta.) Suite 11; Cook & Burkhard: Fall (3 stations) Suite 12 	<ul style="list-style-type: none"> EC(L'Italien): Sp,Su,Fa (9 stations), Suite 3 	<ul style="list-style-type: none"> US EPA Region 2 (Yeh): Sept. (30 stations) Suite 9
1993	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6; DS: Sp? (5-9 sta.) Suite 11 	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6 	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6; DS: Sp? (5-9 sta.) Suite 11 Cleckner et al. (1995): Fall (1 station); mercury 	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6 	<ul style="list-style-type: none"> EC(L'Italien): Sp,Su,Fa (6 stations) Suite 2; EPA: Sp Su (6+ stations) Suite 6
1992	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6; EPA? NOAA? (Eisenreich et al.): Su (5 stations) Suite 7 	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6 	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6 	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6 	<ul style="list-style-type: none"> EC(L'Italien): Sp,Su,Fa, Suite 2, 6 stations; EPA: SP SU (6+stations) Suite 6
1991	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6; DS: Sp? (5-9 sta.) Suite 11 	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6; DS: Sp? (5-9 sta.) Suite 11 	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6; DS: Sp? (5-9 sta.) Suite 11 	<ul style="list-style-type: none"> EPA: Sp Su (6+ stations) Suite 6; DS: Sp? (5-9 sta.) Suite 11 	<ul style="list-style-type: none"> EPA: Sp Su (6+ sta.) Suite 6; DS: Sp? (5-9 sta.) Suite 11; EC: Su (2 sta.) metals (including cadmium)
1990	<ul style="list-style-type: none"> EPA? NOAA? (Eisenreich et al.): Su (6 stations) Suite 7 	<ul style="list-style-type: none"> EPA (McConnell,Bidleman): Su (10 stations) Suite 8 	<ul style="list-style-type: none"> EPA (McConnell,Bidleman): Su (11 stations) Suite 8 	<ul style="list-style-type: none"> EPA (McConnell,Bidleman): Su (18 stations) Suite 8 	<ul style="list-style-type: none"> EC(L'Italien) annual spring cruise, (47 stations) Suite 1; EPA (McConnell,Bidleman): Su (7 stations) Suite 8
1989			<ul style="list-style-type: none"> EPA (McConnell,Bidleman): Sp (3 stations -Green Bay) Suite 8 		
1988	<ul style="list-style-type: none"> EPA? NOAA? (Eisenreich et al.): Su (5 stations) Suite 7 	<ul style="list-style-type: none"> EC(L'Italien): Sp, (63 stations for Hur+Ont) Suite 1 			<ul style="list-style-type: none"> EC(L'Italien): Sp, (63 stations for Hur+Ont) Suite 1
1987	<ul style="list-style-type: none"> EC(L'Italien): Sp, (46 stations for Sup+Hur) Suite 1 	<ul style="list-style-type: none"> EC(L'Italien): Sp, (46 stations for Sup+Hur) Suite 1 		<ul style="list-style-type: none"> Gill and Bruland (1990) Su, (1 nearshore station only?) mercury 	<ul style="list-style-type: none"> Gill and Bruland (1990) Su, (1 nearshore station only?) mercury
1986	<ul style="list-style-type: none"> EC(L'Italien): Sp, (96 total stations for 4 lakes) Suite 1; EC IWD (Stevens&Neilson) Sp (19 stations) Suite 10 NOAA/EPA? (Eisenreich) Su (6 stations) Suite 7; 	<ul style="list-style-type: none"> EC(L'Italien): Sp, (96 total stations for 4 lakes) Suite 1; EC IWD (Stevens&Neilson): Sp (18 stations) Suite 10 		<ul style="list-style-type: none"> EC(L'Italien): Sp (96 total stations for 4 lakes) Suite 1; EC IWD (Stevens&Neilson) Sp (21 stations) Suite 10 	<ul style="list-style-type: none"> EC(L'Italien): Sp (96 total stations for 4 lakes) Suite 1; EC IWD (Stevens&Neilson): Sp (33 stations) Suite 10

Note: Suite 1 indicates that the compounds measured in the cruise were those marked with an x in the Suite 1 column of Table 13; similarly for Suites 2-11.

Abbrev.: DS = Deborah Swackhamer; EC = Envr. Canada; EPA = Envr. Protection Agency; LMMBS = Lk Mich MassBalance Study; IWD = EC Inland Waters Directorate.

Table 13: Compounds Measured in Water Monitoring Programs in the Great Lakes

			Suite 1	Suite 2	Suite 3	Suite 4	Suite 5	Suite 6	Suite 7	Suite 8	Suite 9	Suite 10	Suite 11	Suite 12
			EC(L'Italien) 86-90 (Sup., Hur., Erie, Ont.)	EC(L'Italien) 92-93 Ontario	EC(L'Italien) 94-95 Erie	EC(L'Italien) 96-97 Superior	Lake Mich. Mass Bal. Study (94-95)	EPA Study: 91-97 Great Lakes	Eisenreich et al. 86,88,90,92 Superior	EPA (Bidleman) GL Survey 1989, 1990	US EPA Region II Ontario, 94	EC IWD Stevens Great Lakes 86	Swackhamer Great Lakes 91-97	Cook & Burkhard Michigan 94
chemical name	cas # (1)	(I,II)												
Metals / Organometallics														
cadmium and cadmium compounds		II												
alkylated lead compounds		I												
mercury and mercury compounds		I							x					
tributyl tin	000688-75-3	II												
Organochlorine Biocides														
aldrin	000309-00-2	I	x	x	x	x					x	x		x
atrazine	001912-24-9						x	x(93)			x			x
DEA (atrazine metabolite)							x							
DIA (atrazine metabolite)							x							
Alpha-Chlordane	005103-71-9	I	x	x	x	x					x			x
Gamma-Chlordane	005566-34-7	I	x	x	x	x					x			x
cis-Chlordane		I												x
trans-Chlordane	005103-74-2	I												x
trans-nonachlor	039765-80-5	I					x				x			x
cis-nonachlor	005103-73-1	I									x			x
p,p'-DDT	000050-29-3	I	x	x	x	x		x(92,93)?				x		x
p,p'-DDD(p,p'-TDE)	000072-54-8	I	x	x	x	x		x(92,93)?			x	x		x
p,p'-DDE	000072-55-9	I	x	x	x	x		x(92,93)?			x	x		x
o,p'-DDT	000789-02-6	I	x	x	x	x		x(92,93)?				x		x
dieldrin	000060-57-1	I	x	x	x	x					x	x		x
endrin	000072-20-8	II	x	x	x	x					x	x		x
endrin aldehyde	007421-93-4	II ?		x	x	x								x
Apha-Endosulphan	000959-98-8		x	x	x	x						x		x
Beta-Endosulphan	033213-65-9		x	x	x	x						x		x
heptachlor	000076-44-8	II	x	x	x	x					x	x		x
heptachlor epoxide	001024-57-3	II	x	x	x	x					x	x		x
alpha-hexachlorocyclohexane	000319-84-6	II	x	x	x	x				x	x	x		x
beta-hexachlorocyclohexane	000319-85-7	II												x
delta-hexachlorocyclohexane	000319-86-8	II												x
gamma-hexachlorocyclohexane	000058-89-9	II	x	x	x	x				x	x	x		x
mixed hexachlorocyclohexanes	000319-84-6	II												
methoxychlor	000072-43-5	II	x	x	x	x						x?		x
mirex	002385-85-5	I	x	x	x	x					x	x		x
photomirex		I ?		x	x	x								x
pentachlorophenol	000087-86-5	II												
toxaphene	008001-35-2	I									x		x	x
Industrial/Miscellaneous														
4-bromophenyl phenyl ether	000101-55-3	II												x ?
3,3'-dichlorobenzidene	000091-94-1	II												
hexachloro-1,3-butadiene	000087-68-3	II		x	x	x								x
4,4'-methylene bis(2-chloroaniline)	000101-14-4	II												
octachlorostyrene	029082-74-4	I		x	x	x					x			x
hexachlorocyclopentadiene	000077-47-4			x	x	x								x
phthalates					x	x								
Chlorobenzenes														
1,2-dichlorobenzene	000095-50-1		x	x	x	x						x		
1,3-dichlorobenzene	000541-73-1		x	x	x	x						x		
1,4-dichlorobenzene	000106-46-7	II	x	x	x	x						x		

1,2,3-trichlorobenzene	000087-61-6			x	x	x	x					x	x	x	
1,2,4-trichlorobenzene	000120-82-1			x	x	x	x					x	x	x	
1,3,5-trichlorobenzene	000108-70-3			x	x	x	x					x	x	x	
1,2,3,4-tetrachlorobenzene	000634-66-2			x	x	x	x					x	x	x	
1,2,3,5-tetrachlorobenzene	000634-90-2											x		x	
1,2,4,5-tetrachlorobenzene	000095-94-3											x		x	
pentachlorobenzene	000608-93-5			x	x	x	x					x	x	x	
hexachlorobenzene	000118-74-1			x	x	x	x					x	x	x	
Polychlorinated Dibenzo-P-Dioxins & Dibenzofurans															
2,3,7,8-TCDD	001746-01-6					x	x								x
other PCDD/F congeners															x
Polychlorinated Biphenyls (PCBs)															
All 209 PCB congeners								x							x (all 209?)
A subset of 135 PCB congeners												x			
A subset of 82 PCB congeners									x(88,90,92)						
A subset of 65 PCB congeners								x							
A subset of 35 PCB congeners									x(86)						
Aroclor 1242	053469-21-9				x+								x?		
Aroclor 1254	011097-69-1				x+								x?		
Aroclor 1260	011096-82-5				x+								x?		
Total PCBs *				x	x+?	x+	x	x?	x?	x?			x		
Polycyclic Aromatic Hydrocarbons															
dinitropyrenes (mixed)	078432-19-6														
acenaphthene	000083-32-9			II - A											x
acenaphthylene	000208-96-8			II - A	x(88,90)	x	x	x				x			x
anthracene	000120-12-7			II								x			x
benz [a] anthracene	000056-55-3			II			x	x			x(86)	x			x
dibenz [a,h] anthracene	000053-70-3			II - A							x(86)	x			x
chrysene	000218-01-9			II - A			x	x			x(86)	x			x
fluoranthene	000206-44-0			II - A	x(87,88,90)	x	x	x			x(86)	x			x
benzo [b] fluoranthene	000205-99-2			II - A							x(86)	x			x
benzo [j] fluoranthene	000205-82-3			II - A											
benzo [k] fluoranthene	000207-08-9			II - A							x(86)	x			x
benzo [b and k] fluoranthene				II - A	x(88,90)	x	x	x							
fluorene	000086-73-7			II - A	x(88,90)	x	x	x			x(86)	x			x
indene	000095-13-6			II	x(88,90)	x									
naphthalene	000091-20-3			II - A				x	x			x			x
2-chloronaphthalene	000091-58-7			II	x(88,90)	x	x	x							x
1- methyl naphthalene	000090-12-0			II	x(88,90)	x	x	x							
2- methyl naphthalene	000091-57-6			II	x(88,90)	x	x	x							
1,2,3,4-Tetrahydronaphthalene				II	x(88,90)	x									
perylene	000198-55-0			II											
benzo [g,h,i] perylene	000191-24-2			II	x(88,90)	x	x	x			x(86)	x			x
dibenzo(ah)perylene				II											
phenanthrene	000085-01-8			II	x(88,90)	x	x	x			x(86)	x			x
pyrene	000129-00-0			II - A	x(88,90)	x	x	x			x(86)	x			x
benzo [a] pyrene	000050-32-8			I	x(88,90)	x	x	x			x(86)	x			x
benzo [e] pyrene	000192-97-2			II - A							x(86)	x			
indeno [1,2,3-c,d] pyrene	000193-39-5			II - A	x(88,90)	x	x	x			x(86)	x			x
Suspended Sediment Properties															
Total Suspended Solids					x?	x	x	x	x	x?	x	x	x		x
Dissolved Organic Carbon (DOC)						x	x	x	x	x?	x	x	x		x
Particulate Organic Carbon (POC)						x	x			x?	x	x			x
Total Organic Carbon (TOC)								x	x	x?	x?	x			x
Notes:															
Where an entry contains a year within parentheses, e.g., x(86) this implies that particular compound was measured in 1986 but not in the other years covered in that column.															
Where the entry is x? there is an uncertainty as to whether the exact compound listed has been measured.															
In the column referring to Level, I and II are designations for substances as specified in the Binational Virtual Elimination Strategy (BVES); additional compounds measured are also included in this table															
II - A refers to PAH compounds added to form a group of PAH compounds for this analysis; it is only one of many possible groups of PAH's that could be considered															
* Reference to Total PCBs is usually simply the sum of the subset of congeners measured, not a category to be compared between measurements.															
+ For the cruises in 1992, 93, 94, and 95 there was reportedly PCB contamination on the research vessels, making the PCB measurements uncertain.															
** Additional compounds may also have been measured in Suite 12															

i. Metal/Organometallics

Total mercury and methyl mercury were measured in Lake Michigan during 1994 and 1995 as part of the Lake Michigan Mass Balance Study (U.S. EPA 1995; Mason and Sullivan, 1997). This study was carried out over two years and mercury measurements were made during six cruises [June, August, and October/November 1994 and April, August, and September/October 1995], with approximately 20 samples collected per cruise. Determinations of both dissolved and particulate-phase concentrations were made.

Cleckner et al. (1995) report measurements of mercury at a site in Lake Michigan 6.4 km east of Chicago, where the water depth was about 12 meters. Eight samples at depths of 0.3 and 10 meters, and 7 samples of the surface micro-layer were collected over a 4-day period in September 1993 and analyzed for total mercury.

Gill and Bruland (1990) report measurements of particulate and dissolved mercury in Lake Ontario and Lake Erie (collected near the shores, as reported by Mason and Sullivan, 1997). Analysis of the dissolved fraction for total organo-mercury compounds and total reactive mercury was also carried out.

Reports of measurements of other metals in the Great Lakes were found in the literature. For example, measurements of cadmium in Lake Ontario were reported by Nriagu et al. (1993). In this study, ultra-clean techniques were used, and the authors stated that the "preliminary results obtained using the protocols described cast doubt on most of the published data on trace metal concentrations in the Great Lakes." *Nriagu found a dramatic decrease in cadmium with depth, suggesting that atmospheric deposition may be very important for this compound.* High levels of cadmium were also found near the bottom of the lake, presumably released from metal-rich sediments. Cadmium in the Great Lakes ecosystem — including potential human health effects from eating contaminated fish — was recently discussed by Bernier *et al.* (1995).

No data or reports of measurements for tributyltin or alkylated lead in the water of any of the Great Lakes could be identified through searches of the literature or discussion with government regulatory/monitoring personnel familiar with monitoring programs which have been conducted in the Great Lakes.

ii. Organochlorine Biocides

Stevens and Nielson (1989) report data from a cruise in spring 1986 for Lakes Superior, Huron, Erie and Ontario, carried out under the auspices of Environment Canada's Inland Waters Directorate. The set of organochlorine compounds measured were: aldrin, DDT and metabolites, dieldrin, endrin, α - and β -Endosulphan, heptachlor, heptachlor epoxide, α - and γ -HCHs, methoxychlor, and mirex.

Environment Canada's Inland Waters Directorate has continued to measure a range of compounds in this category (L'Italien 1993, 1996ab, 1997). For all the cruises during the 1986-90 period involving Lake Superior, Huron, Erie and Ontario, as well as the 92-93 study of Ontario, the 94-95 study of Erie, and the 96-97 study of Superior, the following organochlorine biocides have or are being measured: aldrin, α - and γ -chlordane, DDT and metabolites, dieldrin, endrin, α - and β -Endosulphan, heptachlor, heptachlor epoxide, α - and γ -HCHs, methoxychlor, and mirex. In addition, except for the 1986-90 studies, endrin aldehyde and photomirex were also measured.

For 1986-90 Environment Canada work, total water column — dissolved plus particulate — concentrations are reported. In more recent work (1992-95), pollutant concentrations associated with the particulate phase (i.e., suspended sediment) were also attempted. In the studies of Lake Erie (1994-95) and Lake Ontario (1992-93), the levels of suspended sediments were too low to reliably measure particulate-phase concentrations. When measurements could be made, the proportion of many organic pollutants associated with the particulate phase was often on the order of 1% or less. Lake Superior has even lower suspended sediment concentrations than Lake Erie or Ontario, and thus, the fraction associated with suspended sediments may be even lower.

During 1996-97, Environment Canada is conducting 2 cruises (spring and summer) sampling only the surface waters. Sample volumes of 100 liters will be taken, and only the dissolved phase concentrations will be measured (L'Italien, 1997).

Bidleman and McConnell carried out cruises with the EPA in Green Bay in June of 1989 and Lakes Michigan, Huron, Erie and Ontario during August 1990. Organochlorine biocides measured included alpha- and gamma-hexachlorocyclohexanes. Measurements were reported for the dissolved phase only.

In 1992 and 1993, EPA annual cruises measured DDT and its metabolites for all five lakes and atrazine in 1993. The EPA makes reference to total water column concentrations only (SOLEC, 1994). During the Lake Michigan Mass Balance Study (1994-1995), organochlorine biocides measured included atrazine and metabolites and trans-nonachlor.

In 1994, the US EPA Region II utilized *Lake Guardian* research vessel for a September cruise on Lake Ontario. The following organochlorine compounds were measured: aldrin, atrazine, α - and γ -chlordane, trans- and cis-nonachlor, p-p'-DDD, p,p'-DDE, dieldrin, endrin, heptachlor, heptachlor epoxide, α - and γ -HCHs, mirex, and toxaphene. Measurements of pollutant concentrations in the total water-column and suspended sediment phase were made (U.S. EPA Region II, 1994).

iii. Industrial/Miscellaneous

For the two-year studies of Lake Ontario (1992-93), Lake Erie (1994-95) and Lake Superior (1996-97), Environment Canada measurements include the following compounds in this category: hexachloro-1,3 butadiene and octachlorostyrene. Hexachlorocyclopentadiene is/was also measured. For the latter two studies, measurements of a number of phthalates are included.

The only other measurement that could be identified for compounds in this pollutant category were those made for octachlorostyrene by the U.S. EPA Region II, during a September 1994 cruise on Lake Ontario (U.S. EPA Region II, 1994).

iv. Chlorobenzenes

Stevens and Nielson (Envr. Canada Inland Waters Directorate) carried out an annual spring cruise in 1986 for Lakes Superior, Huron, Erie and Ontario. The set of chlorobenzenes measured were: 1,2-, 1,3-, and 1,4-dichlorobenzene, 1,2,3-, 1,2,4-, and 1,3,5-trichlorobenzene, 1,2,3,4-tetrachlorobenzene, pentachlorobenzene and hexachlorobenzene. Environment Canada has continued to measure the same group of chlorobenzenes as part of their cruises on the Great Lakes (L'Italien, 1993, 1996ab, 1997), including the cruises during the 1986-90 period involving Lakes Superior, Huron, Erie and Ontario, the 1992-93 study of Lake Ontario, the 1994-95 study of Lake Erie, and the 1996-97 study of Lake Superior.

U.S. EPA Region II measured several chlorobenzenes during their September 1994 cruise on Lake Ontario: 1,2,3-, 1,2,4-, and 1,3,5-trichlorobenzene, 1,2,3,4-, 1,2,3,5-, and 1,2,4,5-tetrachlorobenzene, pentachlorobenzene and hexachlorobenzene.

v. Polychlorinated Dibenzo-P-Dioxins & Dibenzofurans

There were water measurements made for dioxin compounds for 2,3,7,8-tetrachloro-dibenzo-*p*-dioxin by Serge L'Italien and Environment Canada for Lake Erie in their 1994-1995 study and in their current study of Lake Superior for 1996 and 1997. In the case of Lake Erie (1994-95), they carried out cruises each year in the Spring, Summer and Fall, each with 9 sampling stations. The current project for Lake Superior involves two cruises a year in the Spring and Summer, each with 14 sampling stations.

Cook and Burkhard measured PCDD/F as an adjunct research project of the Lake Michigan Mass Balance Study in water samples from Lake Michigan in 1995 (Cook, 1997).⁴

⁴. Cook and Burkhard report that EPA Method 1613 is being used for the analysis of dioxins and furans in water samples (Cook, 1997). This method includes

vi. Polychlorinated Biphenyls (PCBs)

There are a total of 209 different PCB congeners. The detailed lists of all the individual congener measurements have not been included in Table 13, as it was beyond the scope of this analysis to consider congener-specific details in relation to PCB monitoring programs. Instead, the total number of congeners measured is listed.⁵

Stevens and Nielson (Inland Waters Directorate) report PCB measurements in terms of Aroclors 1242, 1254, and 1260. These measurements were part of an annual spring cruise in 1986 for Lakes Superior, Huron, Erie and Ontario. Since 1986, Environment Canada has made measurements of total PCBs. In the case of the 92-93 study of Ontario, there is an explicit reference to the Aroclors 1242, 1254 and 1260, but it is not clear whether this same combination corresponds to references to total PCBs for the other years. Unfortunately, PCB contamination was discovered on the research vessels for the cruises in 1992, 1993, 1994, and 1995, making the PCB measurements questionable (L'Italien, 1997).

PCB measurements were made for all the cruises in which Eisenreich and others were involved for Lake Superior. In 1986 only 35 PCB congeners were measured but for 1988, 1990 and 1992 a subset of 82 congeners were measured. Jeremiason *et al.* (1994) refers to measurements for both the particulate and dissolved fractions for the 1988, 1990 and 1992 data. Baker and Eisenreich (1990) present data from the 1986 cruise; only dissolved-phase measurements are given and no reference is made to the particulate-phase concentrations.

The EPA has conducted annual cruises for all the Great Lakes since 1991 except for 1994-95 when their efforts were concentrated on the Lake Michigan Mass Balance Study. For the years 1991-1993, the EPA conducted both a spring and summer cruise for at least 6 stations per lake. For 1996 and 1997, there was just a spring cruise for six stations on each lake. For all these cruises, measurements have been made for all 209 PCB congeners, except for the Lake Michigan Mass Balance Study (LMMBS) which measured only 65 PCB congeners. Measurements of both dissolved and particulate-phase concentrations were made in the LMMBS.

measurements of all seventeen 2,3,7,8-substituted tetrachloro- through octachloro-dibenzo-p-dioxins and dibenzofurans (e.g., Telliard, et al.).

⁵. As mentioned as a note to Table 13, reference to "total PCBs" is usually simply the sum of the subset of congeners measured, not the actual total PCB's in the sample. Thus, "total PCB" data from measurement programs analyzing different subsets of the 209 PCB congeners cannot be compared easily.

Finally, U.S. EPA Region II measured a total of 135 PCB congeners during their September 1994 cruise on Lake Ontario.

vii. Polycyclic Aromatic Hydrocarbons (PAHs)

Environment Canada has been regularly measuring a range of PAH's in the Great Lakes. In 1987, the only PAH measured for the cruises on Lake Superior and Huron was fluoranthene. For the cruises during 1988 and 1990 involving Lake Huron, and Ontario, as well as the 1992-93 study of Ontario, the 1994-95 study of Erie, and the 1996-97 study of Superior, the following PAH's have or are being measured: acenaphthylene, fluoranthene, benzo[b+k] fluoranthene, fluorene, 2-chloronaphthalene, 1-methylnaphthalene, 2-methylnaphthalene, benzo[g,h,i] perylene, phenanthrene, pyrene, benzo[a]pyrene, and indeno[1,2,3-c,d] pyrene. For the cruises during 1988 and 1990 involving Lake Huron, and Ontario, as well as the 92-93 study of Ontario, two additional PAH's were measured: indene, and 1,2,3,4-tetrahydronaphthalene. In the most recent cruises on Lake Erie in 1994-95 and Lake Superior in 1996-97, two additional PAH's are being measured: benz[a]anthracene and chrysene/triphenylene.

Eisenreich (sponsored by the EPA) conducted a cruise on Lake Superior in August of 1986 and measured the following PAH's: benz[a]anthracene, dibenz[a,h]anthracene, chrysene/triphenylene, fluoranthene, benzo[b]fluoranthene, benzo[k]fluoranthene, fluorene, benzo[g,h,i]perylene, phenanthrene, pyrene, benzo[a]pyrene, benzo[e]pyrene, and indeno[1,2,3-c,d]pyrene.

The September 1994 cruise on Lake Ontario by the US EPA Region II measured the following PAH's: acenaphthylene, anthracene, benz [a] anthracene, dibenz [a, h] anthracene, chrysene, fluoranthene, benzo [b] fluoranthene, benzo [k] fluoranthene, fluorene, naphthalene, benzo[g,h,i] perylene, phenanthrene, pyrene, benzo[a] pyrene, benzo[e]pyrene, and indeno[1,2,3-c,d] pyrene.

viii. Suspended Sediment Properties

These properties are not part of the BVES compound list but are relevant for much of the work involving water measurements, similar to the importance of particulate measurements in interpreting air concentration data. Although, most cruises measure total suspended solids, we were unable to confirm whether all cruises have done so.

The last three entries in Table 13 are dissolved, particulate and total organic carbon, and again, it was not possible to determine during the course of this study whether such measurements had been made or not for some of the cruises. Information on the nature of suspended sediment measurements that were made (or not made) could not be obtained for the Environment Canada cruises and the Eisenreich et al. cruises in Lake Superior.

B. Summary of Monitoring of Persistent Toxic Substances in the Water of the Great Lakes

The detailed information in Tables 12 and 13 and discussed in the narrative above is summarized in Table 14, below. In this table, the measurement programs for each BVES compound for each Great Lake for the period from 1986 - 1996 are summarized.

It can be noted that for many of the compounds that are being measured in water monitoring programs in the Great Lakes, the measurements are relatively infrequent. This may pose a challenge for the accurate estimate of the direction and rate of gas exchange for those compound for which this phenomenon is relevant. Typically, data from 1 or more years previous must be used when attempting to make such gas-exchange estimates.

Table 14. Summary of Water Monitoring of BVES Persistent Toxic Substances in the Great Lakes (1986-1996)

Notes: The number of measurement locations or stations for each cruise of a particular study is indicated in parentheses, e.g., (14,14) indicates that there were two cruises with 14 stations each.

Abbrev.: Lvl = BVES Level; DS = D. Swackhamer; EC = Envr. Canada; EPA = U.S. Envr. Protection Agency; S&N = Stevens & Nielson; LMMBS = L. Mich Mass Balance Study; C&B = Cook and Burkhard, EPA National Health and Environmental Effects Research Laboratory, Duluth, Mn.

Compound or Group	Lvl	Lake Superior	Lake Huron	Lake Michigan	Lake Erie	Lake Ontario
METALS / ORGANOMETALLICS						
Total Cadmium	II					EC '91 (2) many different depths sampled (Nriagu et al., 1993)
Individual Cadmium Species	II					
Alkylated Lead	I					
Total Mercury	I			LMMBS '94-95, (6 cruises, ~ 20 samples per cruise) Cleckner '93 (1) [Cleckner et al. (1995)]	Gill and Bruland '87 (1? near shore?) [Gill and Bruland (1990)]	Gill and Bruland '87 (1? near shore?) [Gill and Bruland (1990)]
Individual Mercury Species	I			LMMBS '94-95 (6 cruises, ~ 20 samples per cruise) (methyl mercury)	Gill and Bruland '87 (1? near shore?) [Gill and Bruland (1990)] total dissolved organo- Hg and total dissolved reactive Hg	Gill and Bruland '87 (1? near shore?) [Gill and Bruland (1990)] total dissolved organo- Hg and total dissolved reactive Hg
Tributyltin	II					

Table 14. Summary of Water Monitoring of BVES Persistent Toxic Substances in the Great Lakes (1986-1996)

Notes: The number of measurement locations or stations for each cruise of a particular study is indicated in parentheses, e.g., (14,14) indicates that there were two cruises with 14 stations each.

Abbrev.: Lvl = BVES Level; DS = D. Swackhamer; EC = Envr. Canada; EPA = U.S. Envr. Protection Agency; S&N = Stevens & Nielson; LMMBS = L. Mich Mass Balance Study; C&B = Cook and Burkhard, EPA National Health and Environmental Effects Research Laboratory, Duluth, Mn.

Compound or Group	Lvl	Lake Superior	Lake Huron	Lake Michigan	Lake Erie	Lake Ontario
ORGANOCHLORINE BIOCIDES						
Aldrin / Dieldrin	I	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38)	C&B '94 (3)	S&N '86 (21) EC '86 (21) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Chlordane		EC '86 (22) EC '87 (22) EC '96-97 (14,14)	EC '86 (38) EC '87 (38) EC '88 (38)	C&B '94 (3)	EC '86 (21) EC '94-95 (9,9,9)	EC '86,88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
DDT / DDD / DDE	I	S&N '86 (19) EC '86 (22) EC '87 (22) EPA '92-93 (6) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38) EPA '92-93 (6)	EPA '92-93 (6) C&B '94 (3)	S&N '86 (21) EC '86 (21) EPA '92-93 (6) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6) EPA '92-93 (6) EPA Region II '94 (30)
Endrin	II	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38)	C&B '94 (3)	S&N '86 (21) EC '86 (21) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Heptachlor / Heptachlor Epoxide	II	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38)	C&B '94 (3)	S&N '86 (21) EC '86 (21) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Hexachlorocyclohexanes	II	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38) EPA (Bidleman) '90 (10)	EPA (Bidleman) '89 (3- Green Bay) EPA (Bidleman) '90 (11) C&B '94 (3)	S&N '86 (21) EC '86 (21) EPA (Bidleman) '90 (18) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EPA (Bidleman) '90 (7) EC '92-93 (6,6,6) EPA Region II '94 (30)
Methoxychlor	II	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38)	C&B '94 (3)	S&N '86 (21) EC '86 (21) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6)
Mirex	I	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38)	C&B '94 (3)	S&N '86 (21) EC '86 (21) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Pentachlorophenol	II					
Toxaphene	I	DS '91,93,96,97 (~7)	DS '91 (~7)	DS '91,93,94,95,97 (~7) C&B '94 (3)	DS '91 (~7)	DS '91 (~7) EPA Region II '94 (30)

Table 14. Summary of Water Monitoring of BVES Persistent Toxic Substances in the Great Lakes (1986-1996)

Notes: The number of measurement locations or stations for each cruise of a particular study is indicated in parentheses, e.g., (14,14) indicates that there were two cruises with 14 stations each.

Abbrev.: Lvl = BVES Level; DS = D. Swackhamer; EC = Envr. Canada; EPA = U.S. Envr. Protection Agency; S&N = Stevens & Nielson; LMMBS = L. Mich Mass Balance Study; C&B = Cook and Burkhard, EPA National Health and Environmental Effects Research Laboratory, Duluth, Mn.

Compound or Group	Lvl	Lake Superior	Lake Huron	Lake Michigan	Lake Erie	Lake Ontario
INDUSTRIAL / MISCELLANEOUS						
4-Bromophenyl Phenyl Ether	II			C&B '94 (3) (?)		
3,3'-Dichlorobenzidene	II					
Hexachloro-1,3-Butadiene	II	EC '96-97 (14,14)		C&B '94 (3)	EC '94-95 (9,9,9)	EC '92-93 (6,6,6)
4,4'-Methylene bis(2-Chloroaniline)	II					
Octachlorostyrene	I	EC '96-97 (14,14)		C&B '94 (3)	EC '94-95 (9,9,9)	EC '92-93 (6,6,6) EPA Region II '94 (30)
CHLOROBENZENES						
1,4-dichlorobenzene	II	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38)		S&N '86 (21) EC '86 (21) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6)
Tetrachlorobenzenes	II	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38)	C&B '94 (3)	S&N '86 (21) EC '86 (21) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Pentachlorobenzene	II	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38)	C&B '94 (3)	S&N '86 (21) EC '86 (21) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Hexachlorobenzene	I	S&N '86 (19) EC '86 (22) EC '87 (22) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38)	C&B '94 (3)	S&N '86 (21) EC '86 (21) EC '94-95 (9,9,9)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)

Table 14. Summary of Water Monitoring of BVES Persistent Toxic Substances in the Great Lakes (1986-1996)

Notes: The number of measurement locations or stations for each cruise of a particular study is indicated in parentheses, e.g., (14,14) indicates that there were two cruises with 14 stations each.

Abbrev.: Lvl = BVES Level; DS = D. Swackhamer; EC = Envr. Canada; EPA = U.S. Envr. Protection Agency; S&N = Stevens & Nielson; LMMBS = L. Mich Mass Balance Study; C&B = Cook and Burkhard, EPA National Health and Environmental Effects Research Laboratory, Duluth, Mn.

Compound or Group	Lvl	Lake Superior	Lake Huron	Lake Michigan	Lake Erie	Lake Ontario
POLYCHLORINATED DIBENZO-P-DIOXINS & DIBENZOFURANS (PCDD/F'S)						
2,3,7,8-TCDD	I	EC '96-97 (14,14)		C&B '94 (3)	EC '94-95 (9.9.9)	
other PCDD/F congeners	I			C&B '94 (3)		
POLYCHLORINATED BIPHENYLS (PCB'S)						
PCB's	I	S&N '86 (19) Eisenreich, '86,88,90,92 (5 or 6) EC '86 (22) EC '87 (22) EPA '91-93 (6,6) EPA '96-97 (6) EC '96-97 (14,14)	S&N '86 (18) EC '86 (38) EC '87 (38) EC '88 (38) EPA '91-93 (6,6) EPA '96-97 (6)	EPA '91-93 (6,6) C&B '94 (3) LMMBS '94-95, (8 cruises, 41 stations) EPA '96-97 (6)	S&N '86 (21) EPA '91-93 (6,6) EC '94-95 (9.9.9) EPA '96-97 (6)	S&N '86 (33) EC '86,88,90 (55) EC '92-93 (6,6,6) EPA '91-93 (6,6) EPA Region II '94 (30) EPA '96-97 (6)
POLYCYCLIC AROMATIC HYDROCARBONS						
Acenaphthene	II-a			C&B '94 (3)		
Acenaphthylene	II-a	EC '96-97 (14,14)	EC '88 (38)	C&B '94 (3)	EC '94-95 (9.9.9)	EC '88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Anthracene	II			C&B '94 (3)		EPA Region II '94 (30)
Benz [a] Anthracene	II	Eisenreich, '86 (6) EC '96-97 (14,14)		C&B '94 (3)	EC '94-95 (9.9.9)	EPA Region II '94 (30)
Benzo [b]Fluoranthene	II-a	Eisenreich, '86 (6)		C&B '94 (3)		EPA Region II '94 (30)
Benzo [j]Fluoranthene	II-a					
Benzo [k]Fluoranthene	II-a	Eisenreich, '86 (6)		C&B '94 (3)		EPA Region II '94 (30)
Benzo [a] Pyrene	I	Eisenreich, '86 (6) EC '96-97 (14,14)	EC '88 (38)	C&B '94 (3)	EC '94-95 (9.9.9)	EC '88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Benzo [e] Pyrene	II-a	Eisenreich, '86 (6)				EPA Region II '94 (30)
Benzo [g,h,i] Perylene	II	Eisenreich, '86 (6) EC '96-97 (14,14)	EC '88 (38)	C&B '94 (3)	EC '94-95 (9.9.9)	EC '88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)

Table 14. Summary of Water Monitoring of BVES Persistent Toxic Substances in the Great Lakes (1986-1996)

Notes: The number of measurement locations or stations for each cruise of a particular study is indicated in parentheses, e.g., (14,14) indicates that there were two cruises with 14 stations each.

Abbrev.: Lvl = BVES Level; DS = D. Swackhamer; EC = Envr. Canada; EPA = U.S. Envr. Protection Agency; S&N = Stevens & Nielson; LMMBS = L. Mich Mass Balance Study; C&B = Cook and Burkhard, EPA National Health and Environmental Effects Research Laboratory, Duluth, Mn.

Compound or Group	Lvl	Lake Superior	Lake Huron	Lake Michigan	Lake Erie	Lake Ontario
POLYCYCLIC AROMATIC HYDROCARBONS (continued)						
Chrysene	II-a	Eisenreich, '86 (6) EC '96-97 (14,14)		C&B '94 (3)	EC '94-95 (9.9.9)	EPA Region II '94 (30)
Dibenz [a,h] Anthracene	II-a	Eisenreich, '86 (6)		C&B '94 (3)		EPA Region II '94 (30)
Dinitropyrenes	II					
Fluoranthene	II-a	EC '87 (22) Eisenreich, '86 (6) EC '96-97 (14,14)	EC '87-88 (38)	C&B '94 (3)	EC '94-95 (9.9.9)	EC '88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Fluorene	II-a	Eisenreich, '86 (6) EC '96-97 (14,14)	EC '88 (38)	C&B '94 (3)	EC '94-95 (9.9.9)	EC '88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Indeno [1,2,3-cd] Pyrene	II-a	Eisenreich, '86 (6) EC '96-97 (14,14)	EC '88 (38)	C&B '94 (3)	EC '94-95 (9.9.9)	EC '88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Naphthalene	II-a	EC '96-97 (14,14)		C&B '94 (3)	EC '94-95 (9.9.9)	EPA Region II '94 (30)
Phenanthrene	II	Eisenreich, '86 (6) EC '96-97 (14,14)	EC '88 (38)	C&B '94 (3)	EC '94-95 (9.9.9)	EC '88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Perylene	II					
Pyrene	II-a	Eisenreich, '86 (6) EC '96-97 (14,14)	EC '88 (38)	C&B '94 (3)	EC '94-95 (9.9.9)	EC '88,90 (55) EC '92-93 (6,6,6) EPA Region II '94 (30)
Other PAH's		EC '87 (22) EC '96-97 (14,14)	EC '88 (38)	C&B '94 (3)	EC '94-95 (9.9.9)	EC '88,90 (55) EC '92-93 (6,6,6)

5. SPATIAL REPRESENTATIVENESS of AMBIENT MONITORING DATA for the ESTIMATION of the NET LOADING of PERSISTENT TOXIC SUBSTANCES to the GREAT LAKES

It has long been recognized that a fundamental question in relation to the use of ambient monitoring data to estimate loadings to the Great Lakes is the following:

Can data from one or only a few sampling locations in the proximity of a particular Great Lake be used to accurately estimate deposition to the entire Lake?

This question has particular relevance to the use of the data from the Integrated Atmospheric Deposition Network (IADN), which has one master station for each Great Lake. In estimating atmospheric deposition to each of the Great Lakes using IADN data, data from the one master station per Lake is used to represent the atmospheric concentrations of pollutants for the entire Lake (Hoff et al., 1996).

There have been several recent attempts to characterize spatial variability in air and water concentrations in the Great Lakes.

A. Lake Michigan Urban Air Toxics Study

In the Lake Michigan Urban Air Toxics Study (LMUATS) (Keeler, 1994; Pirrone *et al.*, 1995ab), ambient air samples were collected at three sites for one month during the summer of 1991:

- (1) on a building roof at Illinois Institute of Technology in Chicago, Illinois;
- (2) onboard a research ship located on Lake Michigan 5-10 km offshore of Chicago;
- (3) at South Haven, MI, a rural site approximately 130 km northeast of Chicago, about 3 km inland from the opposite (eastern) shore of Lake Michigan

In Table 15 below, the BVES pollutants (Table 1) measured in LMUATS are listed.

Additional biocides (trans-nonachlor, atrazine, chloropyrifos, simazine, metolachlor, and alachlor) and PAH's (fluorenone, retene, cyclopenta(c,d)pyrene, and coronene) were also measured in the organics series.

Water concentrations were not measured for any of the organics during the study.

Table 15. BVES Compounds for which air measurements were made in the Lake Michigan Urban Air Toxics Study

<p>Metals/Organometallics Mercury (total) Lead (total)</p>	<p>PCB's 8 specific congeners homologue group totals, e.g., monochloro-PCB's, dichloro-PCB's, etc.</p>
<p>Organochlorine Biocides α-HCH ; γ-HCH aldrin and dieldrin mirex α- and γ-chlordane and trans-nonachlor 4,4'-DDT, p,p'-DDE, and p,p'-DDD</p>	<p>PAH's Naphthalene Acenaphthylene Acenaphthene Fluorene Phenanthrene Anthracene Fluoranthene Pyrene Benz[a]Anthracene Chrysene Benzofluoranthenes (mixed) Benzo[e]Pyrene Benzo[a]Pyrene Indeno[1,2,3-c,d]Pyrene Dibenz[a,h]Anthracene Benzo[g,h,i]Perylene</p>
<p>Industrial Miscellaneous: hexachlorobutadiene [no data presented in Keeler (1994) or Pirrone <i>et al.</i> (1995b)]</p>	
<p>Chlorobenzenes hexachlorobenzene 1,4-dichlorobenzene [no data presented in Keeler (1994) or Pirrone <i>et al.</i> (1995b)]</p>	

Land-based measurements were made for one month from July 8 to August 9, 1991. Daily 12-hour samples were taken at each land-based site. Over-lake samples 5-10 km offshore of Chicago were taken during two sampling periods (5 days and 3 days in duration). Approximately 17 samples were taken during these periods at the over-lake site. Total (vapor + particle) concentrations of organic compounds were measured.


In Figures 6 and 7, below, the average concentrations of biocides, PCB's, and PAH's measured at the three sampling sites are shown.⁶ For each pollutant, the concentrations at each site have been normalized (i.e., divided by) the average concentration of the pollutant measured at Chicago. The data presented in Figure 6 show that while there were significant differences in ambient air concentrations

⁶. Only a subset of the total number of compounds has been plotted in Figures 2 and 3. The compounds plotted are those for which data were given in Pirrone *et al.* (1995b). Data for additional compounds is given by Keeler (1994), but these were not included in these figures.

between different sites, there was no consistent trend for organochlorine biocides and HCB, i.e., the relative ordering of sites with respect to concentration changes from compound to compound. For the PCB's (also shown in Figure 6) and PAH's (shown in Figure 7), there was a consistent, relatively dramatic difference between the sites. The site in Chicago consistently had the highest concentration. Also, for each individual PAH and for total PCB's, the over-lake site 5-10 km offshore of Chicago had higher concentrations than the South Haven site on the other side of the Lake. For PCB's and PAH's, the relative magnitude of concentrations was generally consistent with the following pattern:

[Chicago] >> [over-lake, 5-10 km off-shore of Chicago] > [South Haven]


Figure 6. Spatial Variations in Air Concentrations of Biocides and PCB's
 Lake Michigan Urban Air Toxics Study (Pirrone, Keeler & Holsen, 1995)



Std. Dev. plotted as "error bar" around a given avg. conc.; Each conc. divided by Chicago avg (i.e., Chicago avg = 1);
 R/V Laurentian: 5-10 km offshore of Chicago; South Haven: rural, ~130 km NE of Chicago, near eastern shore of L. Mich.

Figure 7. Spatial Variations in Air Concentrations of PAH's
 Lake Michigan Urban Air Toxics Study (Pirrone, Keeler & Holsen, 1995)

Air Concentration (relative to Chicago avg.)



Std. Dev. plotted as "error bar" around a given avg. conc.; Each conc. divided by Chicago avg (i.e., Chicago avg = 1);
 R/V Laurentian: 5-10 km offshore of Chicago; South Haven: rural, ~130 km NE of Chicago, near eastern shore of L. Mich.

B. Lake Michigan Mass Balance Study

Results of measurements made in the Lake Michigan Mass Balance Study are being assembled and presented elsewhere, and these data were considered to be outside of the scope of this report. One set of the LMMBS data was obtained in the course of this analysis, however. A brief mention of some of these data will be made here, as they are particularly relevant to the question of spatial variability.

Over-lake air measurements in the northern, central, and southern portions of Lake Michigan during a summer 1994 cruise are summarized in Table 16 (Sweet, 1996).

Table 16. Summary of Air Monitoring Data from a Cruise On Lake Michigan During Summer 1994 (Sweet, 1996)					
Lake Region	Pollutant Concentrations (average values, pg/m ³ , for samples taken at two locations in each region)				
	∑PCB's	DDE	Dieldrin	Benzo(a)Pyrene	Lead
Northern Region	4,800	5	63	28	8,000
Central Region	4,600	8	51	27	9,000
Southern Region	5,100	16	150	180	86,000
Ratio of the Average Concentration in the Southern Region to the Average Central-North Concentration					
	1.1	2.5	2.6	6.5	10.1

It can be seen from the data in the above table that the over-lake concentrations of the reported pollutants were roughly the same in the north and central portions of Lake Michigan (during the cruise), but that for some pollutants, most notably benzo(a)pyrene and lead, the levels in the air in the southern portion of the Lake were relatively much higher.

As has been observed by many, emissions from the heavily populated urban centers surrounding the southern portion of Lake Michigan (e.g., the Chicago metropolitan area) can exert a strong influence on the levels of pollution in nearby regions of the Lake. For pollutants associated with combustion processes in urban areas such as benzo(a)pyrene and lead, this influence is clearly seen in the above

data. In the measurements made during this cruise, large spatial variability was not found for PCB's, DDE, or dieldrin in the air above Lake Michigan.

In another part of this study, air and precipitation measurements were made at nine sites in the vicinity of Lake Michigan. These data are summarized in Table 17 (from Sweet, 1996). The data shown are the average of 12 monthly composite samples. As discussed by Clyde Sweet (1996), these data, together with those of Table 16, show several interesting patterns, including the following:

- The highest values of PCB's, BaP, and lead in air and precipitation were found in Chicago IL.
- The highest concentrations of DDE in air and precipitation were found at South Haven MI, suggesting a localized source (e.g., an area of high past use of DDT).
- There was a distinct south-north trend for dieldrin in both air and precipitation (concentrations higher at the southern sites, lower at the northern sites), consistent with the heavy past use of aldrin in agricultural areas south of Lake Michigan (dieldrin is a persistent breakdown product of aldrin).
- The second highest average measured air concentration of PCB's (750 pg/m^3) was at Beaver Island, a remote island in the northern part of Lake Michigan. The concentration of PCB's was four times higher here than at the other remote sampling site, at Sleeping Bear Dunes (the IADN master station for Lake Michigan). The concentrations of PCB's at Beaver Island in the summer months was particularly high.
- Except for PCB's, concentrations measured at the shoreline sites (Table 17) were similar to those measured at over-lake sites (Table 16). Over-water concentrations were significantly higher at over-water sites than at all shore-line sites except for Chicago.
- The above two results are consistent with a net volatilization of PCB's *from Lake Michigan* during the summer months, increasing the over-water concentrations and the concentration at Beaver Island (air parcels arriving at Beaver Island spent a relatively long time traversing the water of Lake Michigan). [This result is consistent with that of Hornbuckle *et al.* (1993) who found higher concentrations of PCB's over the water surface of Green Bay than over surrounding land areas.]
- Average concentrations of lead in precipitation was fairly uniform at all sampling sites.
- Average concentrations of BaP in precipitation were generally highest in urban areas, with the concentration at Chicago 10 - 100 times greater than at any other location.
- Average concentrations of PCB's in precipitation were relatively uniform at the shoreline sampling stations, although the levels measured at Chicago were somewhat higher than any other location measured.
- Based on the measurements, estimating the overall wet deposition to Lake Michigan using data from Sleeping Bear Dunes would seem to be a relatively accurate approach. However, estimating the overall dry deposition of from data at Sleeping Bear Dunes would underestimate the loading, especially for BaP and lead.

Table 17. Summary of Air and Precipitation Monitoring Data at Nine Sites in the Vicinity of Lake Michigan (Sweet, 1996)

Monitoring Location (all sites near the shore except Bondville)	North-South Orientation Relative to Lake	East-West Orientation Relative to Lake	General Character of Sampling Site	Average Pollutant Concentrations in Air and Precipitation (air concentrations in units of pg/m ³ ; concentrations in precipitation in units of ng/liter)									
				∑PCB's		DDE		Dieldrin		Benzo(a)Pyrene		Lead	
				air (vapor)	precip.	air (vapor)	precip.	air (vapor)	precip.	air (particle phase)	precip.	air (particle phase)	precip.
Manitowoc, WI	central	west	urban	218	5.1	16	0.3	40	1.5	34	1.8	12,000	2,700
Chiwaukee, WI	south-central	west	urban	282	2.4	13	0.3	69	1.0	63	14	15,000	1,900
Chicago, IL	south	west	urban	2400	13.0	33	1.0	247	5.8	809	178	38,000	3,300
Bondville, IL	~200 km south of Chicago		urban	196	1.8	9	0.1	606	--	20	< 1	8,000	1,400
Indiana Dunes, IN	south	east	urban	571	1.7	14	0.1	162	1.7	172	15	17,000	2,200
South Haven, MI	south-central	east	rural	241	5.0	814	2.5	245	2.3	59	3.0	8,000	2,100
Muskegon, MI	central	east	urban	504	3.2	27	0.3	103	0.2	48	4.9	10,000	2,100
Sleeping Bear Dunes, MI	north-central	east	remote	183	2.4	14	0.1	40	0.7	9	1.8	2,000	1,500
Beaver Island, MI	north	center	remote	750	1.7	3	0.3	19	0.5	12	3.4	4,000	1,300

C. Ambient Air Monitoring of Persistent Toxic Substances Conducted by the Michigan Department of Environmental Quality

Organochlorine compounds in the vapor phase were measured at three sites in the region of the upper Great Lakes during 1990-1991 (Monosmith and Hermanson, 1996), and additional work in the region is continuing (Monosmith, 1997; Hermanson *et al.*, 1997). During the 1990-1991 sampling period, the following compounds were measured, simultaneously, at sites near Grand Traverse Bay, Saginaw Bay, and Sault Ste. Marie, Michigan:

- 25 PCB congeners⁷
- Hexachlorobenzene
- p,p'-DDT and p,p'-DDE
- α -Hexachlorocyclohexane (HCH) and γ -HCH

One 48-hour sample was taken each month, from November 1990 through October 1991, at each of the three sites.

Concentrations of all compounds measured tended to be higher in the summer than in the rest of the year. The sampling site at Sault Ste. Marie had the lowest concentrations of PCB's, while the site at Traverse had the highest concentrations of most PCB's measured. For HCB, DDT/DDE, and the HCH's, there was no consistent pattern to the relative concentrations at the three sites, and for some sampling dates, there were large differences in the concentrations at the three sites.

Additional compounds being measured in the continuing work include 16 different PAH compounds, dieldrin, mirex, and 13 trace metals (not including mercury) (Monosmith 1997).

⁷. The 25 congeners chosen for detailed analysis all had relatively high vapor pressures and were expected to exist predominantly in the vapor phase in the atmosphere. The 25 congeners were represented by 14 gas-chromatograph ("GC") peaks; i.e., there were 14 peaks quantified in detail, with each peak either representing a single PCB congener or a group of PCB congeners. Σ PCB, representing the sum of 121 congeners, was also measured.

(this page left intentionally blank)

6. SUMMARY AND CONCLUSIONS

This analysis has considered ambient monitoring programs for a set of persistent toxic substances in the Great Lakes region, and the use of these monitoring data to estimate loadings to the Great Lakes.

The compounds considered were those listed in the Binational Virtual Elimination Strategy (BVES), as shown in Table 1, above.

This analysis focused primarily on the universe of ongoing, systematic, government-sponsored monitoring efforts in the Great Lakes region for air, precipitation, and the water of the Great Lakes. Detailed consideration of the monitoring data themselves or of loading estimates derived from these data was not possible to include given the limited resources available for this analysis.

An overall summary of the coverage of the BVES compounds in the monitoring programs in the Great Lakes region is given in Table 18, below, based on Sections 3 and 4 above.

In considering Table 18, and other issues discussed in this report, the following overall conclusions and observations emerge. Many of the compound-specific issues below are summarized in Table 19, as well.

- Systematic, ambient monitoring programs for air, precipitation, and water exist in the Great Lakes region for many of the BVES compounds. These data can be used to:
 - estimate the overall loading of monitored compounds to the Great Lakes (although uncertainties related to spatial representativeness for some compounds' measurements, e.g., PAH's, is a concern, and, for all of the measured compounds, there are uncertainties in the methodology of estimating the overall loading);
 - provide information on source regions, using back-trajectory modeling;
 - provide validation data for comprehensive atmospheric fate and transport models.
- It would be extremely helpful if a system could be established for making Great Lakes regional air, precipitation, and water monitoring data readily available to the public and to the research community.

- For some of the BVES compounds, there are little or no measurements of ambient air and precipitation concentrations in the Great Lakes region.
- The following BVES compounds are not being monitored comprehensively in air and/or precipitation in the IADN program [see *table 19 for additional notes regarding these and other compounds*]:
 - mercury speciation (i.e., total mercury is being measured at several locations, but, individual species are not being measured, e.g, the proportion of gaseous mercury that is Hg⁰, HgCl₂, HgO, methylmercury, etc.)
 - alkylated lead
 - 4-bromophenyl phenyl ether
 - 3,3'-dichlorobenzidene
 - 1,4-dichlorobenzene
 - PCDD/F's
 - dinitropyrenes
 - hexachloro-1,3-butadiene
 - pentachlorophenol
 - toxaphene
- The addition of 4,4'-methylene bis (2-chloroaniline) and tributyltin compounds to the list of substances monitored might be implemented on a temporary basis, to determine the potential significance of the air pathway to Great Lakes loadings for these compounds.
- Some of the BVES compounds are being monitored in Canada at one or more sites in the Great Lakes region, but not in the U.S. For example, Environment Canada has a recently established a monitoring program for octachlorostyrene, dinitropyrenes, and pentachlorophenol (in conjunction with the COA); no monitoring efforts for these compounds could be found in the U.S. in the Great Lakes region. As another example, the Canadian IADN satellite stations monitor many compounds that are not monitored at any of the U.S. sites, including Endrin, Heptachlor, Methoxychlor, Mirex, and Octachlorostyrene (only at OMOEE IADN satellite sites).
- For some of the BVES compounds, there are limited or no measurements in the water of the Great Lakes. For compounds that exist to a certain extent in the vapor phase in the atmosphere, a lack of water concentration data makes the estimation of loading to the lakes difficult.
- For many of the compounds that are being measured in water monitoring programs in the Great Lakes, the measurements are relatively infrequent. This may pose a challenge for the accurate estimate of the direction and rate of gas

exchange for those compound for which this phenomenon is relevant. Typically, data from 1 or more years previous must be used when attempting to make such gas-exchange estimates. No measurements of any BVES compounds could be identified in Lake Huron within the last 5 years.

- The spatial representativeness of monitoring locations is an issue for at least some of the BVES compounds.
 - For PAH's and other urban-source-associated pollutants, there may be strong influences of urban plumes on the loading to a given Lake, and, the representativeness of rural sampling locations is in question.
 - For PCB's and other pollutants which may be volatilizing from one or more of the Lakes, the representativeness of shoreline stations (as opposed to over-water monitoring locations) is in question.
- In addition to the spatial representativeness issue and the lack of measurements for some compounds, an incomplete understanding and incomplete monitoring of the various mass transfer processes between the atmosphere and the Lakes makes the monitoring-based estimation of the net loadings from the atmosphere to the Great Lakes somewhat uncertain.
- As an example, the deposition velocity for particle-associated material will depend on the particle size distribution, but, chemical specific particle size distributions are infrequently — if ever — measured for BVES pollutants in the Great Lakes region.
- Fog deposition, droplet resuspension, and indirect atmospheric loading (discussed briefly in Section 2) are not characterized well enough at this time to even attempt to include them in the loading estimates.
- The results of recent research, including the Lake Michigan Mass Balance Study and the AEOLOS research program, are eagerly anticipated. These may serve to significantly increase the understanding of atmospheric loading to the Great Lakes.

Table 18. Summary of Air, Precipitation and Water Monitoring Programs for BVES Substances in the Great Lakes Region

column #'s (see notes for descriptions) →	2	3	4	5	6	7	8	9	10	11	12	13	14
		V/P	LRT	Air Monitoring: G.L. Region		Precipitation Monitoring: G.L. Region		Water Monitoring: 1992-1996?					>Loading Estimate Possible?
Compound or Compound Group	Level	Vapor-Particle	Long-Range Air	Integrated Atmos. Dep. Network	Other Programs	Integrated Atmos. Dep. Network	Other Programs	Superior	Michigan	Huron	Erie	Ontario	
MERCURY and MERCURY COMPOUNDS													
Elemental Mercury (Hg ⁰)	I	v	1	-	-	-	-	-	-	-	-	-	0
Divalent Mercury (e.g., HgCl ₂)	I	v	2	-	-	-	-	-	-	-	-	-	0
Monomethyl Mercury	I	v	3	-	-	-	-	-	-	-	-	-	0
Total Gaseous Mercury	I	VV	1-3	-	ksqflr	-	-	-	-	-	-	-	~ 0 (F)
Particulate Mercury	I	PP	2	-	tqflr	-	-	-	-	-	-	-	½ (F)
Total Mercury	I	v	1-2	-	-	-	dsrqflr	-	m	-	-	-	~ 0 (F)
OTHER METALS / ORGANOMETALLICS													
Alkylated Lead	I	(V V)	2	-	-	-	-	-	-	-	-	-	0
Total Cadmium (Cd)	II	PP	2	ceou	tmwv	gnou	-	-	-	-	-	-	1
Individual Cd Species	II	PP	2	-	-	-	-	-	-	-	-	-	0
Tributyltin Compounds	II	v/p	3-B	-	-	-	-	-	-	-	-	-	0
ORGANOCHLORINE BIOCIDES													
Aldrin	I	VV	3-4	ceo	-	gno	-	e	u	-	e	eu	~ 1
Dieldrin	I	VV	3-4	ceou	mw	gnou	-	e	u	-	e	eu	~ 1
Chlordane	I			ceou	w	gnou	-	e	u	-	e	eu	~ 1
DDT / DDD / DDE	I	v/p	2	ceou	mw	gnou	-	eu	um	u	eu	eu	1
Endrin	II	VV	3	ce	-	gn	-	e	u	-	e	eu	~ 1
Heptachlor / Heptachlor Epoxide	II	VV	3-4	ceo	-	gno	-	e	u	-	e	eu	~ 1
Hexachlorocyclohexanes	II	v	1-2	ceou	mw	gnou	-	e	u	-	e	eu	~ 1
Methoxychlor	II	v/p	3	ceo	-	gno	-	e	u	-	e	e	~ 1
Mirex	I	(V V)	2	ceo	-	no	-	e	u	-	e	eu	~ 1
Pentachlorophenol	II	VV	2	-	t	-	-	-	-	-	-	-	0
Toxaphene	I	v/p	2	-	bh	-	-	s	s	-	-	u	~ ½ (C)

INDUSTRIAL / MISCELLANEOUS													
4-Bromophenyl Phenyl Ether	II	V V	3	-	-	-	-	u?	-	-	-	-	0
3,3'-Dichlorobenzidene	II	v / p	2	-	-	-	-	-	-	-	-	-	0
Hexachloro-1,3-Butadiene	II	V V	1	-	t w v	-	-	e	u	-	e	e	~ 1
4,4'-Methylene bis(2-Chloroaniline)	II	v	3 - 4	-	-	-	-	-	-	-	-	-	0
Octachlorostyrene	I	v	2	o	t	o	-	e	u	-	e	eu	~ 1
CHLOROBENZENES													
1,4-dichlorobenzene	II	V V	2	-	t y w v	g n	-	e	-	-	e	e	~ 1
Tetrachlorobenzenes	II	V V	1	o	-	g n o	-	e	u	-	e	eu	~ 1
Pentachlorobenzene	II	V V	1	o	-	g n o	-	e	u	-	e	eu	~ 1
Hexachlorobenzene	I	V V	1	ce ou	t m	g n o u	-	e	u	-	e	eu	~ 1
POLYCHLORINATED DIBENZO-P-DIOXINS & DIBENZOFURANS (PCDD/F'S) and POLYCHLORINATED BIPHENYLS (PCB'S)													
PCDD/F's	I	v / p	2	-	t p v		p (Dorset)	e	u	-	e	-	½ (G)
PCB's	I	v / p	2	ce ou	m v	n o u	-	eu	mu	u	eu	eu	1
POLYCYCLIC AROMATIC HYDROCARBONS													
Acenaphthene	II-a	V V		ce u	t a v	g n u	-	-	u	-	-	-	~ ½ (H)
Acenaphthylene	II-a	V V		ce u	t a m v	g n u	-	e	u	-	e	eu	1
Anthracene	II	V V	3	ce u	t a m v	u	-	-	u	-	-	u	~ ½ (H)
Benz (a) Anthracene	II	v / p	2	ce ou	t a w v	n o u	-	e	u	-	e	u	~ 1
Benzo (b) Fluoranthene	II-a	v / p		ce ou	t a v	n o u	-	-	u	-	-	u	~ 1
Benzo (j) Fluoranthene	II-a	v / p		-	-		-	-	-	-	-	-	0
Benzo (k) Fluoranthene	II-a	v / p		ce ou	t a v	o u	-	-	u	-	-	u	~ 1
Benzo (a) Pyrene	I	v / p	2	ce ou	t a m w v	n o u	-	e	u	-	e	eu	~ 1
Benzo (e) Pyrene	II-a	v / p		ce u	t a	u	-	-	-	-	-	u	~ ½ (H)
Benzo (g,h,i) Perylene	II	v / p	2	ce ou	t a v	o u	-	e	u	-	e	eu	~ 1
Chrysene	II-a	v / p		ce ou	t a w v	n o u	-	e	u	-	e	u	~ 1
Dibenz (a,h) Anthracene	II-a	P P		ce ou	t a v	o u	-	-	u	-	-	u	1
Dinitropyrenes	II	v / p	2	-	t	-	-	-	-	-	-	-	0
Fluoranthene	II-a	v		ce ou	t a m w v	g n o u	-	e	u	-	e	eu	~ 1
Fluorene	II-a	V V		ce ou	t a m v	g n u	-	e	u	-	e	eu	~ 1
Indeno (1,2,3-cd) Pyrene	II-a	p		ce ou	t a v	o u	-	e	u	-	e	eu	~ 1
Naphthalene	II-a	V V		-	t y m w v	-	-	e	u	-	e	u	~ ½
Phenanthrene	II	(V V)	3	ce ou	t a m w v	n o u	-	e	u	-	e	eu	~ 1
Perylene	II	v / p	2	-	t a	-	-	-	-	-	-	-	0
Pyrene	II-a	v		ce ou	t a m w v	g n o u	-	e	u	-	e	eu	~ 1

Column-by-Column Notes for Table 18.	
2	Level I or II compounds [as designated in the Binational Virtual Elimination Strategy (see text)]
3	<p>Vapor/Particle Partitioning Characteristics of Compound(s) ("VP") (based on Cohen, 1997a)</p> <p>P P = Compound is expected to exist almost entirely in the particle phase in the atmosphere (fraction adsorbed \geq 98% under all conditions) p = Compound is expected to exist mostly in the particle phase in the atmosphere (fraction adsorbed \geq 90% under all conditions) v/p = Compound is expected to exist in significant proportions in both the particle phase and the vapor phase as conditions vary v = Compound is expected to exist mostly in the vapor phase in the atmosphere (fraction adsorbed \leq 10% under all conditions) V V = Compound is expected to exist almost entirely in the vapor phase in the atmosphere (fraction adsorbed \leq 2% under all conditions)</p>
4	<p>L.R.T. (Long Range Transport) Potential (Rating)</p> <p>This analysis has not included quantitative, integrated modeling in its scope, due to time limitations. Instead, a qualitative approach to the assessment of long-range atmospheric transport has been taken. In this approach, pollutants have been generally categorized as to the relative importance of various fate mechanisms. Based on these considerations, an attempt has been made to qualitatively estimate the atmospheric lifetimes of each of the pollutants considered in this analysis. A "Long Range Air Transport Potential" rating scale of 1-4 is defined as follows:</p> <ul style="list-style-type: none"> • Rating = 1 The pollutant is extremely long-lived in the atmosphere, with an atmospheric lifetime \sim a year or longer; distribution of the pollutant is essentially global. • Rating = 2 The pollutant is relatively long-lived in the atmosphere, with atmospheric residence times on the order of at least a week to perhaps several months; long range transport can definitely occur over 1000's to 10,000's of kilometers. • Rating = 3 The pollutant is relatively short-lived in the atmosphere, with atmospheric residence times on the order of several hours to a few days; atmospheric transport may occur on regional, mesoscale distances, perhaps of several 100's to perhaps even a 1000 kilometers. • Rating = 4 The pollutant is extremely short-lived in the atmosphere, with atmospheric residence times on the order of seconds to minutes to at most an hour or so; with such pollutants, atmospheric transport of emissions will be limited to the local region around the source.
5 - 8	Air and Precipitation Measurements made in the Integrated Atmospheric Deposition Network (IADN) and other programs. See Table 3 for description of codes.
9-13	<p>Any Measurements in Great Lakes Water in the period between 1992 and 1996?</p> <p>e = measurements by Environment Canada; u = measurements by the U.S. Environmental Protection Agency m = measurements in the Lake Michigan Mass Balance Study (sponsored by U.S. EPA); s = sampling conducted by D. Swackhamer et al. (personal communication)</p>
14	<p>Loading Estimate Possible? [0 = no; 1 = yes; $\frac{1}{2}$ = somewhat possible]</p> <ul style="list-style-type: none"> • This is a somewhat subjective estimate of the relative possibility of making estimates of loading using existing monitoring data. • If the compound is measured at most or all IADN sites (and has some data water data available, if gas exchange may be important) then it is assigned a value of "1", i.e., it is possible to at least try to make an estimate of loadings to the Lakes • If the compound is not measured in the air or precipitation in the Great Lakes region, then it is assumed that the possibility of a loading estimate is "0" • A possibility of "$\frac{1}{2}$" suggests that for various reasons, the estimation feasibility lies somewhere between the two extremes above. • a "~" in front of a value indicates that air/water gas exchange may be an important pathway, but, that water measurements have not been made in all of the Great Lakes. For the lakes in which water measurements are available, rough estimates of loading can be made; for the lakes for which measurement are not available, very rough loading estimates could be made by using, for example, the average water concentrations measured in other lakes.

ADDITIONAL NOTES for Table 18 , relating to columns 2,4, and 14 only.	
II-a	For this analysis, several additional PAH's were considered, consisting of the remaining compounds in the EPA's 16-PAH list and the ATSDR 17-PAH list.
B	Tributyltin existence in significant amounts in the atmosphere is uncertain. If present in the atmosphere, it's overall long range transport rating might be on the order of "3".
C	Measurements of total toxaphene and major congeners are being made at Eagle Harbor and Pt. Petre in a collaborative research project involving Envr Canada & Indiana Univ.
F	Except for Lake Michigan, mercury has not been measured in the water of the Great Lakes; speciation info of Hg in air/water phases limited; thus, estimates are difficult to make. Total gaseous and particulate-phase Hg were measured at the five IADN master stations during 1995 and 1996 in a research program. This program stopped in December 1996. Sampling at Eagle Harbor is continuing, supported by a trust fund. Sampling at Pt. Petre is being conducted for total gaseous mercury, sponsored by Environment Canada. Sampling for total mercury in precipitation is being conducted at two IADN satellite sites (Brule River, Wisconsin and Dorset, Ontario) as part of the Mercury Deposition Network. The reliability of "particulate mercury" measurements reportedly being made by at least some of the programs is not clear. Thus, the rating of "½" for particulate mercury may be an overestimate.
G	Vapor- and particulate-phase PCDD/F is being measured in multi-site monitoring programs in Canada, sponsored by Environment Canada and the Ontario Ministry of Environment and Energy, and at several sites in Vermont. These programs were not designed specifically for estimating loading to the Great Lakes, but, there are several sites in the region.
H	Only water data for one Lake could be found; thus estimates could be made for that Lake. Crude estimates could possibly be made for other Great Lakes, using the water concentrations found in the one Lake, as a first approximation.

Table 19. Summary of Air and Water Monitoring Issues Identified for BVES Compounds in the Great Lakes Region		
Compound or Group	Air and Precipitation Monitoring Issues	Water Monitoring Issues
<ul style="list-style-type: none"> Alkylated Lead 4-Bromophenyl Phenyl Ether 3,3'-Dichlorobenzidene 4,4'-Methylene bis (2-chloroaniline) Tributyltin 	<ul style="list-style-type: none"> Not included in any of the air or precipitation monitoring programs identified 	<ul style="list-style-type: none"> Not included in any of the water monitoring programs identified
<ul style="list-style-type: none"> Pentachlorophenol Dinitropyrenes Perylene 	<ul style="list-style-type: none"> Limited air monitoring identified in Canada only No precipitation monitoring 	<ul style="list-style-type: none"> Not included in any of the water monitoring programs identified
<ul style="list-style-type: none"> PAH's in general 	<ul style="list-style-type: none"> Spatial representativeness issue: PAH's are emitted primarily in urban areas. 	<ul style="list-style-type: none"> No monitoring in Lake Huron in the last five years
<ul style="list-style-type: none"> PCDD/F (dioxins and furans) 	<ul style="list-style-type: none"> Limited number of Great Lakes monitoring stations in Canada only, near Lakes Erie and Ontario; No monitoring identified near Lakes Superior, Michigan, or Huron; Spatial representativeness: monitoring primarily in urban locations, although, e.g., air monitoring at Pt. Petre. Only one site (Dorset) for precipitation monitoring 	<ul style="list-style-type: none"> Monitoring by Envr. Canada for 2,3,7,8-TCDD in Lake Erie (1994, 1995) and Lake Superior (1996, 1997); Monitoring by Cook and Burkhard (US EPA) in Lake Michigan in 1994 No monitoring in Lake Huron or Lake Ontario in the last five years
<ul style="list-style-type: none"> Mercury 	<ul style="list-style-type: none"> Limited number of monitoring location; Little or no gas-phase speciation data being collected 	<ul style="list-style-type: none"> Systematic measurements only identified for Lake Michigan
<ul style="list-style-type: none"> Toxaphene 	<ul style="list-style-type: none"> Monitoring only at 2 sites (Eagle Harbor and Pt. Petre) No current measurements in precipitation could be identified 	<ul style="list-style-type: none"> No monitoring in Lake Huron or Lake Erie in the last five years
<ul style="list-style-type: none"> Aldrin Endrin Heptachlor & Heptachlor Epoxide Methoxychlor Mirex Octachlorostyrene 	<ul style="list-style-type: none"> Measured at some or all Canadian IADN stations, but not at U.S. sampling sites in the Great Lakes Region 	<ul style="list-style-type: none"> No monitoring in Lake Huron in the last five years
<ul style="list-style-type: none"> DDT/DDD/DDE 	<ul style="list-style-type: none"> Spatial representativeness: high concentrations in the air at South Haven — are there other hot spots in the Great Lakes region? 	
<ul style="list-style-type: none"> Hexachloro-1,3-butadiene 	<ul style="list-style-type: none"> Not part of IADN, but, measured in other programs in Can. & U.S. It may be possible to estimate loadings for many of the Lakes; No data near Lake Superior. 	<ul style="list-style-type: none"> No monitoring in Lake Huron in the last five years
<ul style="list-style-type: none"> 1,4-dichlorobenzene tetrachlorobenzenes pentachlorobenzenes 	<ul style="list-style-type: none"> Limited air measurements in the Great Lakes region 	<ul style="list-style-type: none"> For all, no monitoring in Lake Huron in the last five years For 1,4-DCB, none in Lk. Mich. either
<ul style="list-style-type: none"> PCB's 	<ul style="list-style-type: none"> Different sets of PCB's being monitored in different programs Since one or more lakes may be volatilizing PCB's, representativeness of shoreline monitoring stations is in question 	<ul style="list-style-type: none"> Different sets of PCB's being monitored in different programs

7. BIBLIOGRAPHY and REFERENCES

[note: not all of the following references are cited explicitly in the text]

- Achman, D.R., K.C. Hornbuckle, S. J. Eisenreich (1993). "Volatilization of Polychlorinated Biphenyls from Green Bay, Lake Michigan." Environmental Science and Technology **27**(1): 75-87.
- Arimoto, R. (1989). "Atmospheric Deposition of Chemical Contaminants to the Great Lakes." Journal of Great Lakes Research **15**(2): 339-356.
- Baker, J. E. , and S.J. Eisenreich (1990). "Concentrations and Fluxes of Polycyclic Aromatic Hydrocarbons and Polychlorinated Biphenyls across the Air-Water Interface of Lake Superior." Environmental Science and Technology **24**(3): 342-352.
- Bernier, J., P. Brousseau, et al. (1995). "Immunotoxicity Of Heavy-Metals In Relation to Great-Lakes." Environmental Health Perspectives **103**(S9): 23-34.
- Bidleman, T. F. and L. L. McConnell (1995). "A Review Of Field Experiments to Determine Air-Water Gas-Exchange Of Persistent Organic Pollutants." Science Of the Total Environment **159**(2-3): 101-117.
- Bidleman, T. F. (1988). "Atmospheric Processes - Wet and Dry Deposition Of Organic-Compounds Are Controlled By Their Vapor Particle Partitioning." Environmental Science & Technology **22**(4): 361-367.
- Bloxam, R. (1997). Modeling Mercury Atmospheric Transport, Chemistry and Deposition. Ontario Ministry of Environment and Energy. Personal Communication.
- Bloxam, R., S. Wong, P.K. Misra, E. Voldner, W. Schroeder, G. Petersen (1991). "Modeling the Long-Range Transport, Transformation, and Deposition of Mercury in a Comprehensive Eulerian Framework." Proceedings of the 8th International Conference on Heavy Metals in the Environment. CEP Consultants Ltd., Heavy Metals Secretariat, Edinburgh, U.K.
- Bullock, O.R., W.G. Benjey, and M.H. Keating (1997b). "The Modeling of Regional Scale Atmospheric Mercury Transport and Deposition Using RELMAP." Accepted for publication in Atmospheric Deposition to the Great Lakes and Coastal Waters, edited by Joel Baker, to be published by the Society of Environmental Toxicology and Chemistry (forthcoming).
- Burke, J., M. Hoyer, et al. (1995). "Wet Deposition Of Mercury and Ambient Mercury Concentrations At a Site In the Lake Champlain Basin." Water Air and Soil Pollution **80**(1-4): 353-362.
- Chan, C.H., G. Bruce, and B. Harrison (1994). "Wet Deposition of Organochlorine Pesticides and Polychlorinated Biphenyls to the Great Lakes." Journal of Great Lakes Research **20**(3):546-560.
- Clark, T.L. (1992a). "Atmospheric Deposition of Toxic Metals to Lake Michigan: Preliminary Annual Model Calculations." Presented at the Environmental Protection Agency/Air and Waste Management Association International Symposium of Measurement of Toxic and Related Air Pollutants, Durham, NC, May 1992 (EPA/600/A-92/239; NTIS PB93-120954/XAB).
- Clark, T.L. (1992b). "Model Calculations of the Annual Atmospheric Deposition of Toxic Metals to Lake Michigan." Presented at the Annual Meeting of the Air and Waste Management Association (85th), Kansas City, MO, June 23-27, 1992 (EPA/600/A-92/234; NTIS PB93-120905/XAB).

- Cleckner, L. B., E. S. Esseks, et al. (1995). "Mercury Concentrations In Two 'Great Waters'." Water Air and Soil Pollution **80**(1-4): 581-584.
- Cohen, M.D. (1997a). Review of I. The Capability of Specific Persistent Toxic Substances to be Subjected to Long Range Atmospheric Transport, and of II. The Status and Capabilities of Associated Emissions Inventories. Prepared for the International Joint Commission's International Air Quality Advisory Board. IJC, Windsor, Ontario.
- Cohen, M.D. (1997b). Modeling the Atmospheric Transport and Deposition of Persistent Toxic Substances to the Great Lakes. (Phase II of Transportation and Deposition of Persistent Toxic Substances to the Great Lakes Basin: Transport Modeling Element). Prepared for the International Joint Commission's International Air Quality Advisory Board. IJC, Windsor, Ontario.
- Cohen, M., B. Commoner, H. Eisl, P. Bartlett, A. Dickar, C. Hill, J. Quigley, J. Rosenthal (1995). Quantitative Estimation of the Entry of Dioxins, Furans and Hexachlorobenzene into the Great Lakes from Airborne and Waterborne Sources. Center for the Biology of Natural Systems (CBNS), Queens College, Flushing NY.
- Cook, P. (1997). Personal communication. U.S. EPA National Health and Environmental Effects Research Laboratory, Mid Continent Ecology Division, Duluth, Minnesota.
- Draxler, R. (1994). HY-SPLIT Deposition Module, NOAA Air Resources Laboratory, Silver Spring, MD.
- Environment Canada, U.S. Environmental Protection Agency, Ontario Ministry of Environment and Energy (1994). Integrated Atmospheric Deposition Network Quality Assurance Program Plan. Revised Version 0.2, February 28, 1994.
- Eisenreich, S. J., B. B. Looney, et al. (1981). "Airborne Organic Contaminants In the Great-Lakes Ecosystem." Environmental Science & Technology **15**(1): 30-38.
- Eisenreich, S. J. (1980). "Atmospheric Input Of Trace-Metals to Lake-Michigan." Water Air and Soil Pollution **13**(3): 287-301.
- Eisenreich, S. J., G. J. Hollod, et al. (1979). "Accumulation Of Polychlorinated Biphenyls (PCBs) In Surficial Lake-Superior Sediments - Atmospheric Deposition." Environmental Science & Technology **13**(5): 569-573.
- Eisenreich, S. J., A. W. Elzerman, et al. (1978). "Enrichment Of Micronutrients, Heavy-Metals, and Chlorinated Hydrocarbons In Wind-Generated Lake Foam." Environmental Science & Technology **12**(4): 413-417.
- Gatz, D. F., V. C. Bowersox, and J. Su (1989). "Lead and Cadmium Loadings to the Great Lakes From Precipitation." Journal Of Great Lakes Research **15**(2): 246-264.
- Gill, G.A. and K.W. Bruland (1990). "Mercury Speciation in Surface Freshwater Systems in California and Other Areas." Environmental Science and Technology **24**(9): 1392-1400.
- Graney, J. R., A. N. Halliday, et al. (1995). "Isotopic Record Of Lead Pollution In Lake-Sediments From the Northeastern United-States." Geochimica Et Cosmochimica Acta **59**(9): 1715-1728.
- Guthrie, P.D., C.A. Emery, M.P. Ligocki, and G.E. Mansell (1995). Phase I of the Great Waters Impacts Analysis Pilot Study. Final Report. Prepared by Systems Applications International, San Rafael CA, for Dan Axelrad, U.S. EPA (Contract No. 69-D2-0189; Work Assignment No. 2-34).

- Hermanson, M.H., C.L. Monosmith, and M.T. Donnelly-Kelleher (1997). Atmospheric Environment. In Press. (as cited by Monosmith *et al.*, 1996)
- Hicks, B.B., and R.M. Williams (1980). "Transfer and Deposition of Particles to Water Surfaces." Chapter 26 in Atmospheric Sulfur Deposition: Environmental Impact and Health Effects, edited by D.S. Shriner *et al.*, Ann Arbor Science Publishers, Ann Arbor Michigan, pp 237-244.
- Hoff, R. M., W. M. J. Strachan, *et al.* (1996). "Atmospheric Deposition Of Toxic-Chemicals to the Great-Lakes — a Review Of Data Through 1994." Atmospheric Environment **30**(20): 3505-3527.
- Hoff, R.M., T.F. Bidleman, and S.J. Eisenreich (1993). "Estimation of PCC Loadings from the Atmosphere to the Great Lakes." Chemosphere **27**(10): 2047-2055.
- Hoff, R. M., D. C. G. Muir, *et al.* (1992a). "Annual Cycle Of Polychlorinated-Biphenyls and Organohalogen Pesticides In Air In Southern Ontario .1. Air Concentration Data." Environmental Science & Technology **26**(2): 266-275.
- Hoff, R. M., D. C. G. Muir, *et al.* (1992b). "Annual Cycle Of Polychlorinated-Biphenyls and Organohalogen Pesticides In Air In Southern Ontario .2. Atmospheric Transport and Sources." Environmental Science & Technology **26**(2): 276-283.
- Holsen, T. M., K. E. Noll, *et al.* (1993). "Dry Deposition and Particle-Size Distributions Measured During the Lake-Michigan Urban Air Toxics Study." Environmental Science & Technology **27**(7): 1327-1333.
- Honrath, R.E., C.I Sweet, and C.J. Plouff (1997). "Surface Exchange and Transport Processes Governing Atmospheric PCB Levels over Lake Superior." Environmental Science & Technology **31**(3): 842-852.
- Hornbuckle, K.C., C. W. Sweet, R. F. Pearson, D.L. Swackhamer, and S.J. Eisenreich (1995). "Assessing Annual Water-Air Fluxes of Polychlorinated Biphenyls in Lake Michigan." Environmental Science and Technology **29**(4): 869-877.
- Hornbuckle, K.C., D.R. Achman, and S. J. Eisenreich (1993). "Over-Water and Over-Land Polychlorinated Biphenyls in Green Bay, Lake Michigan." Environmental Science and Technology **27**(1): 87-98.
- Hoyer, M., J. Burke, *et al.* (1995). "Atmospheric Sources, Transport and Deposition Of Mercury In Michigan - 2 Years Of Event Precipitation." Water Air and Soil Pollution **80**(1-4): 199-208.
- Jeremiason, J.D., K.C. Hornbuckle, and S.J. Eisenreich (1994). "PCBs in Lake Superior, 1978-1992: Decreases in Water Concentrations Reflect Loss by Volatilization." Environmental Science and Technology **28**(5): 903-914.
- Junge, C. E. (1977). Basic Considerations About Trace Constituents in the Atmosphere as Related to the Fate of Global Pollutants. Fate of Pollutants in the Air and Water Environments. I. H. Suffet. New York, Wiley: 7-22.
- Keeler, G. J. and N. Pirrone (1996). "Atmospheric Transport and Deposition Of Trace-Elements to Lake Erie From Urban Areas." Water Science and Technology **33**(4-5): 259-265.
- Keeler, G., G. Glinsorn, *et al.* (1995). "Particulate Mercury In the Atmosphere - Its Significance, Transport, Transformation and Sources." Water Air and Soil Pollution **80**(1-4): 159-168.

- Keeler, G.J. (1994). Lake Michigan Urban Air Toxics Study. EPA/600/R-94-191(NTIS: PB95-129102).
- L'Italien, S. (1993) Organic Contaminants in the Great Lakes 1986-1990, Environmental Quality Branch, Environment Canada.
- L'Italien, S. (1996a) Organic Contaminants in Lake Ontario 1992-1993, Environmental Conservation Branch, Environment Canada.
- L'Italien, S. (1996b) Organic Contaminants in Lake Erie 1994-1995, Environmental Conservation Branch, Environment Canada.
- L'Italien, S. (1997), (personal communication).
- Mackay, D., S. Paterson, and W.H. Schroeder (1986). "Model Describing the Rates of Transfer Processes of Organic Chemicals between Atmosphere and Water." Environmental Science and Technology **20**(8):810-816.
- Mason, R. P., and K. A. Sullivan (1997). "Mercury in Lake Michigan." Environmental Science and Technology **31**(3): 942-947.
- McConnell, L. L., W E. Cotham, and T.F. Bidleman (1993). "Gas Exchange of Hexachlorocyclohexane in the Great Lakes." Environmental Science and Technology **27**(7): 1304-1311.
- McQueen, J.T., R.A Valigura, and B.J.B. Stunder (1997). "Evaluation of Surface Flux and Turbulence Predictions Over the Chesapeake Bay from a Mesoscale Model." Paper prepared for the American Meteorological Society's 1997 annual conference. Personal communication, J.T. McQueen, NOAA Air Resources Laboratory, Silver Spring, Maryland.
- McQueen, J.T., R.R. Draxler, B.J.B. Stunder, and G.D. Rolph (1996b). An Overview of the Regional Atmospheric Modeling System (RAMS) as Applied at the NOAA Air Resources Laboratory. NOAA Air Resources Laboratory, Silver Spring, Maryland (in press) (as cited by McQueen et al., 1996a).
- McQueen, J.T., R.A. Valigura, and B.J.B. Stunder (1996c). "Evaluation of the Regional Atmospheric Modeling System for Estimating Nitric Acid Pollution Deposition onto the Chesapeake Bay. Submitted to Atmospheric Environment. (as cited by McQueen et al., 1996a).
- Monosmith, C. (1997). Personal Communication. Michigan Department of Natural Resources, Air Quality Division, Lansing Michigan.
- Monosmith, C.L., and M.H. Hermanson (1996). "Spatial and Temporal Trends of Atmospheric Organochlorine Vapors in the Central and Upper Great Lakes." Environmental Science and Technology **30**(12): 3464-3472.
- Nriagu, J. O., G. Lawson, et al. (1993). "A Protocol For Minimizing Contamination In the Analysis Of Trace-Metals In Great-Lakes Waters." Journal Of Great Lakes Research **19**(1): 175-182.
- Petersen, G., J. Munthe, R. Bloxam (1996). "Numerical Modeling of Regional Transport, Chemical Transformations and Deposition Fluxes of Airborne Mercury Species." in Global and Regional Mercury Cycles: Sources, Fluxes and Mass Balances, edited by W Baeyens et al. Kluwer Academic Publishers, The Netherlands. pp 191-217.
- Pirrone, N. and G. J. Keeler (1996a). "The Rouge River Watershed Pollution By Trace-Elements - Atmospheric Depositions and Emission Sources." Water Science and Technology **33**(4-5): 267-275.

- Pirrone, N. and G. J. Keeler (1996b). "A Preliminary Assessment Of the Urban Pollution In the Great-Lakes Region." Science Of the Total Environment **190**: 91-98.
- Pirrone, N., G. J. Keeler, et al. (1996). "Historical Trends Of Airborne Trace-Metals In Detroit From 1971 to 1992." Water Air and Soil Pollution **88**(1-2): 145-165.
- Pirrone, N., G. Glinsorn, et al. (1995). "Ambient Levels and Dry Deposition Fluxes Of Mercury to Lakes Huron, Erie and St Clair." Water Air and Soil Pollution **80**(1-4): 179-188.
- Pirrone, N., G. J. Keeler, et al. (1995a). "Dry Deposition Of Trace-Elements to Lake-Michigan - a Hybrid-Receptor Deposition Modeling Approach." Environmental Science & Technology **29**(8): 2112-2122.
- Pirrone, N., G. J. Keeler, et al. (1995b). "Dry Deposition Of Semivolatile Organic-Compounds to Lake-Michigan." Environmental Science & Technology **29**(8): 2123-2132.
- Pirrone, N., G. J. Keeler, et al. (1995c). "Trends Of Ambient Concentrations and Deposition Fluxes Of Particulate Trace-Metals In Detroit From 1982 to 1992." Science Of the Total Environment **162**(1): 43-61.
- Pirrone, N. and G. J. Keeler (1993). "Deposition Of Trace-Metals In Urban and Rural-Areas In the Lake-Michigan Basin." Water Science and Technology **28**(3-5): 261-270.
- Rea, A. W., G. J. Keeler, et al. (1996). "The Deposition Of Mercury In Throughfall and Litterfall In the Lake Champlain Watershed - a Short-Term Study." Atmospheric Environment **30**(19): 3257-3263.
- Ridal, J.J., B. Kerman, L. Durham, and M.E. Fox (1996). "Seasonality of Air-Water Fluxes of Hexachlorocyclohexanes in Lake Ontario." Environ. Sci. & Tech. **30**(3):852-858.
- SAI (1996). Draft Final User's Guide to the Regulatory Modeling System for Aerosols and Deposition (REMSAD). Prepared by Systems Applications International (SAI), Inc., San Rafael, California. SYSAPP-96\42. This report can be downloaded from the EPA's Technology Transfer Network (TTN) Bulletin Board.
- Scherbatskoy, T., A. Rea, et al. (1995). "Deposition and Watershed Processing Of Mercury In the Lake Champlain Basin." Abstracts Of Papers Of the American Chemical Society **210**(AUG): 77-GEOC.
- Schmidt, J.A. (1982). "Models of Particle Dry Deposition to a Lake Surface and Some Effects of Surface Microlayers." Journal of Great Lakes Research **8**(2):271-280.
- Schroeder, W.H. and D.A. Lane (1988). "The Fate of Toxic Airborne Pollutants." Environmental Science and Technology **22**(3):240-246.
- Shannon, J. D. and E. C. Voldner (1995). "Modeling Atmospheric Concentrations Of Mercury and Deposition to the Great-Lakes." Atmospheric Environment **29**(14): 1649-1661.
- Shannon, J. D. and E. C. Voldner (1982). "Estimation Of Wet and Dry Deposition Of Pollutant Sulfur In Eastern Canada As a Function Of Major Source Regions." Water Air and Soil Pollution **18**(1-3): 101-104.
- Shannon, J.D. (1981). "A Model of Regional Long-Term Average Sulfur Atmospheric Pollution, Surface Removal, and Net Horizontal Flux." Atmospheric Environment **15**: 689-701.

- Slinn, W.G.N., L. Hasse, B.B. Hicks, A.W. Hogan, D. Lal, P.S. Liss, K.O. Munnich, G.A. Sehmel, and O. Vittori (1978). "Some Aspects of the Transfer of Atmospheric Trace Constituents Past the Air Sea Interface." Atmospheric Environment **12**: 2055-2087.
- Slinn, S.A., and W.G.N. Slinn (1980). "Predictions for Particle Deposition on Natural Waters." Atmospheric Environment **14**:1013-1016.
- Stevens, R. J. J. and Nielson, M. A. (1989). "Inter- and Intralake Distributions of Trace Organic Contaminants in Surface Waters of the Great Lakes." Journal of Great Lakes Research, **15** (3): 377-393.
- SOLEC (1994). State of the Lakes Ecosystem Conference: Toxic Contaminants, October 1994. Environment Canada, USEPA.
- Strachan, W., and S. Eisenreich (1988). Mass Balancing of Toxic Chemicals in the Great Lakes: The Role of Atmospheric Deposition. Appendix I from the Workshop on the Estimation of Atmospheric Loadings of Toxic Chemicals to the Great Lakes Basin (International Joint Commission, Windsor, Ontario).
- Sweet, C.W. (1996). "How Well do Measurements from the IADN Network Reflect Atmospheric Deposition to the Great Lakes?" Presentation at the Air and Waste Management Association's Conference on Atmospheric Deposition to the Great Waters, Niagara Falls, NY, October 28-30, 1996. To be published in the Conference Proceedings by the AWMA. A pre-publication manuscript was kindly provided as a personal communication from Clyde W. Sweet, Illinois State Water Survey.
- Sweet, C.W., D.F. Gatz, R.M. Hoff (1995). "Measuring Toxic Atmospheric Deposition to the Great Lakes: The Integrated Atmospheric Deposition Network (IADN)." Manuscript kindly provided as a personal communication from Clyde W. Sweet, Illinois State Water Survey.
- Telliard, W.A., H.B. McCarty, and L.S. Riddick (1993). "Results of the Interlaboratory Validation Study of U.S. EPA Method 1613 for the Analysis of Tetra- Through Octachlorinated Dioxins and Furans by Isotope Dilution GC/MS." Chemosphere 27(1-3):41-46.
- U.S. EPA (1996). Mercury Study Report to Congress. Volume III: An Assessment of Exposure from Anthropogenic Mercury Emissions in the United States. SAB Review Draft. Office of Air Quality Planning and Standards and Office of Research and Development. EPA-452/R-96-001c.
- US EPA (1995). Lake Michigan Mass Budget/Mass Balance Work Plan. Great Lakes National Program Office. [Note: document was undated; we are unsure of the publication date.]
- U.S. EPA (1994). Estimating Exposure to Dioxin-Like Compounds. Vol. I: Executive Summary. Vol. II: Properties, Sources, Occurrence and Background Exposures. Vol. III: Site-Specific Assessment Procedures. External Review Draft. "Not for Citation or Quotation." EPA/600/6-88-005Ca,b & c, June (Washington, DC: Office of Research & Development).
- US EPA Region II (1994). Lake Ontario Quality Assurance Project Plan.
- Valigura, R.A., W.T. Luke, R.S. Artz, B.B. Hicks (1996). Atmospheric Nutrient Input to Coast Areas: Reducing the Uncertainties. NOAA Coastal Ocean Program. Decision Analysis Series No. 9. NOAA Coastal Ocean Office, Silver Spring MD.
- Voldner, E.C., J.D. Shannon, D. Mackay (1993). "Modeled Trends in Atmospheric Deposition of Lead to the Great Lakes and Resulting Lake Mass Balance." Presented at the 9th International Conference on Heavy Metals in the Environment. Graphics from presentation kindly provided as

personal communication from Jack Shannon, Argonne National Laboratory. Abstract in the Proceedings of the 9th International Conference on Heavy Metals in the Environment, p 275. CEP Consultants Ltd., Heavy Metals Secretariat, Edinburgh, U.K.

Voldner, E.C., and M. Alvo (1993). "Estimation of Wet Deposition of Sulfur, Nitrogen, Cadmium and Lead to the Great Lakes." Environmental Science and Technology **27**(2):292-298.

Voldner, E. C. and W. H. Schroeder (1989). "Modeling Of Atmospheric Transport and Deposition Of Toxaphene Into the Great-Lakes Ecosystem." Atmospheric Environment **23**(9):1949-1961.

Wesely, M. (1989). "Parameterization of Surface Resistances to Gaseous Dry Deposition in Regional-Scale Numerical Models." Atmospheric Environment **23**(6):1293-1304.

Williams, R.M. (1982). "A Model for the Dry Deposition of Particles to Natural Water Surfaces." Atmospheric Environment **16**(8):1933-1938.