



Food and Agriculture  
Organization of the  
United Nations

VOLUME 5

# RECARBONIZING GLOBAL SOILS

PRACTICES  
OVERVIEW

A technical manual  
of recommended  
management  
practices



FORESTRY,  
WETLANDS,  
URBAN SOILS





**VOLUME 5**

# **RECARBONIZING GLOBAL SOILS**

**PRACTICES  
OVERVIEW**

**A technical manual  
of recommended  
management  
practices**

An illustration of soil and roots. The top part shows a layer of dark brown soil with green grass blades growing from it. Below the soil, a network of light brown roots is visible, extending across the width of the page. The background is a light, textured grey.

**FORESTRY,  
WETLANDS,  
URBAN SOILS**

**Food and Agriculture Organization of the United Nations  
Rome, 2021**

## Required citation:

**FAO and ITPS.** 2021. *Recarbonizing global soils: A technical manual of recommended management practices. Volume 5: Forestry, Wetlands and Urban Soils- Practices overview.* Rome, FAO. <https://doi.org/10.4060/cb6606en>

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO or ITPS in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO or ITPS.

ISBN 978-92-5-134900-7

© FAO, 2021



Some rights reserved. This work is made available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo/legalcode/legalcode>).

Under the terms of this licence, this work may be copied, redistributed and adapted for non-commercial purposes, provided that the work is appropriately cited. In any use of this work, there should be no suggestion that FAO endorses any specific organization, products or services. The use of the FAO logo is not permitted. If the work is adapted, then it must be licensed under the same or equivalent Creative Commons licence. If a translation of this work is created, it must include the following disclaimer along with the required citation: “This translation was not created by the Food and Agriculture Organization of the United Nations (FAO). FAO is not responsible for the content or accuracy of this translation. The original [Language] edition shall be the authoritative edition.”

Disputes arising under the licence that cannot be settled amicably will be resolved by mediation and arbitration as described in Article 8 of the licence except as otherwise provided herein. The applicable mediation rules will be the mediation rules of the World Intellectual Property Organization <http://www.wipo.int/amc/en/mediation/rules> and any arbitration will be conducted in accordance with the Arbitration Rules of the United Nations Commission on International Trade Law (UNCITRAL).

**Third-party materials.** Users wishing to reuse material from this work that is attributed to a third party, such as tables, figures or images, are responsible for determining whether permission is needed for that reuse and for obtaining permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

**Sales, rights and licensing.** FAO information products are available on the FAO website ([www.fao.org/publications](http://www.fao.org/publications)) and can be purchased through [publications-sales@fao.org](mailto:publications-sales@fao.org). Requests for commercial use should be submitted via: [www.fao.org/contact-us/licence-request](http://www.fao.org/contact-us/licence-request). Queries regarding rights and licensing should be submitted to: [copyright@fao.org](mailto:copyright@fao.org)

# Contents

Managed forests and silviculture	1
Soil disturbance	2
1. Harvest systems that limit soil disturbance and reduced impact logging	2
Soil cover	13
2. Continuous cover forestry and extended rotations	13
3. Residue retention	22
Nutrient management	33
4. Inclusion of N fixing species	33
5. Forest fertilization	51
Forest restoration	66
6. Forest afforestation, reforestation and natural regeneration	66
7. Rehabilitation of forest soils affected by wildfires	88
8. Forest landscape restoration	107
Wetlands	122
Wetland management	123
9. Avoiding conversion and conservation of wetlands	123
10. Wetland restoration (water supplementation and promoting plant growth)	147
Critical wetland ecosystems	161
Peatlands	161
11. Conservation of pristine peatlands and avoiding drainage of peatlands	161
12. Restoration of peatlands	169
13. Paludiculture	179

Mangroves and organic forest soils	190
14. Restoration of mangrove forest	190
15. Restoration of organic coastal and inland freshwater forests	199
Rice paddies	212
16. Water level management in rice paddies	212
17. Straw residue management	229
18. Selection of rice varieties adapted to salinity	246
19. Integrated rice-based farming systems	252
<b>Urban soils and infrastructures</b>	<b>279</b>
Urban infrastructures	280
20. Management of gardens, parks, and lawns	280
21. Bioretention systems	294
22. Green roofs	304
Urban agriculture	313
23. Urban agriculture	313
Urban forestry	328
24. Urban forestry	328

# Tables

<b>Table 1.</b> Soils threats	4
<b>Table 2.</b> Potential barriers to adoption	6
<b>Table 3.</b> Soil threats	15
<b>Table 4.</b> Potential barriers to adoption	17
<b>Table 5.</b> Changes in soil organic carbon stocks reported for harvest residue retention	23
<b>Table 6.</b> Soil threats	24
<b>Table 7.</b> Potential barriers to adoption	27
<b>Table 8.</b> Changes in soil organic carbon stocks reported for addition of N-fixing tree species	35
<b>Table 9.</b> Soil threats	37
<b>Table 10.</b> Soil threats	39
<b>Table 11.</b> Potential barriers to adoption	42
<b>Table 12.</b> Related case studies available in volumes 4 and 6	43
<b>Table 13.</b> Changes in soil organic carbon stocks reported for nutrient additions	53
<b>Table 14.</b> Soil threats	54
<b>Table 15.</b> Soil threats	55
<b>Table 16.</b> Potential barriers to adoption	59
<b>Table 17.</b> Changes in soil organic carbon stocks reported following afforestation	69
<b>Table 18.</b> Soil threats	72
<b>Table 19.</b> Soil threats	75
<b>Table 20.</b> Potential barriers to adoption	77
<b>Table 21.</b> Related case studies available in volumes 4 and 6	79
<b>Table 22.</b> Changes in soil organic carbon stocks reported for soils affected by forest fires after mulching treatments in different environments	91
<b>Table 23.</b> Results of studies about the short-term effects of post-fire straw mulching on soil physical, physicochemical, chemical, biochemical and microbial properties, analyzed in different biomes and climate conditions	94
<b>Table 24.</b> Soil threats	97
<b>Table 25.</b> Potential barriers to adoption	100
<b>Table 26.</b> Related case studies available in volumes 4 and 6	102
<b>Table 27.</b> Example of forest landscape restoration (FLR) activities across the landscape	108
<b>Table 28.</b> Priority FLR areas to improve soil aspects of landscape functionality across climatic regions and geographies	109
<b>Table 29.</b> Changes in soil organic carbon stocks reported for forest landscape restoration projects	110

<b>Table 30.</b> Soil threats	111
<b>Table 31.</b> Potential barriers to adoption	114
<b>Table 32.</b> Changes in soil organic carbon stocks reported for coastal and freshwater conserved wetlands across the world	126
<b>Table 33.</b> Soil threats	130
<b>Table 34.</b> Potential barriers to adoption	134
<b>Table 35.</b> Related case studies available in volumes 4 and 6	137
<b>Table 36.</b> Changes in soil organic carbon stocks reported for wetland restoration	149
<b>Table 37.</b> Soil threats	150
<b>Table 38.</b> Soil threats	152
<b>Table 39.</b> Potential barriers to adoption	154
<b>Table 40.</b> Related case studies available in volumes 4 and 6	156
<b>Table 41.</b> Soil threats	163
<b>Table 42.</b> Potential barriers to adoption	166
<b>Table 43.</b> Soil threats	171
<b>Table 44.</b> Potential barriers to adoption	175
<b>Table 45.</b> Related case studies available in volumes 4 and 6	176
<b>Table 46.</b> Soil threats	181
<b>Table 47.</b> Potential barriers to adoption	184
<b>Table 48.</b> Related case studies available in volumes 4 and 6	186
<b>Table 49.</b> Changes in soil organic carbon stocks reported for restoration of mangrove forests	191
<b>Table 50.</b> Soil threats	192
<b>Table 51.</b> Soil threats	193
<b>Table 52.</b> Potential barriers to adoption	194
<b>Table 53.</b> Related case studies available in volumes 4 and 6	195
<b>Table 54.</b> Potential avoided emissions and changes in soil organic carbon stocks reported for restored degraded peatlands during the duration of study or model run	201
<b>Table 55.</b> Soil threats	203
<b>Table 56.</b> Soil threats	204
<b>Table 57.</b> Potential barriers to adoption	206
<b>Table 58.</b> SOC stocks and changes in soil organic carbon stocks reported for water level management	215
<b>Table 59.</b> Soil threats	217
<b>Table 60.</b> GHG emissions and global warming potential (GWP) according to water level management CW vs AWD or Sprinkler	219
<b>Table 61.</b> Soil threats	221



<b>Table 62.</b> Potential barriers to adoption	223
<b>Table 63.</b> Related case studies available in volumes 4 and 6	225
<b>Table 64.</b> Changes in soil organic carbon stocks reported for straw residue management	232
<b>Table 65.</b> Soil threats	234
<b>Table 66.</b> Soil threats	237
<b>Table 67.</b> GHG emissions and GWP according to residue type incorporated	238
<b>Table 68.</b> Potential barriers to adoption	242
<b>Table 69.</b> Related case studies available in volumes 4 and 6	242
<b>Table 70.</b> Potential barriers to adoption	249
<b>Table 71.</b> Rice based integrated farming system prevalent in different part of the world	259
<b>Table 72.</b> Changes in soil organic carbon stocks reported for rice based integrated farming systems	261
<b>Table 73.</b> Soil threats	263
<b>Table 74.</b> Different climate change adaptation and mitigation options in rice-fish based integrated farming systems	266
<b>Table 75.</b> Soil threats	269
<b>Table 76.</b> Potential barriers to adoption	272
<b>Table 77.</b> Changes in soil organic carbon stocks reported for management of parks, gardens and lawns	282
<b>Table 78.</b> Soil threats	283
<b>Table 79.</b> Soil threats	285
<b>Table 80.</b> Potential barriers to adoption	288
<b>Table 81.</b> Related case studies available in volumes 4 and 6	291
<b>Table 82.</b> Changes in soil organic carbon stocks reported for bioretention systems	296
<b>Table 83.</b> Soil threats	297
<b>Table 84.</b> Soil threats	298
<b>Table 85.</b> Potential barriers to adoption	300
<b>Table 86.</b> Changes in soil organic carbon stocks reported for green roofs	306
<b>Table 87.</b> Soil threats	307
<b>Table 88.</b> Soil threats	308
<b>Table 89.</b> Potential barriers to adoption	310
<b>Table 90.</b> Related case studies available in volumes 4 and 6	310
<b>Table 91.</b> Changes in soil organic carbon stocks reported for two urban agriculture trials	315
<b>Table 92.</b> Soil threats	316
<b>Table 93.</b> Soil threats	318

<b>Table 94.</b> Potential barriers to adoption	321
<b>Table 95.</b> Related case studies available in volumes 4 and 6	324
<b>Table 96.</b> Changes in soil organic carbon stocks reported for urban forests from diverse regions	330
<b>Table 97.</b> Soil threats	331
<b>Table 98.</b> Soil threats	333
<b>Table 99.</b> Potential barriers to adoption	335
<b>Table 100.</b> Related case studies available in volumes 4 and 6	337

# Figures

<b>Figure 1.</b> Digital elevation model (LIDAR based) showing the soil impact of logging operations by a 23 t forwarder compared to logging operations using a cable yarding system	8
<b>Figure 2.</b> The potential mechanisms that regulate the responses of CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O production and consumption to elevated N	57
<b>Figure 3.</b> Change in SOC stocks after afforestation a) in different climatic zones and b) with different types of tree species	68
<b>Figure 4.</b> Temporal context of the rehabilitation and restoration strategies for the recovery of the burnt forest ecosystems	90
<b>Figure 5.</b> Global forest landscape restoration opportunities	109
<b>Figure 6.</b> Effects of greenhouse gas emissions from peatlands following global rewetting scenarios, and related radiative forcing (RF) and global warming effect	173
<b>Figure 7.</b> Relationship between global warming potential of greenhouse gas emissions, mean annual ground water table and land-use practices for temperate peatlands	180
<b>Figure 8.</b> Pathways contributing to resource flows (i.e. from one enterprise to other) portraying their synergy and mutualisms conducive to enhance family foods and income	253
<b>Figure 9.</b> Multitier rice based integrated farming system	255
<b>Figure 10.</b> Symbiotic relationship between rice-fish-duck yielding maximum mutual benefits to all the entities	256
<b>Figure 11.</b> Framework of potential mutualism and synergies among rice-fish-duck-Azolla based co-culture	256
<b>Figure 12.</b> Pictorial views of operational periodicities of rice, fish and ducks in rice-rice cropping system	257
<b>Figure 13.</b> Schematic view of CH <sub>4</sub> and N <sub>2</sub> O emissions from rice-based integrated farming systems and synergistic effects of duck and fish contributions towards reduction of global warming potentials	267
<b>Figure 14.</b> Schematic view of a bioretention system	295
<b>Figure 15.</b> Schematic view of green roof composition	304

# Photos

<b>Photo 1.</b> Soil disturbance (rut formation and soil compaction) from conventional skidder logging in Austria (top left) and France (top right), cable yarding operation in the Austrian Alps (bottom)	7
<b>Photo 2.</b> A mature stand of Douglas fir managed on CCF principles with a developing understorey of mixed conifers, including Douglas fir, western hemlock, western red cedar and grand fir, Coombs Wood, Cumbria, United Kingdom of Great Britain and Northern Ireland	18
<b>Photo 3.</b> Eucalypt plantation stem-only harvest	28
<b>Photo 4.</b> Acacia mangium planted near Bintulu, in Sarawak, Borneo (Malaysia)	43
<b>Photo 5.</b> Successful afforestation involving a mix of species in a lowland rain forest near Rio de Janeiro, Brazil	78
<b>Photo 6.</b> Helicopter straw mulching application after wildfire in Galicia (NW Spain)	101
<b>Photo 7.</b> Forest natural regeneration. California, United States of America	116
<b>Photo 8.</b> Mosaic landscape. Guilin, China	116
<b>Photo 9.</b> Mangrove restoration. Quang Ninh, Vietnam	117
<b>Photo 10.</b> Multistrata agroforestry, shaded coffee. Quindío, Colombia	117
<b>Photo 11.</b> Converted wetlands – drained wetlands for grazing pasture in Queensland, Australia (left) and degraded mangrove during conversion to aquaculture in Myanmar (right)	135
<b>Photo 12.</b> Conservation of tidal wetlands – mangroves ( <i>Ceriops</i> sp., left) and tidal marsh ( <i>Sarcocornia</i> sp., right) in Queensland, Australia	136
<b>Photo 13.</b> Conservation of freshwater wetland ( <i>Juncus</i> sp. and <i>Melaleuca quinquenervia</i> ) in Queensland, Australia	137
<b>Photo 14.</b> Wetland restoration project in Qixinghe National Natural Reserve, Sanjiang Plain, China	155
<b>Photo 15.</b> Wetland restoration project in Niuxintaobao National Wetland Park, Songnen Plain, China	156
<b>Photo 16.</b> Peatland Restoration and reforestation site in Sumatra Kayuagung, South Sumatra	186
<b>Photo 17.</b> Regeneration of vegetation in drained organic forest can occur after short-term farming from seeds in the seed bank, dispersal or replanting (Cache River, Southern Illinois)	207
<b>Photo 18.</b> Removing water in AWD (Valencia-Spain)	224
<b>Photo 19.</b> Flooding conditions in rice fields (Albufera of Valencia - Spain)	224
<b>Photo 20.</b> Dried fields in Mediterranean wetland (Abufer of Valencia - Spain)	225
<b>Photo 21.</b> Piezometer	225
<b>Photo 22.</b> Depicting an improved version of rice-aquaculture system including a livestock component	254

<b>Photo 23.</b> Duck foraging in dry seeded rice fields creating a conducive environment for initial (after 20 days of rice plant establishment) control of weeds and pests, duck dropping for fertilization and loosening of soil for better root growth and subsequent growth of rice plants (left). Rice-fish-duck co-culture system with luxuriant growth of rice (right)	257
<b>Photo 24.</b> Pond-dyke farming system	258
<b>Photo 25.</b> A flower garden in Chicago, Illinois, United States of America	289
<b>Photo 26.</b> Echo Urban Park in Los Angeles, California, United States of America	289
<b>Photo 27.</b> An urban park in Houston, Texas, United States of America	290
<b>Photo 28.</b> Two urban parks in San Francisco, California, United States of America	291
<b>Photo 29.</b> Urban-agriculture community gardens, Commerce, Texas, United States of America	322
<b>Photo 30.</b> Rooftop garden, Chicago Botanical Gardens, Chicago, Illinois, United States of America	322
<b>Photo 31.</b> Urban community garden, Los Angeles, California, United States of America	United States of America 323
<b>Photo 32.</b> Community garden, West Hollywood, California, United States of America	United States of America 323
<b>Photo 33.</b> Community garden, Governors Island, New York City, New York, United States of America	United States of America 324
<b>Photo 34.</b> Tree growing in a containerized bunker in New York City. Beneath the pavement is a subway line	335
<b>Photo 35.</b> Trees surrounded by pervious brick pavement	336
<b>Photo 36.</b> Trees lining an urban park in New York City (Central Park)	336
<b>Photo 37.</b> Urban woodlots	337





# Managed forests and silviculture

# 1. Harvest systems that limit soil disturbance and reduced impact logging

Mathias Mayer<sup>1,2</sup>, Noémie Pousse<sup>3</sup>, Jason James<sup>4</sup>

<sup>1</sup>Forest Soils and Biogeochemistry, Swiss Federal Institute for Forest, Snow and Landscape Research (WSL), Birmensdorf, Switzerland

<sup>2</sup>Institute of Forest Ecology, Department of Forest and Soil Sciences, University of Natural Resources and Life Sciences (BOKU), Vienna, Austria

<sup>3</sup>ONF (Office National des Forêts) - Research, Development, and Innovation Department, Avignon, France

<sup>4</sup>Exponent, Inc., Bellevue, Washington, United States of America

## 1. Description of the practice

Conventional harvesting operations in forests are inevitably associated with the disturbance of the forest soil. The use of ground-based heavy machinery, including skidders, forwarders, or tractors, can severely damage forest soil (Photo 1). Machinery movement through the forest can cause soil mixing, compaction, and erosion with long-lasting consequences for biological, chemical, and physical soil properties (Horn *et al.*, 2007; Cambi *et al.*, 2015). On steep terrain, rut formation can result in drastic losses of humus and mineral soil (Labrière *et al.*, 2015; Naghdi *et al.*, 2016a). On flat terrain, soil compaction and rutting can lead to a decrease in soil aeration and water infiltration (Goutal, Renault and Ranger, 2013; Bonnaud *et al.*, 2019). Soil disturbance due to logging can be limited if alternative harvesting systems are applied, such as, for example, helicopter-, cable yard-, or animal logging (Bockheim, Ballard and Wellington, 1975; Miller and Sirois, 1986; Aust and Lea, 1992; Naghdi *et al.*, 2009). Adverse effects on soil properties may be reduced by changes in machinery configuration as well, such as reducing tire pressure, installing larger diameter tires, and increasing the number of axles (Solgi *et al.*, 2020). Soil disturbance can also be reduced if networks of logging and skidding roads are properly planned prior to operations (e.g. adequate density) and are planned permanently (e.g. for future harvests) (Ampoorter *et al.*, 2012; Picchio *et al.*, 2020), logging is conducted during seasons with frozen and/or dry soil conditions (Ampoorter *et al.*, 2012; Naghdi *et al.*, 2016b), or slash is placed over skid trails as a means to distribute the load of machinery (Agherkakli *et al.*, 2014). In tropical forests, improvements for forest harvesting operations have been proposed under the term “reduced impact logging” (RIL), which includes measures to reduce environmental impacts of harvesting operations (Putz and Pinard, 1993; Healey, Price and Tay, 2000; Putz *et al.*, 2008).



## 2. Range of applicability

Harvesting systems to limit soil disturbance and RIL can be applied globally. Most countries around the world are choosing a combination of spatial (permanent skidding trails or extraction tracks) and temporal (frozen or dry soils) limitations of forest traffic to permanently limit the surface disturbed. Alternative harvesting systems are rarely chosen due to higher costs.

## 3. Impact on soil organic carbon stocks

Most peer-reviewed literature and meta-analyses have focused on the effects of intensive harvesting practices such as whole tree harvest and intensive biomass removal on soil organic carbon (SOC). The effects of alternative harvest systems and RIL systems on SOC losses and -stocks have only rarely been investigated and results are still inconclusive. For a dipterocarp forest in Sabah, Malaysia, RIL reduced SOC loss after logging by 4 tC/ha when compared to conventional logging (Putz and Pinard, 1993). In contrast, no differences in SOC stocks could be shown for conventional logging and RIL in a rain forest in Southern Cameroon (Tchiofo Lontsi *et al.*, 2019). Likewise, no differences in soil organic matter content were found for helicopter and skidder logged stands in Alabama, United States of America (Aust and Lea, 1991). Other studies compared harvesting systems regarding on-site soil surface damage and soil erosion losses. Particularly soil erosion can be associated with local SOC loss (translocation) from the harvest units (Berhe *et al.*, 2018). In a pine forest of Mississippi, United States of America, for instance, Miller and Sirois (1986) report that cable yard logging damaged 16 percent of the soil surface, while ground skidding damaged 31 percent. In boreal forest stands of British Columbia, Canada, tractor logging resulted in 71 percent exposed mineral soil (i.e. organic layer displaced), while helicopter logging resulted only in 5 percent exposed mineral soil (Bockheim, Ballard and Wellington, 1975); highest percentages of exposed mineral soil were determined for steep slopes and shale-rich soils. In a pine forest of Honduras with slopes greater than 30 percent, cable yard- and animal logging led to six to ten times less erosion than ground-based harvesting (tractor) during the rainy season (Rivera, Kershner and Dobrowolski, 2010). In a tropical forest of Malaysia, 17 percent of the area of a logged stand was covered by roads and skid trails, while under RIL guidelines only 6 percent of the area was similarly disturbed (Pinard, Barker and Tay, 2000); additionally, skid trails with disturbance of subsoil were less than 50 percent compared to regularly logged stands. Similarly, RIL resulted in 4 to 6 percent lower soil disturbance than conventional logging in a Brazilian rain forest (Pereira *et al.*, 2002). Worrell, Bolding and Aust (2011) report higher potential soil erosion rates (0.5 t/ha/yr) in steep Appalachian hardwood forests when comparing conventional skidding to cable yarding operations. In Hyrcanian forests of Iran, skidding on steep slopes (> 20 percent) resulted in higher soil disturbance and forest floor mass loss than skidding on less steep slopes (Naghdi *et al.*, 2016a).

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Conventional harvesting systems using ground-based machinery increase soil bulk density, soil CO<sub>2</sub> concentration, waterlogging and runoff, and decrease soil porosity, water infiltration and permeability, air permeability, oxygen supply, and root and tree growth (Cambi *et al.*, 2015). Compared to conventional systems, alternative harvesting systems have been shown to cause smaller changes in chemical and physical soil properties, including saturated hydraulic conductivity, acidity, oxygen concentration, redox potential (Aust and Lea, 1992). More specifically, forwarder movement was shown to result in water logging and to decrease soil aeration in comparison to cable yarding systems (Goutal, Renault and Ranger, 2013; as a consequence, soil CO<sub>2</sub> efflux (Goutal *et al.*, 2012b), CH<sub>4</sub> absorption capacity (Epron *et al.*, 2016), earthworms abundance and diversity (Bottinelli, Capowiez and Ranger, 2014), tree regeneration, growth and rooting depth (Goutal-Pousse, Boc and Ranger, 2014) were reduced.

### 4.2 Minimization of threats to soil functions

Table 1. Soils threats

Soil threats	
<b>Soil erosion</b>	In general, harvesting systems that reduce soil disturbance may mitigate soil erosion. For example, cable yarding reduced potential soil erosion by 0.5 t/ha/yr (Worrell, Bolding and Aust, 2011) or by a factor 6 to 10 (Rivera, Kershner and Dobrowolski, 2010).
<b>Nutrient imbalance and cycles</b>	In general, harvesting systems that reduce soil erosion are beneficial for nutrient balance and cycles. For example, reduced soil erosion rates due to cable yarding were shown to also maintain soil nutrient stocks (Worrell, Bolding and Aust, 2011). Reduced waterlogging maintain soil nutrients cycling (biological activity, rooting intensity and roots activity, redox potential) (Bonnaud <i>et al.</i> , 2019).
<b>Soil biodiversity loss</b>	Conventional harvesting systems have been shown to reduce and alter abundance of the microbiota, decrease macrofauna diversity, and increase bacterial diversity (Bottinelli, Capowiez and Ranger, 2014; Hartmann <i>et al.</i> , 2014).
<b>Soil compaction</b>	Cable yarding system and RIL can decrease compaction (Goutal <i>et al.</i> , 2012a; Tchifo Lontsi <i>et al.</i> , 2019). Animal logging has been shown to decrease compaction and rutting (Horn <i>et al.</i> , 2007). The extent of compaction within a stand during both conventional and alternative harvesting systems may be reduced by operating during dry or frozen conditions, limiting the number of passes, limiting activity to skid trails, and proper training and supervision of the operator (Piccio <i>et al.</i> 2020).



## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

There is no available information on tradeoffs with other soil threats.

### 5.2 Increases in greenhouse gas emissions

For tropical forests, Ellis *et al.* (2019) estimated that selective logging emitted 834 Tg CO<sub>2</sub> in 2015, representing 6 percent of total tropical GHG emissions. A full implementation of RIL measures, including reduced wood waste, narrower haul roads, and lower impact skidding equipment, would reduce emissions from logging by 366 Tg CO<sub>2</sub>/yr (= 44 percent reduction). Based on eddy covariance measurements in tropical rainforest in Brazil, Miller *et al.* (2011) showed RIL to have only minimal effects on total CO<sub>2</sub> efflux to the atmosphere.

## 6. Recommendations before implementation of the practice

No information available

## 7. Potential barriers to adoption

Table 2. Potential barriers to adoption


Barrier	YES/NO	
Social	Yes	In some countries, limited number of cable yarding companies (Magaud, 2020) constrains wider application of this practice. Moreover, many alternative harvesting systems are applicable only under specific circumstances.
Economic	Yes	RIL (Boltz, Holmes and Carter, 2003) and alternative harvesting systems such as cable yarding and animal skidding (Schweier and Ludowicy, 2020) can be less profitable than conventional ground-based harvesting .

Barrier	YES/NO	
Knowledge	Yes	Training and knowledge of individual contractors and operators is imperative to reduce forest damage (Picchio, Mederski and Tavankar, 2020), for proper implementation of RIL (Putz <i>et al.</i> , 2008), and for reducing soil disturbance during conventional cable yarding or conventional harvest (Chase <i>et al.</i> , 2019).

## Photos of the practice



**Photo 1.** Soil disturbance (rut formation and soil compaction) from conventional skidder logging in Austria (top left) and France (top right). Cable yarding operation in the Austrian Alps (bottom)



**Figure 1.** Digital elevation model (LIDAR based) showing the soil impact of logging operations by a 23 t forwarder compared to logging operations using a cable yarding system

Colors yellow and green indicate low and high elevations, respectively. Figure modified after Goutal-Pousse, Boc and Ranger (2014)

## References

- Agherkakli, B., Najafi, A., Sadeghi, S.H. & Zenner, E.** 2014. Mitigating effects of slash on soil disturbance in ground-based skidding operations. *Scandinavian Journal of Forest Research*, 29(5): 499–505. <https://doi.org/10.1080/02827581.2014.919351>
- Ampoorter, E., De Schrijver, A., Van Nevel, L., Hermy, M. & Verheyen, K.** 2012. Impact of mechanized harvesting on compaction of sandy and clayey forest soils: results of a meta-analysis. *Annals of Forest Science*, 69(5): 533-542. <https://doi.org/10.1007/s13595-012-0199-y>
- Aust, W.M. & Lea, R.** 1991. Soil Temperature and Organic Matter in a Disturbed Forested Wetland. *Soil Science Society of America Journal*, 55(6): 1741-1746. <https://doi.org/10.2136/sssaj1991.03615995005500060039x>
- Aust, W.M. & Lea, R.** 1992. Comparative effects of aerial and ground logging on soil properties in a tupelo-cypress wetland. *Forest Ecology and Management*, 50(1-2): 57-73. [https://doi.org/10.1016/0378-1127\(92\)90314-Y](https://doi.org/10.1016/0378-1127(92)90314-Y)
- Berhe, A.A., Barnes, R.T., Six, J. & Marín-Spiotta, E.** 2018. Role of soil erosion in biogeochemical cycling of essential elements: Carbon, nitrogen, and phosphorus. *Annual Review of Earth and Planetary Sciences*, 46: 521-548. <https://doi.org/10.1146/annurev-earth-082517-010018>
- Bockheim, J., Ballard, T. & Wellington, R.** 1975. Soil disturbance associated with timber harvesting in southwestern British Columbia. *Canadian Journal of Forest Research*, 5(2): 285-290. <https://doi.org/10.1139/x75-039>
- Boltz, F., Holmes, T.P. & Carter, D.R.** 2003. Economic and environmental impacts of conventional and reduced-impact logging in Tropical South America: a comparative review. *Forest Policy and Economics*, 5(1): 69-81. [https://doi.org/10.1016/S1389-9341\(01\)00075-2](https://doi.org/10.1016/S1389-9341(01)00075-2)
- Bonnaud, P., Santenoise, P., Tisserand, D., Nourrisson, G. & Ranger, J.** 2019. Impact of compaction on two sensitive forest soils in Lorraine (France) assessed by the changes occurring in the perched water table. *Forest Ecology and Management*, 437: 380-395. <https://doi.org/10.1016/j.foreco.2019.01.029>
- Bottinelli, N., Capowiez, Y. & Ranger, J.** 2014. Slow recovery of earthworm populations after heavy traffic in two forest soils in northern France. *Applied Soil Ecology*, 73: 130-133. <https://doi.org/10.1016/j.apsoil.2013.08.017>
- Cambi, M., Certini, G., Neri, F. & Marchi, E.** 2015. The impact of heavy traffic on forest soils: A review. *Forest Ecology and Management*, 338: 124-138. <https://doi.org/10.1016/j.foreco.2014.11.022>
- Chase, C.W., Reiter, M., Homyack, J.A., Jones, J.E. & Sucre, E.B.** 2019. Soil disturbance and stream-adjacent disturbance from tethered logging in Oregon and Washington. *Forest Ecology and Management*, 454: 117672. <https://doi.org/10.1016/j.foreco.2019.117672>

- Ellis, P.W., Gopalakrishna, T., Goodman, R.C., Putz, F.E., Roopsind, A., Umunay, P.M., Zalman, J., Ellis, E.A., Mo, K., Gregoire, T.G. & Griscom, B.W. 2019. Reduced-impact logging for climate change mitigation (RIL-C) can halve selective logging emissions from tropical forests. *Forest Ecology and Management*, 438: 255–266. <https://doi.org/10.1016/j.foreco.2019.02.004>
- Epron, D., Plain, C., Lerch, T. & Ranger, J. 2016. Les sols forestiers, puits de méthane: un service écosystémique méconnu. *Revue Forestière Française*. <https://doi.org/10.4267/2042/62129>
- Goutal-Pousse, N., Boc, J. & Ranger, J. 2014. Impacts de la circulation d'un porteur forestier sur deux sols sensibles au tassement et dynamique de restauration naturelle. *Rendez-vous techniques*, (43): 33-39.
- Goutal, N., Boivin, P. & Ranger, J. 2012a. Assessment of the natural recovery rate of soil specific volume following forest soil compaction. *Soil Science Society of America Journal*, 76(4): 1426-1435. <https://doi.org/10.2136/sssaj2011.0402>
- Goutal, N., Parent, F., Bonnaud, P., Demaison, J., Nourrisson, G., Epron, D. & Ranger, J. 2012b. Soil CO<sub>2</sub> concentration and efflux as affected by heavy traffic in forest in northeast France. *European Journal of Soil Science*, 63(2): 261-271.
- Goutal, N., Renault, P. & Ranger, J. 2013. Forwarder traffic impacted over at least four years soil air composition of two forest soils in northeast France. *Geoderma*, 193: 29-40. <https://doi.org/10.1016/j.geoderma.2012.10.012>
- Griscom, B.W., Ellis, P.W., Burivalova, Z., Halperin, J., Marthinus, D., Runting, R.K., Shoch, D. & Putz, F.E. 2019. Reduced-impact logging in Borneo to minimize carbon emissions and impacts on sensitive habitats while maintaining timber yields. *Forest Ecology and Management*, 438: 176-185. <https://doi.org/10.1016/j.foreco.2019.02.025>
- Hartmann, M., Niklaus, P.A., Zimmermann, S., Schmutz, S., Kremer, J., Abarenkov, K., Lüscher, P., Widmer, F. & Frey, B. 2014. Resistance and resilience of the forest soil microbiome to logging-associated compaction. *The ISME journal*, 8(1): 226-244. <https://doi.org/10.1038/ismej.2013.141>
- Healey, J.R., Price, C. & Tay, J. 2000. The cost of carbon retention by reduced impact logging. *Forest Ecology and Management*, 139(1-3): 237-255. [https://doi.org/10.1016/S0378-1127\(00\)00385-6](https://doi.org/10.1016/S0378-1127(00)00385-6)
- Horn, R., Vossbrink, J., Peth, S. & Becker, S. 2007. Impact of modern forest vehicles on soil physical properties. *Forest Ecology and Management*, 248(1-2): 56-63.
- Labrière, N., Locatelli, B., Laumonier, Y., Freycon, V. & Bernoux, M. 2015. Soil erosion in the humid tropics: A systematic quantitative review. *Agriculture, Ecosystems & Environment*, 203: 127-139. <https://doi.org/10.1016/j.agee.2015.01.027>
- Magaud, P. 2020. Débardage par câble aérien – guide technique pour gestionnaires forestiers et entreprises. Office national des Forêts. (also available at: <http://www.poleexcellencebois.fr/activites/projets-collaboratifs/5-formicable>).
- Miller, J.H. & Sirois, D.L. 1986. Soil Disturbance by Skyline Yarding vs. Skidding in a Loamy Hill Forest. *Soil Science Society of America Journal*, 50(6): 1579-1583. <https://doi.org/10.2136/sssaj1986.03615995005000060039x>



- Miller, S.D., Goulden, M.L., Hutyra, L.R., Keller, M., Saleska, S.R., Wofsy, S.C., Figueira, A.M.S., Da Rocha, H.R. & De Camargo, P.B.** 2011. Reduced impact logging minimally alters tropical rainforest carbon and energy exchange. *Proceedings of the National Academy of Sciences*, 108(48): 19431-19435. <https://doi.org/10.1073/pnas.1105068108>
- Mori, T., Imai, N. & Kitayama, K.** 2018. A preliminary report: does reduced impact logging (RIL) mitigate non-CO<sub>2</sub> greenhouse gas emissions from natural production forests? *Tropics*, 27(1): 25-31. <https://doi.org/10.3759/tropics.MS17-08>
- Naghdi, R., Lotfalian, M., Bagheri, I. & Jalali, A.M.** 2009. Damages of skidder and animal logging to forest soils and natural regeneration. *Croatian Journal of Forest Engineering: Journal for Theory and Application of Forestry Engineering*, 30(2): 141-149. <https://doi.org/10.13140/2.1.4671.8083>
- Naghdi, R., Solgi, A. & Ilstedt, U.** 2016a. Soil chemical and physical properties after skidding by rubber-tired skidder in Hyrcanian forest, Iran. *Geoderma*, 265: 12-18. <https://doi.org/10.1016/j.geoderma.2015.11.009>
- Naghdi, R., Solgi, A., Zenner, E.K., Tsioras, P.A. & Nikooy, M.** 2016b. Soil disturbance caused by ground-based skidding at different soil moisture conditions in Northern Iran. *International Journal of Forest Engineering*, 27(3): 169-178. <https://doi.org/10.1080/14942119.2016.1234196>
- Pereira, R., Zweede, J., Asner, G.P. & Keller, M.** 2002. Forest canopy damage and recovery in reduced-impact and conventional selective logging in eastern Para, Brazil. *Forest Ecology and Management*, 168(1): 77-89. [https://doi.org/10.1016/S0378-1127\(01\)00732-0](https://doi.org/10.1016/S0378-1127(01)00732-0)
- Picchio, R., Mederski, P.S. & Tavankar, F.** 2020. How and How Much, Do Harvesting Activities Affect Forest Soil, Regeneration and Stands? *Current Forestry Reports*, 6(2): 115-128. <https://doi.org/10.1007/s40725-020-00113-8>
- Pinard, M.A., Barker, M.G. & Tay, J.** 2000. Soil disturbance and post-logging forest recovery on bulldozer paths in Sabah, Malaysia. *Forest Ecology and Management*, 130(1): 213-225. [https://doi.org/10.1016/S0378-1127\(99\)00192-9](https://doi.org/10.1016/S0378-1127(99)00192-9)
- Powers, R.F., Scott, D.A., Sanchez, F.G., Voldseth, R.A., Page-Dumroese, D., Elioff, J.D. & Stone, D.M.** 2005. The North American long-term soil productivity experiment: findings from the first decade of research. *Forest Ecology and Management*, 220(1-3): 31-50. <https://doi.org/10.1016/j.foreco.2005.08.00>
- Putz, F.E. & Pinard, M.A.** 1993. Reduced-Impact Logging as a Carbon-Offset Method. *Conservation Biology*, 7(4): 755-757.
- Putz, F.E., Sist, P., Fredericksen, T. & Dykstra, D.** 2008. Reduced-impact logging: challenges and opportunities. *Forest Ecology and Management*, 256(7): 1427-1433. <https://doi.org/10.1016/j.foreco.2008.03.036>
- Rivera, S., Kershner, J.L. & Dobrowolski, J.P.** 2010. Evaluation of the surface erosion from different timber yarding methods in Honduras. *Revista Árvore*, 34(4): 577-586. <https://doi.org/10.1590/S0100-67622010000400002>

**Schweier, J. & Ludowicy, C.** 2020. Comparison of A Cable-Based and a Ground-Based System in Flat and Soil-Sensitive Area: A Case Study from Southern Baden in Germany. *Forests*, 11(6): 611. <https://doi.org/10.3390/f11060611>

**Solgi, A., Najafi, A., Page-Dumroese, D.S. & Zenner, E.K.** 2020. Assessment of topsoil disturbance caused by different skidding machine types beyond the margins of the machine operating trail. *Geoderma*, 367: 114238. <https://doi.org/10.1016/j.geoderma.2020.114238>

**Tchiofo Lontsi, R., Corre, M.D., van Straaten, O. & Veldkamp, E.** 2019. Changes in soil organic carbon and nutrient stocks in conventional selective logging versus reduced-impact logging in rainforests on highly weathered soils in Southern Cameroon. *Forest Ecology and Management*, 451: 117522. <https://doi.org/10.1016/j.foreco.2019.117522>

**Warlo, H., von Wilpert, K., Lang, F. & Schack-Kirchner, H.** 2019. Black Alder (*Alnus glutinosa* (L.) Gaertn.) on Compacted Skid Trails: A Trade-off between Greenhouse Gas Fluxes and Soil Structure Recovery? *Forests*, 10(9): 726. <https://doi.org/10.3390/f10090726>

**Worrell, W.C., Bolding, M.C. & Aust, W.M.** 2011. Potential soil erosion following skyline yarding versus tracked skidding on bladed skid trails in the Appalachian region of Virginia. *Southern Journal of Applied Forestry*, 35(3): 131-135. <https://doi.org/10.1093/sjaf/35.3.131>

## 2. Continuous cover forestry and extended rotations

David Paré<sup>1</sup>, Laurent Augusto<sup>2</sup>

<sup>1</sup>Canadian Forest Service, Laurentian Forestry Centre, Québec, Canada

<sup>2</sup>INRAE, Bordeaux Sciences Agro, Villenave d'Ornon, France

### 1. Description of the practice

Continuous cover forestry (CCF; see the full description in Helliwell and Wilson, 2012) includes many silvicultural systems which all involve continuous and uninterrupted maintenance of forest cover and which avoid clearcutting. It implies forest management that works with the characteristics of the site and with tree species that are well adapted to the location. It respects the processes inherent to the site, rather than imposing artificial uniformity, and will normally involve a mixture of tree species and ages. Management is based on the selection and favouring of individual trees (of all sizes) rather than the creation of areas of uniform tree size and spacing, and record keeping is based on periodic recording of stem diameters on sample areas, rather than by age and area of stands. Stand structure will be permanently irregular, although the process of transformation to an uneven-aged condition might involve temporary even-aged elements, possibly including small-scale clearfells, and group or irregular shelterwoods (Helliwell and Wilson, 2012). CCF could minimize soil disturbance because a larger portion of tree roots are preserved following wood harvesting, and because no soil preparation – such as ploughing– is done. On the other hand, it involves a greater number of soil trampling events as interventions are more frequent. CCF also requires more frequent and more technical interventions of the forest managers. CCF may limit changes to the soil microclimate due to smaller openings comparatively to clear-cutting with potential influence on soil organic matter decomposition. However, positive as well as negative impact of large canopy openings have been observed on organic matter decay rates (Mayer *et al.*, 2020).

## 2. Range of applicability

Applicability is worldwide, wherever even-aged forestry is practiced. Typically, over the past two centuries, conventional forest management approaches have favored the plantation of even-aged, single-species stands. Interest in alternative management approaches that involve continuous and uninterrupted maintenance of forest cover have greatly increased in many regions, particularly in developed economies (Puetzman *et al.*, 2015).

## 3. Impact on soil organic carbon stocks

Although precise data on SOC changes are scarce, meta-analyses have revealed that clearcut harvesting results in reductions of < 10 percent of the soil C in the entire soil profile with greatest loss of the forest floor (Johnson, 1992; Johnson and Curtis, 2001; Achat *et al.*, 2015). In two meta-analyses of studies in temperate forests, forest harvesting reduced total soil C by an average of 6–8 percent: C storage declined by 22–30 percent in the forest floor, whereas the mineral horizons showed no significant overall change (Nave *et al.*, 2010; Achat *et al.*, 2015). Evidence that CCF reduces soil C losses in comparison to clear cutting is scant. Mayer *et al.* (2020) reported on the following studies: In Norway-spruce-dominated stands in Austria, single-tree-selection management resulted in 11 percent greater soil C stocks in the upper mineral soil compared to conventional even age-class management (Pötzelsberger and Hasenauer, 2015). However, short-term losses were observed in shelterwood cuts in Chilean Patagonia (Klein *et al.*, 2008). In an oak-hardwood forest in New England, Warren and Ashton (2014) reported a decrease in the soil C stocks in the mineral soil, but neutral effects in the litter layer following shelterwood harvest. Others have found little or no difference between effects of partial, selection, shelterwood, and clearcut harvesting on soil C stocks (Hoover, 2011; Christophel *et al.*, 2015; Puhlick *et al.*, 2016). When differences in SOC content were observed, they were higher under CCF but of low magnitude (Pötzelsberger and Hasenauer, 2015; Jonard *et al.*, 2017). Similarly, two meta-analyses (Liao *et al.* 2010, 2012) showed a systematic loss of SOC in planted even-aged forests compared to naturally regenerated forests, but this difference in SOC storage could be linked to the fact that the naturally regenerating forests in these studies are partly primary forests, with SOC stocks probably at high level. In summary, information is too fragmentary to attribute any soil C changes with the adoption of CCF in replacement of a traditional even-aged silviculture system (Powers *et al.*, 2011). Local information on the effect of this practice on soil erosion, soil disturbance, as well as on impact on forest composition would be factors to consider due to their potential impact on soil C stocks. It is noteworthy that CCF systems involve light but more frequent interventions that could make changes in the soil C stocks at the whole rotation scale difficult to detect statistically.

## 4. Other benefits of the practice

### 4.1 Minimization of threats to soil functions

Table 3. Soil threats

Soil threats	
<b>Soil erosion</b>	On sites sensitive to erosion, maintaining a forest canopy as well as tree root systems could prevent erosion.
<b>Nutrient imbalance and cycles</b>	Especially if the adoption of this practice promotes mixed species stands.
<b>Soil acidification</b>	The absence of clearcut in CCF might slightly reduce the losses of cations induced by water leaching enhanced by clearcuts in sites prone to such losses (i.e. mainly soils with a high drainage regime but with a low buffering capacity of pH).
<b>Soil biodiversity loss</b>	In general, positive impact on biodiversity noting that some forest species need open canopy conditions or high disturbance levels. But the latter (i.e. ruderal species) are usually not a concern for biodiversity (Puetman <i>et al.</i> 2015).
<b>Soil compaction</b>	In general, the maintenance of the tree root systems might increase the resistance of soils to compaction. However, the current knowledge about this possible effect is scarce.
<b>Soil water management</b>	CCF enables better regulation of the water fluxes at the watershed scale.

### 4.2 Increases in production (e.g. food/fuel/feed/timber)

Some studies indicate equivalent or lower productivity (-20 percent) rates in CCF systems (reviewed by Lundmark *et al.*, 2016).

### 4.3 Mitigation of and adaptation to climate change

There is no strong evidence that soil C stock differs between CCF and clear-cut systems. Local situations should be examined carefully. In a broader perspective, mitigation benefits from the outflow of forest products that substitute the use of materials generating greater GHG emissions in addition to those related to the changes in C stocks, both in forest ecosystems and in wood products need to be considered. Lundmark *et al.* (2016) indicated that for Norway spruce in Sweden, biomass growth and yield is more important than the choice of silvicultural system per se for generating long term climate mitigation benefits associated with CO<sub>2</sub> emissions and C stock changes. In Canada, Paradis, Thiffault and Achim (2018) indicated that forest management systems

that produce trees of greater size should increase the proportion of long-lived wood products, suggesting that the quality of the timber produced also has implication on GHG mitigation.

There is evidence that multi-species and multi-cohorts forest stands are more resilient to climate-change and to other threats, especially in the long-term (reviewed by Puettmann *et al.*, 2015). However, unexpected mortality of residual trees may occur with CCF, especially when foresters have little experience with such practice (Puettmann *et al.*, 2015).

Due to climate change and associated effects, it is envisaged to reduce these rotation durations to mitigate the risks associated with storms, fires or pathogen attacks (Roux *et al.*, 2017). This amounts to shifting from carbon sequestration in the ecosystem to carbon storage in products and increasing the share of substitution (Fortin *et al.*, 2012). From an ecosystem perspective, shortening rotations can have an impact on soil fertility and SOC with a general decreasing trend (Achat *et al.*, 2018). As the stand is renewed more often over the same period of time, biomass and nutrient exports are greater and the effects of soil preparation during forest soil regeneration on SOC are also amplified. Thus, the longer the duration of rotation, the more likely it is that SOC will increase, although the effect of very long rotations (*i.e.* several centuries) remains poorly known with variable results depending on the studies (Ji *et al.*, 2017; Leuschner *et al.*, 2014; Zhou *et al.*, 2006). Thus, simultaneously, the risk of climate change-related hazards increases (and may contribute to the reduction of the SOC stock in trees) with the length of the rotation, whereas the shorter the rotation, the greater the SOC losses related to forest management may be (Seely, Welham and Blanco, 2010). However, to date, there are no studies where forest stands are monitored longitudinally (not in chronosequence) over a longer timeframe and have experienced intensive silviculture (several short revolutions) on the one hand or extensive silviculture on the other (long rotation over the same time period). Only model-based studies can address the effect of the length of rotation on SOC (Johnson, Scatena and Pan, 2010; Wang *et al.*, 2013; Seely, Welham and Kimmins, 2002; Achat *et al.*, 2018). Numerical simulations are generally concordant and suggest a decrease in SOC stocks with a shortening of rotations (e.g. -15 to -20 percent after 360 years; Johnson *et al.*, 2010).

## 4.4 Socio-economic benefits

Continuous-cover forestry could generate more uniform cash flows (Puettmann *et al.* 2015); successful natural regeneration avoids the cost of plantation establishment.

Improvement of landscape visual quality and enhanced recreational opportunities (Puettmann *et al.* 2015).

## 5. Potential drawbacks to the practice

### 5.1 Decreases in production (e.g. food/fuel/feed/timber)

Some studies indicate equivalent or lower productivity (-20 percent) rates in CCF systems (reviewed in Lundmark *et al.*, 2016).

## 6. Recommendations before implementation of the practice

The greatest risk is the enhanced mortality of residual trees due to damages to roots and stems during operation or to greater exposure of residual trees to wind, drought or insects. It is advisable to start at small scale with a good knowledge of species autecology and with clear stand-density management goals.

## 7. Potential barriers to adoption

Table 4. Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Implementation is not always easy depending on the current forest composition and structure, may need several steps.
Economic	Yes	Yields are less certain in regions with no tradition of CCF.

## Photo of the practice



**Photo 2.** A mature stand of Douglas fir managed on CCF principles with a developing understorey of mixed conifers, including Douglas fir, western hemlock, western red cedar and grand fir, Coombs Wood, Cumbria, United Kingdom of Great Britain and Northern Ireland



## References

- Achat, D.L., Fortin, M., Landmann, G., Ringeval, B. & Augusto, L.** 2015. Forest soil carbon is threatened by intensive biomass harvesting. *Scientific Reports*, 5: 15 991. <https://doi.org/10.1038/srep15991>
- Achat, D.L., Martel, S., Picart, D., Moisy, C., Augusto, L., Bakker, M.R. & Loustau, D.** 2018. Modelling the nutrient cost of biomass harvesting under different silvicultural and climate scenarios in production forests. *Forest Ecology and Management*, 429: 642-653. <http://dx.doi.org/10.1016/j.foreco.2018.06.047>
- Christophel, D., Höllerl, S., Prietzel, J. & Steffens, M.** 2015. Long-term development of soil organic carbon and nitrogen stocks after shelterwood- and clear-cutting in a mountain forest in the Bavarian Limestone Alps. *European Journal of Forest Research*, 134: 623–640. <https://doi.org/10.1007/s10342-015-0877-z>
- Fortin, M., Ningre, F., Robert, N. & Mothe, F.** 2012. Quantifying the impact of forest management on the carbon balance of the forest-wood product chain: A case study applied to even-aged oak stands in France. *Forest Ecology and Management*, 279: 176-188. <http://dx.doi.org/10.1016/j.foreco.2012.05.031>
- Helliwell, R. & Wilson, E.R.** 2012. Continuous cover forestry in Britain: challenges and opportunities. *Quarterly Journal of Forestry*, 106(3): 214-224.
- Ji, Y.H., Guo, K., Fang, S.B., Xu, X.N., Wang, Z.G. & Wang, S.D.** 2017. Long-term growth of temperate broadleaved forests no longer benefits soil C accumulation. *Scientific Reports*, 7. <http://dx.doi.org/10.1038/srep42328>
- Johnson, D.W.** 1992. Effects of forest management on soil organic carbon. *Water Air and Soil Pollution*, 64: 83-120.
- Johnson, K., Scatena, F.N. & Pan, Y.D.**, 2010. Short- and long-term responses of total soil organic carbon to harvesting in a northern hardwood forest. *Forest Ecology and Management*, 259 (7): 1262-1267. <http://dx.doi.org/10.1016/j.foreco.2009.06.049>
- Jonard, M., Nicolas, M., Coomes, D.A., Caignet, I., Saenger, A. & Ponette, Q.** 2017. Forest soils in France are sequestering substantial amounts of carbon. *Science of the Total Environment*, 574: 616-628. <http://dx.doi.org/10.1016/j.scitotenv.2016.09.028>
- Leuschner, C., Wulf, M., Bauchler, P. & Hertel, D.** 2014. Forest continuity as a key determinant of soil carbon and nutrient storage in beech forests on sandy soils in northern Germany. *Ecosystems*, 17(3): 497-511. <http://dx.doi.org/10.1007/s10021-013-9738-0>
- Liao, C.Z., Luo, Y.Q., Fang, C.M., Chen, J.K. & Li, B.** 2012. The effects of plantation practice on soil properties based on the comparison between natural and planted forests: a meta-analysis. *Global Ecology and Biogeography*, 21(3): 318-327. <http://dx.doi.org/10.1111/j.1466-8238.2011.00690.x>

- Liao, C.Z., Luo, Y.Q., Fang, C.M. & Li, B. 2010. Ecosystem carbon stock influenced by plantation practice: implications for planting forests as a measure of climate change mitigation. *Plos One*, 5(5). <http://dx.doi.org/10.1371/journal.pone.0010867>
- Lundmark, T., Bergh, J., Nordin, A., Fahlvik, N. & Poudel, B.C. 2016. Comparison of carbon balances between continuous-cover and clearcut forestry in Sweden. *Ambio*, 45: 203–213. <https://doi.org/10.1007/s13280-015-0756-3>
- Mayer, M., Prescott, C.E., Abaker, W.E.A., Augusto, L., Cécillon, L., Ferreira, G.W.D., James, J., Jandl, R., Katzensteiner, K., Laclau, J.-P., Laganière, J., Nouvellon, Y., Paré, D., Stanturf, J.A., Vanguelova, E.I. & Vesterdal, L. 2020. Tamm review: influence of forest management activities on soil organic carbon stocks: a knowledge synthesis. *Forest Ecology and Management*, 466: 118127. <https://doi.org/10.1016/j.foreco.2020.118127>
- Nave, L.E., Vance, E.D., Swanston, C.W. & Curtis, P.S. 2010. Harvest impacts on soil carbon storage in temperate forests. *Forest Ecology and Management*, 259: 857–866. <https://doi.org/10.1016/j.foreco.2009.12.009>
- Paradis, L., Thiffault, E. & Achim, A. 2019. Comparison of carbon balance and climate change mitigation potential of forest management strategies in the boreal forest of Quebec (Canada). *Forestry*, 92: 264–277. <https://doi.org/10.1093/forestry/cpz004>
- Pötzelsberger, E. & Hasenauer, H. 2015. Soil change after 50 years of converting Norway spruce dominated age class forests into single tree selection forests. *Forest Ecology and Management*, 338: 176–182. <https://doi.org/10.1016/j.foreco.2014.11.026>
- Powers, M., Kolka, R., Palik, B., McDonald, R. & Jurgensen, M. 2011. Long-term management impacts on carbon storage in Lake States forests. *Forest Ecology and Management*, 262(3): 424–431. <http://dx.doi.org/10.1016/j.foreco.2011.04.008>
- Puettmann, K.J., Wilson, S. McG., Baker, S.C., Donoso, P.J., Drössler, L., Amente, G., Harvey, B.D., Knoke, T., Lu, Y., Nocentini, S., Putz, F.E., Yoshida, T. & Bauhus, J. 2015. Silvicultural alternatives to conventional even-aged forest management - what limits global adoption? *Forest Ecosystems*, 2. <https://doi.org/10.1186/s40663-015-0031-x>
- Roux, A., Dhôte, J.-F. (Coordinators), Achat, D., Bastick, C., Colin, A., Bailly, A., Bastien, J.-C., Berthelot, A., Bréda, N., Cauria, S., Carnus, J.-M., Gardiner, B., Jactel, H., Leban, J.-M., Lobianco, A., Loustau, D., Meredieu, C., Marçais, B., Martel, S., Moisy, C., Pâques, L., Picart-Deshors, D., Rigolot, E., Saint-André, L. & Schmitt, B. 2017. *Quel rôle pour les forêts et la filière forêt-bois françaises dans l'atténuation du changement climatique? Une étude des freins et leviers forestiers à l'horizon 2050* (Doctoral dissertation, Institut National de la Recherche Agronomique (INRA); Institut Géographique National (IGN)).
- Seely, B., Welham, C. & Blanco, J.A., 2010. Towards the application of soil organic matter as an indicator of forest ecosystem productivity: Deriving thresholds, developing monitoring systems, and evaluating practices. *Ecological Indicators*, 10(5): 999–1008. <http://dx.doi.org/10.1016/j.ecolind.2010.02.008>

**Seely, B., Welham, C. & Kimmins, H.**, 2002. Carbon sequestration in a boreal forest ecosystem: results from the ecosystem simulation model, FORECAST. *Forest Ecology and Management*, 169 (1-2): 123-135. [http://dx.doi.org/10.1016/s0378-1127\(02\)00303-1](http://dx.doi.org/10.1016/s0378-1127(02)00303-1)

**Wang, W.F., Wei, X.H., Liao, W.M., Blanco, J.A., Liu, Y.Q., Liu, S.R., Liu, G.H., Zhang, L., Guo, X.M. & Guo, S.M.** 2013. Evaluation of the effects of forest management strategies on carbon sequestration in evergreen broad-leaved (*Phoebe bournei*) plantation forests using FORECAST ecosystem model. *Forest Ecology and Management*, 300: 21-32. <http://dx.doi.org/10.1016/j.foreco.2012.06.044>

**Zhou, G.Y., Liu, S.G., Li, Z., Zhang, D.Q., Tang, X.L., Zhou, C.Y., Yan, J.H. & Mo, J.M.**, 2006. Old-growth forests can accumulate carbon in soils. *Science*, 314(5804): 1417-1417. <http://dx.doi.org/10.1126/science.1130168>

## 3. Residue retention

Gabriel W.D. Ferreira<sup>1</sup>, Cindy E. Prescott<sup>2</sup>

<sup>1</sup>Savannah River Ecology Laboratory, University of Georgia, United States of America

<sup>2</sup>Faculty of Forestry, University of British Columbia, Vancouver, Canada

### 1. Description of the practice

Residue retention consists of leaving on-site all tree components with low merchantable value as well as litter following wood harvesting (i.e. removing only the stems of merchantable trees). Depending on the intensity of harvesting, forest harvest residues can be composed of leaves/needles, branches, twigs, low-quality or small-diameter stems, bark, dead wood, and roots. The amount of residue generated during harvest varies with forest type, productivity, climate, and rotation length. Retention of residues and litter provides carbon and nutrients to be recycled back into the soil. Residues are retained when stem-only harvesting is practiced, and the resulting residues are distributed back over the site. Residues are lost from the site when whole-tree harvesting is employed, when residues are collected for use in energy production, or when residues are piled and burned.

### 2. Range of applicability

Applicable globally.

### 3. Impact on soil organic carbon stocks

Removal of forest harvest residue removal has variable effects on SOC stocks (Table 5). Some meta-analyses (Clarke *et al.*, 2015; James and Harrison, 2016; Hume *et al.*, 2018) and reviews (Johnson and Curtis, 2001; Thiffault *et al.*, 2011; Clarke *et al.*, 2015) found no clear evidence of reduced SOC following the removal of forest residues, while others have reported significant reductions in SOC (Johnson and Curtis 2001; Achat *et al.* 2015a, 2015b). The benefits of residue retention for SOC are greatest in the surface organic layer (Clarke

*et al.*, 2015; Wan *et al.* 2018), in soils that are coarse-textured (Oliveira *et al.*, 2018; Wan *et al.* 2018) and/or poor in organic matter (Thiffault *et al.*, 2011), in temperate rather than boreal forests (Achat *et al.*, 2015b), in conifer forests (Johnson and Curtis 2001), and at high harvest intensity (Achat *et al.*, 2015b). Very high SOC losses with whole-tree harvesting (up to 50 percent in the 0–5 cm soil layer) have been reported in short-rotation Eucalyptus forests in the Democratic Republic of the Congo and Brazil (Epron *et al.*, 2015; Rocha *et al.*, 2018). In contrast, on peaty soils in the UK, SOC stocks were higher under whole-tree harvesting compared to stem-only harvesting (Vanguelova *et al.*, 2010).

Few data are available on the long-term responses of harvest residue manipulation in tropical forests compared to temperate or boreal forests (Achat *et al.*, 2015b; Mayer *et al.* 2020).

**Table 5.** Changes in soil organic carbon stocks reported for harvest residue retention

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Years)	More information	Reference
Various	Temperate and subtropical	Various	NA	None	1 - 20+ after harvest	Review (Compilation of 53 studies)	Thiffault <i>et al.</i> (2011)
Finland	Continental	NA	NA	0.69	10	Forest floor + 0 - 10 cm mineral soil	Kaarakka <i>et al.</i> (2014)
Australia	Mediterranean	Podzol; Ferralsol	54.7 - 72.7	0.62	5	0 - 20 cm	Mendham <i>et al.</i> (2003; 2002)
Canada	Continental	Inceptisols; Spodosols	46 - 65	-0.08 - 0.06	20	0 - 20 cm (14 sites)	Morris <i>et al.</i> (2019)
Finland	Continental			0.52	10 - 11	Forest floor + 0 - 10 cm mineral soil (6 sites)	Smolander <i>et al.</i> (2015)
Brazil	Tropical	Acrisols	18.2 - 23.1	0.04 - 1.82	3	0 - 10 cm	Oliveira <i>et al.</i> (2018)

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

By retaining nutrient-rich material such as leaves and bark on site, residue retention minimizes nutrient losses associated with harvesting (Achat *et al.*, 2015a; Paré and Thiffault, 2016). The increased organic matter with residue retention increases soil biological activity and foster earlier establishment of a stable microbial community structure (Achat *et al.*, 2015a; Baumann *et al.*, 2009; Smolander *et al.*, 2013). Retention of logging residues reduces topsoil compaction caused by heavy-machinery traffic during clear-cutting and logging operations (Achat *et al.*, 2015a), thereby mitigating possible breakdown of soil aggregates, and increases in soil bulk density and soil penetration resistance (Ampoorter *et al.*, 2007; Cambi *et al.*, 2015; Carter *et al.*, 2006).

### 4.2 Minimization of threats to soil functions

Table 6. Soil threats

Soil threats	
Soil erosion	Provide soil cover after harvesting, and protect soil from compaction, which could increase erosion susceptibility, particularly in steep terrains (Cambi <i>et al.</i> , 2015).
Nutrient imbalance and cycles	Minimize nutrient loss at harvesting and recycle nutrients back into the soil (Achat <i>et al.</i> , 2015a; Paré and Thiffault, 2016).
Soil acidification	Slightly increases soil pH and base cations, and reduces exchange acidity and exchangeable H and Al (Achat <i>et al.</i> , 2015a; Iwald <i>et al.</i> , 2013; Johnson <i>et al.</i> , 1991).
Soil biodiversity loss	Increases organic matter input to the soil, which increases soil biological activity; provides habitat for dead-wood-dependent organisms (Ranius <i>et al.</i> , 2018).
Soil compaction	Reduce topsoil compaction caused by heavy-machinery traffic during harvesting (Achat <i>et al.</i> , 2015a; Cambi <i>et al.</i> , 2015).
Soil water management	Residues act as surface mulch, which reduces evaporation and regulates soil temperature after harvesting (Thiffault <i>et al.</i> , 2011). Increases water infiltration by reducing soil compaction.

## 4.3 Increases in production (e.g. food/fuel/feed/timber)

The influence of retaining harvest residues on forest production is site-dependent. Positive impacts on tree growth of retaining harvest residues are observed particularly where the practice has promoted forest soil quality (Achat *et al.*, 2015a; Laclau *et al.*, 2010; Rocha *et al.*, 2018).

## 4.4 Mitigation of and adaptation to climate change

Improvement of soil quality and prevention of soil degradation through residue retention may reduce the need for fertilizer. Fossil fuels are used in the production, transport and application of fertilizer.

## 4.5 Socio-economic benefits

Residue retention reduces fertilizer requirements by reducing nutrient losses associated with forest harvest. By alleviating soil compaction, residue retention reduces the need for mitigative measures to improve soil health.

## 4.6 Additional benefits to the practice

Retaining logging residues may provide other ecological functions related to wildlife habitat, understory vegetation diversity, and water quality (Vance *et al.*, 2018).

# 5. Potential drawbacks to the practice

## 5.1 Increases in greenhouse gas emissions

Retention of harvest residues that could otherwise be used for bioenergy production and thereby reduce fossil fuel use could negatively affect GHG emissions. Soil organic C losses related to the removal of harvest residues has been argued to be negligible in comparison with the greenhouse mitigation benefit of avoided fossil-fuel emissions (Cowie *et al.*, 2006). In boreal forests, replacing fossil fuels by forest residues reduces the climate impacts associated with energy production by 7 percent to 62 percent (Repo *et al.*, 2012; Repo, Tuomi and Liski, 2011). Potential benefits of residue retention for GHG, such as increased C sequestration in tree biomass and SOC and potential reductions in fertilization must also be considered in calculations of net GHG emissions.

## 5.2 Conflict with other practice(s)

By creating obstacles, retention of harvest residues may make site preparation untenable or may restrict the machinery that can be used. Abundant residues can also make access difficult for mechanized or manual planters and reduce the availability of suitable planting spots. Retaining residues may increase fire risk in wildfire-prone forest ecosystems by increasing fuel load (Vance *et al.*, 2018).

## 5.3 Decreases in production (e.g. food/fuel/feed/timber)

In some cases, retention of harvest residues may impede forest regeneration, particularly in situations where residue promotes disease, insects, or wildfire susceptibility (Cleary *et al.*, 2013; Vance *et al.*, 2018). On cold sites, surface residues may reduce temperatures and growing season length, but may also reduce frost damage (Thiffault *et al.* 2011). On wet sites surface residues can hinder evaporation and make growing conditions less favorable. Residue retention can also slow down tree regeneration and early stand development by preventing soil preparation (Mayer *et al.*, 2020).

## 5.4 Other conflicts

On-site retention of woody residues may represent an opportunity cost in regions where there is an established market for using woody residues for bioenergy generation (He *et al.*, 2016; Repo *et al.*, 2015).

# 6. Recommendations before implementation of the practice

Decisions on whether to retain or remove residue are context-dependent. With regards to SOC, residue retention is most likely to have a beneficial effect in short-rotation intensive forestry plantations on soils that are coarse-textured and poor in SOC or nutrients. Residue retention is recommended in situations in which residue removal is uneconomical, operationally unfeasible, or environmentally unsustainable. Therefore, wood final use, market conditions and transport costs, machinery availability, requirement for other management practices such as site preparation, tree planting or fertilization), susceptibility of forest to wildfire, disease and pest risks, and interaction with should be considered when deciding about forest residue retention or removal. Policies, regulations, certification schemes, national guidelines and practical guides have been developed to ensure that forest biomass harvesting is sustainable (Stupak *et al.*, 2007). Regarding GHG emissions, the benefits of harvesting residues for bioenergy (i.e. replacing fossil fuels) must be weighed against the potential benefits of retaining residues (e.g. increased SOC, improved soil quality, reduced fertilizer requirement).



## 7. Potential barriers to adoption

**Table 7.** Potential barriers to adoption

Barrier	
Biophysical	Land size (smallholder cannot invest in specialized machinery requirements); Topography can create restrictions for mechanized wood processing; Wildfire-prone forests because of accumulation of fuel load (Vance <i>et al.</i> , 2018).
Cultural	Total land clearing is adopted across multiple production sites (forestry and agricultural) and changes require long-term knowledge construction.
Social	Residues can be perceived as untidy and difficult access to forests; residues removal makes forest monuments more visible; On the other hand, stump harvesting can be perceived as negative due to great interference in the ground in a “uncaring” fashion (Ranius <i>et al.</i> , 2018).
Economic	Harvesting residues increase revenue in some locations; access to specialized machinery is required to process wood in the field and manage woody-debris during replanting operations.
Institutional	Lack of economic incentives and support from governments, including subsidies and payment for ecosystem services.
Legal (Right to soil)	Lack of policy regulation and standards in some regions.
Knowledge/capacity	Requires regional evaluations; practice complexity (multiple levels) and interactions with other management practices can discourage adoption.

## Photo of the practice



**Photo 3.** Eucalypt plantation stem-only harvest

Wood has been piled for site removal, and harvest residues (branches, treetops, and roots) are left on site. January 2014, Minas Gerais, southeast Brazil

## References

- Achat, D.L., Deleuze, C., Landmann, G., Pousse, N., Ranger, J. & Augusto, L. 2015a. Quantifying consequences of removing harvesting residues on forest soils and tree growth – A meta-analysis. *Forest Ecology and Management*, 348: 124–141. <https://doi.org/10.1016/j.foreco.2015.03.042>
- Achat, D.L., Fortin, M., Landmann, G., Ringeval, B. & Augusto, L. 2015b. Forest soil carbon is threatened by intensive biomass harvesting. *Scientific Reports*, 5(1): 15991. <https://doi.org/10.1038/srep15991>
- Ampoorter, E., Goris, R., Cornelis, W.M. & Verheyen, K. 2007. Impact of mechanized logging on compaction status of sandy forest soils. *Forest Ecology and Management*, 241(1–3): 162–174. <https://doi.org/10.1016/j.foreco.2007.01.019>
- Baumann, K., Marschner, P., Smernik, R.J. & Baldock, J.A. 2009. Residue chemistry and microbial community structure during decomposition of eucalypt, wheat and vetch residues. *Soil Biology and Biochemistry*, 41(9): 1966–1975. <https://doi.org/10.1016/j.soilbio.2009.06.022>
- Cambi, M., Certini, G., Neri, F. & Marchi, E. 2015. The impact of heavy traffic on forest soils: A review. Elsevier B.V. <http://dx.doi.org/10.1016/j.foreco.2014.11.022>
- Carter, M.C., Dean, T.J., Wang, Z. & Newbold, R.A. 2006. Impacts of harvesting and postharvest treatments on soil bulk density, soil strength, and early growth of *Pinus taeda* in the Gulf Coastal Plain: A Long-Term Soil Productivity affiliated study. *Canadian Journal of Forest Research*, 36(3): 601–614. <https://doi.org/10.1139/x05-248>
- Clarke, N., Gundersen, P., Jönsson-Belyazid, U., Kjønåas, O.J., Persson, T., Sigurdsson, B.D., Stupak, I. & Vesterdal, L. 2015. Influence of different tree-harvesting intensities on forest soil carbon stocks in boreal and northern temperate forest ecosystems. *Forest Ecology & Management*, 351: 9–19.
- Cleary, M.R., Arhipova, N., Morrison, D.J., Thomsen, I.M., Sturrock, R.N., Vasaitis, R., Gaitnieks, T. & Stenlid, J. 2013. Stump removal to control root disease in Canada and Scandinavia: A synthesis of results from long-term trials. *Forest Ecology and Management*, 290: 5–14. <https://doi.org/10.1016/j.foreco.2012.05.040>
- Cowie, A.L., Smith, P. & Johnson D. 2006. Does soil carbon loss in biomass production systems negate the greenhouse benefits of bioenergy? *Mitigation and Adaptation Strategies for Global Change*, 11: 979–1002.
- Epron, D., Mouanda, C., Mareschal, L. & Koutika, L.-S. 2015. Impacts of organic residue management on the soil C dynamics in a tropical eucalypt plantation on a nutrient-poor sandy soil after three rotations. *Soil Biology and Biochemistry*, 85: 183–189.
- He, L., English, B.C., Menard, R.J. & Lambert, D.M. 2016. Regional woody biomass supply and economic impacts from harvesting in the southern U.S. *Energy Economics*, 60: 151–161. <https://doi.org/10.1016/j.eneco.2016.09.007>

- Hume, A.M., Chen, H.Y. & Taylor A.R.** 2018. Intensive forest harvesting increases susceptibility of northern forest soils to carbon, nitrogen and phosphorus loss. *Journal of Applied Ecology*, 55: 246-255.
- Iwald, J., Löfgren, S., Stendahl, J. & Karlton, E.** 2013. Acidifying effect of removal of tree stumps and logging residues as compared to atmospheric deposition. *Forest Ecology and Management*, 290(2013): 49–58. <https://doi.org/10.1016/j.foreco.2012.06.022>
- James, J. & Harrison, R.** 2016. The effect of harvest on forest soil carbon: A meta-analysis. *Forests*, 7: 308
- Johnson, D.W. & Curtis, P.S.** 2001. Effects of forest management on soil C and N storage: meta analysis. *Forest Ecology and Management*, 140: 227–238.
- Johnson, C.E., Johnson, A.H. & Siccama, T.G.** 1991. Whole-Tree Clear-Cutting Effects on Exchangeable Cations and Soil Acidity. *Soil Science Society of America Journal*, 55(2): 502. <https://doi.org/10.2136/sssaj1991.03615995005500020035x>
- Kaarakka, L., Tamminen, P., Saarsalmi, A., Kukkola, M., Helmisaari, H.S. & Burton, A.J.** 2014. Effects of repeated whole-tree harvesting on soil properties and tree growth in a Norway spruce (*Picea abies* (L.) Karst.) stand. *Forest Ecology and Management*, 313: 180–187. <https://doi.org/10.1016/j.foreco.2013.11.009>
- Laclau, J.P., Levillain, J., Deleporte, P., Nzila, J.D.D., Bouillet, J.P., Saint André, L., Versini, A., Mareschal, L., Nouvellon, Y., Thongo M'Bou, A. & Ranger, J.** 2010. Organic residue mass at planting is an excellent predictor of tree growth in Eucalyptus plantations established on a sandy tropical soil. *Forest Ecology and Management*, 260(12): 2148–2159. <https://doi.org/10.1016/j.foreco.2010.09.007>
- Mayer, M., Prescott, C.E., Abaker, W.E.A., Augusto, L., Cécillon, L., Ferreira, G.W.D., James, J., Jandl, R., Katzensteiner, K., Laclau, J.-P., Laganière, J., Nouvellon, Y., Paré, D., Stanturf, J.A., Vanguelova, E.I. & Vesterdal, L.** 2020. Tamm Review: Influence of forest management activities on soil organic carbon stocks: A knowledge synthesis. *Forest Ecology and Management*, 466(March): 118127. <https://doi.org/10.1016/j.foreco.2020.118127>
- Mendham, D.S., O'Connell, A.M., Grove, T.S. & Rance, S.J.** 2003. Residue management effects on soil carbon and nutrient contents and growth of second rotation eucalypts. *Forest Ecology and Management*, 181(3): 357–372. [https://doi.org/10.1016/S0378-1127\(03\)00007-0](https://doi.org/10.1016/S0378-1127(03)00007-0)
- Mendham, D.S., Sankaran, K. V, Connell, A.M.O. & Grove, T.S.** 2002. Eucalyptus globulus harvest residue management effects on soil carbon and microbial biomass at 1 and 5 years after plantation establishment. *Soil Biology and Biochemistry*, 34: 1903–1912. [https://doi.org/10.1016/S0038-0717\(02\)00205-5](https://doi.org/10.1016/S0038-0717(02)00205-5)
- Morris, D.M., Hazlett, P.W., Fleming, R.L., Kwiaton, M.M., Hawdon, L.A., Leblanc, J.-D., Primavera, M.J. & Weldon, T.P.** 2019. Effects of Biomass Removal Levels on Soil Carbon and Nutrient Reserves in Conifer-Dominated, Coarse-Textured Sites in Northern Ontario: 20-Year Results. *Soil Science Society of America Journal*, 83(s1): S116. <https://doi.org/10.2136/sssaj2018.08.0306>

- Paré, D. & Thiffault, E.** 2016. Nutrient Budgets in Forests Under Increased Biomass Harvesting Scenarios. *Current Forestry Reports*, 2(1): 81–91. <https://doi.org/10.1007/s40725-016-0030-3>
- Oliveira, F.C.C., Silva, I.R., Ferreira, G.W.D., Soares, E.M.B., Silva, S.R. & Silva, E.F.** 2018. Contribution of eucalyptus harvest residues and nitrogen fertilization to carbon stabilization in ultisols of southern bahia. *Revista Brasileira de Ciência do Solo*, 42: 1–15. <https://doi.org/10.1590/18069657rbcs20160340>
- Ranius, T., Hämäläinen, A., Egnell, G., Olsson, B., Eklöf, K., Stendahl, J., Rudolphi, J., Sténs, A. & Felton, A.** 2018. The effects of logging residue extraction for energy on ecosystem services and biodiversity: A synthesis. *Journal of Environmental Management*, 209: 409–425. <https://doi.org/10.1016/j.jenvman.2017.12.048>
- Repo, A., Ahtikoski, A. & Liski, J.** 2015. Cost of turning forest residue bioenergy to carbon neutral. *Forest Policy and Economics*, 57: 12–21. <https://doi.org/10.1016/j.forpol.2015.04.005>
- Repo, A., Känkänen, R., Tuovinen, J.-P., Antikainen, R., Tuomi, M., Vanhala, P. & Liski, J.** 2012. Forest bioenergy climate impact can be improved by allocating forest residue removal. *Global Change Biology Bioenergy*, 4: 202–212. <https://doi.org/10.1111/j.1757-1707.2011.01124.x>
- Repo, A., Tuomi, M. & Liski, J.** 2011. Indirect carbon dioxide emissions from producing bioenergy from forest harvest residues. *Global Change Biology Bioenergy*, 3: 107–115. <https://doi.org/10.1111/j.1757-1707.2010.01065.x>
- Rocha, J.H.T., Gonçalves, J.L. de M., Brandani, C.B., Ferraz, A. de V., Franci, A.F., Marques, E.R.G., Arthur Junior, J.C. & Hubner, A.** 2018. Forest residue removal decreases soil quality and affects wood productivity even with high rates of fertilizer application. *Forest Ecology and Management*, 430(April): 188–195. <https://doi.org/10.1016/j.foreco.2018.08.010>
- Smolander, A., Kitunen, V., Kukkola, M. & Tamminen, P.** 2013. Response of soil organic layer characteristics to logging residues in three Scots pine thinning stands. *Soil Biology and Biochemistry*, 66: 51–59. <https://doi.org/10.1016/j.soilbio.2013.06.017>
- Smolander, A., Saarsalmi, A. & Tamminen, P.** 2015. Response of soil nutrient content, organic matter characteristics and growth of pine and spruce seedlings to logging residues. *Forest Ecology and Management*, 357: 117–125. <https://doi.org/10.1016/j.foreco.2015.07.019>
- Stupak, I., Asikainen, A., Jonsell, M., Karlton, E., Lunnan, A., Mizaraité, D., Pasanen, K., Pärn H., Raulund-Rasmussen, K. & Röser D.** 2007. Sustainable utilisation of forest biomass for energy—possibilities and problems: policy, legislation, certification, and recommendations and guidelines in the Nordic, Baltic, and other European countries. *Biomass Bioenergy*, 31: 666–684.
- Thiffault, E., Hannam, K.D., Paré, D., Titus, B.D., Hazlett, P.W., Maynard, D.G. & Brais, S.** 2011. Effects of forest biomass harvesting on soil productivity in boreal and temperate forests—A review. *Environmental Reviews*, 19(1): 278–309. <https://doi.org/10.1139/a11-009>

- Vance, E.D., Prisley, S.P., Schilling, E.B., Tatum, V.L., Wigley, T.B., Lucier, A.A. & Van Deusen, P.C.** 2018. Environmental implications of harvesting lower-value biomass in forests. *Forest Ecology and Management*, 407(May 2017): 47–56. <https://doi.org/10.1016/j.foreco.2017.10.023>
- Vanguelova, E., Pitman, R., Luro, J. & Helmisaari H.-S.** 2010. Long term effects of whole tree harvesting on soil carbon and nutrient sustainability in the UK. *Biogeochemistry*, 101: 43-59.
- Wan, X., Xiao, L., Vadeboncoeur, M.A., Johnson, C.E. & Huang, Z.** 2018. Response of mineral soil carbon storage to harvest residue retention depends on soil texture: A meta-analysis. *Forest Ecology and Management*, 408(October 2017): 9–15. <https://doi.org/10.1016/j.foreco.2017.10.028>

## 4. Inclusion of N fixing species

Cindy E. Prescott, Sue J. Grayston

Faculty of Forestry, University of British Columbia, Vancouver, Canada

### 1. Description of the practice

Some tree species have evolved an association with N-fixing bacteria, which infect the tree roots, stimulating the formation of nodules in which the bacteria proliferate, and consume energy compounds from the plant (Binkley and Fisher, 2019). The bacteria convert atmospheric N into ammonia and then amino acids, some of which is exported to the tree roots and incorporated into tree biomass. The fixed N is liberated to the soil via root or mycorrhizal exudates or as tree litter, which has higher concentrations of N than litter of non-N-fixing tree species. Tree genera with N-fixing root associates (hereafter referred to as “N-fixing tree species”) include several genera of leguminous trees (such as *Acacia*, *Robinia*, *Falcataria*, *Albizia*), and “actinorhizal” species such as *Alnus*, *Eleagnus*, and *Casuarina* (Binkley and Fisher, 2019). Inputs of N from N-fixing tree species vary among species and sites but average about 75 kgN/ha/yr, which is more than an order of magnitude greater than typical annual rates of N input from atmospheric deposition (Binkley and Fisher, 2019).

Nitrogen-fixing tree species have multiple uses in forestry, both in pure stands and in mixtures with other species. In pure stands they are used to produce timber (Huong *et al.*, 2020), restore degraded soils (Mailly and Margolis 1992; Frouz *et al.*, 2009) and generate fuelwood (Mailly and Margolis, 1992). Nitrogen-fixing tree species are also planted to improve soil prior to planting commercial tree species (Voigtlaender *et al.*, 2012). Some N-fixing species are used for forest restoration (such as *Acacia koa*, Gugger *et al.*, 2018) or are used as nurse species to foster the regeneration of other native tree species (Chaer *et al.*, 2011). Nitrogen-fixing trees are also grown in mixtures with other commercial tree species such as Eucalyptus, pine and Douglas-fir (Forrester *et al.*, 2006). Potential ecological benefits of such mixtures include improved nutrient cycling, soil fertility, biomass production and carbon sequestration. Other benefits include reduced requirement for fertilizer, diversification of products, improved risk management and protection from pests and diseases (Forrester *et al.*, 2006).

## 2. Range of applicability

Nitrogen-fixing trees occur in tropical and temperate zones but are most abundant in subtropical and tropical latitudes (Mayer *et al.*, 2019). Species of *Acacia* are most often used for forestry in the tropics, and species from the *Leucaena*, *Casuarina*, *Albizia* or *Enterolobium* genera are also occasionally used. *Acacia mangium* is used in Eucalypt plantations in several tropical areas to increase the yield of Eucalypt without resorting to inorganic fertilizers. *Acacia* are also planted in monoculture, especially in South East Asia (Koutika and Richardson, 2019). Plantations of *Acacia mangium* and *Acacia auriculiformis* have been particularly important for small-holder forestry, providing important income for disadvantaged farmers (Huong *et al.*, 2020). *Acacia mangium* and *Erythrophleum fordii* and have been widely used for restoration of degraded soils in subtropical China (Luo *et al.*, 2016). In temperate latitudes, N-fixing tree species mostly belong to the *Robinia* or *Alnus* genera, and more rarely to the *Caragana* genus (Marron and Epron, 2019).

## 3. Impact on soil organic carbon stocks

Several studies in temperate and tropical regions have shown that soil C sequestration in forest soils is greater under N-fixing than under non-N-fixing species, given similar aboveground productivities (Forrester *et al.*, 2013; Binkley and Fisher, 2019). Comparisons of adjacent stands of N-fixing and non-N-fixing tree species have reported 20 percent–100 percent more soil C under the N-fixers; this would equate to 0.5–1.2 tC/ha/yr greater soil C accumulation in N-fixer forests than in comparable non-N-fixer forests (Resh, Binkley and Parotta, 2002). Analyzing 19 case studies with N-fixing species in temperate and tropical forests, Binkley (2005) reported a mean rate of C accretion in soils of 0.87 tC/ha/yr relative to non-N-fixing tree species. Overall, these soils accumulated about 12 to 15 g of C for every g of N accumulated. In a meta-analysis for north-temperate forests, Nave *et al.* (2009) reported a significant increase (+12 percent) in mineral SOC storage in response to N-fixing vegetation (Table 8).

Nitrogen-fixing trees increase SOC stocks much faster than other species on degraded soils such as post-mining sites (Frouz *et al.*, 2009), eroded soils (Zhang *et al.*, 2018), or after afforestation of savannas (Tang and Li, 2013). Inclusion of N-fixing *Acacia* trees into plantations of *Eucalyptus* can increase SOC stocks after a single rotation (Forrester *et al.*, 2013; Koutika *et al.*, 2014; Voigtlaender *et al.*, 2019). The greater rate of soil C increase under N-fixing trees results from a combination of greater stabilization of the organic matter added by the N-fixing trees, and reduced decomposition rates of older soil organic matter (Kaye *et al.*, 2002; Resh, Binkley and Parotta, 2002).



**Table 8.** Changes in soil organic carbon stocks reported for addition of N-fixing tree species

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Year)	Depth	More information	Reference
Global	Tropical and temperate	Various		0.87 0.25–1.80	-	-	19 case studies	Binkley (2005)
British Columbia (BC), Canada and Washington State, United States of America	Cold temperate moist	Haplorthod, Xerochrept	NA	1.13	23	O + top 50 cm	<i>Alnus rubra</i> mixed vs pure <i>Pseudotsuga menziesii</i> , infertile site	Binkley (1983)
				0.53	23	O + top 50 cm	<i>Alnus rubra</i> mixed vs pure <i>Pseudotsuga menziesii</i> , fertile site	
Haplumbrept		0.43–0.48		58	O + top 90 cm	<i>Alnus rubra</i> mix vs pure <i>Pseudotsuga menziesii</i>	Binkley <i>et al.</i> (1992)	
Boistfort series on Eocene basalt		0.75		40	O + top 20 cm	<i>Alnus rubra</i> vs adjacent <i>Pseudotsuga menziesii</i>	Bormann and DeBell (1981)	
Haplorthod = Humo Ferric Podzol		0.63		23	O + top 50 cm	<i>Alnus sinuata</i> mix vs pure <i>Pseudotsuga menziesii</i>	Binkley, Louisier and Cromack, (1984)	
Pointe-Noire, Democratic Republic of the Congo	Tropical moist	Ferralsols	15.9	0.27	7	Top 25 cm	<i>Acacia</i> mixed vs pure <i>Eucalyptus</i>	Koutika <i>et al.</i> (2014)
Brazil		Ferralsols		-0.22	6	O + top 15 cm		Voigtlaender <i>et al.</i> (2012)
		Andic Haplumbrepts	138	1.17	19	O + top 45 cm	<i>Alnus rubra</i> mixed with <i>Pseudotsuga menziesii</i>	Rothe <i>et al.</i> (2002)

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Year)	Depth	More information	Reference
Coastal Oregon, United States of America	Warm temperate moist	Dystrandept	35.6	1.19-1.81	12	Top 30 cm	<i>Ceanothus vs Pseudotsuga menziesii</i>	Binkley, Cromack and Fredriksen, (1982)
Victoria, Australia		Sodosol, Dermosol, Tenosol, Rudosol, Dermosol	31 - 43	1.00 to 1.44	9-17	Top 10 cm	<i>Acacia mearnsii</i>	Kasel <i>et al.</i> (2011)
Southeastern Australia		NA	NA	0.26	14	Top 20 cm	<i>Acacia dealbata vs Eucalyptus</i>	Hoogmoed <i>et al.</i> (2014)
		-0.07		<i>Acacia implexa vs Eucalyptus</i>				
Senegal	Tropical dry	Regosol	NA	0.32	34	O + top 100 cm	<i>Casuarina</i>	Maily and Margolis (1992)
Hawai'i	Tropical wet	Hydrudands		0.77	15	Top 40 cm	<i>Albizia vs Eucalyptus</i>	Resh, Binkley and Parotta, (2002)
				1.40		Top 50 cm		
		Hydrudands Andisol	128	1.1	16	Top 100 cm		Kaye <i>et al.</i> (2000)
		Hydrandepts	8	0.12-0.55	12	Top 20 cm		Garcia-Montiel and Binkley (1998)
Puerto Rico, United States of America	Tropical moist	Haplusterts	NA	1.25	16	Top 40 cm	<i>Casuarina vs Eucalyptus</i>	Resh <i>et al.</i> (2002)
		Troposamments		0.42	7			
				1.62	16		<i>Leucaena vs Eucalyptus</i>	
				0.71	7			
Victoria, Australia	Warm temperate	Acrisols Dermosols	68	2.0	8	Top 30 cm	<i>Acacia mix with Eucalyptus</i>	Forrester <i>et al.</i> (2013)

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

When used for afforestation and reforestation of lands with degraded soils, N-fixing tree species have the potential to increase nutrient content and cycling (Koutika *et al.*, 2017; Voigtlaender *et al.*, 2019), reduce surface runoff and increase water infiltration (Paula *et al.*, 2019), and increase earthworm density (Zou, 1993) and aggregate formation (Garay *et al.*, 2004). In plantations in southeastern Australia, the macroaggregate SOC pool under *Acacia mearnsii* increased at least four-fold faster than that of *Eucalyptus* and other *Acacia* species (Kasel *et al.*, 2011). Colonization of *Eucalyptus* roots by AMF and acid and alkaline phosphatase activities in soil were significantly higher when grown in mixed systems with *Acacia mangium* (Bini *et al.*, 2018).

### 4.2 Minimization of threats to soil functions

Table 9. Soil threats

Soil threats	
<b>Soil erosion</b>	Afforestation and reforestation with N-fixing trees contributes to erosion control; (Vitková <i>et al.</i> , 2017); <i>Acacia</i> used to control gully erosion in Brazil (Chaer <i>et al.</i> , 2011). <i>Casuarina equisetifolia</i> used for coastal protection and to stabilize drifting sand dunes (Maily and Margolis, 1992).
<b>Nutrient imbalance and cycles</b>	N-fixing tree species increase N content and availability especially in N-poor soils (Binkley and Fisher, 2019). <i>Alnus</i> also increases soil P cycling (Giardina <i>et al.</i> , 1995). Actinorrhizal N-fixing trees can accelerate rock weathering, and thereby enhance their access to multiple rock-derived nutrients (Perakis and Pett-Ridge, 2019).
<b>Soil biodiversity loss</b>	Higher bacteria, lower fungi, more earthworms under N-fixer (Zou, 1993). More litter-transforming macroarthropods, especially millipedes under <i>Acacia mangium</i> than <i>Eucalyptus</i> (Pellens and Garay, 1999; Zagato <i>et al.</i> , 2020). Higher soil macrofauna abundance and diversity in mixed plantations of <i>Eucalypt</i> and <i>Acacia</i> , than pure plantations of either species (Zagato <i>et al.</i> , 2020). Shifts in soil bacteria composition (Pereira <i>et al.</i> , 2017).
<b>Soil water management</b>	N-fixers used in afforestation and land restoration moderate water flows.

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Mixed forest plantations involving N-fixing species and other species such as eucalypt have been proposed to increase productivity in regions with N-deficient soils (Paula *et al.*, 2019). In a meta-analysis of 148 case studies from 34 plantations, mixed forest plantations involving at least equal proportions of N-fixing and non-N-fixing species are generally more productive than monospecific plantations (Marron and Epron, 2019). On average, mixed-tree plantations were 18 percent more productive than the non-N fixing monocultures. The size of the increase in the mixed stand was greater on lower productivity sites (Marron and Epron, 2019).

A rotation of N-fixing trees may improve site fertility prior to planting the more valuable commercial tree species. For example, twenty-year-old plots of pure Eucalyptus and pure *Falcataria* in Hawaii were harvested, and the plots replanted with the same or the opposite species. Plots with soils enriched in N by *Falcataria* in the first generation had twice as much available N and almost double the growth of *Eucalyptus* compared with soils that had Eucalyptus for both generations (Binkley and Fisher, 2019). Even intercropping *Acacia mangium* with *Eucalyptus* in Brazil led to much higher amounts of soil N in only one crop rotation (Voigtlaender *et al.*, 2019; Paula *et al.*, 2019).

### 4.4 Mitigation of and adaptation to climate change

Use of N-fixing tree species reduces the need for nitrogenous fertilizers and so eliminates GHG emissions associated with the production, distribution and application of N fertilizers (Paula *et al.*, 2019).

Mixed plantations with non-N-fixing trees and N-fixing species can sequester more C than pure plantations of non-N-fixing trees, such as Eucalyptus (Forrester *et al.*, 2006, 2013).

### 4.5 Socio-economic benefits

Mixed forest plantations provide a greater diversity of products than monospecific forest plantations (Paula *et al.*, 2019). Pairs of mixtures of fast-growing exotic species such as *Eucalyptus* sp. or *Pinus* sp. with N-fixing trees such as *Acacia mangium* or *A. mearnsii* have the potential to offer a wide range of products in the same area, including timber, firewood, charcoal, tannins, resins, and essential oils (Paula *et al.*, 2019). Intercropped plantations with fast-growing N-fixing trees and *Eucalyptus* assists the restoration of degraded areas and increases the economy of small- and medium-sized farmers in the *Cerrado* and Amazon regions of Brazil (Paula *et al.*, 2019). N-fixing trees such as *Erythrina* spp. are used to produce shade-grown cacao in Brazil (Gama-Rodrigues, 2020).

The tradition of using *Robinia* for afforestation in Central Europe has led to it being an important part of the economy in some countries. In addition to its valuable resistant wood, *Robinia* is used for honey-making and more recently for production of dead wood (Vítková *et al.*, 2017).

## 4.6 Additional benefits to the practice

Mixed-species plantations (including those containing N-fixing tree species) may be better protected from pests and diseases than are monocultures (Marron and Epron, 2019). Mixing tree species reduces the impact of specialist insect herbivores on individual susceptible tree species and the impact of specialist pathogens on host tree species (Bauhus *et al.*, 2017).

Afforestation and reforestation with N-fixing species can facilitate the recovery of natural forests on deforested and degraded soils, by creating conditions conducive to the establishment and growth of more nutrient-demanding native tree species (Yang *et al.*, 2009; Root-Bernstein *et al.*, 2017). For example, Dipterocarps have been grown in gaps of *Acacia* trees in the tropics (Norisada and Kojima, 2005). In the State of Amazônia, northern Brazil, biomass and richness of native plant species regenerating in the understory of planted N-fixing tree species was higher than in areas planted with non-N-fixing species (Chaer *et al.*, 2011).

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

Changes due to N-fixing species can be similar to those of atmospheric N deposition or N fertilization – soil acidification, elevated nitrate, and increased N losses through nitrate leaching and N<sub>2</sub>O emission.

Addition of N through N<sub>2</sub> fixation can create or exacerbate P deficiencies, especially on soils with low organic matter and P reserves or with high P adsorption capacity (Sitters, Edwards and Olde Venterink, 2013; de Carvalho Balieiro *et al.*, 2020).

**Table 10.** Soil threats

Soil threats	
<b>Nutrient imbalance and cycles</b>	N fixation increases demand for the soil nutrients P and Mo, so fertilization may be necessary to realize growth potential (Augusto <i>et al.</i> , 2013; Dynarski, Pett-Ridge and Perakis, 2020; Mailly and Margolis, 1992) P supply depleted within 1 rotation of <i>Albizia</i> in Hawai'i (Binkley <i>et al.</i> , 2000). Elevated soil nitrate persists for a decade after removal of N-fixer (Nsikani <i>et al.</i> , 2017).
<b>Soil acidification</b>	Greater soil N associated with declines in soil pH and exchangeable Ca, Mg, and K (Yamashita, Ohta and Hardiono, 2008; Binkley and Fisher, 2019).
<b>Soil water management</b>	Greater soil N from <i>Alnus</i> associated with increased nitrate leaching and aluminum mobilization (Perakis <i>et al.</i> , 2013) and associated pollution of waterbodies.

## 5.2 Increases in greenhouse gas emissions

In temperate regions, N-fixation by *Alnus* species increased N<sub>2</sub>O emissions compared to non-N-fixing species across 12 sites in an elevational gradient in Switzerland (Buhlmann *et al.*, 2017). Several studies have found greater concentrations of N, mainly NO<sub>3</sub>, in soil solution under alder (Binkley *et al.*, 1992; Buhlmann, Körner and Hiltbrunner, 2016), and increased denitrification under alder versus non-N-fixing vegetation ( Mogge, Kaiser, and Munch, 1998). Varying rates of N<sub>2</sub>O emissions have been found in *Alnus glutinosa* forests in Switzerland (0.2-1.7 kg N<sub>2</sub>O-N/ha/season; Buhlmann *et al.*, 2017), southern Germany (0.5-1.0 kg N<sub>2</sub>O-N/ha/yr; Eickenscheidt *et al.* 2014; and 3.1 Kg N<sub>2</sub>O-N ha/yr (Warlo *et al.*, 2019) and northern Germany (0.8-7.3 kg N<sub>2</sub>O-N/ha/yr; Mogge, Kaiser, and Munch, 1998). Higher rates of N<sub>2</sub>O emissions are associated with: Buhlmann *et al.* (2017) estimated stands of *Alnus viridis* in Switzerland would release 130 t N<sub>2</sub>O-N per growing season (Mid-June–mid-Oct), which represents 1.5 percent total Swiss emissions of this GHG (FOEN, 2015).

In the tropics, leguminous plantations raise N availability in the soil, making these plantations a possible N<sub>2</sub>O source (IPCC, 2003). *Acacia* are important leguminous trees for industrial plantations because of rapid growth and tolerance of acidic, nutrient-poor soils and more than 8 million ha of *Acacia* were planted in the tropics and sub-tropics by 2000, 96 percent in Asia (FAO, 2001). Stimulation of N<sub>2</sub>O emissions has been documented in leguminous plantations in the Asian tropics (Arai *et al.* 2008), where these trees are not a dominant component of native forests (Primack and Corlett, 2005). Annual N<sub>2</sub>O fluxes in *Acacia* plantations or tropical secondary forests containing native N-fixing species are eight times higher (2.56 Kg N<sub>2</sub>O/ha/yr) than those from secondary forests without N-fixing species in Indonesia (Arai *et al.*, 2008; Konda *et al.*, 2008; Erickson *et al.*, 2001). In southern China, N<sub>2</sub>O emissions from *Acacia auriculiformis* plantations (2.3 kg N<sub>2</sub>O-N/ha/yr) were significantly greater than from *Eucalyptus urophylla* (1.9 kg N<sub>2</sub>O-N/ha/yr) and was further stimulated by N fertilization to 3.1 Kg N<sub>2</sub>O-N/ha/yr (Zhang *et al.*, 2014). This demonstrates that the elevated N<sub>2</sub>O emissions result from the input of fixed N, and further suggests that elevated atmospheric N deposition will increase N<sub>2</sub>O emissions from leguminous tree plantations in the tropics. Addition of P, as well as N, to *Acacia* stands mitigated the increased N<sub>2</sub>O fluxes (reduced from 3.1 to 2.7 Kg N<sub>2</sub>O-N) due to increased N uptake by vegetation in the presence of increased P availability (Zhang *et al.*, 2014). There are pronounced seasonal fluctuations in N<sub>2</sub>O emissions in tropical forests, with higher emissions in the wet season than the dry season. Denitrification is accelerated in the wet season due to greater prevalence of anaerobic soil conditions (Werner *et al.*, 2007, Arai *et al.*, 2008, Konda *et al.*, 2010).

## 5.3 Conflict with other practice(s)

Relative to monocultures of commercial tree species, mixtures of commercial species with N-fixing tree species can be more challenging to manage.

## 5.4 Decreases in production (e.g. food/fuel/feed/timber)

When grown in mixtures, there is potential for N-fixing tree species to reduce productivity of the commercial tree species through competition for resources other than N (Lavery *et al.*, 2004). The success of mixed-tree plantations (i.e. when the mixture is more productive than the monoculture) is variable and difficult to predict, but most likely to succeed where N is limiting productivity (Marron and Epron, 2019).

## 5.5 Other conflicts

The introduction of N-fixing species outside of their natural ranges carries the risks associated with introduction of any non-native species, more so because the competitive advantage of N-fixing species can make them invasive, such as *Robinia pseudoacacia* in central Europe (Vítková *et al.*, 2017), *Acacia mangium* and *Falcataria moluccana* in tropical and subtropical regions (Hughes *et al.*, 2013; Koutika and Richardson, 2019). Although negative impacts of invasive N-fixing species have been reported in tropical and subtropical rangelands around the world, exotic N-fixing tree species (including *Robinia* and *Acacia* spp.) are used in many areas to meet the needs of people without major risk of invasion (Richardson *et al.*, 2011). A stratified approach, combining tolerance in some areas with strict eradication at some sites, has been recommended to allow for the sustainable use of N-fixing tree species (Vítková *et al.*, 2017). Invasion risk assessments are recommended prior to new plantings of N-fixing species (Koutika and Richardson, 2019).

# 6. Recommendations before implementation of the practice

- ◆ Assess likelihood of species becoming invasive;
- ◆ Investigate potential end uses and markets for candidate tree species;
- ◆ Survey local peoples' perception of species;
- ◆ Determine the N-fixing tree species and genotype best suited for the site and, if grown in mixture, for the companion tree species;
- ◆ Affirm that the productivity of the site is primarily limited by N;
- ◆ Assess likelihood of deficiencies of nutrients other than N;
- ◆ Mix or rotate species to avoid consequences of excess N relative to other nutrients

## 7. Potential barriers to adoption

Table 11. Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Site productivity must be N-limited; site conditions must be suitable for N-fixing species.
Cultural	Yes	N-fixing tree species may not be traditionally used.
Social	Yes	Risk of species becoming invasive must be weighed; local peoples' perception of species desirability.
Economic	Yes	Potential declines in overall productivity, as well as additional management costs. Development of market for products of N-fixing tree species needed.
Knowledge	Yes	Impacts on productivity of commercial tree species needs to be better understood, as well as implications on drought, disease and pest damage.



## Photo of the practice



Photo 4. *Acacia mangium* planted near Bintulu, in Sarawak, Borneo (Malaysia)

Table 12. Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Soil fertility improvement of nutrient-poor and sandy soils in the Congolese coastal plains</i>	Africa	7	6	1

## References

- Arai, S., Ishizuka, S., Ohta, S., Ansori, S., Tokuchi, N., Tanaka, N. & Hardjono, A. 2008. Potential N<sub>2</sub>O emissions from leguminous tree plantation soils in the humid tropics. *Global Biogeochemical Cycles*, 22: CB2028. <https://doi.org/10.1029/2007GB002965>.
- Augusto, L., Delerue, F., Gallet-Budynek, A. & Achat, D.L. 2013. Global assessment of limitation to symbiotic nitrogen fixation by phosphorus availability in terrestrial ecosystems using a meta-analysis approach. *Global Biogeochemical Cycles*, 27: 1–12. <https://doi.org/10.1002/gbc.20069>
- Bauhus J., Forrester D.I., Gardiner B., Jactel H., Vallejo, R. & Pretzsch, H. 2017. Ecological stability of mixed-species forests. In Pretzsch, H., Forrester, D. & Bauhus, J. (Eds.) *Mixed-Species Forests*. Springer, Berlin, Heidelberg. [https://doi.org/10.1007/978-3-662-54553-9\\_7](https://doi.org/10.1007/978-3-662-54553-9_7)
- Bini, D., Santos, C.A., Silva, M.C.P., Bonfim, J.A. & Cardoso, E.J.B.N. 2018. Intercropping *Acacia mangium* stimulates AMF colonization and soil phosphatase activity in *Eucalyptus grandis*. *Scientia Agricola*, 75(2): 102–110. <https://doi.org/10.1590/1678-992x-2016-0337>
- Binkley, D. 1983. Interaction of site fertility and red alder on ecosystem production in Douglas-fir plantations. *Forest Ecology and Management*, 5: 215–227. [https://doi.org/10.1016/0378-1127\(83\)90073-7](https://doi.org/10.1016/0378-1127(83)90073-7)
- Binkley, D. 2005. How nitrogen-fixing trees change soil carbon. In Binkley D. & Menyailo, O. (Eds.) *Tree species effects on soils: implications for global change*. pp 155–164.
- Binkley, D.J., Lousier, D. & Cromack, K. 1984. Ecosystem effects of sitka alder in a Douglas-fir plantation. *Forest Science*, 30(1): 26–35. <https://doi.org/10.1093/forestscience/30.1.26>
- Binkley, D., Cromack, K. Jr. & Fredriksen, R.L. 1982. Nitrogen accretion and availability in some snowbrush ecosystems. *Forest Science*, 28(4): 720–724. <https://doi.org/10.1093/forestscience/28.4.720>
- Binkley, D., Sollins, P., Bell, R., Sachs, D. & Myrold, D. 1992. Biogeochemistry of adjacent conifer and alder/conifer ecosystems. *Ecology*, 73: 2022–2034. <https://doi.org/10.2307/1941452>
- Binkley, D. & Fisher, R.F. 2019. *Ecology and Management of Forest Soils*. Fifth Edition ISBN: 978-1-119-45565-3 Wiley-Blackwell 456 pp.
- Binkley, D., Giardina, C. & Bashkin, M.A. 2000. Soil phosphorus pools and supply under the influence of *Eucalyptus saligna* and nitrogen-fixing *Albizia facaltaria*. *Forest Ecology and Management*, 128: 241–247. [https://doi.org/10.1016/S0378-1127\(99\)00138-3](https://doi.org/10.1016/S0378-1127(99)00138-3)
- Bormann, B.T. & DeBell, D.S. 1981. Nitrogen content and other soil properties related to age of red alder stands. *Soil Science Society of America Journal*, 45: 428–432. <https://doi.org/10.2136/sssaj1981.03615995004500020038x>
- Bühlmann, T., Körner, C. & Hiltbrunner, E. 2016. Shrub Expansion of *Alnus viridis* Drives Former Montane Grassland into Nitrogen Saturation. *Ecosystems*, 19(6): 968–985. <https://doi.org/10.1007/s10021-016-9979-9>

- Bühlmann, T., Caprez, R., Hiltbrunner, E., Körner, C. & Niklaus, P.A.** 2017. Nitrogen fixation by *Alnus* species boosts soil nitrous oxide emissions. *European Journal of Soil Science*, 68(5): 740-748. <https://doi.org/10.1111/ejss.12457>
- Chaer, G.M., Resende, A.S., Campello, E.F.C., Miana de Faria, S. & Boddey, R.M.** 2011. Nitrogen-fixing legume tree species for the reclamation of severely degraded lands in Brazil. *Tree Physiology*, 31(2): 139–149. <https://doi.org/10.1093/treephys/tpq116>
- de Carvalho Balieiro, F., Cesário, F.V. & Santos, F.M.** 2020. Litter decomposition and soil carbon stocks in mixed plantations of Eucalyptus spp. and nitrogen-fixing trees. In Nogueira Cardoso, E. Bran, Gonçalves, J., Balieiro, F. & Franco, A. (Eds.) *Mixed plantations of Eucalyptus and leguminous trees*. pp. 57-90 Springer, Cham. 280 pp. [https://doi.org/10.1007/978-3-030-32365-3\\_4](https://doi.org/10.1007/978-3-030-32365-3_4)
- Dynarski, K.A., Pett-Ridge, J.C. & Perakis, S.S.** 2020. Decadal-scale decoupling of soil phosphorus and molybdenum cycles by temperate nitrogen-fixing trees. *Biogeochemistry*, 149: 355–371. <https://doi.org/10.1007/s10533-020-00680-9>
- Eickenscheidt, T., Heinichen, J., Augustin, J., Freibauer, A. & Drösler, M.** 2014. Nitrogen mineralization and gaseous nitrogen losses from waterlogged and drained organic soils in a black alder (*Alnus glutinosa* (L.) Gaertn.) forest. *Biogeosciences*, 11, 2961–2976. <https://doi.org/10.5194/bg-11-2961-2014>
- Erickson, H., Keller, M. & Davidson, E.A.** 2001. Nitrogen oxide fluxes and nitrogen cycling during postagricultural succession and forest fertilization in the humid tropics. *Ecosystems*, 4: 67-84. <https://doi.org/10.1007/s100210000060>
- FAO.** 2001. *Global Forest Resources Assessment 2000*. 479 pp. Food and Agriculture Organization of the United Nations, Rome.
- FOEN.** 2015. *Switzerland's Greenhouse Gas Inventory 1990–2013: National Inventory Report, Including Reporting Elements Under the Kyoto Protocol*. Submission of 15 April 2015 under the United Nations Framework Convention on Climate Change and under the Kyoto Protocol. Federal Office for the Environment FOEN, Bern.
- Forrester, D.I., Bauhus, J., Cowie, A.L. & Vanclay, J.K.** 2006. Mixed-species plantations of Eucalyptus with nitrogen-fixing trees: A review. *Forest Ecology and Management*, 233: 211–230. <https://doi.org/10.1016/j.foreco.2006.05.012>
- Forrester, D.I., Pares, A., O'Hara, C., Khanna, P.K. & Bauhus, J.** 2013. Soil organic carbon is increased in mixed-species plantations of Eucalyptus and nitrogen-fixing Acacia. *Ecosystems*, 16: 123–132. <https://doi.org/10.1007/s10021-012-9600-9>
- Frouz, J., Pižl, V., Cienciala, E. & Kalčík, J.** 2009. Carbon storage in post-mining forest soil, the role of tree biomass and soil bioturbation. *Biogeochemistry*, 94: 111-121. <https://doi.org/10.1007/s10533-009-9313-0>

- Gama-Rodrigues A.C.** 2020. Multifunctional mixed-forest plantations: The use of Brazilian native leguminous tree species for sustainable rural development. In Bran Nogueira Cardoso, E., Gonçalves, J., Balieiro, F. & Franco, A. (Eds.) *Mixed plantations of Eucalyptus and leguminous trees*. pp. 241-256 Springer, Cham. [https://doi.org/10.1007/978-3-030-32365-3\\_12](https://doi.org/10.1007/978-3-030-32365-3_12)
- Garay, I., Pellens, R., Kindel, A., Barros, E. & Franco, A.I.A.** 2004. Evaluation of soil conditions in fast growing plantations of *Eucalyptus grandis* and *Acacia mangium* in Brazil: A contribution to the study of sustainable land use. *Applied Soil Ecology*, 27: 177–187. <https://doi.org/10.1016/j.apsoil.2004.03.007>
- Garcia-Montiel, D. & Binkley, D.** 1998. Effect of *Eucalyptus saligna* and *Albizia falcataria* on soil processes and nitrogen supply in Hawaii. *Oecologia*, 113: 547–556. <https://doi.org/10.1007/s004420050408>
- Giardina, C.P., Huffman, S., Binkley, D. & Caldwell, B.A.** 1995. Alders increase soil phosphorus availability in a Douglas fir plantation. *Canadian Journal of Forest Research*, 25: 1652–1657. <https://doi.org/10.1139/x95-179>
- Gugger, P.F., Liang, C.T., Sork, V.L., Hodgkiss, P & Wright, J.W.** 2018. Applying landscape genomic tools to forest management and restoration of Hawaiian koa (*Acacia koa*) in a changing environment. *Evolutionary Applications*, 11(2): 231-242. <https://doi.org/10.1111/eva.12534>
- Hoogmoed, M., Cunningham, S.C., Baker, P.J., Beringer, J. & Cavagnaro, T.R.** 2014. Is there more soil carbon under nitrogen-fixing trees than under non-nitrogen-fixing trees in mixed-species restoration plantings? *Agriculture, Ecosystems & Environment*, 188: 80–84. <https://doi.org/https://doi.org/10.1016/j.agee.2014.02.013>
- Hughes R.F., Johnson. M.T. & Uowolo, A.** 2013. *The invasive alien tree Falcataria moluccana: Its impacts and management. Proceedings of the XIII International Symposium on Biological Control of Weeds*, Waikoloa, Hawaii, USA, 11-16 September, 2011. pp.218-223. USDA Forest Service, Pacific Southwest Research Station, Institute of Pacific Islands Forestry, Hilo, USA.
- Huong, V.D., Nambiar, E.K.S., Hai, N.X., Ha, K.M. & Dang, N.V.** 2020. Sustainable management of *Acacia auriculiformis* plantations for wood production over four successive rotations in South Vietnam. *Forests*, 11, 550.
- Intergovernmental Panel on Climate Change (IPCC).** 2003, Good Practice Guidance for Land Use, Land-Use Change and Forestry. Penman, J. *et al.*, (Eds.). 599 pp., Institute for Global Environmental Strategies for the IPCC, Kanagawa.
- Kasel, S. Singh, S., Sanders, G.J. & Bennett, L.T.** 2011. Species-specific effects of native trees on soil organic carbon in biodiverse plantings across north-central Victoria, Australia. *Geoderma*, 161(1-2): 95-106. <https://doi.org/10.1016/j.geoderma.2010.12.014>
- Kaye, J.P., Resh, S.C., Kaye, M.W. & Chimmer, R.A.** 2000. Nutrient and carbon dynamics in a replacement series of Eucalyptus and Albizia trees. *Ecology*, 81: 3267-3273. [https://doi.org/10.1890/0012-9658\(2000\)081\[3267:NACDIA\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2000)081[3267:NACDIA]2.0.CO;2)

- Kaye, J.P., Binkley, D., Zou, X. & Parrotta, J.A.** 2002. Non-labile soil <sup>15</sup>Nitrogen retention beneath three tree species in a tropical plantation. *Soil Science Society of America Journal*, 66: 612-619. <https://doi.org/10.2136/sssaj2002.6120>
- Konda, R., Ohta, S., Ishizuka, S., Arai, S., Ansori, S., Tanaka, N. & Hardjono, A.** 2008. Spatial structures of N<sub>2</sub>O, CO<sub>2</sub>, and CH<sub>4</sub> fluxes from *Acacia mangium* plantation soils during a relatively dry season in Indonesia. *Soil Biology & Biochemistry*, 40: 3020-3030.
- Konda, R., Ohta, S., Ishizuka, S., Heriyanto, J. & Wicaksono, A.** 2010. Seasonal changes in the spatial structures of N<sub>2</sub>O, CO<sub>2</sub>, and CH<sub>4</sub> fluxes from *Acacia mangium* plantation soils in Indonesia. *Soil Biology & Biochemistry*, 42: 1512-1522. <https://doi.org/10.1016/j.soilbio.2010.05.022>
- Koutika, L.S., Epron, D., Bouillet, J.P. & Mareschal, L.** 2014. Changes in N and C concentrations, soil acidity and P availability in tropical mixed acacia and eucalypt plantations on a nutrient-poor sandy soil. *Plant and Soil*, 379: 205–216.
- Koutika, L.S., Tchichelle, S.V., Mareschal, L. & Epron, D.** 2017. Nitrogen dynamics in a nutrient-poor soil under mixed-species plantations of eucalypts and acacias. *Soil Biology & Biochemistry*, 108: 84–90. <https://doi.org/10.1016/j.soilbio.2017.01.023>
- Koutika, L.S. & Richardson, D.M.** 2019. *Acacia mangium* Willd: benefits and threats associated with its increasing use around the world. *Forest Ecosystems*, 6(2). <https://doi.org/10.1186/s40663-019-0159-1>
- Lavery, J.M., Comeau, P.G. & Prescott, C.E.** 2004. The influence of red alder patches on light, litterfall, and soil nutrients in adjacent conifer stands. *Canadian Journal of Forest Research*, 34: 56-64. <https://doi.org/10.1139/x03-194>
- Luo, D., Cheng, R., Shi, Z., Wang, W., Xu, G. & Liu, S.** 2016. Impacts of nitrogen-fixing and non-nitrogen-fixing tree species on soil respiration and microbial community composition during forest management in subtropical China. *Ecological Research*, 31: 683–693. <https://doi.org/10.1007/s11284-016-1377-4>
- Mailly, D. & Margolis, H. A.** 1992. Forest floor and mineral soil development in *Casuarina equisetifolia* plantations on the coastal sand dunes of Senegal. *Forest Ecology and Management*, 55: 259-278. [https://doi.org/10.1016/0378-1127\(92\)90105-I](https://doi.org/10.1016/0378-1127(92)90105-I)
- Marron, N. & Epron, D.** 2019. Are mixed-tree plantations including a nitrogen-fixing species more productive than monocultures? *Forest Ecology and Management*, 441: 242–252. <https://doi.org/https://doi.org/10.1016/j.foreco.2019.03.052>
- Mayer, M., Prescott, C.E., Abaker, W.E.A., Augusto, L., Cécillon, L., Ferreira, G.W.D., James, J., Jandl, R., Katzensteiner, K., Laclau, J.-P., Laganière, J., Nouvellon, Y., Paré, D., Stanturf, J.A., Vanguelova, E.I. & Vesterdal, L.** 2020. Tamm Review: Influence of forest management activities on soil organic carbon stocks: A knowledge synthesis. *Forest Ecology and Management*, 466: 118127. <https://doi.org/10.1016/j.foreco.2020.118127>

- Mogge, B., Kaiser, E.A. & Munch, J.C.** 1998. Nitrous oxide emissions and denitrification N-losses from forest soils in the Bornhoved Lake Region (Northern Germany). *Soil Biology & Biochemistry*, 30: 703–710. [https://doi.org/10.1016/S0038-0717\(97\)00205-8](https://doi.org/10.1016/S0038-0717(97)00205-8)
- Nave, L., Vance, E., Swanston, C. & Curtis, P.** 2009. Impacts of elevated N inputs on north temperate forest soil C storage, C/N, and net N-mineralization. *Geoderma*, 153: 231-240. <https://doi.org/10.1016/j.geoderma.2009.08.012>
- Norisada, M. & Kojima, K.** 2005. Photosynthetic characteristics of dipterocarp species planted on degraded sandy soils in southern Thailand. *Photosynthetica*, 43: 491–499. <https://doi.org/10.1007/s11099-005-0080-4>
- Nsikani, M.M., Novoa, A., van Wilgen, B.W., Keet, J.-H. & Gaertner, M.** 2017. *Acacia saligna*'s soil legacy effects persist up to 10 years after clearing: Implications for ecological restoration. *Austral Ecology*, 42: 880-889. <https://doi.org/10.1111/acc.12515>
- Paula R.R., de Oliveira I.R., Gonçalves J.L.M. & de Vicente Ferraz A.** 2020. Why mixed forest plantation? In Bran Nogueira Cardoso, E., Gonçalves, J., Balieiro, F. & Franco, A. (Eds.) *Mixed plantations of Eucalyptus and leguminous trees*. pp. 1-13. Springer, Cham. [https://doi.org/10.1007/978-3-030-32365-3\\_1](https://doi.org/10.1007/978-3-030-32365-3_1)
- Pellens, R. & Garay, I.** 1999. Edaphic macroarthropod communities in fast-growing plantations of *Eucalyptus grandis* Hill ex Maid (Myrtaceae) and *Acacia mangium* Wild (Leguminosae) in Brazil. *European Journal of Soil Biology*, 35: 77-89. [https://doi.org/10.1016/S1164-5563\(99\)00209-5](https://doi.org/10.1016/S1164-5563(99)00209-5)
- Perakis, S.S. & Pett-Ridge, J.C.** 2019. Nitrogen-fixing red alder trees tap rock-derived nutrients. *Proceedings of the National Academy of Sciences*, 116(11): 5009-5014. <https://doi.org/10.1073/pnas.1814782116>
- Perakis, S.S., Sinkhorn, E.R., Catricala, C.E., Bullen, T.D., Fitzpatrick, J.A., Hynicka, J.D. & Cromack, K.** 2013. Forest calcium depletion and biotic retention along a soil nitrogen gradient. *Ecological Applications*, 23: 1947–1961. <https://doi.org/10.1890/12-2204.1>
- Pereira, A.P.dA., de Andrade, P.A.M., Bini, D., Durrer, A., Robin, A., Bouillet, J.P., Fernando, D.A. & Bran Nogueira Cardoso, E.J.** 2017. Shifts in the bacterial community composition along deep soil profiles in monospecific and mixed stands of *Eucalyptus grandis* and *Acacia mangium*. *PLoS ONE*, 12(7): e0180371. <https://doi.org/10.1371/journal.pone.0180371>
- Primack, R. & Corlett, R.** 2005. *Tropical Rain Forests – An Ecological and Biogeographical Comparison*. p. 421. Blackwell, Malden, Massachusetts, USA.
- Resh, S.C., Binkley, D. & Parrotta, J.A.** 2002. Greater soil carbon sequestration under nitrogen-fixing trees compared with eucalyptus species. *Ecosystems*, 5: 217–231. <https://doi.org/10.1007/s10021-001-0067-3>

- Richardson, D.M., Carruthers, J., Hui, C., Impson, F.A., Miller, J.T., Robertson, M.P., Rouget, M., Le Roux, J.J. & Wilson, J.R.** 2011. Human-mediated introductions of Australian acacias – a global experiment in biogeography. *Diversity Distribution*, 17: 771–787. <https://doi.org/10.1111/j.1472-4642.2011.00824.x>
- Root-Bernstein, M., Valenzuela, R., Huerta, M., Armesto, J. & Jaksic, F.** 2017. *Acacia caven* nurses endemic sclerophyllous trees along a successional pathway from silvopastoral savanna to forest. *Ecosphere*, 8: e01667. <https://doi.org/10.1002/ecs2.1667>
- Rothe, A., Cromack, K., Jr., Resh, S.C., Makineci, E. & Son, Y.** 2002. Soil carbon and nitrogen changes under Douglas-fir with and without red alder. *Soil Science Society of America Journal*, 66: 1988–1995. <https://doi.org/10.2136/sssaj2002.1988>
- Sitters, J., Edwards, P.J. & Olde Venterink, H.** 2013. Increases of soil C, N, and P pools along an *Acacia* tree density gradient and their effects on trees and grasses. *Ecosystems*, 16: 347–357. <https://doi.org/10.1007/s10021-012-9621-4>
- Tang, G. & Li, K.** 2013. Tree species controls on soil carbon sequestration and carbon stability following 20 years of afforestation in a valley-type savanna. *Forest Ecology and Management*, 291: 13–19. <https://doi.org/10.1016/j.foreco.2012.12.001>
- Vítková, M., Müllerová, J., Sádlo, J., Pergl, J. & Pyšek, P.** 2017. Black locust (*Robinia pseudoacacia*) beloved and despised: A story of an invasive tree in Central Europe. *Forest Ecology and Management*, 384: 287–302. <https://doi.org/10.1016/j.foreco.2016.10.057>
- Voigtlaender, M., Brandani, C., Caldeira, D., Tardy, F., Bouillet, J.-P., Gonçalves, J.L.M., Moreira, M.Z., Leite, F.P., Brunet, D. & Paula, R.R.** 2019. Nitrogen cycling in monospecific and mixed-species plantations of *Acacia mangium* and *Eucalyptus* at 4 sites in Brazil. *Forest Ecology and Management*, 436: 56–67. <https://doi.org/10.1016/j.foreco.2018.12.055>
- Voigtlaender M., Laclau, J.-P., Gonçalves, J.L.M., Piccolo, M.C., Moreira, M.Z., Nouvellon, Y., Ranger, J. & Bouillet, J.-P.** 2012. Introducing *Acacia mangium* trees in *Eucalyptus grandis* plantations: consequences for soil organic matter stocks and nitrogen mineralization. *Plant and Soil*, 352: 99–111. <https://doi.org/10.1007/s11104-011-0982-9>
- Warlo, H., von Wilpert, K., Lang, F. & Schack-Kirchner, H.** 2019. Black Alder (*Alnus glutinosa* (L.) Gaertn.) on compacted skid trails: a trade-off between greenhouse gas fluxes and soil structure recovery? *Forests*, 10: 726. <https://doi.org/10.3390/f10090726>
- Werner, C., Kiese, R. & Butterbach-Bahl, K.** 2007. Soil-atmosphere exchange of N<sub>2</sub>O, CH<sub>4</sub>, and CO<sub>2</sub> and controlling environmental factors for tropical rain forest sites in western Kenya. *Journal of Geophysical Research Atmospheres*, 112: D03308. <https://doi.org/10.1029/2006JD007388>
- Yamashita, N., Ohta, S. & Hardjono, A.** 2008. Soil changes induced by *Acacia mangium* plantation establishment: Comparison with secondary forest and *Imperata cylindrica* grassland soils in South Sumatra, Indonesia. *Forest Ecology and Management*, 254: 362–370. <https://doi.org/10.1016/j.foreco.2007.08.012>

**Yang, L., Liu, N., Ren, H. & Wang, J.** 2009. Facilitation by two exotic Acacia: *Acacia auriculiformis* and *Acacia mangium* as nurse plants in South China. *Forest Ecology and Management*, 257: 1786–1793. <https://doi.org/10.1016/j.foreco.2009.01.033>

**Zagatto M.R.G., Oliveira Filho, L.C.I., Pompeo, P.N., Niva C.C., Baretta, D. & Bran Nogueira Cardoso, E.J.** 2020. Mesofauna and macrofauna in soil and litter of mixed plantations. In Bran Nogueira Cardoso, E., Gonçalves, J., Balieiro, F. & Franco, A. (Eds.) *Mixed plantations of Eucalyptus and leguminous trees*. pp.155-172. Springer, Cham. [https://doi.org/10.1007/978-3-030-32365-3\\_8](https://doi.org/10.1007/978-3-030-32365-3_8)

**Zhang, H., Duan, H., Song, M. & Guan, D.** 2018. The dynamics of carbon accumulation in Eucalyptus and Acacia plantations in the Pearl River delta region. *Annals of Forest Science*, 75(2): 40. <https://doi.org/10.1007/s13595-018-0717-7>

**Zhang, W., Zhu, X., Luo, Y., Rafique, R., Chen, H., Huang, J. & Mo, J.** 2014. Responses of nitrous oxide emissions to nitrogen and phosphorus additions in two tropical plantations with N-fixing vs. non- N-fixing tree species. *Biogeosciences*, 11: 4941–4951. <https://doi.org/10.5194/bg-11-4941-2014>

**Zou, X.** 1993. Species effects on earthworm density in tropical tree plantations in Hawaii. *Biology and Fertility of Soils*, 15: 35–38. <https://doi.org/10.1007/BF00336285>



## 5. Forest fertilization

Cindy E. Prescott, Sue J. Grayston

Faculty of Forestry, University of British Columbia, Vancouver, Canada

### 1. Description of the practice

Forest fertilization refers to the periodic addition of nutrients considered to limit productivity, and is employed in order to increase the production of stemwood biomass of forests. Globally, the most common nutrients added to forest are nitrogen (N) and phosphorus (P); others such as potassium (K), sulphur (S), boron (Bo), calcium (Ca) or magnesium (Mg) may be added on certain sites or when secondary deficiencies of these nutrients may develop following addition of N and P (Binkley and Fisher, 2019). Single-nutrient deficiencies occur in one-third of cases, in the remainder two or more nutrients are co-limiting or arise as secondary deficiencies (Harpole *et al.*, 2011). Nitrogen is usually applied as urea or ammonium nitrate granules, and P as triple superphosphate or rock phosphate. Organic materials such as compost or municipal biosolids or industrial residuals such as wood waste or ash are occasionally applied as fertilizers. Under intensive forest management, a suite of elements demonstrated to yield the most profitable growth responses are applied every year or few years during a rotation. Under extensive forest management regimes, single-element fertilizers (usually N) are operationally applied once or twice during a rotation (Binkley and Fisher, 2019). Fertilization of nutrient-deficient trees leads to increases in photosynthesis rates, leaf biomass, stemwood biomass and coarse-root biomass, but not fine-root biomass, which may decline. The decline in fine-root biomass is associated with declines in root exudation and allocation to root symbionts (Maier *et al.*, 2004). This has implications for SOC, as in some forests root- and mycorrhiza-derived C constitutes the majority of long-lived soil C (Clemmensen *et al.*, 2013; Sokol *et al.*, 2019). Fertilization also accelerates stand growth and self-thinning and shortens economic rotation times (Binkley and Fisher, 2019).

### 2. Range of applicability

Globally, the productivity of most forests is limited by N; native forests in both temperate and tropical regions increase growth by about 20 percent when N fertilizer is added (Binkley and Fisher, 2019; LeBauer and

Treseder, 2008). Phosphorus deficiency is common in some temperate regions, especially on highly weathered soils, and is widespread in fast-growing plantations in the tropics (Binkley and Fisher, 2019). In weathered tropical soils, fertilization with N, P and K is necessary to increase primary productivity and wood production (Gonçalves *et al.*, 2008; Laclau *et al.*, 2009, Wright *et al.*, 2018). In areas typically N-limited that receive substantial atmospheric N deposition, primary deficiencies of P may become the norm (Hedwall, Bergh and Brunet, 2017).

In boreal conifer forests, N is applied to forests on mineral soils whereas P and K are used in peatland forests. Boron may be added to forests that had been under slash and burn cultivation in the past. A typical nitrogen (N) dose is 150 kg ha<sup>-1</sup> and the growth response to this dose is 20–25 m<sup>3</sup>/ha. Most of the response occurs within 5 years and the N-fertilization effect is over in 10–12 years (Pukkala, 2017).

Addition of N to Douglas-fir forests of western North America increases growth rates by 2–4 m<sup>3</sup>/ha/yr for 8–15 years. In intensively managed pine plantations in the south-eastern U.S. average response to one-time fertilization with 225 kg N/ha plus 30 kg P/ha is 3.5 m<sup>3</sup>/ha/yr over 8 years (Binkley and Fisher, 2019). Most conifer plantations in New Zealand and Australia are fertilized at least once in a rotation. Australian plantations of eucalypts and pine, are typically fertilized with N (50–70 kg/ha), P (25–40 kg/ha), and sometimes, K, S, copper and zinc, at planting, early-rotation, and mid-rotation (Binkley and Fisher, 2019). Average growth response is 3–10 m<sup>3</sup>/ha/yr. Growth responses after fertilization with N, P, and K in South Africa are commonly 6–8 m<sup>3</sup>/ha/yr (Binkley and Fisher, 2019).

Almost all tropical forest plantations are fertilized with P, and most are also fertilized with other elements including N, Ca, Mg, and K. Typical growth responses to fertilizer additions in Brazilian eucalypt plantations are 4–8 m<sup>3</sup>/ha/yr for five years or longer (Gonçalves *et al.*, 1997).

### 3. Impact on soil organic carbon stocks

Increased stocks of SOC are often encountered in forests amended with nitrogenous fertilizers or simulated N deposition. In a meta-analysis of the impacts of elevated N inputs (including N fertilisation) on the storage of C in forest soils based on 72 experimental sites, Nave *et al.* (2009) found that N inputs increased total SOC stocks (combined forest floor and mineral soil) by 7.7 percent. Stocks of organic C increased predominantly in the mineral soil (by 12.2 percent). In the review by Johnson and Curtis (2001), N fertilization was the only forest management practice that had a clear positive effect on the SOC pool. Greater accumulations of humus are often noted following N fertilization of boreal forests (Nohrstedt, 1990; Mälikönen and Kukkola, 1991; Mäkipää, 1995; Olsson *et al.*, 2005) and simulated N deposition in temperate forests (Lovett *et al.*, 2013), as have greater SOC concentrations in mineral soil (Pregitzer *et al.*, 2008; Cusack *et al.*, 2011; Huang *et al.*, 2011). Added N may also increase accumulation of C in occluded particulate organic matter (Zak *et al.*, 2017). Increased mineral-associated C stocks in response to N fertilization have also been reported in temperate and tropical forests, even in the absence of a detectable increase in bulk SOC (Hagedorn, Spinnier and Siegwolf, 2003; Cusack *et al.*, 2011).

**Table 13.** Changes in soil organic carbon stocks reported for nutrient additions

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Years)	Depth (cm)	Fertilization rates	Reference
Sweden and Finland	Boreal	NA	NA	0.1 - 1.2	14-30	O horizon + O -10	Meta-analysis; Repeated additions of N or NPK; Cumulative amount 600-2340 kgN/ha	Hyvönen <i>et al.</i> (2008)
Sweden		NA	32.3	0.12 - 0.25	18-21	O horizon	-	Nohrstedt <i>et al.</i> (1989)
Massachusetts, United States of America	Cool temperate	Typic Dystrudepts	1.3	0.28	20 N dep	0-40	50 and 150 kgN/ha/yr	Frey <i>et al.</i> (2014)
New York State, United States of America		Lithic Dystrachrepts	NA	0.3-1.3	6	O horizon	50 kgN/ha/yr	Lovett <i>et al.</i> (2013)
Washington, United States of America		Vitrixerands, Hapludands, Dystraxept	175	3.7	21-29	0-100	672-896 kgN/ha in 3 or 4 additions	Adams <i>et al.</i> (2005)
Pennsylvania, United States of America		Alfisols	0.5	0.42	22 N dep	14-21	100 kgN/ha/yr	Wang <i>et al.</i> (2019)
Michigan, United States of America		Typic haplorthod	1.8	0.7	10 N dep	0-70	30 kgN/ha/yr	Pregitzer <i>et al.</i> (2007)
China	Subtropical moist	Oxisols	NA	1.5 (50 kgN/yr) 1.3 (100kgN/yr) 0.9 (150kgN/yr)	14 N dep	0-50	50-150 kgN/ha/yr	Yu <i>et al.</i> (2020)
Puerto Rico	Tropical wet	Ultisols and Oxisols	0.9	0.005	7	0-25	300 kgN/ha/yr + PK	Li, Xu and Zou, (2006)
			79	5.7	3 N dep	0-40	50 kgN/ha/yr	Cusack <i>et al.</i> (2011)

## 4. Other benefits of the practice

### 4.1 Minimization of threats to soil functions

Table 14. Soil threats

Soil threats	
Nutrient imbalance and cycles	Addition of specific nutrients can balance overall nutrient supply.
Soil acidification	Urea can be applied to increase pH; ammonium nitrate to reduce pH.

### 4.2 Increases in production (e.g. food/fuel/feed/timber)

Intensive forest management, including fertilization, increases the rate of stemwood production and reduces the economic rotation time for the stand. In the southeast U.S., the average productivity of commercial *Pinus taeda* plantations is more than 4-fold higher than of natural stands, and 17 percent of this gain has been attributed to fertilizer amendments (Noormets *et al.*, 2015). The productivity of eucalypts in Brazil has nearly doubled in the last 2 decades, owing to intensive management techniques, including fertilization (Goncalves *et al.*, 2013). Average gains from fertilization in other types of plantations are given in Section 2. In addition to the value of the timber to the landholder, the potential to produce timber faster means that less land is needed to supply the same amount of timber. This could increase the amount of land available for other purposes such as agriculture, recreation or conservation.

Forest fertilization may alter the understory vegetation towards a flora indicative of more fertile sites – increases of grasses and herbs, declines in some bryophytes, lichens and dwarf shrubs. As such fertilization can improve forage production and quality for livestock in forest grazing or silvopastoral systems. Forest fertilization can also increase browse quality, which increases populations of herbivore and their predators, and can also lead to tree damage (Sullivan and Sullivan, 2018).

### 4.3 Mitigation of and adaptation to climate change

The production, transport and application of chemical fertilizers all emit greenhouse gases. However, the additional biomass produced following fertilization sequesters CO<sub>2</sub> equivalents greater than those emitted by at least an order of magnitude. Fertilization of mid-rotation pine stands in the southeastern U.S. with N and P sequestered 6.2 million tons of CO<sub>2</sub> in additional stem growth whereas the CO<sub>2</sub> equivalents corresponding to fertilizer production, transport, and application were 231,000 tons (Albaugh *et al.*, 2019). In a short-rotation Eucalyptus plantation in Brazil, for every 1 kg of CO<sub>2</sub>-eq. emitted, 43.15 kg of CO<sub>2</sub>-eq. were sequestered in

the (utilized) biomass (Quartucci, Schweier and Jaeger, 2015). The net positive effects of fertilization on overall C balance may be even larger if one factors in 1) the additional carbon sequestered in coarse root, branch, and foliage growth, 2) changes in soil CO<sub>2</sub> evolution and ecosystem carbon storage, 3) reductions in rotation length that increase carbon sequestration in the system, 4) increased tree size, which increases the proportion of dimension lumber that retain the sequestered carbon in the system for a longer time than smaller size trees, which may be used to make pulp and paper (Albaugh *et al.*, 2012), and 5) if the additional growth is used for bioenergy so substitutes for coal, oil and natural gas (Bergh and Hedwall 2013). For example, fertilizing 10 percent of Swedish forest land could reduce annual GHG emission by 11.9 million or 18.1 million t CO<sub>2</sub>eq if the reference fossil fuel is fossil gas or coal, respectively (Sathre, Gustavsson and Bergh, 2010).

As forest fertilization can be applied within present management systems and will give immediate effects, it is probably the most efficient tool of forest management to affect the carbon cycle, at least in a short term. The increased amount of biomass can be used as a growing carbon stock or to replace fossil fuels (Hedwall *et al.*, 2014).

## 4.4 Socio-economic benefits

The economic benefit of forest fertilization to the landowner depends on the magnitude of response, its predictability, the time required before the benefit is realized (rotation length), and the risk of forest loss or damage during the rotation. Forest fertilization can offer socio-economic benefits on a national level – for example an assessment of the potential benefits of forest fertilization in Latvia, applying 436 kgN/ha to 2000 ha of forests annually would generate an additional 142 000 m<sup>3</sup> of wood. After 10 years this would generate 3.2 million USD per year and contribute to CO<sub>2</sub>-eq sequestration of 12.7 million (Lazdiņš *et al.*, 2018).

# 5. Potential drawbacks to the practice

## 5.1 Tradeoffs with other threats to soil functions

Table 15. Soil threats

Soil threats	
<b>Nutrient imbalance and cycles</b>	Growth response from addition of one nutrient can create deficiency of another (Binkley and Fisher 2019). Chronic addition of N could reduce or alter the cycling of other nutrients (Wang <i>et al.</i> , 2019).
<b>Soil acidification</b>	nitrification induced by N fertilization may acidify soil and soil solution Alters pH depending on formulation and dosage (Binkley and Fisher, 2019)
<b>Soil biodiversity loss</b>	Fertilization usually reduces belowground C fluxes and microbial biomass and alters the saprotrophic community (Noormets <i>et al.</i> , 2015).

<b>Soil threats</b>	
	N and P fertilization reduce proportional allocation to roots and root symbionts (Maier <i>et al.</i> , 2004), and often reduces soil microbial biomass and activity (Högberg <i>et al.</i> , 2003) and changes soil microbial community (Morrison <i>et al.</i> , 2016).
<b>Soil water management</b>	Plant growth response to fertilization can increase water demand (Ward <i>et al.</i> , 2015). Increase in leaf biomass from fertilization can increase water usage and loss via transpiration (Ward <i>et al.</i> , 2015). Addition of more N or P than soils can retain risks eutrophication of waterbodies (Laclau <i>et al.</i> , 2010).


## 5.2 Increases in greenhouse gas emissions

Fertilization may have detrimental effects on GHG emissions (Liu and Greaver, 2009). Forest soils are sinks for 5 percent of global annual methane emissions, through methane oxidation (Schlesinger, 1997) and potential sources, or sinks for nitrous oxide through microbial nitrification and denitrification (Robertson and Groffman, 2007). Methane has 25 times, and N<sub>2</sub>O has 298 times the global warming potential of CO<sub>2</sub> (Forster *et al.*, 2007). Although N addition increases the terrestrial C sink in forests, the CO<sub>2</sub> reduction may be offset by N stimulation of N<sub>2</sub>O emissions and inhibition of CH<sub>4</sub> oxidation.

Fertilization effects on GHG fluxes are complex and vary with timing and chemistry of fertilizers, environmental variables and ecosystem properties (Figure 2). N fertilization can potentially increase N<sub>2</sub>O emissions through microbial nitrification and denitrification (Levy-Booth, Prescott and Grayston, 2014). N fertilization can either stimulate CH<sub>4</sub> oxidation by supplying a limiting nutrient (N) to CH<sub>4</sub>-oxidizing bacteria (Bodelier and Laanbroek, 2004) or by supplying NH<sub>4</sub> to ammonia-oxidizing bacteria that can also oxidize CH<sub>4</sub> (Jiang and Bakken, 1999). N fertilization can also inhibit CH<sub>4</sub> oxidation due to NH<sub>4</sub> competing with CH<sub>4</sub> for active sites on the methane monooxygenase enzyme (Dunfield and Knowles, 1995), or through suppression of methane monooxygenase enzyme due to alternative labile C sources being available to CH<sub>4</sub> oxidizers in N-amended soils (Fender *et al.*, 2012).

In a meta-analysis of the effects of N addition on the flux of three major GHGs: CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O (Liu and Greaver, 2009), N addition increased ecosystem carbon content of forests by 6 percent. Across all ecosystems, N addition increased CH<sub>4</sub> emission by 97 percent, reduced CH<sub>4</sub> uptake by 38 percent and increased N<sub>2</sub>O emission by 216 percent. Nitrogen addition ranging from 25 to 200 kg N/ha/yr applied to forest ecosystems for 6–15 years increased ecosystem carbon by an average of 6 percent. Coniferous and deciduous forests with chronic anthropogenic N inputs show 30–50 percent reduced CH<sub>4</sub> uptake and 2–3-fold increases in N<sub>2</sub>O emissions. Nitrogen enrichment of tropical forests, in particular, increased N<sub>2</sub>O emissions by 739 percent. Contrastingly, in unpolluted temperate coniferous forests, N fertilization had no effects on GHG fluxes (Basiliko *et al.*, 2009).

In addition, volatilization of  $\text{NH}_3$  from surface-applied urea may occur when soils are moist and relative humidity and air temperatures are high (Elliot and Fox, 2014). Enhanced efficiency fertilizers have been developed to minimize losses through  $\text{NH}_3$  volatilization (Raymond *et al.*, 2016).



**Figure 2.** The potential mechanisms that regulate the responses of  $\text{CO}_2$ ,  $\text{CH}_4$  and  $\text{N}_2\text{O}$  production and consumption to elevated N. ANPP, aboveground net primary productivity; BNPP, belowground net primary productivity; SOC, soil organic carbon; DOC, dissolved organic carbon; DIN, dissolved inorganic nitrogen; DON, dissolved organic nitrogen. Source: adapted from Liu and Greaver (2009)

### 5.3 Conflict with other practice(s)

Fertilizers are expensive so the growth response and consequent economic benefits of fertilization versus other stand-tending practices must be predictable. Fertilization may stimulate growth of non-crop vegetation and necessitate vegetation control to realize the full benefit of fertilization (Binkley and Fisher, 2019).

### 5.4 Decreases in production (e.g. food/fuel/feed/timber)

Fertilization can intensify drought stress (Ward *et al.*, 2015) as a consequence of greater water demand and lower proportion of fine roots (Noormets *et al.*, 2015), especially in heavily fertilized plantations (Battie-Laclau *et al.*, 2014).

### 5.5 Other conflicts

The potential benefits of N fertilization for increasing tree growth and SOC stocks must be weighed against the associated environmental costs, as the production, transport and application of synthetic fertilizers all entail fossil fuel combustion and emission of CO<sub>2</sub>.

There may be public concerns over potential eutrophication of drainage waters, although N fertilizer in forestry generally leads to small and transient increases in N concentrations in stream water (Binkley, Burnham and Allen, 1999; Smethurst, 2010). Leaching losses can be reduced by applying fertilizer in small doses rather than a single large application (Bergh *et al.*, 2008; Albaugh *et al.*, 2019).

## 6. Recommendations before implementation of the practice

Addition of N is not advisable in ecosystems where N is plentiful due to atmospheric N deposition or previous agricultural use, as it could stimulate losses as nitrate in water or as greenhouse gases through denitrification (Gao *et al.*, 2015).

Site water balance must be considered to prevent drought stress caused by increased leaf biomass and productivity without concomitant increases in root biomass.

In degraded soils with low organic matter and clay contents, N would be better added in organic forms such as composts or municipal biosolids that would increase retention of the N in the soil (Larney and Angers, 2012).

Investment in research is necessary to predict growth response to addition of nutrients. Across a landscape this involves determining relationships between growth response and site factors, soil or foliar nutrient concentrations, and remotely sensed factors such as leaf area index (LAI) and normalized difference vegetation index (NDVI) (Blinn *et al.*, 2019).



Possible effects of N addition on the fluxes of CH<sub>4</sub> and N<sub>2</sub>O must be given careful consideration.

Given that equivalent SOC gains are achievable through incorporation of N-fixing tree species (Nave *et al.* 2010), this may be preferable to chemical fertilizers in many situations.

## 7. Potential barriers to adoption

**Table 16.** Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	In degraded or coarse-textured soils there may be insufficient retention of added fertilizer, and organic fertilizers would be more effective.
Cultural	Yes	N is popularly viewed negatively as a pollutant due to association with atmospheric N deposition. Both N and P are associated with degradation of water quality resulting from excessive applications in agriculture.
Social	Yes	May be public concerns about effects on water quality and biodiversity, also association of chemical fertilizers with fossil-fuel use.
Economic	Yes	Cost of fertilizer mandates good ability to predict growth response, which requires research into relationships with site factors and leaf area (Albaugh <i>et al.</i> , 2019).
Knowledge	Yes	Belowground effects of nutrient addition need to be better understood before effects on SOC can be predicted (Noormets <i>et al.</i> , 2015)

## References

- Adams, A.B., Harrison, R.B., Sletten, R.S., Strahm, B.D., Turnblom, E.C. & Jensen, C.M. 2005. Nitrogen-fertilization impacts on carbon sequestration and flux in managed coastal Douglas-fir stands of the Pacific Northwest. *Forest Ecology and Management*, 220(1-3): 313-325.  
<https://doi.org/https://doi.org/10.1016/j.foreco.2005.08.018>
- Albaugh, T.J., Fox, T.R., Cook, R.L., Raymond, J.E., Rubilar, R.A. & Campoe, O.C. 2019. Forest fertilizer applications in the southeastern United States from 1969 to 2016. *Forest Science*, 65(3): 355-362.  
<https://doi.org/10.1093/forsci/fxy058>
- Albaugh, T.J., Vance, E.D., Gaudreault, C., Fox, T.R., Allen, H.L., Stape, J.L. & Rubilar, R.A. 2012. Carbon emissions and sequestration from fertilization of pine in the southeastern United States. *Forest Science*, 58: 419-429. <https://doi.org/10.5849/forsci.11-050>
- Basiliko, N., Khan, A., Prescott, C.E., Roy, R. & Grayston, S.J. 2009. Soil greenhouse gas and nutrient dynamics in fertilized western Canadian plantation forests. *Canadian Journal of Forest Research*, 39(6): 1220-1235. <https://doi.org/10.1139/X09-043>
- Battie-Laclau, P., Laclau, J.P., Domec, J.C., Christina, M., Bouillet, J.P., Piccolo, M.C., Gonçalves, J.L.M., Moreira, R.M., Krusche, A.V., Bouvet, J.M. & Nouvellon, Y. 2014. Effects of potassium and sodium supply on drought-adaptive mechanisms in *Eucalyptus grandis* plantations. *New Phytologist*, 203: 401-413. <https://doi.org/10.1111/nph.12810>
- Bergh, J. & Hedwall, P.O. 2013. Fertilization in boreal and temperate forests and the potential for biomass production. In: S. Kellomäki, A. Kilpeläinen & A. Alam, eds. *Forest BioEnergy Production*. Springer, New York, NY.
- Binkley, D., Burnham, H. & Allen, H.L. 1999. Water quality impacts of forest fertilization with nitrogen and phosphorus. *Forest Ecology and Management*, 121: 191-213.
- Binkley, D. & Fisher, R.F. 2019. *Ecology and Management of Forest Soils*, 5th Edition. Wiley-Blackwell, New York, NY. 456 Pages. ISBN: 978-1-119-45565.
- Blinn, C.E., House, M.N., Wynne, R.H., Thomas, V.A., Fox, T.R. & Sunnall, M. 2019. Landsat 8 based leaf area index estimation in loblolly pine plantations. *Forests*, 10(3): 222.  
<https://doi.org/10.3390/f10030222>
- Bodelier, P.L.E. & Laanbroek, H.J. 2004. Nitrogen as a regulatory factor of methane oxidation in soils and sediments. *FEMS Microbiology Ecology*, 47: 265-277. [https://doi.org/10.1016/S0168-6496\(03\)00304-0](https://doi.org/10.1016/S0168-6496(03)00304-0)
- Clemmensen, K.E., Bahr, A., Ovaskainen, O., Dahlberg, A., Ekblad, A., Wallander, H., Stenlid, J., Finlay, R.D., Wardle, D.A. & Lindahl, B.D. 2013. Roots and associated fungi drive long-term carbon sequestration in boreal forest. *Science*, 339: 1615-1618. <https://doi.org/10.1126/science.1231923>
- Cusack, D.F., Silver, W.L., Torn, M.S. & McDowell, W.H. 2011. Effects of nitrogen additions on above- and belowground carbon dynamics in two tropical forests. *Biogeochemistry*, 104: 203-225.

- Dunfield, P. & Knowles, R.** 1995. Kinetics of inhibition of methane oxidation by nitrate, nitrite, and ammonium in a humisol. *Applied Environmental Microbiology*, 61: 3129–3135. PMID:16535109.
- Elliot, J.R. & Fox, T.R.** 2014. Ammonia volatilization following fertilization with urea or ureaform in a thinned loblolly pine plantation. *Soil Science Society of America Journal*, 78(4): 1469–1473.  
<https://doi.org/10.2136/sssaj2013.12.0512n>
- Fender, A., Pfeiffer, B., Gansert, D., Leuschner, C., Daniel, R. & Jungkunst, H.F.** 2012. The inhibiting effect of nitrate fertilisation on methane uptake of a temperate forest soil is influenced by labile carbon. *Biology and Fertility of Soils*, 48(6): 621–631. <https://doi.org/10.1007/s00374-011-0660-3>
- Forster, P., Ramaswamy, V., Artaxo, P., Berntsen, T., Betts, R., Fahey, D.W., Haywood, J., Lean, J., Lowe, D.C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M. & Van Dorland, R.** 2007. Changes in atmospheric constituents and in radiative forcing. In S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, M. Averyt, M. Tignor & M. Miller (Eds.) *Climate change 2007: the physical science basis*. Contribution of Working Group I to the 4th Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK.
- Frey, S.D., Ollinger, S., Nadelhoffer, K., Bowden, R., Brzostek, E., Burton, A., Caldwell, B.A., Crow, S., Goodale, C.L., Grandy, A.S., Finzi, A., Kramer, M.G., Lajtha, K., LeMoine, J., Martin, M., McDowell, W.H., Minocha, R., Sadowsky, J.J., Templer, P.H. & Wickings, K.** 2014. Chronic nitrogen additions suppress decomposition and sequester soil carbon in temperate forests. *Biogeochemistry*, 121: 305–316. <https://doi.org/10.1007/s10533-014-0004-0>
- Gao, W., Yang, H., Kou, L. & Li, S.** 2015. Effects of nitrogen deposition and fertilization on N transformations in forest soils: a review. *Journal of Soils and Sediments*, 15: 863–879.  
<https://doi.org/10.1007/s11368-015-1064-z>
- Gonçalves, J.L.M., Stape, J.L., Laclau, J.-P., Bouillet, J.-P. & Ranger, J.** 2008. Assessing the effects of early silvicultural management on long-term site productivity of fast-growing eucalypt plantations: the Brazilian experience. *Southern Forests*, 70: 105–118.  
<https://doi.org/10.2989/SOUTH.FOR.2008.70.2.6.534>
- Gonçalves, J.L.M., Barros, N.F., Nambiar, E.K.S. & Novais, R.F.** 1997. Soil and stand management for short-rotation plantations. In E.K.S. Nambiar and A.G. Brown (Eds.) *Management of Soil, Nutrients, and Water in Tropical Plantation Forests*. ACIAR Monograph #43. Canberra: ACIAR. pp. 379–417.
- Hagedorn, F., Spinnler, D. & Siegwolf, R.** 2003. Increased N deposition retards mineralization of old soil organic matter. *Soil Biology & Biochemistry*, 35: 1683–1692.
- Harpole, W.S., Ngai, J.T., Cleland, E.E., Seabloom, E.W., Borer, E.T., Bracken, M.E.S., Elser, J., Gruner, D.S., Hillebrand, H., Shurin, J.B. & Smith, J.E.** 2011. Nutrient co-limitation of primary producer communities. *Ecology Letters*, 14: 852–862. <https://doi.org/10.1111/j.1461-0248.2011.01651.x>
- Hedwall P.-O., Gong, P., Ingerslev, M. & Bergh, J.** 2014. Fertilization in northern forests – biological, economic and environmental constraints and possibilities. *Scandinavian Journal of Forest Research*, 29(4): 301–311. <https://doi.org/10.1080/02827581.2014.926096>

- Hedwall, P. O., Bergh, J. & Brunet, J.** 2017. Phosphorus and nitrogen co-limitation of forest ground vegetation under elevated anthropogenic nitrogen deposition. *Oecologia*, 185(2): 317–326. <https://doi.org/10.1007/s00442-017-3945-x>
- Högberg, M.N., Bååth, E., Nordgren, A., Arnebrant, K. & Högberg, P.** 2003. Contrasting effects of nitrogen availability on plant carbon supply to mycorrhizal fungi and saprotrophs – a hypothesis based on field observations in boreal forest. *New Phytologist*, 160: 225–238. <https://doi.org/10.1046/j.1469-8137.2003.00867.x>
- Huang, Z., Clinton, P.W., Baisden, W.T. & Davis, M.R.** 2011. Long-term nitrogen additions increased surface soil carbon concentration in a forest plantation despite elevated decomposition. *Soil Biology & Biochemistry*, 43: 302–307. <https://doi.org/10.1016/j.soilbio.2010.10.015>
- Hyvönen, R., Persson, T., Andersson, S., Olsson, B., Ågren, G.I. & Linder, S.** 2008. Impact of long-term nitrogen addition on carbon stocks in trees and soils in northern Europe. *Biogeochemistry*, 89: 121–137. <https://doi.org/10.1007/s10533-007-9121-3>
- Jiang, Q.Q. & Bakken, L.R.** 1999. Nitrous oxide production and methane oxidation by different ammonia-oxidizing bacteria. *Applied Environmental Microbiology*, 65: 2679–2684. <https://doi.org/10.1128/AEM.65.6.2679-2684.1999>
- Johnson, D. & Curtis, P.** 2001. Effects of forest management on soil C and N storage: meta-analysis. *Forest Ecology and Management*, 140: 227–238. [https://doi.org/10.1016/S0378-1127\(00\)00282-6](https://doi.org/10.1016/S0378-1127(00)00282-6)
- Laclau, J.-P., Almeida, J.C.R., Goncalves, J.L.M., Saint-Andre, L., Ventura, M., Ranger, J., Moreira, R.M. & Nouvellon, Y.** 2009. Influence of nitrogen and potassium fertilization on leaf lifespan and allocation of above-ground growth in Eucalyptus plantations. *Tree Physiology*, 29: 111–124. <https://doi.org/10.1093/treephys/tpn010>
- Laclau, J.-P., Ranger, J., Gonçalves, J.L.M., Maquère, V., Krusche, A.V., M'Bou, A.T., Nouvellon, Y., Saint-André, L., Bouillet, J.-P., Piccolo, M.C. & Deleporte, P.** 2010. Biogeochemical cycles of nutrients in tropical Eucalyptus plantations: Main features shown by intensive monitoring in Congo and Brazil. *Forest Ecology and Management* 259:1771–1785. <https://doi.org/10.1016/j.foreco.2009.06.010>
- Larney, F.J. & Angers, D.A.** 2012. The role of organic amendments in soil reclamation: a review. *Canadian Journal of Soil Science*, 92: 19–38. <https://doi.org/10.4141/cjss2010-064>
- Lazdiņš, A., Okmanis, M., Makovskis, K. & Petaja, G.** 2018. Forest fertilization: Economic effect and impact on GHG emissions in Latvia. *Baltic Forestry*, 24(1):9–16.
- LeBauer, D.S. & Treseder, K.K.** 2008. Nitrogen limitation of net primary productivity in terrestrial ecosystems is globally distributed. *Ecology*, 89: 371–379. <https://doi.org/10.1890/06-2057.1>
- Levy-Booth, D.J., Prescott, C.E. & Grayston, S.J.** 2014. Microbial functional genes involved in nitrogen fixation, nitrification and denitrification in forest ecosystems. *Soil Biology & Biochemistry*, 75: 11–25. <https://doi.org/10.1016/j.soilbio.2014.03.021>

- Li, Y., Xu, M. & Zou, X.** 2006. Effects of nutrient additions on ecosystem carbon cycle in a Puerto Rican tropical wet forest. *Global Change Biology*, 12: 284-293. <https://doi.org/10.1111/j.1365-2486.2005.01096.x>
- Liu, L. & Greaver, T.L.** 2009. A review of nitrogen enrichment effects on three biogenic GHGs: the CO<sub>2</sub> sink may be largely offset by stimulated N<sub>2</sub>O and CH<sub>4</sub> emission. *Ecology Letters*, 12: 1103–1117. <https://doi.org/10.1111/j.1461-0248.2009.01351.x>
- Lovett, G.M., Arthur, M.A., Weathers, K.C., Fitzhugh, R.D. & Templer, P.H.** 2013. Nitrogen addition increases carbon storage in soils, but not in trees, in an eastern U.S. deciduous forest. *Ecosystems*, 16: 980–1001. <https://doi.org/10.1007/s10021-013-9662-3>
- Maier, C.A., Albaugh, T.J., Allen, H.L. & Dougherty, P.M.** 2004. Respiratory carbon use and carbon storage in mid-rotation loblolly pine (*Pinus taeda* L.) plantations: the effect of site resources on the stand carbon balance. *Global Change Biology*, 10: 1335–1350. <https://doi.org/10.1111/j.1529-8817.2003.00809.x>
- Mäkipää, R.** 1995. Effect of nitrogen input on carbon accumulation of boreal forest soils and ground vegetation. *Forest Ecology and Management*, 79: 217–226. [https://doi.org/10.1016/0378-1127\(95\)03601-6](https://doi.org/10.1016/0378-1127(95)03601-6)
- Mälkönen, E. & Kukkola, M.** 1991. Effect of long-term fertilization on the biomass production and nutrient status of Scots pine stands. *Fertilizer Research*, 27: 113–127. <https://doi.org/10.1007/BF01048614>
- Morrison, E.W., Frey, S.D., Sadowsky, J.J., van Diepen, L.T.A., Thomas, W.K. & Pringle, A.** 2016. Chronic nitrogen additions fundamentally restructure the soil fungal community in a temperate forest. *Fungal Ecology*, 23:48–57. <https://doi.org/10.1016/j.funeco.2016.05.011>
- Nave, L., Vance, E., Swanston, C. & Curtis, P.** 2009. Impacts of elevated N inputs on north temperate forest soil C storage, C/N, and net N-mineralization. *Geoderma*, 153: 231–240. <https://doi.org/10.1016/j.geoderma.2009.08.012>
- Nohrstedt, H.-Ö., Arnebrant, K., Bååth, E. & Söderström, B.** 1989. Changes in carbon content, respiration rate, ATP content, and microbial biomass in nitrogen-fertilized pine forest soils in Sweden. *Canadian Journal of Forest Research*, 19(3): 323–328. <https://doi.org/10.1139/x89-048>
- Nohrstedt, H.Ö.** 1990. Effects of repeated nitrogen fertilization with different doses on soil properties in a *Pinus sylvestris* stand. *Scandinavian Journal of Forest Research*, 5(1-4): 3–15. <https://doi.org/10.1080/02827589009382588>
- Noormets, A., Epron, D., Domec, J.C., McNulty, S.G., Fox, T., Sun, G. & King, J.S.** 2015. Effects of forest management on productivity and carbon sequestration: A review and hypothesis. *Forest Ecology and Management*, 355: 124–140. <https://doi.org/10.1016/j.foreco.2015.05.019>
- Olsson, P., Linder, S., Giesler, R. & Högberg, P.** 2005. Fertilization of boreal forest reduces both autotrophic and heterotrophic soil respiration. *Global Change Biology*, 11: 1745–1753. <https://doi.org/10.1111/j.1365-2486.2005.001033.x>

- Pregitzer, K.S., Burton, A.J., Zak, D.R. & Talhelm, A.F.** 2008. Simulated chronic nitrogen deposition increases carbon storage in Northern Temperate forests. *Global Change Biology*, 14: 1-12.  
<https://doi.org/10.1111/j.1365-2486.2007.01465.x>
- Pukkala, T.** 2017. Optimal nitrogen fertilization of boreal conifer forest. *Forest Ecosystems*, 4(3).  
<https://doi.org/10.1186/s40663-017-0090-2>
- Quartucci, F., Schweier, J. & Jaeger, D.** 2015. Environmental analysis of Eucalyptus timber production from short rotation forestry in Brazil. *International Journal of Forest Engineering*, 26(3): 1-15.  
<https://doi.org/10.1080/14942119.2015.1099813>
- Raymond, J.E., Fox, T.R., Strahm, B.D. & Zerpa, J.** 2016. Ammonia volatilization following nitrogen fertilization with enhanced efficiency fertilizers and urea in loblolly pine (*Pinus taeda* L.) plantations of the southern United States. *Forest Ecology and Management*, 376: 247–255.  
<https://doi.org/https://doi.org/10.1016/j.foreco.2016.06.015>
- Robertson, G.P. & Groffman, P.M.** 2007. Nitrogen transformations. In E.A. Paul (Ed.) *Soil Biology and Biochemistry*. 3<sup>rd</sup> ed. Academic Press, Oxford, UK. pp. 341–364.
- Sathre, R., Gustavsson, L. & Bergh, J.** 2010. Primary energy and greenhouse gas implications of increasing biomass production through forest fertilization. *Biomass and Bioenergy*, 34(4): 572-581.  
<https://doi.org/https://doi.org/10.1016/j.biombioe.2010.01.038>
- Schlesinger, W.H.** 1997. *Biogeochemistry: An Analysis of Global Change*. 2<sup>nd</sup> ed. Academic Press, San Diego, Calif.
- Smethurst, P.J.** 2010. Forest fertilization: trends in knowledge and practice compared to agriculture. *Plant and Soil*, 335: 83–100. <https://doi.org/10.1007/s11104-010-0316-3>
- Sokol, N.W., Kuebbing, S.E., Karlsen-Ayala, E. & Bradford, M.A.** 2019. Evidence for the primacy of living root inputs, not root or shoot litter, in forming soil organic carbon. *New Phytologist*, 221: 233-246.  
<https://doi.org/10.1111/nph.15361k>
- Sullivan, T.P. & Sullivan, D.S.** 2017. Influence of nitrogen fertilization on abundance and diversity of plants and animals in temperate and boreal forests. *Environmental Reviews*. <https://doi.org/10.1139/er-2017-0026>
- Wang, J.-J., Bowden, R.D., Lajtha, K., Washko, S.E., Wurzbacher, S.J. & Simpson, M.J.** 2019. Long-term nitrogen addition suppresses microbial degradation, enhances soil carbon storage, and alters the molecular composition of soil organic matter. *Biogeochemistry*, 142: 299–313.  
<https://doi.org/10.1007/s10533-018-00535-4>
- Ward, C., Pothier, D. & Paré, D.** 2014. Do boreal forests need fire disturbance to maintain productivity? *Ecosystems*, 17: 1053–1067. <https://doi.org/10.1007/s10021-014-9782-4>
- Wright, S.J., Turner, B.L., Yavitt, J.B., Harms, K.E., Kaspari, M., Tanner, E.V.J., Bujan, J., Griffin, E.A., Mayor, J.R., Pasquini, S.C., Sheldrake, M. & Garcia, M.N.** 2018. Plant responses to fertilization experiments in lowland, species- rich, tropical forests. *Ecology*, 99: 1129–1138.  
<https://doi.org/10.1002/ecy.2193>

**Yu, M., Wang, Y.-P., Baldock, J.A., Jiang, J., Mo, J., Zhou, G. & Yan, J.** 2020. Divergent responses of soil organic carbon accumulation to 14 years of nitrogen addition in two typical subtropical forests. *Science of the Total Environment*, 707: 136104. <https://doi.org/10.1016/j.scitotenv.2019.136104>

**Zak, D.R., Freedman, Z.B., Upchurch, R.A., Steffens, M. & Kögel-Knabner, I.** 2017. Anthropogenic N deposition increases soil organic matter accumulation without altering its biochemical composition. *Global Change Biology*, 23: 933–944. <https://doi.org/10.1111/gcb.13480>

# 6. Forest afforestation, reforestation and natural regeneration

Cindy E. Prescott<sup>1</sup>, Yann Nouvellon<sup>2</sup>

<sup>1</sup>Faculty of Forestry, University of British Columbia, Vancouver, Canada

<sup>2</sup>Cirad, Research Unit Eco&sols, Kasetsart University, Bangkok, Thailand

## 1. Description of the practice

Afforestation is the conversion from other land uses into forest, or the increase of the canopy cover to above 10 percent (FAO, 2000). Afforestation is the reverse of deforestation and includes areas that are actively converted from other land uses. Afforestation includes conversion to forest through silvicultural measures or natural transitions into forest, for example on abandoned agricultural land or in burnt-over areas that have not been classified as forest during the barren period. The conversion should be long-term, *i.e.* the transition into forest is expected to last more than ten years. Local climatological conditions, land use contexts or the purpose of the analysis may however justify use of a longer time frame. If the area had been temporarily deforested, re-establishment of forest is termed “reforestation” (FAO, 2000).

Afforestation may occur passively (*i.e.* with no human intervention) following land abandonment, or actively through planting of tree seedlings or sowing seeds. It may also occur via assisted natural regeneration (ANR), in which specific actions are taken to reduce barriers to natural forest regeneration, such as control of grazing or weed competition, amelioration of soil or microclimate conditions or seed dispersal.

## 2. Range of applicability

Afforestation (and reforestation) is applicable in locations where the climate dictates that the natural vegetation would be forest (see Hotspot: Forests). Afforestation is not recommended in areas that would naturally be grassland (Silveira *et al.*, 2020) or other non-forest biomes.



### 3. Impact on soil organic carbon stocks


Afforestation increases SOC content especially of soils that have been degraded through mineral extraction or unsustainable cropping (Table 17). The rate of SOC accumulation is greatest in soils that have been depleted in C (Shi *et al.*, 2013; Wang and Huang 2020). Afforestation on lands made barren by mining increased SOC stocks by 15-fold (average 173 percent; meta-analysis by Nave *et al.* 2013). SOC stocks significantly increased within 15-25 years of afforestation and continued through subsequent decades. SOC accumulation on afforested mine sites averaged 2.46 tC/ha/yr in the first 10 years and 0.87 tC/ha/yr over the first 40 years (Frouz *et al.*, 2014). Afforestation on former cropland (that is, land used for cultivation of crops; FAO 2017) may result in a significant increase in SOC stocks (Mayer *et al.* 2020). In some studies, no new steady-state levels were reached within 100 years (Poepflau *et al.*, 2011; Bárcena *et al.*, 2014) while in others, modest decadal increases culminated in a ~15 percent net increase in SOC stock by the end of the first century (Nave *et al.*, 2013). Afforestation of degraded croplands under China's Grain for Green program averaged 0.26 tC/ha/yr (Deng *et al.*, 2014). In Ethiopia, afforestation of degraded croplands with *Eucalyptus* elevated SOC stocks to nearly 70 percent of levels found in a natural dry subtropical montane forest within 30 years (Assefa *et al.*, 2017).

In contrast to cropped lands, afforestation of grasslands and pastures has little effect on SOC stocks (Guo and Gifford 2002; Mayer *et al.* 2020), and often results in SOC losses (Poepflau *et al.* 2011). SOC also accumulates faster on highly degraded soils restored to grassland compared to those afforested (Wei *et al.* 2012; Frouz *et al.*, 2014). The high stocks of SOC in grasslands and pastures have been attributed to much greater fine root length and water content than in forest soils (Berhongaray and Alvarez, 2019; Deng *et al.* 2014).

Immediately following afforestation, SOC stocks may decrease due to site disturbance, soil erosion, and low NPP and C inputs from the young vegetation (Paul *et al.*, 2002). The decline usually lasts about 10 years when sites are actively managed (Laganière, Angers and Paré, 2010; Deng *et al.*, 2014; Li *et al.*, 2012; Nave *et al.*, 2013) but can last up to 35 years following agricultural abandonment (Paul *et al.*, 2002). Following this, SOC accumulates rapidly for a few decades (Laganière, Angers and Paré, 2010; Deng *et al.*, 2014; Li *et al.*, 2012; Nave *et al.*, 2013). SOC accumulates quickly in a surface organic layer and more slowly in mineral layers (Li *et al.*, 2012; (Bárcena *et al.*, 2014; Ledo *et al.*, 2020). The incorporation rate slows (Vinduskova and Frouz, 2013; Frouz *et al.*, 2014) as the SOC content approaches saturation or equilibrium with the site conditions (soil, climate, organic inputs, and fertility; Saeur *et al.*, 2012).

The SOC sequestration rate after afforestation is lowest in cold climates; the meta-analysis of Laganière, Angers and Paré (2010) reported average SOC losses of 1.5 percent following afforestation in the boreal zone results, compared with gains ranging from 7 percent to 17 percent in the other climate zones (the highest gains were in the temperate maritime zone (Figure 3). In a global meta-analysis, planted forests in the warm temperate zone had the highest rate of SOC accumulation (0.96 tC/ha/yr), whereas the cold temperate zone had the lowest rate (0.21 tC/ha/yr), and the tropical climatic zone (0.56 tC/ha/yr) was close to the global average (0.50 tC/ha/yr) (Wang and Huang, 2020). The proportion of ecosystem C accumulated in the soil was highest in the warm temperate zone (27.0 percent) and lowest in the tropical zone (10.7 percent), with the cold temperate zone (17.9 percent) coming close to the global average (14.1 percent); the median proportion of ecosystem C accumulated in the soil was highest in the warm temperate zone. In tropical climates, mineral SOC accumulates faster and without the initial lag period often reported under temperate conditions (Don, Schumacher and Freibauer, 2011; Bárcena *et al.*, 2014a), but the greater productivity is countered by higher SOC losses. In a

meta-analysis of afforestation in China, SOC accumulation rate was most strongly related to MAP, followed by MAT, stand age, and soil clay content (Wang and Huang, 2020).



**Figure 3.** Change in SOC stocks after afforestation a) in different climatic zones and b) with different types of tree species

Error bars are the standard errors of the mean. Different letters indicate significant difference at  $P < 0.10$ . The number of observations is indicated in parentheses. The mean age of plantation is 22.9 years and the mean depth of sampling is 34.7 cm. BO, boreal; TC, temperate continental; TM, temperate maritime; STR, subtropical; TR, tropical. SOC, soil organic carbon; Eucal, *Eucalyptus* spp.; Conif, coniferous (excluding pine); Broad, broadleaf (excluding *Eucalyptus* spp.). Source: Laganière, Angers and Paré (2010)

Soil clay content is positively related to SOC change rate (Shi *et al.*, 2013; Wang and Huang 2020). Soils with high clay content (>33 percent) accumulated approximately 25 percent more C upon afforestation than coarse-textured soils (Laganière, Angers and Paré, 2010). The rate of increase in SOC following afforestation is greatest in organic layer and declines with soil depth (Shi *et al.*, 2013). Afforestation of shallow and deep peat soils may reduce SOC stocks as a consequence of drainage (Simola, Pitkänen and Turunen, 2012).

Cropland conversion to deciduous forests show faster increase in soil C stocks than to evergreen forest (Deng *et al.*, 2014; Guo and Gifford, 2002; Li *et al.*, 2012). In a global meta-analysis C, during the 2-3 decades following afforestation of agricultural soils, the average increase in SOC on sites afforested with broadleaf tree species was 25 percent, compared with 2 percent with conifers (and 12 percent with *Eucalyptus* or *Pinus* spp.; Laganière, Angers and Paré, 2010). In a global meta-analysis, soil C stock increased after afforestation with hardwoods such as *Eucalyptus*, but did not change after afforestation with softwoods such as pine (Li *et al.* 2012). On post-mining sites in the northern temperate zone, sites with deciduous forests accumulated SOC faster and deeper in the profile than sites with coniferous forests (Frouz *et al.*, 2014, Vinduskova and Frouz, 2013). The faster SOC accumulation under deciduous tree species is attributed to the high N and Ca concentrations in their leaf litter, which encourage bioturbation by earthworms (Morris *et al.*, 2007; Frouz *et al.*, 2013). Evergreen broadleaf species have higher average SOC sequestration rates (0.73 t/ha/yr) than the average for deciduous trees (0.42 t/ha/yr) or all evergreen trees (0.43 t/ha/yr), in a global quantitative review of afforestation studies (Hou *et al.*, 2020). On post-mining sites, SOC levels increase faster under trees with N-fixing root associates than under other tree species (Frouz *et al.*, 2009; Schiavo *et al.*, 2009; Kuznetsova *et al.*, 2011).

**Table 17.** Changes in soil organic carbon stocks reported following afforestation

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Years)	Depth (cm)	More information	Reference
Global meta-analyses								
Global	Various	Various	NA	0.50	Various	0-30	Afforestation meta-analysis	Wang and Huang (2020)
				0.49 total 0.34 organic layer 0.15 mineral layer		Standardized to top 100 cm		Li <i>et al.</i> (2012)
				0.42 0.46 0.15 0.09 0.05		Organic 0-20 20-40 40-60 60-100		Shi <i>et al.</i> (2013)
Regional meta-analyses and reviews								
Warm temperate zone	Warm temperate	Various	NA	0.96		0-30	Afforestation meta-analysis	Wang and Huang (2020)
				0.56				
Tropics	Tropical		60	0.44	28	NA	Afforestation of grassland meta-analysis	Don, Schumacher and Freibauer (2011)
		70	1.04	32	Afforestation of cropland meta-analysis			

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Years)	Depth (cm)	More information	Reference
			NA	1.3 0.41	20 100		Afforestation of agricultural land - review	Silver, Ostertag and Lugo (2000)
Cold temperate	Cold temperate			0.21 (0-30 cm)		0-30	Afforestation meta-analysis	Wang <i>et al.</i> (2020)
<b>National and local studies</b>								
South Carolina, United States of America	Warm temperate moist	Ultisols	32.5	1.0	40	0-60	Afforestation of abandoned agricultural land	Richter <i>et al.</i> (1999)
Southern and eastern Australia	Warm temperate moist & dry			0.57	6-45	0-30		England <i>et al.</i> (2016)
Southern Quebec, Canada		Brunisols, Gleysols, Podzols, Regosols	80	0.18 (sandy) 0.86 (loamy)	22	0-100 (mineral soil)		Ouimet <i>et al.</i> (2007)
Michigan, United States of America	Cold temperate moist	Typic Hapludalfs	51.8	0.35 deciduous 0.26 coniferous		NA	Afforestation of cropland	Morris <i>et al.</i> (2007)
Iowa, United States of America		mesic Typic Hapludolls	NA	0.56	15-50	0-30		Sauer <i>et al.</i> (2012)
Poland		Various	NA	0.34	10-50	0-100		Smal <i>et al.</i> (2019)
China	Various	Various	NA	0.13 0.26	40	0-20 0-100		Cropland conversion Grain for Green

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Years)	Depth (cm)	More information	Reference		
Post-mining restoration										
Temperate zone of N. Hemisphere	Various	Various	NA	2.46	0-10	0-10 + organic	NA	Frouz <i>et al.</i> (2014)		
				0.87	0-40					
				0.81 coniferous 2.32 deciduous	30					
United States of America				0.48-1.51	27-50	0-150			Amichev, Burger and Rodrigue (2008)	
India	Tropical dry			66.7	2.6	8		0-30		Ahirwal, Maiti and Singh (2017)
Ohio, United States of America				22.3	1.93 deciduous	NA				Akala and Lal (2001)
				4.86	1.19	31				Lorenz and Lal (2007)
				23.0	1.35-3.62	4-38		0-10 + organic		Sourkova <i>et al.</i> (2005)
Czechia	Cold temperate moist			NA	0.16-1.32	22-31		0-20		Frouz <i>et al.</i> (2009)
			0.65 active restoration 0.93 passive restoration		40	0-10 + organic		Frouz <i>et al.</i> (2014)		
			0.62		29	0-10 + organic		Frouz and Kalčík (2006)		
				0.42-1.21	15-37	0-12 + organic		Karu <i>et al.</i> (2009)		
Estonia				1.4-1.66	29-34	0-25		Reintam, Kar and Rooma (2002)		

## 4. Other benefits of the practice

Afforestation of degraded soils has positive effects on a range of ecosystem services such as enhanced water quality, regulation of water flow, reduced erosion and avalanche risk, habitat for forest-related biodiversity, recreation, and production of timber and non-timber wood products.

### 4.1. Improvement of soil properties

By increasing rooting and organic matter content, afforestation of degraded soils improves physical properties such as bulk density, structure, porosity, permeability, and water-holding capacity (Feng *et al.*, 2011; Sauer *et al.*, 2012). Depending on the tree species planted, afforestation can raise or lower soil pH (Hong *et al.*, 2018).

### 4.2 Minimization of threats to soil functions

Table 18. Soil threats

Soil threats	
<b>Soil erosion</b>	Forestation reduced water erosion of soils within 20 years (Bonnesoeur <i>et al.</i> , 2019). Windbreaks reduce wind speeds and soil loss from agricultural land (Brandle, Hodges and Wight, 2000).
<b>Nutrient imbalance and cycles</b>	Afforestation increases content and cycling of nutrients (Prescott <i>et al.</i> 2019). N-fixing species increase N and other nutrients (Perakis and Pett-Ridge, 2019).
<b>Soil salinization and alkalinization</b>	Afforestation of saline soils with salt-tolerant trees can lower the water table, promote the downward movement of salts in the soil profile and rehabilitate these soils (Bell, 1999; Byers <i>et al.</i> , 2006; Wicke <i>et al.</i> , 2011). Mangroves desalinate water (Reef and Lovelock, 2015; also see Hotspot: Mangroves).
<b>Soil contamination/pollution</b>	Some tree species can be used for phytoremediation of contaminated sites (Mleczek <i>et al.</i> , 2017).
<b>Soil acidification</b>	Afforestation neutralizes soil pH: it lowers pH in relatively alkaline soil but raises pH in relatively acid soil (Hong <i>et al.</i> , 2018)
<b>Soil biodiversity loss</b>	Restores soil microbial and faunal communities (Prescott <i>et al.</i> , 2019).
<b>Soil sealing</b>	Some trees species adapted to these soil conditions (Prescott, Katzensteiner and Weston, 2020).

Soil threats	
<b>Soil compaction</b>	Bulk density declines with increasing SOC (Sauer <i>et al.</i> , 2012)  Planting tree species that tolerate prolonged anoxic soil conditions can improve soil structure and aeration (Prescott, Katzensteiner and Weston, 2020).
<b>Soil water management</b>	Forestation reduced risk of moderate floods within 20 years (Bonnesoeur <i>et al.</i> , 2019). It can also decrease stream salinity (Ruprecht <i>et al.</i> , 2019).

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Afforestation of areas used for crop production reduces the area available for agricultural production and the food production potential (Sauer *et al.*, 2012). Effects on food production can be mitigated through forest landscape restoration mosaic approaches<sup>1</sup> in which sloping and marginal lands are afforested, leaving more productive land for agriculture. Agroforestry practices that promote food such as nuts and fruit from trees, incorporate trees into agricultural lands (e.g. intercropping, enhanced fallows, fertilizer trees, shade cropping) or integrate crops into the forest (e.g. forest gardens, taungya) provide benefits of trees while maintaining food production (e.g. Huang *et al.*, 2020; Giudice Badari *et al.*, 2020). Afforestation can include trees beneficial for fuel or fodder production. Afforestation with species of high value for timber production can be accomplished through monoculture plantations but these are less effective at storing C (Lewis *et al.*, 2019). Timber production can be balanced with other benefits of afforestation by mixing timber species with native species providing other ecosystem services (Amazonas *et al.*, 2013) or by including both in a multifunctional landscape approach (Stanturf *et al.*, 2015).

### 4.4 Mitigation of and adaptation to climate change

Afforestation and reforestation (i.e. reforesting unforested land) can contribute to climate change mitigation by increasing stocking density in forests, carbon sequestration in soils, and wood use in construction, or as a substitute of fossil carbon (e.g. coke) in the steel industry or energy facilities, thus avoiding fossil C-CO<sub>2</sub> emissions (Fallot *et al.*, 2009). The IPCC's Fifth Assessment Report (IPCC, 2014) ranked afforestation as moderate for technical mitigation potential and high for both immediacy and ease. The global climate mitigation potential of reforestation (i.e. transition from non-forest to forest at a 30 percent tree-cover threshold) has been estimated at 3-10 Pg CO<sub>2</sub>eq/y (Griscom *et al.*, 2017) and 3-18 billion tonnes of CO<sub>2</sub> per year (The Royal Society, 2018). If all areas that would naturally support forest and woodland were reforested, the extra 0.9 billion ha canopy cover would store 205 gigatons of C (Bastin *et al.*, 2019). A large-scale afforestation program that was economically, politically, and technically feasible would cover about 345 million ha and would sequester about 104 Gt of carbon (Nilsson and Schopfhauser, 1995).

<sup>1</sup> Also factsheet n° 8 “Forest Landscape Restoration”, this volume.

Afforestation and reforestation also generate changes in albedo (i.e. surface reflectivity of light). Afforestation leads to darker surfaces (lower albedo), especially at high latitudes (e.g. boreal areas); the associated alteration of radiative forcing can weaken the benefits from increased C storage (Kirschbaum *et al.*, 2011). Afforestation can also influence local and regional climate by evaporative cooling (Bonan *et al.*, 2008; Locatelli *et al.*, 2015; Alkama and Cescatti, 2016) and increasing rainfall by recycling it via transpiration (Ellison *et al.*, 2017). Overall, in afforested areas, the warming effect of decreased albedo dominates the cooling effect of increased surface roughness and evaporation at high latitudes, leading to warmer local climate, while the reverse occurs in tropical areas where surface temperatures of afforested lands are generally lower than surface temperatures of grasslands or croplands (Jackson *et al.*, 2008; Bonan, 2016; Peng *et al.*, 2014).

## 4.5 Socio-economic benefits

In addition to eventual products from forests (timber, fuel, fodder, food), requirements for seedlings can generate business opportunities. For example, the widespread adoption by farmers in Java of small rural woodlots has led to the development of local seedling vendors and the creation of processing industries to use the timbers (FAO, 2005).

Payment for ecosystem services such as C sequestration can also generate socio-economic benefits from afforestation. For example, the International Small Group & Tree Planting Program (TIST; <https://program.tist.org/>) encourages small groups of subsistence farmers to improve their local environment and farms by planting and maintaining trees on degraded land. Over 93 000 farmers in 4 countries have successfully planted more than 19 000 000 trees and captured over 5 500 000 metric tons of carbon dioxide to date. As the trees grow, carbon captured is quantified and verified and certified greenhouse gas credits are sold in the global carbon market. TIST farmers receive annual carbon pre-payments for each tree established and 70 percent of the net profit when credits are sold. Smallholder farmers also derive significant non-carbon related benefits verified to exceed USD 8 per tree. Trees provide fruit, fodder, fuel, windbreaks, shade and stabilize riverbanks plus participants have access to health information and training and safe cooking stoves.

Stakeholders interested in the potential benefits of tree planting beyond just the commercial timber companies include farmers and rural communities, water managers and hydroelectric power generating agencies and protected area managers (FAO, 2005).



## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

Table 19. Soil threats

Soil threats	
<b>Soil erosion</b>	Erosion is usually low on afforested lands, except under some tropical plantations that do not have an understorey and/or a thick litter floor (e.g. some rubber and teak plantations; Liu <i>et al.</i> , 2016; Lacombe <i>et al.</i> , 2018; Neyret <i>et al.</i> , 2020).
<b>Nutrient imbalance and cycles</b>	Changes in SOM inputs and pH can lead to differences in soil N mineralization and nitrification (Li <i>et al.</i> , 2014).
<b>Soil salinization and alkalinization</b>	In sub-humid areas (i.e. aridity index between 0.5 and 0.65) replacing open arable lands/grasslands by planted forests can lower the water table and increase salt concentration in soils and groundwater (Toth <i>et al.</i> , 2014)
<b>Soil acidification</b>	Trees can acidify soil (Berhongaray <i>et al.</i> , 2013; Ritter, Vesterdal and Gundersen, 2003), especially N <sub>2</sub> -fixing species (Russell, Hall and Raich, 2017; Dubiez <i>et al.</i> , 2019).
<b>Soil water management</b>	Higher water use of forest may reduce total water supply (Bonnesoeur <i>et al.</i> , 2019). In arid regions, afforestation can exacerbate water shortages (Cao <i>et al.</i> , 2011).

### 5.2 Increases in greenhouse gas emissions

Afforestation of wet sites can increase CO<sub>2</sub> flux from soil if drainage is required. Replacement of grassland and pasture can cause loss of C from soil.

### 5.3 Conflict with other practice(s)

Conversion of biomes such as grasslands and savanna can reduce soil C stocks (Guo and Gifford, 2002; Shi *et al.*, 2013) and compromise other ecosystem services, such as soil nutrient cycles (Berthrong, Jobbágy and Jackson, 2009), hydrological regulation, erosion mitigation and water yield (Bonnesoeur *et al.*, 2019) and reduce biodiversity (Bond, 2016; Veldman *et al.*, 2015).

## 5.4 Decreases in production (e.g. food/fuel/feed/timber)

Afforestation with native species and/or mixed species may be less cost-effective than monocultures of exotic species. This can be mitigated by growing them in mixtures (e.g. Eucalypts and native species; Amazonas *et al.*, 2013) or by underplanting native species in plantations of exotic species (Parrotta, Turnbull and Jones, 1997).

## 5.5 Other conflicts

Unless carefully planned to suit local situations, afforestation can have unintended negative ecological and social consequences – including reduced water supply, destruction of native grasslands, spread of invasive tree species, increased social inequity, displacement of farmland, and increased deforestation elsewhere (Holl and Brancalion, 2020).

Afforestation can create conflict where ownership is unclear or disputed, or where the land is already being used by local communities for other purposes. Attempts to reforest land subject to conflicting land-ownership claims are unlikely to be successful because of deliberate vandalism by disadvantaged parties. Collaborative and participative approaches involving local communities are necessary under such situations (Stanturf *et al.*, 2015; FAO, 2005).

Many sites available for afforestation are poor or degraded, with the better land usually being used for agriculture. Rehabilitation of the site and soil may be necessary prior to planting trees (Prescott, Katzensteiner and Weston, 2020).

Afforestation can be expensive and subject to considerable risk (e.g. fires, droughts, disease and changing markets), with long periods before any financial return is possible. Direct subsidies, joint ventures between landowners and an industrial partner, low-interest loans, micro-credit, tax concessions or payment for ecosystem services schemes may be necessary (FAO, 2005). Contracting, open bidding, and other market-based mechanisms for carrying out various operations can support development of local business (Yin, Sedjo and Liu, 2010).

Focus on acreage expansion and neglecting forest management can result in low stocking levels of both natural and plantation forests (Yin, Sedjo and Liu, 2010). Some subsidy programs lead to over-reliance on single species, poor stand quality and reversion to cropland if subsidies end (Yin, Sedjo and Liu, 2010). Emphasis should be on ‘tree-growing’ rather than ‘tree-planting’ (Lyons, 2019).

Limited silvicultural knowledge is a common impediment to successful reforestation and most foresters currently rely on a handful of well-known species for plantation development. Often, little is now known of the identity, ecology, silviculture or site requirements of many indigenous species (FAO, 2005).

## 6. Recommendations before implementation of the practice

- ◆ Engage local communities to develop strong local support. Resolve land ownership and tenure conflicts.
- ◆ Ensure institutional support and long-term funding to incentivize and support afforestation programs.
- ◆ Clarify distribution of costs and benefits, and responsibilities and rewards, among all parties involved in program.
- ◆ Develop appropriate methods for tree species to be used in afforestation programs.
- ◆ Assess site conditions and ameliorate to improve likelihood of trees survival and growth.
- ◆ Identify tree species that are able to tolerate conditions at degraded sites available for afforestation and will not become invasive or negatively influence the environment.
- ◆ Have a monitoring plan in place and adaptive management protocols to prevent plantation failure.
- ◆ Train local silviculturalists in site preparation, weed control, species and provenance choice, nutrition, stand tending and fire prevention.
- ◆ Consider agroforestry or forest landscape restoration approaches to accommodate food production and restoration demands.

## 7. Potential barriers to adoption

Table 20. Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	If site has inadequate stability, fertility, soil depth, water supply or water-holding capacity, rehabilitation measures are needed prior to planting.
Cultural	Yes	Tree species should be familiar to locals and useful/desired. Trees may not be desired where landscape has been deforested for generations.
Social	Yes	Planting must be supported by local community.
Economic	Yes	Funding needed to develop capacity to produce seedlings of right species and genotype locally, for site preparation and establishment, and long-term monitoring and maintenance. Economic return only at the end of the rotation, which can last several years or decades.

Barrier	YES/NO	
Institutional	Yes	Successful reforestation programs have occurred when national governments have made a serious and prolonged effort over a number of years by providing such supportive incentives and policies.
Legal (Right to soil)	Yes	Local community should have right to use products or be subsidized.
Knowledge	Yes	Knowledge of appropriate species for environment and for sustainable use by locals, and knowledge of the best management practices.

## Photo of the practice



**Photo 5.** Successful afforestation involving a mix of species in a lowland rain forest near Rio de Janeiro, Brazil  
 The site was previously farmland. This is part of a major recovery effort for the Atlantic Forest of Brazil

**Table 21.** Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Agroforestry, silvopastoral systems and water funds initiatives contribute to improve soil capacity to remove and store carbon in Colombia</i>	Latin America and the Caribbean	9, 20 and 40	4	36
<i>Soil fertility improvement of nutrient-poor and sandy soils in the Congolese coastal plains</i>	Africa	7	6	1
<i>Afforestation of a former farmland in Japan</i>	Asia	5	6	2
<i>Reforestation of highlands in Javor Mountain, Republic of Srpska, Bosnia and Herzegovina</i>	Europe	15	6	5
<i>Natural afforestation of abandoned mountain grasslands along the Italian peninsula</i>	Europe	23 to 72	6	6
<i>Afforestation of vineyards in Italy</i>	Europe	Up to 30	6	7
<i>Afforestation by planting in bench terraces: Kalimanska watershed, Grdelica gorge, Southeastern Serbia</i>	Europe	60	6	8

## References

- Ahirwal, J., Maiti, S.K. & Singh, A.K. 2017. Changes in ecosystem carbon pool and soil CO<sub>2</sub> flux following post-mine reclamation in dry tropical environment, India. *Science of The Total Environment*, 583: 153-162. <https://doi.org/10.1016/j.scitotenv.2017.01.043>
- Akala, V.A. & Lal, R. 2001 Soil organic carbon pools and sequestration rates in reclaimed minesoils in Ohio. *Journal of Environmental Quality*, 30(6): 2098–2104. <https://doi.org/10.2134/jeq2001.2098>
- Alkama, R. & Cescatti, A. 2016. Climate change: Biophysical climate impacts of recent changes in global forest cover. *Science*, 351: 600-604. <https://doi.org/10.1126/science.aac8083>
- Amazonas, N.T., Forrester, D.I., Silva, C.C., Almeida, D.R.A., Rodrigues, R.R. & Brancalion, P.H.S. 2018. High diversity mixed plantations of Eucalyptus and native trees: An interface between production and restoration for the tropics. *Forest Ecology and Management*, 417: 247-256. <https://doi.org/10.1016/j.foreco.2018.03.015>
- Amichev, B.Y., Burger, J.A. & Rodrigue, J.A. 2008. Carbon sequestration by forests and soils on mined land in the Midwestern and Appalachian coalfields of the US. *Forest Ecology and Management*, 256(11): 1949–1959. <https://doi.org/10.1016/j.foreco.2008.07.020>
- Assefa, D., Rewald, B., Sandén, H., Rosinger, C., Abiyu, A., Yitferu, B. & Godbold, D.L. 2017. Deforestation and land use strongly effect soil organic carbon and nitrogen stock in Northwest Ethiopia. *Catena*, 153: 89–99. <https://doi.org/10.1016/j.catena.2017.02.003>
- Bárcena, T.G., Kiar, L.P., Vesterdal, L., Stefánsdóttir, H.M., Gundersen, P. & Sigurdsson, B.D. 2014. Soil carbon stock change following afforestation in Northern Europe: a meta-analysis. *Global Change Biology*, 20: 2393-2405. <https://doi.org/10.1111/gcb.12576>
- Bastin, J-F, Finegold, Y., Garcia, C., Mollicone, D., Rezende, M., Routh, D., Zohner, C.M. & Crowther, T.W. 2019. The global tree restoration potential. *Science*, 365(6448): 76-79. <https://doi.org/10.1126/science.aax0848>
- Bell, D.T. 1999. Australian trees for the rehabilitation of waterlogged and salinity-damaged landscapes. *Australian Journal of Botany*, 47(5): 697-716. <https://doi.org/10.1071/BT96110>
- Berhongeray, G. & Alvarez, R. 2019. Soil carbon sequestration of Mollisols and Oxisols under grassland and tree plantations in South America - A review. *Geoderma Regional*, 18: e00226. <https://doi.org/10.1016/j.geodrs.2019.e00226>
- Berhongeray, G., Alvarez, R., De Paepe, J., Caride, C. & Cantet, R. 2013. Land use effects on soil carbon in the Argentine Pampas. *Geoderma*, 192:97-110. <https://doi.org/10.1016/j.geoderma.2012.07.016>
- Berthrong, S.T., Jobbágy, E.G. & Jackson, R.B. 2009. A global meta-analysis of soil exchangeable cations, pH, carbon, and nitrogen with afforestation. *Ecological Applications*, 19: 2228–2241. <https://doi.org/10.1890/08-1730.1>
- Bonan, G.B. 2008. Forests and climate change: Forcings, feedbacks, and the climate benefits of forests. *Science*, 320(5882): 1444-1449. <https://doi.org/10.1126/science.1155121>

- Bonan, G.B.** 2016. Forests, Climate, and Public Policy: A 500-Year Interdisciplinary Odyssey. *Annual Review of Ecology, Evolution, and Systematics*, 47: 97–121. <https://doi.org/10.1146/annurev-ecolsys-121415-032359>
- Bond, W.J.** 2016. Ancient grasslands at risk. *Science*, 351(6269): 120–122. <https://doi.org/10.1126/science.aad5132>
- Bonnesoeur, V., Locatelli, B., Guariguata, M.R., Ochoa-Tocachi, B.F., Vanacker, V., Mao, Z., Stokes, A. & Mathez-Stiefel, S.-L.** 2019. Impacts of forests and forestation on hydrological services in the Andes: A systematic review. *Forest Ecology and Management*, 433: 569–584. <https://doi.org/10.1016/j.foreco.2018.11.033> *et al.* 2019
- Brandle, J.R., Hodges, L. & Wight, B.** 2000. Windbreak practices. In Garrett, H.E., Rietveld, W.J., Fisher, R.F. (Eds.) *North American Agroforestry: An Integrated Science and Practice*. American Society of Agronomy, pp. 79–118.
- Byers, J.E., Cuddington, K., Jones, C.G., Talley, T.S., Hastings, A., Lambrinos, J.G., Crooks, J.A. & Wilson, W.G.** 2006. Using ecosystem engineers to restore ecological systems. *Trends in Ecology and Evolution*, 21: 493–500. <https://doi.org/10.1016/j.tree.2006.06.002>
- Cao, S.X., Chen, L., Shankman, D., Wang, C.M., Wang, X.B. & Zhang, H.** 2011. Excessive reliance on afforestation in China’s arid and semi-arid regions, Lessons in ecological restoration. *Earth-Science Reviews*, 104: 240–245. <https://doi.org/10.1016/j.earscirev.2010.11.002>
- Deng, L., Liu, G. & Shangguan, Z.** 2014. Land-use conversion and changing soil carbon stocks in China’s ‘Grain-for-Green’ Program: a synthesis. *Global Change Biology*, 20: 3544–3556. <https://doi.org/10.1111/gcb.12508>
- Don, A., Schumacher, J. & Freibauer, A.** 2011. Impact of tropical land-use change on soil organic carbon stocks – a meta-analysis. *Global Change Biology*, 17: 1658–1670. <https://doi.org/10.1111/j.1365-2486.2010.02336.x>
- Dubiez, E., Freycon, V., Marien, J.N., Peltier, R. & Harmand, J.M.** 2019. Long term impact of *Acacia auriculiformis* woodlots growing in rotation with cassava and maize on the carbon and nutrient contents of savannah sandy soils in the humid tropics (Democratic Republic of Congo). *Agroforestry Systems*, 93: 1167–1178. <https://doi.org/10.1007/s10457-018-0222-x>
- Ellison, D., Morris, C.E., Locatelli, B., Sheil, D., Cohen, J., Murdiyarso, D., Gutierrez, V., Noordwijk, M. van, Creed, I.F., Pokorny, J., Gaveau, D., Spracklen, D.V., Tobella, A.B., Ilstedt, U., Teuling, A.J., Gebrehiwot, S.G., Sands, D.C., Muys, B., Verbist, B., Springgay, E., Sugandi, Y. & Sullivan, C.A.** 2017. Trees, forests and water: Cool insights for a hot world. *Global Environmental Change*, 43: 51–61. <https://doi.org/10.1016/j.gloenvcha.2017.01.002>
- England, J.R., Paul, K.I., Cunningham, S.C., Madhavan, D.B., Baker, T.G., Read, Z., Wilson, B.R., Cavagnaro, T.R., Lewis, T. Perring, M.P., Herrman, T. & Polglase, P.J.** 2016. Previous land use and climate influence differences in soil organic carbon following reforestation of agricultural land with mixed-species plantings. *Agriculture, Ecosystems and Environment*, 227: 61–72. <https://doi.org/10.1016/j.agee.2016.04.026>

**Fallot, A., Saint-André, L., Le-Maire, G., Laclau, J.P., Nouvellon, Y., Marsden, C., Bouillet, J.P., Silva, T., Pikett, M.G. & Hamel, O.** 2009. Biomass sustainability, availability and productivity. *Revue de Metallurgie*, 106(10): 410-418. <https://doi.org/10.1051/metal/2009072>

**FAO.** 2000. *FRA 2000: On definitions of Forest and Forest Change*. (also available at: [www.fao.org/3/ad665e/ad665e04.htm](http://www.fao.org/3/ad665e/ad665e04.htm))

**FAO.** 2005. *Helping Forests Take Cover*. Regional Centre For Asia and the Pacific RAP Publication 2005/13 Bangkok. (also available at: [www.fao.org/3/ac945e/ac945e00.htm](http://www.fao.org/3/ac945e/ac945e00.htm))

**FAO.** 2017. Land Use, Irrigation and Agricultural Practices – Definitions. (also available at: [www.fao.org/fileadmin/templates/ess/ess\\_test\\_folder/Definitions/Land\\_Use\\_Definitions\\_FAOSTAT](http://www.fao.org/fileadmin/templates/ess/ess_test_folder/Definitions/Land_Use_Definitions_FAOSTAT))

**Frouz, J. & Kalčík, J.** 2006. Accumulation of soil organic carbon in relation to other soil characteristic during spontaneous succession in non-



- Hong, S., Piao, S., Chen, A., Liu, Y., Liu, L., Peng, S., Sardans, J., Sun, Y., Peñuelas, J. & Zeng, H. 2018. Afforestation neutralizes soil pH. *Nature Communications*, 9: 520. <https://doi.org/10.1038/s41467-018-02970-1>
- Hou, G., Deland, C.O., Lu, X. & Gao, L. 2020 Grouping tree species to estimate afforestation-driven soil organic carbon sequestration. *Plant and Soil*, 455: 507-518. <https://doi.org/10.1007/s11104-020-04685-z>
- Huang, J., Pan, J., Zhou, L., Zheng, D., Yuan, S., Chen, J., Li, J., Gui, Q. & Lin, W. 2020. An improved double-row rubber (*Hevea brasiliensis*) plantation system increases land use efficiency by allowing intercropping with yam bean, common bean, soybean, peanut, and coffee: A 17-year case study on Hainan Island, China. *Journal of Cleaner Production*, 263, 121493. <https://doi.org/10.1016/j.jclepro.2020.121493>
- IPCC. 2014. Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth In Core Writing Team, Pachauri, R.K. & Meyer, L.A. (Eds.) *Assessment Report of the Intergovernmental Panel on Climate Change*. IPCC, Geneva, Switzerland, 151 pp.
- Jackson, R.B., Randerson, J.T., Canadell, J.G., Anderson, R.G., Avissar, R., Baldocchi, D.D., Bonan, G.B., Caldeira, K., Diffenbaugh, N.S., Field, C.B., Hungate, B.A., Jobbágy, E.G., Kueppers, L.M., Nosoetto, M.D. & Pataki, D.E. 2008. Protecting climate with forests. *Environmental Research Letters*, 3(4). <http://dx.doi.org/10.1088/1748-9326/3/4/044006>
- Karu, H., Szava-Kovats, R., Pensa, M. & Kull, O. 2009. Carbon sequestration in a chronosequence of Scots pine stands in a reclaimed opencast oil shale mine. *Canadian Journal of Forest Research*, 39(8): 1507–1517. <https://doi.org/10.1139/X09-069>
- Kirschbaum, M.U.F., Whitehead, D., Dean, S.M., Beets, P.N., Shepherd, J.D. & Ausseil, A.G.E. 2011. Implications of albedo changes following afforestation on the benefits of forests as carbon sinks. *Biogeosciences*, 8: 3687-3696. <https://doi.org/10.5194/bg-8-3687-2011>
- Lacombe, G., Valentin, C., Sounyafong, P., de Rouw, A., Soulléuth, B., Silvera, N., Pierret, A., Sengtaheuanghong, O. & Ribolzi, O. 2018. Linking crop structure, throughfall, soil surface conditions, runoff and soil detachment: 10 land uses analyzed in Northern Laos. *Science of the Total Environment*, (616-617): 1330-1338. <https://doi.org/10.1016/j.scitotenv.2017.10.185>
- Laganière, J., Angers, D.A. & Paré, D. 2010. Carbon accumulation in agricultural soils after afforestation: A meta-analysis. *Global Change Biology*, 16(1): 439-453. <https://doi.org/10.1111/j.1365-2486.2009.01930.x>
- Ledo, A., Smith, P., Zerihun, A., Whitaker, J., Vicente-Vicente, J.L., Qin, Z., McNamara, N.P., Zinn, Y.L., Llorente, M., Liebig, M., Kuhnert, M., Dondini, M., Don, A., Diaz-Pines, E., Datta, A., Bakka, H., Aguilera, E. & Hillier, J. 2020. Changes in soil organic carbon under perennial crops. *Global Change Biology*, 26: 4158-4168. <https://doi.org/10.1111/gcb.15120>
- Lewis, S.L., Wheeler, C.E., Mitchard, E.T.A. & Koch, A. 2019. Regenerate natural forests to store carbon. *Nature*, 568(7750): 25-28. <https://doi.org/10.1038/d41586-019-01026-8>

- Li, D., Niu, S. & Luo, Y.** 2012. Global patterns of the dynamics of soil carbon and nitrogen stocks following afforestation: a meta-analysis. *New Phytologist*, 195: 172–181. <https://doi.org/10.1111/j.1469-8137.2012.04150.x>
- Li, M., Zhou, X., Zhang, Q. & Cheng, X.** 2014. Consequences of afforestation for soil nitrogen dynamics in central China. *Agriculture, Ecosystems & Environment*, 183: 40–46. <https://doi.org/10.1016/j.agee.2013.10.018>
- Liu, H., Blagodatsky, S., Giese, M., Liu, F., Xu, J. & Cadisch, G.** 2016. Impact of herbicide application on soil erosion and induced carbon loss in a rubber plantation of Southwest China. *Catena*, 145: 180–192. <https://doi.org/10.1016/j.catena.2016.06.007>
- Locatelli, B., Catterall, C.P., Imbach, P., Kumar, C., Lasco, R., Marín-Spiotta, E., Mercer, B., Powers, J.S., Schwartz, N. & Uriarte, M.** 2015. Tropical reforestation and climate change: Beyond carbon. *Restoration Ecology*, 23: 337–343. <https://doi.org/10.1111/rec.12209>
- Lorenz, K. & Lal, R.** 2007. Stabilization of organic carbon in chemically separated pools in reclaimed coal mine soils in Ohio. *Geoderma*, 141(3–4): 294–301. <https://doi.org/10.1016/j.geoderma.2007.06.008>
- Lyons, D.** 2019. Forest tree planting, start tree growing [online]. [Cited 16 October 2020] <https://forestsnews.cifor.org/61174/forget-tree-planting-start-tree-growing?fnl=en>
- Mayer, M., Prescott, C.E., Abaker, W.E.A., Augusto, L., Cécillon, L., Ferreira, G.W.D., James, J., Jandl, R., Katzensteiner, K., Laclau, J.-P., Laganière, J., Nouvellon, Y., Paré, D., Stanturf, J.A., Vanguelova, E.I. & Vesterdal, L.** 2020. Tamm Review: Influence of forest management activities on soil organic carbon stocks: A knowledge synthesis. *Forest Ecology and Management*, 466: 118127. <https://doi.org/10.1016/j.foreco.2020.118127>
- Mleczek, M., Goliński, P., Krzesłowska, M., Gąsecka, M., Magdziak, Z., Rutkowski, P., Budzyńska, S., Waliszewska, B., Kozubik, T., Karolewski, Z. & Niedzielski, P.** 2017. Phytoextraction of potentially toxic elements by six tree species growing on hazardous mining sludge. *Environmental Science and Pollution Research*, 24: 22183–22195. <https://doi.org/10.1007/s11356-017-9842-3>
- Morris, S.J., Bohm, S., Haile-Mariam, S. & Paul, E.A.** 2007. Evaluation of carbon accrual in afforested agricultural soils. *Global Change Biology*, 13: 1145–1156. <https://doi.org/10.1111/j.1365-2486.2007.01359.x>
- Nave, L.E., Swanston, C.W., Mishra, U. & Nadelhoffer, K.J.** 2013. Afforestation effects on soil carbon storage in the United States: A synthesis. *Soil Science Society of America Journal*, 77: 1035–1047. <https://doi.org/10.2136/sssaj2012.0236>
- Neyret, M., Robain, H., de Rouw, A., Janeau, J.L., Durand, T., Kaewthip, J., Trisophon, K. & Valentin, C.** 2020. Higher runoff and soil detachment in rubber tree plantations compared to annual cultivation is mitigated by ground cover in steep mountainous Thailand. *Catena*, 189: 104472. <https://doi.org/10.1016/j.catena.2020.104472>
- Nilsson, S. & Schopfhauser, W.** 1995. The carbon-sequestration potential of a global afforestation program. *Climatic Change*, 30: 267–293. <https://doi.org/10.1007/BF01091928>

- Parrotta, J.A., Turnbull, J.W. & Jones, N.** 1997. Catalyzing native forest regeneration on degraded tropical lands. *Forest Ecology and Management*, 99: 1–7. [https://doi.org/10.1016/S0378-1127\(97\)00190-4](https://doi.org/10.1016/S0378-1127(97)00190-4)
- Paul, K.I., Polglase, P.J., Nyakuengama, J.G. & Khanna, P.K.** 2002. Change in soil carbon following afforestation. *Forest Ecology and Management*, 168(1-3): 241-257. [https://doi.org/10.1016/S0378-1127\(01\)00740-X](https://doi.org/10.1016/S0378-1127(01)00740-X)
- Peng, S.S., Piao, S., Zeng, Z., Ciais, P., Zhou, L., Li, L.Z.X., Myneni, R.B., Yin, Y. & Zeng, H.** 2014. Afforestation in China cools local land surface temperature. *Proceedings of the National Academy of Sciences of the United States of America*, 111: 2915-2919. <https://doi.org/10.1073/pnas.1315126111>
- Perakis, S.S. & Pett-Ridge, J.C.** 2019. Nitrogen-fixing red alder trees tap rock-derived nutrients. *Proceedings of the National Academy of Sciences*, 116(11): 5009-5014. <https://doi.org/10.1073/pnas.1814782116>
- Poepplau, C., Don, A., Vesterdal, L., Leifeld, J., Van Wesemael, B., Schumacher, J. & Gensior, A.** 2011. Temporal dynamics of soil organic carbon after land-use change in the temperate zone – carbon response functions as a model approach. *Global Change Biology*, 17: 2415-2427. <https://doi.org/10.1111/j.1365-2486.2011.02408.x>
- Prescott, C.E., Frouz, J., Grayston, S.J., Quideau, S.A. & Straker, J.** 2019. Rehabilitating forest soils after disturbance. *Developments in Soil Science*, 36: 309-343. <https://doi.org/10.1016/B978-0-444-63998-1.00013-6>
- Prescott, C.E., Katzensteiner, K. & Weston C.** 2020. Soils and Restoration of Forested Landscapes. *In* Stanturf, J.A & Callahan, M. (Eds.) *Soils and Restoration of Forested Landscape*. Elsevier (in press).
- Reef, R. & Lovelock, C.E.** 2015. Regulation of water balance in mangroves, *Annals of Botany*, 115(3): 385–395. <https://doi.org/10.1093/aob/mcu174>
- Reintam, L., Kaar, E. & Rooma, I.** 2002. Development of soil organic matter under pine on quarry detritus of open-cast oil-shale mining. *Forest Ecology and Management*, 171(1–2): 191–198. [https://doi.org/10.1016/S0378-1127\(02\)00472-3](https://doi.org/10.1016/S0378-1127(02)00472-3)
- Richter, D., Markewitz, D., Trumbore, S. & Wells, C.G.** 1999. Rapid accumulation and turnover of soil carbon in a re-establishing forest. *Nature*, 400: 56–58. <https://doi.org/10.1038/21867>
- Ritter, E., Vesterdal, L. & Gundersen, P.** 2003. Changes in soil properties after afforestation of former intensively managed soils with oak and Norway spruce. *Plant and Soil*, 249: 319–330. <https://doi.org/10.1023/A:1022808410732>
- Ruprecht, J., Sparks, T., Liu, N., Dell, B. & Harper, R.** 2019. Using reforestation to reverse salinisation in a large watershed. *Journal of Hydrology*, 577: 123976. <https://doi.org/10.1016/j.jhydrol.2019.123976>
- Russell, A.E., Hall, S.J. & Raich, J.W.** 2017. Tropical tree species traits drive soil cation dynamics via effects on pH: a proposed conceptual framework. *Ecological Monographs*, 87(4): 685-701. <https://doi.org/10.1002/ecm.1274>

- Sauer, T.J., James, D.E., Cambardella, C.A. & Hernandez-Ramirez, G.** 2012. Soil properties following reforestation or afforestation of marginal cropland. *Plant and Soil*, 360: 375–390.  
<https://doi.org/10.1007/s11104-012-1258-8>
- Shi, S., Zhang, W., Zhang, P., Yu, Y. & Ding, F.** 2013. A synthesis of change in deep soil organic carbon stores with afforestation of agricultural soils. *Forest Ecology and Management*, 296: 53–63.  
<https://doi.org/10.1016/j.foreco.2013.01.026>
- Silveira, F.A.O., Arruda, A.J., Bond, W., Durigan, G., Fidelis, A., Kirkman, K., Oliveira, R.S., Overbeck, G.E., Sansevero, J.B.B., Siebert, F., Siebert, S.J., Young, T.P. & Buisson, E.** 2020. Myth-busting tropical grassy biome restoration. *Restoration Ecology*, 28(5): 1067–1073.  
<https://doi.org/10.1111/rec.13202>
- Silver, W.L., Ostertag, R. & Lugo, A.E.** 2000. The potential for carbon sequestration through reforestation of abandoned tropical agricultural and pasture lands. *Restoration Ecology*, 8: 394–407.  
<https://doi.org/10.1046/j.1526-100x.2000.80054.x>
- Simola, H., Pitkänen, A. & Turunen, J.** 2012. Carbon loss in drained forestry peatlands in Finland, estimated by re-sampling peatlands surveyed in the 1980s. *European Journal of Soil Science*, 63: 798–807.  
<https://doi.org/10.1111/j.1365-2389.2012.01499.x>
- Smal, H., Ligeza, S., Pranagal, J., Urban, D. & Pietruczyk-Popławska, D.** 2019. Changes in the stocks of soil organic carbon, total nitrogen and phosphorus following afforestation of post-arable soils: A chronosequence study. *Forest Ecology and Management*, 451: 117536.  
<https://doi.org/10.1016/j.foreco.2019.117536>
- Stanturf, J.A., Kant, P., Barnekow Lillesø, J.-P., Mansourian, S., Kleine, M., Graudal, L. & Madsen, P.** 2015. *Forest Landscape Restoration as a Key Component of Climate Change Mitigation and Adaptation*. IUFRO World Series Volume 34. Vienna 72 p.
- The Royal Society.** 2018. *Greenhouse Gas Removal*. (also available at: <https://royalsociety.org/-/media/policy/projects/greenhouse-gas-removal/royal-society-greenhouse-gas-removal-executive-summary-2018.pdf>)
- Tóth T., Balog, K., Szabó, A., Pásztor, L., Jobbágy, E.G., Noretto, M.D. & Gribovszki, Z.** 2014. Influence of lowland forests on subsurface salt accumulation in shallow groundwater areas, *AoB PLANTS*, 6: plu054. <https://doi.org/10.1093/aobpla/plu054>
- Vindušková, O. & Frouz, J.** 2013. Soil carbon accumulation after open-cast coal and oil shale mining in Northern Hemisphere: a quantitative review. *Environmental Earth Science*, 69: 1685–1698.  
<https://doi.org/10.1007/s12665-012-2004-5>
- Wang, S. & Huang, Y.** 2020. Determinants of soil organic carbon sequestration and its contribution to ecosystem carbon sinks of planted forests. *Global Change Biology*, 26: 3163–3173.  
<https://doi.org/10.1111/gcb.15036>
- Wei, J., Cheng, Y., Li, W. & Liu, W.** 2012. Comparing the effect of naturally restored forest and grassland on carbon sequestration and vertical distribution in the Chinese Loess Plateau. *PLoS ONE*, 7(7): e40123.  
<https://doi.org/10.1371/journal.pone.0040123>

**Wicke, B., Smeets, E., Dornburg, V., Vashev, B., Gaiser, T., Turkenburg, W. & Faaij, A.** 2011. The global technical and economic potential of bioenergy from salt-affected soils. *Energy and Environmental Science*, 4: 2669-2681. <https://doi.org/10.1039/C1EE01029H>

**Yin, R., Sedjo, R. & Liu, P.** 2010. The potential and challenges of sequestering carbon and generating other services in China's forest ecosystems. *Environmental Science & Technology*, 44(15): 5687-5688. <https://doi.org/10.1021/es1015636>

# 7. Rehabilitation of forest soils affected by wildfires

Montserrat Díaz-Raviña<sup>1</sup>, María Teresa Fontúrbel-Lliteras<sup>2</sup>, Ángela Martín<sup>1</sup>,  
Cristina Fernández<sup>2</sup>

<sup>1</sup>Departamento de Bioquímica del Suelo, Instituto de Investigaciones Agrobiológicas de Galicia (IIAG-CSIC), Santiago de Compostela, Spain

<sup>2</sup>Centro de Investigación Forestal-Lourizán, Consellería do Medio Rural, Xunta de Galicia, Pontevedra, Spain

## 1. Description of the practice

Fires are one of the most frequent and important disturbances of forest ecosystems. When, after a severe fire, heavy rainfall occurs in a burnt area of pronounced relief, there is a potential risk of strong alterations in the hydrological behavior of the affected basins. This can lead to large increases in surface runoff and intense erosive episodes (Vega, Fernández and Fontúrbel, 2018). These phenomena favor soil degradation and can cause floods and landslides, threatening human life, infrastructure and various valuable resources within and outside the burnt area.

Traditionally, hydrological-forestry restoration after fires has been mainly focused on the recovery of the destroyed vegetation cover and on the reduction of soil and sediment losses after fire that, in most of the environments, occurred during the first year (Fernández and Vega, 2016b). Therefore, it is necessary to apply emergency soil stabilization measures to try to reduce: a) runoff and soil erosion risk and consequently, the maintenance/conservation of the quality of water and aquatic habitats and b) the degradation of soil, which is an essential element for the recovery of the affected ecosystem. These measures favor indirectly the maintenance/recovery of most physical, chemical, biochemical and microbiological soil properties, which are related to the soil organic carbon (SOC) stock and hence to the maintenance/conservation of the quality of the burnt soil. Given the risk of large-scale hydrological events, these actions allow, above all, to protect human life and a set of valuable resources that can be critically threatened in a very short time after the fire events. To this end, their objectives are to protect the burnt soil, to limit its disintegration and subsequent loss of C and nutrient stocks, uprooting and transport, as well as to reduce runoff, while stabilizing the watercourses, where appropriate. The most efficient treatment to achieve that objective is the application of a mulching of different plant materials (straw, wood strands, wood chips) over the burnt soil surface (see the reports by Robichaud,

Beyers and Neary, 2000, Vega *et al.*, 2013a and Fernández *et al.*, 2019a). This reproduces the natural conditions in a pine forest affected by a low or medium severity wildfire, when the fall of pine needles forms a mulch that protects the soil from erosion; in this case, the implementation of these emergency soil stabilization measures is not necessary.

## 2. Range of applicability

The use of soil stabilization techniques is applicable worldwide, especially in fire-prone areas. However, apart from United States of America and NW Spain, post-fire soil stabilization measures are not being widely implemented in other fire-prone regions. During the last decade, a protocol of these urgent measures, applied by land forest managers in the integrated fight against wildfires, has been elaborated and annually implemented by forest managers in the temperate humid zones (Galicia, NW Spain) (Vega *et al.*, 2013a). The most effective and widely used technique is wheat straw mulching applied (Robichaud, Beyers and Neary, 2000; Vega *et al.*, 2013a; Fernández *et al.*, 2019a), which has shown an immediate effectiveness in increasing ground cover and reducing soil erosion losses during the first months after the fire in areas burned at high severity. Straw mulching can be spread over the soil surface via ground (hand) or aerial (helicopter) applications (Figure 4). All soil erosion mitigation measures are costly treatments (more than 3000 €/ha, 2020), but straw mulching has the highest benefit-cost ratio and its use is limited to a reduced extension of burnt surface areas, that are highly susceptible to suffer soil erosion (for example, high severity fires, great extension of burnt areas, sloping terrain, abundant high-intensity rainfall events following wildfires, proximity to surface and subsurface waters). This protocol implemented in Galicia is specific for this temperate humid zone (NW Spain). Following these emergency measures, medium- and long-term rehabilitation and restoration strategies for the recovery of the burnt forest ecosystems (soil – microorganism - tree vegetation) should apply. A detailed description of this long-term process of the recovery of the burnt forest ecosystems as well as the temporal context of measures implemented is shown in Figure 4 (Vega *et al.*, 2013a).




Figure 4. Temporal context of the rehabilitation and restoration strategies for the recovery of the burnt forest ecosystems

Source: modified from Vega *et al.* (2013a)

### 3. Impact on soil organic carbon stocks

Degraded burnt forest soils located in the temperate humid zone (with low levels of OC) can contribute notably to global C sequestration since they have a high potential for storage of organic C when they are subjected to soil recovery techniques to restore their pre-fire quality. Emergency strategies such as mulching application on the burnt soil surface are the first step to stabilize the soil, to avoid C loss and later on to increase C storage following medium- and long-term rehabilitation and restoration strategies. It should be noted, however, that soil potential for C sequestration is finite and limited by soil depth (marked effects only in the first 0–10 cm). Moreover, specific environmental conditions (soil type, farming or forestry system and climate) are also determinants in C sequestration because the soil organic matter (OM) content at equilibrium depends on the interaction of factors as OM inputs, rates of endogenous SOM and exogenous OM mineralization, soil texture and climate (Johnston, Poulton and Coleman, 2009).

Available data on SOC changes in the short term after the fire and mulching application cover are still scarce and provided by local studies (e.g. Díaz-Raviña *et al.*, 2012; Fontúrbel *et al.*, 2012; Berryman *et al.*, 2014; Fernández-Fernández *et al.*, 2016; Lucas-Borja *et al.*, 2019a). The summary of the results of the experiments longer than 1 year are shown in Table 22. In most cases, the absence of short-term effects of mulch application on soil carbon content is the most common result. However, most of the changes in carbon stocks as a consequence of wildfire depend on fire severity and initial carbon content (Vega *et al.*, 2013b) and there are not consistent data about the effects of mulch application on high-severity burnt soils on soil carbon storage. The most determinant short-term effect of mulching treatments on the carbon cycle is related to the significant reduction in soil erosion losses (Robichaud *et al.*, 2013b; Fernández, Vega and Fontúrbel, 2016a) of up to 95 percent and in consequence the reduction in the loss of soil carbon. However, following the fire the C losses by



erosion have rarely been quantified. Gómez-Rey *et al.* (2013b) measured 10-fold lower carbon losses in mulched soils compared to untreated one in a moderately burnt soil in NW Spain. Pierson *et al.* (2019) reported that mulching treatments reduced C and N losses by up to 75 percent compared with untreated areas in different areas burnt at high-severity in the western United States of America. The information on carbon sequestration on vegetation is still very scarce and inconclusive (Fernández, 2021).

**Table 22.** Changes in soil organic carbon stocks reported for soils affected by forest fires after mulching treatments in different environments

Location	Climate zone	Soil type	Control C content OC percent (sd)	Additional C storage OC percent (sd)	Duration (Years)	Depth (cm)	Treatment after fire	Reference
SE Washington (United States of America)	Cool temperate moist	Ashy silt loam (Limberjim Series) Alfic Udivitrand	2.39 (0.12)	3.07 (0.91) p<0.01	4	0-10	Wheat straw mulching	Berryman <i>et al.</i> (2014)
				3.40 (0.27) p<0.01		0-10	Wood strands mulching	
Galicia (NW Spain)	Temperate	Udorthent	12.1 (1.3)	15.4 (0.2) nd		0-2	Seeding	Díaz Raviña <i>et al.</i> (2018)
				17.1 (0.1) nd		0-2	Straw mulching	
			11.6 (0.9)	12.9 (0.2) nd	2-5	Seeding		
				13.6 (0.2) nd	2-5	Straw mulching		

\*burnt soil, no treatment after fire; nd: no statistical differences compared with the control; sd: standard deviation into brackets

Only those experiences longer than 1 year are reported

The SOC sequestration during the whole process of the rehabilitation and restoration of burned forest ecosystems is long and depends on the persistence of negative soil effects induced by wildfires (more than 10 years) (Prieto-Fernández, Acea and Carballas, 1998) as well as the age of the pre-fire vegetation (around 5 years in a shrubland, 20-40 years in a pine forest or more than 100 years in an oak forest). In fact, according to our knowledge, there are no studies about SOC sequestered in restored burned forest ecosystems after such a long time period. In addition, the high spatial variation in OC content values observed for forest ecosystems (Hope *et al.*, 2015; Díaz-Raviña *et al.*, 2018) makes the quantification of SOC sequestered difficult. We hypothesize that the soil potential of additional C storage in the temperate humid zone during the very long restoration and rehabilitation process of these burnt soils can contribute notably to global soil C storage. This is coincident with the findings of Barreiro, Bååth and Díaz-Raviña (2016) in a laboratory study performed with a burnt acidic soil following the incorporation of different mulching plant materials with a high C/N ratio (72-680, milled materials) into the soil by mixing. The data showed clearly an increased microbial activity, especially the growth of fungi, which were negatively affected immediately after the fire and have a higher potential for C sequestration

than bacteria. Under field conditions, we are unaware of the existence of experimental plots where these long-term experiments can be carried out. Nevertheless, the study of Díaz-Raviña *et al.* 2018 (Table 22) allows us to quantify the maximum potential of organic C storage for the burnt soil using as reference the organic C of unburnt control soil (climax vegetation, 26.6 percent and 19.5 percent organic C in 0-2 cm and 2-5 cm soil depth, respectively). Thus, in principle, if the adequate rehabilitation and restoration strategies are implemented, around 14.4 percent and 7.9 percent organic C could be theoretically sequestered in 0-2 cm and 2-5 cm soil depth, respectively.

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

The available information on the effects of mulching treatments on soil properties in field studies is still scarce. In addition, monitoring of the recovery of the burnt ecosystem is limited to a short-time period (up to one year when straw mulching still remains over the soil surface and it is not incorporated into the soil and decomposed). The availability of information for the medium- and long-term is even more scarce (Díaz-Raviña *et al.*, 2018). In different ecosystems, straw mulching has been demonstrated to have favorable short-term effects on soil physical, chemical and microbial properties (Bautista *et al.*, 1996, 2009; Kribeche *et al.*, 2013; Lucas-Borja *et al.*, 2019a), although some negative effects or absence of changes have been also indicated (Lucas-Borja *et al.*, 2019a). These beneficial effects are related to the ability of mulch residues to form a soil cover surface that conserves soil water and attenuates temperature changes, provides stable organic matter and nutrients, improves soil structure, and thus stimulates microbial communities. After fires, the effects of mulching on soil properties depend on soil type and previous soil properties, vegetation type and cover, fire severity and climate and environmental conditions. Table 23 shows the changes in soil properties that have been analyzed in different biomes and conditions.

Studies conducted in central and southeastern Spain, in arid and semi-arid Mediterranean ecosystems affected by wildfire reported increases of soil moisture and infiltration capacity and reductions of soil penetration resistance between one and two years after mulching treatments (Bautista, Bellot and Vallejo, 1996; Bautista, Robichaud and Blade, 2009; Kribeche *et al.*, 2013; Santana, Alday and Baeza, 2014). Prats *et al.* (2013) also report increases in soil moisture and soil shear strength and reductions in soil water repellency in a burnt area in central Portugal treated with hydromulching. In contrast, in severely burnt soils in NW Spain, no significant effect of straw mulch or wood strands mulch was observed in soil shear strength and soil penetration resistance (Fernández *et al.*, 2011; Fernández and Vega, 2021). In this region, and after a high severity fire, Díaz-Raviña *et al.* (2012), also found no change in aggregate stability, soil moisture, water-holding capacity or water repellency in soils treated with mulching and seeding.

The study of Pereira *et al.* (2018) summarizes the impacts of post-fire soil rehabilitation treatments on soil chemical and microbial properties during the first year after experimental and wildland fires in NW Spain, when the soil is most vulnerable to disturbance. The authors indicate that, in general, the effects of mulch application under field conditions are barely evident in most cases (Díaz-Raviña *et al.*, 2012, 2018; Fontúrbel *et al.* 2012; Gómez-Rey *et al.*, 2013a; Gómez-Rey and González-Prieto, 2014, 2015; Barreiro *et al.*, 2015; Lombao *et al.*,

2015). The fires considered were of low-moderate severity, except in Díaz-Raviña *et al.* (2012, 2018) and Gómez-Rey and González-Prieto (2014, 2015) studies, which were of high severity. However, other studies conducted after moderate to high-severity fires in different environmental conditions report positive effects of post-fire rehabilitation treatments. Thus, Kribeche *et al.* (2013) observed increases in soil respiration consistent with the improvement of physical properties under pine forests in SE Spain, one year after seeding and mulching treatments. Berryman *et al.* (2014) reported increases in total N and microbial respiration of wood stakes in a mixed-conifer forest in the United States of America treated with wheat straw and wood strands. Lucas-Borja *et al.* (2019a) also observed increases in soil C, pH, basal respiration, microbial biomass and enzyme activities after mulch application in a Mediterranean pine forest.

The above results highlight the importance of site characteristics and environmental conditions in the response of soil to post-fire rehabilitation treatments. In arid and semi-arid environments, the application of mulch or other forest residue materials can improve post-fire microenvironmental conditions more effectively than in temperate ecosystems, which typically have higher vegetation cover. However, in all cases there is still an urgent need of long-term assessments.

In the temperate humid zone (NW Iberian Peninsula) field studies performed at a longer time scale (4-8 years) showed that significant changes after straw mulching cover in both soil quality as well as vegetation recovery (vegetation cover and species composition) were not detectable (Díaz-Raviña *et al.*, 2018; Fernández, 2021). However, Morgan *et al.* 2014 in a field study performed with a burnt forest ecosystem located in Washington (United States of America) observed that mulch treatment apparently influenced plant cover and diversity up to six years following its application. At long-term, as was mentioned previously, a higher positive effect of the mulching cover on soil quality and hence on soil C storage is expected to be observed in humid conditions than in arid and semiarid ones. Maintenance/conservation of soil quality due to a reduction in soil losses after pre-emergence soil stabilization measures will improve soil quality if these measures are accompanied for those soil restoration and rehabilitation strategies at medium- and long term (Cerdá and Robichaud, 2009). This is supported by the findings of Barreiro *et al.* (2016) in a previous laboratory experiment.

**Table 23.** Results of studies about the short-term effects of post-fire straw mulching on soil physical, physicochemical, chemical, biochemical and microbial properties, analyzed in different biomes and climate conditions

Ecosystem/vegetation	Soil parameter	Change (respect to burnt untreated)	Reference
Physical properties			
SE Spain, semiarid ecosystem/pine forest	Moisture Penetration resistance	Increase Decrease	Bautista, Bellot and Vallejo (1996) Bautista, Robichaud and Blade (2009)
NW Spain, oceanic climate/shrubland	Shear strength	None	Fernández <i>et al.</i> (2011)
NW Spain, Mediterranean climate/pine forest	Aggregate stability Moisture Water repellency Water-holding capacity	None None None None	Díaz-Raviña <i>et al.</i> (2012, 2018)
Western United States of America, continental climates/coniferous forest	Water repellency	None and Decrease	Robichaud <i>et al.</i> (2013a)
SE Spain, semiarid ecosystem/pine forest	Infiltration capacity Penetration resistance	Increase Decrease	Kribeche <i>et al.</i> (2013)
SE Spain, Mediterranean climate/ Abandoned old-field terraces	Moisture Soil temperature	Increase Decrease	Santana, Alday and Baeza (2014)
NW Spain, oceanic climate/pine stands, shrubland	Aggregate stability (dry mean weight diameter)	None	Fernández <i>et al.</i> (2016a)
Central Spain, semiarid ecosystem	Moisture Hydraulic conductivity	Increase Decrease	Lucas-Borja <i>et al.</i> (2019b)
NW Spain	Penetration resistance Shear strength	None None	Fernández and Vega (2021)
Physicochemical properties			
NW Spain, Mediterranean and temperate climate / shrublands and pine stands	pH	None	Barreiro <i>et al.</i> (2015); Díaz-Raviña <i>et al.</i> (2012, 2018); Fontúrbel <i>et al.</i> (2012);

Ecosystem/vegetation	Soil parameter	Change (respect to burnt untreated)	Reference
			Gómez-Rey <i>et al.</i> (2013a); Gómez-Rey and González-Prieto (2014); Lombao <i>et al.</i> (2015)
Central Spain, semiarid ecosystem/pine forest	pH	None	Lucas-Borja <i>et al.</i> (2019a)
NW Spain, Mediterranean and temperate climate / Pine stands and shrublands	Electric conductivity	None	Barreiro <i>et al.</i> (2015); Díaz-Raviña <i>et al.</i> (2012, 2018); Lombao <i>et al.</i> (2015)
Central Spain, semiarid ecosystem/pine forest	Electric conductivity	Increase	Lucas-Borja <i>et al.</i> (2019a)
Chemical properties			
NW Spain, temperate climate /pine plantation and/or shrublands	Organic carbon Total nitrogen	None None	Barreiro <i>et al.</i> (2015); Díaz-Raviña <i>et al.</i> (2018); Gómez-Rey <i>et al.</i> (2013a); Lombao <i>et al.</i> (2015)
Central Spain, semiarid ecosystem/pine forest	Organic carbon Total nitrogen	Increase Increase	Lucas-Borja <i>et al.</i> (2019a)
SE Washington, United States of America, cool temperate moist / mixed-conifer and grand fir	Total carbon Total nitrogen	None Increase	Berryman <i>et al.</i> (2014)
NW Spain, temperate climate/pine plantation and/or shrubland	Gross N mineralization, Ammonium immobilization, Nitrification, Nitrate immobilization	None None None Decrease	Gómez-Rey and González-Prieto (2015); Fernández-Fernández <i>et al.</i> (2016)
NW Spain, temperate climate/pine plantation and shrublands	Labile C pools (water soluble C and carbohydrates)	None	Lombao <i>et al.</i> (2015); Díaz-Raviña <i>et al.</i> (2018)
NW Spain, temperate climate/pine stands and shrublands	Macronutrients Trace elements	None None	Gómez-Rey and González-Prieto (2014); Fernández-Fernández <i>et al.</i> (2016)
Biochemical and microbial properties			

Ecosystem/vegetation	Soil parameter	Change (respect to burnt untreated)	Reference
NW Spain, temperate climate / pine stands and/or shrublands	Microbial biomass, Soil respiration, <i>Enzyme activities:</i> --glucosidase, acid phosphatase, urease	None None None None None	Fontúrbel <i>et al.</i> (2012); Lombao <i>et al.</i> (2015); Díaz-Raviña <i>et al.</i> (2018)
SE Spain, semiarid ecosystem/pine forest	Soil respiration	Slight Increase	Kribeche <i>et al.</i> (2013)
Central Spain, semiarid ecosystem/pine forest	Microbial biomass, Soil respiration, <i>Enzyme activities:</i> --glucosidase, acid phosphatase, urease	Increase Increase Increase Increase Increase	Lucas-Borja <i>et al.</i> (2019a)
SE Washington, United States of America, temperate and humid ecosystem/ mixed-conifer and grand fire	Soil respiration of a standard wood substrate	Increase	Berryman <i>et al.</i> (2014)
NW Spain, temperate climate / pine stands and shrublands	Bacterial activity	None	Díaz-Raviña <i>et al.</i> (2018)
NW Spain, temperate climate / pine stands and/or shrublands	Microbial structure (PLFAs pattern) Specific microbial groups biomass (bacteria, fungi, actinobacteria, G <sup>+</sup> and G <sup>-</sup> bacteria)	None None	Barreiro <i>et al.</i> (2015); Díaz-Raviña <i>et al.</i> (2018)
NW Spain/temperate climate/pine forest and/or shrubland	Microbial biomass Microbial functional diversity	Slight increase Slight increase	Fontúrbel <i>et al.</i> (2016)

## 4.2 Minimization of threats to soil functions

Table 24. Soil threats

Soil threats	
Soil erosion and related soil C and nutrient losses in the top layer (0–2 cm, 2–5 cm)	A significant reduction in soil erosion, organic C and nutrient stocks losses is the most important effect of mulch application in burnt soils.
Modification of nutrient cycling and subsequent unbalance of these elements Loss of C and nutrients (macro- and micro-nutrients) in the eroded sediments	No short and long-term effect on N availability was found in burnt soils after mulch application (Gómez-Rey <i>et al.</i> , 2013b; Gómez-Rey <i>et al.</i> , 2014; Jonas <i>et al.</i> , 2019).
Soil biodiversity loss	Small changes in soil microbial structure or microbial community composition. Information about soil biodiversity is scarce and there is not information respect loss biodiversity. In contrast, an increase of soil biodiversity is expected at long-term with increasing organic C content.
Soil water management	In climates with a strong summer hydric stress the mulch cover maintains/increases soil moisture (Santana, Alday and Baeza, 2014; Lucas-Borja <i>et al.</i> , 2019b).

## 4.3 Increases in production (e.g. food/fuel/feed/timber)

By reducing both soil erosion and loss of organic C and nutrients, maintaining thus soil quality, the mulching techniques help to preserve site productivity. Increased plant regrowth after mulching has been reported in dry environments and related to an increase in soil moisture under the mulch cover (Fernández and Vega, 2014; Fernández *et al.*, 2016b). However, information concerning forest productivity is not available and that related to biomass accumulation after fire and mulching is still inconclusive (Fernández, 2021). In the long term, the positive changes in soil quality can be accompanied by changes in forest production (fuel, timber, food).

## 4.4 Mitigation of and adaptation to climate change

These measures have a positive impact on C sequestration when vegetation cover is enhanced and soil carbon sinks are maintained. At long-term, increased organic C and an improvement of all soil properties related to it (water retention, C and nutrient availability, soil structure...) is expected (Barreiro, Bååth and Díaz-Raviña,

2016). The magnitude of this increase in C storage should be strongly related with the initial pre-fire soil quality and effectiveness of restoration and rehabilitation strategies implemented (Cerdá and Robichaud, 2009). These positive changes in soil quality would be accompanied by changes in forest production (for example, fiber, fuel, food, honey, mushrooms, aromatic plants, nuts, truffles, forest berries) and other activities related to soil ecological functions such as C sequestration in vegetation.

## 4.5 Socio-economic benefits

The application of a mulch cover as an urgent measure to protect the burnt soil avoid significantly soil erosion losses, protecting water bodies and aquatic habitats. Mulching treatment tends to mitigate the floods provoked by post-fire water erosion and avoid eroded sediments reaching the aquatic ecosystems. If the latter process takes place, the burnt sediments can contaminate the surface and underground water and hence diminish the quality of water reservoirs and affect negatively the sea water quality, which may have important consequences on the shellfish industry (death of bivalves by anaerobiosis). Besides that, damage to infrastructures is also avoided. The restoration and rehabilitation techniques of forest ecosystems allow us to restore the value of the forest as a place of leisure and recreation (biodiversity, landscape, mountain villages, culture, gastronomy, *appellation d'origine* products). This is essential for people's health and in nowadays has a great economic potential to combat work stress through different activities related to ecotourism (natural parks, forest with different plants and animals, mountain sports, skiing, climbing, trekking, walks...).

## 4.6 Additional benefits to the practice

The soil, in addition to the production function, also performs ecological functions such as the C sequestration and mitigation of climate change, mentioned in the previous section, water purification and soil contaminant reduction, climate and flood regulation, nutrient cycling, habitat for organisms, source of biodiversity. Therefore, since these urgent measures to stabilize burnt soil try to maintain/improve soil quality of burnt soils, these ecological soil functions are recovered with the subsequent important benefits to maintain the life in the planet.

# 5. Potential drawbacks to the practice

## 5.1 Increases in greenhouse gas emissions

The possible modification of the equilibrium between mineralization (source of C) and humification (sink or C) processes of the organic matter can affect the GHE emissions (CO<sub>2</sub>, CH<sub>4</sub>, NO<sub>2</sub>).



## 5.2 Conflict with other practice(s)

Mulching application after a fire can also reduce the possible impacts of salvage logging (Fernández and Vega, 2016b). Besides that, logging residues can also be used locally as a mulch cover (Fernández *et al.*, 2007), especially when non-commercial stands are affected (Fernández *et al.*, 2019b).

## 5.3 Decreases in production (e.g. food/fuel/feed/timber)

When the application rates of the straw mulch is not adequate, some delay in the recovery of the vegetative cover can occur (Dodson and Peterson, 2010). Inappropriate forestry practices that reduce soil quality will have a negative impact on production.

## 5.4 Other conflicts

These urgent measures need to be accompanied with further good forestry practices for the sustainable burnt soil management. It should be in accordance with current legislation.

# 6. Recommendations before implementation of the practice

Initially, it is necessary to identify the burnt areas susceptible to suffer post-fire erosion and apply the selected criteria to delineate the burnt area for implementation of these measures. To achieve the treatment goals, at least 75 percent of the burnt soil needs to be covered with the mulch material. It is important to check the availability of different materials taking into account that the amount of material to reach the desired cover is variable depending on the residue (from 2.5 t/ha with wheat straw to 10 t/ha with wood strands). The wheat straw is the best option for using as a mulch treatment because it generally had no significant effects on soil quality, but reduced significantly erosion, organic C and nutrient stock losses compared to that of burnt soil control (70-90 percent reduction values) (Vega *et al.* 2013a). In addition, straw mulching has a better cost/benefit ratio that observed in soils with other plant materials mulching (wood strands, wood chips).

## 7. Potential barriers to adoption

Table 25. Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	The accessibility to the soil surface affected by the wildfire (mountainous area, high slope, absence of roads...). In this case, mulching can be applied by helicopter (heli-mulching).
Cultural	Yes	All sectors of society need to be made aware of the serious economic and ecological damage caused by wildfires and hence of the urgent need for soil protection against post-fire erosion. They should also know that the wheat straw mulching, applied manually or by helicopter, is the best option to mitigate these damages. This information has to be disseminated among the population (press, radio, television, web pages ...).
Social	Yes	The perception of the necessity of soil protection is increasing worldwide. A shift into emergency measures after fire is still needed.
Economic	Yes	Availability of funds for urgent measures implementation
Institutional		The support of the administration is necessary: the elaboration of an operating protocol specific for the climate region, the formation of specialized forest managers and the inclusion of these measurements in the integrated fight against forest wildfires (prevention, extinction, impacts, recovery).
Legal (Right to soil)	Yes	The implementation of these practices must be in accordance with current legislation.
Knowledge	Yes	Due the urgent character of these measures immediately after fire, a fire severity analysis is necessary to apply these measures only in high-severity affected soils.
Other	Yes	The adverse effects of wildfires can extend far beyond those areas directly affected by the fire and cause significant ecological and economic damage affecting many sectors of society. Therefore, scientists, administration, forest managers, forest land owners and representatives of both the productive sectors should collaborate in the evaluation of environmental impact, risk assessment and in the preparation of the emergency measures protocol accompanied by medium- and long-term restoration and rehabilitation strategies. Long-term field studies monitoring the C sequestration following the rehabilitation and restoration strategies for the recovery of the burnt forest ecosystems are needed.

## Photos of the practice



Photo 6. Helicopter straw mulching application after wildfire in Galicia (NW Spain)

**Table 26.** Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Straw mulch and biochar application in recently burned areas of Algarve (Portugal) and Andalusia (Spain)</i>	Europe	1	6	10

## References

- Barreiro, A., Fontúrbel, M.T., Lombao, A., Martín, A., Vega, J.A., Fernández, C., Carballas, T. & Díaz-Raviña, M.** 2015. Using phospholipid fatty acid and community level physiological profiling techniques to characterize soil microbial communities following an experimental fire and different stabilization treatments. *Catena*, 135: 419-429. <https://doi.org/10.1016/j.catena.2014.07.011>
- Barreiro, A., Bååth, E. & Díaz-Raviña, M.** 2016. Bacterial and fungal growth in burnt acid soils amended with different C/N mulch materials. *Soil Biology and Biochemistry*, 97: 102-111. <https://doi.org/10.1016/j.soilbio.2016.03.009>
- Bautista, S., Bellot, J. & Vallejo, V.R.** 1996. Mulching treatment for post-fire soil conservation in a semiarid ecosystem. *Arid Soil Research and Rehabilitation*, 10: 235-242. <https://doi.org/10.1080/15324989609381438>
- Bautista, S., Robichaud P.R. & Blade, C.** 2009. Post-fire mulching. In A. Cerda, P.R. Robichaud (Eds.) *Fire effects on soils and restoration strategies*. Science Publishers, Enfield, NH, USA, pp. 353-372.
- Berryman, E.M., Morgan, P., Robichaud, P.R. & Page-Dumroese, D.** 2014. Post-fire erosion control mulches alter belowground processes and nitrate reductase activity of a perennial forb, heartleaf arnica (*Arnica cordifolia*). *Res. Note RMRS-RN-69*. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 10 p., 69. <https://doi.org/10.2737/RMRS-RN-69>
- Cerdá, A. & Robichaud P.R. (Eds.)** 2009. *Fire effects on soils and restoration strategies*. Science Publishers, Enfield, NH, USA.
- Díaz-Raviña, M., Lombao, A., Barreiro, A., Martín, A., Iglesias, L., Díaz-Fierros, F., & Carballas, T.** 2018. Medium term impact of post-fire emergency rehabilitation treatments on a forest ecosystem in Galicia (NW Spain). *Spanish Journal of Soil Science*, 8(3): 322-335. <https://doi.org/10.3232/SJSS.2018.V8.N3.03>
- Díaz-Raviña, M., Martín, A., Barreiro, A., Lombao, A., Iglesias, L., Díaz-Fierros, F. & Carballas, T.** 2012. Mulching and seeding treatments for post-fire stabilisation in N.W. Spain: short-term effects and effectiveness. *Geoderma*, 191: 31-39. <https://doi.org/10.1016/j.geoderma.2012.01.003>
- Dodson, E.K. & Peterson, D.W.** 2010. Mulching effects on vegetation recovery following high severity wildfire in north-central Washington State, USA. *Forest Ecology and Management*, 260: 1816-1823. <https://dx.doi.org/10.1016/j.foreco.2010.08.026>
- Fernández, C.** 2021. Medium-term effects of straw helimulching on post-fire vegetation recovery in shrublands in north-west Spain. *International Journal of Wildland Fire*. <https://doi.org/10.1071/WF20092>
- Fernández, C. & Vega, J.A.** 2014. Efficacy of bark strands and straw mulching after wildfire in NW Spain: Effects on erosion control and vegetation recovery. *Ecological Engineering*, 63: 50-57. <https://doi.org/10.1016/j.ecoleng.2013.12.005>

- Fernández, C. & Vega, J.A.** 2016a. Effects of mulching and post-fire salvage logging on soil erosion and vegetative regrowth in NW Spain. *Forest Ecology and Management*, 375: 46-54.  
<https://doi.org/10.1016/j.foreco.2016.05.024>
- Fernández, C. & Vega, J.A.** 2016b. Modelling the effect of soil burn severity on soil erosion at hillslope scale in the first year following wildfire in NW Spain. *Earth Surface Processes and Landforms*, 41: 928-935.  
<https://doi.org/10.1002/esp.3876>
- Fernández, C. & Vega, J.A.** 2021. Is wood strand mulching a good alternative to helimulching to mitigate the risk of soil erosion and favour the recovery of vegetation in NW Spain? *Landscape and Ecological Engineering*. <https://doi.org/10.1007/s11355-020-00439-2>
- Fernández, C., Vega J.A., Fontúrbel, M.T., Pérez-Gorostiaga, P., Jiménez, E. & Madrigal, J.** 2007. Effects of wildfire, salvage logging and slash treatments on soil degradation. *Land Degradation and Development*, 38(6): 591-607. <https://doi.org/10.1002/ldr.797>
- Fernández, C., Vega, J.A., Jiménez, E. & Fontúrbel, M.T.** 2011. Effectiveness of three post-fire treatments at reducing soil erosion in Galicia (NW Spain). *International Journal of Wildland Fire*, 20: 104-114. <https://doi.org/10.1071/WF09010>
- Fernández, C., Vega, J.A. & Fontúrbel, T.** 2016a. Reducing post-fire soil erosion from the air: Performance of heli-mulching in a mountainous area on the coast of NW Spain. *Catena*, 147: 489-495.  
<https://doi.org/10.1016/j.catena.2016.08.005>
- Fernández, C., Vega, J.A., Fontúrbel, M.T., Barreiro, A., Lombao, A., Gómez-Rey, M.X., Díaz-Raviña, M. & González-Prieto, S.** 2016b. Effects of straw mulching on initial post-wildfire vegetation recovery. *Ecological Engineering*, 95: 138-142.
- Fernández, C., Vega, J.A., Arbones, P. & Fontúrbel, M.T.** 2019a. Eficacia de los tratamientos de estabilización del suelo después de incendio en Galicia. CIF Lourizan. Xunta de Galicia: Santiago de Compostela. (also available at: [https://lourizan.xunta.gal/sites/w\\_forlou/files/libro.pdf](https://lourizan.xunta.gal/sites/w_forlou/files/libro.pdf))
- Fernández, C., Fernández-Alonso, J.M. & Vega, J.A.** 2019b. Effects of mastication of burned non-commercial *Pinus pinaster* Ait. trees on soil compaction and vegetation response. *Forest Ecology and Management*, 449: 117457. <https://doi.org/10.1016/j.foreco.2019.117457>
- Fernández-Fernández, M., Vieites-Blanco, C., Gómez-Rey, M.X. & González-Prieto, S.J.** 2016. Straw mulching is not always a useful post-fire stabilization technique for reducing soil erosion. *Geoderma*, 284: 122-131. <https://doi.org/10.1016/j.geoderma.2016.09.001>
- Fontúrbel, M.T., Barreiro, A., Vega, J.A., Martín, A., Jiménez, E., Carballas, T., Fernández, C. & Díaz-Raviña, M.** 2012. Effects of an experimental fire and post-fire stabilisation treatments on soil microbial communities. *Geoderma*, 191: 51-60. <https://doi.org/10.1016/j.geoderma.2012.01.037>
- Gómez-Rey, M.X., Couto-Vázquez, A., García-Marco, S. & González-Prieto, S.J.** 2013a. Impact of fire and post-fire management techniques on soil chemical properties. *Geoderma*, 195-196: 155-164.  
<https://doi.org/10.1016/j.geoderma.2012.12.005>

- Gómez-Rey, M.X., Couto-Vázquez, A., García-Marco, S., Vega, J.A. & González-Prieto, S.J.** 2013. Reduction of nutrient losses with eroded sediments by post-fire soil stabilisation techniques. *International Journal of Wildland Fire*, 22(5): 696–706. <https://doi.org/10.1071/WF12079>
- Gómez-Rey, M.X. & González-Prieto, S.J.** 2014. Short and medium-term effects of a wildfire and two emergency stabilisation treatments on the availability of macronutrientes and trace elements in topsoil. *Science of Total Environment*, 493: 251–261. <https://doi.org/10.1016/j.scitotenv.2014.05.119>
- Gómez-Rey, M.X. & González-Prieto, S.J.** 2015. Soil gross N transformation rates after a wildfire and straw mulch application for burned soil emergency stabilisation. *Biology and Fertility of Soils*, 51: 493–505. <https://doi.org/10.1007/s00374-015-0997-0>
- Hope, G., Jordan, P., Winkler, R., Giles, T., Curran, M., Soneff, K. & Chapman, B.** 2015. Post-wildfire Natural Hazard Risk Analysis in British Columbia. Prov. B.C., Victoria, B.C. Land Management Handbook, 69. (also available at: <https://www.for.gov.bc.ca/hfd/pubs/docs/lmh/lmh69.pdf>)
- Johnston, A.E., Poulton, P.R. & Coleman, K.** 2009. Soil organic matter: its importance in sustainable agriculture and carbon dioxide fluxes. In Sparks, D.L. (Ed.) *Advances in Agronomy*, 101, pp. 1–57. Academic Press. [https://doi.org/10.1016/S0065-2113\(08\)00801-8](https://doi.org/10.1016/S0065-2113(08)00801-8)
- Jonas, J.L., Berryman, E., Wolk, B., Morgan, P. & Robichaud, P.R.** 2019. Post-fire wood mulch for reducing erosion potential increases tree seedlings with few impacts on understory plants and soil nitrogen. *Forest Ecology and Management*, 453: 117567. <https://doi.org/10.1016/j.foreco.2019.117567>
- Kribeche, L., Bautista, S., Gimeno, T., Blade, C. & Vallejo, V.R.** 2013. Evaluating the Effectiveness of Post Fire Emergency Rehabilitation Treatments on Soil Degradation and Erosion Control in Semi-Arid Mediterranean Areas of the Spanish South East. *Arid Land Research and Management*, 27: 361–376. <https://doi.org/10.1080/15324982.2013.771229>
- Lliteras, M.T.F., Filgueira, C.F. & Hidalgo, J.A.V.** 2016. Efectos a medio plazo de tratamientos de rehabilitación post-incendio en propiedades microbiológicas del suelo. *Cuadernos de la Sociedad Española de Ciencias Forestales*, (42): 111–128.
- Lombao, A., Díaz-Raviña, M., Martín, A., Barreiro, A., Fontúrbel, M.T., Vega, J.A., Fernández, C. & Carballas T.** 2015. Influence of straw mulch application on the properties of a soil affected by a forest wildfire. *Spanish Journal of Soil Science*, 5: 26–40. <https://doi.org/10.3232/SJSS.2015.V5.N1.03>
- Lucas-Borja, M.E., Plaza-Álvarez, P.A., Ortega, R., Miralles, I., González-Romero, J., Sagra, J., Moya, D., Zema, D.A. & de las Heras, J.** 2019a. Short-term changes in soil functionality after wildfire and straw mulching in a *Pinus halepensis* M. forest. *Forest Ecology and Management*, 457: 117700. <https://doi.org/10.1016/j.foreco.2019.117700>
- Lucas-Borja, M. E., Zema, D.A., Gianmarco Carrá B., Cerdá, A., Plaza-Álvarez, P.A., Sagra, J., González-Romero, J., Moya, D. & de las Heras, J.** 2019b. Short-term changes in infiltration between straw mulched and non-mulched soils after wildfire in Mediterranean forest ecosystems. *Ecological Engineering*, 122: 27–31. <https://doi.org/10.1016/j.ecoleng.2018.07.018>

- Morgan, P., Moy, M., F, Droske CH.A., Lentile, L.B., Lewis, S.A., Robichaud, P.R. & Hudak, A.T.** 2014. Vegetation response after post-fire mulching and native grass seeding. *Fire Ecology*, 10: 49-62. <https://doi.org/10.4996/fireecology.1003049>
- Pereira, P., Francos, M., Brevik, E.C., Ubeda, X. & Bogunovic, I.** 2018. Post-fire soil management. *Current Opinion in Environmental Science & Health*, 5: 26–32. <https://doi.org/10.1016/j.coesh.2018.04.002>
- Pierson, D.N., Rochibaud, P.C., Rhoades, C.C. & Brown, R.E.** 2019. Soil carbon and nitrogen eroded after severe wildfire and erosion mitigation treatments. *International Journal of Wildland Fire*, 28: 814-822. <https://doi.org/10.1071/WF18193>
- Prats, S.A., Malvar, M.C., Vieira, D.C.S., MacDonald, L. & Keizer, J.J.** 2013. Effectiveness of hydromulching to reduce runoff and erosion in a recently burnt pine plantation in central Portugal. *Land Degradation & Development*, 27(5): 1319–1333. <https://doi.org/10.1002/ldr.2236>
- Prieto-Fernández, A., Acea, M.J. & Carballas, T.** 1998. Soil microbial and extractable C and N after fire. *Biology and Fertility of Soils*, 27: 132-142. <https://doi.org/10.1007/s003740050411>
- Robichaud, P.R., Beyers, J.L. & Neary, D.G.** 2000. *Evaluating the Effectiveness of Postfire Rehabilitation treatments*. Gen. Tech. Rep. RMRS-GTR-63. Fort Collins: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 85 p.
- Robichaud P.R., Lewis S.A., Wagenbrenner J.W., Ashum, L.E. & Brown R.E.** 2013a. Post-fire mulching for runoff and erosion mitigation Part I: Effectiveness at reducing hillslope erosion rates. *Catena*, 105: 75-92. <https://doi.org/10.1016/j.catena.2012.11.015>
- Robichaud, P.R., Wagenbrenner, J.W., Lewis, S.A, Ashmun, L.E, Brown, R.E. & Wohlgemuth, P.M.** 2013b. Post-fire mulching for runoff and erosion mitigation Part II: Effectiveness in reducing runoff and sediment yields from small catchments. *Catena*, 105: 93-111. <https://doi.org/10.1016/j.catena.2012.11.016>
- Santana, V.M., Alday, J.G. & Bacza, M.J.** 2014. Mulch application as post-fire rehabilitation treatment does not affect vegetation recovery in ecosystems dominated by obligate seeders. *Ecology Engineering*, 71: 80–86. <https://doi.org/10.1016/j.ecoleng.2014.07.037>
- Vega, J.A., Fontúrbel, M.T., Fernández, C., Arellano, A., Díaz-Raviña, M., Carballas, M.T., Martín, A., González-Prieto, S., Merino, A. & Benito, E.** 2013a. Acciones urgentes contra la erosión en áreas forestales quemadas: Guía para su planificación en Galicia. ISBN: 978-84-8408-716-8. pp. 139. [http://fuegored.weebly.com/uploads/2/2/2/8/22283836/guia\\_planificacion\\_galicia.pdf](http://fuegored.weebly.com/uploads/2/2/2/8/22283836/guia_planificacion_galicia.pdf)
- Vega, J.A., Fontúrbel, M.T., Merino, A., Fernández, C., Ferreiro, A. & Jiménez, E.** 2013b. Testing the ability of visual indicators of soil burn severity to reflect changes in soil chemical and microbial properties in pine forests and shrubland. *Plant and Soil*, 369: 73-91. <https://doi.org/10.1007/s11104-012-1532-9>
- Vega, J.A., Fernández, C. & Fontúrbel, M.T.** 2018. Medidas de atenuación de los daños post-incendio en Galicia. In Díaz-Fierros F (Ed.) *Incendios Forestales: Reflexiones desde Galicia*. Hércules de Ediciones, Santiago de Compostela, pp 136-174.



## 8. Forest landscape restoration

Blanca Bernal

Winrock International, Arlington, United States of America

### 1. Description of the practice

Forest Landscape Restoration (FLR) is the process of regaining ecological integrity and functionality to enhance human well-being and livelihoods across deforested or degraded landscapes, bringing back through restoration the biological productivity of an area (IUCN and WRI, 2014; Lamb, Stanturf and Madsen, 2012; Schultz, Jedd and Beam, 2012). FLR does not necessarily seek to convert deforested and degraded lands to forests, but to balance landscape functions by integrating forests and woodlands strategically in a mosaic of land uses (Mansourian, Vallauri and Dudley, 2005), often assessing present, past, and reference land uses to assess the feasibility of restoring landscape functionality (Schulz and Schröder, 2017). A successful FLR practice would address the drivers of deforestation and landscape degradation, which often includes implementation of sustainable management of non-forest landscape units such as improved management of pastures or conservative agriculture (Mansourian, Vallauri and Dudley, 2005; Stanturf, Mansourian and Kleine, 2017). This landscape-wide approach to restoration thereby covers multifunctional and interdependent land uses and socioeconomic activities that are reintegrated as a mosaic landscape (Photo 8) to recover degraded lands with the purpose of balancing the provision of goods and ecosystem services (Hobbs, 2002; Mansourian, Vallauri and Dudley, 2005; Schulz and Schröder, 2017), thereby differing from traditional site-specific restoration efforts and *ad hoc* treatments (Lamb, Stanturf and Madsen, 2012; Stanturf *et al.*, 2015).

FLR has guiding principles that define its dynamic nature (IUCN and WRI, 2014), namely to manage adaptively for long-term resilience, tailor to local conditions using a variety of approaches, focus on the landscape level, maintain and enhance natural ecosystems, restore ecosystem multi-functionality, allow for multiple benefits, and involve stakeholders to support participatory landscape governance (Zhou *et al.*, 2008).

FLR practices are diverse and can include one or more of the following activities (IUCN and WRI, 2014; Maginnis, Rietbergen-McCracken and Sarre, 2007):

**Table 27.** Example of forest landscape restoration (FLR) activities across the landscape

In forest land	In agricultural land	In protective land and buffers
<p><b>Afforestation/reforestation:</b> Planting of forests and woodlots.</p> <p><b>Natural forest regeneration:</b> Passive/assisted planting on degraded forests and marginal agricultural sites.</p> <p><b>Silviculture and improved forest management:</b> Rehabilitation of degraded primary forests, management of secondary forests.</p>	<p><b>Agroforestry and on-farm trees:</b> Multi-strata crops, live fences and windbreakers, intercropping, agrosilvopastoral systems, tree gardens.</p> <p><b>Improved fallow:</b> establishment/management of trees or shrubs on fallow and shifting cultivation land.</p> <p><b>Improved rangeland and cropland:</b> sustainable management to increase soil carbon and ecosystem functionality.</p>	<p><b>Mangrove restoration:</b> natural regeneration through hydrological restoration, and/or mangrove planting.</p> <p><b>Revegetation for protection and erosion control:</b> creation of riparian buffers, floodplain reconnection, slope revegetation.</p>

FLR is therefore more than planting trees and restoring forests or implementing agroforestry, yet these activities are often a key component of FLR approaches. A focused review of Forest Restoration (i.e. Afforestation, Reforestation, Assisted Natural Regeneration, and Restoration of Mangroves) is available in Factsheet n° 6 of this volume, whereas in Volume 3, factsheets n° 38, 39 and 40 reviews Agroforestry practices. Improved agriculture practices such as crop rotation and diversification, soil organic cover, or other conservative agriculture practices are described in detail in Volume 3 of this manual.

## 2. Range of applicability

There are 2.2 billion hectares of degraded land around the world that have FLR potential (Figure 5), particularly mosaic landscapes (Minnemeyer *et al.*, 2011; Stanturf *et al.*, 2015) and strategic areas where restoration maximizes ecological gains (Lamb, Stanturf and Madsen, 2012) and/or where diverse landscape functions can be enhanced (Schultz, Jedd and Beam, 2012).

Numerous studies show multiple benefits associated to FLR practices (Chavez-Tafur and Zagt, 2014; Corbeels *et al.*, 2019; Gourevitch *et al.*, 2016; Lal, 2004; Nave *et al.*, 2013; Schultz, Jedd and Beam, 2012; Stanturf *et al.*, 2015; Stanturf, Lamb and Madsen, 2012) due to the increase in vegetative cover and the restoration of landscape functionality across a variety of landscapes and climates. The benefits of this practice are also identified in the restored soils, namely the enhancement of carbon sequestration and storage, improvement of water quality on the watershed, increase of ecosystem health and productivity, and increase of biodiversity. These are described in more detail in section 4 below. Based on these soil benefits, priority FLR locations to improve soil conditions and increase landscape functionality are summarized in Table 28.




Figure 5. Global forest landscape restoration opportunities

Source: Stanturf *et al.* (2015), adapted from Bastin *et al.* (2019). "Remote restoration" refers to areas with less than one person per km<sup>2</sup> in a 500 km radius; "wide-scale restoration" refers to areas with over

Table 28. Priority FLR areas to improve soil aspects of landscape functionality across climatic regions and geographies

Priority FLR site	Potential soil-based FLR benefits
<b>Steep slopes</b>	Protect erosion-prone soils, improve soil structure and aggregation
<b>Riparian strips</b>	Protect erosion-prone soils, filter and trap runoff and sediments, improve soil structure and aggregation, improve soil water retention
<b>Areas prone to sheet erosion and with compacted soils</b>	Protect erosion-prone soils and increase infiltration capacity, increase soil aeration, improve soil structure, improve soil biodiversity
<b>Groundwater recharge areas</b>	Increase evapotranspiration and water table depth
<b>Coastal protection zones</b>	Decrease soil loss and erosion, increase sedimentation and soil aggregation
<b>Agrosilvopastoral systems</b>	Increase land productivity, reduce erosion, increase water holding capacity, improve soil structure

Source: Adapted from Lamb, Stanturf and Madsen (2012)

### 3. Impact on soil organic carbon stocks

**Table 29.** Changes in soil organic carbon stocks reported for forest landscape restoration projects

Location	Climate zone	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (years)	More information	Reference
Jiangxi, China	Subtropical moist	26.6, top 40 cm	0.77	19	Broadleaf afforestation in degraded land	Zhou <i>et al.</i> (2008)
			0.35		Conifer afforestation in degraded land	
Copenhagen, Denmark	Cool temperate moist	15.0, top 5 cm	0.36	29	Spruce afforestation in abandoned cropland	Vesterdal, Ritter and Gundersen (2002)
Guangdong, China	Tropical moist	4.0, top 20 cm	0.17	56	Eucalyptus sp. plantation in degraded land	Zhang <i>et al.</i> (2019)
			0.81		Broadleaf mixed secondary forest in degraded land	
Perth, Australia	Warm temperate dry	43.3, top 30 cm	0.15	26	Eucalyptus sp. reforestation in former cropland	Harper <i>et al.</i> (2012)
Nebraska, United States of America		36.2, top 15 cm	0.11	35	Windbreak agroforestry	Sauer, Cambardella and Brandle (2007)
Ontario, Canada	Cool temperate moist	68.5, top 20 cm	1.34	13	Alley cropping agroforestry	Peichl <i>et al.</i> (2006)
Limón, Costa Rica	Tropical wet	94, top 100 cm	4.63	16	Silvopastoral system	Amézquita <i>et al.</i> (2004)
Puntarenas, Costa Rica	Tropical dry	129, top 100 cm	0.10	10		
Florida, United States of America	Subtropical moist	0.6, top 30 cm	2.44	19	Planted mangroves in barren land	Osland <i>et al.</i> (2012)
Louisiana, United States of America		19.3, top 10 cm	0.12	17	Mangrove natural regeneration in marsh	Henry and Twilley (2013)

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Implementing FLR in degraded landscapes ultimately increases organic matter (SOM) inputs to the soil in the long term, and a decrease in losses of soil and organic matter (Sanderman and Baldock, 2010). This net SOM increase drives FLR's improvement of soil properties, specifically by increasing nutrients and fertility, water holding capacity, and soil health and biodiversity (Lal, 2004; van Noordwijk *et al.*, 1997). Further, SOM improves soil aggregation, which in turn reduces erosion and soil loss (Harper *et al.*, 2012; van Noordwijk *et al.*, 1997; Zhou *et al.*, 2008), an important FLR benefit at the landscape level. Although difficult to quantify, an study estimated that 50 percent of the aggregate carbon formed in just 2.5 decades in Midwestern United States of America (Wick, Ingram and Stahl, 2009), while a 1 percent increase of SOM resulted in 18 percent higher average aggregate size in a West Indies site (Blanchart *et al.*, 2004). Moreover, Canada's government soil data (Stone and Hilborn, 2012) shows that increases in SOM content produce proportional changes in the erodibility factor  $K$  (average soil loss in tons per hectare) of the Universal Soil Loss Equation (USLE).

### 4.2. Minimization of threats to soil functions

Table 30. Soil threats

Soil threats	
<b>Soil erosion</b>	SOM favors aggregation, reducing risks of erosion. Vegetation cover reduces the impact and damage to the soil of erosion agents such as runoff or wind (Harper <i>et al.</i> , 2012; Lal, 2004; Zhou <i>et al.</i> , 2008).
<b>Nutrient imbalance and cycles</b>	Organic matter is rich in nutrients, fertilizing the soil. In the soil, it increases both the fixation of ions or nutrients and soil microbial diversity, fostering nutrient cycling (Lal, 2004; Osland <i>et al.</i> , 2012).
<b>Soil salinization and alkalinization</b>	FLR improves water balance in the watershed, reducing risks of salinization (Harper <i>et al.</i> , 2012). Vegetation reduces soil temperature and thus evaporation. On occasion, however, trees can accelerate surface water loss (Zhang and Shangguan, 2016) and could potentially increase soil salinity.
<b>Soil contamination / pollution</b>	Similar to the nutrient cycling, vegetation and SOM increases cycling and transformation of elements, filtering pollutants and reducing their accumulation in the soil (Lal, 2004; Zhang <i>et al.</i> , 2019).
<b>Soil biodiversity loss</b>	FLR improves soil health and biodiversity thanks to increased organic inputs and active rhizospheres (Gourevitch <i>et al.</i> , 2016; Harper <i>et al.</i> , 2012).
<b>Soil sealing</b>	Vegetation aerates the soil and reduces ponding (Zhang and Shangguan, 2016) and intercepts runoff and sediments (Lal, 2004), decreasing the risk of soil sealing with FLR.

Soil threats	
Soil compaction	Belowground biomass and SOM increases soil aeration and porosity, reducing soil compaction (Harper <i>et al.</i> , 2012; Nair <i>et al.</i> , 2009).
Soil water management	FLR increases water infiltration capacity (Lamb, Stanturf and Madsen, 2012) and improves water cycling as trees increase evapotranspiration and SOM increases water holding capacity (Lal, 2004; Zhang and Shangguan, 2016).

### 4.3. Increases in production (e.g. food/fuel/feed/timber)

FLR, albeit diverse and context-dependent, can increase the production of wood products, non-timber forest products, bioenergy materials, and game abundance and diversity (Stanturf, Mansourian and Kleine, 2017). The selection of species to bring back to the landscape is driven by the function(s) that are wanted to recover, the local customs on what species to grow, and the economic limitations on materials and/or species available. The functions to recover do not necessarily compete but they often do; for example, FLR for timber production (e.g. woodlots) is not best suited for food production or biodiversity support (Daoxiang *et al.*, 2015), while agroforestry, on the other hand, has great potential to combine fuelwood production with food and feed production (Nair *et al.*, 2009; Stanturf, Mansourian and Kleine, 2017). Because FLR looks at the entire landscape or watershed unit, a mosaic of interventions that cover a variety of functions allows to combine benefits and goods generated beyond a single field site, e.g. integrating woodlots, crops, pastures, and riparian buffers or natural forests with improved management practices, among a myriad of options.

### 4.4. Mitigation of and adaptation to climate change

Unsustainable removal of trees and other vegetation from the landscape or halting vegetation growth will inevitably yield a net emission of GHG if they result in deforestation and/or degradation (Stanturf, Mansourian and Kleine, 2017). Because FLR entails the recovery of efficient carbon-sequestering vegetation and the concomitant increase in soil carbon sink capacity, it represents an important climate change mitigation strategy, as enhancing vegetation cover in the landscape increases carbon capture from the atmosphere and sediment retention in the watershed.

Studies show that SOM increases its recalcitrance under FLR when woody species like trees and shrubs are integrated in the landscape (van Noordwijk *et al.*, 1997; Zhang *et al.*, 2019), a process that allows FLR to enhance SOM's long term carbon sink capacity and ability to be a carbon pool more resilient than biomass (Nair *et al.*, 2009; Vesterdal, Ritter and Gundersen, 2002). FLR is a key climate change adaptation strategy, particularly in coastal areas and floodplains where tree coverage reduces flooding as well as the impact of severe storms and accelerated sea-level rise (IUCN and WRI, 2014; Stanturf *et al.*, 2015; Stanturf, Mansourian and Kleine, 2017). By balancing land uses at the landscape level, FLR increases the resilience of the land and the communities that live and depend on it.

## 5. Potential drawbacks to the practice

FLR is usually in conflict with conventional agriculture and urban expansion (Minnemeyer *et al.*, 2011), land uses usually more economically productive whose focus is rarely environmental enhancement. While implementing FLR would have negligible negative environmental impacts compared to these more profitable land uses, replacing FLR with conventional agriculture (e.g. tillage, fertilizer application, and/or irrigation) would result in the soil threats listed above (Section 4) no longer being prevented or minimized. Developed lands can provide more economic benefits to a region in the short-term than FLR, yet smallholders might not necessarily experience a significant income increase with development expansion. Furthermore, a land use that can be defined as “profitable” because it provides high immediate economic benefits despite its environmental impact is likely to have significant economic costs on the long term, unless it is adapted to become resilient (Wei *et al.*, 2020). The landscape and its communities are therefore usually less resilient without FLR (IUCN and WRI, 2014; Stanturf, Mansourian and Kleine, 2017). However, where conventional agriculture is replaced by climate smart or conservative agriculture, which can include agroforestry implementation, some of the soil and ecosystem benefits FLR brings to the landscape would be provided. Natural components in the landscape replaced with urban settings, on the other hand, will virtually eliminate most FLR benefits even under urban forestry implementation. To balance both environmental and socioeconomic needs across the landscape FLR proposes a landscape view that goes beyond a single site, where trade-offs are assessed and land uses are placed strategically in the landscape to maximize multifunctionality (Mansourian, Vallauri and Dudley, 2005; Schulz and Schröder, 2017).

Alternative and more profitable land uses or practices that remove or degrade carbon stocks in the landscape will result in a net GHG emission and a loss of that sink capacity, unless the carbon in the landscape is maintained and enhanced using alternative approaches or balancing land uses.

## 6. Recommendations before implementation of the practice

Because of the multiple factors driving landscape degradation and the diverse actors to involve in a successful FLR implementation, wide-scale recovery of degraded landscapes is challenging. Some of these barriers for adoption and recommendations to overcome them are described below (see Section 7). In addition to these specific ones, there are a number of cross-cutting recommendations to implement FLR:

- ◆ Clearly define the scale of the intervention, functions to recover, target land uses or landscape units, and overall goal of the restoration (Hobbs, 2002).
- ◆ Monitor and evaluate outcomes to implement an adaptive management approach (Lamb, Stanturf and Madsen, 2012).
- ◆ Map stakeholders, intervention trade-offs, and multifunctional landscape hotspots to identify priority approaches that maximize beneficiaries (Schulz and Schröder, 2017).

- ◆ Engage the multiple stakeholders and actors of the landscape to address early their interests facilitates collaborative management and balancing land use trade-offs (Maginnis, Rietbergen-McCracken and Sarre, 2007; Zhou *et al.*, 2008).
- ◆ Ensure capacity to manage conflict and negotiate is available, to successfully integrate and generate consensus around diverse perspectives of what a functional landscape is under the given circumstances and idiosyncrasies (Mansourian, Vallauri and Dudley, 2005).
- ◆ Smallholders, key for FLR in fragmented landscapes, face often implementation challenges; ensuring adequate supply of resources (e.g. seedlings, equipment, guidance) and creating partnerships to support production and distribution of resources facilitates smallholder engagement and efficient implementation of best management practices (Stanturf *et al.*, 2015).

## 7. Potential barriers to adoption

Table 31 describes the main types of barriers to adopt FLR, namely biophysical, social, economic, institutional, legal, and technical, and proposed approaches to overcome them.

**Table 31.** Potential barriers to adoption

Barrier	Explanation	Options to overcome barriers
<b>Biophysical</b>	Herbivory, wildfires, extreme weather, historical drivers of site degradation, and severity of soil degradation prior FLR implementation are barriers to address in early implementation stages (Gong <i>et al.</i> , 2013; Stanturf <i>et al.</i> , 2015; Stanturf, Lamb and Madsen, 2012).	Assessment of current and historical landscape conditions give insights on the drivers of degradation and on the feasibility and potential success of restoration approaches (Hobbs, 2002; Schulz and Schröder, 2017). Integrating potential future scenarios under altered climate can improve the resilience of the FLR strategy.
<b>Social</b>	Involvement of local entities and communities is key for FLR implementation (Zhou <i>et al.</i> , 2008). Smallholders face challenges engaging in restoration when institutional support is scarce and where conflict of interest with unclear management and rights arise (Corbeels <i>et al.</i> , 2019).	Community engagement in planning and implementation, as well as benefit-sharing, can reduce unsustainable resource exploitation and its resulting degradation (Hawkins <i>et al.</i> , 2010; Stanturf, Lamb and Madsen, 2012). Stakeholders need to be mapped to include all perspectives in a trade-offs assessment (Lamb, Stanturf and Madsen, 2012; Mansourian, Vallauri and Dudley, 2005).



Barrier	Explanation	Options to overcome barriers
<b>Economic</b>	Funds beyond planting are often necessary until the ecosystem can self-regenerate (Chavez-Tafur and Zagt, 2014). Furthermore, there can be competing demands for the land that make alternatives to FLR more economically profitable or apparently better suited to support livelihoods (Schedlbauer and Kavanagh, 2008; Stanturf <i>et al.</i> , 2015).	Implementation of FLR plans that alleviate food insecurity and poverty would have a direct positive economic impact at the local level (Smith, 2008). A trade-offs assessment can help identify economic constraints of stakeholders and incentives to overcome them (Mansourian, Vallauri and Dudley, 2005). Potential options to cover implementation and adoption costs are payment for ecosystem services schemes, private sector investment, and/or government support, among others.
<b>Institutional</b>	Successful FLR implementation typically requires engagement of multiple stakeholders, participatory planning, and coordination of multiple institutional levels and agencies (IUCN and WRI, 2014; Stanturf <i>et al.</i> , 2015).	Government engagement in adopting FLR targets can help in the creation of decision-making structures and taskforces to coordinate efforts across institutions or agencies and develop integral strategies (Stanturf, Mansourian and Kleine, 2017).
<b>Legal</b>	Insecure land tenure is frequent in mosaic landscapes and can threaten the success of FLR and lead to conflict.	Clear governance and oversight needs government engagement and support to reduce legal insecurities and favor fair conflict resolution.
<b>Knowledge</b>	Due to the diverse nature of FLR, limited standardized guidelines and implementation protocols are available (Corbeels <i>et al.</i> , 2019; IUCN and WRI, 2014; Stanturf <i>et al.</i> , 2019), challenging the ability to increase capacity and thus implementation.	International, national, and subnational institutions can facilitate the dissemination of FLR guidance materials, training resources, and standard operating procedures. ‘Trainer Of Trainers’ approaches can improve in-country capacity building and knowledge management, further engaging local communities.

## Photos of the practice



Photo 7. Forest natural regeneration. California, United States of America

This conifer forest in Southern California was burned by wildfires. The forest, under protection, and was allowed to regenerate naturally without assisted planting. Burned trees, standing or downed, were left on the landscape to maintain the ecology of the site while young trees mature.



Photo 8. Mosaic landscape. Guilin, China

This heterogeneous landscape in rural China presents a mosaic of land uses combining biophysical and socioeconomic components, integrating landscape productivity to sustain local livelihoods through rice paddies, crops, and farms, with tree coverage along crops or as forested areas.



Photo 9. Mangrove restoration. Quang Ninh, Vietnam

Shrimp and aquaculture farming along coastlines is a major driver of mangrove deforestation around the world. The picture shows mangroves thriving after restoring the biophysical setting of the abandoned aquaculture pond area.



Photo 10. Multistrata agroforestry, shaded coffee. Quindío, Colombia

Crops like coffee or cocoa, among others, maintain productivity and quality when grown under the shade of trees. Coffee shrubs in this plantation increased their productivity thanks to the ability of trees to maintain optimal coffee growth temperatures by shading the crop.

## References

- Amézquita, M.C., Ibrahim, M., Llanderal, T., Buurman, P. & Amézquita, E. 2004. Carbon Sequestration in Pastures, Silvo-Pastoral Systems and Forests in Four Regions of the Latin American Tropics. *Journal of Sustainable Forestry*, 21(1): 31–49. [https://doi.org/10.1300/J091v21n01\\_02](https://doi.org/10.1300/J091v21n01_02)
- Bastin, J.-F., Finegold, Y., Garcia, C., Mollicone, D., Rezende, M., Routh, D., Zohner, C.M. & Crowther, T.W. 2019. The global tree restoration potential. *Science*, 365(6448): 76–79. <https://doi.org/10.1126/science.aax0848>
- Blanchart, E., Albrecht, A., Brown, G., Decaens, T., Duboisset, A., Lavelle, P., Mariani, L. & Roose, E. 2004. Effects of tropical endogeic earthworms on soil erosion. *Agriculture, Ecosystems and Environment*, 104: 303–315.
- Chavez-Tafur, J. & Zagt, R.J. 2014. Towards productive landscapes. European Tropical Forest Research Network and Tropenbos International, Wageningen, the Netherlands.
- Corbeels, M., Cardinael, R., Naudin, K., Guibert, H. & Torquebiau, E. 2019. The 4 per 1000 goal and soil carbon storage under agroforestry and conservation agriculture systems in sub-Saharan Africa. *Soil and Tillage Research*, 188: 16–26. <https://doi.org/10.1016/j.still.2018.02.015>
- Daoxiong, C., Wenfu, G., Zhilong, L. & Dongjing, S. 2015. Transforming China's forests. *Unasylva*, 245(55): 74–81.
- Gong, X., Liu, Y., Li, Q., Wei, X., Guo, X., Niu, D., Zhang, W., Zhang, J. & Zhang, L. 2013. Sub-tropic degraded red soil restoration: Is soil organic carbon build-up limited by nutrients supply. *Forest Ecology and Management*, 300: 77–87. <https://doi.org/10.1016/j.foreco.2012.12.002>
- Gourevitch, J.D., Hawthorne, P.L., Keeler, B.L., Beatty, C.R., Greve, M. & Verdone, M.A. 2016. Optimizing investments in national-scale forest landscape restoration in Uganda to maximize multiple benefits. *Environmental Research Letters*, 11(11): 114027. <https://doi.org/10.1088/1748-9326/11/11/114027>
- Harper, R.J., Okom, A.E.A., Stilwell, A.T., Tibbett, M., Dean, C., George, S.J., Sochacki, S.J., Mitchell, C.D., Mann, S.S. & Dods, K. 2012. Reforesting degraded agricultural landscapes with Eucalypts: Effects on carbon storage and soil fertility after 26years. *Agriculture, Ecosystems & Environment*, 163: 3–13. <https://doi.org/10.1016/j.agee.2012.03.013>
- Hawkins, S., To, P.X., Phuong, P.X., Thuy, P.T., Tu, N.D., Cuong, C.V., Brown, S., Dart, P. & Robertson, S. 2010. Roots in the Water: Legal Frameworks for Mangrove PES in Vietnam. Katoomba Group's Legal Initiative Country Study Series. Forest Trends: Washington, DC.
- Henry, K.M. & Twilley, R.R. 2013. Soil Development in a Coastal Louisiana Wetland during a Climate-Induced Vegetation Shift from Salt Marsh to Mangrove. *Journal of Coastal Research*, 29(6): 1273. <https://doi.org/10.2112/JCOASTRES-D-12-00184.1>
- Hobbs, R.J. 2002. The ecological context: a landscape perspective. *Handbook of Ecological Restoration*, pp. 24–45. Principles of Restoration. Cambridge, UK., Cambridge University Press.

- IUCN & WRI. 2014. *A guide to the Restoration Opportunities Assessment Methodology (ROAM): Assessing forest landscape restoration opportunities at the national or sub-national level*. 125 pp.
- Lal, R. 2004. Soil carbon sequestration to mitigate climate change. *Geoderma*, 123(1–2): 1–22. <https://doi.org/10.1016/j.geoderma.2004.01.032>
- Lamb, D., Stanturf, J. & Madsen, P. 2012. What Is Forest Landscape Restoration? In J. Stanturf, D. Lamb & P. Madsen, eds. *Forest Landscape Restoration*, pp. 3–23. World Forests. Dordrecht, Springer Netherlands. (also available at [http://link.springer.com/10.1007/978-94-007-5326-6\\_1](http://link.springer.com/10.1007/978-94-007-5326-6_1)).
- Maginnis, S., Rietbergen-McCracken, J. & Sarre, A. 2007. *The Forest Landscape Restoration Handbook*. New York, NY, Earthscan, Taylor & Francis. 192 pp. (also available at [https://books.google.com/books?id=jq8eBAAAQBAJ&printsec=frontcover&source=gbs\\_ge\\_summary\\_r&cad=0#v=onepage&q&f=false](https://books.google.com/books?id=jq8eBAAAQBAJ&printsec=frontcover&source=gbs_ge_summary_r&cad=0#v=onepage&q&f=false)).
- Mansourian, S., Vallauri, D. & Dudley, N. 2005. *Forest Restoration in Landscapes: Beyond Planting Trees*. New York, NY, Springer New York. (also available at <http://link.springer.com/10.1007/0-387-29112-1>).
- Minnemeyer, S., Laestadius, L., Sizer, N., Saint-Laurent, C. & Potapov, P. 2011. A world of opportunity. World Resource Institute, Washington, DC.
- Nair, P.K.R., Nair, V.D., Kumar, B.M. & Haile, S.G. 2009. Soil carbon sequestration in tropical agroforestry systems: a feasibility appraisal. *Environmental Science & Policy*, 12(8): 1099–1111. <https://doi.org/10.1016/j.envsci.2009.01.010>
- Nave, L.E., Swanston, C.W., Mishra, U. & Nadelhoffer, K.J. 2013. Afforestation Effects on Soil Carbon Storage in the United States: A Synthesis. *Soil Science Society of America Journal*, 77(3): 1035–1047. <https://doi.org/10.2136/sssaj2012.0236>
- van Noordwijk, M., Cerri, C., Woomer, P.L., Nugroho, K. & Bernoux, M. 1997. Soil carbon dynamics in the humid tropical forest zone. *Geoderma*, 79(1–4): 187–225. [https://doi.org/10.1016/S0016-7061\(97\)00042-6](https://doi.org/10.1016/S0016-7061(97)00042-6)
- Osland, M.J., Spivak, A.C., Nestlerode, J.A., Lessmann, J.M., Almario, A.E., Heitmuller, P.T., Russell, M.J., Krauss, K.W., Alvarez, F., Dantin, D.D., Harvey, J.E., From, A.S., Cormier, N. & Stagg, C.L. 2012. Ecosystem Development After Mangrove Wetland Creation: Plant–Soil Change Across a 20-Year Chronosequence. *Ecosystems*, 15(5): 848–866. <https://doi.org/10.1007/s10021-012-9551-1>
- Peichl, M., Thevathasan, N.V., Gordon, A.M., Huss, J. & Abohassan, R.A. 2006. Carbon Sequestration Potentials in Temperate Tree-Based Intercropping Systems, Southern Ontario, Canada. *Agroforestry Systems*, 66(3): 243–257. <https://doi.org/10.1007/s10457-005-0361-8>
- Sauer, T.J., Cambardella, C.A. & Brandle, J.R. 2007. Soil carbon and tree litter dynamics in a red cedar–scotch pine shelterbelt. *Agroforestry Systems*, 71(3): 163–174. <https://doi.org/10.1007/s10457-007-9072-7>
- Schedlbauer, J.L. & Kavanagh, K.L. 2008. Soil carbon dynamics in a chronosequence of secondary forests in northeastern Costa Rica. *Forest Ecology and Management*, 255(3–4): 1326–1335. <https://doi.org/10.1016/j.foreco.2007.10.039>

- Schultz, C.A., Jedd, T. & Beam, R.D.** 2012. The Collaborative Forest Landscape Restoration Program: A History and Overview of the First Projects. *Journal of Forestry*, 110(7): 381–391. <https://doi.org/10.5849/jof.11-082>
- Schulz, J.J. & Schröder, B.** 2017. Identifying suitable multifunctional restoration areas for Forest Landscape Restoration in Central Chile. *Ecosphere*, 8(1): e01644. <https://doi.org/10.1002/ecs2.1644>
- Smith, P.** 2008. Land use change and soil organic carbon dynamics. *Nutrient Cycling in Agroecosystems*, 81(2): 169–178. <https://doi.org/10.1007/s10705-007-9138-y>
- Stanturf, J., Lamb, D. & Madsen, P., eds.** 2012. *Forest Landscape Restoration: Integrating Natural and Social Sciences*. World Forests. Dordrecht, Springer Netherlands. (also available at <http://link.springer.com/10.1007/978-94-007-5326-6>).
- Stanturf, J., Mansourian, S. & Kleine, M.** 2017. IMPLEMENTING FOREST LANDSCAPE RESTORATION. *International Union of Forest Research Organizations, Special Programme for Development of Capacities (IUFRO-SPDC)*. Vienna, Austria. 128 p.: 132.
- Stanturf, J.A., Kant, P., Barnekow Lillesø, J.-P., Mansourian, S., Kleine, M., Graudal, L. & Madsen, P.** 2015. Forest Landscape Restoration as a Key Component of Climate Change Mitigation and Adaptation. International Union of Forest Research Organizations (IUFRO).
- Stanturf, J.A., Kleine, M., Mansourian, S., Parrotta, J., Madsen, P., Kant, P., Burns, J. & Bolte, A.** 2019. Implementing forest landscape restoration under the Bonn Challenge: a systematic approach. *Annals of Forest Science*, 76(2): 50. <https://doi.org/10.1007/s13595-019-0833-z>
- Stone, R.P. & Hilborn, D.** 2012. OMAFRA Factsheet, Universal Soil Loss Equation (USLE). Ontario Ministry of Agriculture, Food, and Rural Affairs. <http://www.omafra.gov.on.ca/english/engineer/facts/12-051.htm#:~:text=The%20Universal%20Soil%20Loss%20Equation,crop%20system%20and%20management%20practices.&text=Five%20major%20factors%20are%20used,loss%20for%20a%20given%20site>
- Vesterdal, L., Ritter, E. & Gundersen, P.** 2002. Change in soil organic carbon following afforestation of former arable land. *Forest Ecology and Management*, 169(1–2): 137–147. [https://doi.org/10.1016/S0378-1127\(02\)00304-3](https://doi.org/10.1016/S0378-1127(02)00304-3)
- Wei, Y.-M., Han, R., Wang, C., Yu, B., Liang, Q.-M., Yuan, X.-C., Chang, J., Zhao, Q., Liao, H., Tang, B., Yan, J., Cheng, L. & Yang, Z.** 2020. Self-preservation strategy for approaching global warming targets in the post-Paris Agreement era. *Nature Communications*, 11(1): 1624. <https://doi.org/10.1038/s41467-020-15453-z>
- Wick, A.F., Ingram, L.J. & Stahl, P.D.** 2009. Aggregate and organic matter dynamics in reclaimed soils as indicated by stable carbon isotopes. *Soil Biology & Biochemistry*, 41: 201–209. <https://doi.org/10.1016/j.soilbio.2008.09.012>
- Zhang, H., Deng, Q., Hui, D., Wu, J., Xiong, X., Zhao, J., Zhao, M., Chu, G., Zhou, G. & Zhang, D.** 2019. Recovery in soil carbon stock but reduction in carbon stabilization after 56-year forest restoration in degraded tropical lands. *Forest Ecology and Management*, 441: 1–8. <https://doi.org/10.1016/j.foreco.2019.03.037>

**Zhang, Y. & Shangguan, Z.** 2016. The coupling interaction of soil water and organic carbon storage in the long vegetation restoration on the Loess Plateau. *Ecological Engineering*, 91: 574–581.

<https://doi.org/10.1016/j.ecoleng.2016.03.033>

**Zhou, P., Luukkanen, O., Tokola, T. & Nieminen, J.** 2008. Vegetation Dynamics and Forest Landscape Restoration in the Upper Min River Watershed, Sichuan, China. *Restoration Ecology*, 16(2): 348–358.

<https://doi.org/10.1111/j.1526-100X.2007.00307.x>



# Wetlands



# 9. Avoiding conversion and conservation of wetlands

Valerie Hagger, Catherine E. Lovelock

The University of Queensland, St Lucia, Queensland, Australia

## 1. Description of the practice

Avoiding conversion, which includes avoiding drainage of wetlands for use in agriculture, aquaculture and other land-uses (

Photo 11), and conservation of wetlands in their natural state (Photo 12 and Photo 13) maintains ecosystem function of wetlands. Ecosystem functions, including carbon sequestration, nutrient cycling, water purification, flood mitigation, habitat for fisheries and biodiversity, and coastal protection provide essential ecosystem services that support productive landscapes and human well-being (Barbier *et al.*, 2011; Zedler, 2003). Intact wetlands also trap sediments and build soils, thereby maintaining coastal elevation (Shepard *et al.*, 2011). Wetland types included in this practice include all marine and coastal wetlands, and inland wetlands, classified as Lacustrine, Riverine, Palustrine, Marine and Estuarine (Ramsar Convention on Wetlands, 2018), excluding peatlands (see Chapter 4.2.1 Peatland) and human-made wetlands. Hydrological modifications of wetlands (e.g. drainage or excavations for aquaculture or salt ponds), land clearing, fertilizing, grazing, and any other factors that cause degradation of wetland vegetation and soils result in decreases in ecosystem functions. Particularly important is managing hydrology which if altered can result in changes in plant and fauna communities. Additionally, activities like drainage and construction of pond walls to prevent water flows can lead to increases in aeration of soils which results in CO<sub>2</sub> emissions as organic matter in soils decomposes (IPCC, 2014). Soil structure can also be damaged resulting in erosion and subsidence, and thereby increased vulnerability to sea level rise, severe storms, and other impacts of climate change. Loss of soil organic carbon (SOC) as a consequence of wetland damage varies regionally with climate, and locally with soil salinity and texture, and the organic matter available (Lovelock *et al.*, 2017). Identifying the location, extent and ecological character of wetlands, and the ecosystem services they provide to people, such as their SOC sequestration and storage, are important for valuing the socio-economic benefits from avoiding conversion and conservation of wetlands and to support decision-making in conservation and sustainable management actions.

## 2. Range of applicability

The most recent estimate of global natural wetland area is  $15 \times 10^6 \text{ km}^2$  –  $16 \times 10^6 \text{ km}^2$ , of which about 91 percent is inland and only 9 percent is coastal and marine (Davidson and Finlayson, 2019). Decline in global wetlands is occurring across almost all classes of inland and marine or coastal natural wetlands (Davidson and Finlayson, 2018). The Satellite-based Wetland Observation Service ([www.swos-service.eu](http://www.swos-service.eu)) has been developed to assist countries with mapping and monitoring changes in their wetlands (Weise *et al.*, 2020).

Despite their small global area compared to terrestrial forests and grasslands (Griscom *et al.*, 2017), wetlands have high importance in the global carbon cycle and greenhouse gas (GHG) emissions (Duarte *et al.*, 2013; Moomaw *et al.*, 2018) and in nutrient cycling (Jickells *et al.*, 2016; Reddy *et al.*, 1999, also see Hotspot “Wetlands”). Conservation of wetlands is an important management practice because of their rapid global degradation (Davidson, 2014; Gedan *et al.*, 2009; Millennium Ecosystem Assessment, 2005; Valiela *et al.*, 2001; Waycott *et al.*, 2009), which has led to significant CO<sub>2</sub> emissions (Pendleton *et al.*, 2012) as well as losses in other ecosystem services. An increasing number of countries are motivated to develop sustainable management actions for wetlands in order to conserve and restore wetlands (Finlayson, 2012). For example, by listing wetlands of international importance under the Ramsar Convention on Wetlands and managing for sustainable use. Although the ecological-character of Ramsar listed wetlands have been found to be significantly better than those of wetlands generally, overall there has been a widespread deterioration of wetlands (Davidson *et al.*, 2020), and there are calls for further action to reverse ongoing wetland loss and degradation (Finlayson *et al.*, 2019). Monitoring of change in wetland extent and condition and the drivers of change (e.g. Goldberg *et al.*, (2020) for mangroves) can assist in the design of management strategies to avoid losses of wetlands and thereby support their conservation. Priorities for conservation are areas where losses have been substantial, such as saltmarsh conversion in Europe (Gedan and Bertness 2009), mangrove conversion in south-east Asia (Thomas *et al.*, 2017), or where biodiversity is particularly vulnerable, for example wetland sites of international significance for migratory birds (Waliczky *et al.*, 2019). Wetland conservation is also important where the provision of ecosystem services to communities is highly valued, for example flood protection from tidal marshes and mangroves in cyclone risk areas (Hochard *et al.*, 2019; Narayan *et al.*, 2017), water purification within watersheds (Zedler, 2003), or where carbon stocks are particularly high and therefore where degradation leads to high levels of CO<sub>2</sub> emissions (e.g. Sasmito *et al.*, (2020) and Serrano *et al.*, (2019) for mangroves, tidal marshes, and seagrasses).

The management of suitable hydrological regimes and nutrient levels are necessary to maintain wetland plant communities and avoid SOC losses (Alhassan *et al.*, 2018; Millennium Ecosystem Assessment, 2005) and high levels of methane and nitrous oxide emissions (Beaulieu *et al.*, 2019; Dalal and Allen, 2008; Ma *et al.*, 2018). Conservation of wetlands can encompass sustainable use of wetlands to support livelihoods and generate income such as bio-charcoal production, timber and non-timber product creation, sustainable fisheries and aquaculture, and tourism (Gosling *et al.*, 2017; Thompson and Friess, 2019).

### 3. Impact on soil organic carbon stocks

Conservation of wetlands maintains SOC, avoiding CO<sub>2</sub> emissions, and secures carbon sequestration due to long-term SOC accumulation (Bridgham *et al.*, 2006; McLeod *et al.*, 2011; Nahlik and Fennessy 2016) and in the case of saline tidal wetlands, is associated with low levels of methane and nitrous oxide emissions (Kroeger *et al.*, 2017). SOC stocks and sequestration rates varies with wetland type (Table 32), being generally higher in tidal coastal wetlands than freshwater wetlands per unit area, and is influenced by latitude and precipitation (Atwood *et al.*, 2017; Hinson *et al.*, 2019), hydro-geomorphology (Sasmito *et al.*, 2020) and within-wetland variation in vegetation (Pearse *et al.*, 2018). A review of key data sources (Table 32) shows that overall more data on SOC stocks and accumulation are needed for tidal marshes and freshwater wetlands in tropical environments. Globally the area of near pristine wetlands – up to 1.15 million km<sup>2</sup> of coastal wetlands (seagrass, mangrove, tidal marsh), and 3.6 million km<sup>2</sup> of freshwater wetlands, sequesters 0.86 and 0.72 Gt (gigaton) CO<sub>2</sub> per year in the soil respectively (Bridgham *et al.*, 2014; McLeod *et al.*, 2011). There is high uncertainty in estimates of wetland extent in many nations because of the limited number of national inventories; although governments and non-government organizations are improving inventories (e.g. see Canada Wetland Inventory, <https://maps.ducks.ca/cwi/>). The global degradation of coastal wetlands is likely to have emitted an estimated 0.15-1.02 Gt CO<sub>2</sub> per year (Pendleton *et al.*, 2012), equivalent to 3-19 percent of global anthropogenic CO<sub>2</sub> emissions due to deforestation (van der Werf *et al.*, 2009). Reducing impacts on coastal wetlands could result in 141-466 Tg (metric ton) of avoided CO<sub>2</sub> emissions per year associated with decomposition of above and below ground biomass and SOC (Griscom *et al.*, 2017). Sediments in coastal wetlands can accumulate 1.9-3.9 mm/year (Breithaupt *et al.*, 2012), which contributes to their carbon sink capacity. Accumulation of SOC in coastal wetlands is likely to increase with sea level rise (Rogers *et al.*, 2019), although thresholds of SLR are likely to be 6.1-7.6 mm/year (Saintilan *et al.*, 2020). Wetland conservation is therefore particularly important given the high levels of carbon stocks in soils that may have accumulated over 100 to 1000s of years, which if disturbed liberate CO<sub>2</sub> to the atmosphere (Lovelock *et al.*, 2017) (see also [Mangrove Hotspots and Wetland Hotspots presented in volume 2 of this manual](#)).

**Table 32.** Changes in soil organic carbon stocks reported for coastal and freshwater conserved wetlands across the world

Wetland & Location*	Climate zone	Baseline SOC stock (tC/ha)	Additional SOC storage (tC/ha/yr)	Duration	Depth (cm)	More information	Reference
Coastal and inland wetlands							
Tidal marsh, mangrove and freshwater, Global	Temperate, subtropical, tropical	NA	Means: All wetlands 1.85; Tidal marsh 2.48; Mangrove 2.30; Freshwater marsh 1.97; Peatland 0.77	Literature to 2017	NA	473 soil/sediment cores from various wetlands	Cheng <i>et al.</i> (2020)
Coastal and inland wetlands, United States	Temperate	Means ( $\pm$ SE): Eastern Mountains and Upper Midwest ( $478 \pm 58$ ); Interior Plains ( $195 \pm 25$ ); Coastal Plains ( $198 \pm 21$ ); West ( $216 \pm 30$ ); Tidal saline wetlands (340); Freshwater inland wetlands (295)	NA	2011	120	967 soil pits from wetland sites across the US	Nahlik and Fennessy (2016)
Coastal wetlands							
Tidal marsh, mangrove, seagrass, Global	Temperate, subtropical, tropical	NA	Mean ( $\pm$ SE): Tidal marsh $2.18 \pm 0.24$ (range 0.18–17.13);	Literature and unpublished data 2003–2010	NA	Data from 96 tidal marsh sites, 34 mangrove sites and 123 seagrass sites	McLeod <i>et al.</i> (2011)

Wetland & Location*	Climate zone	Baseline SOC stock (tC/ha)	Additional SOC storage (tC/ha/yr)	Duration	Depth (cm)	More information	Reference
			Mangrove $2.26 \pm 0.39$ (range 0.2–9.49); Seagrass $1.38 \pm 0.38$ (range 0.45–1.90)				
Mangrove, Global		Mean ( $\pm$ SD): $283 \pm 193$	NA	Literature from 1994–2016	100	1 230 sampling locations from 48 countries. Gives variation across countries	Atwood <i>et al.</i> (2017)
		Mean ( $\pm$ SD): $361 \pm 136$ (range 86 –729)	NA	Literature and unpublished data from 1990–2016	100	Spatial modelling based on global mangrove SOC database (1 812 soil profiles from 47 countries). Gives variation across countries	Sanderman <i>et al.</i> (2018)
		Means ( $\pm$ SE): Global $333.7 (\pm 11.2)$ ; West Africa $278.4 (\pm 16.5)$ ; Asia $294.8 (\pm 20.4)$ ; Middle East $110.8 (\pm 11.6)$ ; Southeast Asia $375.6 (\pm 17.5)$ ; Oceania $447.9 (\pm 19.0)$ ; Americas $350.9 (\pm 19.9)$ ; Central America $401.9 (\pm 20.0)$ ; South America $154.9 (\pm 11.7)$	NA	2007–2017	100	Soil cores from 190 sites across five continents	Kauffman <i>et al.</i> (2020)
		NA	Mean ( $\pm$ SD): $2.31 \pm 2.09$	Literature and unpublished data from 1989–2012	NA	65 sediment cores from Brazil, Colombia, Malaysia, Indonesia, China, Japan, Vietnam, and Thailand, and	Breithaupt <i>et al.</i> (2012)

Wetland & Location*	Climate zone	Baseline SOC stock (tC/ha)	Additional SOC storage (tC/ha/yr)	Duration	Depth (cm)	More information	Reference
						additional data from Mexico and the US	
Tidal marsh and mangrove, Brazil	Tropical	Mean: Tidal marsh 257; Mangrove 341	NA	unknown	135- >300	Field surveys 9 mangrove and 3 tidal marsh sites mouth of Amazon River	Kauffman <i>et al.</i> (2018)
Tidal marsh, mangrove, seagrass, Australia	Arid, semi-arid, temperate, subtropical, and tropical	Mean ( $\pm$ SD): Tidal marsh 168 $\pm$ 127; Mangrove 251 $\pm$ 155; Seagrass 112 $\pm$ 88	Mean ( $\pm$ SD): Tidal marsh 0.39 $\pm$ 0.3; Mangrove 1.26 $\pm$ 0.9; Seagrass 0.36 $\pm$ 0.3	Literature and unpublished data	100	Database of 1 553 wetland sites (593 tidal marsh, 323 mangrove, 637 seagrass) on SOC stocks (1103 cores), and SOC sequestration rates (352 cores). Gives variation across regions	Serrano <i>et al.</i> (2019)
Tidal marsh, mangrove, seagrass, South-eastern Australia	Temperate	Means ( $\pm$ SD): Tidal marsh 57.96 $\pm$ 2.90 (range 23.33 - 291.18); Mangroves 50.64 $\pm$ 1.35 (range 23.34 - 77.81); Seagrass 23.48 $\pm$ 0.57 (range 23.33 - 73.42)	NA	2014	30	Spatial modelling based on 287 sediment cores from 96 coastal wetlands across Victoria (125 tidal marsh, 60 mangrove, and 102 seagrass)	Lewis <i>et al.</i> (2020)
Tidal marsh, mangrove, tidal freshwater wetlands, United States		Mean 270	NA	Literature and unpublished data 1998-2017	100	Spatial modelling based on 1 959 soil cores from 49 studies across US	Holmquist <i>et al.</i> (2018)
Freshwater wetlands							

Wetland & Location*	Climate zone	Baseline SOC stock (tC/ha)	Additional SOC storage (tC/ha/yr)	Duration	Depth (cm)	More information	Reference
Freshwater wetlands, Global	Temperate and tropics	NA	Tropical: mean 1.29 (range 0.42-3.06); Temperate: 1.43.	2004–2009	50–300	9 soil cores each in 5 wetland sites in northern Ohio, eastern and western Costa Rica, and Botswana	Mitsch <i>et al.</i> (2013)
Freshwater wetlands, South-eastern Australia	Temperate	Alpine wetlands 290 ± 180; Shallow freshwater marsh 200 ± 200; Saline wetlands 64 ± 48; Freshwater meadow 130 ± 100; Deep freshwater marsh 230 ± 190; Permanent open freshwater 110 ± 120; All wetlands 186 ± 176	Permanent open freshwater sites 2.3 ± 0.7; Shallow freshwater marshes 0.91 ± 0.27; deep freshwater marsh 1.6 ± 0.5; All wetlands 1.9 ± 0.4	2015–2016	50–100	>1 600 samples across 103 temperate, alpine, and semi-arid wetland sites	Carnell <i>et al.</i> (2018)
Saline and freshwater marshes, North-east China		NA	Freshwater marsh: mean 2.03; Saline marsh: mean 0.62; Overall mean 1.65 ± 0.67.	2011–2012	40	12 wetland sites in Heilongjiang and Jilin Provinces	Zhang <i>et al.</i> (2016)
Freshwater wetlands, North America		NA	Mean 0.17 (range 0–6.16)	From 1996	NA	Literature and published sources	Bridgham <i>et al.</i> (2006) Bridgham <i>et al.</i> (2014)

\*Soil types are rarely published in studies and therefore have not been included in the table.

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Avoiding conversion and conservation of wetlands contributes to sediment accretion, nutrient retention, biodiversity of soil fauna (including bioturbators), maintenance of soil structure suitable for plant recruitment and high rates of water infiltration, and avoids soil compaction (Spivak *et al.*, 2019). Conserved riparian forests in Brazil (i.e. those with low degradation) have been shown to have higher soil carbon contributing to nutrient supply and soil structure, and maintaining water quality and stream habitat in riverine wetlands (Celentano *et al.*, 2017).

### 4.2. Minimization of threats to soil functions

Table 33. Soil threats

Soil threats	
<b>Soil erosion</b>	Wetland vegetation cover is crucial for avoiding soil erosion and compaction. For example, conversion of saltmarshes to agriculture has resulted in soil loss and up to 9m subsidence in the Sacramento-San Joaquin River Delta in California (Ingebritsen <i>et al.</i> , 2000).
<b>Nutrient imbalance and cycles</b>	Intact wetlands retain nutrients and help purify surface waters (Alongi and McKinnon 2005; Saderne <i>et al.</i> , 2020). For example, conversion of wetlands in the River Thames has contributed to nutrient pollution in the estuary (Jickells <i>et al.</i> , 2016). Similarly, water quality declined in the Upper Midwestern region of the US, when about 60 percent of wetlands were drained, mostly for agriculture (Zedler 2003).
<b>Soil salinization and alkalinization</b>	Coastal wetlands can protect adjacent lands from tidal inundation through attenuation of tidal flows and storm surge and waves (Gedan <i>et al.</i> , 2011; Silliman <i>et al.</i> , 2019; Temmerman <i>et al.</i> , 2013).
<b>Soil contamination/pollution</b>	Intact wetlands retain metal and organic pollutants (Barbier <i>et al.</i> , 2011; Rabaoui <i>et al.</i> , 2020). In many cases wetlands are constructed to assist with treatment of pollutants (Vymazal and Bfezinova 2015).
<b>Soil acidification</b>	Wetlands tend to be acid due to the biogeochemical processes associated with organic matter decomposition (Spivak <i>et al.</i> , 2019). However, coastal wetlands can export alkalinity which contributes to the net CO <sub>2</sub> sink capacity of wetlands (Maher <i>et al.</i> , 2018). Drainage of wetlands can also cause wetlands to become highly acidic, because of acid sulphate soils being exposed to air, releasing sulphuric acid (Cook <i>et al.</i> , 2000).



Soil threats	
<b>Soil biodiversity loss</b>	Conservation of wetlands maintains biodiversity of soil fauna and microbes which contribute to the unique biogeochemistry of wetlands (Spivak <i>et al.</i> , 2019) and strongly influence soil organic matter decomposition (Jackson <i>et al.</i> , 2017). Benthic fauna in coastal wetlands are important in food webs that support coastal fisheries (Abrantes <i>et al.</i> , 2015).
<b>Soil compaction</b>	Conservation of wetlands avoids compaction, which reduces ecosystem functions of water infiltration as well as recruitment of seedlings and root growth (Ola <i>et al.</i> , 2020).
<b>Soil water management</b>	Intact wetlands have unique hydrological regimes to maintain soil conditions and native plant communities (Spivak <i>et al.</i> , 2019), which prevents dieback (Duke <i>et al.</i> , 2017) and loss of SOC (Alhassan <i>et al.</i> , 2018).

### 4.3. Increases in production (e.g. food/fuel/feed/timber)

Many wetlands are important for subsistence activities of local communities which can be compatible with conservation if managed appropriately. These activities include extraction of timber (e.g. fuel wood, charcoal, building materials) and non-timber forest products (e.g. honey, waxes, plant fibres), fishing and collection of crabs, and ecotourism (Gosling *et al.*, 2017; Uddin *et al.*, 2013). Coastal wetlands are also important nursery grounds and food sources for fisheries (Whitfield, 2017), and support commercial fisheries (Barbier *et al.*, 2011). Wetland conservation can support food security and incomes of communities contributing to multiple sustainable development goals (Friess *et al.*, 2019).

### 4.4. Mitigation of and adaptation to climate change

Wetland conservation contributes directly to climate change mitigation and adaptation because wetlands are significant carbon sinks, and provide coastal protection from flooding and erosion, buffering the impacts of sea level rise, increased storm surges, and wave activity associated with climate change (Duarte *et al.*, 2013; Menéndez *et al.*, 2020). Avoiding conversion and drainage of wetlands prevents CO<sub>2</sub> emissions, and also prevents increases in methane and nitrous oxide emissions from conversion to rice, aquaculture, other agricultural land uses or freshwater wetlands (IPCC, 2014, 2019; Moomaw *et al.*, 2018), which can contribute to global warming.

## 4.5. Socio-economic benefits

Avoiding conversion of coastal and inland wetlands secures provision of their ecosystem services which have initially been valued globally at US \$13,165 billion per year, around 60 percent of which is from coastal wetlands (Costanza *et al.*, 1997). Intact wetlands provide a number of key ecosystem services to communities that include coastal protection from storms (valued at US \$8,240 per ha annually in the US (Costanza *et al.*, 2008)), fisheries (valued at AU \$25,741 and AU \$5,297 per ha annually for saltmarsh and mangrove respectively in Australian estuaries (Taylor *et al.*, 2018)), flood mitigation (valued at \$65 billion per year for mangroves (Menéndez *et al.*, 2020)), erosion control, water purification (pollutant control), forest products, and carbon sequestration (Barbier *et al.*, 2011; Zedler 2003). In Thailand, mangroves have an aggregate value of \$19,000 per ha which was estimated for provision of coastal protection, wood products and habitat-fishery linkages (Barbier *et al.*, 2008). They also support tourism valued at US \$42,000 per year for the Sundarbans mangrove reserve in Bangladesh (Uddin *et al.*, 2013), but much higher (US \$104 million per year) for Can Gio Mangrove Biosphere Reserve in Vietnam (Kuenzer and Tuan 2013), or when considering the number of visitors attracted to mangroves worldwide (Spalding and Parrett 2019). The conversion of coastal wetlands globally, and associated CO<sub>2</sub> emissions, has been estimated to result in economic damages attributable to climate change of US \$6–42 billion annually (Pendleton *et al.*, 2012).

## 5. Potential drawbacks to the practice

### 5.1. Increases in greenhouse gas emissions

Wetlands are carbon sinks, but can emit methane (IPPC, 2014), particularly when polluted and at higher temperatures (Al-Haj and Fulweiler, 2020). Methane emissions decrease with increased salinity, and have been found to decrease as salinity in mangrove and tidal marsh soils increase (Al-Haj and Fulweiler 2020; Poffenbarger *et al.*, 2011; Purvaja and Ramesh, 2001). Overall methane emissions from tidally influenced mangrove and tidal marsh are low, and even lower in seagrass (Al-Haj and Fulweiler, 2020; Negandhi *et al.*, 2019). When balancing carbon sequestration and methane emissions, seagrasses have been found to remain as net GHG sinks (Al-Haj and Fulweiler, 2020). At low elevations, mangroves and tidal marsh also remain net GHG sinks (Negandhi *et al.*, 2019; Rosentreter *et al.*, 2018), however may become net GHG sources in some systems (Al-Haj and Fulweiler, 2020). Over a long time period, freshwater wetlands have also been found to become net GHG sinks (Mitsch *et al.*, 2013). Whilst drainage of wetlands or vegetation degradation generally reduces methane emissions, nitrous oxide emissions are higher, especially near human settlements and agriculture regions (Dalal and Allen, 2008; Ma *et al.*, 2018). Methane and nitrous oxide fluxes have large spatial and temporal variability in wetlands (Dalal and Allen, 2008; Kayranli *et al.*, 2010) and further measurements are required to fully account for GHG fluxes in wetlands.

## 5.2. Conflict with other practice(s)

Avoiding conversion and conservation of wetlands conflicts with aquaculture and agricultural land uses. In coastal regions there are conflicts with shrimp farming, rice production, palm oil and salt production (Goldberg *et al.*, 2020; Richards and Friess, 2016; Thomas *et al.*, 2017; Valiela *et al.*, 2001).

## 5.3. Decreases in production (e.g. food/fuel/feed/timber)

Whilst wetland conservation might be perceived to have a negative impact on seafood production and export incomes, many aquaculture ponds are disused in south-east Asia due to low water quality and disease (Primavera *et al.*, 2011). Therefore, productivity and incomes from low productivity aquaculture can be similar to those from maintaining natural ecosystems for fishing (Thompson *et al.*, 2017). In many nations, community-based forest agreements allow local communities to sustainably manage their local forests, including wetlands, to support their livelihoods from collection of fuelwood, charcoal, construction materials, and fishing (Datta *et al.*, 2010; Feurer *et al.*, 2018; Roy *et al.*, 2012). Furthermore, rice and other agriculture on coastal floodplains are at risk of salinization with subsidence and rising sea levels, and increased pollution, which is negatively impacting food security (Boretti, 2020). Wetland conservation and restoration enables the provision of ecosystem services necessary to maintain food and forest production (see Sections 4.3 and 4.5). For example, managed realignment to restore saltmarshes in Europe reduces the costs of coastal defence against rising sea levels whilst providing environmental benefits (Watts *et al.*, 2003). In addition, rehabilitation of abandoned aquaculture ponds enhances carbon storage and coastal protection in the coastal zone (Duncan *et al.*, 2016).

# 6. Recommendations before implementation of the practice

Identifying the location, extent and ecological character of wetlands, and the ecosystem services they provide to people, such as their carbon sequestration and storage, are important to quantify the socio-economic benefits from the conservation and sustainable management of wetlands and support decision-making. Quantifying the potential payments for ecosystem services from wetlands are also necessary to enable wetland conservation and restoration to be considered as economically productive systems, alongside other land uses (Millennium Ecosystem Assessment, 2005). The maintenance of suitable hydrological regimes and nutrient and sediment levels should be prioritized to maintain native plant communities and avoid SOC losses and high levels of methane and nitrous oxide emissions (Al-Haj and Fulweiler, 2020; Alhassan *et al.*, 2018; Ma *et al.*, 2018; Millennium Ecosystem Assessment, 2005). An ecosystem-based approach to wetland conservation should be applied that encompasses connectivity among multiple wetlands (i.e. basin-scale management, or integrated coastal zone management) and considers the trade-offs between wetland ecosystem services are more likely to maximize the provision of environmental and socio-economic benefits (Thorslund *et al.*, 2017). In addition, climate change impacts on hydrology, wetland extent and carbon cycling should be considered to adapt conservation measures to allow long-term wetland SOC storage (Moomaw *et al.*, 2018).

## 7. Potential barriers to adoption

Table 34. Potential barriers to adoption

Barrier	YES/NO	
<b>Biophysical</b>	<b>Yes</b>	Changes in rainfall and temperatures and rising sea levels due to climate change, reductions in water flows and sediment transport due to damming upstream, freshwater withdrawal for irrigation, excessive nutrient loads, and introduction of invasive species (Millennium Ecosystem Assessment 2005).
<b>Cultural</b>	<b>Yes</b>	Conflicts with traditional and/or cultural land-use practices that result in overharvesting and overexploitation.
<b>Social</b>	<b>Yes</b>	Social perceptions of the threats to and value of wetland ecosystem services (Boulton <i>et al.</i> , 2016).
<b>Economic</b>	<b>Yes</b>	Conflicts with productive aquaculture and agricultural practices, and national economic development plans for commodities.
<b>Institutional</b>	<b>Yes</b>	Lack of funds, resources and governance to regulate unsustainable practices and promote conservation or sustainable management.
<b>Legal (Right to soil)</b>	<b>Yes</b>	Contested land tenure, and lack of recognition of Indigenous land tenure rights (Fa <i>et al.</i> , 2020; Lovelock and Brown 2019).
<b>Knowledge</b>	<b>Yes</b>	Lack of knowledge sharing between scientists and decision-making by land owners and managers, and integrating the knowledge of Indigenous communities in wetland conservation and management. Lack of information on the trade-offs between different wetland ecosystem services (Boulton <i>et al.</i> , 2016).
<b>Other</b>	<b>Yes</b>	Clearing and drainage of inland wetlands for agricultural expansion and irrigation, and conversion of coastal wetlands for urban expansion, infrastructure development, agriculture and aquaculture (Goldberg <i>et al.</i> , 2020; Millennium Ecosystem Assessment 2005). Coastal squeeze also prevents coastal wetland expansion and conservation under climate change (Luo, 2018).

## Photos of the practice



**Photo 11.** Converted wetlands – drained wetlands for grazing pasture in Queensland, Australia (left) and degraded mangrove during conversion to aquaculture in Myanmar (right)



Photo 12. Conservation of tidal wetlands – mangroves (*Ceriops* sp., left) and tidal marsh (*Sarcocornia* sp., right) in Queensland, Australia

SOC stocks and sequestration rates are generally higher in tidal wetlands than freshwater wetlands (Photo 13) per unit area



Photo 13. Conservation of freshwater wetland (*Juncus* sp. and *Melaleuca quinquenervia*) in Queensland, Australia

Table 35. Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n <sup>o</sup>
<i>Management of Common Reed (Phragmites australis) in Mediterranean wetlands, Spain</i>	Europe	Unknown	6	18
<i>Preserving Soil Organic Carbon in Prairie Wetlands of Central North America</i>	North America	Various	6	19
<i>Maintenance of Marshlands in Urban Tidal Wetlands in New York City, United States</i>	North America	100	6	31

## References

- Abrantes, K.G., Barnett, A., Baker, R. & Sheaves, M. 2015. Habitat-specific food webs and trophic interactions supporting coastal-dependent fishery species: an Australian case study. *Reviews in Fish Biology and Fisheries*, 25(2): 337–363. <https://doi.org/10.1007/s11160-015-9385-y>
- Al-Haj, A.N. & Fulweiler, R.W. 2020. A synthesis of methane emissions from shallow vegetated coastal ecosystems. *Global Change Biology*, 26(5): 2988–3005. <https://doi.org/10.1111/gcb.15046>
- Alhassan, A.-R.M., Ma, W., Li, G., Jiang, Z., Wu, J. & Chen, G. 2018. Response of soil organic carbon to vegetation degradation along a moisture gradient in a wet meadow on the Qinghai–Tibet Plateau. *Ecology and Evolution*, 8(23): 11999–12010. <https://doi.org/10.1002/ece3.4656>
- Alongi, D.M. & McKinnon, A.D. 2005. The cycling and fate of terrestrially-derived sediments and nutrients in the coastal zone of the Great Barrier Reef shelf. *Marine Pollution Bulletin*, 51(1): 239–252. <https://doi.org/10.1016/j.marpolbul.2004.10.033>
- Atwood, T.B., Connolly, R.M., Almahasheer, H., Carnell, P.E., Duarte, C.M., Lewis, C.J.E., Irigoien, X., Kelleway, J.J., Lavery, P.S., Macreadie, P.I., Serrano, O., Sanders, C.J., Santos, I., Steven, A.D.L. & Lovelock, C.E. 2017. Global patterns in mangrove soil carbon stocks and losses. *Nature Climate Change*, 7(7): 523+. <https://doi.org/10.1038/nclimate3326>
- Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C. & Silliman, B.R. 2011. The value of estuarine and coastal ecosystem services. *Ecological Monographs*, 81(2): 169–193. <https://doi.org/10.1890/10-1510.1>
- Barbier, E.B., Koch, E.W., Silliman, B.R., Hacker, S.D., Wolanski, E., Primavera, J., Granek, E.F., Polasky, S., Aswani, S., Cramer, L.A., Stoms, D.M., Kennedy, C.J., Bael, D., Kappel, C.V., Perillo, G.M.E. & Reed, D.J. 2008. Coastal ecosystem-based management with nonlinear ecological functions and values. *Science*, 319(5861): 321–323. <https://doi.org/10.1126/science.1150349>
- Beaulieu, J.J., DelSontro, T. & Downing, J.A. 2019. Eutrophication will increase methane emissions from lakes and impoundments during the 21st century. *Nature Communications*, 10(1): 1375. <https://doi.org/10.1038/s41467-019-09100-5>
- Boretti, A. 2020. Implications on food production of the changing water cycle in the Vietnamese Mekong Delta. *Global Ecology and Conservation*, 22. <https://doi.org/10.1016/j.gecco.2020.e00989>
- Boulton, A.J., Ekebom, J. & Gislason, G.M. 2016. Integrating ecosystem services into conservation strategies for freshwater and marine habitats: a review. *Aquatic Conservation–Marine and Freshwater Ecosystems*, 26(5): 963–985. <https://doi.org/10.1002/aqc.2703>
- Breithaupt, J.L., Smoak, J.M., Smith, T.J., Sanders, C.J. & Hoare, A. 2012. Organic carbon burial rates in mangrove sediments: Strengthening the global budget. *Global Biogeochemical Cycles*, 26. <https://doi.org/10.1029/2012gb004375>



- Bridgham, S.D., Megonigal, J.P., Keller, J.K., Bliss, N.B. & Trettin, C.** 2006. The carbon balance of North American wetlands. *Wetlands*, 26(4): 889–916. [https://doi.org/10.1672/0277-5212\(2006\)26\[889:Tcbona\]2.0.Co;2](https://doi.org/10.1672/0277-5212(2006)26[889:Tcbona]2.0.Co;2)
- Bridgham, S.D., Moore, T.R., Richardson, C.J. & Roulet, N.T.** 2014. Errors in greenhouse forcing and soil carbon sequestration estimates in freshwater wetlands: a comment on Mitsch *et al.*, (2013). *Landscape Ecology*, 29(9): 1481–1485. <https://doi.org/10.1007/s10980-014-0067-2>
- Celentano, D., Rousseau, G.X., Engel, V.L., Zelarayán, M., Oliveira, E.C., Araujo, A.C.M., de Moura, E.G.** 2017. Degradation of Riparian Forest Affects Soil Properties and Ecosystem Services Provision in Eastern Amazon of Brazil. *Land Degradation & Development* 28, 482–493. <https://doi.org/10.1002/ldr.2547>
- Cheng, C.F., Li, M., Xue, Z.S., Zhang, Z.S., Lyu, X.G., Jiang, M. & Zhang, H.R.** 2020. Impacts of Climate and Nutrients on Carbon Sequestration Rate by Wetlands: A Meta-analysis. *Chinese Geographical Science*, 30(3): 483–492. <https://doi.org/10.1007/s11769-020-1122-3>
- Cook, F.J., Hicks, W., Gardner, E.A., Carlin, G.D., Froggatt, D.W.** 2000. Export of Acidity in Drainage Water from Acid Sulphate Soils. *Marine Pollution Bulletin* 41, 319–326. [https://doi.org/10.1016/S0025-326X\(00\)00138-7](https://doi.org/10.1016/S0025-326X(00)00138-7)
- Costanza, R., d'Arge, R., deGroot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., Oneill, R.V., Paruelo, J., Raskin, R.G., Sutton, P. & vandenBelt, M.** 1997. The value of the world's ecosystem services and natural capital. *Nature*, 387(6630): 253–260. <https://doi.org/10.1038/387253a0>
- Costanza, R., Perez-Maqueo, O., Martinez, M.L., Sutton, P., Anderson, S.J. & Mulder, K.** 2008. The value of coastal wetlands for hurricane protection. *Ambio*, 37(4): 241–248. [https://doi.org/10.1579/0044-7447\(2008\)37\[241:Tvowcf\]2.0.Co;2](https://doi.org/10.1579/0044-7447(2008)37[241:Tvowcf]2.0.Co;2)
- Dalal, R.C. & Allen, D.E.** 2008. Greenhouse gas fluxes from natural ecosystems. *Australian Journal of Botany*, 56(5): 369–407. <https://doi.org/10.1071/BT07128>
- Damien T. Maher, Mitchell Call, Isaac R. Santos & Christian J. Sanders.** 2018. Beyond burial: lateral exchange is a significant atmospheric carbon sink in mangrove forests. *Biology Letters*, 14(7): 20180200. <https://doi.org/doi:10.1098/rsbl.2018.0200>
- Datta, D., Guha, P. & Chattopadhyay, R.N.** 2010. Application of criteria and indicators in community based sustainable mangrove management in the Sunderbans, India. *Ocean & Coastal Management*, 53(8): 468–477. <https://doi.org/10.1016/j.ocecoaman.2010.06.007>
- Davidson, N.C.** 2014. How much wetland has the world lost? Long-term and recent trends in global wetland area. *Marine and Freshwater Research*, 65(10): 934–941. <https://doi.org/10.1071/mf14173>
- Davidson, N.C., Dinesen, L., Fennessy, S., Finlayson, C.M., Grillas, P., Grobicki, A., McInnes, R.J. & Stroud, D.A.** 2020. Trends in the ecological character of the world's wetlands. *Marine and Freshwater Research*, 71(1): 127–138. <https://doi.org/10.1071/mf18329>

- Davidson, N.C. & Finlayson, C.M.** 2018. Extent, regional distribution and changes in area of different classes of wetland. *Marine and Freshwater Research*, 69(10): 1525–1533. <https://doi.org/10.1071/MF17377>
- Davidson, N.C. & Finlayson, C.M.** 2019. Updating global coastal wetland areas presented in Davidson and Finlayson (2018). *Marine and Freshwater Research*, 70(8): 1195–1200. <https://doi.org/10.1071/mf19010>
- Duarte, C.M., Losada, I.J., Hendriks, I.E., Mazarrasa, I. & Marbà, N.** 2013. The role of coastal plant communities for climate change mitigation and adaptation. *Nature Climate Change*, 3(11): 961–968. <https://doi.org/10.1038/nclimate1970>
- Duke, N.C., Kovacs, J.M., Griffiths, A.D., Preece, L., Hill, D.J.E., van Oosterzee, P., Mackenzie, J., Morning, H.S. & Burrows, D.** 2017. Large-scale dieback of mangroves in Australia’s Gulf of Carpentaria: a severe ecosystem response, coincidental with an unusually extreme weather event. *Marine and Freshwater Research*, 68(10): 1816–1829. <https://doi.org/10.1071/MF16322>
- Duncan, C., Primavera, J.H., Petteorelli, N., Thompson, J.R., Loma, R.J.A. & Koldewey, H.J.** 2016. Rehabilitating mangrove ecosystem services: A case study on the relative benefits of abandoned pond reversion from Panay Island, Philippines. *Marine Pollution Bulletin*, 109(2): 772–782. <https://doi.org/10.1016/j.marpolbul.2016.05.049>
- Fa, J.E., Watson, J.E., Leiper, I., Potapov, P., Evans, T.D., Burgess, N.D., Molnár, Z., Fernández-Llamazares, Á., Duncan, T., Wang, S., Austin, B.J., Jonas, H., Robinson, C.J., Malmer, P., Zander, K.K., Jackson, M.V., Ellis, E., Brondizio, E.S. & Garnett, S.T.** 2020. Importance of Indigenous Peoples’ lands for the conservation of Intact Forest Landscapes. *Frontiers in Ecology and the Environment*, n/a(n/a). <https://doi.org/10.1002/fee.2148>
- Feurer, M., Gritten, D. & Than, M.M.** 2018. Community Forestry for Livelihoods: Benefiting from Myanmar’s Mangroves. *Forests*, 9(3). <https://doi.org/10.3390/f9030150>
- Finlayson, C.M.** 2012. Forty years of wetland conservation and wise use. *Aquatic Conservation-Marine and Freshwater Ecosystems*, 22(2): 139–143. <https://doi.org/10.1002/aqc.2233>
- Finlayson, C.M., Davies, G.T., Moomaw, W.R., Chmura, G.L., Natali, S.M., Perry, J.E., Roulet, N. & Sutton-Grier, A.E.** 2019. The Second Warning to Humanity - Providing a Context for Wetland Management and Policy. *Wetlands*, 39(1): 1–5. <https://doi.org/10.1007/s13157-018-1064-z>
- Friess, D.A., Aung, T.T., Huxham, M., Lovelock, C., Mukherjee, N. & Sasmito, S.** 2019. SDG 14: Life below Water – Impacts on Mangroves. In C.J. Pierce Colfer, G. Winkel, G. Galloway, P. Pacheco, P. Katila & W. de Jong, eds. *Sustainable Development Goals: Their Impacts on Forests and People*, pp. 445–481. Cambridge, Cambridge University Press.
- Gedan, K.B. & Bertness, M.D.** 2009. Experimental warming causes rapid loss of plant diversity in New England salt marshes. *Ecology Letters*, 12(8): 842–848. <https://doi.org/10.1111/j.1461-0248.2009.01337.x>

- Gedan, K.B., Kirwan, M.L., Wolanski, E., Barbier, E.B. & Silliman, B.R.** 2011. The present and future role of coastal wetland vegetation in protecting shorelines: answering recent challenges to the paradigm. *Climatic Change*, 106(1): 7–29. <https://doi.org/10.1007/s10584-010-0003-7>
- Goldberg, L., Lagomasino, D., Thomas, N. & Fatoyinbo, T.** 2020. Global declines in human-driven mangrove loss. *Global Change Biology*, n/a(n/a). <https://doi.org/10.1111/gcb.15275>
- Gosling, A., Shackleton, C.M. & Gambiza, J.** 2017. Community-based natural resource use and management of Bigodi Wetland Sanctuary, Uganda, for livelihood benefits. *Wetlands Ecology and Management*, 25(6): 717–730. <https://doi.org/10.1007/s11273-017-9546-y>
- Griscom, B.W., Adams, J., Ellis, P.W., Houghton, R.A., Lomax, G., Miteva, D.A., Schlesinger, W.H., Shoch, D., Siikamaki, J.V., Smith, P., Woodbury, P., Zganjar, C., Blackman, A., Campari, J., Conant, R.T., Delgado, C., Elias, P., Gopalakrishna, T., Hamsik, M.R., Herrero, M., Kiesecker, J., Landis, E., Laestadius, L., Leavitt, S.M., Minnemeyer, S., Polasky, S., Potapov, P., Putz, F.E., Sanderman, J., Silvius, M., Wollenberg, E. & Fargione, J.** 2017. Natural climate solutions. *Proceedings of the National Academy of Sciences of the United States of America*, 114(44): 11645–11650. <https://doi.org/10.1073/pnas.1710465114>
- Hinson, A.L., Feagin, R.A. & Eriksson, M.** 2019. Environmental Controls on the Distribution of Tidal Wetland Soil Organic Carbon in the Continental United States. *Global Biogeochemical Cycles*, 33(11): 1408–1422. <https://doi.org/10.1029/2019gb006179>
- Hochard, J.P., Hamilton, S. & Barbier, E.B.** 2019. Mangroves shelter coastal economic activity from cyclones. *Proceedings of the National Academy of Sciences*, 116(25): 12232–12237. <https://doi.org/10.1073/pnas.1820067116>
- Ingebritsen, S.E., Ikehara, M.E., Galloway, D.L. & Jones, D.R.** 2000. Delta subsidence in California: The sinking heart of the state. Fact Sheet No. 005–00. Reston, VA.
- IPCC.** 2014. 2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands. Gland, Switzerland, IPCC.
- IPCC.** 2019. 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Chapter 7 Wetlands. Gland, Switzerland, IPCC.
- Jackson, R.B., Lajtha, K., Crow, S.E., Hugelius, G., Kramer, M.C. & Pineiro, G.** 2017. The Ecology of Soil Carbon: Pools, Vulnerabilities, and Biotic and Abiotic Controls. In D.J. Futuyma, ed. *Annual Review of Ecology, Evolution, and Systematics*, Vol 48, pp. 419–445. Annual Review of Ecology Evolution and Systematics
- Jickells, T.D., Andrews, J.E. & Parkes, D.J.** 2016. Direct and Indirect Effects of Estuarine Reclamation on Nutrient and Metal Fluxes in the Global Coastal Zone. *Aquatic Geochemistry*, 22(4): 337–348. <https://doi.org/10.1007/s10498-015-9278-7>
- K. Bromberg Gedan, B. R. Silliman & M. D. Bertness.** 2009. Centuries of Human-Driven Change in Salt Marsh Ecosystems. *Annual Review of Marine Science*, 1(1): 117–141. <https://doi.org/10.1146/annurev.marine.010908.163930>

**Kauffman, J.B., Adame, M.F., Arifanti, V.B., Schile-Beers, L.M., Bernardino, A.F., Bhomia, R.K., Donato, D.C., Feller, I.C., Ferreira, T.O., Garcia, M.D.J., MacKenzie, R.A., Megonigal, J.P., Murdiyarso, D., Simpson, L. & Trejo, H.H.** 2020. Total ecosystem carbon stocks of mangroves across broad global environmental and physical gradients. *Ecological Monographs*, 90(2).

<https://doi.org/10.1002/ecm.1405>

**Kauffman, J.B., Bernardino, A.F., Ferreira, T.O., Giovannoni, L.R., Gomes, L.E.d.O., Romero, D.J., Jimenez, L.C.Z., Ruiz, F.** 2018. Carbon stocks of mangroves and salt marshes of the Amazon region, Brazil. *Biology Letters* 14, 20180208. <https://doi.org/10.1098/rsbl.2018.0208>

**Kayranli, B., Scholz, M., Mustafa, A. & Hedmark, Å.** 2010. Carbon Storage and Fluxes within Freshwater Wetlands: a Critical Review. *Wetlands*, 30(1): 111–124. <https://doi.org/10.1007/s13157-009-0003-4>

**Kroeger, K.D., Crooks, S., Moseman-Valtierra, S. & Tang, J.** 2017. Restoring tides to reduce methane emissions in impounded wetlands: A new and potent Blue Carbon climate change intervention. *Scientific Reports*, 7(1): 11914. <https://doi.org/10.1038/s41598-017-12138-4>

**Kuenzer, C. & Tuan, V.Q.** 2013. Assessing the ecosystem services value of Can Gio Mangrove Biosphere Reserve: Combining earth-observation- and household-survey-based analyses. *Applied Geography*, 45: 167–184. <https://doi.org/10.1016/j.apgeog.2013.08.012>

**Lovelock, C.E., Atwood, T., Baldock, J., Duarte, C.M., Hickey, S., Lavery, P.S., Masque, P., Macreadie, P.I., Ricart, A.M., Serrano, O. & Steven, A.** 2017. Assessing the risk of carbon dioxide emissions from blue carbon ecosystems. *Frontiers in Ecology and the Environment*, 15(5): 257–265. <https://doi.org/10.1002/fee.1491>

**Lovelock, C.E. & Brown, B.M.** 2019. Land tenure considerations are key to successful mangrove restoration. *Nature Ecology & Evolution*, 3(8): 1135–1135. <https://doi.org/10.1038/s41559-019-0942-y>

**Ma, W.W., Alhassan, A.R.M., Wang, Y.S., Li, G., Wang, H. & Zhao, J.M.** 2018. Greenhouse gas emissions as influenced by wetland vegetation degradation along a moisture gradient on the eastern Qinghai-Tibet Plateau of North-West China. *Nutrient Cycling in Agroecosystems*, 112(3): 335–354. <https://doi.org/10.1007/s10705-018-9950-6>

**McLeod, E., Chmura, G.L., Bouillon, S., Salm, R., Bjork, M., Duarte, C.M., Lovelock, C.E., Schlesinger, W.H. & Silliman, B.R.** 2011. A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO<sub>2</sub>. *Frontiers in Ecology and the Environment*, 9(10): 552–560. <https://doi.org/10.1890/110004>

**Menéndez, P., Losada, I.J., Torres-Ortega, S., Narayan, S. & Beck, M.W.** 2020. The Global Flood Protection Benefits of Mangroves. *Scientific Reports*, 10(1): 4404. <https://doi.org/10.1038/s41598-020-61136-6>

**Millennium Ecosystem Assessment.** 2005. Ecosystems and human well-being: wetlands and water synthesis. Washington DC, World Resources Institute.

- Mitsch, W.J., Bernal, B., Nahlik, A.M., Mander, U., Zhang, L., Anderson, C.J., Jorgensen, S.E. & Brix, H. 2013. Wetlands, carbon, and climate change. *Landscape Ecology*, 28(4): 583–597. <https://doi.org/10.1007/s10980-012-9758-8>
- Moomaw, W.R., Chmura, G.L., Davies, G.T., Finlayson, C.M., Middleton, B.A., Natali, S.M., Perry, J.E., Roulet, N. & Sutton-Grier, A.E. 2018. Wetlands In a Changing Climate: Science, Policy and Management. *Wetlands*, 38(2): 183–205. <https://doi.org/10.1007/s13157-018-1023-8>
- Nahlik, A.M. & Fennessy, M.S. 2016. Carbon storage in US wetlands. *Nature Communications*, 7(1): 13835. <https://doi.org/10.1038/ncomms13835>
- Narayan, S., Beck, M.W., Wilson, P., Thomas, C.J., Guerrero, A., Shepard, C.C., Reguero, B.G., Franco, G., Ingram, J.C. & Trespalacios, D. 2017. The Value of Coastal Wetlands for Flood Damage Reduction in the Northeastern USA. *Scientific Reports*, 7. <https://doi.org/10.1038/s41598-017-09269-z>
- Negandhi, K., Edwards, G., Kelleway, J.J., Howard, D., Safari, D. & Saintilan, N. 2019. Blue carbon potential of coastal wetland restoration varies with inundation and rainfall. *Scientific Reports*, 9. <https://doi.org/10.1038/s41598-019-40763-8>
- Ola, A., Staples, T.L., Robinson, N. & Lovelock, C.E. 2020. y Plasticity in the Above- and Below-Ground Development of Mangrove Seedlings in Response to Variation in Soil Bulk Density. *Estuaries and Coasts*, 43(1): 111–119. <https://doi.org/10.1007/s12237-019-00660-9>
- Pearse, A.L., Barton, J.L., Lester, R.E., Zawadzki, A. & Macreadie, P.I. 2018. Soil organic carbon variability in Australian temperate freshwater wetlands. *Limnology and Oceanography*, 63: S254–S266. <https://doi.org/10.1002/lno.10735>
- Pendleton, L., Donato, D.C., Murray, B.C., Crooks, S., Jenkins, W.A., Sifleet, S., Craft, C., Fourqurean, J.W., Kauffman, J.B., Marba, N., Megonigal, P., Pidgeon, E., Herr, D., Gordon, D. & Baldera, A. 2012. Estimating Global ‘Blue Carbon’ Emissions from Conversion and Degradation of Vegetated Coastal Ecosystems. *Plos One*, 7(9). <https://doi.org/10.1371/journal.pone.0043542>
- Poffenbarger, H.J., Needelman, B.A. & Megonigal, J.P. 2011. Salinity Influence on Methane Emissions from Tidal Marshes. *Wetlands*, 31(5): 831–842. <https://doi.org/10.1007/s13157-011-0197-0>
- Primavera, J.H., Rollon, R.N. & Samson, M.S. 2011. 10.10 - The Pressing Challenges of Mangrove Rehabilitation: Pond Reversion and Coastal Protection. In E. Wolanski & D. McLusky, eds. *Treatise on Estuarine and Coastal Science*, pp. 217–244. Waltham, Academic Press.
- Purvaja, R. & Ramesh, R. 2001. Natural and Anthropogenic Methane Emission from Coastal Wetlands of South India. *Environmental Management*, 27(4): 547–557. <https://doi.org/10.1007/s002670010169>
- Rabaoui, L., Cusack, M., Saderne, V., Krishnakumar, P.K., Lin, Y.-J., Shemsi, A.M., El Zrelli, R., Arias-Ortiz, A., Masqué, P., Duarte, C.M. & Qurban, M.A. 2020. Anthropogenic-induced acceleration of elemental burial rates in blue carbon repositories of the Arabian Gulf. *Science of the Total Environment*, 719: 135177. <https://doi.org/10.1016/j.scitotenv.2019.135177>
- Ramsar Convention on Wetlands. 2018. Global Wetland Outlook: State of the World’s Wetlands and their Services to People. Gland, Switzerland, Ramsar Convention Secretariat.

- Reddy, K.R., Kadlec, R.H., Flaig, E. & Gale, P.M. 1999. Phosphorus retention in streams and wetlands: A review. *Critical Reviews in Environmental Science and Technology*, 29(1): 83–146.  
<https://doi.org/10.1080/10643389991259182>
- Richards, D.R. & Friess, D.A. 2016. Rates and drivers of mangrove deforestation in Southeast Asia, 2000–2012. *Proceedings of the National Academy of Sciences*, 113(2): 344–349.  
<https://doi.org/10.1073/pnas.1510272113>
- Rogers, K., Kelleway, J.J., Saintilan, N., Megonigal, J.P., Adams, J.B., Holmquist, J.R., Lu, M., Schile-Beers, L., Zawadzki, A., Mazumder, D. & Woodroffe, C.D. 2019. Wetland carbon storage controlled by millennial-scale variation in relative sea-level rise. *Nature*, 567(7746): 91–95.  
<https://doi.org/10.1038/s41586-019-0951-7>
- Rosentreter, J.A., Maher, D.T., Erler, D.V., Murray, R.H. & Eyre, B.D. 2018. Methane emissions partially offset “blue carbon” burial in mangroves. *Science Advances*, 4(6): eaao4985.  
<https://doi.org/10.1126/sciadv.aao4985>
- Roy, A.K.D., Alam, K. & Gow, J. 2012. A review of the role of property rights and forest policies in the management of the Sundarbans Mangrove Forest in Bangladesh. *Forest Policy and Economics*, 15: 46–53.  
<https://doi.org/10.1016/j.forpol.2011.08.009>
- Saderne, V., Cusack, M., Serrano, O., Almahasheer, H., Krishnakumar, P.K., Rabaoui, L., Qurban, M.A. & Duarte, C.M. 2020. Role of vegetated coastal ecosystems as nitrogen and phosphorous filters and sinks in the coasts of Saudi Arabia. *Environmental Research Letters*, 15(3). <https://doi.org/10.1088/1748-9326/ab76da>
- Saintilan, N., Khan, N.S., Ashe, E., Kelleway, J.J., Rogers, K., Woodroffe, C.D. & Horton, B.P. 2020. Thresholds of mangrove survival under rapid sea level rise. *Science*, 368(6495): 1118–1121.  
<https://doi.org/10.1126/science.aba2656>
- Sanderman, J., Hengl, T., Fiske, G., Solvik, K., Adame, M.F., Benson, L., Bukoski, J.J., Carnell, P., Cifuentes-Jara, M., Donato, D., Duncan, C., Eid, E.M., zu Ermgassen, P., Lewis, C.J.E., Macreadie, P.I., Glass, L., Gress, S., Jardine, S.L., Jones, T.G., Nsombo, E.N., Rahman, M.M., Sanders, C.J., Spalding, M. & Landis, E. 2018. A global map of mangrove forest soil carbon at 30 m spatial resolution. *Environmental Research Letters*, 13(5). <https://doi.org/10.1088/1748-9326/aabe1c>
- Sasmito, S.D., Sillanpaa, M., Hayes, M.A., Bachri, S., Saragi-Sasmito, M.F., Sidik, F., Hanggara, B.B., Mofu, W.Y., Rumbiak, V.I., Hendri, Taberima, S., Suhaemi, Nugroho, J.D., Pattiasina, T.F., Widagti, N., Barakalla, Rahajoe, J.S., Hartantri, H., Nikijuluw, V., Jowey, R.N., Heatubun, C.D., zu Ermgassen, P., Worthington, T.A., Howard, J., Lovelock, C.E., Friess, D.A., Hutley, L.B. & Murdiyarso, D. 2020. Mangrove blue carbon stocks and dynamics are controlled by hydrogeomorphic settings and land-use change. *Global Change Biology*, 26(5): 3028–3039. <https://doi.org/10.1111/gcb.15056>
- Serrano, O., Lovelock, C.E., B. Atwood, T., Macreadie, P.I., Canto, R., Phinn, S., Arias-Ortiz, A., Bai, L., Baldock, J., Bedulli, C., Carnell, P., Connolly, R.M., Donaldson, P., Esteban, A., Ewers Lewis, C.J., Eyre, B.D., Hayes, M.A., Horwitz, P., Hutley, L.B., Kavazos, C.R.J., Kelleway, J.J., Kendrick, G.A., Kilminster, K., Lafratta, A., Lee, S., Lavery, P.S., Maher, D.T., Marbà, N., Masque, P., Mateo, M.A., Mount, R., Ralph, P.J., Roelfsema, C., Rozaimi, M., Ruhon, R., Salinas, C., Samper-Villarreal, J.,

- Sanderman, J., J. Sanders, C., Santos, I., Sharples, C., Steven, A.D.L., Cannard, T., Trevathan-Tackett, S.M. & Duarte, C.M. 2019. Australian vegetated coastal ecosystems as global hotspots for climate change mitigation. *Nature Communications*, 10(1): 4313. <https://doi.org/10.1038/s41467-019-12176-8>
- Shepard, C.C., Crain, C.M. & Beck, M.W. 2011. The Protective Role of Coastal Marshes: A Systematic Review and Meta-analysis. *Plos One*, 6(11). <https://doi.org/10.1371/journal.pone.0027374>
- Silliman, B.R., He, Q., Angelini, C., Smith, C.S., Kirwan, M.L., Daleo, P., Renzi, J.J., Butler, J., Osborne, T.Z., Nifong, J.C. & van de Koppel, J. 2019. Field Experiments and Meta-analysis Reveal Wetland Vegetation as a Crucial Element in the Coastal Protection Paradigm. *Current Biology*, 29(11): 1800-1806.e3. <https://doi.org/10.1016/j.cub.2019.05.017>
- Spalding, M. & Parrett, C.L. 2019. Global patterns in mangrove recreation and tourism. *Marine Policy*, 110: 103540. <https://doi.org/10.1016/j.marpol.2019.103540>
- Spivak, A.C., Sanderman, J., Bowen, J.L., Canuel, E.A. & Hopkinson, C.S. 2019. Global-change controls on soil-carbon accumulation and loss in coastal vegetated ecosystems. *Nature Geoscience*, 12(9): 685–692. <https://doi.org/10.1038/s41561-019-0435-2>
- Taylor, M.D., Gaston, T.F. & Raoult, V. 2018. The economic value of fisheries harvest supported by saltmarsh and mangrove productivity in two Australian estuaries. *Ecological Indicators*, 84: 701–709. <https://doi.org/10.1016/j.ecolind.2017.08.044>
- Temmerman, S., Meire, P., Bouma, T.J., Herman, P.M.J., Ysebaert, T. & De Vriend, H.J. 2013. Ecosystem-based coastal defence in the face of global change. *Nature*, 504(7478): 79–83. <https://doi.org/10.1038/nature12859>
- Thomas, N., Lucas, R., Bunting, P., Hardy, A., Rosenqvist, A. & Simard, M. 2017. Distribution and drivers of global mangrove forest change, 1996–2010. *Plos One*, 12(6): e0179302. <https://doi.org/10.1371/journal.pone.0179302>
- Thompson, B.S. & Friess, D.A. 2019. Stakeholder preferences for payments for ecosystem services (PES) versus other environmental management approaches for mangrove forests. *Journal of Environmental Management*, 233: 636–648. <https://doi.org/10.1016/j.jenvman.2018.12.032>
- Thompson, B.S., Primavera, J.H. & Friess, D.A. 2017. Governance and implementation challenges for mangrove forest Payments for Ecosystem Services (PES): Empirical evidence from the Philippines. *Ecosystem Services*, 23: 146–155. <https://doi.org/10.1016/j.ecoser.2016.12.007>
- Thorslund, J., Jarsjo, J., Jaramillo, F., Jawitz, J.W., Manzoni, S., Basu, N.B., Chalov, S.R., Cohen, M.J., Creed, I.F., Goldenberg, R., Hysin, A., Kalantari, Z., Koussis, A.D., Lyon, S.W., Mazi, K., Mard, J., Persson, K., Pietron, J., Prieto, C., Quin, A., Van Meter, K. & Destouni, G. 2017. Wetlands as large-scale nature-based solutions: Status and challenges for research, engineering and management. *Ecological Engineering*, 108: 489–497. <https://doi.org/10.1016/j.ecoleng.2017.07.012>
- Uddin, M.S., de Ruyter van Steveninck, E., Stuip, M. & Shah, M.A.R. 2013. Economic valuation of provisioning and cultural services of a protected mangrove ecosystem: A case study on Sundarbans Reserve Forest, Bangladesh. *Ecosystem Services*, 5: 88–93. <https://doi.org/10.1016/j.ecoser.2013.07.002>

**Valiela, I., Bowen, J.L. & York, J.K.** 2001. Mangrove forests: One of the world's threatened major tropical environments. *Bioscience*, 51(10): 807–815. [https://doi.org/10.1641/0006-3568\(2001\)051\[0807:mfootw\]2.0.co;2](https://doi.org/10.1641/0006-3568(2001)051[0807:mfootw]2.0.co;2)

**Vymazal, J. & Bfezinova, T.** 2015. The use of constructed wetlands for removal of pesticides from agricultural runoff and drainage: A review. *Environment International*, 75: 11–20. <https://doi.org/10.1016/j.envint.2014.10.026>

**Waliczky, Z., Fishpool, L.D.C., Butchart, S.H.M., Thomas, D., Heath, M.F., Hazin, C., Donald, P.F., Kowalska, A., Dias, M.P. & Allinson, T.S.M.** 2019. Important Bird and Biodiversity Areas (IBAs): their impact on conservation policy, advocacy and action. *Bird Conservation International*, 29(2): 199–215. <https://doi.org/10.1017/s0959270918000175>

**Watts, C.W., Tollhurst, T.J., Black, K.S. & Whitmore, A.P.** 2003. In situ measurements of erosion shear stress and geotechnical shear strength of the intertidal sediments of the experimental managed realignment scheme at Tollesbury, Essex, UK. *Estuarine, Coastal and Shelf Science*, 58(3): 611–620. [https://doi.org/10.1016/S0272-7714\(03\)00139-2](https://doi.org/10.1016/S0272-7714(03)00139-2)

**Waycott, M., Duarte, C.M., Carruthers, T.J.B., Orth, R.J., Dennison, W.C., Olyarnik, S., Calladine, A., Fourqurean, J.W., Heck, K.L., Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Short, F.T. & Williams, S.L.** 2009. Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proceedings of the National Academy of Sciences of the United States of America*, 106(30): 12377–12381. <https://doi.org/10.1073/pnas.0905620106>

**van der Werf, G.R., Morton, D.C., DeFries, R.S., Olivier, J.G.J., Kasibhatla, P.S., Jackson, R.B., Collatz, G.J. & Randerson, J.T.** 2009. CO<sub>2</sub> emissions from forest loss. *Nature Geoscience*, 2(11): 737–738. <https://doi.org/10.1038/ngeo671>

**Weise, K., Hofer, R., Franke, J., Guelmami, A., Simonson, W., Muro, J., O'Connor, B., Strauch, A., Flink, S., Eberle, J., Mino, E., Thulin, S., Philipson, P., van Valkengoed, E., Truckenbrodt, J., Zanderg, F., Sanchez, A., Schroder, C., Thonfeld, F., Fitoka, E., Scott, E., Ling, M., Schwarz, M., Kunz, I., Thumer, G., Plasmeijer, A., Hilarides, L.** 2020. Wetland extent tools for SDG 6.6.1 reporting from the Satellite-based Wetland Observation Service (SWOS). *Remote Sensing of Environment*, 247. <https://doi.org/10.1016/j.rse.2020.111892>

**Whitfield, A.K.** 2017. The role of seagrass meadows, mangrove forests, salt marshes and reed beds as nursery areas and food sources for fishes in estuaries. *Reviews in Fish Biology and Fisheries*, 27(1): 75–110. <https://doi.org/10.1007/s11160-016-9454-x>

**Zedler, J.B.** 2003. Wetlands at your service: reducing impacts of agriculture at the watershed scale. *Frontiers in Ecology and the Environment*, 1(2): 65–72. [https://doi.org/10.1890/1540-9295\(2003\)001\[0065:Waysri\]2.0.Co;2](https://doi.org/10.1890/1540-9295(2003)001[0065:Waysri]2.0.Co;2)



# 10. Wetland restoration (water supplementation and promoting plant growth)

Shangqi Xu<sup>1</sup>, Guodong Wang<sup>1</sup>, Bolong Wen<sup>1</sup>, Xia Liu<sup>2</sup>, Xiujun Li<sup>1</sup>, Chunjie Tian<sup>1</sup>

<sup>1</sup>Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, Changchun, China

<sup>2</sup>Jilin Province Science and Technology Information Institute, Changchun, China

## 1. Description of the practice

For wetlands other than peatland, the most recommended practices in wetland restoration for C sequestration are water supplementation (such as rewetting of drained wetland) and the promotion of plant growth (Carnell *et al.*, 2018). Water supplementation can build an anaerobic environment and reduce soil C loss (Villa and Bernal, 2018). Promoting plant growth can provide more additional organic C to soil.

Water supplementation means rewetting the drained wetland or raising the water table of the degraded wetland. The most recommended water level is about ~10~30 cm above ground (Nadeau and Conway, 2015; Wang *et al.*, 2017). A water level that is too low cannot provide the anaerobic environment, while a very deep water table may restrain plant growth. Therefore, the best way is maintaining a lower water level in the growing season to promote planting growth and maintaining a deep water level in the non-growing season to sequester C (Wang *et al.*, 2017).

Promoting plant growth aims to add more organic C to soil. Wetland vegetation can be restored through natural succession via its seed bank in the soil if it is still in place (Wang *et al.*, 2015). Sowing or planting native wetland plants is also recommended (Renzi *et al.*, 2019). But there should have a diversity of plants, only having one or two species may reduce the overall biomass and biodiversity. In some wetlands (e.g. salt marsh), the soil in its natural state can limit plants growth, plants that can grow well in the barren soil should be selected for C sequestration (Doherty *et al.*, 2011). In addition, in degraded wetland with barren soil, at the beginning of wetland restoration soil improvement practices such as fertilization and tillage could be a choice to promote vegetation recovery. However, these practices should be used carefully and get adequate assessment. For

example, although fertilization can promote plant growth and increase soil C in most conditions (Xu *et al.*, 2020), this practice should be used with the proper dosages for the wetland environment in order to prevent potential pollution. Furthermore, these practices, especially tillage and other soil mechanical disturbance, should be strictly prohibited after 1-2 years of restoration (Xu *et al.*, 2020).

Apart from the mentioned practices above, conventional agricultural cultivation and other human activities that lead to wetland degradation should be completely banned (Xu *et al.*, 2019). If the degradation continues, the ecosystem services provide by wetland such as water cleaning, water storage and biological habitat would lost.

## 2. Range of applicability

In general, wetlands degraded due to use for rainfed farming can restore better than that for other land-uses. Wetland restoration benefits soil C sequestration in boreal, subarctic, and temperate degraded wetlands, but it shows less results in term of increasing C sequestration in tropical degraded wetlands (Xu *et al.*, 2019). The restoration of seasonal wetlands and freshwater wetlands increases soil C sequestration more than that of tidal wetlands and salt wetlands, in which the restoration may have no positive effect to C sequestration (Xu *et al.*, 2019). Restoration practices should be conducted as soon as possible, as if the wetlands had been degraded more than 15 years, the restoration will be much less effective (Xu *et al.*, 2019).

## 3. Impact on soil organic carbon stocks

Wetland restoration has been implemented worldwide and is considered to be an effective way to recover the soil C lost as a result of wetland cultivation (Crooks *et al.*, 2011). Wetland restoration can accelerate carbon sequestration and greatly contribute to mitigating climate change if proper measures are utilized (Table 36). Most restored wetlands have an soil C content that is higher than that of cultivated wetlands but still lower than that of natural wetlands (Xu *et al.*, 2019).

**Table 36.** Changes in soil organic carbon stocks reported for wetland restoration

Location	Climate zone	Soil type	Baseline C stock or content	Additional C storage or content	Duration (Years)	Depth (cm)	More information	Reference
Global	NA	NA	NA	9% increase (95%CI: -3%~23%)	NA	NA	Meta-analysis; Measured data; Wetland restoration	Xu <i>et al.</i> (2019)
Eastern England	NA	NA	59 ( $\pm 10$ SE) tC/ha	0.65-1.04 tC/ha/yr	0-100	0-30	Model combined with field chronosequence; Saltmarsh restoration after degradation from agriculture	Burden <i>et al.</i> (2019)
Nebraska, United States of America	NA	Mollisols; Fillmore, Scott, and Massie series	96.4 ( $\pm 1.56$ SE) tC/ha	5.1 ( $\pm 2.72$ SE) tC/ha	20-30	0-50	Measured; total carbon; Freshwater marsh restoration (sediment removal)	Daniel <i>et al.</i> (2017)
Yellow River Delta, China	Warm temperate continental monsoon climate	NA	3.54-3.86 g/kg	1.4-6.89 g/kg	7	0-20	Coastal marsh restored with freshwater supplement	Wang <i>et al.</i> (2011)
Yangtze River estuary, China	Subtropical monsoon climate	NA	26.44 ( $\pm 2.69$ SE) tC/ha	2.99 ( $\pm 0.57$ SE) tC/ha	6	0-10	Coastal marsh restored with agricultural abandon	Li <i>et al.</i> (2012)
Sangjiang Plain, China	Monsoon climate of medium latitudes	NA	33.41( $\pm 1.45$ SE) g/kg	25.65-42.12 g/kg	3-12	0-20	Freshwater marsh restoration with agricultural abandon	Song <i>et al.</i> (2012)
Prairie Pothole region, Canada	Temperate continental climate	Mineral soil	116 tC/ha	2.5-6.1 tC/ha/yr	1-12	0-30	Freshwater mineral soil wetlands	Badiou <i>et al.</i> (2011)

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Wetland restoration has positive effects on soil properties. One of the main benefits is the increase of biological diversity and activity, enzymatic activity and biogeochemical cycles, which are important for the restoration of wetland ecological functions (Meli *et al.*, 2014; Zedler and Kercher, 2005). Soil bulk density of restored wetlands is usually lower than that of degraded wetlands, and the restored soil can hold more water with higher proportion of porosity and organic matter (Suir *et al.*, 2019). The effects of wetland restoration on soil pH or CEC mainly depend on the original soil condition and type of wetland.

### 4.2 Minimization of threats to soil functions

Table 37. Soil threats

Soil threats	
<b>Soil erosion</b>	Plant growth can reduce the soil wind erosion, as well as water erosion. Raised water level can reduce soil wind erosion (Luo <i>et al.</i> , 2015).
<b>Nutrient imbalance and cycles</b>	Restored wetlands are effective at removing N, whereas P can be released for several years after restoration (Audet <i>et al.</i> , 2020; Wang <i>et al.</i> , 2015).
<b>Soil salinization and alkalinization</b>	Wetland restoration can decrease the pH of salt marsh with increasing soil C. The salinization could be decreased with increasing water as the salt can drift with the water (Wang <i>et al.</i> , 2017).
<b>Soil contamination/pollution</b>	Wetland restoration is an important path to control pollution. Plant growth can reduce the pollution, and the wetland environment can retain contamination (Su <i>et al.</i> , 2019).
<b>Soil acidification</b>	Wetland restoration can remediate drained acid sulfate soil with pH increase and sulfur reduction (Johnston <i>et al.</i> , 2014).
<b>Soil biodiversity loss</b>	The soil biodiversity loss will be halted with wetland restoration. Soil microbial diversity and fauna diversity will increase after wetland restoration (Li <i>et al.</i> , 2020; Xu <i>et al.</i> , 2017).
<b>Soil compaction</b>	Wetland restoration can reduce the soil bulk density and relieve the soil compaction caused by wetland degradation (Suir <i>et al.</i> , 2019).
<b>Soil water management</b>	Wetland restoration can increase the ability to store water and regulate runoff.

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

The plants in wetland could be food, fibre, fuel, fodder, herb or timber depending on the vegetation type and community in the restored wetland. In addition, some wetlands can provide aquatic products such as fish, crab and shrimp, as well as increase bird population (Ducks Unlimited Canada, 2020).

### 4.4 Mitigation of and adaptation to climate change

Wetland restoration can transform degraded wetland to a C sink and can reduce GHG emissions. The restoration of coastal wetland in northeast China has the potential to reduce CO<sub>2</sub> emission from 609 mg CO<sub>2</sub>/m/h to 278 mg CO<sub>2</sub>/m/h (Chen *et al.*, 2018; Olsson *et al.*, 2015). For degraded wetland without drainage, the restoration may decrease the emission of CH<sub>4</sub>, as well as N<sub>2</sub>O, with a decrease of more than 20 percent (Gleason *et al.*, 2009).

### 4.5 Socio-economic benefits

Wetlands give us natural places to play, learn and explore. They are destinations for hiking, hunting, canoeing, photography and more (Ducks Unlimited Canada, 2020). Restored wetlands can be a tourism resource particularly with more wetland parks being built around the world. Typical wetland parks based on restored wetlands include Everglades National Park in United States of America, Jiuli Lake Wetland in Xuzhou in China, Tianjin Lingang Ecological Wetland Park in China and so on.

### 4.6 Additional benefits to the practice

Wetlands can regulate microclimate, clean water and store flood water, especially in a city or suburb, which can benefit to intensive landscapes and human settlements (McLaughlin and Cohen, 2013). Wetlands protect wildlife by providing hundreds of species with safe places to eat, sleep and raise young (Ducks Unlimited Canada, 2020).

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

Table 38. Soil threats

Soil threats	
Soil erosion	In certain conditions, e.g. riparian, water level increase may increase water flow and lead to water erosion (Thi and Minh, 2019).
Nutrient imbalance and cycles	Wetland restoration may lead to P limitation with increasing restored duration (Smith <i>et al.</i> , 2011).
Soil salinization and alkalinization	For some cultivated coastal salt marsh for farming, the restoration of tidal hydrology may result in soil salinization (Li <i>et al.</i> , 2012).
Soil water management	Wetland restoration may influence the regional water management due to its water demand (Hodge and McNally, 2000).

### 5.2 Increases in greenhouse gas emissions

The restoration of drained wetland with water level increase always results in higher CH<sub>4</sub> emission as the anaerobic environment can promote CH<sub>4</sub> emission, while upland soil often was a slight CH<sub>4</sub> sink (Tangen *et al.*, 2015). For cultivated wetland without drainage in the Prairie Pothole Region, the CO<sub>2</sub> emission may increase after restoration (Gleason *et al.*, 2009). The effects of wetland restoration on GHG are context dependent and may not be significant in certain environments (Gleason *et al.*, 2009).

### 5.3 Conflict with other practice(s)

Wetland restoration is in conflict with agricultural activities such as tillage, crop planting and applications of pesticide, especially the rainfed farming. In addition, other human activities, including intensive livestock grazing, infrastructure development etc., are in conflict with restoration as they may cause degradation.

## 5.4 Decreases in production (e.g. food/fuel/feed/timber)

Wetland restoration may decrease the production of grains or other agricultural products as the agricultural activity lead to wetland degradation should be stopped.

## 6. Recommendations before implementation of the practice

First, wetlands restoration may bring economic loss to landowners as the production activities lead to wetlands degradation must stopped, and they must reestablish livelihood-generating activities that are not in conflict with restoration. Thus, conflicts between landowner and restoration performer must be fully solved before restoration.

A proper plan based on the field condition is required. The restoration should take in a full consideration of the recovery effects and costs, which depend on the original land use of the wetland, recovery area, water demand, the hydrological regulation method, revegetation method and so on.

Before wetland restoration, the applicability of the restore practices should be evaluated. A pilot project can provide much useful information for the conduction of wetland restoration before the large-scale restoration.

## 7. Potential barriers to adoption

Table 39. Potential barriers to adoption

Barrier	YES/NO	
<b>Biophysical</b>	<b>Yes</b>	It is hard to restore degraded wetlands to their natural status, because restoration is a long term process and the environment or climate in a large-scale may change (Moreno-Mateos <i>et al.</i> , 2012).
<b>Cultural</b>	<b>Yes</b>	Depending on the emphasis degree of public on natural environments (Zhu <i>et al.</i> , 2016), and raising awareness can be a useful strategy to foster public support for wetland restoration (Scholte <i>et al.</i> , 2016).
<b>Social</b>	<b>Yes</b>	Urban expansion, agriculture, aquaculture and industrial developments are main reasons of wetland losses and they may prevent wetland restoration (Zedler and Kercher, 2005), especially in developing countries in which the conflicts between development and ecological conservation are more prominent (Marambanyika and Beckedahl, 2016).
<b>Economic</b>	<b>Yes</b>	Wetland restoration may lead to landowners suffer economic losses due to the original producing activities on the degraded wetlands had to be stopped. The loss of land or income decrease landowners' willingness to participate in wetland restoration (Zhu <i>et al.</i> , 2016).
<b>Institutional</b>	<b>Yes</b>	Land ownership, e.g. private land ownership, in some region may be barriers for wetland restoration as wetland restoration needs the support of landowners (Marambanyika and Beckedahl, 2016; Zhu <i>et al.</i> , 2016).
<b>Legal (Right to soil)</b>	<b>Yes</b>	Incomplete environmental protection act and land act, as well as poor implementation of legislation, may affect the restoration, because the restoration project may be hard to conduct when legal support lacks (Marambanyika and Beckedahl, 2016).
<b>Knowledge</b>	<b>Yes</b>	Wetland restoration requires professional knowledge to assess the ecosystem conditions, the preferred practices, the potential effects and risk (Zhu <i>et al.</i> , 2016).
<b>Other</b>	<b>Yes</b>	Some wetlands may be difficult to restore, e.g., the seriously polluted wetland that significantly affects plant growth, and the lost coastal wetland due to sea level rise (Zedler, 2004).




## Photos of the practice



**Photo 14.** Wetland restoration project in Qixinghe National Natural Reserve, Sanjiang Plain, China

The left photo shows the cultivated wetland before restoration (in 2014), which had been reclaimed for corn for about 30 years. The right photo shows the wetland after restoration (in 2018). Restorations consisted of reflooding by removal of ditches and tile lines. The restored wetland site was dominated by *Phragmites australis* and *Carex species*



**Photo 15.** Wetland restoration project in Niuxintaobao National Wetland Park, Songnen Plain, China

The left photo shows the degraded wetland before restoration (in 2010). The right photo shows the wetland after restoration (in 2013). Restoration practices included water supplement in combined with the promotion of plant growth and reproduction, e.g., raking and plant (or rhizome) transplantation. The restored wetland site was dominated by *Phragmites australis*

**Table 40.** Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Preserving Soil Organic Carbon in Prairie Wetlands of Central North America</i>	North America	Various	6	19
<i>Maintenance of Marshlands in Urban Tidal Wetlands in New York City, United States</i>	North America	100	6	31

## References

- Audet, J., Zak, D., Bidstrup, J. & Hoffmann, C.C.** 2020. Nitrogen and phosphorus retention in Danish restored wetlands. *Ambio*, 49(1): 324-336. <https://doi.org/10.1007/s13280-019-01181-2>
- Badiou, P., McDougal, R., Pennock, D. & Clark, B.** 2011. Greenhouse gas emissions and carbon sequestration potential in restored wetlands of the Canadian prairie pothole region. *Wetlands Ecology and Management*, 19(3): 237–256. <https://doi.org/10.1007/s11273-011-9214-6>
- Burden, A., Garbutt, A. & Evans, C.D.** 2019. Effect of restoration on saltmarsh carbon accumulation in Eastern England. *Biology Letters*, 15(1): 20180773. <https://doi.org/10.1098/rsbl.2018.0773>
- Carnell, P.E., Windecker, S.M., Brenker, M., Baldock, J., Masque, P., Brunt, K. & Macreadie, P.I.** 2018. Carbon stocks, sequestration, and emissions of wetlands in south eastern Australia. *Global Change Biology*, 24(9): 4173–4184. <https://doi.org/10.1111/gcb.14319>
- Chen, Q.F., Guo, B.B., Zhao, C.S. & Xing, B.X.** 2018. Characteristics of CH<sub>4</sub> and CO<sub>2</sub> emissions and influence of water and salinity in the Yellow River delta wetland, China. *Environmental Pollution*, 239: 289–299. <https://doi.org/10.1016/j.envpol.2018.04.043>
- Crooks, S., Herr, D., Tamelander, J., Laffoley, D. & Vandever, J.** 2011. Mitigating climate change through restoration and management of coastal wetlands and near-shore marine ecosystems : Challenges and opportunities. Washington, DC. (also available at: <https://openknowledge.worldbank.org/handle/10986/18318>)
- Daniel, D.W., Smith, L.M. & McMurry, S.T.** 2017. Effects of sediment removal and surrounding land use on carbon and nitrogen storage in playas and watersheds in the Rainwater Basin region of Nebraska. *Soil and Tillage Research*, 174: 169-176. <https://doi.org/10.1016/j.still.2017.07.001>
- Doherty, J.M., Callaway, J.C. & Zedler, J.B.** 2011. Diversity-function relationships changed in a long-term restoration experiment. *Ecological Applications*, 21(6): 2143-2155. <https://doi.org/10.1890/10-1534.1>
- Ducks Unlimited Canada.** 2020. Our work, impact areas, Wetlands. [online]. [Cited 28 October 2020]. <https://www.ducks.ca/our-work/wetlands/>
- Gleason, R.A., Tangen, B.A., Browne, B.A. & Euliss, N.H.** 2009. Greenhouse gas flux from cropland and restored wetlands in the Prairie Pothole Region. *Soil Biology & Biochemistry*, 41(12): 2501-2507. <https://doi.org/10.1016/j.soilbio.2009.09.008>
- Hodge, I. & McNally, S.** 2000. Wetland restoration, collective action and the role of water management institutions. *Ecological Economics*, 35(1): 107-118. [https://doi.org/10.1016/S0921-8009\(00\)00171-3](https://doi.org/10.1016/S0921-8009(00)00171-3)
- Johnston, S.G., Burton, E.D., Aaso, T. & Tuckerman, G.** 2014. Sulfur, iron and carbon cycling following hydrological restoration of acidic freshwater wetlands. *Chemical Geology*, 371: 9-26. <https://doi.org/10.1016/j.chemgeo.2014.02.001>
- Li, W., Dou, Z., Cui, L., Zhao, X., Zhang, M., Zhang, Y., Gao, C., Yang, Z., Lei, Y. & Pan, X.** 2020. Soil fauna diversity at different stages of reed restoration in a lakeshore wetland at Lake Taihu, China. *Ecosystem Health and Sustainability*, 6(1): 1722034. <https://doi.org/10.1080/20964129.2020.1722034>

- Li, X.R., Xiao, Y.P., Ren, W.W., Liu, Z.F., Shi, J.H. & Quan, Z.X.** 2012. Abundance and composition of ammonia-oxidizing bacteria and archaea in different types of soil in the Yangtze River estuary. *J Zhejiang Univ Sci B*, 13(10): 769-782. <https://doi.org/10.1631/jzus.B1200013>
- Luo, Z., Deng, L. & Yan, C.** 2015. Soil erosion under different plant cover types and its influencing factors in Napahai Catchment, Shangri-La County, Yunnan Province, China. *International Journal of Sustainable Development and World Ecology*, 22(2): 135-141. <https://doi.org/10.1080/13504509.2014.924448>
- Ma, M., Zhou, X., Ma, Z. & Du, G.** 2012. Composition of the soil seed bank and vegetation changes after wetland drying and soil salinization on the Tibetan Plateau. *Ecological Engineering*, 44: 18-24. <https://doi.org/10.1016/j.ecoleng.2012.03.017>
- Marambanyika, T. & Beekedahl, H.** 2016. The missing link between awareness and the implementation of wetland policy and legislation in communal areas of Zimbabwe. *Wetlands Ecology and Management*, 24(5): 545-563. <https://doi.org/10.1007/s11273-016-9486-y>
- McLaughlin, D.L. & Cohen, M.J.** 2013. Realizing ecosystem services: wetland hydrologic function along a gradient of ecosystem condition. *Ecological Applications*, 23(7): 1619-1631. <https://doi.org/10.1890/12-1489.1>
- Meli, P., Rey Benayas, J.M., Balvanera, P. & Martinez Ramos, M.** 2014. Restoration enhances wetland biodiversity and ecosystem service supply, but results are context-dependent: a meta-analysis. *Plos One*, 9(4): e93507. <https://doi.org/10.1371/journal.pone.0093507>
- Moreno-Mateos, D., Power, M.E., Comin, F.A. & Yockteng, R.** 2012. Structural and functional loss in restored wetland ecosystems. *Plos Biology*, 10(1):e1001247. <https://doi.org/10.1371/journal.pbio.1001247>
- Nadeau, C.P. & Conway, C.J.** 2015. Optimizing water depth for wetland-dependent wildlife could increase wetland restoration success, water efficiency, and water security. *Restoration Ecology*, 23(3):292-300. <https://doi.org/10.1111/rec.12180>
- Olsson, L., Ye, S., Yu, X., Wei, M., Krauss, K.W. & Brix, H.** 2015. Factors influencing CO<sub>2</sub> and CH<sub>4</sub> emissions from coastal wetlands in the Liaohe Delta, Northeast China. *Biogeosciences*, 12(16): 4965-4977. <https://doi.org/10.5194/bg-12-4965-2015>
- Renzi, J.J., He, Q. & Silliman, B.R.** 2019. Harnessing positive species interactions to enhance coastal wetland restoration. *Frontiers in Ecology and Evolution*, 7: 131. <https://doi.org/10.3389/fevo.2019.00131>
- Scholte, S.S.K., Todorova, M., van Teeffelen, A.J.A. & Verburg, P.H.** 2016. Public support for wetland restoration: what is the link with ecosystem service values? *Wetlands*, 36(3): 467-481. <https://doi.org/10.1007/s13157-016-0755-6>
- Smith, C.S., Serra, L., Li, Y.C., Inglett, P. & Inglett, K.** 2011. Restoration of disturbed lands: the Hole-in-the-Donut restoration in the Everglades. *Critical Reviews in Environmental Science and Technology*, 41: 723-739. <https://doi.org/10.1080/10643389.2010.530913>

- Song, Y.Y., Song, C.C., Yang, G.S., Miao, Y.Q., Wang, J.Y. & Guo, Y.D.** 2012. Changes in labile organic carbon fractions and soil enzyme activities after marshland reclamation and restoration in the Sanjiang Plain in Northeast China. *Environmental Management*, 50(3): 418-426. <https://doi.org/10.1007/s00267-012-9890-x>
- Su, H., Guo, P., Zhang, Y., Deng, J., Wang, M., Sun, Y. & Wu, Y.** 2019. Effects of planting patterns on the concentration and bioavailability of heavy metals in soils during wetland restoration. *International Journal of Environmental Science and Technology*, 16(2): 853-864. <https://doi.org/10.1007/s13762-018-1724-9>
- Suir, G.M., Sasser, C.E., DeLaune, R.D. & Murray, E.O.** 2019. Comparing carbon accumulation in restored and natural wetland soils of coastal Louisiana. *International Journal of Sediment Research*, 34(6): 600-607. <https://doi.org/10.1016/j.ijsrc.2019.05.001>
- Tangen, B.A., Finocchiaro, R.G. & Gleason, R.A.** 2015. Effects of land use on greenhouse gas fluxes and soil properties of wetland catchments in the Prairie Pothole Region of North America. *Science of the Total Environment*, 533: 391-409. <https://doi.org/10.1016/j.scitotenv.2015.06.148>
- Thi, T.D. & Minh, D.D.** 2019. Riverbank stability assessment under river water level changes and hydraulic erosion. *Water*, 11(12): 2598. <https://doi.org/10.3390/w11122598>
- Villa, J.A. & Bernal, B.** 2018. Carbon sequestration in wetlands, from science to practice: An overview of the biogeochemical process, measurement methods, and policy framework. *Ecological Engineering*, 114: 115-128. <https://doi.org/10.1016/j.ecoleng.2017.06.037>
- Wang, G., Wang, M., Lu, X. & Jiang, M.** 2015. Effects of farming on the soil seed banks and wetland restoration potential in Sanjiang Plain, Northeastern China. *Ecological Engineering*, 77: 265-274. <https://doi.org/10.1016/j.ecoleng.2015.01.039>
- Wang, H., Wang, R.Q., Yu, Y., Mitchell, M.J. & Zhang, L.J.** 2011. Soil organic carbon of degraded wetlands treated with freshwater in the Yellow River Delta, China. *Journal of Environmental Management*, 92(10): 2628-2633. <https://doi.org/10.1016/j.jenvman.2011.05.030>
- Wang, X., Zhang, D., Guan, B., Qi, Q. & Tong, S.** 2017. Optimum water supplement strategy to restore reed wetland in the Yellow River Delta. *Plos One*, 12(5): e0177692. <https://doi.org/10.1371/journal.pone.0177692>
- Xu, S., Liu, X., Li, X. & Tian, C.** 2019. Soil organic carbon changes following wetland restoration: A global meta-analysis. *Geoderma*, 353: 89-96. <https://doi.org/10.1016/j.geoderma.2019.06.027>
- Xu, S., Sheng, C. & Tian, C.** 2020. Changing soil carbon: influencing factors, sequestration strategy and research direction. *Carbon Balance and Management*, 15(1): 2. <https://doi.org/10.1186/s13021-020-0137-5>
- Xu, S., Zhang, B., Ma, L., Hou, A., Tian, L., Li, X. & Tian, C.** 2017. Effects of marsh cultivation and restoration on soil microbial communities in the Sanjiang Plain, Northeastern China. *European Journal of Soil Biology*, 82: 81-87. <https://doi.org/10.1016/j.ejsobi.2017.08.010>

**Zedler, J.B.** 2004. Compensating for wetland losses in the United States. *Ibis*, 146: 92-100.

<https://doi.org/10.1111/j.1474-919X.2004.00333.x>

**Zedler, J.B. & Kercher, S.** 2005. Wetland resources: Status, trends, ecosystem services, and restorability. *Annual Review of Environment and Resources*, 30: 39-74.

<https://doi.org/10.1146/annurev.energy.30.050504.144248>

**Zhu, H., Guan, Z. & Wei, X.** 2016. Factors influencing farmers' willingness to participate in wetland restoration: evidence from China. *Sustainability*, 8(12): 1325. <https://doi.org/10.3390/su8121325>

# 11. Conservation of pristine peatlands and avoiding drainage of peatlands

Laura Villegas<sup>1</sup>, Felix Beer<sup>2</sup>, Maria Nuutinen<sup>1</sup>, Kai Milliken<sup>1</sup>

<sup>1</sup>Food and Agriculture Organization of the United Nations (FAO), Rome, Italy

<sup>2</sup>University of Greifswald, partner in the Greifswald Mire Centre, Greifswald, Germany

## 1. Description of the practice

Conservation of peatlands, especially the avoidance of draining areas, means to refrain from all actions that disrupt natural ecosystem functions. Activities like drainage, fertilizing, tree cutting, slash and burn, infrastructure development (e.g. construction of roads, trails), and other factors may cause the lowering of natural water table levels, leading to SOC loss by aerobic degradation and dissolution of SOC in water (DOC). Degradation of the peat layer results in negative effects on land, causing subsidence (i.e. lowering of the peat surface) GHG emissions, and decreases community resilience to climate change (See [Hotspots: Peatlands](#)).

For conservation it is essential to map peatlands, develop management and monitoring plans, and ensure that local populations have access to livelihood sources that do not threaten peatlands. Knowing the location, extent, and carbon storage of peat deposits, as well as understanding the ecosystem services they provide (see sections 3 and 4 below), are necessary steps towards establishing suitable conservation strategies. Conservation is often a very cost-effective land-use management option (Joosten, Tapio-Biström and Tol, 2012) that still allows a range of non-invasive, livelihood-development activities (e.g. collection of berries, honey production, fibre, feed, small-scale livestock rearing, or development of non-degrading ecotourism).

## 2. Range of applicability

Pristine peatlands occur in more than 160 countries (see [Hotspots: Peatlands](#)) and currently many initiatives are working to better map and assess peatlands (FAO, 2020). An increasing number of countries are interested in knowing the actual location and extent of their peatlands, and to develop sustainable, drainage-free livelihood options. The assessment of peatlands is the starting point for avoiding their drainage as their conservation can be better realized. Conservation as a management practice should be applied to all pristine peatlands globally, in particular and most urgently in zones with high pressure for peatland drainage and vegetation removal.

Maintaining the hydrological conditions and the nutrient status of a peatland ought to be the objective to avoid SOC losses. In order to achieve this, it is critical to maintain typical biotic communities, natural succession, and linkages between peatlands within a landscape. On the other hand, drainage and nutrient discharge should be avoided. However, peatland-specific analysis of the situation and threats can better inform the prioritization of conservation activities. In some cases, ecotourism and non-timber product extraction can be compatible with conservation and restoration purposes, while contributing to the creation of incentives to promote conservation.

## 3. Impact on soil organic carbon stocks

Peatland conservation maintains SOC in the peat, avoiding anthropogenic GHG emissions, and securing carbon sequestration due to long-term peat accumulation. Globally the area of near-pristine peatlands – more than three million km<sup>2</sup> – are estimated to sequester 0.37 Gt CO<sub>2</sub> per year in the soil. Similarly, degrading peatlands worldwide emit estimated 1.3 Gt CO<sub>2</sub> per year, equivalent to at least 5 percent of global anthropogenic CO<sub>2</sub> emissions (IPCC, 2014). An estimated 336–644 Gt SOC can be potentially conserved in the peat soil if peatlands are conserved, and drainage avoided. Peat usually accumulates slowly, at the rate of about a millimetre per year, which makes peatlands slow carbon sinks. Also, their potential for SOC maintenance lies in the avoided carbon emissions of the carbon stored, and not on the yearly sequestration rate, which is only considerable over long temporal scales (Soares *et al.*, 2015; Donato *et al.*, 2011).

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Avoiding drainage of pristine peatlands contributes to land stabilization, nutrient balance, soil biodiversity and moisture regimes, allowing the peat to continuously accumulate SOC.



## 4.2 Minimization of threats to soil functions

Table 41. Soil threats

Soil threats	
<b>Soil erosion</b>	Avoiding drainage, conservation of the peat layer and pristine vegetation cover is crucial for avoiding peat erosion. Pristine peatlands regulate water run-off, whereas dry and bare peat is exposed to oxidation and is easily eroded, even in flat areas, causing quick surface lowering (land loss or subsidence), water and carbon losses (Joosten, 2015).
<b>Nutrient imbalance and cycles</b>	Intact peatlands retain nutrients and help purify surface waters (Succow and Joosten, 2001).
<b>Soil salinization and alkalinization</b>	Conserved coastal peat swamps act as a buffer between salt- and freshwater systems, preventing saline intrusion into coastal lands (Silvius, Joosten and Opdam, 2008) and the eventual loss of adjacent land (Joosten, 2015).
<b>Soil contamination/pollution</b>	Intact peatlands retain pollutants and help filtering and purifying surface water (Wichtmann, Schröder, and Joosten, 2016).
<b>Soil acidification</b>	Depending on peatland type: ombrotrophic peatlands are already acidic (Succow and Joosten, 2001).
<b>Soil biodiversity loss</b>	Maintaining the pH, SOC (Mandic-Mulec <i>et al.</i> , 2014), plant cover and specific physicochemical characteristics of peat (Opelt <i>et al.</i> , 2007) might be determinant of the composition of a microbial community.
<b>Soil compaction</b>	Compaction caused by soil decomposition and mineralization is avoided when peatlands are wet.
<b>Soil water management</b>	Intact peatlands have unique and vulnerable soil moisture regimes (Dommain, Couwenberg and Joosten, 2010). Maintenance of the natural regime prevents irreversible drying – when peat soils are exposed to prolonged, intensive drainage, the peat becomes hydrophobic, rendering it very difficult to rewet and restore given that the peat soil diminishes its capacity to retain water. Peats reaching this stage lose water holding capacity (up to 40–75 percent) (Andriess, 1988) compared to wet peat soil.

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Although intensive productive activities should be prevented, some subsistence and income-generating activities related, for example, to extraction of non-timber products, like honey, and other activities, like fishing, and hunting, can be compatible with conservation activities. In fact, conserved peatlands ensure, in many cases, local food security and can regulate water supply for irrigation of downstream productive sites outside the peatlands.

### 4.4 Mitigation of and adaptation to climate change

Peatland conservation and avoiding drainage contributes directly to climate change mitigation as well as increasing adaptive capacity. Preventing drying caused by drainage or other activities avoids new GHG emissions from degradation and from fires. It also allows conserved peatlands to protect the coastal areas from extreme weather events, while regulating water supply and reducing fire risks in the catchment (see Hotspot “Peatlands” in volume 2 of this manual).

### 4.5 Socio-economic benefits

Peatland conservation and avoiding drainage are the most cost-effective ways to manage peatlands, and lower the social and climate cost of peatland degradation (Joosten *et al.*, 2012). These activities secure the provision of ecosystem services that support economies and human well-being (see Hotspot “Peatlands” in volume 2 of this manual).

## 5. Potential drawbacks to the practice

### 5.1 Increases in greenhouse gas emissions

None are known. Pristine peatlands are climate neutral, and in most cases are slow carbon sinks (see above section 3 on impact on SOC sequestration). Conservation would also make peatlands less vulnerable to drying and other climatic changes, helping to reduce non-anthropogenic GHGs (Swindles *et al.*, 2019).

### 5.2 Conflict with other practice(s)

Avoiding drainage of peatlands and promoting their conservation reduces options to establish other practices on peatlands, except in some cases ecotourism and extraction of non-timber products on a small scale (Crump, 2017). However, this is justified by the long-term benefits of conservation and avoided drainage in the landscape, ecosystem services provision, and reduction of environmental risk to communities.

### 5.3 Decreases in production (e.g. food/fuel/feed/timber)

In the short term, preserving pristine peatlands implies reduced availability of peatlands for degradation and conversion into drainage-based production systems (e.g. oil palm, timber plantations). Intensive monoculture agricultural practices result in long-term land loss and degradation, disruption of ecosystem services, increased emissions and loss of adaptive capacity of the landscape, including surrounding communities. Considering that peatlands represent only three percent of the earth's surface, refraining from using these lands as productive areas will secure long-term ecosystem services and a positive long-term impact on the regions where they are located.

## 6. Recommendations before implementation of the practice

Peatland location and boundaries should be clearly demarcated in the land-use planning maps:

- ◆ Each peatland should be considered as one hydrological unit, meaning that draining the water in a part of a peatland, will affect the groundwater table in the whole peatland. Therefore, management measures should address the entire peatland entity to reap maximum benefits in terms of biodiversity or carbon conservation. The maintenance of high water tables should be prioritized to conserve the SOC in peat soils, noting that changes in hydrology of the landscape outside the peatland area may affect peatland hydrology.
- ◆ Conservation of the peatland vegetation cover is important to secure peat formation and therefore long-term carbon storage.
- ◆ Due to their relevance for ecosystem services, hydrology, and the adaptive capacity of surrounding communities and landscapes, peatlands should be excluded from drainage-based activities, i.e. set aside as conservation areas.

## 7. Potential barriers to adoption

Table 42. Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Potential changes in the rainfall due to climatic changes and land cover changes, such as deforestation.
Cultural	Yes	Conflicts with traditional and cultural land-use practices that include drainage for production/extraction.
Social	Yes	Social acceptance of prohibition/limitation of activities on the peatlands; social perceptions of different conservation interventions (Harrison <i>et al.</i> , 2019).
Economic	Yes	Conflict with activities that generate short-term revenue but cause important losses in the longer term (Harrison <i>et al.</i> , 2019).
Institutional	Yes	Lack of funds and incentives to promote conservation over drainage-based alternatives (Harrison <i>et al.</i> , 2019).
Legal (Right to soil)	Yes	Frequent lack of clarity regarding wetlands' legal status and responsibility for different areas and activities, conflicting/unclear laws and ineffective law enforcement (Harrison <i>et al.</i> , 2019).
Knowledge	Yes	Lack of data and knowledge to assess conservation potential and fire impacts, the effect of conservation efforts, specific dynamics and characteristics of each peatland type and location (Harrison <i>et al.</i> , 2019).
Other: Drainage-based livelihoods	Yes	Drainage-based peatland management such as grazing, cropping, plantations and forestry activities often prevent conservation and the maintenance of the natural ground water level (FAO, 2012).

## References

- Andriessse, J.P.** 1988. Irreversible drying. In J.P. Andriessse (Ed.) *Nature and management of tropical peat soils*. pp. 27. FAO Soils Bulletin 59. Rome, Italy.
- Crump, J.** 2017. *Smoke on water – Countering global threats from peatland loss and degradation*. United Nations Environment Programme and GRID-Arendal. Nairobi and Arendal.
- Dommain, R., Couwenberg, J. & Joosten, H.** 2010. Hydrological self-regulation of domed peatlands in south-east Asia and consequences for conservation and restoration. *Mires and Peat*, 6(05): 1–17.
- Donato, D., Kauffman, J., Murdiyarso, D., Kurnianto, S., Stidham, M. & Kanninen, M.** 2011. Mangroves among the most carbon-rich forests in the tropics. *Nature Geoscience*, 4: 293–297. <https://doi.org/10.1038/ngeo1123>
- FAO.** 2020. *Peatlands mapping and monitoring – Recommendations and technical overview*. Rome, Italy. (also available at: <http://www.fao.org/3/CA8200EN/CA8200EN.pdf>)
- FAO.** 2012. *Peatlands – guidance for climate change mitigation through conservation, rehabilitation and sustainable use*. Rome, Italy. (also available at: <http://www.fao.org/3/an762e/an762e.pdf>)
- Harrison, M.E., Ottay, J.B., D’Arcy, L.J., Cheyne, S.M., Anggodo, Belcher, C., Cole, L., Dohong, A., Ermiasi, Y., Feldpausch, T., Gallego-Sala, A., Gunawan, A., Höing, A., Husson, S.J., Kulu, I.P., Soebagio, S.M., Mang, S., Mercado, L., Morrogh-Bernard, H.C., Page, S.E., Priyanto, R., Capilla, B.R., Rowland, L., Santos, E.M., Schreer, V., Sudyana, I.N., Taman, S.B.B., Thornton, S.A., Upton, C., Wich, S.A. & Veen, F.J.F. van.** 2020. Tropical forest and peatland conservation in Indonesia: Challenges and directions. *People and Nature*, 2(1): 4–28. <https://doi.org/10.1002/pan3.10060>
- IPCC.** 2014. 2013 *Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands*. T. Hiraiishi, T. Krug, K. Tanabe, N. Srivastava, B. Jamsranjav, M. Fukuda & T. Troxler (Eds.) Switzerland.
- Joosten, H., Tapio-Biström, M. & Tol, S.** 2012. *Peatlands – guidance for climate change mitigation through conservation, rehabilitation and sustainable use*. FAO and Wetlands International. Rome, Italy. (also available at: <http://www.fao.org/3/an762e/an762e.pdf>)
- Joosten, H.** 2015. Peatlands, climate change mitigation and biodiversity conservation. Policy brief for the Nordic Council of Ministers. Copenhagen, Denmark. (also available at: [https://www.ramsar.org/sites/default/files/documents/library/ny\\_2.\\_korrektur\\_anp\\_peatland.pdf](https://www.ramsar.org/sites/default/files/documents/library/ny_2._korrektur_anp_peatland.pdf))
- Mandic-Mulec, I., Ausec, L., Danevcic, T., Levicnik-Höfferle, S., Jerman, V. & Kraigher, B.** 2014. Microbial Community Structure and Function in Peat Soil. *Food Technol. Biotechnol.*, 52(2): 180–187.
- Opelt, K., Berg, C., Schonman, S., Eberl, L. & Berg, G.** 2007. High specificity but contrasting biodiversity of Sphagnum-associated bacterial and plant communities in bog ecosystems independent of the geographical region. *ISME J*, 1(6): 502–516. <https://doi.org/10.1038/ismej.2007.58>

**Silvius, M.J., Joosten, H. & Opdam, S.** 2008. Peatlands and people. *In* F. Parish, A. Sirin, D. Charman, H. Joosten, T. Minayeva, M.J. Silvius & L. Stringer (Eds.) *Assessment on Peatlands, Biodiversity and Climate Change: Main Report*. pp. 20–28. Global Environment Centre and Wetlands International. Wageningen, the Netherlands. (also available at:

[http://www.imcg.net/media/download\\_gallery/books/assessment\\_peatland.pdf](http://www.imcg.net/media/download_gallery/books/assessment_peatland.pdf))

**Soares, P., Zuchello, F., Anjos, L., Pereira, M. & Oliveira, A.** 2015. Soil attributes and C and N variation in histosols under different agricultural usages in the state of Rio de Janeiro, Brazil. *Bioscience Journal*, 31: 1349–1362. <https://doi.org/10.14393/BJ-v31n5a2015-26365>

**Swindles, G.T., Morris, P.J., Mullan, D.J., Payne, R., Roland, T., Amesbury, M.J., Lamentowicz, M., Turner, E., Gallego-Sala, A., Sim, T., Barr, L.D., Blaauw, M., Blundell, A., Chambers, F.M., Charman, D.J., Feurdean, A., Galloway, J.M., Gałka, M., Green, S.M., Kajukalo, K., Karofeld, E., Korhola, A., Lamentowicz, L., Langdon, P., Marcisz, K., Mauquoy, D., Mazei, Y.A., McKeown, M.M., Mitchell, E.A.D., Novenko, E., Plunkett, G., Roe, H.M., Schoning, K., Sillasoo, Ü., Tsyganov, A.N., van der Linden, M., Väliranta, M. & Warner, B.** 2019. Widespread drying of European peatlands in recent centuries. *Nature Geosciences*, 12: 922–928. <https://doi.org/10.1038/s41561-019-0462-z>

**Succow, M. & Joosten, H.** 2001. *Landschaftsökologische Moorkunde, 2nd edition [Landscape ecology of mires]*. Schweizerbart. Stuttgart, Germany.

**Wichtmann, W., Schröder, C. & Joosten, H.** 2016. *Paludiculture – productive use of wet peatlands*. Schweizerbart Science Publishers. Stuttgart, Germany

## 12. Restoration of peatlands

Felix Beer<sup>1</sup>, Laura Villegas<sup>2</sup>, Maria Nuutinen<sup>2</sup>, Fahmuddin Agus<sup>3</sup>

<sup>1</sup>University of Greifswald, partner in the Greifswald Mire Centre, Germany

<sup>2</sup>Food and Agriculture Organization of the United Nations, Rome, Italy

<sup>3</sup>Indonesian Soil Research Institute, Bogor, Indonesia

### 1. Description of the practice

The restoration of degraded peatlands consists of two equally important steps: i) rewetting of the peatland and ii) revegetation with peat-generating species of ideally economic viability, if needed (FAO, 2014; FAO, 2020).

Rewetting consists of raising the mean annual groundwater table of the whole hydrological unit of a drained peatland, ideally back to surface level (Jauhiainen, Page and Vasander, 2016; Tannenberger *et al.*, 2020). This way, the oxygen-deficient soil conditions are re-established and further oxidation of the peat is minimized, which also reduces subsidence<sup>2</sup> (Jauhiainen, Page and Vasander, 2016) and the loss of SOC as greenhouse gas emissions (Günther *et al.*, 2020; Wakhid *et al.*, 2017) and as organic carbon dissolved in water (DOC). Moreover, with restoration, the risk of peat fires is minimized. For effective results, and following the greenhouse gas emission reduction method of the Wetland Supplement (IPCC, 2014), the whole peatland hydrological unit needs to be rewetted. Rewetting is done by:

- ◆ blocking drainage canals and ditches (e.g. with peat collected at site);
- ◆ raising overflow heights of weirs, “dams” or “canal blockings” and sluices;
- ◆ establishing and allowing obstructions in water courses (trees, rocks, vegetation growth, beaver dams);
- ◆ removing subsurface drainage pipes;
- ◆ reducing evapotranspiration from tree growth in the peatland (only in originally treeless peatlands);

---

<sup>2</sup> Subsidence is the lowering of the surface caused by peat compaction and oxidation (see e.g. FAO, 2020).

- ◆ establishing hydrological buffer zones with higher water levels; and/ or
- ◆ in low-lying coastal peatland areas, such as ‘polders’<sup>3</sup> in the Netherlands, which are protected by dikes or other artificial barriers and drained through active pumping, the pumping must be reduced to establish close-to-surface water tables (Couwenberg, 2018).

Revegetation is the re-establishment of vegetation that is adapted to wet soil conditions. Revegetation can happen through natural regeneration or active sowing or planting of the site’s native wetland plants. (FAO, 2014; FAO, 2020). Rewetting addresses the negative effects of drainage, while revegetation helps regulate water balance and is necessary for the peatland to recover, and potentially become a carbon sink again.

## 2. Range of applicability

The reasons for peatland restoration are growing and include the achievement of peatland biodiversity and/or recuperation of water regulation and other ecosystem functions and services, such as avoidance of greenhouse gas emissions and other carbon losses (FAO, 2014). The potential and need of restoring peat soils for SOC preservation has been developed in degraded boreal, subarctic, temperate, as well as tropical peatlands (Joosten, Tanneberger and Moen, 2017; Giesen and Nirmala, 2018).

## 3. Impact on soil organic carbon stocks

Peat soils consist of partly preserved dead plant material that built up often over thousands of years at an average rate of 0.5–1 mm per year in temperate and boreal regions, and up to 2.5 mm per year in tropical zones (Page *et al.*, 2010). Vegetation<sup>4</sup> cover of native peatland species is crucial to re-establish SOC and restoring peat accumulation over time (Jauhiainen, Page and Vasander, 2016). That said, carbon sequestration in peat soils is a slow process that happens over centuries. When the IPCC Wetlands Supplement was published (2014) there was a lack of data on soil carbon sequestration rates in years or decades following peatland rewetting (IPCC, 2014). However, a recent study from temperate climates illustrated that the carbon sink function of a restored peatlands can be approximately 1 tonne C per ha per year, depending on the vegetation and water regime (Swenson *et al.*, 2019).

---

<sup>3</sup> A polder is a low-lying tract of land that forms an artificial hydrological entity, enclosed by embankments known as dikes. In polders, ground-water tables need to be kept artificially low through permanent pumping.

<sup>4</sup> To note: revegetation (including replanting) is necessary only when natural plant regeneration is not possible due to e.g. burning of the soil seed bank or over 5 km distance from peatland areas with endemic wetland vegetation (see e.g. FAO, 2020).



## 4. Other benefits of the practice

### 4.1. Minimization of threats to soil functions

Peatland restoration positively affects soil characteristics. Water run-off of restored peatlands can be reduced by 25 percent compared to a peatland with no restoration efforts (Shantz and Price, 2006) due to increased water storage capacities and despite increased hydraulic conductivity of newly regrown soil vegetation (Wallage and Holden, 2011). Reduced bulk density increases water and nutrient movement, reduces soil moisture fluctuations, and prevents soil erosion (Wichtmann *et al.*, 2016). However, the establishment of pre-drainage hydrological and soil conditions necessarily takes place over a similar temporal scale as peat formation itself and, in some cases, restoration efforts will never achieve the pristine state found prior to drainage due to changed site conditions.

### 4.2 Minimization of threats to soil functions

Table 43. Soil threats

Soil threats	
<b>Soil erosion</b>	Erosion by surface water run-off is minimized through canal blocking and managed grazing. Wind erosion from bare peat is minimized by permanent vegetation cover (Wichtmann, Schröder and Joosten, 2016).
<b>Nutrient imbalance and cycles</b>	Restored peatlands can regain nutrient retention and water purification functions (Succow and Joosten, 2001).
<b>Soil salinization and alkalinization</b>	In coastal regions: peatland restoration stops subsidence, and could prevent saltwater intrusion and/or flooding (Hooijer <i>et al.</i> , 2015).
<b>Soil contamination/pollution</b>	Peatlands can act as water filtering systems and may retain nutrients and pollutants.
<b>Soil acidification</b>	Depending on peatland type: ombrotrophic peatlands already are acidic (Succow and Joosten, 2001). Rewetting of acid sulphate peat soil reduces the risk of acidification (Wösten <i>et al.</i> , 2006).
<b>Soil biodiversity loss</b>	Loss of non-peatland soil organisms (earthworms, moles, rodents) is a positive sign of successful peatland restoration.
<b>Soil compaction</b>	Compaction caused by soil decomposition and mineralization is halted, wetland plant establishment recreates original topsoil conditions with low bulk density, high hydraulic conductivity, etc. (Wichtmann, Schröder and Joosten, 2016).

<b>Soil threats</b>	
<b>Soil water management</b>	Restored peatlands with wetland vegetation self-regulate water run-off and provide a hydrological buffer function (Dommain, Couwenberg and Joosten, 2010; Wallage and Holden, 2011).

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Rewetting establishes conditions for the growth of wetland plants that can be used for a range of purposes: energy production, fibre for artisanal and construction materials, fodder, horticultural substrates (Wichtmann, Schröder and Joosten, 2016), and also for food (Giesen and Nirmala, 2018). Productivity depends on species and site characteristics but can be high compared to conventional drainage-based agriculture (Zerbe *et al.*, 2013).

### 4.4 Mitigation of and adaptation to climate change


Rewetting of peatlands minimizes CO<sub>2</sub> emissions immediately (Carlson *et al.*, 2015; Couwenberg, 2018). Methane emissions may increase in the first years after rewetting (Günther *et al.*, 2020). Models of global warming show the potential trade-offs between the CO<sub>2</sub> and N<sub>2</sub>O (long-lived GHGs) emissions of drained peatlands versus increased CH<sub>4</sub> (a short-lived GHG) emissions of rewetted peatlands and suggest that rewetting would greatly reduce the global warming potential of these ecosystems in decades to come (Günther *et al.*, 2020; Wilson *et al.*, 2016; Figure 6). Thus, rewetting all or half of global peatlands now would decrease global heating more than rewetting later, or never rewetting peatlands (Günther *et al.*, 2020).

If bringing the water table back to surface level entirely is not practicable (e.g. due to economic constraints) partial rewetting, as applied on drained peatlands in Indonesia and Malaysia, reduces SOC losses and fire risk, and thus, can partially help but not solve issues related to subsidence, fire and degradation (Carlson, Goodman and May-Tobin, 2015; Wakhid *et al.*, 2017, see figure 1 in factsheet n°13 “Paludiculture”, this volume). However, blocking only a part of drainage system only slows down soil degradation, and should be regarded as an intermediary step towards full rewetting.

Additionally, drained soils are at increased risk of catastrophic peat fire and losses of carbon stores (Turetsky *et al.*, 2015). Peatland restoration thus decreases the risk of dramatic negative impacts on human livelihoods as well as on the global carbon cycle from peat fires (Page and Hooijer, 2016).

**Note:** The authors of this chapter acknowledge the contributions from Beth A. Middleton, Eric Ward and Lorenzo Menichetti in section 4.4 on climate change mitigation and adaptation.

## 4.5 Socio-economic benefits



**Figure 6.** Effects of greenhouse gas emissions from peatlands following global rewetting scenarios, and related radiative forcing (RF) and global warming effect

The gray area shows the period of rewetting. Source: Günther *et al.*, 2020

Restoration halts peatland subsidence – lowering of the peat surface – and related loss of coastal and riverine areas and helps to avoid peatlands becoming undrainable over time (Hooijer *et al.*, 2015), therefore also protecting lives, infrastructure and livelihoods. Negative health and economic impacts of fire and haze are prevented with restoration measures. Restored peatlands can provide sustainable livelihoods with a range of options for wet biomass production and other economic activities (see factsheet n° 13 on paludiculture, this volume).

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

None are known.

### 5.2 Increases in greenhouse gas emissions

Whilst rewetting of peatlands immediately reduces CO<sub>2</sub> emissions, the CH<sub>4</sub> emissions can increase depending on the water level for a few years (Wilson *et al.*, 2016; Jauhiainen, Page and Vasander, 2016). As a consequence, restored peatlands should not be flooded, but it is recommended to manage and maintain the medium annual ground water table close to or at the surface (Wichtmann, Schröder and Joosten, 2016). Furthermore, methane emissions are not as important as CO<sub>2</sub> emissions for long term climate impact, as methane breaks down quicker in the atmosphere compared to long-lived CO<sub>2</sub> (Günther *et al.*, 2020; Figure 6). Wetland plants (e.g. reed or cattail) in the temperate zone lead to decreased CH<sub>4</sub> emissions compared to rewetted bare peat soils (Swenson *et al.*, 2019; Vroom *et al.*, 2018), highlighting the importance of revegetation for preserving SOC and reducing the global warming potential.

### 5.3 Conflict with other practice(s)

Invasive activities (e.g. intensive agriculture or plantations), which would include tillage, drainage, annual crops, or high livestock density with a significant trampling impact, cannot be aligned with restoration of peatlands.

### 5.4 Decreases in production (e.g. food/fuel/feed/timber)

Restoration of peatlands is not compatible with conventional drainage-based land-use. There are established traditional peatland uses for various biomass extraction, and similarly alternative productive systems based on biomass production in wet peatland condition (Paludiculture, see factsheet n° 13 of this volume) are being studied through pilot projects (see case studies).

## 6. Recommendations before implementation of the practice

- ◆ Peatland restoration should involve and consult all concerned stakeholders, e.g. farmers, landowners, users of waterways, and neighbouring communities.
- ◆ It is necessary to properly plan the restoration activity suited to the local characteristics of sites, including hydrological balance. Proper preparation requires also a verified map of the peat extent and the whole hydrological unit, where the peatland is embedded.
- ◆ Policy makers should explore potential incentives and pricing instruments (i.e. subsidies) that compensate for lost drainage-based productivity. Some schemes may reward practices that transition from conventional peatland use to peatland restoration.

## 7. Potential barriers to adoption

Table 44. Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Restoration may not lead to original pre-drainage peatland conditions, depending on the duration and intensity of peatland degradation.
Cultural	Yes	Peatlands are wetlands that often are difficult to access, and sometimes host insects that may spread malaria, and may be perceived as unpleasant or threatening environments.
Social	Yes	Resistance to restoration is often present as it requires abandonment of conventional drainage-based land-uses.
Economic	Yes	Peatland restoration may not be compatible with drainage-based production practices, resulting in short-term economic losses if no policies are in place to incentivize restoration (see chapter 4.1.3 Paludiculture) and reduced economical viability of the land utilization.
Institutional	Yes	Drainage-based utilization is institutionalized in many countries as the most familiar use, which sometimes also benefits of incentives.
Legal (Right to soil)	Yes	Landowners need to agree to change hydrological conditions of the area, which may stop conventional land-use and thus reduce the lands economical potential.
Knowledge	Yes	Restoration requires peatland-specific knowledge, that is lacking in many regions of the world.

**Table 45.** Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Biomass from reeds as a substitute for peat in energy production in Lida region, Grodno oblast, Belarus</i>	Eurasia	Unknown	6	20
<i>Sphagnum farming for replacing peat as horticultural growing media, Lower Saxony, Germany</i>	Europe	10	6	21

## References

- Carlson, K.M., Goodman, L.K. & May-Tobin, C.C.** 2015. Modeling relationships between water table depth and peat soil carbon loss in Southeast Asian plantations. *Environmental Research Letters*, 10(7): 074006. <https://doi.org/10.1088/1748-9326/10/7/074006>
- Couwenberg, J.** 2018. *Some facts on submerged drains in Dutch peat pastures*. IMCG bulletin June/July 2018, International Mire Conservation Group. (also available at: [http://www.imcg.net/media/2018/imcg\\_bulletin\\_1806.pdf](http://www.imcg.net/media/2018/imcg_bulletin_1806.pdf))
- Dommain, R., Couwenberg, J. & Joosten, H.** 2010. Hydrological self-regulation of domed peatlands in south-east Asia and consequences for conservation and restoration. *Mires & Peat*, 6: 1-17
- FAO.** 2020. *Peatland mapping and monitoring. Recommendations and technical overview*. Italy, Rome. (also available at: <http://www.fao.org/3/CA8200EN/CA8200EN.pdf>)
- FAO.** 2014. *Towards climate-responsible peatland management*. 100 pp. Italy, Rome. (also available at: <http://www.fao.org/3/a-i4029e.pdf>)
- Giesen, W. & Nirmala Sari, E.N.** 2018. *Tropical Peatland Restoration Report: The Indonesian Case. Berbak Green Prosperity Partnership*. Millenium Challenge Account Indonesia, Euroconsult Mott MacDonald, Jakarta.
- Günther, A., Barthelmes, A., Huth, V., Joosten, H., Jurasinski, G., Koebisch, F. & Couwenberg, J.** 2020. Prompt rewetting of drained peatlands reduces climate warming despite methane emissions. *Nature Communications*, 11(1): 1644. <https://doi.org/10.1038/s41467-020-15499-z>
- Hooijer, A., Vernimmen, R., Visser, M. & Mawdsley, N.** 2015. Flooding projections from elevation and subsidence models for oil palm plantations in the Rajang Delta peatlands, Sarawak, Malaysia. Deltares report 1207384, 76pp. (also available at: [https://www.preventionweb.net/files/45060\\_45060rajangdeltapeatlandsubsidencecf.pdf](https://www.preventionweb.net/files/45060_45060rajangdeltapeatlandsubsidencecf.pdf))
- IPCC.** 2014. 2013 *Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands*. Hiraishi, T., Krug, T., Tanabe, K., Srivastava, N., Jamsranjav, B., Fukuda, M. & Troxler, T. (Eds.) Switzerland.
- Jauhainen, J., Page, S. & Vasander, H.** 2016. Greenhouse gas dynamics in degraded and restored tropical peatlands. *Mires & Peat*, 17: 1-12.
- Joosten, H., Tanneberger, F. & Moen, A.** 2017. *The mire and peatlands of Europe*. Schweizerbart Science Publishers, Stuttgart, 780 pages
- Leifeld, J. & Menichetti, L.** 2018. The underappreciated potential of peatlands in global climate change mitigation strategies. *Nature Communications*, 9(1): 1071. <https://doi.org/10.1038/s41467-018-03406-6>
- Page, S.E., Wüst, R. & Banks, C.** 2010. Past and present carbon accumulation and loss in Southeast Asian peatlands. *PAGES news*, 18: 25-27. (also available at: [http://pages-140.unibe.ch/download/docs/newsletter/2010-1/Special%20Section/Page\\_2010-1\(25-27\).pdf](http://pages-140.unibe.ch/download/docs/newsletter/2010-1/Special%20Section/Page_2010-1(25-27).pdf))

- Page, S.E. & Hooijer, A.** 2016. In the line of fire: the peatlands of Southeast Asia. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371. <https://doi.org/10.1098/rstb.2015.0176>
- Shantz, M.A., & Price, J.S.** 2006. Characterization of surface storage and runoff patterns following peatland restoration, Quebec, Canada. *Hydrological Processes*, 20(18): 3799–3814. <https://doi.org/10.1002/hyp.6140>
- Succow, M. & Joosten, H.** 2001. *Landschaftsökologische Moorkunde [Landscape ecology of mires]*. Schweizerbart'sche, p. 622.
- Swenson, M.M., Regan, S., Bremmers, D.T.H., Lawless, J., Saunders, M. & Gill, L.W.** 2019. Carbon Balance of a restored and cutover raised bog: implications for restoration and comparison to global trends. *Biogeosciences*, 16: 713-731. <https://doi.org/10.5194/bg-16-713-2019>
- Tannenberger, F., Appulo, L., Ewert, S., Lakner, S., O’Brolcháin, N., Peters, J. & Wichtmann, W.** 2020. The Power of Nature-based Solutions: How Peatlands Can Help Us to Achieve Key EU Sustainability Objectives. *Advanced Sustainability Systems*. <https://doi.org/10.1002/adsu.202000146>
- Turetsky, M.R., Benscoter, B., Page, S., Rein, G., Van Der Werf, G.R. & Watts, A.** 2015. Global vulnerability of peatlands to fire and carbon loss. *Nature Geoscience*, 8(1): 11–14. <https://doi.org/10.1038/ngeo2325>
- Vroom, R. J. E., Xie, F., Geurts, J. J. M., Chojnowska, A., Smolders, A. J. P., Lamers, L. P. M., & Fritz, C.** 2018. Typha latifolia paludiculture effectively improves water quality and reduces greenhouse gas emissions in rewetted peatlands. *Ecological Engineering*, 124: 88–98. <https://doi.org/10.1016/j.ecoleng.2018.09.008>
- Wakhid, N., Hirano, T., Okimoto, Y., Nurzakiah, S. & Nursyamsi, D.** 2017. Soil carbon dioxide emissions from a rubber plantation on tropical peat. *Science of the Total Environment*, 581–582: 857–865. <https://doi.org/10.1016/j.scitotenv.2017.01.035>
- Wallage, Z.E. & Holden, J.** 2011. Near-surface macropore flow and saturated hydraulic conductivity in drained and restored blanket peatlands. *Soil Use and Management*, 27(2): 247–254. <https://doi.org/10.1111/j.1475-2743.2011.00336.x>
- Wilson, D., Blain, D., Couwenberg, J., Evans, C.D., Murdiyarso, D., Page, S.E., Renou-Wilson, F., Rieley, J.O., Sirin, A., Strack, M. & Tuittila E.-S.** 2016. Greenhouse gas emission factors associated with rewetting of organic soils. *Mires & Peat*, 17: 1-28. <https://doi.org/10.19189/MaP.2016.OMB.222>
- Wichtmann, W., Schröder, C. & Joosten, H.** 2016. *Paludiculture - productive use of wet peatlands. Climate protection, biodiversity, regional economic benefits*. Stuttgart, Schweizerbart Science Publishers, 272
- Wösten, H., Hooijer, A., Siderius C., Satriadi Rais, D., Idris, A. & Rieley, J.** 2006. Tropical Peatland water management modelling of Air Hitam Laut catchment in Indonesia. *International Journal of river basin Management*, 4: 233-244. <https://doi.org/10.1080/15715124.2006.9635293>
- Zerbe, S., Steffenhagen, P., Parakenings, K., Timmermann, T., Frick, A., Gelbrecht, J. & Zak, D.** 2013. Ecosystem Service Restoration after 10 Years of Rewetting Peatlands in NE Germany. *Environmental Management*, 51: 1194-1209. <https://doi.org/10.1007/s00267-013-0048-2>



## 13. Paludiculture

Felix Beer<sup>1</sup>, Wendelin Wichtmann<sup>1</sup>, Laura Villegas<sup>2</sup>, Fahmuddin Agus<sup>3</sup>

<sup>1</sup>University of Greifswald, partner in the Greifswald Mire Centre, Germany

<sup>2</sup>Food and Agriculture Organization of the United Nations, Rome, Italy

<sup>3</sup>Indonesian Soil Research Institute, Bogor, Indonesia

### 1. Description of the practice

Paludiculture produces biomass from wet or rewetted peatlands under conditions that maintain the peat integrity, facilitating peat accumulation and ensuring the provision of peatland ecosystem services (FAO, 2014). Spontaneous vegetation or artificially establishing crops on rewetted sites (see chapter 4.1.2 on Peatland Restoration) require adapted machinery for harvesting. Aside from producing traditional agricultural products such as food, feed, fiber and fuel, the biomass can be used as a raw material for industrial biochemistry, for production of construction materials, substrate for horticulture, high-quality liquid or gaseous biofuels, extraction and synthesizing of pharmaceuticals and cosmetics, among other purposes.

Under paludiculture, water tables need to be established at or close to the surface throughout the year (Jauhiainen *et al.*, 2016; Tanneberger *et al.*, 2020; Wichtmann, Schröder and Joosten, 2016). Paludiculture is currently considered the most efficient solution to balance restored peatland use for livelihood purposes and to increase or to allow SOC maintenance and sequestration in the long term.

When considering wet peatland use, management methods must be well adapted to site-specific conditions. Peatland type, natural vegetation, soil conditions, such as nutrient availability and acidity and hydrology are some factors that need to be considered (Wichtmann, Schröder and Joosten, 2016). In temperate zones, herbaceous *Sphagnum* sp., *Drosera* sp., *Typha latifolia*, *Phragmites australis*, *Alnus glutinosa*, are some of the potential paludiculture species (Abel, Couwenberg and Joosten, 2012). In Southeast Asia, the palm *Metroxylon sagu* and trees *Dyera polyphylla*, *Shorea* ssp., are amongst a whole range of known species used for food, timber and other non-timber products (Van der Meer and Karyanto, 2013; Giesen and Nirmala, 2018). Species that are not able to cope with permanent wet soil conditions with high water tables, cannot be considered for paludiculture purposes (Giesen and Nirmala, 2018; FAO, 2014; Tata, 2019).

## 2. Range of applicability

The need for a shift from drainage to wet-based peatland utilization is being promoted internationally (e.g. Crump, 2017; FAO, 2014), and it is urgent due the recognition of the severe impacts of drainage-based peatland use on climate, biodiversity loss (Dohong, Aziz and Dargusch, 2017), sustainable livelihoods, and other important ecosystem services they provide (see chapter 4.1.1 and 4.1.2). Paludiculture approaches and techniques with different plant species, and for a range of purposes are being explored in temperate regions (Wichtmann, Schröder and Joosten, 2016) as well as in tropical Southeast Asia (Giesen and Nirmala, 2018).

## 3. Impact on soil organic carbon stocks

Paludiculture can reduce the loss of SOC as GHG emissions (Figure 7) and/or dissolved organic carbon (DOC), and in the long-term the carbon sequestration function will potentially be restored. The magnitude of the effect of paludiculture<sup>5</sup> practices on the carbon sequestration potential and the temporal scale needed are not well understood and further long-term research is needed.




Figure 7. Relationship between global warming potential of greenhouse gas emissions, mean annual ground water table and land-use practices for temperate peatlands

Source: adapted from Tanneberger *et al.*, 2020

<sup>5</sup>Also see factsheet n° 13 on Paludiculture, this volume.

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

See factsheet n° 12 on Peatland Restoration, this volume.

### 4.2 Minimization of threats to soil functions

Table 46. Soil threats

Soil threats	
<b>Soil erosion</b>	Surface lowering (land loss or subsidence) is prevented by avoiding drainage (Joosten, 2015). Wind erosion which is a severe problem of ploughed, dry organic soils is eliminated (Succow and Joosten, 2001).
<b>Nutrient imbalance and cycles</b>	Intact peatlands retain nutrients and help purify surface waters (Succow and Joosten, 2001).
<b>Soil salinization and alkalization</b>	Coastal peat swamps act as a buffer between salt- and freshwater systems, preventing saline intrusion into coastal lands (Silvius, Joosten and Opdam, 2008).
<b>Soil contamination / pollution</b>	Intact peatlands retain pollutants and help purify surface water.
<b>Soil acidification</b>	Contingent on peatland type: ombrotrophic peatland are already acidic (Succow and Joosten, 2001).
<b>Soil compaction</b>	Compaction caused by soil decomposition and mineralization is halted (Schröder <i>et al.</i> , 2015).
<b>Soil water management</b>	Intact peatlands with wetland vegetation self-regulate water run-off and release, thus having a buffer function (Dommain, Couwenberg and Joosten, 2010).

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Biomass production on rewetted peatlands can be equal or even exceeding that of drainage-based agriculture and other biomass production (Wichtmann, Schröder and Joosten, 2016). Bioenergy production from paludiculture can replace fossil fuels (Wichtmann *et al.*, 2014; Wichtmann *et al.*, 2019).

### 4.4 Mitigation of and adaptation to climate change

Paludiculture is the only climate-neutral, SOC preserving land-use option for peatland use. Transforming drainage-based peatland-use such as oil palm plantations into paludiculture (e.g. sago or illipe nut) cultivation can reduce anthropic greenhouse gas emissions between 70 to 117 tonnes CO<sub>2</sub>eq per ha per year with CO<sub>2</sub> responsible for approximately 60 percent and N<sub>2</sub>O responsible for approximately 40 percent of the emissions (Cooper *et al.*, 2020; IPCC, 2014). In Central Europe, reported anthropic GHG emission reduction potential is 20–60 tonnes CO<sub>2</sub>eq per hectare per year (Couwenberg *et al.*, 2011; Tanneberger *et al.*, 2020; Figure 7). Furthermore, paludiculture has the potential to reduce CH<sub>4</sub> emissions – which increase after peatland rewetting – through improved topsoil properties by wetland plant root chemistry (Vroom *et al.*, 2018). In the tropics however, methane dynamics in drained and rewetted areas are yet poorly understood (Sakabe *et al.*, 2018; Deshmukh *et al.*, 2020). If biomass from paludiculture is used to replace fossil fuel energy production, the climate change mitigation potential is further increased (Wichtmann, Couwenberg and Kowatsch, 2009). Climate change adaptation benefits include a net cooling effect on regional climate by keeping water in the landscape, thus playing an important role in water availability for communities, in arid regions in particular. Furthermore, benefits for disaster risk reduction include inhibited future flooding risks by preventing subsidence (Hooijer *et al.*, 2015) and reducing fire risks (Page and Hooijer, 2016).

### 4.5 Socio-economic benefits

Paludiculture provides long-term sustainable livelihoods for people, by halting subsidence and thus avoiding loss of riverine and coastal land through subsequent flooding (see “Hotspots: Peatlands”), by preventing hazardous environments as well as substantial economic losses due to fires and haze (Page and Hooijer, 2016). Paludiculture also has the potential to offer viable income sources from production on wet peatlands, particularly when viable value chains are established (Giesen and Nirmala, 2018; Wichtmann, Schröder and Joosten, 2016; Uda, 2019).

### 4.6 Additional benefits to the practice

Further benefits include nutrient retention, water regulation and purification (FAO, 2014; Walton *et al.*, 2020), and maintenance and improvement of biodiversity (Wichtmann, Schröder and Joosten, 2016).

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

None are known.

### 5.2 Increases in greenhouse gas emissions

See factsheet n° 12 of this volume and Peatlands chapter of volume 2.

### 5.3 Conflict with other practice(s)

Raising the water table can lead to land use conflicts with conventional land-use practices such as grasslands in Europe or oil palm plantations in Southeast Asia, causing negative short-term economic effects (Wichtmann, Schröder and Joosten, 2016). Furthermore, land-use conflicts between paludiculture and conservation may occur after peatland restoration. Adaptive guidelines and spatial land-use planning is required to guide and manage transition from traditional drainage-based land-use towards sustainable paludiculture on peatlands (Joosten *et al.*, 2015; Tanneberger *et al.*, 2018).

### 5.4 Decreases in production (e.g. food/fuel/feed/timber)

At present, paludiculture is not as economically viable as drainage-based agriculture. Moreover, it cannot compete with food production in terms of productivity (Leifeld and Menichetti, 2018). That said, conventional cost benefit analysis needs to account for the long-term, highly negative impacts of drainage-based production systems.

### 5.5 Other conflicts

In Central Kalimantan, Indonesia, the establishment of paludiculture practices has encountered several challenges, such as low local community participation (Syahaza, Bakce and Irianti, 2019). The concept of wet agriculture or forestry in peatlands is often a novel concept to local communities and requires guidance and education for general acceptance and successful implementation. There are, however, also traditional, wet production systems that are non-disruptive to peatland ecosystems, such as traditional fishing using 'beje', or collection of rattan - both within the limits of natural regeneration.

## 6. Recommendations before implementation of the practice

- ◆ Study available cases of paludiculture (e.g. FAO, 2014; Giesen and Nirmala, 2018; Wichtmann, Schröder and Joosten, 2016) and connect with the community or practitioners,
- ◆ Prepare the land for wet cultivation, including restoration of high water tables,
- ◆ Assessing suitable plant species in participatory processes with local communities, according to climate, peat type, hydrology and regional market conditions,
- ◆ Establish or strengthen value chains and develop a market for paludiculture products,
- ◆ Establish (plant) nurseries to ensure sufficient plant provision,
- ◆ Depending on the production system, investment into new machinery suitable for soft and wet peat soils may be needed (Schröder *et al.*, 2015); and
- ◆ Promote the simplification of spatial planning processes to speed up rewetting procedures and the establishment of paludiculture.

## 7. Potential barriers to adoption

Table 47. Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Access to rewetted peat soils is reduced and requires adapted machinery (Wichtmann, Schröder and Joosten, 2016).
Cultural	Yes	Practitioners and consumers' unfamiliarity with wetland products.
Social	Yes	Paludiculture may reduce the number of workers per ha (e.g. if biomass for energy is produced instead of meat) but this could also be increased if more labour-intensive management of paludiculture sites and/or products (e.g. sago) requires new processing chains, which results in increased added value.
Economic	Yes	Limited paludiculture establishment, market development and participation in Payment for Ecosystem Services mechanisms reduces competitiveness against drainage-based production systems.

Barrier	YES/NO	
Institutional	Yes	Slow planning processes may hinder the implementation (Peters <i>et al.</i> , 2020).
Legal (Right to soil)	Yes	In the EU, wetland plants are not yet considered agricultural crops - nor are there agricultural subsidies available for paludiculture (Peters <i>et al.</i> , 2020). Political frameworks that support the development of paludiculture are not known from any part of the world.
Knowledge	Yes	Worldwide, productive use of wet peatlands is a nascent field of peatland management. Hence, knowledge about paludiculture and its potential for ecosystem services is still limited. Finding profitable commodities suitable for rewetted peatland cultivation that can compete with conventional commodities is the research challenge. Salient solutions would allow for more voluntary implementation of paludiculture systems.

## Photo of the practice



Photo 16. Peatland Restoration and reforestation site in Sumatra Kayuagung, South Sumatra

Table 48. Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Biomass from reeds as a substitute for peat in energy production in Lida region, Grodno oblast, Belarus</i>	Eurasia	Unknown	6	20
<i>Sphagnum farming for replacing peat as horticultural growing media, Lower Saxony, Germany</i>	Europe	10	6	21



## References

- Abel, S., Couwenberg, J. & Joosten, H. 2012. Towards More Diversity in Paludiculture – A Literature Review of Useful Wetland Plants. International Peat Congress, 14(299). <https://doi.org/10.13140/2.1.1590.4963>
- Cooper, H.V., Evers, S., Aplin, P., Crout, N., Dahalan, M.P.B. & Sjogersten, S. 2020. Greenhouse gas emissions resulting from conversion of peat swamp forest to oil palm plantation. *Nature Communications*, 11(1): 407. <https://doi.org/10.1038/s41467-020-14298-w>
- Couwenberg, J., Thiele, A., Tanneberger, F., Augustin, J., Bärtsch, S., Dubovik, D., Liashchynskaya, N., Michaelis, D., Minke, M., Skuratovich, A. & Joosten, H. 2011. Assessing greenhouse gas emissions from peatlands using vegetation as a proxy. *Hydrobiologia*, 674(1): 67–89. <https://doi.org/10.1007/s10750-011-0729-x>
- Crump, J (Ed.). 2017. *Smoke on Water – Countering Global Threats From Peatland Loss and Degradation*. A UNEP Rapid Response Assessment. United Nations Environment Programme and GRID-Arendal, Nairobi and Arendal. (also available at: [https://gridarendal-website-live.s3.amazonaws.com/production/documents/:s\\_document/376/original/RRapeatland\\_revised\\_jan.pdf?1515398975](https://gridarendal-website-live.s3.amazonaws.com/production/documents/:s_document/376/original/RRapeatland_revised_jan.pdf?1515398975))
- Deshmukh, C.S., Julius, D., Evans, C.D., Nardi, Susanto, A.P., Page, S.E., Gauci, V., Laurén, A., Sabiham, S., Agus, F., Asyhari, A., Kurnianto, S., Suardiwerianto, Y. & Desai, A.R. 2020. Impact of forest plantation on methane emissions from tropical peatland. *Global Change Biology*, 1–19. <https://doi.org/10.1111/gcb.15019>
- Dohong, A., Aziz, A.A. & Dargusch, P. 2017. A review of the drivers of tropical peatland degradation in South-East Asia. *Land Use Policy*, 69: 349–360. <https://doi.org/10.1016/j.landusepol.2017.09.035>
- Dommain, R., Couwenberg, J. & Joosten, H. 2010. Hydrological self-regulation of domed peatlands in south-east Asia and consequences for conservation and restoration. *Mires & Peat*, 6: 1-17
- FAO. 2014. Towards climate-responsible peatland management. Rome, Italy. (also available at: <http://www.fao.org/3/a-i4029e.pdf>)
- Giesen, W. & Nirmala Sari, E.N. 2018. *Tropical Peatland Restoration Report: The Indonesian Case*. Berbak Green Prosperity Partnership. Millenium Challenge Account – Indonesia, Euroconsult Mott MacDonald, Jakarta. (also available at: <https://luk.staff.ugm.ac.id/rawa/GiesenNirmala2018TropicalPeatlandRestorationReportIndonesiaForBRG.pdf>)
- Hooijer, A., Vernimmen, R., Visser, M. & Mawdsley, N. 2015. *Flooding projections from elevation and subsidence models for oil palm plantations in the Rajang Delta peatlands, Sarawak, Malaysia*. Deltares report 1207384.
- IPCC. 2014. *2013 Supplement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories: Wetlands*. T. Hiraishi, T. Krug, K. Tanabe, N. Srivastava, B. Jamsranjav, M. Fukuda & T. Troxler (Eds.). Switzerland.

- Joosten, H.** 2015. Managing Soil Carbon in Europe. Paludiculture as a new perspective for Peatlands. In Steven A Banwart, Elke Noellemeyer, Eleanor Milne (Eds.) *Soil Carbon: Science, Management and Policy for Multiple Benefits*. CAB International. Oxfordshire, UK
- Leifeld, J. & Menichetti, L.** 2018. The underappreciated potential of peatlands in global climate change mitigation strategies. *Nature Communications*, 9(1): 1071. <https://doi.org/10.1038/s41467-018-03406-6>
- Page, S.E. & Hooijer, A.** 2016. In the line of fire: the peatlands of Southeast Asia. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 371. <https://doi.org/10.1098/rstb.2015.0176>
- Peters, J., Hirschelmann, S., Krüger, S., Hedden-Dunkhorst, H., Salathé, T. & Kopansky, D.** 2020. *Peatland Strategies in Europe*. Greifswald Moor Centrum
- Sakabe, A., Itoh, M., Hirano, T. & Kusin, K.** 2018. Ecosystem-scale methane flux in tropical peat swamp forest in Indonesia. *Global Change Biology*, 1-14. <https://doi.org/10.1111/gcb.14410>
- Schröder, C., Dahms, T., Paulitz, J., Wichtmann, W. & Wichmann, S.** 2015. Towards large-scale paludiculture: addressing the challenges of biomass harvesting in wet and rewetted peatlands. *Mires & Peat*, 17
- Silvius, M.J., Joosten, H. & Opdam, S.** 2008. Peatlands and people. In F. Parish, A. Sirin, D. Charman, H. Joosten, T. Minayeva, M.J. Silvius & L. Stringer (Eds.) *Assessment on Peatlands, Biodiversity and Climate Change: Main Report*. pp. 20–28. Global Environment Centre and Wetlands International. Wageningen, the Netherlands. (also available at: [http://www.imcg.net/media/download\\_gallery/books/assessment\\_peatland.pdf](http://www.imcg.net/media/download_gallery/books/assessment_peatland.pdf))
- Succow, M. & Joosten, H. (Ed).** 2001. Landschaftsökologische Moorkunde [*Landscape ecology of mires*]. Schweizerbart'sche, p. 622
- Syahaza, A., Bakce, D. & Irianti, M.** 2019. Improved Peatlands Potential for Agricultural Purposes to Support Sustainable Development in Bengkalis District, Riau Province, Indonesia. *Journal of Physics: Conference Series*, 1351. <https://doi.org/10.1088/1742-6596/1351/1/012114>
- Tata, H.** 2019. Mixed farming systems on peatlands in Jambi and Central Kalimantan provinces, Indonesia: should they be described as paludiculture? *Mires and Peat*, 25: 1–17.
- Tanneberger, F., Schröder, C., Hohlbein, M., Wichmann, S., Wichtmann, W. & Permien, T.** 2018. Putting paludiculture into practice - How can we avoid land use conflicts? In *Proceedings from the 20th EGU General Assembly*. p.18555. EGU2018, 4-13 April 2018, Vienna, Austria.
- Tannenberger, F., Appulo, L., Ewert, S., Lakner, S., O'Brolcháin, N., Peters, J. & Wichtmann, W.** 2020. The Power of Nature-based Solutions: How Peatlands Can Help Us to Achieve Key EU Sustainability Objectives. *Advanced Sustainability Systems*. <https://doi.org/10.1002/adss.202000146>
- Uda, S.K.** 2019. *Sustainable Peatland Management in Indonesia: Towards better understanding of socio-ecological dynamics in tropical peatland management*. PhD thesis, University of Wageningen. p. 232

- van der Meer, P. & Karyanto, O.** 2013. *Opportunities for sustainable forestry in peatlands of Indonesia. Quick Assessment and Nationwide Screening (QANS) of Peat and Lowland Resources and Action Planning for the Implementation of a National Lowland Strategy.* Report on QANS Component 3.
- Vroom, R.J.E., Xie, F., Geurts, J.J.M., Chojnowska, A., Smolders, A.J.P., Lamers, L.P.M. & Fritz, C.** 2018. *Typha latifolia* paludiculture effectively improves water quality and reduces greenhouse gas emissions in rewetted peatlands. *Ecological Engineering*, 124: 88–98.  
<https://doi.org/10.1016/j.ecoleng.2018.09.008>
- Walton, C.R., Zak, D., Audet, J., Petersen, R.J., Lange, J., Oehmke, C., Wichtmann, W., Kreyling, J., Grygoruk, M., Jabłońska, E., Kotowski, W., Wiśniewska, M.M., Ziegler, R. & Hoffmann, C.C.** 2020. Wetland buffer zones for nitrogen and phosphorus retention: Impacts of soil type, hydrology and vegetation. *Science of the Total Environment*. <https://doi.org/10.1016/j.scitotenv.2020.138709>
- Wichtmann, W., Couwenberg, J. & Kowatsch, A.** 2009. Climate protection by reed cultivation. (in german). *Ökologisches Wirtschaften*, 24.
- Wichtmann, W., Oehmke, C., Bärish, S., Deschan, F., Malashevich, V. & Tanneberger, F.** 2014. Characteristics of Biomass from wet fens in Belarus and their potential to substitute peat briquettes as a fuel. *Mires & Peat*, 13: 10.
- Wichtmann, W., Schröder, C. & Joosten, H.** 2016. *Paludiculture - productive use of wet peatlands. Climate protection, biodiversity, regional economic benefits.* Stuttgart, Schweizerbart Science Publishers, 272 p.
- Wichtmann, W., Bork, L., Dahms, T., Körner, N., Kabengele, G.R., Oehmke, C. Wenzel, M. & Barz, M.** 2019. Das Projekt Bonamoor. Biomasseproduktion und Optimierung auf nassen Moorstandorten und deren thermische Verwertung (in german). In *Proceedings zum 13. Rostocker Bioenergieforum. Schriftenreihe Umweltingenieurwesen*. Band 87. Universität Rostock. S. 135 – 145.

# 14. Restoration of mangrove forest

Beth A. Middleton, Eric J. Ward

U.S. Geological Survey, Wetland and Aquatic Research Center, Lafayette, LA, United States of America

## 1. Description of the practice

Mangrove forests occur worldwide along tropical coasts in inundated soils where primary production and anaerobic conditions contribute to the building of soil organic matter (Also see [Mangroves Hot-spot, Volume 2](#)). Note that peat may accumulate in certain coastal mangrove (Middleton and McKee, 2001). The actual amount of soil organic matter stored in these wetlands depends on the balance between primary production and decomposition processes (Middleton and McKee, 2001; Kolka *et al.*, 2018; Middleton, 2020). The restoration of mangroves can increase carbon stocks both in soil and aboveground biomass (Wickland *et al.*, 2013; Chimner *et al.*, 2017; Friess *et al.*, 2019). While tropical inland peatland forests may have higher carbon sequestration rates than mangrove swamps, methane (CH<sub>4</sub>) emissions are generally lower in mangrove swamp, so that these wetlands have greater carbon sequestration potential (Kolka *et al.*, 2018; Al-Haj and Fulweiler, 2020). Mangroves and forested boreal and temperate peatlands tend to store more carbon than non-forested peatlands (Kolka *et al.*, 2018).

Mangrove forest regeneration will occur naturally if the environment suits their life history requirements and propagules are deposited at appropriate elevations (Lewis, 1994; Middleton, 1999). Furthermore, the overall functional equivalence of restored wetlands to natural wetlands is a matter of debate (Kolka *et al.*, 2018).

In mangroves, if the soils of the restored mangrove swamps were not originally damaged, total carbon stocks of restored and pristine forests are similar. CO<sub>2</sub> sequestration is substantial after restoration (Sharma *et al.*, 2020). Restored mangrove swamps hold an amount of carbon equivalent to undisturbed ones after 25-30 years (e.g. Cambodia; Sharma *et al.*, 2020) so that restoration programs have a high potential for carbon sequestration (Hutchison *et al.*, 2014; Chimner *et al.*, 2017). The success of mangrove restoration also depends on the environmental context, e.g. former aquaculture ponds often have higher success rates than seafront plantings (Friess *et al.*, 2019).

## 2. Range of applicability

Mangroves are restricted primarily to the tropics, with poleward ranges set by freezing-induced damage and mortality (Osland *et al.*, 2017).

## 3. Impact on soil organic carbon stocks

**Table 49.** Changes in soil organic carbon stocks reported for restoration of mangrove forests

Location	Context	Cseq/Additional C storage	More information	Reference
Mangrove; northern Cambodia	Restored ecosystem C storage after 25 years	949.4 ± 64.4 tC/ha	Restored and pristine carbon stocks similar	Sharma <i>et al.</i> (2020)
Worldwide	Mangrove Restoration Potential Map	0.069 Gt carbon in above ground biomass; 0.296 Gt carbon in top m of soil (over an area of 8 120 km <sup>2</sup> of restorable mangroves worldwide)	Potential increase in carbon after restoration	Worthington and Spalding (2018)

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Peat accretion via organic matter from roots (Middleton and McKee, 2001) is an important component of maintaining coastal elevation with sea level rise (McKee *et al.*, 2007). Peat elevation is most likely to be stable if the vegetation is healthy, and collapse can occur if vegetation is damaged (Chambers, Steinmuller and Breithaupt, 2019).

## 4.2 Minimization of threats to soil functions

Table 50. Soil threats

Soil threats	
Soil erosion	Restored mangroves protect from coastal erosion from tsunamis and tropical storms (Alongi, 2012; Hutchison <i>et al.</i> , 2014).
Soil salinization and alkalinization	Restored mangroves in high salinity environments have decreased CO <sub>2</sub> flux and root productivity (Troxler <i>et al.</i> , 2015).
Soil contamination / pollution	Restored mangroves purify water (Hutchison <i>et al.</i> , 2014).
Soil acidification	Restoration reduces acid sulfate soils in aquaculture ponds (Alongi, 2002).
Soil biodiversity loss	Plant-soil-microbial relationships are understudied (Alongi, 2002).
Soil water management	Raising water table for restoration will decrease GHG loss by slowing decomposition of soil organic matter (Kolka <i>et al.</i> , 2018).

## 4.3 Increases in production (e.g. food/fuel/feed/timber)

Mangrove restoration increase production of materials from natural forests, as opposed to conventional agriculture and silviculture (Middleton, 1999). Mangroves are invaluable for supporting both primary and secondary production related to biodiversity and wildlife (Alongi, 2002).

## 4.4 Mitigation of and adaptation to climate change

Mangrove swamps have high productivity and contribute to coastal geomorphology and sediment deposition (Barbier *et al.*, 2011) through peat accumulation via roots (Middleton and McKee, 2001; McKee *et al.*, 2007).

## 4.5 Socio-economic benefits

Carbon sink function is maximized and atmospheric GHG concentrations are reduced, which minimize global shifts in climate temperature and precipitation (Moomaw *et al.*, 2018).

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

Table 51. Soil threats

Soil threats	
Soil erosion	Aquaculture and other practices less useful in coastal protection (Alongi, 2002).
Soil contamination/ pollution	Shrimp farming results in nitrogen eutrophication of the pond (Burford and Longmore, 2001).
Soil acidification	Pond aquaculture can lead to acid sulfate soils (Alongi, 2002).
Soil biodiversity loss	Seed banks of natural swamp lost during farming (Middleton, 1999).
Soil water management	Lowering water table for farming or forestry will increase CO <sub>2</sub> loss through decomposition of soil organic matter (Kolka <i>et al.</i> , 2018).

### 5.2 Increases in greenhouse gas emissions

The rehabilitation of mangrove forests ultimately reduces net CO<sub>2</sub> emissions (Wickland *et al.*, 2013; Cameron *et al.*, 2019). Freshwater wetland ecosystems are a noted source of methane (CH<sub>4</sub>) emissions worldwide, though uncertainty is high given the rarity of long-term, continuous flux data, especially in tropical regions (Knox *et al.*, 2019; Zhang *et al.*, 2017), with emissions generally decreasing with salinity in coastal ecosystems (Poffenbarger, Needelman and Megonigal, 2011), including mangroves (Al-Haj and Fulweiler, 2020). Thus, CH<sub>4</sub> emission rates are often lower in mangroves than inland peatland forests, which, coupled with lower restoration costs, makes mangrove restoration generally a more cost-effective natural carbon sequestration method (Taillardat *et al.*, 2020).

### 5.3 Conflict with other practice(s)

Restoration conflicts with agriculture (rice, oil palm), infrastructure development (Friess *et al.*, 2019), and aquaculture ponds, which are productive for only 3-10 years (Cameron *et al.*, 2019).

## 5.4 Decreases in production (e.g. food/fuel/feed/timber)

Restoration negatively impacts production from agriculture and aquaculture (Cameron *et al.*, 2019) and other extractive activities (e.g., peat mining, logging). At the same time restoration provides opportunities for the development of useful products from the natural swamp e.g. food and medicine (Alongi, 2012).

## 6. Recommendations before implementation of the practice

Secure buy-in from local stakeholders through their participation in the restoration activities, as well as their long-term usage of restored mangrove resources.

## 7. Potential barriers to adoption

**Table 52.** Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Environment may have shifted in the region, and physical conditions at the site may no longer be suitable for the desired target vegetation type (Middleton, 1999).
Cultural	Yes	Stakeholders who have used the land previously may object to a change in the land-use.
Social	Yes	See “Cultural Barrier” above.
Economic	Yes	Management of freshwater for coastal wetland restoration could potentially flood coastal communities (Meeder <i>et al.</i> , 2018); Harvest from aquaculture ponds supplies fish resources and reduces the chance of overharvesting from wild caught fisheries (Cameron <i>et al.</i> , 2019). Economic loss from land uses such as agriculture and other infrastructure.
Institutional	Yes	Legal barriers to restoration. Land tenure issues may direct restoration efforts to sub-optimal locations (Friess <i>et al.</i> , 2019).



Barrier	YES/NO	
Legal (Right to soil)		Securing land for the rehabilitation of mangroves may be difficult when the target property spans several human communities under various legal and land tenure restrictions (Cameron <i>et al.</i> , 2019).
Knowledge	No	Restoration approaches are generally well known (Middleton, 1999).
Other	Yes	Rewetting of former agricultural land results in net annual removals of CO <sub>2</sub> in the majority of organic soil classes and a lowering of net GHG emissions (Wilson <i>et al.</i> , 2016).

**Table 53.** Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Mangrove restoration in abandoned ponds in Bali, Indonesia</i>	Asia	10	6	17

## References

- Al-Haj, A.N. & Fulweiler, R.W. 2020. A synthesis of methane emissions from shallow vegetated coastal ecosystems. *Global Change Biology*, 26(5): 2988–3005. <https://doi.org/10.1111/gcb.15046>
- Alongi, D.M. 2002. Present state and future of the world's mangrove forests. *Environmental Conservation*, 29(3): 331–349. <https://doi.org/10.1017/S0376892902000231>
- Alongi, D.M. 2012. Carbon sequestration in mangrove forests. *Carbon Management*, 3: 313–322.
- Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C. & Silliman, B.R. 2011. The value of estuarine and coastal ecosystem services. *Ecological Monographs*, 81: 169–193. <https://doi.org/10.1890/10-1510.1>
- Burford, M.A. & Longmore, A.R. 2001. High ammonium production from sediments in hypereutrophic shrimp ponds. *Marine Ecology Progress Series*, 224: 187–195.
- Cameron, C., Hutley, L.B., Friess, D.A. & Brown, B. 2019. High greenhouse gas emission mitigation benefits from mangrove rehabilitation in Sulawesi, Indonesia. *Ecosystem Services*, 40: 101035. <https://doi.org/10.1016/j.ecoser.2019.101035>
- Chambers, L.G., Steinmuller, H.E. & Breithaupt, J.L. 2019. Toward a mechanistic understanding of “peat collapse” and its potential contribution to coastal wetland loss. *Ecology*, 100(7): e02720. <https://doi.org/10.1002/ecy.2720>
- Chinner, R. A., Cooper, D. J., Wurster, F. C. & Rochefort, L. 2017. An overview of peatland restoration in North America: where are we after 25 years? *Restoration Ecology*, 25(2): 283–292. <https://doi.org/10.1111/rec.12434>
- Friess, D. A., Rogers, K., Lovelock, C. E., Krauss, K. W., Hamilton, S. E., Lee, S. Y., Lucas, R., Primavera, J., Rajkaran, A. & Shi, S. 2019. The state of the world's mangrove forests: past, present, and future. *Annual Review of Environment and Resources*, 44: 89–115.
- Hutchison, J., Manica, A., Swetnam, R., Balmford A. & Spalding, M. 2014. Predicting global patterns in mangrove forest biomass. *Conservation Letters*, 7: 233–240. <https://doi.org/10.1111/conl.12060>
- Knox, S. H., Jackson, R. B., Poulter, B., McNicol, G., Fluet-Chouinard, E., Zhang, Z., Papale, D., Chu, H., Keenan, T.F., Baldocchi, D., Torn, M.S., Mammarella, I., Trotta, C., Aurela, M., Bohrer, G., Campbell, D.I., Cescatti, A., Chamberlain, S., Chen, J., Chen, W., Dengel, S., Desai, A.R., Euskirchen, E., Friborg, T., Gasbarra, D., Goded, I., Goeckede, M., Heimann, M., Helbig, M., Hirano, T., Hollinger, D.Y., Iwata, H., Kang, M., Klatt, J., Krauss, K.W., Kutzbach, L., Lohila, A., Mitra, B., Morin, T.H., Nilsson, M.B., Niu, S., Noormets, A., Oechel, W.C., Peichl, M., Peltola, O., Reba, M.L., Richardson, A.D., Runkle, B.R.K., Ryu, Y., Sachs, T., Schäfer, K.V.R., Schmid, H.P., Shurpali, N., Sonntag, O., Tang, A.C.I., Ueyama, M., Vargas, R., Vesala, T., Ward, E.J., Windham-Myers, L., Wohlfahrt, G. & Zona, D. 2019. FLUXNET-CH<sub>4</sub> synthesis activity: Objectives, observations, and future directions. *Bulletin of the American Meteorological Society*, 100(12): 2607–2632. <https://doi.org/10.1175/BAMS-D-18-0268.1>

- Kolka, R., Trettin, C., Tang, W., Krauss, K.W., Bansal, S., Drexler, J.Z., Wickland, K.P., Chinner, R.A., Hogan, D.M., Pindilli, E.J., Benscoter, B., Tangen, B., Kane, E.S., Bridgham, S.D. & Richardson, C.J. 2018. Terrestrial wetlands. pp. 507–567. No. 13. (also available at <http://pubs.er.usgs.gov/publication/70201054>).
- Field, C.D. 1999. Mangrove rehabilitation: choice and necessity. In R.S. Dodd (Ed.) *Diversity and Function in Mangrove Ecosystems*. pp. 47–52. Developments in Hydrobiology. Paper presented at, 1999, Dordrecht.
- McKee, K.L., Cahoon, D. & Feller, I.C. 2007. Caribbean mangroves adjust to rising sea level through biotic controls on soil elevation change. *Global Ecology and Biogeography*, 16(5): 545–556. <https://doi.org/10.1111/j.1466-8238.2007.00317.x>
- Middleton, B.A. 1999. *Wetland restoration, flood pulsing and disturbance dynamics*. John Wiley and Sons, New York.
- Middleton, B.A. 2020. Trends of decomposition and soil organic matter stocks in *Taxodium distichum* swamps of the southeastern United States. *PLoS One*, 15(1): e0226998. <https://doi.org/10.1371/journal.pone.0226998>
- Middleton, B.A. & McKee, K.L. 2001. Degradation of mangrove tissues and implications for peat formation in Belizean island forests. *Journal of Ecology*, 89: 818–828. <https://doi.org/10.1046/j.0022-0477.2001.00602.x>
- Meeder, J.F., Ross, M.S., Parkinson, R.W., Castaneda, S. 2020. Enhancing coastal wetland resilience to SLF: just add water? *Solutions*, 9(3). <https://www.thesolutionsjournal.com/article/enhancing-coastal-wetland-resilience-slr-just-add-water/>
- Moomaw, W.R., Chmura, G., Davies, G., Finlayson, M., Middleton, B.A., Nutali, S.M., Perry, J.E., Roulet, N. & Sutton-Grier, A. 2018. Brinson Review: Wetlands in a changing climate: science, policy and management. *Wetlands*, 38: 183–205. <https://doi.org/10.1007/s13157-018-1023-8>
- Osland, M.J., Day, R.H., Hall, C.T., Brumfield, M.D., Dugas, J.L. & Jones, W.R. 2017. Mangrove expansion and contraction at a poleward range limit: climate extremes and land-ocean temperature gradients. *Ecology*, 98(1): 125–137. <https://doi.org/10.1002/ecy.1625>
- Poffenbarger, H.J., Needelman, B.A., Megonigal, J.P. 2011. Salinity influence on methane emissions from tidal marshes. *Wetlands*, 31(5): 831–842. <https://doi.org/10.1007/s13157-011-0197-0>
- Sharma, S., MacKenzie, R.A., Tieng, T., Soben, K., Tulyasuwan, N., Resanond, A., Blate, G. & Litton, C.M. 2020. The impacts of degradation, deforestation and restoration on mangrove ecosystem carbon stock across Cambodia. *Science of the Total Environment*, 706. <https://doi.org/10.1016/j.scitotenv.2019>
- Taillardat, P., Thompson, B.S., Carneau, M., Trottier, K. & Friess, D.A. 2020. Climate change mitigation potential of wetlands and the cost-effectiveness of their restoration. *Interface Focus*, 10(5). <http://dx.doi.org/10.1098/rsfs.2019.0129>
- Troxler, T.G., Barr, J.G., Fuentes, J.D., Engel, V., Anderson, G., Sanchez, C., Lagomasino, D., Price, R. & Davis, S.E. 2015. Component-specific dynamics of river mangrove CO<sub>2</sub> efflux in the Florida coastal

Everglades. *Agricultural and Forest Meteorology*, 213: 273–282.

<https://doi.org/10.1016/j.agrformet.2014.12.012>

**Wickland, K.P., Krusche, A.V., Kolka, R.K., Kishimoto-Mo, A.W., Chimner, R.A., Ogle, S. & Srivastava, N.** 2013. Inland wetland mineral soils. *In 2013 Supplement to the 2006 IPCC guidelines for national greenhouse gas inventories: wetlands*. Intergovernmental Panel on Climate Change, Geneva, Switzerland.

**Wilson, D., Blain, D., Couwenberg, J., Evans, C.D., Murdiyarso, E., Page, S.E., Wilson, Renou-Wilson, Rieley, J.O., Sirin, A., Strack, M. & Tuittila, E.-S.** 2016. Greenhouse gas emission factors associated with rewetting or organic soils. *Mires and Peat*, 17: 1–28.

**Worthington, T. & Spalding, M.** 2018. *Mangrove restoration potential. A global map highlighting a critical opportunity*. (also available at: <https://www.iucn.org/sites/dev/files/content/documents/mangrove-tnc-report-final.31.10.lowspreads.pdf>)

**Zhang, Z., Zimmermann, N. E., Stenke, A., Li, X., Hodson, E. L., Zhu, G., Huang, C. & Poulter, B.** 2017. Emerging role of wetland methane emissions in driving 21<sup>st</sup> century climate change. *Proceedings of the National Academy of Sciences*, 114(36): 9647–9652. <https://doi.org/10.1073/pnas.1618765114>

# 15. Restoration of organic coastal and inland freshwater forests

Beth A. Middleton<sup>1</sup>, Eric Ward<sup>1</sup>, Lorenzo Menichetti<sup>2</sup>

<sup>1</sup>US Geological Survey, Wetland and Aquatic Research Center, Lafayette, United States of America.

<sup>2</sup>Now Ecology, SLU (Sveriges Lantbruksuniversitet), Uppsala, Sweden

**Note:** This document provides a specific perspective with regards to chapter 4.2.1 on peatlands restoration. For a complete overview of the practice, readers may also refer to chapter 4.2.1.

## 1. Description of the practice

Peatland forests occur worldwide in inundated soils where primary production and anaerobic conditions contribute to the building of soil organic matter (Günther *et al.*, 2020). Greenhouse gas emissions (GHG) can be substantial from drained freshwater forests with organic soils. Therefore, rewetting peat via hydrologic restoration (see factsheet n° 12 on Peatland restoration, this volume) can restore the function of these forests as carbon sinks and reduce their emission of certain components of GHG (Wilson *et al.*, 2016). While the drainage of forests with organic soil is often a part of the process of agriculture, forestry, and peat harvesting, drying of peat can contribute to GHG emissions (Wilson *et al.*, 2016; Günther *et al.*, 2020). Reflooding of organic forest soils to restore hydrology can lead to an increase in tree health, production and organic matter accumulation (Middleton, 1999, 2020a), and a considerable overall reduction in CO<sub>2</sub> and N<sub>2</sub>O emissions (Wilson *et al.*, 2016). Depending on the duration and nature of the previous land-use, forested peatland restoration can be successful from seeds remaining in the seed bank or deposited via flood-pulsed dispersal (Middleton 1999, 2000, 2003). It is important to consider the nutrient status, hydrology and salinity of disturbed inland peat soils in peatland forest restoration (Chimner *et al.*, 2017). Furthermore, the overall functional equivalence of restored wetlands to natural wetlands is a matter of debate (Kolka *et al.*, 2018).

## 2. Range of applicability

Peatland forests are widely distributed across continents and exist in at least 180 countries (Parish *et al.*, 2008) in coastal, inland and high mountain settings. These wetlands are highly variable in hydrology and fertility. Forests include riparian, groundwater-fed and precipitation-fed types such as tropical freshwater hardwood and palm forests on floodplains (Dargie *et al.*, 2017), temperate swamp forest (Middleton, 2020a, 2020b), boreal bog forests, temperate bottomland forests and others (Kolka *et al.*, 2018). Though widespread on the landscape, peatlands cover only three percent of the land surface with an overall land area of approximately 4.23 km<sup>2</sup> (Xu *et al.*, 2018).

The rewetting of degraded worldwide peatland forests has a positive effect on GHG mitigation (Kolka *et al.*, 2018) in restored agricultural, forestry and excavated peatland. Natural forested peatlands vs. drained peatlands have lower levels of CO<sub>2</sub> emission, acidity, and compaction (Warren *et al.*, 2016; Silviu, 2014; Runkle and Kutzbach, 2014, respectively). Carbon losses associated with land clearing and cultivation are often very high in these ecosystems and recovery is slow (Warren *et al.*, 2016). Model results predict only about one third of the carbon lost during a single 25-year oil palm rotation is recovered 75 years after the restoration of tropical peatland forests (Warren *et al.*, 2016).

Recovery of forests can be difficult following farming. In freshwater bald cypress (*Taxodium distichum*) swamps of the southeastern United States, bald cypress does not rapidly reestablish in abandoned farmland because of the loss of soil seed banks and microbes during farming, and lack of dispersal in highly fragmented landscapes (Middleton, 1999, 2003). After bald cypress forest removal, recalcitrant cypress wood does not readily build carbon in the soil if forests are slow to reestablish (Middleton, 2020a). These considerations may be the main reasons that certain wet forested soils can be slow to build carbon after disturbance removes the trees.

## 3. Impact on soil organic carbon stocks

The rewetting of forested peatland has the potential of increasing carbon storage in the soil, and above ground plant biomass in tropical, temperate and boreal settings (Warren *et al.*, 2015; Wilson *et al.*, 2016, Table 54). Above ground carbon stocks represented by trees and above ground roots (e.g. knees) can contribute to large amounts of “teal carbon” soil organic matter, especially in anaerobic conditions with tree species with slow decomposing wood (Middleton 2020a, 2020b).

**Table 54.** Potential avoided emissions and changes in soil organic carbon stocks reported for restored degraded peatlands during the duration of study or model run

Location	Context <sup>1</sup>	Avoided emissions and/or C sequestration	Duration	Methodology	More information	Reference
Tropical degraded peatlands	All peatland (forested and not forested) Restored vs. Degraded	Avoided emissions*: 16.7 (0.5–31.5) tC/ha/yr CO <sub>2</sub> eq	Based on 2005–2006 land use	Modeled emissions from global mapping of degraded peatlands	Avoided emissions include CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O and DOC	Leifeld and Meni-chetti (2018)
Temperate degraded peatlands		Avoided emissions*: 4.1 (2.6–5.4) tC/ha/yr CO <sub>2</sub> eq				
Boreal degraded peatlands		Avoided emissions*: 4.6 (2.8–6.3) tC/ha/yr CO <sub>2</sub> eq				
Tropical peat swamp forest; Southeast Asia (region-wide)	Reflooding of oil palm forests to restore swamp forest	Avoided emissions: 45, 110 and 140 tC/ha in dry, moderate and wet climate models	75 years	Modeled CO <sub>2</sub> emissions only	One year of oil palm cultivation offset by 6–20 years of restoration	Warren <i>et al.</i> (2016)
	Eliminating surface fires between oil palm rotations	Avoided emissions: 545.8 tC/ha	100 years		Estimate based on lack of fire only	
Tropical peatland forest; Southeast Asia (region-wide)	Restored swamp after 25-year oil palm rotation	C sequestration rate: 0.6 to 1.7 tC/ha/yr; Avoided emissions: 12 tC/ha/yr	25 years		Biomass not included; model simulation	

Location	Context <sup>1</sup>	Avoided emissions and/or C sequestration	Duration	Methodology	More information	Reference
Tropical peat swamp forest; Southeast Asia (Indonesian Borneo)	Simulated reflooding of degraded peatland forest	Avoided emissions: 2.6 tC/ha/yr CO <sub>2</sub> eq	2004-2006	Model	Model based on chamber measurements of CO <sub>2</sub> and CH <sub>4</sub>	Page <i>et al.</i> (2009)
Tropical peat swamp forest; Malaysia, Brunei Darussalam & Indonesia	Converted oil palm plantation	Avoided emissions: 19–32 tC/ha/yr CO <sub>2</sub> eq	1 year	Field study	Chamber measurements of CO <sub>2</sub> , CH <sub>4</sub> and N <sub>2</sub> O	Cooper <i>et al.</i> (2020)
Boreal forest, organic soil; global	Forest rewetting	Avoided emissions: 0.33-0.42 tC/ha/yr CO <sub>2</sub> eq	Various	Review	Emission avoided, CO <sub>2</sub> , CH <sub>4</sub> , N <sub>2</sub> O & DOC	Wilson <i>et al.</i> (2016)
	Cropland rewetting	Avoided emissions: 9.13 t CO <sub>2</sub> tC/ha/yr CO <sub>2</sub> eq				
Temperate forest, organic soil; global	Forest rewetting; nutrient poor soils <sup>1</sup>	Avoided emissions: 2.31 tC/ha/yr CO <sub>2</sub> eq				
	Cropland rewetting	Avoided emissions: 6.99 tC/ha/yr CO <sub>2</sub> eq				
Tropical forest, organic soil; global		Avoided emissions: 14.97 tC/ha/yr CO <sub>2</sub> eq				

Nutrient “poor” refers to rainwater fed settings with little external input of nutrients. \*Area-weighted mean (Range)

Baseline emissions of these systems are rarely known



## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

After the snowmelt period, runoff on restored black spruce peatland is 25 percent of unrestored sites (Shantz and Price, 2006).

### 4.2 Minimization of threats to soil functions

Table 55. Soil threats

Soil threats	
<b>Soil erosion</b>	Water runoff reduced after restoration (Shantz and Price, 2006).
<b>Soil salinization and alkalinization</b>	Very low levels of soil salinity can cause a reduction in production, survival and regeneration with sea level rise, storm surge and freshwater extraction. Minimizing freshwater over-usage helps prevent vegetation damage (Middleton and Montagna, 2018).
<b>Soil acidification</b>	Reflooding likely reduces the soil acidification that occurs in drained organic forest soils (Silvius, 2014).
<b>Soil biodiversity loss</b>	Improve biodiversity with seed bank transfer, dispersal via flood pulsing or planting (Photo 17; Middleton, 1999)
<b>Soil compaction</b>	Reflooding may decrease compaction (Runkle and Kutzbach, 2014).
<b>Soil water management</b>	Reflooding can reduce regional flooding, particularly on floodplains at least partially managed for nature conservation (Middleton, 1999). Wet peatlands often regulate water flow. Evapotranspiration by trees on tropical floodplains may reduce water (Joosten, Tapio-Biström and Tol, 2012). Above-ground root systems of certain tropical and temperate floodplain trees maintain hydrology (Joosten, Tapio-Biström and Tol, 2012, Mirosław-Swiątek and Amataya, 2017, Middleton, 2020b).
<b>Soil organic matter accretion</b>	Reflooding reduces organic matter decomposition (Wilson <i>et al.</i> , 2016a; Middleton, 2020a).

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Rewetting drained forests with organic soils increases production of trees and associated forest products (Middleton, 1999).

### 4.4 Mitigation of and adaptation to climate change

Reflooding drained forested peatland decreases CO<sub>2</sub> emission, but may increase CH<sub>4</sub> emission (Haddaway *et al.*, 2014), particularly in the years immediately following rewetting (Wilson *et al.*, 2016). It is important to note here that recent eddy covariance studies conclude that natural peatland forests may have higher CH<sub>4</sub> emissions than previously thought (Deshmukh *et al.*, 2020; Wong *et al.*, 2020), possibly because previous chamber studies did not capture the efflux of CH<sub>4</sub> via tree stems (Pangala *et al.*, 2013; Covey and Megonigal, 2019). On the other hand, N<sub>2</sub>O emissions tend to decrease with rewetting (Wilson *et al.*, 2016).

### 4.5 Socio-economic benefits

Carbon sink function is maximized, so that rewetted organic forests can reduce global heating (Moomaw *et al.*, 2018; Günther *et al.*, 2019), and emissions, air quality impacts and risk to property from fire (Turetsky *et al.*, 2015; Warren *et al.*, 2016).

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

Table 56. Soil threats

Soil threats	
Soil biodiversity loss	Suitable plant material may not be available because forested wetland seeds die after drainage for farming (Photo 17; Middleton, 1999).
Soil water management	Reflooding reduces carbon release to soil water and surface water (Wilson <i>et al.</i> , 2016).

## 5.2 Increases in greenhouse gas emissions

Previous land use likely has a large influence on GHG emissions after rewetting. For example, CH<sub>4</sub> emissions following the reflooding of abandoned farmland can be very high (Hendriks *et al.*, 2007; Harpenslager *et al.*, 2015). In contrast, rewetted boreal peatlands after tree removal have much lower levels of CH<sub>4</sub> emission (Tuittila *et al.*, 2000; Waddington, Tóth and Bourbonniere, 2008), but more studies are needed on different land use transitions (Wilson *et al.*, 2016). See 4.4 above regarding trade-offs between CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions.

## 5.3 Conflict with other practice(s)

Raised water levels decrease value for agriculture, forestry and peat extraction (Haddaway *et al.*, 2014), and obstruct canals and ditches sometimes used as waterways for transportation. In particular, adjacent land usage that relies on drainage may be threatened regionally (Joosten, 2014) including the entire peatland and surrounding areas. Only species adapted to high water tables would thrive after thorough rewetting.

## 5.4 Decreases in production (e.g. food/fuel/feed/timber)

Tree species less adapted to flooding will have lower productive capacity after reflooding (Middleton, 1999), with a subsequent decrease in soil organic matter accumulation depending on the decomposition rate or organic matter (Middleton, 2020a). However, the productive capacity of wetland species increases (see e.g., the high productive capacity of mangrove forests that often generate peat soils).

## 5.5 Other conflicts

Soil carbon is a balance of production levels and decomposition rates. Higher primary production occurs in low levels of flooding, even though the lowest rates of decomposition occur in more deeply flooded sites (Middleton, 2020a). While deeper water may be easier to maintain in impoundments, natural water regimes including at least occasional periods of drawdown during the growing season maximizes the production of most wetland tree species (Middleton, 1999, 2020a).

## 6. Recommendations before implementation of the practice

Subsidence in farmed tropical peatland eventually may make farming impossible (e.g. drained peatland in Southeast Asia (Hooijer *et al.*, 2014)), so that restoration may become an attractive land use alternative for carbon mitigation (see factsheet n° 12 on Peatland restoration, this volume). The assumption is that hydrologic restoration restores the function of the abandoned farmland as a carbon sink over time, and that CO<sub>2</sub> removal in rewetted organic soil is higher than in drained soils (Wilson *et al.*, 2016). However, carbon loss from peat during farming can greatly exceed the capacity of this type of peatland to recapture atmospheric carbon after restoration. Models suggest that 75 years after hydrologic restoration, only as little as 1/3 of carbon is recaptured that was released to the atmosphere during 25 years of oil palm cultivation (Warren *et al.*, 2016). For example, over 100 years of cultivation consisting of four oil palm rotations with burning, more carbon was lost from peat than is naturally accumulated by these peatlands over 3000 years (Warren *et al.*, 2016). Furthermore, drained and degraded peatlands that are not hydrologically restored continue to lose soil carbon and have an increased fire risk for many years. This situation is particularly problematic if the abandoned farm site is not actively managed (Turetsky *et al.*, 2015; Warren *et al.*, 2016; FAO, 2012).

## 7. Potential barriers to adoption

Table 57. Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Permanent changes in hydrology e.g. downcutting of channel feeding floodplain (Middleton, 1999).
Cultural	Yes	Farmers, peat miners, and other land managers may object to a change in land use to restored wetland because of traditional, drainage-based usage.
Social	Yes	A change to one part of the watershed may affect the entire region (Joosten, 2014). Stakeholders may view rewetting as irreversible although it would be possible to re-dig ditches or canals, or remove dams or drainage obstacles.
Economic	Yes	Flooding in urban areas (Meeder <i>et al.</i> , 2018). Decreased flooding downstream impacts agriculture (Silvius, 2014).
Institutional	Yes	May require public funding and planning (Joosten, 2014).

Barrier	YES/NO	
Legal (Right to soil)	Yes	Legal barriers to restoration e.g. property flooding; land regulations; traditional land use may not be possible after restoration so that land is no longer as valuable.
Knowledge	Yes	Success of vegetative restoration is elusive (Middleton, 1999). The capacity of rewetted peatlands to act as carbon sinks is not well known for tropical peat swamps (Wilson <i>et al.</i> , 2016).
Other	Yes	Rewetting of former agricultural land can produce high levels of CH <sub>4</sub> emission (Wilson <i>et al.</i> , 2016).

## Photo of the practice



**Photo 17.** Regeneration of vegetation in drained organic forest can occur after short-term farming from seeds in the seed bank, dispersal or replanting (Cache River, Southern Illinois)

## References

- Bussell, J., Jones, D.L., Healey, J.R. & Pullin, A.** 2010. How do draining and re-wetting affect carbon stores and greenhouse gas fluxes in peatland soils? CEE Review 08-012 (SR49). *Collaboration for Environmental Evidence*. (also available at: [www.environmentalevidence.org/SR49.html](http://www.environmentalevidence.org/SR49.html))
- Chimner, R.A., Cooper, D.J., Wurster, F.C. & Rochefort, L.** 2017. An overview of peatland restoration in North America: where are we after 25 years? *Restoration Ecology*, 25(2): 283–292. <https://doi.org/10.1111/rec.12434>
- Cooper, H.V., Evers, S.L., Aplin, P., Crout, N., Dahalan, M.P.B. & Sjogersten, S.** 2020. Greenhouse gas emissions resulting from conversion of peat swamp forest to oil palm plantation. *Nature Communications*, 11: 407. <https://doi.org/10.1038/s41467-020-14298-w>
- Covey, K.R. & Megonigal, J.P.**, 2019. Methane production and emissions in trees and forests. *New Phytologist*, 222(1): 35–51. <https://doi.org/10.1111/nph.15624>
- Dargie, G.C., Lewis, S.L., Lawson, I.T., Mitchard, E.T.A., Page, S.E., Bocko, Y.E. & Ifo, S.A.** 2017. Age, extent and carbon storage of the central Congo Basin peatland complex. *Nature*, 542: 86–90. <https://doi.org/10.1038/nature21048>
- Deshmukh, C.S., Julius, D., Evans, C.D., Susanto, A.P., Page, S.E., Gauci, V., Laurén, A., Subiham, S., Agus, F., Ashyhari, A., Kurnianto, S., Suardiwerianto, Y. & Desai, A.R.** 2020. Impact of forest plantation on methane emissions from tropical peatland. *Global Change Biology*, 26(4): 2477–2495. <https://doi.org/10.1111/gcb.15019>
- Günther, A., Barthelmes, A., Huth, V., Joosten, H., Jurasinski, G., Koebisch, F. & Couwenberg, J.** 2020. Prompt rewetting of drained peatlands reduces climate warming despite methane emissions. *Nature Communications*, 11: 1644. <https://doi.org/10.1038/s41467-020-15499-z>
- Haddaway, N.R., Burden, A., Evans, C.D., Healey, J.R., Jones, D.L., Dalrymple, S.E. & Pullin, A.S.** 2014. Evaluating effects of land management on greenhouse gas fluxes and carbon balances in boreo-temperate lowland peatland systems. *Environmental Evidence*, 3: 5. <https://doi.org/10.1186/2047-2382-3-5>
- Harpenslager, S.F., van den Elzen, E., Kox, M.A.R., Smolders, A.J.P., Ettwig, K.F. & Lamers, L.P.M.** 2015. Rewetting former agricultural peatlands: Topsoil removal as a prerequisite to avoid strong nutrient and greenhouse gas emissions. *Ecological Engineering*, 84: 159–168. <https://doi.org/10.1016/j.ecoleng.2015.08.002>
- Hendriks, D.M.D., van Huissteden, J., Dolman, A.J. & van den Molen, M.K.** 2007 The full greenhouse gas balance of an abandoned peat meadow. *Biogeosciences*, 4: 411–424. <https://doi.org/10.5194/bg-4-411-2007>

- Hooijer, A. Triadi, B., Karyanto, O., Page, S.E., van der Vat & M. Erkens, G.** 2012. Subsidence in drained coastal peatland in SE Asia: implications for sustainability. *In Proceedings of the 14th International Peat Congress*. (also available at: <https://peatlands.org/assets/uploads/2019/06/Hooijer-176.pdf>)
- Joosten, H., Tapio-Biström, M.-L. & Tol, S.** 2012. *Peatlands – guidance for climate change mitigation through conservation, rehabilitation and sustainable use*. Mitigation of Climate Change in Agriculture Series 5. Food and Agriculture Organization of the United Nations (FAO) and Wetlands International, Rome, Italy. (also available at: <http://www.fao.org/docrep/015/an762e/an762e.pdf>)
- Joosten, H.** 2014. Rewetting of drained peatland. *In* Biancalani, R. and Avagyan, A. (Eds.) *Mitigation of climate change in agriculture*, pp. 38–40. Series 9. Food and Agriculture Organization of the United Nations (FAO), Rome, Italy.
- Kolka, R., Trettin, C., Tang, W., Krauss, K., Bansal, S., Drexler, J., Wickland, K., Chimner, R., Hogan, D., Pindilli, E.J., Benscoter, B., Tangen, B., Kane, E., Bridgham, S., Richardson, C.** 2018. Chapter 13. Terrestrial wetlands. *In* N. Cavallaro, G. Shrestha, R. Birdsey, R.G. Mayes, Najjar, S.C. Reed, P. Romero-Lankao & Z. Zhu (Eds.) *Second state of the carbon cycle report (SOCCR2): a sustained assessment report*, pp. 507–546. U.S. Global Change Research Program, Washington, DC, USA. <https://doi.org/10.7930/SOCCR2.2018>.
- Leifeld, J. & Menichetti, L.** 2018. The underappreciated potential of peatlands in global climate change mitigation strategies. *Nature Communications*, 9: 1081. <https://doi.org/10.1038/s41467-018-03406-6>.
- Meeder, J., Ross, M.S., Parkinson, R.W. & Castaneda, S.** 2018. Enhancing coastal wetland resilience to SLR: just add water. *Solutions*, 9(3): <https://www.thesolutionsjournal.com/article/enhancing-coastal-wetland-resilience-slr-just-add-water/>
- Middleton, B.A.** 1999. *Wetland restoration, flood pulsing and disturbance dynamics*. John Wiley and Sons, New York.
- Middleton, B.A.** 2003. Soil seed banks and the potential restoration of forested wetland after farming. *Journal of Applied Ecology*, 40:1025–1034. <https://doi.org/10.1111/j.1365-2664.2003.00866.x>
- Middleton, B.A.** 2020a. Trends of decomposition and soil organic matter stocks in *Taxodium distichum* swamps of the southeastern United States. *PLoS One*, 15(1): e0226998. <https://doi.org/10.1371/journal.pone.0226998>
- Middleton B.A.** 2020b. Carbon stock trends of baldcypress knees along climate gradients of the Mississippi River Alluvial Valley using allometric methods. *Forest Ecology and Management*, 461: 117969. <https://doi.org/10.1016/j.foreco.2020.117969>
- Middleton, B.A. & Montagna, P.** 2018. Turning on the faucet to coastal wetlands. *Solutions* 9(3). (also available at: <https://www.thesolutionsjournal.com/article/turning-faucet-healthy-coast/>)
- Mirosław-Swiątek, D. & Amatya, D.M.** 2017. Effects of cypress knee roughness on flow resistance and discharge estimates of the Turkey Creek watershed. *Annals of Warsaw University of Life Sciences – SGGW. Land Reclamation*, 49: 179–199. <https://doi.org/10.1515/sggw-2017-0015>

- Moomaw, W.R., Chmura, G., Davies, G., Finlayson, M., Middleton, B.A., Nutali, S.M., Perry, J.E., Roulet, N. & Sutton-Grier, A.** 2018. Brinson Review: Wetlands in a changing climate: science, policy and management. *Wetlands*, 38: 183-205. <https://doi.org/10.1007/s13157-018-1023-8>
- Page, S., Hoscilo, A., Wösten, H., Jaujjainen, J., Silviu, M., Rieley, J., Ritzema, H., Tansey, K., Graham, L., Vasander, H. & Limin, S.** 2009. Restoration ecology of lowland tropical peatlands in Southeast Asia: current knowledge and future research directions. *Ecosystems*, 12: 888–905. <https://doi.org/10.1007/s10021-008-9216-2>
- Pangala, S.R., Moore, S., Hornibrook, E.R. & Gauci, V.** 2013. Trees are major conduits for methane egress from tropical forested wetlands. *New Phytologist*, 197(2): 524–531. <https://doi.org/10.1111/nph.12031>
- Parish, F., Sirin, A., Charman, D., Joosten, H., Minaeva, T. & Silviu, M.** eds. 2008. *Assessment on peatlands, biodiversity and climate change*. Kuala Lumpur, Global Environment Centre and Wageningen, Wetlands International. 179 pp.
- Runkle, B.R.K. & Kutzbach, L.** 2014. Peatland characterization. In Biancalani, R. & Avagyan, A. (Eds.) *Mitigation of climate change in agriculture*, pp. 6–11. Series 9. Food and Agriculture Organization of the United Nations (FAO), Rome, Italy.
- Silviu, M.** 2014. Plantation expansion and peatland conversion. In Biancalani, R. & Avagyan, A. (Eds.) *Mitigation of climate change in agriculture*, pp. 53–57. Series 9. Food and Agriculture Organization of the United Nations (FAO), Rome, Italy.
- Tuittila, E.-S., Komulainen, V.-M., Vasander, H., Nykänen, H., Martikainen, P.J. & Laine, J.** 2000. Methane dynamics of a restored cut-away peatland. *Global Change Biology*, 6: 569–581. <https://doi.org/10.1046/j.1365-2486.2000.00341.x>
- Turetsky, M.R., Benschoter, B., Page, S., Rein, G., Van Der Werf, G.R. & Watts, A.** 2015. Global vulnerability of peatlands to fire and carbon loss. *Nature Geoscience*, 8(1): 11–14. <https://doi.org/10.1038/ngeo2325>
- Waddington, J.M., Tóth, K. & Bourbonniere, R.A.** 2008. Dissolved organic carbon export from a cutover and restored peatland. *Hydrological Processes*, 22: 2215–2224. <https://doi.org/10.1002/hyp.6818>
- Warren, M., Frolking, S., Dai, Z., & Kurnianto, S.** 2016. Impacts of land use, restoration, and climate change on tropical peat carbon stocks in the twenty-first century: implications for climate mitigation. *Mitigation and Adaptation Strategies for Global Change*, 22: 1041-1061. <https://doi.org/10.1007/s11027-016-9712-1>
- Wilson, D., Blain, D., Couwenberg, J., Evans, C.D., Murdiyarso, E., Page, S.E., Wilson, Renou-Wilson, F., Rieley, J.O., Sirin, A., Strack, M. & Tuittila, E.-S.** 2016. Greenhouse gas emission factors associated with rewetting of organic soils. *Mires and Peat*, 17(4): 1–28. <https://doi.org/10.19189/MaP.2016.OMB.222>



**Wong, G.X., Hirata, R., Hirano, T., Kiew, F., Aeries, E.B., Musin, K.K., Waili, J.W., San Lo, K. & Melling, L.** 2018. Micrometeorological measurement of methane flux above a tropical peat swamp forest. *Agricultural and Forest Meteorology*, 256: 353–361. <https://doi.org/10.1016/j.agrformet.2018.03.025>

**Xu, J., Morris, P.J., Liu, J. & Holden, J.** 2018. PEATMAP: refining estimates of global peatland distribution based on a meta-analysis. *Catena*, 160: 134–140. <https://doi.org/10.1016/j.catena.2017.09.010>

**Zhu, Y., Purdy, K.J., Eyice, Ö., Shen, L., Harpenslager, S.F., Yvon-Durocher, G., Dumbrell, J.J. & Trimmer, M.** 2020. Disproportionate increase in freshwater methane emissions induced by experimental warming. *Nature Climate Change*, 10: 685–690. <https://doi.org/10.1038/s41558-020-020-0824-y>

# 16. Water level management in rice paddies

Sara Ibáñez Asensio

Plant Production Department, Universitat Politècnica de València, Valencia, Spain

## 1. Description of the practice

Water level (or water sheet) management is a key to allow SOC sequestration and the GHG emissions balance in wetland-cultivated areas. The water sheet level along the soil profile governs the emissions of CH<sub>4</sub>, N<sub>2</sub>O and CO<sub>2</sub>: during the flooding period, the anaerobic soil conditions lead to increased CH<sub>4</sub> emissions (due to the activity of anaerobic soil bacteria) and decreased CO<sub>2</sub> emissions (due to the limited mineralization from soil aerobic bacteria). This also generally leads to increased SOC sequestration rates. On the contrary, during dry periods, the aerobic soil conditions lead to decreased CH<sub>4</sub> emissions and increased CO<sub>2</sub> and N<sub>2</sub>O emissions.

In the case of rice fields, the water sheet position in the soil profile can also determine crop yield. For example, a submergence or high water sheet allows a better weed control and a higher efficiency of fertilizer use. On the other hand, drainage is important to a correct soil-plant-atmosphere exchange of substances such as ammonia-N, hydrogen sulphide and other harmful substances that are produced under reductive conditions during submergence. The global impact of the management practices chosen by rice farmers in each agricultural region is by no means a minor issue, since rice is the major cereal crop for more than half of the world's population and about 750 million tons of rice are grown worldwide each year. Rice fields cover 1.29 million km<sup>2</sup>, just over 10 percent of the world's wetlands (Ramsar, 2018).

Rice fields (paddies) can be cultivated in several water regime conditions:

- ◆ Continuous waterlogging (CW): flooding is permanent, except when soil conditioning, fertilization, and rice harvesting tasks are performed, and;
- ◆ Alternate wetting and drying (AWD): the most commonly used water regime that consists of alternation of flooding with drainage until the soil is dry, or even only damp, and flooding again

(Photo 18, Photo 19 and Photo 20). It aims to control the water sheet height, raising and lowering it to different depths, but always at levels that ensure an adequate humidity for the development of rice plant. Different actions can be undertaken to manage the water table height, such as:

- Mid-season drainage (MSD): waterlogged soil is drained for 30 days just 21 days after transplanting; during this time the soil is dry or damp, and remains flooded the rest of the time;
  - Alternative wetting (AW): waterlogged soil is drained several times throughout the rice cycle, shortening times intervals between drying acts;
  - Intermittent irrigation (II): repetition of drainage and irrigation that decreases the amount of water used and ensures longer aerobic conditions;
  - Controlled irrigation (CI): during the crop cycle the soil remains more dry or damp than waterlogged, i.e. higher deep flooding of 10 cm or more in height of water layer, leaving paddy soil dry at 60–80 percent during the growing season (without flooding after the re-greening of rice seed).
- ◆ Rainfed irrigation with sprinkler (S): soil is not flooded at any time during the rice growing cycle but irrigated to ensure that soil moisture is maintained at the levels required by the crop.

In a context of climate change and increasing water scarcity in many wetlands around the world, the irrigation system used becomes very relevant. The use of sprinklers in rice as it can be a smart water management that contributes effectively to reduce water consumption. Both systems (AWD and S) also reduce costs, another of the great challenges of modern agriculture.

## 2. Range of applicability

Water sheet management can be applied in several climate conditions, from tropical and sub-humid to Mediterranean areas. The key for establishing in a simple way the suitability of the AWD strategy is to take into consideration basic environmental aspects such as the climatic parameters (temperature, rain and potential evapotranspiration) and some soil properties (percolation potential and texture). Based on their combination, the farmer must select the best strategy to drain while maintaining a level that does not reduce crop yield and allow agricultural work. In the Philippines, for example, 96.7 percent of paddy areas are moderately or highly adequate to AWD during the dry season; however during the wet season the abundant rains typical of the monsoon period and a moderate to high clay content leads to excessive water amounts that can be difficult to drain in wide rice areas (Sander *et al.*, 2017). In drier climates (e.g. Mediterranean climate) AWD can be applied successfully when the rice fields conditions are adequate, reducing methane emissions and water consumption without decreasing rice yield. An important factor is that if there is a scarcity of water due to low rainfall, the management regime significantly influences the physicochemical characteristics of the soil and therefore increases the importance of observing their long-term effects (Fangueiro *et al.*, 2017; Meijide *et al.*, 2017).

### 3. Impact on soil organic carbon stocks

Studies in areas with a traditional rotation of rice management show that long waterlogging enhances a slow accumulation of organic carbon in wet soil conditions due to the organic-mineral associations developed during the first 100 years of soil development. A meta-analysis carried out in China associates the highest storage capacity of C with the pedogenic subgroups of Fe-accumulating and Fe-leaching paddy soils in the long term (Pan *et al.*, 2003).

This thesis explains why cultivation in traditional flood conditions shows a greater contribution of organic carbon than non-flooded highland crops. For example, values in eastern China for A horizons range from 17.8 to 30 g/kg C in paddy rice under cultivation for 50 to 2000 years, while they only reach 11 g/kg C in non-paddy soils (Wissing *et al.*, 2014); or 15-16 g/kg C in recent paddy soils (Xiong *et al.*, 2015; Chen *et al.*, 2017). It would appear, then, that in comparison with rainfed cultivation, irrigated rice cultivation in China led to increased SOC storage in paddy soils. Pan *et al.* (2003) reported that paddy cultivation induced total C sequestration to half of China's total annual CO<sub>2</sub> emission in the 1990s.

AWD can have negative effects on carbon sequestration as the time that soil is dry increases, but these can be counteracted by implementing different cultivation techniques (e.g. no-till, biomass incorporation, improved varieties). Table 58 shows SOC values in rice fields cultivated in CW and AWD; during various periods of time, with incorporation of biomass (CB) or rice straw (RS) and without incorporation (conventional tillage, CT). As a general idea, values in the Table 58 indicate:

- ◆ Lower OC accumulation in short duration experiences regardless of water level management (CW vs. AWD) (Huque *et al.*, 2017 *vs* Zhang *et al.*, 2010, Chen *et al.*, 2017 and Xionghu *et al.*, 2011);
- ◆ Importance of time on OC accumulation in relation to the change in water management. Haque *et al.* (2017) reported minimal differences for a one-year duration while Chen *et al.* (2017) showed an evolution of carbon stocks associated with drying cycles in paddy during more than 100 years
- ◆ Positive effect of rice straw incorporation (RS). Zhang *et al.* (2010) and Xionghu *et al.* (2011) showed similar records of stocks even though the number of crop years are very different (several thousand years *vs* 28 years)
- ◆ Positive and similar effect of cover crop biomass (B) incorporation between WC or AWD management (Haque *et al.*, 2017)

**Table 58.** SOC stocks and changes in soil organic carbon stocks reported for water level management

Location (Reference)	Climate (MAR, MAP)	Soil type	Crop and soil management	Water management	SOC stock or concentration	Depth (cm)	Duration	Fertilizer additions (t/ha)
Tai Lake plain, Yangsu, Eastern China (Zhang <i>et al.</i> , 2010, data recalculated)	Subtropical monsoon (1177 mm, 15.7 °C)	Entic Hapludept 39% clay content 24 g C/kg 1 g/cm <sup>3</sup> pH: 6.5	Rice variety: <i>cv. Wuyunjing 7</i>	CW	34.4 tC/ha	0-15	paddy for several thousand years	Inorganic fertilizer + urea
Wnagcheng Hunan Province, Southern China (Xionghui <i>et al.</i> , 2012)	1370 mm, 17°C	Reddish Yellow Paddy Soil pH: 6.6	- Rice variety: <i>Conventional and Hybrid (Xiang 67)</i> ; - Double rice (DR) - Typical local management: rice straw incorporation (RS) and conventional tillage (CT)	CW	33.2 tC/ha Additional C storage: 0.075 tC/ha/yr	NA	28 years	RS: 2 x 2.63 t/ha NPK : <i>Early+late</i> N: 0.15+0.18 t/ha P: 2 x 0.038 t/ha k: 2 x 0.0996 t/ha
Taoyuan County Hunan Province,	Tropical Humid monsoon	Soils derived from		CW	42.9 tC/ha	0-20	paddy for over <u>100 years</u>	

Location (Reference)	Climate (MAR, MAP)	Soil type	Crop and soil management	Water management		SOC stock or concentration	Depth (cm)	Duration	Fertilizer additions (t/ha)
Southern China, Asia (Chen <i>et al.</i> , 2017)	(1450 mm, 16.5 °C)	quaternary red clay		AWD Wet season: wetting and dry cycles Dry season: dryness		35.5 tC/ha		Double rice under conventional tillage weeds during 8 years abandonment	
Jinju, South Democratic People's Republic of Korea (Haque <i>et al.</i> , 2017)	Continental monsoon (1528 mm, 13.5 °C)	Silty loam 20.4 g OC/kg	- Rice variety: <i>v. Japonica</i> - Rice under different cover crop biomass incorporation (CBO, CB1, CB2 and CB3)	5–7 cm water when flooded	CW	<i>CBO: 4.7 gC/kg</i> <i>CB1: 5.99 gC/kg</i> <i>CB2: 7.67 gC/kg</i> <i>CB3: 5.78 gC/kg</i>	NA	1 year	Rates of cover crop biomass incorporation and NPK (t/ha): CBO: 0; CB1: 3; CB2: 6; CB3: 12 NPK: N:0.09; P:0.02; K: 0.048
					AWD (MSD): drain for 30 days, 21 days after transplanting	<i>CBO: 4.59 gC/kg</i> <i>CB1: 5.87 gC/kg</i> <i>CB2: 7.61 gC/kg</i> <i>CB3: 5.72 gC/kg</i>			

CW: continuous waterlogging; AWD: alternate wetting and drying; MSD: midseason drainage; DR: double rice; CT: conventional tillage; RS: rice straw incorporation

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

The main improvement derived from lowering water sheet through drainage and increase the drying periods is that translate into aerobic conditions that improve the oxidative condition of the soil and reduce the redox processes of Fe, improving the permeability of the soil in the next fallow season (Chen *et al.*, 2017; Shiratori *et al.*, 2007).

Mid-season drainage is important to cut-off the supply of ammonia-N and to secure some desirable plant characteristics in relation, for example, to upper leaves, lower inter-node, ear formation, and healthy root growth. In addition, this action removes hydrogen sulphide and other harmful substances, which are produced by microbial action under reductive conditions of submergence (Amin, Rowshon and Aimrun, 2011).

During the time that the water sheet is elevated and paddy rice is flooded, anaerobic conditions increase the total and hot water extractable carbon in soil, decrease bulk density and induce soil aggregates stabilization (Shiratori *et al.*, 2007; Wissing *et al.*, 2014).

### 4.2 Minimization of threats to soil functions

Table 59. Soil threats

Soil threats	
<b>Soil salinization and alkalization</b>	In coastal zones, after flooding, salinization is reduced by dilution effect; in Thailand, for example, water drainage reduced soil salinity by 50 percent (Maeght <i>et al.</i> , 2005).
<b>Soil pollution / contamination</b>	Maintenance of the water sheet below the topsoil reduces absorption of contaminants in soils watered with water enriched or pollution, i.e. arsenic (Spanu <i>et al.</i> , 2012).

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Management of the water sheet can condition rice yield. In Spain, Fanguero *et al.* (2017) found that an increase in drying time leads to a decrease in production that can reach up to 50 percent when CW is replaced by rainfed irrigation and conventional tillage. With abundant rainfall and under NT such as in subtropical Chinese rice fields, the substitution of CW for AWD implies a drop in production of only 10 percent (from around 7.5 to about 6.5 t/ha) (Xu *et al.*, 2015). On the contrary, when NT conditions are used in semi-arid areas of Spain, the change of the irrigation system even implies a slight increase in production.

### 4.4 Mitigation of and adaptation to climate change

In general, implementing AWD management translates into lower emissions as compared to CW practices. Collected data in different climatic conditions corroborate that widen soil dry periods reduces CH<sub>4</sub> and in consequence the Global Warming Potential (GWP) and the intensity of greenhouse gases (Sriphrom *et al.*, 2019; Nelson, 2015). In addition, Shiratori *et al.* (2007) showed that installing subsurface drains beneath poorly drained clayey soil rice fields (0.6–0.8 m below) also aid in reducing in wet season methane emissions up to 70 percent. In Haque *et al.* (2017), biomass incorporation is translated in increased emissions due to mineralization, affecting the GWP (Table 60).



**Table 60.** GHG emissions and global warming potential (GWP) according to water level management CW vs AWD or Sprinkler

Location (Reference)	Climate (MAP, MAT)	Soil type	Duration (Years)	Crop and soil management	Fertilizer additions	Water management	GHG emissions (t/ha/yr)			GWP tCO <sub>2</sub> eq/ha/yr	
							CH <sub>4</sub>	N <sub>2</sub> O	CO <sub>2</sub>		
Po Valley, Italy, (Mejjide <i>et al.</i> , 2017)	Mediterranean (650 mm, 12.3 °C (2009))	Loam to clay loam, Calcic Gleysol	2 years CW (2009) AWD/MSD (2010)	Non cultivated after harvest	N (inorganic fertilizer) 2009: 0.13 t/ha 2010: 0.12 t/ha	CW	0.37	0.0008	NA	11.48	
						Midseason drainage (AWD/MSD)	0.21	0.0011		2.89	
Extremadura region, Spain (Fangueiro <i>et al.</i> , 2017)	Mediterranean (< 480 mm, 16.8 °C)	Loam Hydragic Anthrosol	3	Conventional tillage	N inorganic fertilizer: 0.14 x 2 = 0.28 t/ha	10 cm water when flooded	CW	0.35	0.011	6.7	19.4
				No-till			Sprinkler	0.004	0.008	10.2	12.4
							CW	0.12	0.014	5.3	12.5
							Sprinkler	0.001	0.006	5.8	7.39
Lucknow, India (Tyagi, Kumari and Singh,2010)	Humid Subtropical (1001 mm, 25.2 °C)	Sandy loam	1	Rice variety: <i>v. Somali 4006</i>	NA	CW vs. Midseason drainage (AWD/MSD)  In all cases 4-8 cm water when flooded  Drainage: MSD1: once at vegetative cycle MSD2: 7 days, 70 DAP  AW: 3 days, 21 and 77 DAP	<i>Daily average (kg/ha/day)</i>  CW: 3.5 MSD1: 3.15 MSD2: 2.2 AW: 2.04	NA	NA	CW: 8.15 MSD1: 7.41 MSD2: 5.1 AW: 4.82	

Location (Reference)	Climate (MAP, MAT)	Soil type	Duration (Years)	Crop and soil management	Fertilizer additions	Water management	GHG emissions (t/ha/yr)			GWP tCO <sub>2</sub> eq/ha/yr	
							CH <sub>4</sub>	N <sub>2</sub> O	CO <sub>2</sub>		
Hubei, lowland Central China  (Xu <i>et al.</i> , 2015)	Subtropical monsoon  (1500 mm, 17.5 °C)	Silty clay loam	1 (drought year in study area)	No-till  Rice varieties: <i>v. HY3</i> (Drought- resistant) and <i>v.</i> <i>HY299</i> (Typical)	NA	CW	<i>v. HY3</i> : 0.96 <i>v. HY299</i> : 0.92	NA	<i>v. HY3</i> : 9.25 <i>v. HY299</i> : 7.45	<i>v. HY3</i> : 35.55 <i>v. HY299</i> : 32.26	
						Alternative wetting (AWD/AW)  Drainage: AW1: soil always damp  AW2: let soil dry	AW1 <i>v. HY3</i> : 0.37 <i>v. HY299</i> : 0.38  AW2: <i>v. HY3</i> : 0.18 <i>v. HY299</i> : 0.15		AW1: <i>v. HY3</i> : 12.14 <i>v. HY299</i> : 15.44  AW2: <i>v. HY3</i> : 18.05 <i>v. HY299</i> : 17.83	AW1 <i>v. HY3</i> : 23.99 <i>v. HY299</i> : 27.99  AW2 <i>v. HY3</i> : 25.50 <i>v. HY299</i> : 24.49	
Jinju, South Democratic People's Republic of Korea  (Haque <i>et al.</i> 2017)	Continental monsoon  (1 528 mm, 13.5 °C)	Silty loam	1	Rice variety : <i>v.</i> <i>Japonica</i>	N : 0.09 t/ha P : 0.02 t/ha K : 0.048 t/ha  Cover crop biomass incorporation (t/ha): CB0: 0 CB1: 3 CB2: 6 CB4: 12	5–7 cm water when flooded	CW	NA	NA	CB0: -1 CB1: 5 CB2: 6 CB4: 12	CB0: 5 CB1: 8 CB2: 23 CB4: 40
							Midseason drainage (MSD/AWD)  Drainage for 30 days, 21 DAP			CB0: -1.8 CB1: 2 CB2: 5 CB4: 11	CB0: 4 CB1: 6 CB2: 16 CB4: 21

CW: continuous waterlogging; AWD: alternate wetting and drying; MSD: midseason drainage; AW: alternative wetting; DAP: days after planting; CB: cover biomass incorporation

## 4.5 Socio-economic benefits

On one hand, an efficient water sheet management is a good measure for saving resources in areas with water shortages where it is necessary to purify it, or extract from wells, and therefore the economic cost is higher. For example, for wetter weather conditions as in the Philippines, the generalized application of AWD would save 30 percent of water (Sander *et al.*, 2017; Nelson *et al.*, 2015). On other hand, fertilizer consumption can be reduced because the midseason drainage of the otherwise flooded field can outcompete the effect of organic fertilizer application regarding CH<sub>4</sub>. In addition, in most cases implies lower evapotranspiration rate (Meijide *et al.*, 2017).

## 4.6 Additional benefits to the practice

Studies of water level fluctuations in rice fields show that increases in anaerobiosis time can favors the growth of other plant species. An example is explained by Nishio *et al.*, (2006) who established that the fluctuations of water level in flood plains and the artificial management of water level in the rivers around rice paddies influenced the spatial distribution of *Ud. nipponensis*.

# 5. Potential drawbacks to the practice

## 5.1 Tradeoffs with other threats to soil functions

Table 61. Soil threats

Soil threats	
<b>Soil salinization and alkalization</b>	In sensible coastal wetlands, shortening flood periods or decreasing the height of the sheet of water above the ground can exacerbate soil salinization due to marine intrusion (Moreno-Ramón <i>et al.</i> , 2015).
<b>Soil compaction</b>	In aerobic conditions, the faster decomposition of OM tends to limit soil aggregates stabilization (Wissing <i>et al.</i> , 2014).
<b>Soil biodiversity loss</b>	With minimum water use the soil condition are very different and changes in biota and weed can be important (Nielsen <i>et al.</i> , 2009).

## 5.2 Increases in greenhouse gas emissions

AWD management often results in an increment, sometimes important, of CO<sub>2</sub> and N<sub>2</sub>O emissions. As in concerned Mediterranean areas, drainage of the water table in the middle may not significantly increase fluxes of N<sub>2</sub>O with respect to the absence of drainage (Table 60).

## 5.3 Decreases in production (e.g. food/fuel/feed/timber)

In general, in any weather and soil conditions, mid-season drainage, rain-fed and other water-saving techniques resulting in reduced the yield grain (Meijide *et al.* 2017). For avoiding the decrease of yield under AWD management, some complementary measures can be used, such as timely irrigation, coordinating irrigation with fertilization and weed control (Zhi, 2000). For example, field experiences of Fangueiro *et al.* (2017) indicate that plots with NT and S irrigation obtain a grain yield (8.2 t/ha), which is similar to the yield of fields with CW and CT (between 6.7 and 8.9 t/ha)

# 6. Recommendations before and during the practice implementation

The success of the implementation of water saving practices is based in simultaneous improving agronomic practices. It is therefore necessary to know soil properties such as EC, texture, bulk density and pH, and climatic conditions, which determine water movement in soil and its fertility (Amin *et al.*, 2011; Fangueiro *et al.*, 2017). The main available strategies include fertilization and other agronomic management, developing improved varieties, changing the crop planting date, optimum use of rainfall, supplementary irrigation in rain-fed fields, improving water distribution, and water reuse or recycling. For example, mid-summer drainage, intermittent irrigation, and subsurface drainage systems can accelerate the leaching of reducible iron, to results in degradation of paddy fields, so application of iron materials is a possible countermeasure for increasing soil oxidation capacity.

In general, a good option could be the incorporation of organic matter amendments, although always without exceeding a certain threshold level from which production decreases again (Haque *et al.*, 2017). In any case, we must not forget that more amendment implies more SOC and usually more emission.

A common recommended practice for many regions of Asia and that is working well in Philippines, Vietnam, and Bangladesh (Nelson *et al.*, 2015) consists in starting the dry phase during the growing season, about 2–3 weeks after transplanting (or 3–4 weeks after sowing). Generally, the field dries until the saturated soil zone reaches a level of approximately 10–15 cm below the soil surface. At that time, it is irrigated again until the standing water reaches 3–5 cm on the field. The whole process can be controlled by piezometers (Photo 21). Depending on the soil and rain regime of the area, soil-drying time can take up to 10 days. In any case, the choice of the time of irrigation, with the soil dry or still somewhat humid, is relevant because grain yield and aboveground biomass are less if you wait for is fully dry (Xu *et al.*, 2015). One of the things to keep in mind with these practices is that high levels of water are required after transplanting and proper rooting, as well as during

the flowering stage. On the other hand, low water levels are required in tillering, panicle development and maturation stages. About 5 cm water depth is needed at milk stage for translocation of nutrients stored in plant body to ear or panicle for healthy development of developing grain or spikelet (Amin *et al.*, 2010).

A recommended practice applied in the South Democratic People’s Republic of Korea (Haque *et al.*, 2017) consists of incorporation of cover crop biomass (3 t/ha), which maintained an optimal and increasing level of rice paddy yield compared to CW. The yield differences between both systems (AWD and CW) did not show significant differences between them, but AWD reduced water inputs and greenhouse gas emissions. Therefore, in rainfed areas, the use of complementary methodologies such as the application of a thin transparent polyethylene film, or the incorporation of straw, can increase the efficiency of this management; especially if what is sought is to apply irrigation to keep the soil moist without standing water.

## 7. Potential barriers to adoption

**Table 62.** Potential barriers to adoption

Barrier	YES/NO	
Cultural	Yes	Tendency among farmers to flood the field as much as possible (Howell, Shrestha and Dodd, 2015).
Economic	No	Cost of supplementary measures (drainage in the wet season, weed control, preliminary tests of management viability, etc.)
Knowledge	Yes	Individual farmer may not be aware that AWD practice can save water for his or her fellow farmers’ field without compromising the yield on his/her field. Some farmers perceived that AWD will increase weed infestation (Howell, Shrestha and Dodd, 2015). Hence, building awareness will be necessary.

## Photos of the practice



Photo 18. Removing water in AWD (Valencia-Spain)



Photo 19. Flooding conditions in rice fields (Albufera of Valencia - Spain)



Photo 20. Dried fields in Mediterranean wetland (Abuferas of Valencia - Spain)



Photo 21. Piezometer

Table 63. Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Water regimes in rainfed rice-paddies in Indonesia and Thailand</i>	Asia	1 and 3	6	16

## References

- Amin, M.S.M., Rowshon, M.K. & Aimrun, W. 2011. Paddy Water Management for Precision Farming of Rice. In Uhlig, U. (Ed.). *Current Issues of Water Management*. <https://doi.org/10.5772/28883>
- Chen, A., Xie, X., Ge, T., Hou, H., Wang, W., Wei, W. & Kuzyako, W. 2017. Rapid decrease of soil carbon after abandonment of subtropical paddy fields. *Plant Soil*, 415: 203-214. <https://doi.org/10.1007/s11104-016-3154-0>
- Fangueiro, D., Becerra, D., Albarrán, A., Peña, D., Sanchez-Llerena, J., Rato-Nunes, J.M., López-Piñero, A. 2017. Effect of tillage and water management on GHG emissions from Mediterranean rice growing ecosystems. *Atmospheric Environment*, 150: 303-312. <https://doi.org/10.1016/j.atmosenv.2016.11.020>
- Haque, M., Biswas, J.C., Kim, S.W. & Kim P.J. 2017. Intermittent drainage in paddy soil: ecosystem carbon budget and global warming potential. *Paddy Water Environ*, 15: 403-411. <https://doi.org/10.1007/s10333-016-0558-7>
- Howell, K.R., Shrestha, P. & Dodd, I.C. 2015. Alternate wetting and drying irrigation maintained rice yields despite half the irrigation volume, but is currently unlikely to be adopted by smallholder lowland rice farmers in Nepal. *Food and Energy Security*, 4(2): 144–157. <https://doi.org/10.1002/fes3.58>
- Maeght, J.L., Grungerger, O., Hammecker, C., Sukchan, S., Hartmann, C. & Wiriyakinateekul, W. 2005. Salinity control by farmers practices in sandy soil. In *Proceedings of Management of Tropical Sandy Soils for Sustainable Agriculture: A holistic approach for sustainable development of problem soils in the tropics*. 27<sup>th</sup> November – 2<sup>nd</sup> December 2005, Khon Kaen, Thailand
- Mejjide, A., Gruening, C., G., Goded, I., Seufert, G. & Cescatti, A. 2017. Water management reduces greenhouse gas emissions in a Mediterranean rice paddy field. *Agriculture, Agri. Ecosys. Environ.*, 238: 168-178. <https://doi.org/10.1016/j.agee.2016.08.017>
- Moreno-Ramón, H., Marqués-Mateu, A., Ibáñez-Asensio, S. & Gisbert, J.M. 2015 Wetland soils under rice management and seawater intrusion: characterization and classification. *Spanish Journal of Soil Science*, 5(2): 111–129.
- Nelson, A., Wassmann, R., Sander, B.O. & Palao, L.K. 2015. Climate-Determined Suitability of the Water Saving Technology "Alternate Wetting and Drying" in Rice Systems: A Scalable Methodology demonstrated for a Province in the Philippines. *PLOS ONE*, 10(12): e0145268. <https://doi.org/10.1371/journal.pone.0145268>
- Nielsen, D. & Brock, M.A. 2009. Modified water regime and salinity as a consequence of climate change: prospects for wetlands of Southern Australia. *Climatic Change*, 95: 523-533. <https://doi.org/10.1007/s10584-009-9564-8>
- Nishio, M., Tanaka, H., Tanaka, D., Kawakami, R., Edo, K. & Yamazaki, Y. 2016. Managing Water Levels in Rice Paddies to Conserve the Itasenpara Host Mussel *Unio douglasiae nipponensis*. *Journal of Shellfish Research*, 35(4): 857-863. <https://doi.org/10.2983/035.035.0414>



Pan, G., Li, L., Zhang, X., Dai Jingyu, Z.Y. & Zhang, P. 2003. Soil organic carbon storage of China and the sequestration dynamics in agricultural lands. *Advances in Earth Science*, 18(4): 609–618.

Ramsar Convention on Wetlands. 2018. *Global Wetland Outlook: State of the World's Wetlands and their Services to People 2018*. Gland, Switzerland, Ramsar Convention Secretariat. (also available at: [https://static1.squarespace.com/static/5b256c78e17ba335ea89fe1f/t/5b9ffd2e0e2c7277f629eb8f/1537211739585/RAMSAR+GWO\\_ENGLISH\\_WEB.pdf](https://static1.squarespace.com/static/5b256c78e17ba335ea89fe1f/t/5b9ffd2e0e2c7277f629eb8f/1537211739585/RAMSAR+GWO_ENGLISH_WEB.pdf))

Sander, B.O., Wassmann, R., Palao, L.K. & Nelson, A. 2017. Climate-based suitability assessment for alternate wetting and drying water management in the Philippines: a novel approach for mapping methane mitigation potential in rice production. *Carbon Management*, 8(4): 331–342. <https://doi.org/10.1080/17583004.2017.1362945>

Spanu, A., Daga, L., Orlandoni, A.M. & Sanna, G. 2012. The role of irrigation techniques in arsenic bioaccumulation in rice (*Oryza sativa* L.). *Environ. Sci. Technol*, 46: 8333–8340. <https://doi.org/10.1021/es300636d>

Sriphrom, P., Chidthaisong, A. & Towprayoon, S. 2019. Effect of alternate wetting and drying water management on rice cultivation with low emissions and low water used during wet and dry season. *Journal of Cleaner Production*, 223: 980 – 988. <https://doi.org/10.1016/j.jclepro.2019.03.212>

Shiratori, Y., Watanabe, H., Furukawa, Y., Tsuruta, H. & Inubushi, K. 2007. Effectiveness of a subsurface drainage system in poorly drained paddy fields on reduction of methane emissions. *Soil Sci. Plant Nutr*. 53: 387–400. <https://doi.org/10.1111/j.1747-0765.2007.00171.x>

Tyagi, L., Kumari, B. & Singh, S.N. 2010. Water management: A tool for methane mitigation from irrigated paddy fields. *Sci. Total Environ.*, 408: 1085–1090. <https://doi.org/10.1016/j.scitotenv.2009.09.010>

Wissing, L., Kölbl, A., Schad, P., Bräuer, T., Cao, Z.H. & Kögel-Knabner, I. 2014. Organic carbon accumulation on soil mineral surfaces in paddy soils derived from tidal wetlands. *Geoderma*, 228: 90–103. <https://doi.org/10.1016/j.geoderma.2013.12.012>

Xionghui, J., Jiamei, W., Hua, P., Lihong, S., Zhenhua, Z., Zhaobing, L., Faxiang, T., Liangjie, H. & Jian, Z. 2012. The effect of rice straw incorporation into paddy soil on carbon sequestration and emissions in the double cropping rice system. *Journal of the Science of Food and Agriculture*, 92(5): 1038–1045. <https://doi.org/10.1002/jsfa.5550>

Xiong, Z., Liu, Y., Wu, Z., Zhang, X., Liu, P. & Huang, T. 2015. Differences in net global warming potential and greenhouse gas intensity between major rice-based cropping systems in China. *Scientific Reports*, 5(1): 17774. <https://doi.org/10.1038/srep17774>

Xu, Y., Ge, J., Tian, S., Li, S., Nguy-Robertson, A.L., Zhan, M. & Cao, C. 2015. Effects of water-saving irrigation practices and drought resistant rice variety on greenhouse gas emissions from a no-till paddy in the central lowlands of China. *Science of The Total Environment*, 505: 1043–1052. <https://doi.org/10.1016/j.scitotenv.2014.10.073>

**Ying, X., Junzhu, G., Shaoyang, T., Shuya, L., Nguy-Robertson, A., Ming, Z., Cougui, C.** 2015. Effects of water-saving irrigation practices and drought resistant rice variety on greenhouse gas emissions from a no-till paddy in the central lowlands of China. *The Science of the total environment*, 505: 1043-1052

**Zhang, A., Cui, L., Pan, G., Li, L., Hussain, Q., Zhang, X., Zheng, J. & Crowley, D.** 2010. Effect of biochar amendment on yield and methane and nitrous oxide emissions from a rice paddy from Tai Lake plain, China. *Agriculture, Ecosystems & Environment*, 139(4): 469–475.

<https://doi.org/10.1016/j.agee.2010.09.003>

**Zhi, M.** 2000. Water-efficient irrigation regimes of rice for sustainable increases in water productivity. *In Proceedings of International Rice Research Conference*, International Rice Research Institute, Laguna, Philippines

# 17. Straw residue management

Sara Ibáñez-Asensio

Department of plant production, Universitat Politècnica de València, Valencia, Spain

## 1. Description of the practice

Possible alternatives for the management of rice residues are: i) incorporation ii) burning, or iii) elimination. Stubble burning has been widely used in the past, and refers to the deliberate act of applying fire to the straw stubble that remains after the rice is harvested. Burning straw can show positive and negative effects on soil and crop yield in the short and long term, depending on the burning technique, the regional climatic and topographic conditions. It is necessary to be careful with a high intensity fires, high slope topographies or torrential rains. In general, burning a crop in the field is not the best recommended choice as it results in air pollution and high CO<sub>2</sub> emissions in nearby areas with people, and may damage soils (Whitbread *et al.*, 2003). However, the scientific community and the public acknowledge that burning residues through the use of slash-and-burn agriculture, some communities do not rely on external inputs based on fertilizers, pesticides and irrigation with the use of fossil energies. Those agroecosystems are irreplaceable and ecologically acceptable for that type of communities that generally are located in developing countries (Kleinman *et al.*, 1995). Another straw management that is used by farmers, is to cut and remove rice straw for another use outside the field (e.g. for livestock feeding, biosolids).

Residues can also be left on the soil after harvest, and incorporated into the upper soil layer, leading to a slow plant biochemical degradation. Specific machinery used in rice paddies in rotation allows the cutting and removal of rice straw, in order to sow the next crop (wheat or another cereal) on the bare soil, and after planting the deposition of straw as mulch. Rice straw can also be used in compost preparations and, be returned to the soil as a source of stabilized organic matter. Both options can also be carried out with other plant residues such as corn or wheat, in areas where rice is grown in rotation. Another recent possibility is the use of biochar, which is a C-rich by-product obtained by biomass pyrolysis under limited oxygen conditions (Also see factsheet n° 17 on Biochar, volume 3). In any of the cases (fresh residue incorporation, compost or biochar), its incorporation can be carried out-with or without tillage or accompanied or not by inorganic N or NPK fertilizers applications. Rice straw incorporation is considered as a good agronomic practice as long as site-specific management practices are implemented to improve soil properties and increase rice production (Asai *et al.*, 2009; Jiang *et*

*al.*, 2019). Regardless of how the organic matter is applied in the paddy fields (with or without N or NPK fertilizer application, with or without tillage, or simple deposition as mulch), it is widely documented that incorporating rice straw to the soil seems to increase SOC stocks but it negatively affects GHG emissions. Recent studies conclude that the intensity of the effects (positives and negatives) will depend on: i) the climate; ii) the type and the amount of fertilizer; iii) the amendment (type, amount and the frequency); iv) the initial content of nitrogen; v) organic carbon and other soil properties; and vi) duration of the period of soil saturation (see sections 4 and 5).

## 2. Range of applicability

The agronomic benefits derived from the incorporation practices have shown to be important in all type of soils and rainfall situations, as is evidenced by different meta-analyses carried out in many paddy rice areas with different environmental conditions in China and India (Huang *et al.*, 2013; Srinivasarao *et al.*, 2014). Specifically in terms of soils, positive effects have been registered from depleted soils (oxisols) to poorly developed soils (inceptisols), including also very heavy soils with silt layer (vertisols) and from low-nitrogen content soils (<1) to high N content soils (> 1.5 kgN per kg of soil). In reference to the climatic conditions, these studies focus on paddy areas with annual mean precipitation from < 1 000 mm to more than 1 500 mm and from cool annual mean temperatures (< 10 °C) to high temperatures (> 20 °C).

Incorporation of rice straw is already common practice in large areas of the world's most important rice growing areas. In this sense, alternatives that are more efficient are being explored recently in relation to the specific environmental conditions of the crop. For example, critical C inputs requirements for maintaining SOC stock are highest in coarse-textured soils and highest rates of rainfall. Consequently, in the cold northeast China region the return of soil fertility due to the addition of straw is very slow, and biochar is a possible solution for a much faster restitution of SOC (Sui *et al.*, 2016).

In addition, in semi-arid tropics (where rainfall exceeds potential evapotranspiration only two or four months of year), if rice-wheat residues are pyrolyzed before soil addition, about 50 percent of the C in biomass can be returned to soil (Srinivasarao *et al.*, 2014). The straw addition technique is also applied in arid and semi-arid Mediterranean environments where traditional burning was replaced by the incorporation of the fresh residue (Jégou and Sanchis-Ibor, 2019).

### 3. Impact on soil organic carbon stocks

As indicated before, compared to the removal option, crop residue retention is always a good option for increasing soil C storage. And as for the type of residue, in principle biochar would be the best alternative, since it can be known as an indicative value that the carbon content of the rice straw biochar can double the total C content of the initial vegetable residue. In traditional paddy rice fields in northern China, biochar addition of 1.78 t/ha increases soil C by up to between 6 and 12 g/kg. In the case of straw rice incorporation (5 t/ha), increases reaches 4 g/kg of C. While it may sometimes seem that the rice straw amendment causes minimal changes in the total C content of the soil, a meta-analysis from China suggests that soil C saturation would occur after 12 years of straw C input (Zhu *et al.*, 2014).

In Chinese clay loam soils, straw reintroduction led to lower SOC concentration increases than in sandy loam and silty soils (Liu *et al.*, 2014). In addition, rotary tillage rice straw (6 t/ha) would lead to higher SOC increases than conventional tillage. The net increase was higher at 14-21 cm depth (18.4 percent) than 7-14 cm (8.7 percent) and 0-7 cm depths (7 percent) (Zhu *et al.*, 2014). At last, the application of biochar doses above 30 t/ha triggers carbon immobilization well above the increase that can be achieved with the addition of non-composted straw (50-90 percent for biochar versus 10-20 percent for straw), regardless of whether it has been used nitrogen fertilization (usually urea). Lower doses of biochar do not seem to enhance carbon sequestration much more than that obtained by incorporating straw residue rates of 3-6 t/ha (Table 64).

**Table 64.** Changes in soil organic carbon stocks reported for straw residue management

Location	Climate (MAR, MAP)	Soil type	SOC stocks (tC/ha)		Annual Cseq (tC/ha/yr)	Depth (cm)	Duration (Years)	Crop and soil management	NPK Additions (t/ha)	Biomass incorporation (t/ha)	Reference
			*C: 40.7 *RS:44.7 *B1: 46 *B2:48.9 *B3: 62.7	C: 39.7 RS:44.3 B1: 44 B2:44.4 B3:76.5							
Shenyang, Liaoning Province, Northeast China	Semi-humid temperate, continental monsoon (500 mm, 8.3 °C)	pH: 6.7 BD: 1.31 g/cm <sup>3</sup> C/N: 9.17	NA		NA	0-20	2	- Rice variety: <i>Japonica Shennong 265</i>  - Upland crop-single rice	With urea : N: 0.21 P <sub>2</sub> O <sub>5</sub> : 0.62 K <sub>2</sub> O: 0.2	RS: 5.05 (370 gC/kg; 7 gN/kg) incorporated before transplanting at 5 cm; B1: 1.78 B2:14.8 B3: 29.6  Biochar composition: 671 gC/kg; 8.1 gN/kg	Sui <i>et al.</i> (2016)  data recalculated
			*C: 23.6 *RS:28.2 *B1: 29 *B2: 30 *B3: 33.7	C: 21.93 RS: 27.5 B1: 26 B2: 27.7 B3: 35.9		20-40					
Tai Lake plain, Yiangsu, Eastern China	Subtropical monsoon (1177 mm; 15.7 °C)	Entic Hapludept 24 gC/kg BD= 1 g/cm <sup>3</sup>	C: 34.4 B1: 39.02 B2: 48.06	NA	NA	0-15	1	- Rice variety: <i>cv. Wuyunjing 7</i>  - Water management: AWD Flooding-Drainage-re-Flooding -Moist	With urea: N: 0.3 P <sub>2</sub> O <sub>5</sub> : 0.13 K <sub>2</sub> O: 0.13	B1: 10 B2: 40	Zhang <i>et al.</i> (2010)  data recalculated
Wnagcheng and Changsha County, Hunan Province,	1370 mm, 17°C	pH: 6.6 20.03 gC/kg 2.1 gTN/kg	*C: 33 NPK: 34.6 RS+NPK:36.3	*C: -0.07 NPK: 0.08 RS+NPK:0.1	NA	NA	28	- 28-year trial: Conventional and Hybrid ( <i>Xiang 67</i> ), conventional tillage	N: 0.15 ( <i>v.early</i> ) + (0.18 <i>v.late</i> ) P: 38.7 x 2 K: 99.6 x 2	RS: 2.63	Xionghui <i>et al.</i> (2012)

Location	Climate (MAR, MAP)	Soil type	SOC stocks (tC/ha)	Annual Cseq (tC/ha/yr)	Depth (cm)	Duration (Years)	Crop and soil management	NPK Additions (t/ha)	Biomass incorporation (t/ha)	Reference
Southern China	1500 mm, 17.1°C	pH: 5.5 12.7 gC/kg 2.1 gTN/kg	C: 20.8 RS: 21.3	C: -0.02 RS: 0.16		1	- 1-year trial: Early and late rice, no-till - Double rice in both trials	N: 0.15 (v. early) + 0.18 (v. late) P: 0.9 + 0.45 K: 0.9 + 0.112	RS: 4.5	
Yangtze Delta Plain, Yiangsu, Eastern China	Subtropical monsoon (1050 mm, 15.7 °C)	CEC: 15 cmol/kg BD: 1.28 g/cm <sup>3</sup>	NA	C: 0.08 RS1: 0.94 RS2: 1.77	0-25	3	Upland crop-single rice: wheat-rice; Short flood	With urea: N: 0.45 P <sub>2</sub> O <sub>5</sub> : 0.06 K <sub>2</sub> O: 0.12	RS1: 3 RS2: 6	Xiong <i>et al.</i> (2015)
				C: 0.47 RS1: 1.43 RS2: 2.42			Double rice (oil-rape-rice-Rice); Long Flood			
Cuttack, Odisha, NE India	Tropical (1500 mm)	Sandy clay loam (Aeric Endoaquept) TC: 4.9 g/kg TN: 0.5 g/kg BD: 1.41 g/cm <sup>3</sup>	*C: 8.08 U: 8.88 RS+U: 9.42 RS+GM: 9.35	*C: - 0.41 U: 0.07 RS+U: 0.35 RS+GM: 0.22	0-15	4	- Rice variety: <i>Cv Gayatri</i> - Continuous flooding and conventional tillage	0.6 tN/ha	RS: 383 gC/kg GM: 370 gC/kg U: 0.6 tN/ha RS+U: 0.3+0.3 RS+GM: 0.3+0.3	Bhattacharyya <i>et al.</i> (2012)

\*Without fertilizer nor organic amendment

+ According to the dose of urea

C: control (straw removal); RS: rice straw (incorporation); B: biochar (incorporation); GM: green manure incorporation; U: urea; CW: continuous flooding; AWD: alternate wetting and drying; BD: bulk density; NA: non available

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Regardless of their form of application (fresh, compost or biochar) and nitrogen fertilization (with or without), applying straw residues on the first centimeters of soil helps to improve nitrogen contents, porosity, capillary porosity, and air-filled porosity (Sui *et al.*, 2016). In addition, long-term incorporation of straw increases rice root biomass (Jiang *et al.*, 2019). Soils treated with rice husk biochar generally show a higher cation exchange capacity (Wang *et al.*, 2011), and in most studies, it has been found that microbial biomass increases, with significant changes in the microbial community component (Lehmann *et al.* 2011).

### 4.2 Minimization of threats to soil functions

Table 65. Soil threats

Soil threats	
<b>Nutrient imbalance and cycles</b>	The addition of biochar increases nutrient retention and decreases its leaching. It increases fertility, especially on tropical soils (Noguera <i>et al.</i> , 2010). In addition, it is estimated that rice straw contains approximately 40 percent of the nitrogen absorbed by rice, 30 percent of phosphorus and 80 percent of potassium, and can therefore be reincorporated into the soil and can be reintroduced into soil (Chivenge <i>et al.</i> , 2020).
<b>Soil acidification</b>	Biochar addition increases by 6 to 12 percent in acid (4.6 pH) and neutral soils (6.2 pH) respectively (Liu <i>et al.</i> , 2012)
<b>Soil biodiversity loss and soil compaction</b>	Straw addition increases the amount of soil macro-aggregates because: i) its decomposition increases colloids to binding micro-aggregates to macro, and ii) increases microbial biomass and therefore the production of microbial-derived binding agents (Liu <i>et al.</i> , 2014)  Long-term straw incorporation increases soil methanotrophic abundance and root size (Jiang <i>et al.</i> , 2019).
<b>Soil water management</b>	Biochar improves the saturated hydraulic conductivity of topsoil and xylem sap flow of the rice plant (Asai <i>et al.</i> , 2009).



### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Benefits on productivity resulting from the incorporation to the soil of any form of organic matter varies with the temperature and rainfall of each area. This is because in hot and rainy places processes of decomposition of organic matter and the washing of salts from the soil are rapid, while the drop in temperature and rainfall slows them down.

For example, for Indian semiarid tropics rice paddy areas, each t/ha increase in SOC stock in the root zone lead to an increase of 0.16 t/ha in grain rice yield. In the semi-arid and sub-humid areas with low rainfall (average annual rainfall < 1 000 mm) and therefore poorly developed soils (Inceptisols), an increase in rice production of between 15 to 20 percent as response to additional return of rice straw about 5 to 9 t/ha/yr was noted. When the rain reaches 1 000 mm, leaching increases and with it the impoverishment of the soil. In these conditions adding organic matter means an increase of yield about 40 percent (in Vertisols) and 50 percent (in Oxisols) (Srinivasarao *et al.*, 2014).

In China, rice straw incorporation or biochar show more discreet outcomes (Liu *et al.*, 2012; Xiong *et al.*, 2015). Specifically, Huang *et al.*, 2013 in their meta-analysis show that, in general and regardless of the soil N content, crop residues retention increased rice yields by 5.2 percent as average value. The magnitude of the increase was significant at any average annual temperature but the highest increase was 7.2 percent for an area with average temperatures <10°C and the smallest was by 3.3 percent for 10-15 °C. Finally, the meta-analysis show that straw incorporation increases rice yield with an increase in the duration of the application (4.7 percent for 3 years and 9.7 percent for > 10 years).

### 4.4 Mitigation of and adaptation to climate change

In general, when rice residue has been removed, addition of biochar does not increase GHG emissions. For the management conditions analysed in China (with plots where nitrogen fertilization has been applied), incorporate low rates of biochar (about 2 t/ha) decrease CH<sub>4</sub> emissions while medium rates (around 10 or 30 t/ha) produce little changes. However, high rates (above 40 t/ha) increase emissions (Sui *et al.* 2016; Zhang *et al.* 2010). Regardless, experiences in field trials in a hot and humid tropic of the Philippines (Knoblauch *et al.*, 2010) show that the incorporation of large amounts of charred rice husks, only significant increased methane emissions in the first season, since in the following years it decreases rapidly.

On one hand, in the climatic and management rice conditions of China, Sui *et al.* (2016) report that biochar addition decreases CO<sub>2</sub> emissions only if rates of application are medium or large (above 10 t/ha in the cases analysed). On the other hand, the studies related by Liu *et al.* (2012) and Zhang *et al.* (2010) confirm that there is a significant reduction of N<sub>2</sub>O emissions, which will be greater as more biochar is incorporated into the soil. For example, with rates of about 40 t/ha, nitrous oxide emissions fell by about half. By reducing the dose of biochar, the drop is approximately 30 percent.

Data from China indicate that rice straw management only reduces nitrous oxide and CO<sub>2</sub> emissions if low amounts of residues are incorporated. In the eastern (Xiong *et al.*, 2015) with rates below 6 t/ha, the recorded reductions of nitrous oxide do not exceed 10 percent but if the contribution of the straw is made for the second time in the year (double rice management), then emissions increase. As for CO<sub>2</sub>, for the conditions analyzed by Sui *et al.* (2016) in northeast the addition of 5 t/ha of straw reduces the emissions by half compared to the control if urea is not supplied. On contrary, if nitrogen fertilizer is applied, the emissions are very similar to the control.

As to global warming, to date there is limited knowledge regarding biochar incorporation. Study conducted in northeast China (Sui *et al.*, 2016) in paddy rice fields without nitrogen fertilization indicate that straw incorporation had a global warming potential nearly 1.5 times more than that of 29.6 t/ha biochar amendment. In the same vein, they also found that a large biochar application with N fertilization markedly decreased GWP when compared with rice straw incorporation, nevertheless the improvement in relation to the complete removal of the straw from the fields and the role that nitrogen incorporation plays in all this was not clear. Prendergast-Miller *et al.* (2014) attribute the positive effect of biochar to increased oxygen released by rice roots, which stimulates methanotrophs and can suppress methanogens in the rhizosphere.

If, on the contrary, the net GWP is considered, the effect of applying straw does not seem so negative. In a two-year study in the south Democratic People's Republic of Korea (Lee *et al.*, 2020) showed that the sum of methane (CH<sub>4</sub>) and nitrous oxide (N<sub>2</sub>O) fluxes and net ecosystem carbon budget (difference between C input and output) is greater in plots with straw removal. The reported values were 12.65 t CO<sub>2</sub>eq/ha the first year and 6.92 the second in plots without straw; 6.02 t CO<sub>2</sub>eq/ha the first year and 3.46 t CO<sub>2</sub>eq/ha the second when straw is mixed with soil; and finally, 13.36 and 7.38 t CO<sub>2</sub>eq/ha respectively where straw is left over soil. To do the GWP net balance calculation, C input source is the straw incorporated plus net primary production, and harvest C removal and heterotrophic respiration C loss are counted as C output.

## 4.5 Socio-economic benefits

Of the three possible alternatives for the management of rice residues, incorporation is the most beneficial alternative for the soil and the environment (as supported by the data provided in the previous sections 4.1, 4.2 and 4.3 and the later: 5.4 and 5.5). For example, in arid and semi-arid Mediterranean environments, mulch use maintains soil moisture during dry season and can be an important complement as water sheet management is implemented when soil surfaces are dry.

Regarding fertilizer, the most widespread practice in paddy management is to apply inorganic NPK fertilization, at rates that can reach around 200 kgN/ha (with urea), 125 kgP<sub>2</sub>O<sub>5</sub>/ha and 125 kgK<sub>2</sub>O/ha. However, it is possible that crop residues application can substitute a part of these inorganic fertilizers. In reality, rice yield in China and the Lao People's Democratic Republic is not adversely affected as a consequence of the rates reduction of inorganic N, P, and K fertilizers by averages between 8.3 and 29.4 percent. (Wang *et al.*, 2011; Asai *et al.*, 2009). In the case of tropical areas, rice straw incorporation or biochar show more limited outcomes (Liu *et al.*, 2012; Xiong *et al.*, 2015), but even so a cover without tillage is associated with a high-efficiency practice because it allows decrease tillage and inorganic fertilizers input.

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

Table 66. Soil threats

Soil threats	
<b>Nutrient imbalance and cycles</b>	Biochar addition produces N immobilization and a decrease in plant N uptake due to the effect of C in the C/N ratio (Lehmann <i>et al.</i> , 2002; Asai <i>et al.</i> , 2009).
<b>Soil water management</b>	Prolonging the waterlogging time causes increase on global warming. On at same subtropical monsoon climate, the increase due to rice straw incorporation can range values from 6-12 percent by upland crop single rice to 107-180 percent by double rice in flooding conventional (Xiong <i>et al.</i> 2015).

### 5.2 Increases in greenhouse gas emissions

Under the most common rice growing conditions, incorporation of straw worsens forecasts on global warming with respect to the option of disposing of rice straw. However, depending on agricultural management conditions important differences can be observed in the magnitude of these increases. In general, the studies realized in China by Sui *et al.* (2016) show that even when rates are low or moderate (beneath 6 t/ha/yr), to complete with nitrogen fertilization is recommended because adding only rice straw increase GWP by 60 percent (Sui *et al.*, 2016).

In general, in rice production area in Asia (China and India), methane emission increases with increasing amounts of rice straw incorporation regardless both nitrogen fertilization and water management. The increases in emissions are also biggest when rice is flooded during long periods, and the magnitude of increases can even reach 900 percent. However, the long-term addition of straw rice makes CH<sub>4</sub> and CO<sub>2</sub> emissions stabilize. The maximum values are registered after the first year of application and went on decreasing season after season during three years (Philippines, Knoblauch *et al.*, 2010; the South Democratic People's Republic of Korea, Luyima *et al.*, 2019). In this aspect, emissions in double and single rice system in Chinese rice production areas (with a straw rate of approximately 6.6 and 4.2 Mt/ha respectively) by plowing during 20 years were, on average, 48 percent lower than CH<sub>4</sub> emissions IPCC estimates (Jiang *et al.*, 2019).

About biochar, as seen in section 4.4, the impact of its incorporation depends largely on the agricultural practices carried out. In China (Sui, *et al.*, 2016), low rates incorporation produces higher CO<sub>2</sub> emissions than if the residue were completely removed. Furthermore, when there is no N fertilization, methane emissions are higher than when there is, regardless of the amount of biochar incorporated. Lastly, despite the few available studies on N<sub>2</sub>O emissions, it is indicated the possibility of a threshold value of incorporated stubble from which the emissions go from being negative increases (-30 percent) to positive (+10 percent), so more GWP is produced (Xiong *et al.*, 2015; Das and Adhya, 2014).

**Table 67.** GHG emissions and GWP according to residue type incorporated

Location and reference	Climate and soil type	More information	GHG emissions (t/ha/yr)				GWP (tCO <sub>2</sub> eq/ha/yr)		Duration (Years)	Reference
			CH <sub>4</sub>		CO <sub>2</sub>		CH <sub>4</sub> + CO <sub>2</sub>			
Shenyang, Liaoning Province, Northeast China	See Table 64	UR only; For biomass and fertilizer additions (Table 64)	*C: 0.045	C: 0.075	*C: 8.46	C: 20.9	*C: 9.6	C: 22.8	2	Sui <i>et al.</i> (2016)
			*RS: 0.424	RS: 0.237	*RS: 4.81	RS: 19.04	*RS: 15.4	RS: 25		
Tai Lake plain, Yangsu, Eastern China	See Table 64	For biomass and fertilizer additions (Table 64)	B1*: 0.102	B1: 0.031	*B1: 8.78	B1: 21.45	*B1: 11.3	B1: 22.2	1	Zhang <i>et al.</i> (2010)
			B2*: 0.073	B2: 0.066	*B2: 7.99	B2: 29.43	*B2: 9.8	B2: 31		
Wnagcheng and Changsha County, Hunan Province, Southern China	See Table 64	GHG emissions and GWP: CH <sub>4</sub> only For biomass and fertilizer additions (Table 64)	*C: 0.64				*C: 15.96		28	Xionghui <i>et al.</i> (2012)
			NPK: 0.49				NPK: 12.16			
			RS+NPK: 1.23				RS+NPK: 30.87			
			C: 0.21				C: 5.31		1	
			RS: 0.32				RS: 8.05			

Location and reference	Climate and soil type	More information	GHG emissions (t/ha/yr)		GWP (tCO <sub>2</sub> eq/ha/yr)	Duration (Years)	Reference
			CH <sub>4</sub>	N <sub>2</sub> O			
Yangtze Delta Plain, Yiangsu, Eastern China		All cases: NPK (with U) UR: wheat-rice; short flood DR: oil-rape-rice-rice; CW For biomass and fertilizer additions (Table 64)	UR and DR	UR	CH <sub>4</sub> + N <sub>2</sub> O	3	Xiong <i>et al.</i> (2015)
			C: 0.104 RS1: 0.208 RS2: 0.302	C: 0.00226 RS1: 0.00208 RS2: 0.00219 DR C: 0.00277 RS1: 0.00313 RS2: 0.00304	C: 4.32 RS1: 4.6 RS2: 4.8 DR C: 5.87 RS 1: 12.2 RS 2: 16		
Yangtze river, Jiangsu Province, China	Monsoon (MAT: 23.7 °C)	GHG emissions CH <sub>4</sub> only; rice-wheat rotation WC, CT; NPK additions (t/ha): N: 0.3; P <sub>2</sub> O <sub>5</sub> : 0.15; K <sub>2</sub> O: 0.24	C: 0.12 WS: 0.66 RS: 0.13 RS+ WS: 0.64	NA	NA	2	Hou <i>et al.</i> (2013)
			N <sub>2</sub> O			1	Liu <i>et al.</i> (2012)

Location and reference	Climate and soil type	More information	GHG emissions (t/ha/yr)		GWP (tCO <sub>2</sub> eq/ha/yr)	Duration (Years)	Reference
Sichuan, Hunan and Jiangxi, South China	humid to semihumid climate; acid to neutral soils	B (wheat straw 450-550 °C): B1: 20 t/ha B2: 40 t/ha Urea: 0.24 <sup>(1)</sup> to 0.3 <sup>(2)</sup> t/ha	C*: 0.00146 <sup>(1)</sup> - 0.00188 <sup>(2)</sup> B1*: 0.00079 <sup>(1)</sup> - 0.00133 <sup>(2)</sup> B2*: 0.00068 <sup>(1)</sup> - 0.00087 <sup>(2)</sup> *according to the dose of urea				
Cuttack, Odisha, Northeast India	See table 64	For biomass and fertilizer additions (Table 64)	CH <sub>4</sub>	N <sub>2</sub> O	CH <sub>4</sub> + N <sub>2</sub> O	4	Bhattacharyya <i>et al.</i> (2012)
			*C: 0.069 U: 0.093 RS+U: 0.115 RS+GM: 0.127	*C: 0.0002 U: 0.001 RS+U: 0.0008 RS+GM: 0.00072	*C: 5.86 U: 8.08 RS+U: 9.42 RS+GM: 10.2		
Cuttack, Odisha, NE India	Tropical (MAP: 1500 mm) Sandy clay loam (Aeric Endoaquept)	Rice variety: <i>v IR 36</i> CW, CT Urea additions: 120 kgN/ha	CH <sub>4</sub>	N <sub>2</sub> O	CH <sub>4</sub> + N <sub>2</sub> O	1	Das and Adhya (2014)
			*C: 0.11 U: 0.15 RS+U: 0.21	*C: 0.00016 U: 0.00076 RS+U: 0.00057	*C: 2.9 U: 3.96 RS + U: 5.34		

\*without fertilizer or organic amendment

C: control (straw removal); RS: rice straw (incorporation); B: biochar (incorporation); WS: wheat straw (incorporation); U: urea; GM: green manure (incorporation); NT: no tillage; CT: conventional tillage; CW: continuous flooding

### 5.3 Conflict with other practice(s)

Rice vegetative cycle and the tillage operations carried out in each phase of the crop largely determine the GHG emission. The CH<sub>4</sub> flux for example, increases after rice transplantation and dropped quickly during the midseason drainage; after reflooding, it increases again to an emission peak and then decreases gradually to a negligible amount until harvest. In addition, the peak emission at the end of the reproductive stage has been observed in all fields receiving rice straw (Naser *et al.*, 2007).

### 5.4 Decreases in production (e.g. food/fuel/feed/timber)

A medium rate of biochar application (about 16 t/ha) increases N immobilization and so can leading to low grain yields (Asai *et al.*, 2009). The addition of straw also results in effective nitrogen immobilization in the soil, which can amount to up to 2-4 percent of the applied nitrogen (Said-Pullicino *et al.*, 2014)

### 5.5 Other conflicts

From an environmental point of view, there is a fundamental difference between the three possible alternatives for the management of rice residues. Burning residues in the field results in air pollution affecting nearby populated areas and high CO<sub>2</sub> emissions and can damage soil. Cao *et al.* (2006) estimated that in China, emissions from agriculture due to burning in field during 2000 were 100.0 Gg of black carbon and 395.8 Gg of organic carbon, contributing to the 6.8 percent and 9.8 percent respectively of total country emissions.

At this point, other economic considerations come in relation to the other two options of removing or incorporating crop residues. In fields with difficult-to-drain dammed soils, the withdrawal implies the use of machinery adapted to the soil waterlogging conditions, which greatly increases costs. In the case of transforming the stubble into biochar before returning it to the field, the cost of its subsequent incineration must be included (Srinivasarao *et al.*, 2014).

## 6. Recommendations before implementation of the practice

Soil water management also plays a relevant role in the potential for greenhouse gas emissions that straw residue incorporations represent (Xiong *et al.*, 2016). According to the meta-analysis on the Chinese rice paddy, a good soil and water management strategy to decrease the production of greenhouse gases is to avoid excessive C accumulation in soil, increase P availability, and decrease available of Fe content (Wang *et al.*, 2017).

In this sense, an increasingly common practice in many rice regions is to change the traditional water level management, replacing the permanent flooding (Continuous Waterlogging, CW) with the alternation of flooding with drainage until the soil is dried, or even only damp wetting (Alternate Wetting and Drying, AWD).

This system, in addition to reducing water consumption and saving on the costs associated with its handling, reduces CH<sub>4</sub> and in consequence the Global Warming Potential (GWP) and the intensity of greenhouse gases (Sriphrom *et al.*, 2019; Tiag *et al.*, 2010; Xu *et al.*, 2017). Sprinkler irrigation experiences carried out in the Mediterranean regions of Spain and Italy corroborate this trend and show very positive results (Fangueiro *et al.*, 2017; Mejide *et al.*, 2017)

For last, a common recommended practice for China (Huang *et al.*, 2013), is to apply inorganic N fertilizer in the early vegetative stage. SOC increase is independently by residue types (legume and non-legume) or labor type (tillage and non-tillage)

## 7. Potential barriers to adoption

**Table 68.** Potential barriers to adoption

Barrier	
Economic	See section 5.5

**Table 69.** Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Management of Rice straw in Mediterranean wetlands, Spain</i>	Europe	7	6	11
<i>Conservation Agriculture in intensive rice-based cropping systems in the Eastern Gangetic Plain</i>	Asia	5	6	12
<i>Organic rice cultivation with internal nutrient cycling in Japanese Andosols</i>	Asia	4, 8 and 12	6	14



## References

- Asai, H., Samson, B.K., Stephan, H.M., Songyikhangsuthor, K., Homma, K., Kiyono, Y., Inoue, Y., Shiraiwa, T. & Horie, T. 2009. Biochar amendment techniques for upland rice production in Northern Laos: 1. Soil physical properties, leaf SPAD and grain yield. *Field Crops Research*, 111(1-2): 81-84. <https://doi.org/10.1016/j.fcr.2008.10.008>
- Bhattacharyya, P., Roy, K.S., Neogia, S., Adhya, T.K., Rao, K.S. & Manna, M.C. 2012. Effects of rice straw and nitrogen fertilization on greenhouse gas emissions and carbon storage in tropical flooded soil planted with rice. *Soil and Tillage Research*, 124: 119-130. <https://doi.org/10.1016/j.still.2012.05.015>
- Cao, G., Zhang, X. & Zheng, F. 2006. Inventory of black carbon and organic carbon emissions from China. *J. Atmos. Envi.*, 40: 6516-6527. <https://doi.org/10.1016/j.atmosenv.2006.05.070>
- Chivenge, P., Rubianes, F., Van Chin, D., Van Thach, T., Khang, V.T., Romasanta, R.R., Van Hung, N. & Van Trinh, M. 2020. Rice Straw Incorporation Influences Nutrient Cycling and Soil Organic Matter. *In* M. Gummert, N.V. Hung, P. Chivenge & B. Douthwaite (Eds.) *Sustainable Rice Straw Management*, pp. 131–144. Cham, Springer International Publishing. [https://doi.org/10.1007/978-3-030-32373-8\\_8](https://doi.org/10.1007/978-3-030-32373-8_8)
- Das, T. & Adhya, K. 2014. Effect of combine application of organic manure and inorganic fertilizer on methane and nitrous oxide emissions from a tropical flooded soil planted to rice. *Geoderma*, 213: 185-192. <https://doi.org/10.1016/j.geoderma.2013.08.011>
- Fangueiro, D., Becerra, D., Albarrán, A., Peña, D., Sanchez-Llerena, J., Rato-Nunes, J.M., López-Piñeiro, A. 2017. Effect of tillage and water management on GHG emissions from Mediterranean rice growing ecosystems. *Atmospheric Environment*, 150: 303-312. <https://doi.org/10.1016/j.atmosenv.2016.11.020>
- Hou, P., Li, G., Wang, S., Jin, X., Yang, Y., Chen, X., Ding, C., Liu, Z. & Ding, Y. 2013. Methane emissions from rice fields under continuous straw return in the middle-lower reaches of the Yangtze River. *Journal of Environmental Sciences*, 25(9): 1874-1881. [https://doi.org/10.1016/S1001-0742\(12\)60273-3](https://doi.org/10.1016/S1001-0742(12)60273-3)
- Huang, S., Zeng, Y., Wu, J., Shi, Q. & Pan, X. 2013. Effect of crop residue retention on rice yield in China: A meta-analysis. *Field Crops Research*, 154: 188-194. <https://doi.org/10.1016/j.fcr.2013.08.013>
- Jégou, A. & Sanchis-Ibor, C. 2019. The opaque lagoon. Management and governance of water in l'Albufera wetland of Valencia (Spain). *Limnetica*, 38(1): 503-515.
- Jiang, Y., Qian, H., Huang S., Zhang X., Wang, L., Zhang, L., Shen, M., Xiao, X., Chen, F., Zhang, H., Lu, C., Li, C., Zhang, J., Deng, A., van Groenigen, K.J. & Zhang, W. 2019. Acclimation of methane emissions from rice paddy fields to straw addition. *Science Advances*, 5(1): 1-9. <https://doi.org/10.1126/sciadv.aau9038>
- Kleinman, P.J.A., Pimentel, D. & Bryant R.B. 1995. The ecological sustainability of slash-and-burn agriculture. *Agriculture, Ecosystems & Environment*, 52(2-3): 235.-249. [https://doi.org/10.1016/0167-8809\(94\)00531-I](https://doi.org/10.1016/0167-8809(94)00531-I)

- Knoblauch, C., Maarifat, A.A., Pfeiffer, E.M. & Haefele, S.M.** 2010. Degradability of black carbon and its impact on trace gas fluxes and carbon turnover in paddy soils. *Soil Biol. Biochem.*, 43(9): 1768-1778. <https://doi.org/10.1016/j.soilbio.2010.07.012>
- Lehmann, J., Rillig, M.C., Thies, J., Masiello, C.A., Hockaday, W.C. & Crowley, D.** 2011. Biochar effects on soil biota - A review. *Soil Biology & Biochemistry*, 43: 1812-1836. <https://doi.org/10.1016/j.soilbio.2011.04.022>
- Lehmann, J.P., Da Silva Jr., Rondon, M., Steiner, C., Nehls, T., Zech, W. & Glaser, B.** 2002. Nutrient availability and leaching in an archaeological Anthrosol and a Ferralsol of the Central Amazon basin: fertilizer, manure and charcoal amendments. *Plant and Soil*, 249: 343-357.
- Liu, X., Qu, J., Li, L., Zhang, A., Jufeng, Z., Zheng, J. & Pan, G.** 2012. Can biochar amendment be an ecological engineering technology to depress N<sub>2</sub>O emission in rice paddies?—A cross site field experiment from South China. *Ecological Engineering*, 42: 168–173. <https://doi.org/10.1016/j.ecoleng.2012.01.016>
- Liu, C., Lu, M., Cui, J., Li, B. & Fang, C.** 2014. Effects of straw carbon input on carbon dynamics in agricultural soils: a meta-analysis. *Global Change Biology*, 20: 1366-1381. <https://doi.org/10.1111/gcb.12517>
- Luyima, D., Jeong, H., Lee, J. H. & Kim, S. H.** 2019. Effects of Straw Incorporation Time on Rice Yield and Methane Emissions from Sandy Loam Paddy Fields. *Journal- Faculty of Agriculture Kyushu University*, 64(2): 213-218. <https://doi.org/10.5109/2339106>
- Mejjide, A., Gruening, C., Goded, I., Seufert, G. & Cescatti, A.** 2017. Water management reduces greenhouse gas emissions in a Mediterranean rice paddy field. *Agriculture, Ecosystems & Environment*, 238: 168–178. <https://doi.org/10.1016/j.agee.2016.08.017>
- Naser, H.M., Nagata, O., Tamura, S. & Hatano, R.** 2007. Methane emissions from five paddy fields with different amounts of rice straw application in central Hokkaido, Japan. *Soil Science & Plant Nutrition*, 53(1): 95–101. <https://doi.org/10.1111/j.1747-0765.2007.00105.x>
- Noguera, D., Rondón, M., Laossi, K.-R., Hoyos, V., Lavelle, P., Cruz de Carvalho, M.H. & Barot, S.** 2010. Contrasted effect of biochar and earthworms on rice growth and resource allocation in different soils. *Soil Biology and Biochemistry*, 42(7): 1017–1027. <https://doi.org/10.1016/j.soilbio.2010.03.001>
- Prendergast-Miller, M. T., Duvall, M. & Sohi, S.** 2014. Biochar–root interactions are mediated by biochar nutrient content and impacts on soil nutrient availability. *Eur. J. Soil Sci.*, 65: 173-185. <https://doi.org/10.1111/ejss.12079>
- Said-Pullicino, D. Cucu, M.A., Sodano, M., Birk, J.J, Glaser, B. & Celi, L.** 2014. Nitrogen immobilization in paddy soils as affected by redox conditions and rice straw incorporation. *Geoderma*, 228-229: 44-53. <https://doi.org/10.1016/j.geoderma.2013.06.020>
- Sriphrom, P., Chidthaisong, A. & Towprayoon, S.** 2019. Effect of alternate wetting and drying water management on rice cultivation with low emissions and low water used during wet and dry season. *Journal of Cleaner Production*, 223: 980–988. <https://doi.org/10.1016/j.jclepro.2019.03.212>

- Sui, Y., Gao, J., Liu, C., Zhang, W., Lan, Y., Li, S., Meng, J., Xua, Z. & Tang, L. 2016. Interactive effects of straw-derived biochar and N fertilization on soil C storage and rice productivity in rice paddies of Northeast China. *Science of the Total Environment*, 544: 203–210.  
<https://doi.org/10.1016/j.scitotenv.2015.11.079>
- Srinivasarao, Ch., Lal, R., Kundu, S., Babu, M.B.B.P., Venkateswarlu, B. & Singh, A.K. 2014. Soil carbon sequestration in rainfed production systems in the semiarid tropics of India. *Science of The Total Environment*, 487: 587–603. <https://doi.org/10.1016/j.scitotenv.2013.10.006>
- Tyagi, L., Kumari, B. & Singh, S.N. 2010. Water management – A tool for methane mitigation from irrigated paddy fields. *Science of The Total Environment*, 408(5): 1085–1090.  
<https://doi.org/10.1016/j.scitotenv.2009.09.010>
- Wang, J., Zhang, M., Xiong, Z., Liu, P. & Pan, G. 2011. Effects of biochar addition on N<sub>2</sub>O and CO<sub>2</sub> emissions from two paddy soils. *Biology and Fertility of Soils*, 47: 887–896.  
<https://doi.org/10.1007/s00374-011-0595-8>
- Wang, C., Sardans, J., Wang, C., Zeng, C., Tong C., Asensio, D. & Peñuelas, J. 2017. Relationships between the potential production of the greenhouse gases CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O and soil concentrations of C, N and P across 26 paddy fields in southeastern China. *Atmospheric Environment*, 164: 458–467.  
<https://doi.org/10.1016/j.atmosenv.2017.06.023>
- Whitbread, A., Blair, G., Konboon, Y., Lefroy, R. & Naklang, K. 2003. Managing crop residues, fertilizers and leaf litters to improve soil C, nutrient balances, and the grain yield of rice and wheat cropping systems in Thailand and Australia. *Agriculture, Ecosystems & Environment*, 100(2): 251–263.  
[https://doi.org/10.1016/S0167-8809\(03\)00189-0](https://doi.org/10.1016/S0167-8809(03)00189-0)
- Xiong, Z., Liu, Y., Wu, Z., Zhang, X., Liu, P. & Huang, T. 2015. Differences in net global warming potential and greenhouse gas intensity between major rice-based cropping systems in China. *Scientific Reports*, 5(1): 17774. <https://doi.org/10.1038/srep17774>
- Xionghui, J., Jiamei, W., Hua, P., Lihong, S., Zhenhua, Z., Zhaobing, L., Faxiang, T., Liangjie, H. & Jian, Z. 2012. The effect of rice straw incorporation into paddy soil on carbon sequestration and emissions in the double cropping rice system. *Journal of the Science of Food and Agriculture*, 92(5): 1038–1045.  
<https://doi.org/10.1002/jsfa.5550>
- Zhang, A., Cui, L., Pan, G., Li, L., Hussain, Q., Zhang, X., Zheng, J. & Crowley, D. 2010. Effect of biochar amendment on yield and methane and nitrous oxide emissions from a rice paddy from Tai Lake plain, China. *Agriculture, Ecosystems & Environment*, 139(4): 469–475.  
<https://doi.org/10.1016/j.agee.2010.09.003>
- Zhu, L., Hu, N., Yang, M., Zhan, X. & Zhang, Z. 2014. Effects of Different Tillage and Straw Return on Soil Organic Carbon in a Rice-Wheat Rotation System. *PLOS ONE*, 9(2): e88900.  
<https://doi.org/10.1371/journal.pone.0088900>

# 18. Selection of rice varieties adapted to salinity

Sara Ibáñez Asensio, Héctor Moreno-Ramón

Plant Production Department, Universitat Politècnica de València, Valencia, Spain

## 1. Description of the practice

Rice is classified as a relatively salt-sensitive crop (FAO, 1988) and several studies have estimated that rice yield decreases by 50 percent in salt-affected soil (when the electrical conductivity (EC) reaches 6.9 dS/m) (Grattan *et al.*, 2002). When the water-soluble salt content of the soil exceeds a certain threshold value, salinity degrades the soil physical and chemical properties, affecting plant growth. The effect occurs at the osmotic, oxidative and ionic levels, and results in the interruption of the metabolic functions of the cell due to ionic toxicity, as well as the appearance of psychopathologies in roots, leaf and fruit, among others (Flowers, 2004; Singh *et al.*, 2016).

Decrease in yield entails a lower biomass production and in theory a lower carbon immobilization in the soil. Faced with this situation, farmers tend to increase the inputs in chemical fertilizers and organic amendments, generally increasing Global Warming potential (GWP) (Wang *et al.*, 2012; Yuan *et al.*, 2018). A good alternative to maintain soil-plant-atmosphere balance are rice varieties that are more resistant to salinity conditions, making it one of the most important lines in the fight against the loss of rice production in areas affected by this type of land degradation. The new rice cultivars developed show a wide range of variations in salt tolerance.

Pokkali, Getu, Nona Brokra, Cherireruppu, FL478, CSR13, CSR43, PSBRC50, BRRI Dhan 54, SR86, IR65192-4B-10-13 are some of the most tolerant cultivars for white rice, whereas for black rice, Niewdam Gs.no.00621, Niewdam Gs.no.21629 and KKU-LLR-065 are the most salt-tolerant varieties. Some of these cultivars have been developed by genetic engineering, as in the case of Pokkali, a traditional variety that has been used as a donor in many breeding programs (e.g. FL478 is a cross between Pokkali and IR 29).

## 2. Range of applicability

Coastal wetlands in arid and semiarid environments are the main areas at risk of soil salinization due to water scarcity, poor quality water inputs or the presence of a saline water table near the soil surface. These situations cause a great concentration of salts in root area of soils and therefore can affect rice production. Although this kind of issue is mainly related to arid and semiarid areas, it can appear in other areas.

In coastal areas (deltas, plains coast, coastline, and lagoons), the interaction between freshwater and seawater intrusion can be a source of salts in soil. Farmers in general should avoid bare soil in paddy fields, because evapotranspiration can produce salts that rise from deep zones and appear on soil surface, affecting plants. Frequently, management of the sheet of water over the rice fields triggers intrusion of seawater since the loading of the sheet of water is non-existent. Due to these situations, the water table rises and causes losses in rice production, affecting it in the vegetative development. Germination stage is considered as the most tolerant period, while seedling and reproductive stages are the periods where the greatest saline stress occurs (Castillo *et al.*, 2007; Ebrahimi *et al.*, 2011).

In general, salinization is related to geographical position or climate, but land management can end up salinizing fields that initially did not present such problems. Water scarcity is one of those situations and has to do with poor quality water inputs. Irrigation time, electrical conductivity and the amount of water are factors that must be controlled by farmers. The accumulation of salts in soil profile has different treatments in rainfed agriculture if we compare with rice in flooding conditions. To avoid soil salinization and loss of rice yield, a higher dose of water irrigation can be applied to wash salts and extract them through drains from the root area.

Finally, salinization problems may be associated with soil type or original rock of rice fields. If salinity problems are defined by geological material the problem can be complicated to manage because the problem is in the origin of the soils. In addition, rice paddies are also normally found in places with low gradient or low transmissivity aquifers, so artificial underground drainage is not possible. At the end, a recurring option nowadays is the use of resistant varieties, or those genetically modified based on local species.

## 3. Impact on soil organic carbon stocks

General data of SOC and soil salinity can be found in several studies, but there is no study about the SOC production according to paddy varieties (resistant and non-resistant salinity).

Although no direct relationship can be evidenced, it is known that salinity affects the biogeochemical cycles that support C storage and therefore usually leads to a reduction of SOC sequestration due a low microbiological activity. For example, Morrissey *et al.* (2014) concluded that organic matter content decreased 21 percent when salinity changed from fresh to brackish (0.03-1.85 ppt) in the Chesapeake Bay in Virginia (United States of America). From this point of view, it is therefore reasonable to assume that using varieties that are less affected by soil salinity should help prevent this decline.

However, in contrast, sometimes microbial activities help to counteract this negative behavior of saline soils. Various studies conducted in coastal rice paddies have also revealed that increasing soil salinity has an inhibitory effect on organic carbon mineralization rates (Rao and Pathak, 1996; Weston, Dixon and Joye, 2006; Weston *et al.*, 2011, Vepraskas and Lindbo, 2012; Moreno-Ramón *et al.*, 2015; Luo *et al.*, 2019). In this way, this positive effect would be enhanced to the use of resistant varieties, resulting in an increase in SOC in the medium and long term.

Overall, environmental conditions and the degree of salinity of soils are the main factors that determine these behaviors.

## 4. Other benefits of the practice

### 4.1 Minimization of threats to soil functions

In South Asia (alluvial Indo-Gangetic plains Ganges delta), sodic soils can be used for rice cultivation with salt-tolerant varieties (Singh *et al.*, 2016). It is therefore possible to use degraded soils with a combining cost-effective crop and nutrient management and maximize the productivity and profitability of sodic soils.

### 4.2 Increases in production (e.g. food/fuel/feed/timber)

The use of salinity-tolerant varieties improve yield in rice fields. Islam and Gregorio (2013) reveals that BRRI Dhan 54 in wet season in Bangladesh showed the highest yield (6 t/ha) versus IR77674-B-25-1-2-1-3-12-4-AJY, a tolerant variety, which registered the lowest value (4 t/ha). BRRI Dhan 41 developed the highest plant development (height) in wet season, although the yield was quite low (4.2 t/ha). In addition, for a dry season BRRI dhan47 (IR63307-4B-4-3) and BINA dhan8 (IR669463R-149-1-1) can tolerate soil EC 12 to 14 dS/m during the seedling stage and EC 6 dS/m during all the cycle. In that situation, tolerant varieties obtained a yield potential between 2.8 to 8.1 t/ha.

In the same trend, Singh *et al.* (2016) concluded that CSR43 variety produced about 0.5 t/ha additional grain yields over current varieties in sodic soils located in the alluvial plains of India and in the salt-affected areas of the Ganges delta.

### 4.3 Mitigation of and adaptation to climate change

Currently, there are no specific data on how the implementation of salinity-resistant varieties affects greenhouse gas emissions. There is no specific study comparing tolerant and non-tolerant varieties under salinity and in the same conditions (soil, climate, management, etc.). In this sense, to determine the effect of salt-tolerant and non-tolerant rice varieties on GWP associated only with the plant and its metabolic processes it would be necessary more specific studies. Rice paddy greenhouse gas emission data exist and it is evident in the other sections of

this technical document, but there is a clear lack of data in the case of the emission of tolerant varieties versus non-tolerant ones.

In general, a multitude of evidence indicates that the addition of chemical fertilizers and rice straw with traditional varieties can double the gas emissions (CH<sub>4</sub> and N<sub>2</sub>O). In that regard, the farmer's aim will be to obtain the minimum incidence of toxicities or nutritional deficiencies in rice plants, in order to avoid the use of fertilizers or the incorporation of carbon (straw) because both worsen the forecasts on global warming. The magnitude of these increases can range depending climatic and management conditions. For example, in the case of rice flooding conditions with incorporation of straw and conventional tillage, emissions increase between 108 to 180 percent compared to no straw incorporation and tilling (Xiong *et al.*, 2015). On the other hand, Sui *et al.* (2016) found 60 percent more emissions with the incorporation of straw only, being around 34 percent when only urea is added (Bhattacharyya *et al.*, 2012).

The use of salinity-resistant varieties as a way of maintaining the yield of rice at optimum levels without the need to increase fertilization or the incorporation of straw, therefore seems a good option to avoid increasing the emission of greenhouse gases

## 5. Potential drawbacks to the practice

The use of genetic modified varieties or other salt-tolerant varieties can displace local varieties and the biodiversity can be reduced.

## 6. Potential barriers to adoption

**Table 70.** Potential barriers to adoption

Barrier	YES/NO	
Cultural	Yes	Reticence to abandon the use of traditional varieties in the area.
Economic	Yes	The increase of production costs due to the rise in the price of rice seeds of new varieties.
Knowledge	Yes	Ignorance of the most appropriate management for new varieties.

## References

- Bhattacharyya, P., Roy, K.S., Neogia, S., Adhya, T.K., Rao, K.S. & Manna, M.C.** 2012. Effects of rice straw and nitrogen fertilization on greenhouse gas emissions and carbon storage in tropical flooded soil planted with rice. *Soil and Tillage Research*, 124: 119-130. <https://doi.org/10.1016/j.still.2012.05.015>
- Castillo, E., To Phuc, G., Abdelbaghi, M.A. & Kazuyuki, I.** 2007. Response to salinity in rice: comparative effects of osmotic and Ionic stress. *Plant Prod. Sci.*, 10(2): 159-170. <https://doi.org/10.1626/pps.10.159>
- Ebrahimi H., Ared, F., Rezai, M., Amin, E. & Khaledin, M.R.** 2011. The effects of salinity at different growth stage on rice yield. *Ecology, Environment & Conservation Paper*, 17(2): 111-117.
- Grattan, S.R., Zeng, L., Shannon, M.C. & Roberts, S.R.** 2002. Rice is more sensitive to salinity than previously thought. *California Agriculture*, 56: 189-195.
- FAO.** 1988. *Salt-Affected Soils and their Management*. FAO soils bulletin 39. Food and Agriculture Organization of the United Nations. Rome, 1988. <http://www.fao.org/3/x5871e/x5871e00.htm>
- Flowers, T.J.** 2004. Improving crop salt tolerance. *Journal of Experimental Botany*, 55(396): 307-319. <https://doi.org/10.1093/jxb/erh003>
- Islam, M.R. & Gregorio, G.B.** 2013. Progress of salinity tolerant rice variety development in Bangladesh. *SABRAO Journal of Breeding and Genetics*, 45(1): 21-30.
- Luo, M., Huang, J.-F., Zhu, W.-F. & Tong, C.** 2019. Impacts of increasing salinity and inundation on rates and pathways of organic carbon mineralization in tidal wetlands: a review. *Hydrobiologia*, 827(1): 31-49. <https://doi.org/10.1007/s10750-017-3416-8>
- Moreno-Ramón, H., Marqués-Mateu, A., Ibáñez-Asensio, S. & Gisbert, J.M.** 2015. Wetland soils under rice management and seawater intrusion: characterization and classification. *Spanish Journal of Soil Science*, 5(2): 111-129. <https://doi.org/10.3232/SJSS.2015.V5.N2.02>
- Morrissey, E.M., Gillespie, J.L., Morina, J.C. & Franklin, R.B.**, 2014. Salinity affects microbial activity and soil organic matter content in tidal wetlands. *Global Change Biology*, 20: 1351-1362. <https://doi.org/10.1111/gcb.12431>
- Rao, D.L.N. & Pathak, H.** 1996. Ameliorative influence of organic matter on biological activity of salt-affected soils. *Arid Soil Research and Rehabilitation*, 10(4): 311-319. <https://doi.org/10.1080/15324989609381446>
- Singh, Y.P., Mishra, V.K., Singh, S., Sharma, D.K., Singh, D., Singh, U.S., Singh, R.K., Haefele, S.M. & Ismail, A.M.** 2016. Productivity of sodic soils can be enhanced through the use of salt tolerant rice varieties and proper agronomic practices. *Field Crops Research*, 190: 82-90. <https://doi.org/10.1016/j.fcr.2016.02.007>
- Sui, Y., Gao, J. Liu, C., Zhang, W., Lan, Y. Li, S., Meng, J., Xua, Z. & Tang, L.** 2016. Interactive effects of straw-derived biochar and N fertilization on soil C storage and rice productivity in rice paddies of Northeast



China. *Science of the Total Environment*, 544: 203–210.

<https://doi.org/10.1016/j.scitotenv.2015.11.079>

**Vepraskas, M.J. & Lindbo, D.L.** 2012. Redoximorphic features as related to soil hydrology and hydric soils. In Lin, H (Ed.) *Hydropedology: Synergistic Integration of Soil Science and Hydrology*. Whatlham, Academic Press, Elsevier. pp. 143-172.

**Wang, J., Chen, Z., Ma, Y., Sun, L., Xiong, Z., Huang, Q. & Sheng, Q.** 2013. Methane and nitrous oxide emissions as affected by organic–inorganic mixed fertilizer from a rice paddy in southeast China. *Journal of Soils and Sediments*, 13(8): 1408–1417. <https://doi.org/10.1007/s11368-013-0731-1>

**Weston, N.B., Dixon, R.E. & Joye, S.B.** 2006. Ramifications of increased salinity in tidal freshwater sediments: Geochemistry and microbial pathways of organic matter mineralization. *Journal of Geophysical Research: Biogeosciences*, 111(G1). <https://doi.org/10.1029/2005JG000071>

**Weston, N.B., Vile, M.A., Neubauer, S.C. & Velinsky, D.J.** 2011. Accelerated microbial organic matter mineralization following salt-water intrusion into tidal freshwater marsh soils. *Biogeochemistry*, 102: 135–151. <https://doi.org/10.1007/s10533-010-9427-4>

**Xiong, Z., Liu, Y., Wu, Z., Zhang, X., Liu, P. & Huang, T.** 2015. Differences in net global warming potential and greenhouse gas intensity between major rice-based cropping systems in China. *Scientific Reports*, 5(1): 17774. <https://doi.org/10.1038/srep17774>

**Yuan, J., Yuan, Y., Zhu, Y. & Cao, L.** 2018. Effects of different fertilizers on methane emissions and methanogenic community structures in paddy rhizosphere soil. *Science of The Total Environment*, 627: 770–781. <https://doi.org/10.1016/j.scitotenv.2018.01.233>


# 19. Integrated rice-based farming systems

Prafulla K. Nayak, Amaresh K. Nayak, Bipin B. Panda

ICAR, National Rice Research Institute, Cuttack, Odisha, India

## 1. Description of the practice

Rice is the staple food of about 50 percent of the global population and is cultivated on 160 million ha (Mha) mainly concentrated in Asia (China, India, Indonesia, Bangladesh, Vietnam, Thailand, Myanmar, Pakistan, Philippines, the Democratic People's Republic of Korea and Japan). Rice cultivation constitutes the vital source of income of about 140 million of rice-farming households and rural poor (Pathak, Samal and Shahid, 2018). Rice production is intricately linked with water and land ecosystems, and its intensification in the last decades led to soil, water and environmental degradations and increased greenhouse gases emissions (GHG), reducing its societal benefits (Pathak, Samal and Shahid, 2018; Kopittke *et al.*, 2019). As a response, rice-based integrated farming systems (RIFS) aim to combine rice cultivation with diverse practices such as livestock, aquaculture, agroforestry, agri-horticulture, beekeeping, mushroom, vermicomposting and/or other crops including pulses and cereals. In such systems, the synergies, mutualism and by-products generated from one component become potential inputs for others (Figure 8) (Hu *et al.*, 2016; Bashir *et al.*, 2020; Nayak *et al.*, 2020a).



**Figure 8.** Pathways contributing to resource flows (i.e. from one enterprise to other) portraying their synergy and mutualisms conducive to enhance family foods and income

RIFS include a large diversity of combinations of practices that mostly depend on various factors, such as the size of land holding, the facilities of obtaining credits, potential marketing, awareness among the stakeholders including their level of training and knowledge, among others. The various combinations of RIFS are categorised under different schemes:

### A. Rice–crop–animal husbandry systems

These systems include crops as cereals or pulses. In conventional rice-wheat and/or rice-pulses cropping systems, crop residues are usually dumped, thrown or burnt, leading in the long-term to soil degradation such as nutrient imbalances or the increased use of fertilizers. The inclusion of an animal component allows to make use of the crop residues that serve as animal feed, while the animals can also be used for traction. The animal waste (e.g. dung and slurries) is returned to fields leading to improvements in soil health and fertility (Bhatt *et al.*, 2016; Adarsh, Jacob, and Giffy, 2019). In between fallow periods, fodder crops can be taken up. In addition, some parts of the land (preferably the irrigated ones) can be permanently assigned for fodder and forage crop cultivation. The introduction of pulses (such as green gram (*Vigna radiata*), black gram (*Vigna mungo*), lentil (*Lens culinaris*), horse gram (*Macrotyloma uniflorum*) or chickpea (*Cicer arietinum*), either as rotational,

intercrops, relay crops or cover crops in a rice-pulse cropping system supports the improvement of soil health and provides a better feed for animals.

## B. Rice-aquaculture systems

These systems support the means of livelihoods of millions of smallholder farmers worldwide (Halwart and Gupta, 2004; FAO, 2019a). Many forms of integrated rice-aquaculture systems exist (Nayak *et al.*, 2020a; Bashir *et al.*, 2020). These are categorised as:

**B1.** Rice-fish systems (e.g. rice-cum-fish, rice-fish-vegetables, rice-fish-livestock, Photo 22), where fish is grown together with or in alternance with rice cultivation. Depending on the regional availability and consumers demand, the main fish species used are the Indian major carp, the exotic carp, the common carp, tilapia, silver carp, minor carps, nutritionally important small fish and also crabs, shrimps, crayfishes or catfishes (Halwart and Gupta, 2004; Hu *et al.*, 2016; FAO, 2019a; Nayak *et al.*, 2020a). Besides fish, various vegetables (dyke or main fields) and animal components (livestock or poultry) can be introduced. In these systems, fish is placed in the rice fields only after the establishment of rice plants (in dry seeding rice: after attaining six inches growth of rice plants, and transplanted rice: mostly after 20 days of planting to avoid damage to rice plants); and for a period of 7-10 months or more, in the case where ridges and water refuges are created for water storage. The harvested pond water can also be used for irrigation in addition of fish culture. The application of organic manures (cow dung and fertilizer) stimulates the natural growth of fish food organisms (e.g. planktons, benthos) that support fish growth. The introduction of *Azolla* causes twin benefits, as it can be used as feed for fish and as a nitrogen source for rice.



**Photo 22.** Depicting an improved version of rice-aquaculture system including a livestock component

This was enabled after a land reshaping to create wide bunds (or dyke) of 2-4 m wide all around the site. The pond (or water refuge) is connected with two sides trenches. Rice cultivation covers 65 percent of the total area

**B2.** Rice-fish-agroforestry-horticultural-duck/poultry systems (multitier farming systems): these systems include the use of improved rice varieties, vegetables, tuber crops (*Amorphophallus*, *Yam*, *Colocasia*, *Turmeric*, *Ginger*), fruit crops (e.g. papaya, coconut, arecanut, banana, guava, mango), fodder (e.g. napier, gunia grass, legume fodder, cowpea/lobia), agroforestry (e.g. *Acacia mangium*, *A. auriculiformis*, *Eucalyptus globulus*), floriculture, apiculture along with animal components (e.g. fish, prawn, poultry, duckery, goatry) and additional activities (e.g. beekeeping, mushroom cultivation) (FAO, 2019a; Nayak *et al.*, 2020a) (Figure 9).




Figure 9. Multitier rice based integrated farming system

The land shaped creates the rice ecosystem of an upland (Tier I and Tier II, 15 percent of the field area), rainfed lowland (Tier III, 20 percent, up to 50 cm water depth), deep water (Tier IV, 20 percent, up to 50 - 100 cm water depth), micro-watershed (20 percent area) and raised wide bund (25 percent) surrounding the entire fields area. Different components as rice, fish and prawn, dry season crops, horticultural plants and agroforestry components need to be suitably cultivated in different tiers of land. The duck and poultry houses are constructed on the bund having projection to facilitate dropping fall directly in the pond water. The goat house is constructed on the bund using bamboo, wood and wire net with straw thatching or asbestos top. Source: Authors' own compilation and analysis

**B3.** Rice-cum duck farming/poultry farming (Photo 23; Figure 10, Figure 11 and Figure 12) (Nayak *et al.*, 2018; Nayak *et al.*, 2020a; Li *et al.*, 2019). The main used duck species are Khaki campbell (egg layer) or White pekin (meat type), but the use of local species more adapted to the diversity of regions is encouraged.




Figure 10. Symbiotic relationship between rice-fish-duck yielding maximum mutual benefits to all the entities

Source: Authors' own compilation and analysis




Figure 11. Framework of potential mutualism and synergies among rice-fish-duck-Azolla based co-culture

Azolla is a free-floating aquatic fern widely distributed in tropical India, and has nitrogen fixing capabilities through symbiotic cyanobionts (around 1100 kg N/ha/year) considered as one of the best bio-fertilizer for rice plants and feed for livestock. Source: Nayak *et al.* (2020a)




Figure 12. Pictorial views of operational periodicities of rice, fish and ducks in rice-rice cropping system

The figure indicates periods of foraging and restrictions in foraging of fish and ducks within the rice fields to protect rice plants from potential damage during foraging i.e. 20 days after rice plants establishments, during rice flowering to harvest (30 days only for ducks) in rice-rice cropping systems. Strict implementation of foraging restrictions is one of the characteristic features for rice-fish-duck co-culture technology



Photo 23. Duck foraging in dry seeded rice fields creating a conducive environment for initial (after 20 days of rice plant establishment) control of weeds and pests, duck dropping for fertilization and loosening of soil for better root growth and subsequent growth of rice plants (left). Rice-fish-duck co-culture system with luxuriant growth of rice (right)

**B4.** Pond-dike farming systems (Photo 24): Their type mostly depends on the type of dyke planted crops (e.g. mulberry dyke-fishpond, sugar cane dyke-fishpond, banana dyke-fishpond, vegetable dyke fishpond). Input and output materials are suitably recycled, and the system energy flows are fundamentally balanced. For example, in integrated mulberry dyke-fishpond systems, the mulberry leaves are used to feed silkworms, the generated by-products from silkworm are used as feed for fish and the fertile mud (bottom of pond) fertilizes the mulberry plants at dyke. The systems can be suitably integrated with raising of vegetables, livestock, and other components (Karim and Little, 2018; Babu *et al.*, 2019).



Photo 24. Pond-dyke farming system

**B5.** Plant/fish rotations (forage or compost): where rice grows in rotation with fish. After the rice harvest (i.e. during fallow periods), fish are introduced for three to six-month. For example, in Indonesia *sawah tambak* (rice field pond brackish water) is very popular in the coastal Java region. Fish are grown as if they were a second crop (*palawija ikan system*) during the fallow-season after rice harvest, by using a hoe (water depth of 30–40 cm). In the western coastal region of India (low-lying coastal rice lands), after rice harvest (in September) rice fields are flooded with tidal water and used to raise shrimps (Halwart and Gupta, 2004).

**B6.** New emerging systems (e.g. rice-crayfish, rice-snail or rice-crab): In brackish and freshwater rice-rice systems shrimp cultivation (e.g. *Penaeus monodon*, *Penaeus merguensis*, *Penaeus indicus*, *Metapenaeus ensis*; *Macrobrachium rosenbergii* in freshwater) are gaining importance. Among the rice-crab co-culture, freshwater crabs (*Oziotelphusa senex senex* or *Parathelphusa hydrodromus*), brackish water crab like the Chinese mitten crab (*Eriocheir sinensis*) and mud crab (*Scylla serrata*, cultured throughout the world) are used. In rice-crayfish co-culture the two common species are *Procambarus clarkii* and *Cherax quadricarinatus*. During the rice harvest, the rice field water is drained or receded, to help crayfish to make burrows for shelter. After harvest, the re-growth of rice stubbles serves as food for crayfish. Further, watering rice stubbles decomposes and facilitates the growth of fish food organisms (i.e. planktons, benthos, insects, worms and mollusks) which helps the growth of crayfish (Si *et al.*, 2017; FAO, 2019a. Although the Golden apple snail (*Pomacea canaliculate*) is considered as a major pest to rice, rice-snail co-culture is used, as snail meat is used as a cheap source of feed in the culture of prawn, shrimp and rabbitfish (Visca and Palla, 2018).



## 2. Range of applicability

**Table 71.** Rice based integrated farming system prevalent in different part of the world

Type of RIFS	Sub-type of RIFS
Rice-crop-animal husbandry systems	<b>Rice-wheat-animal husbandry:</b> Applicable in South and East Asian countries, Sub Saharan Africa and South America but mostly concentrated in tropical and sub-tropical climatic conditions (Dixon, Gibbon and Gulliver, 2001).
	<b>Rice-pulses-Animal husbandry:</b> Prominent farming system in rainfed areas across the tropical and sub-tropical regions. The rice-pulses and animal husbandry systems are prevalent in rainfed upland and lowland areas (Dixon, Gibbon and Gulliver, 2001; Erenstein <i>et al.</i> , 2007).
Rice-aquaculture integration: Traditionally in Asian countries but adapted worldwide, even in deserts and arid lands like Egypt or Oman, thanks to the efficient use of water.	<b>Rice-Fish integration:</b> Already practised in the ancient India and China, and later adopted in most of the Asian countries (Bangladesh, Vietnam, Indonesia, Philippines, Malaysia, Thailand and Japan). These systems are mostly suitable in lowland rice area including coastal areas, where water retain in the fields even after rice harvest (Halwart and Gupta, 2004; Lu and Li, 2006; Hu <i>et al.</i> , 2016). Under plain, medium lowland and rainfed conditions, rice and fish can be grown at the same time, while in deep water and coastal lowland areas, fish either grown simultaneously with rice and/or also with off season fish rearing and seed raising (FAO, 2019a; FAO-SHOU, 2020). In hilly regions, rice-fish integration depends on suitable designs of counter bunding and water storage ponds. The traditional system of rice-fish co-culture has evolved with time, with the introduction of higher economically important aquatic species or to integrated-aquaculture and agriculture systems.
	<b>Rice-fish-livestock-horticultural-duck/poultry multitier system:</b> Suitable for all kinds of rice ecologies (i.e. upland, medium lowland and deep-water ecologies).

Type of RIFS	Sub-type of RIFS
	<b>Rice-livestock-horticultural and agroforestry-based IFS:</b> Mostly prevalent in Asia, Europe, South America and Africa. Suitable to medium deep or deep-water lowlands, free from heavy flooding having clay soil and with prolonged water retention capacity (Nayak <i>et al.</i> , 2020a)
	<b>Rice-fish-duck integration:</b> Mostly prevalent in Asia and African countries. Well adapted to medium deep- or deep-water lowland rice ecologies, free from heavy flooding and having clay soil and with prolonged water retention capacity (Pernollet <i>et al.</i> , 2015; Nayak <i>et al.</i> , 2018; Nayak <i>et al.</i> , 2020a; Li <i>et al.</i> , 2020).
	<b>Pond-dyke farming system:</b> Mostly prevalent in Asia, Africa and South America region (Gongfu, 1990; Babu <i>et al.</i> , 2019).
	<b>Plant/fish rotations (forage or compost):</b> Popular in South Asia including China, India, Indonesia and Cambodia. This type of rice-fish systems is very popular in low-laying coastal areas of Indonesia and India (Halwart and Gupta, 2004).
	<b>New emerging systems (e.g rice-crayfish, rice-snail or rice-crab):</b> Practised in Asia, Australia and the United States of America.

### 3. Impact on soil organic carbon stocks

Suitable combination of system components (crop, aquaculture, livestock, agroforestry, and horticultural components) in a rice-based integrated system contribute positively to SOC sequestration (Oliveira *et al.* 2018; Nayak *et al.* 2018; Li *et al.* 2019). Some examples of rice based integrated systems that show an enhancement of soil carbon stocks are mentioned in Table 72.

**Table 72.** Changes in soil organic carbon stocks reported for rice based integrated farming systems

Rice-based IFS	Location	Climate Zone	Soil type	Baseline C stocks (tC/ha)	Additional C storage (tC/ha/year)	Duration (Years)	Depth (cm)	More information	References
ICLF tree-based farming system	South Brazil 10°38'13" S, 55°42'32" W	Tropical Moist	Kaolinitic oxisol	Pasture = 16.5	Pasture = 1.37 ICLF = 1.91	12	0-30	IFS with trees promotes SOC accumulation, even on short periods (3 yrs), if there is no soil fertility constraint (N deficiency).	Oliveira <i>et al.</i> (2018)
ICL and ICLF	South-East Brazil 21°57'42" S 47°50'28" W	Tropical Moist	Oxisol	Extensive grazing = 1.45 ICL = 1.48 ICLF = 1.55	Extensive grazing = 1.68 ICL = 1.96 ICLF = 1.74	6	0-40	Land intensification increases C stocks however, converting pasture (extensive grazing) to ICL and ICLF increases soil C stocks at the rate of 0.28 MgC/ha/yr	Bieluczyk <i>et al.</i> (2020)
Rice-fish IFS	Cuttack, Odisha, India	Tropical Moist	Aeric Endoaquept sandy clay loam	Rice monoculture = 0.15	0.18	4	0-15	Increase in C stock over rice monocropping.	Nayak <i>et al.</i> (2018)
Rice duck IFS				0.23					

Rice-based IFS	Location	Climate Zone	Soil type	Baseline C stocks (tC/ha)	Additional C storage (tC/ha/year)	Duration (Years)	Depth (cm)	More information	References
Rice-fish-duck IFS					0.30			Increase in C stock over rice monocropping. Enhanced soil fertility and biodiversity	
Rice duck IFS	South China, 23°14'N, 113°38'E	Tropical Wet	Sandy loam soil	0.18	0.25	1	0-15	Rice-duck system enhanced carbon stocks	Li <i>et al.</i> (2019)
Rice-crayfish IFS	Hubei Province, China	Tropical Moist		Rice monoculture = 0.20	0.27	10	0-10	The C fractions like Microbial biomass carbon, dissolve organic carbon and particulate organic carbon are also increased	Si <i>et al.</i> (2017)
Rice-carb co-culture	Liaoning province, China 40°51'N, 122°13'E	Tropical Moist	Heavy clay of alluvial origin	0.25 (conventional rice monoculture)	conventional rice crab culture = 0.29 organic manure rice-crab culture = 0.35	5	0-20	Enhancement of bacteria contribution to SOM turnover	Yan <i>et al.</i> (2014)

ICLF: integrated crop-livestock-forestry system; ICL: integrated crop-livestock system; IFS: integrated farming system; SOC (t/ha) = SOC x BD x Depth cm x 10<sup>-1</sup>, where BD, bulk density (g/cm<sup>3</sup>), SOC (g/kg), soil organic carbon;

The SOC sequestration rate was calculated by dividing the changes/accumulation of SOC stock respect to treatment/establishment by the number of years i.e. SOC sequestration rate (t/ha/year) = SOC stock change/ accumulation/ storage (tC/ha) / nos. of year.

## 4. Other benefits of the practice

### 4.1 Improvement of soil properties

Improvements in land use managements include applying compost, conservation agriculture including cover crop, crop rotation, perennial crops, minimum or zero tillage and inclusion agroforestry and fodder crops with practices of rice-animal co-culture. These practices potentially lead to increased build up of soil organic matter, SOC and available nitrogen, phosphorus and potassium contents, which subsequently influence physico-chemical properties: for instance, in the long term, an increase of clay content of soil, possibly due to augmentation of rate of biological weathering (Bot and Benites, 2005; Teng *et al.*, 2016; Nayak *et al.*, 2018; Li *et al.*, 2019). Higher SOM reduces the bulk density and readily dispersible clay content and increases microbial activity (Gajda, Czyż and Dexter, 2016). The continuous addition of excreta of livestock components enhances SOC and available nitrogen, phosphorus and potassium content, and additionally, introduction of *Azolla* contributes substantial amount of nitrogen fertilizers for rice growth (Nayak *et al.*, 2020a) Long-term rotation of rice-shrimp farming leading to improvements in soil's physical and chemical properties including the upsurging of soil nutrients (Cai *et al.*, 2019). Duck activities enhances aeration (bioturbations). The use of leguminous crops increases available nitrogen and soil organic carbon content (Erenstein *et al.*, 2007; Adarsh, Jacob, and Giffy, 2019).

### 4.2 Minimization of threats to soil functions

**Table 73.** Soil threats

Soil threats	
<b>Soil erosion</b>	The diversity of practices included in RIFS reduces and protects soil from erosion, through increased cover crops areas and time periods, cultivation of perennial forage and other perennial crops, and site-specific inclusion of components as e.g. hedges, ponds, ditches, trees, agroforestry, suitable management of livestock and farm residues, addition of organic manure; management practices as e.g. terrace management, increased use of conservation agriculture and reduced tillage including living-plant windbreaks. Further, the enhancement of SOC increases soil aggregation stability which prevents erosion processes (Bots and Benites, 2005; FAO, 2019b).
<b>Nutrient imbalance and cycles</b>	Rice-fish, rice-duck, rice-fish-duck, crop-livestock-agroforestry and horticultural systems improve soil nutrients (NPK), enhance SOC build up and enhance nutrient recycling with augmentation of microbial diversities (Nayak <i>et al.</i> , 2018; Li <i>et al.</i> , 2019; Masciandaro <i>et al.</i> , 2018). Crops rotation and diversification, use of <i>Azolla</i> , and the addition of organic manures increase the soil available nitrogen levels and enhance the N use efficiency. In lowland ecology, rice-fish-duck co-culture leads to enhancement of total nitrogen (121 percent), available nitrogen (50 percent), available phosphorus (67 percent) and potassium (150 percent) respectively, compared to conventional rice farming (Nayak <i>et al.</i> , 2018).

Soil threats	
<b>Soil salinization and alkalization</b>	The maintenance of a suitable soil moisture, adequate drainage and increased SOC and aggregate stability prevent from salinization and alkalization. RIFS provisioned for increased use of organics and FYM leading to reduction in the root zone accumulation of salt (Kaledhonkar, Meena and Sharma, 2019).
<b>Soil contamination / pollution</b>	RIFS lead to optimized and reduced application of synthetic fertilizers, pesticides and herbicides which ultimately prevents soil and water contamination and enhanced the water quality (Long <i>et al.</i> , 2013; Nayak <i>et al.</i> , 2020a).
<b>Soil acidification</b>	Judicious and reduced rates of application of agricultural chemicals and high soil organic matter prevent soil acidification (Long <i>et al.</i> , 2013). Amendments like lime, organic and farmyard manure mostly used in RIFS resulting increase in pH leading to amelioration of acid sulphate soil (Halim <i>et al.</i> , 2018).
<b>Soil biodiversity loss</b>	Higher SOC stocks enhance soil fauna, flora and microbial population. RIFS increase soil biodiversity and biological soil quality index (SQI <sub>biol</sub> ) as compared to conventional farming (Kremen, Iles and Bacon, 2012; Nayak <i>et al.</i> , 2020a).
<b>Soil water management</b>	The retention of available moisture and an effective water drainage system when saturated leads to higher water use efficiency (Ahmed, Ward and Saint, 2014).

### 4.3. Increases in production (e.g. food/fuel/feed/ timber)

RIFS increase water use efficiency and system productivity (Nayak *et al.* 2018; Li *et al.* 2019; FAO, 2019a), since it is a multi-enterprising farming system where the substantial part of food, fuel and fibre requirements are provisioned from the system itself. The potential enhancements of productivity from co-culture have been reported from many countries (Bashir *et al.*, 2020), such as China where rice with fish, turtle, crayfish and crabs produce higher average production (9.3-12.0 t/ha of rice and 1.9-2.5 t/ha of fish) (Zhang *et al.*, 2016); Bangladesh where rice-fish (shrimp, prawn, fish) yielded 3.8-5.0 t/ha rice and 1.8 t/ha of fish (Islam, Barman and Murshed-e-Jahan, 2015; Ahmed, Ward and Saint, 2014); India, rice-fish (fish, prawn, crabs) integration produces 3.0-5.0 t/ha rice and 0.7-2.0 t/ha fish (Das, Sarkar and Prasad, 2014; Nayak *et al.*, 2018; Nayak *et al.*, 2020a); Indonesia where rice-fish integration produces 6.5-7.8 t/ha rice and 0.3-0.89 t/ha fish (Dwiyanana and Mendoza, 2006); Vietnam yielded 4.2-5.7 t/ha rice and 2.2 t/ha fish (Berg *et al.*, 2017); and the African continent (Melaku and Natarajan, 2019).

## 4.4 Mitigation of and adaptation to climate change

CH<sub>4</sub> is a potent greenhouse gas (28 times more than CO<sub>2</sub>) emitted from submerged rice ecosystems due to the anaerobic degradation of organic matter by the soil microorganism (IPCC, 2007). When rice cultivation is integrated with aquatic animals (e.g. fish, shrimp, shellfish, crayfish, crab, turtle, frog and ducks), bioturbations (paddling, scooping, trampling of soil and water) and foraging activities (resulting in reduction of weeds and aquatic organisms leading to a decline in oxygen demand for their respiration) occur. These activities enhance the available dissolved oxygen levels, which leads to better soil and water aerations, and results in an acceleration of CH<sub>4</sub> oxidative processes (by methanotrophic bacteria) and a reduction of CH<sub>4</sub> emissions (Nayak *et al.*, 2018; Nayak *et al.*, 2020a; Nayak *et al.*, 2020b; Xu *et al.*, 2017; Zhang *et al.*, 2017). However, seasonal cumulative CH<sub>4</sub> emissions are potentially higher in co-culture systems (rice-fish, rice-duck and rice-frog) (Frei *et al.*, 2007; Datta *et al.*, 2009; Bhattacharyya *et al.*, 2013; Xu *et al.*, 2017; Fang *et al.*, 2019; Wang *et al.*, 2019); nevertheless, the GHG intensity estimated per unit output (expressed in rice equivalent yield) is much less. In rice monoculture, at some point of growing period the crop is subjected to alternate wetting and drying causing wet spell and dry spell but in rice-fish system the water remains standing throughout the growing period which explains the differential CH<sub>4</sub> emission. GWP potential is reduced in rice-fish-duck integrated farming, possibly, due to the reduced agri-chemicals/fertilization and better aerated environments within the paddy ecosystem (Nayak *et al.*, 2020a).

Paddy cultivation is also an important anthropogenic source of N<sub>2</sub>O (with a global warming potential 298 times higher than CO<sub>2</sub>). N<sub>2</sub>O emissions depend mostly on the intensity and methods of N fertilization (synthetic and organic), water management and drainage, and temperature. In addition to submerged anaerobic conditions, microbial functions as nitrification and denitrification can potentially be disturbed (IPCC, 2014; Wu *et al.*, 2018). Significant reduction of N<sub>2</sub>O emissions (Figure 13) from rice-animal co-culture (rice-fish, rice-crab, rice-crayfish, rice-duck, and rice-fish-duck) have been reported (Frei *et al.*, 2007; Datta *et al.*, 2009; Bhattacharyya *et al.*, 2013; Xu *et al.*, 2017; Fang *et al.*, 2019; Wang *et al.*, 2019; Nayak *et al.*, 2020a).

RIFS enhances the system's resilience and adaptive capacity and provides mutualism and flexibility to reduce trade-offs and competitiveness among the system's components, all of which offer adaptation options to overcome the vulnerability to climate-induced disturbances. In a four-year experiment on rice fish-duck co-culture, the SOC stocks doubled (+106 percent) and the GWP decreased by 11 percent compared to conventional rice farming at Cuttack, India (Nayak *et al.*, 2018; Nayak *et al.*, 2020 a). Different adaptation and mitigation strategies of RIFS are discussed in

Table 74 and Figure 13.



**Table 74.** Different climate change adaptation and mitigation options in rice-fish based integrated farming systems

Integrated system	Component	Climate change adaptation and mitigation option	Emission potentials
Integrated Rice-fish system	Crops	<ul style="list-style-type: none"> <li>Reduction in synthetic fertilizers, pesticides and herbicides</li> </ul>	Reduction in GHG emission
	Fish production	<ul style="list-style-type: none"> <li>Lesser feed requirements</li> <li>More efficient water use</li> <li>Ducks and fish bioturbation (rapid movement) and presence of <i>Azolla</i> in rice ecosystems enhance the concentration of dissolved oxygen in water, resulting in aerobic conditions, which decrease methanogens bacterial activity and subsequently decreases the GHG emissions.</li> </ul>	
Crop-livestock-agroforestry based IFS	Crops	<ul style="list-style-type: none"> <li>Organic manure used for plant growth.</li> <li>Reduction of synthetic chemicals (fertilizer and pesticides), their production, transport and application.</li> <li>Reduced area for feed crops with efficient land use.</li> <li>Recycling of crop residue, manure and nutrient recycling.</li> </ul>	Adoption and mitigation of global warming
	Livestock	<ul style="list-style-type: none"> <li>Quality feed for livestock (ruminants, pig, duck and poultry can eat crop residue and by-products) lower enteric methane emissions.</li> <li>Sound manure management reduces GHG emissions.</li> <li>Efficient use of land area management.</li> </ul>	
	Agro-forestry	<ul style="list-style-type: none"> <li>Inclusion of agroforestry component in RIFS significantly reduces the effects of global warming potential. Higher carbon sequestration in biomass and soil.</li> <li>Improved soil health and water infiltration and retention capacity.</li> <li>Fodder availability throughout the year.</li> <li>Improved thermal comfort, welfare, health and production of animals</li> </ul>	




Figure 13. Schematic view of CH<sub>4</sub> and N<sub>2</sub>O emissions from rice-based integrated farming systems and synergistic effects of duck and fish contributions towards reduction of global warming potentials

Source: modified and reconstructed from Xu *et al.* (2017) and Nayak *et al.* (2020a)

## 4.5 Socio-economic benefits

RIFS offer higher socio-economic benefits leading to improved livelihoods and socio-economic standing of the rural farmers (FAO, 2019a; FAO-SHOU, 2020).

Integration with animals (fish, shrimp, ducks, etc.) supports productivity and economic returns (Bashir *et al.*, 2020; Nayak *et al.*, 2018; Nayak *et al.*, 2020a). In the Lowland ecology of India, higher economic returns (expressed in terms of benefit cost ratio) were observed in rice-ornamental fish culture (2.5), crop-livestock-agroforestry system (2.9-3.4), rice-fish-duck (2.5-2.8), rice-fish-duck-*Azolla* (2.7-3.0), and multi-tier rice-fish horticultural system (2.0-2.5). The profit margins are mostly dependent on the integrated components and their effective managements (Nayak *et al.*, 2020a). Integrated aquaculture-agriculture (IAA) systems significantly increase farming households incomes (Ahmed, Ward and Saint, 2014), and rice-animal (rice-fish, rice-crab and rice-crayfish) co-culture plays a significant role in promoting rice ecosystem efficiency and enhancing farmers' incomes in many countries (FAO, 2019a). The rainwater harvesting model (land shaping IAA) and paddy-cum-fish culture contribute to enhanced livelihood security levels of farm families in terms of creation asset (i.e. farm pond, adoption of multiple cropping and aquaculture). It increases their resilience, productivity and income, generates employment, facilitates access to market price, extension services and institutions making them self-reliant and enhancing their social status (Kumaran *et al.*, 2020). Pond dyke integration provides better economic return as compared to rice monoculture (Karim and Little, 2018; Babu *et al.*, 2019). Eventually, the introduction of high value aquatic species generates higher economic return and profits, which in turn generates additional employment and is helpful in improving national economy.

Enhancing the quality of life (food security, balanced nutrition, employment generation and gender equity) along with preserving people's social and cultural needs is a challenging task in most developing countries. Pond and rice fields can help achieving several social benefits (Halwart and Gupta, 2004). Indeed, pond based IAA produce a year-round food production, deliver diversified healthy foods, generate employability, and address gender issues (women have equal level of resources access) (Halwart and Gupta, 2004; FAO, 2019a). Accumulation of farm wastes can create environmental problems, but RIFS relies on recycling of the waste generated in the system, thereby, helping in maintaining sanitation and environment safety in farm families and its surroundings. Crop-livestock-agroforestry based integrated farming generate additional employments (400-500-man days/ha/year) depending upon the extend and type of integration (Nayak *et al.*, 2020a). The two important public health vectors such as mosquitos (malaria and dengue fever) and snails (Schistosomiasis and liver cirrhosis or common liver fluke caused by *Fasciola hepatica*) used rice fields as a breeding ground which potentially caused health hazards for humans are naturally controlled by the adoption of rice-fish/animal co-culture (rice-fish, rice-duck) (Halwart and Gupta, 2004; Singh, 2011).

## 4.6 Other benefits of the practice

### Increased water use efficiency

The rice-aquatic animal co-culture enhances blue water use efficiency with intensifying and diversifying cropping pattern (Ahmed, Ward and Saint, 2014). In rainfed areas, IAA and rice-fish-duck integrated farming with rainwater harvesting and storage facilities allow to reuse water in dyke-farming or emergency lifesaving irrigation for other crops (Ahmed, Ward and Saint, 2014; Nayak *et al.*, 2020a).

### Increased nutrient recycling and biodiversity

Rice-animal (fish, duck or fish-duck) co-culture improve water quality. The addition of faecal matter, continuous movement and activities (scooping, churning and trampling of soil and water) of fish and ducks increase dissolved oxygen levels and increase the aquatic biological diversity, including planktons (phyto- and zoo-plankton), soil benthic fauna's and microbial populations (Halwart, 2008; Nayak *et al.*, 2018). The biodiversity index scoring of rice-fish, rice-fish-duck and crop-livestock-agroforestry integration system (in respect to planned vegetative richness, intensity of cropping, richness of land scape elements, microbial, plankton and benthic richness and livestock richness etc.) is significantly higher than in conventional systems (Nayak *et al.*, 2020a).

### Bio-control prospecting of weed and pests

Significant reduction in weed density and weed biomass with increase in weed control efficiency was observed in rice-fish, rice-duck and rice-fish-duck integrated farming. The weed biodiversity (species richness; Simpson's index) and species diversity (Shannon-Weiner index) declined significantly with increase in Pielou evenness community index<sup>6</sup> in rice-fish-duck integration, signifying highly diversified weed community composition with reduction of formerly dominant weeds (Nayak *et al.*, 2020b). The presence of fish and ducks

---

<sup>6</sup> Species evenness refers to how close in numbers each species in an environment is. Mathematically it is defined as a diversity index, a measure of biodiversity which quantifies how equal the community is numerically.

enhance bio-control efficiency of rice-insect pest (leaf roller, brown plant hopper, zig zag leaf hopper and stem borer etc.), and thus application of pesticides/herbicides can be reduced or avoided (Li *et al.*, 2019; Nayak *et al.*, 2020a).

### Energy efficient system

Intensification of agriculture with extensive use of chemical fertilizers and pesticides/herbicides and large-scale use of mechanized farming operations are progressively making modern agricultural practices becoming less energy efficient. The co-culture and/or mixed farming (crop-dairy-fish-poultry) is more energy efficient and uses more renewable energy as compared to conventional rice farming (Paramesh *et al.*, 2019; Nayak *et al.*, 2020a).

### Conservation of natural resources

The potential reduction in application of agri-chemicals (fertilizers and pesticides lesser 24 percent and 68 percent, respectively) in China (Xie *et al.*, 2015; Long *et al.*, 2013), supports greater diversity of aquatic flora and fauna that supports resources and ecosystems conservation (Halwart, 2008). Additionally, rice ecosystems provide habitat and breeding ground for many other aquatic species (Halwart and Gupta, 2004; Halwart, 2008).

### Enhancing knowledge and skills

RIFS supports the development of skills and aims to exploit the available resources (crops, livestock and genetic potentials) to make more resilient systems. Multi-enterprising systems require improved knowledge and skills in respect to their specific managements. Gender empowerment and capacity building are the main framework in RIFS and demands gender specific managements and knowledge skills (Halwart and Gupta, 2004).

## 5. Potential drawbacks to the practice

### 5.1 Trade-offs with other threats to soil functions

RIFS minimizes and prevent the soil threat, however, if not adopted and implemented properly sometimes leading to trade-offs to other soil threats.

**Table 75.** Soil threats

Soil threats	
<b>Soil erosion</b>	In heavy rainfall areas, possible risks of flooding leading to water flows-based soil erosion. In arid and semi-arid areas, there can be risks of wind-based soil erosion. Cover crops in RIFS to be suitably managed with plantation of annual and perennial crops to avoid erosion (Zhang <i>et al.</i> , 2011).

Soil threats	
<b>Nutrient imbalance and cycles</b>	In case of unsustainable management, possible negative effect on depletion of SOM, nutrients (mainly N, P, K, S and micronutrients) resulting in severe limitations to nutrient available forms and Cation exchange capacity (CEC) which reduce the water and nutrient use efficiency (Nayak <i>et al.</i> , 2018; Li <i>et al.</i> , 2019).
<b>Soil contamination / pollution</b>	Excessive use of inputs like water and agricultural chemicals may lead to soil and water contamination and pollution which degrades soil and water quality and biodiversity (Ongley, 1996). In RIFS, any chemical (insecticides, pesticides, herbicides, growth hormones, antibiotics) used in any component of the system, naturally reaches the other components and enhances bioconcentrations.
<b>Soil water management</b>	In RIFS, if soil and water are not correctly managed, there is a risk of soil erosion, leaching and waterlogging leading to soil acidification and loss of biodiversity.

## 5.2 Increases in greenhouse gas emissions

Adopting suitable manure management in crop-livestock integration reduces nutrient requirements (lesser import of agri-chemicals) and improves animal health and herd management (with efficient digestible feeds) ultimately, resulting in reduced GHG emissions (Soussana *et al.*, 2015; Mottet *et al.*, 2017).

## 5.3 Conflict with other practice(s)

RIFS needs to be properly implemented to avoid conflict with other agricultural practices:

- ◆ Selection of components, extent of integration and management practices should be carefully planned (compatibility and synergy), to avoid damage to other cultivated components of the RIFS. For example, in rice-animal (ducks) co-culture food shortage or scarcity in rice ecosystem, sometimes leads to predation and foraging of planted vegetables and other plants.
- ◆ Intensification of rice cultivation with high degree of mechanisation is affected due to implementation of RIFS.

## 5.4 Decreases in production (e.g. food/fuel/feed/timber)

RIFS has no negative impact on production and productivity, however, without judicious introduction of the different components, there is a risk of disturbing the balance and synergies of the system that may lead to competition between the different system's components.

## 5.5 Other conflicts

RIFS involving animals' co-culture may cause accumulation of pathogenic bacteria that may trigger human health hazards (Singh, 2011).

# 6. Recommendations before implementation of the practice

- ◆ The system's components selection must be based on the existence of mutualism and synergies. The choice of fish, livestock, poultry introduction or selection of crops must not contradict or be competitive.
- ◆ Rice-animal co-culture needs to be suitably fenced to protect animals from predation and prevent them to reach adjacent fields (may cause damage to other field crops) and escape from the field.
- ◆ Farmers need to understand the concept and management practices of each combination (crops, animals). The system should be planned before shaping land. It is necessary to understand precisely the amount of manure needed in order to improve yields, while preserving water quality and avoiding water contaminations that may be harmful.
- ◆ The implementation of RIFS initially needs higher investments, but provides labour intensive cultivation with diversified products and incomes, which in turn need special marketing skills, otherwise products may be perished and wasted. Thus, special governmental incentives might be helpful for achieving overall sustainability and environmental safety with involving farmer's active participations.
- ◆ The multi-enterprising rice based integrated system compete with the intensive mono-cropping practices as well as limited farm resources.

## 7. Potential barriers to adoption

**Table 76.** Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Animals and crops can be damaged during extreme events, such as flooding, where fish, livestock and plants may sweep away or hurt, and drought and/or hot spells (water deficiency).
Cultural	Yes	Practice of some components like pig husbandry may compete with socio-economic and religious belief.
Economic	Yes	<p>Implementation costs: The initial costs of implementation like land shaping and cost of various types of inputs (seeds, planting material, fish fingerlings), animal components (duckling, chicks, etc.) require higher investments.</p> <p>Lack of suitable financial incentive, availability and subsidized inputs for farmers make RIFS less attractive for small and marginal farmers.</p> <p>Poor access to insurance, credit and markets: Limited access of farmers to credit and finance and heavy insurance procedures and gain access to markets largely undermines the economic viability and motivation towards adoption of RIFS.</p> <p>Lack of suitable added-value of product and long-term storage and market transport facilities limits economic successes of the systems.</p>
Institutional	Yes	Lack of coordination among sectors and producers: In many countries, RIFS fall under agriculture, environment and forestry departments and animal husbandry, thereby no single institutional mechanism taking a lead role in the advancement of adoption of IFS.
Legal (Right to soil and water)	Yes	Insecure tenure and small holding: Many small farmers have landless or holding very small size of land which act as major barriers towards adoption. Without formal land title, farmers are not interested to plant trees and horticultural plants.
Knowledge	Yes	RIFS include combination of different components and the production systems are knowledge intensive. This requires suitable access to information and technical support (e.g. extension services). Farmers are mostly reluctant to take risks and fear production losses. Sustained efforts are needed to strengthen extension and institutional functionaries to create adequate awareness and knowledge by organising training and demonstration for farmers/entrepreneur's motivation.
Natural resource	Yes	Fragmentation of land holding size may restrict farmer to implement profit-oriented farming systems.
Other	Yes	Theft, poaching and presence of predators can threaten the good implementation of RIFS.

## References

- Adarsh, S., Jacob, J. & Giffy, T. 2019. Role of pulses in Cropping Systems: A Review. *Agricultural Reviews*, 40(3): 185-191. <https://doi.org/10.18805/ag.R-1888>
- Ahmed, N., Ward, J.D. & Saint, C.P. 2014. Can integrated aquaculture-agriculture (IAA) produce “more crop per drop”? *Food Security*, 6(6): 767–779. <https://doi.org/10.1007/s12571-014-0394-9>
- Babu, S., Das, A., Mohapatra, K.P., Yadav, G.S., Singh, R., Tahashildar, M., Devi, M.T., Das, S., Panwar, A.S. & Prakash, N. 2019. Pond dyke utilization: An innovative means for enhancing productivity and income under Integrated Farming System in North East Hill Region of India. *Indian Journal of Agricultural Sciences*, 89(1):117-122.
- Bashir, M.A., Liu, J., Geng, Y., Wang, H., Pan, J., Zhang, D., Rehim, A., Aon, M. & Liu, H. 2020. Co-culture of rice and aquatic animals: An integrated system to achieve production and environmental sustainability. *Journal of Cleaner Production*, 249: 119310. <https://doi.org/10.1016/j.jclepro.2019.119310>
- Berg, H., Ekman Söderholm, A., Söderström, A.-S. & Tam, N.T. 2017. Recognizing wetland ecosystem services for sustainable rice farming in the Mekong Delta, Vietnam. *Sustainability Science*, 12(1): 137–154. <https://doi.org/10.1007/s11625-016-0409-x>
- Bhatt, R., Kukal, S.S., Busari, M.A., Arora, S. & Yadav, M. 2016. Sustainability issues on rice–wheat cropping system. *International Soil and Water Conservation Research*, 4(1): 64-74. <https://doi.org/10.1016/j.iswcr.2015.12.001>
- Bhattacharyya, P., Sinhababu, D.P., Roy, K.S., Dash, P.K., Sahu, P.K., Dandapat, R., Neogi, S. & Mohanty, S. 2013. Effect of fish species on methane and nitrous oxide emission in relation to soil C, N pools and enzymatic activities in rainfed shallow lowland rice–fish farming system. *Agric. Ecosyst. Environ.* 176: 53–62. <https://doi.org/10.1016/j.agee.2013.05.015>
- Bieluczyk, W., Piccolo, M. de C., Pereira, M.G., Moraes, M.T. de, Soltangheisi, A., Bernardi, A.C. de C., Pezzopane, J.R.M., Oliveira, P.P.A., Moreira, M.Z., Camargo, P.B. de, Dias, C.T. dos S., Batista, I. & Cherubin, M.R. 2020. Integrated farming systems influence soil organic matter dynamics in southeastern Brazil. *Geoderma*, 371: 114368. <https://doi.org/10.1016/j.geoderma.2020.114368>
- Bot, A. & Benites, J. 2005. *The importance of soil organic matter: Key to drought-resistant soil and sustained food production*. FAO Soils Bulletin No. 80. Food and Agriculture Organization of the United Nations, Rome, Italy. (also available at: <http://www.fao.org/3/a0100e/a0100e00.htm>)
- Cai, C., Li, G., Zhu, J.Q., Peng, L., Li, J.F. & Wu, Q.X. 2019. Effects of Rice–crawfish Rotation on Soil Physicochemical Properties in Jiangnan Plain. *Acta Pedol. Sin.* 56: 220–230. <https://doi.org/10.11766/trxb201804020127>
- Das, T., Sarkar, P. & Prasad, N. 2014. Exploring the potential for concurrent rice–fish culture in wetlands of Assam, North East India. *International Research Journal of Biological Sciences*, 3: 60e69



Datta, A., Nayak, D.R., Sinhababu, D.P. & Adhya, T.K. 2009. Methane and nitrous oxide emissions from an integrated rainfed rice–fish farming system of Eastern India. *Agriculture, Ecosystems & Environment*, 129(1-3): 228-237.

Oliveira, J. de M., Madari, B.E., Carvalho, M.T. de M., Assis, P.C.R., Silveira, A.L.R., de Leles Lima, M., Wruck, F.J., Medeiros, J.C. & Machado, P.L.O. de A. 2018. Integrated farming systems for improving soil carbon balance in the southern Amazon of Brazil. *Regional Environmental Change*, 18(1): 105–116. <https://doi.org/10.1007/s10113-017-1146-0>

Dixon, J.A., Gibbon, D.P. & Gulliver, A. 2001. *Farming systems and poverty: improving farmers' livelihoods in a changing world*. Food and Agriculture Organization of the United Nations, World Bank. Rome, Washington. (also available at: <http://www.fao.org/3/a-ac349e.pdf>)

Dwiyana, E. & Mendoza, T.C. 2006. Comparative Productivity, Profitability and Efficiency of Rice Monoculture and Rice-Fish Culture Systems. *Journal of Sustainable Agriculture*, 29(1): 145–166. [https://doi.org/10.1300/J064v29n01\\_11](https://doi.org/10.1300/J064v29n01_11)

Erenstein, O., Thorpe, W., Singh, J. & Varma, A. 2007. *Crop-livestock interactions and livelihoods in the Indo-Gangetic Plains, India: A Regional Synthesis*. CIMMYT, ILRI, RWC. (also available at: [https://cgspace.cgiar.org/bitstream/handle/10568/276/CLISS\\_Synthesis.pdf?sequence=1&isAllowed=y](https://cgspace.cgiar.org/bitstream/handle/10568/276/CLISS_Synthesis.pdf?sequence=1&isAllowed=y))

Fang, K., Yi, X., Dai, W., Gao, H. & Cao, L. 2019. Effects of Integrated Rice–Frog Farming on Paddy Field Greenhouse Gas Emissions. *International journal of environmental research and public health*, 16(11), p.1930.

FAO. 2019a. *Report of the Special Session on Advancing Integrated Agriculture Aquaculture through Agroecology*. Montpellier, France, 25 August 2018. FAO Fisheries and Aquaculture Report No. 1286. Rome. (also available at: <http://www.fao.org/3/ca7209en/CA7209EN.pdf>)

FAO. 2019b. *Soil erosion: the greatest challenge to sustainable soil management*. Rome. 100 pp. (also available at: <http://www.fao.org/3/ca4395en/ca4395en.pdf>)

FAO-SHOU. 2020. *Report of the FAO – SHOU International Promotion Programme Workshop on Social Impact of Rice-Fish Farming*. Shanghai, China, 4-8 December 2018. FAO Fisheries and Aquaculture Report No. 1317, Rome. <https://doi.org/10.4060/ca9907en>

Frei, M., Khan, M.A.M., Razzak, M.A., Hossain, M.M., Dewan, S. & Becker, K. 2007. Effects of a mixed culture of common carp, *Cyprinus carpio* L., and Nile tilapia, *Oreochromis niloticus* (L.), on terrestrial arthropod population, benthic fauna, and weed biomass in rice fields in Bangladesh. *Biological Control*, 41(2): 207–213. <https://doi.org/10.1016/j.biocontrol.2007.02.001>

Gajda, A.M., Czyż, E.A. & Dexter, A.R. 2016. Effects of long-term use of different farming systems on some physical, chemical and microbiological parameters of soil quality. *International Agrophysics*, 30(2): 165–172. <https://doi.org/10.1515/intag-2015-0081>

Gongfu, Z. 1990. The types, structure and results of the dike-pond system in South China. *GeoJournal*, 21(1-2): 83-89. <https://doi.org/10.1007/BF00645312>

- Halim, A., Sa'adah, N., Abdullah, R., Karsani, S.A., Osman, N., Panhwar, Q.A. & Ishak, C.F. 2018. Influence of soil amendments on the growth and yield of rice in acidic soil. *Agronomy*, 8(9): 165. <https://doi.org/10.3390/agronomy8090165>
- Halwart, M. 2008. Biodiversity, nutrition and livelihoods in aquatic rice-based ecosystems. *Biodiversity*, 9(1-2): 36-40. <https://doi.org/10.1080/14888386.2008.9712879>
- Halwart, M. & Gupta, M.V. (eds.) 2004. Culture of fish in rice fields. FAO and The World Fish Center, 83 p. (also available at: <http://pubs.iclarm.net/Pubs/CultureOfFish/Culture-of-Fish.pdf>)
- Hu, L., Zhang, J., Ren, W., Guo, L., Cheng, Y., Li, J., Li, K., Zhu, Z., Zhang, J., Luo, S., Cheng, L., Tang, J. & Chen, X. 2016. Can the co-cultivation of rice and fish help sustain rice production? *Scientific Reports*, 6. <https://doi.org/10.1038/srep28728>
- IPCC. 2007. *Climate Change 2007: the Physical Science Basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, United Kingdom and New York, USA.
- IPCC. 2014. *Climate change 2014: impacts, adaptation, and vulnerability*. Cambridge University Press, Cambridge and New York
- Islam, A.H.M.S., Barman, B.K. & Murshed-e-Jahan, K. 2015. Adoption and impact of integrated rice-fish farming system in Bangladesh. *Aquaculture*, 447: 76-85. <https://doi.org/10.1016/j.aquaculture.2015.01.006>
- Kaledhonkar, M.J., Meena, B.L. & Sharma, P.C. 2019. Reclamation and Nutrient Management for Salt-affected Soils. *Indian J. of Fertilizer*, 15(5): 566-575.
- Karim, M. & Little, D.C. 2018. The impacts of integrated homestead pond-dike systems in relation to production, consumption and seasonality in central north Bangladesh. *Aquaculture Research*, 49(1): 313-334. <https://doi.org/10.1111/are.13462>
- Kopittke, P.M., Menzies, N.W., Wang, P., McKenna, B.A. & Lombi, E. 2019. Soil and the intensification of agriculture for global food security. *Environment International*, 132: 105078. <https://doi.org/10.1016/j.envint.2019.105078>
- Kremen, C., Iles, A. & Bacon, C. 2012. Diversified Farming Systems: An Agroecological, Systems-based Alternative to Modern Industrial Agriculture. *Ecology and Society*, 17(4). <https://doi.org/10.5751/ES-05103-170444>
- Kumaran, M., Ghoshal, T.K., De, D., Biswas, G., Raja, R.A., Anand, P.S., Panigrahi, A. & Vijayan, K.K. 2020. Aquaculture-based production systems for the livelihood security of coastal farm families in the risk-prone agro-ecosystem of India: an appraisal. *Aquaculture International*, 28(2): 805-814. <https://doi.org/10.1007/s10499-019-00495-y>
- Li, M., Li, R., Zhang, J., Liu, S., Hei, Z. & Qiu, S. 2019. A combination of rice cultivar mixed-cropping and duck co-culture suppressed weeds and pests in paddy fields. *Basic and Applied Ecology*, 40: 67-77. <https://doi.org/10.1016/j.baec.2019.09.003>

- Li, M.J., Li, R.H., Zhang, J.E., Guo, J., Zhang, C.X., Liu, S.W., Hei, Z. & Qiu, S.Q.** 2020. Integration of mixed-cropping and rice-duck co-culture has advantages on alleviating the non-point source pollution from rice, (*Oryza sativa* L.) production. *Applied Ecology and Environmental Research*, 18(1): 1281-1300. [http://dx.doi.org/10.15666/aecr/1801\\_12811300](http://dx.doi.org/10.15666/aecr/1801_12811300)
- Long, P., Huang, H., Liao, X., Fu, Z., Zheng, H., Chen, A. & Chen, C.** 2013. Mechanism and capacities of reducing ecological cost through rice-duck cultivation. *Journal of the Science of Food and Agriculture*, 93(12): 2881-2891. <https://doi.org/10.1002/jsfa.6223>
- Lu, J. & Li, X.** 2006. Review of rice-fish-farming systems in China – One of the Globally Important Ingenious Agricultural Heritage Systems (GIAHS). *Aquaculture*, 260(1): 106–113. <https://doi.org/10.1016/j.aquaculture.2006.05.059>
- Masciandaro, G., Macci, C., Peruzzi, E. & Doni, S.** 2018. Chapter 1 - Soil Carbon in the World: Ecosystem Services Linked to Soil Carbon in Forest and Agricultural Soils. In C. Garcia, P. Nannipieri & T. Hernandez (Eds.) *The Future of Soil Carbon*, pp. 1–38. Academic Press. <https://doi.org/10.1016/B978-0-12-811687-6.00001-8>
- Melaku, S. & Natarajan, P.** 2019. Status of integrated aquaculture - Agriculture systems in Africa. *International J. of Fisheries and Aquaculture Studies*, 7(4): 263-269.
- Mottet, A., Henderson, B., Opio, C., Falcucci, A., Tempio, G., Silvestri, S., Chesterman, S. & Gerber, P.J.** 2017. Climate change mitigation and productivity gains in livestock supply chains: insights from regional case studies. *Regional Environmental Change*, 17(1): 129-141. <https://doi.org/10.1007/s10113-016-0986-3>
- Nayak, P.K., Nayak, A.K., Panda, B.B., Lal, B., Gautam, P., Poonam, A., Shahid, M., Tripathi, R., Kumar, U., Mohapatra, S.D. & Jambhulkar, N.N.** 2018. Ecological mechanism and diversity in rice based integrated farming system. *Ecological Indicators*, 91: 359-375. <https://doi.org/10.1016/j.ecolind.2018.04.025>
- Nayak, P.K., Nayak, A.K., Kumar, A., Kumar, U., Panda, B.B., Satapathy, B.S., Poonam, A., Mohapatra, S.D., Tripathi, R., Shahid, M., Chaterjee, D., Panerselvam, P., Mohanty S. & Pathak, H.** 2020a. Rice based integrated farming system in eastern India: A viable technology for productivity and ecological security. NRRI Research Bulletin, No. 24, ICAR-National Rice Research Institute, Cuttack-753006, Odisha, India. pp 44. (also available at: <https://icar-nrri.in/wp-content/uploads/2020/05/NRRI-Research-Bulletin-24.pdf>)
- Nayak, P.K., Panda, B.B., Das, S.K., Rao, K.R., Kumar, U., Kumar, A., Munda, S., Satapathy, B.S. & Nayak, A.K.** 2020b. Weed control efficiency and productivity in rice-fish-duck integrated farming system. *Indian J. Fish.*, 67(3): 62-71, <https://doi.org/10.21077/ijf.2020.67.3.94309-07>
- Ongley, E.D.** 1996. *Control of water pollution from agriculture*. Vol. 55. Food and Agriculture Organization of the United Nations.

- Pathak, H., Samal, P. & Shahid, M.** 2018. Revitalizing rice-systems for enhancing productivity, profitability and climate resilience. *In* Pathak, H., Nayak, A.K., Jena, M., Singh, O.N., Samal, P. & Sharma, S.G. (Eds.) *Rice Research for Enhancing Productivity, Profitability and Climate Resilience*. ICAR-National Rice Research Institute, Cuttack, Odisha, India, pp. 452.
- Paramesh, V., Parajuli, R., Chakurkar, E.B., Sreekanth, G.B., Kumar, H.C., Gokuldas, P.P., Mahajan, G.R., Manohara, K.K., Viswanatha, R.K. & Ravisankar, N.** 2019. Sustainability, energy budgeting, and life cycle assessment of crop-dairy-fish-poultry mixed farming system for coastal lowlands under humid tropic condition of India. *Energy*, 188: 116101. <https://doi.org/10.1016/j.energy.2019.116101>
- Pernollet, C.A., Simpson, D., Gauthier-Clerc, M. & Guillemain, M.** 2015. Rice and duck, a good combination? Identifying the incentives and triggers for joint rice farming and wild duck conservation. *Agriculture, Ecosystems & Environment*, 214: 118–132. <https://doi.org/10.1016/j.agee.2015.08.018>
- Si, G., Peng, C., Yuan, J., Xu, X., Zhao, S., Xu, D. & Wu, J.** 2017. Changes in soil microbial community composition and organic carbon fractions in an integrated rice–crayfish farming system in subtropical China. *Scientific reports*, 7(1): 1-10. <https://doi.org/10.1038/s41598-017-02984-7>
- Singh, B.R.** 2011. Environmental health risks from integrated farming system (IFS). *In* *Environmental Health: Human and Animal Risk Mitigation*, pp.373-383.
- Soussana, J-F., Muriel, T., Philippe, L., & Bertrand, D.** 2015. Agroecology: integration with livestock. FAO, 2015, Rome, Italy. (also available at: <http://www.fao.org/3/a-i4729e.pdf>)
- Teng, Q., Hu, X.F., Cheng, C., Luo, Z., Luo, F., Xue, Y., Jiang, Y., Mu, Z., Liu, L. & Yang, M.** 2016. Ecological effects of rice–duck integrated farming on soil fertility and weed and pest control. *Journal of soils and sediments*, 16(10): 2395-2407. <https://doi.org/10.1007/s11368-016-1455-9>
- Visca Jr, M.D. & Palla, S.Q.** 2018. Golden apple snail, *Pomacea canaliculata* meal as protein source for rabbitfish, *Siganus guttatus* culture. *Aquaculture, Aquarium, Conservation & Legislation*, 11(2): 533-542.
- Wang, A., Ma, X., Xu, J. & Lu, W.** 2019. Methane and nitrous oxide emissions in rice–crab culture systems of northeast China. *Aquaculture and Fisheries*, 4: 134-141. <https://doi.org/10.1016/j.aaf.2018.12.006>
- Wu, S., Hu, Z., Hu, T., Chen, J., Yu, K., Zou, J. & Liu, S.** 2018. Annual methane and nitrous oxide emissions from rice paddies and inland fish aquaculture wetlands in southeast China. *Atmos. Environ.* 175: 135-144. <https://doi.org/10.1016/j.atmosenv.2017.12.008>
- Xie, Y.Q., Zhang, J.F., Jiang, H.M., Yang, J.C., Deng, S.H., Li, X., Guo, J.M., Li, L.L., Liu, X. & Zhou, G.Y.** 2015. Effects of different fertilization practices on greenhouse gas emissions from paddy soil (In Chinese). *J. Agro-Environ. Sci.* 34: 578–584.
- Xu, G., Liu, X., Wang, Q., Yu, X. & Hang, Y.** 2017. Integrated rice–duck farming mitigates the global warming potential in rice season. *Science of the Total Environment*, 575: 58-66. <https://doi.org/10.1016/j.scitotenv.2016.09.233>

**Yan, Y., Liu, M., Yang, D., Zhang, W., An, H., Wang, Y., Xie, H. & Zhang, X.** 2014. Effect of Different Rice-Crab Coculture Modes on Soil Carbohydrates. *Journal of Integrative Agriculture*, 13(3): 641–647.  
[https://doi.org/10.1016/S2095-3119\(13\)60722-4](https://doi.org/10.1016/S2095-3119(13)60722-4)

**Zhang, Y., Li, Y., Jiang, L., Tian, C., Li, J. & Xiao, Z.** 2011. Potential of perennial crop on environmental sustainability of agriculture. *Procedia Environmental Sciences*, 10:1141-1147.  
<https://doi.org/10.1016/j.proenv.2011.09.182>

**Zhang, J., Guoming, Q., Benliang, Z., Kaiming, L. & Zhong, Q.** 2017. Rice-Duck Co-culture in China and its Ecological Relationships and Functions. *In Agroecology in China: Science, Practice, and Sustainable Management*.

**Zhang, J., Hu, L., Ren, W., Guo, L., Tang, J., Shu, M. & Chen, X.** 2016. Rice-soft shell turtle coculture effects on yield and its environment. *Agriculture, Ecosystems & Environment*, 224: 116-122.  
<https://doi.org/10.1016/j.agee.2016.03.045>



# Urban soils and infrastructures

# 20. Management of gardens, parks, and lawns

John M. Galbraith<sup>1</sup>, David Yocca<sup>2</sup>, Mary Pat McGuire<sup>3</sup>

<sup>1</sup>Virginia Tech, Blacksburg, VA, United States of America

<sup>2</sup>Solutions in the Land, and, Green Infrastructure Foundation, United States of America

<sup>3</sup>University of Illinois, Urbana Champaign, United States of America

## 1. Description of the practice/concept

Urban planners, landscape architects, and other urban designers create and maintain gardens, parks, plazas, forest- and ecological -preserves, recreation areas (such as golf courses), streetscapes, and other landscapes in cities. The ground surfaces are composed of mosaics of trees, woodlands, shrubs, grasses, perennials, edibles, lawns, and waterbodies as well as paved areas, such as sidewalks, roads, and parking areas. Landscapes, including pleasure gardens, edible gardens, lawns, and park settings, are also designed and maintained on private land. Locations range from small residential lots to larger homesites and institutional properties, such as campuses and corporations. Taking this variety into account for purposes of this practice, gardens and parks are urban areas created and managed for public space, recreation, and ecological diversity. Together, gardens and parks cover a range of site scale and vegetation complexity and therefore soil extents.

## 2. Range of applicability

The history of development of public gardens and parks in cities all over the world varies by ecological and cultural context. Historical land-uses are drivers for the potential to accumulate soil organic carbon (SOC). The content of SOC varies widely. The highest SOC storage was recorded in wetland soils followed by forest soils and then lawn soils (Bae and Ryu, 2015). The range of accumulation is affected by local climate; geological and topographical formation; landscape typologies deployed within these features; native soil types; urban and infrastructural development; cultural attitudes; and landscape management and maintenance practices. Pouyat *et al.* (2002) compared carbon densities in different urban land types and found that low-density residential and institutional areas had 44 and 38 percent higher organic carbon densities, respectively, than soils of commercial areas.

## 3. Impact on soil organic carbon stocks

The potential organic carbon sequestration in urban areas varies by land use and distance from the urban core (Table 77). The quantity of SOC varies based on vegetative cover, maintenance intensity and history, and regulations for topsoil restoration (Brown, Miltner, and Cogger, 2012). Studies reporting carbon sequestration must be carefully read to distinguish the source (aboveground, root, or soil) and type (organic, inorganic, or total carbon). Eighty-three percent of carbon stocks are stored in soil, 16 percent in trees and shrubs, and 0.6 percent in herbal vegetation (Jo and McPherson, 1995). In most studies, organic carbon is reported. However, increases in soil inorganic carbon have been documented in arid regions as well along a rural-urban transect (Koerner and Klopatek, 2010).

The highest density of soil carbon is in the topsoil and in leaf litter in woodlands. Construction, excavation and mass grading, and infrastructure development in urban areas usually result in clearing of trees and removal or burial of topsoil (Logsdon, Sauer and Cambardella, 2017). In mass grading, topsoil is stripped and either stockpiled for re-spreading, where it oxidizes SOC, or sold by developers as an income source (Pouyat *et al.*, 2010). Compaction by foot and vehicle traffic results in dense soil and lower plant production. All of these factors may lead to lower soil organic matter, carbon stocks, and carbon inputs.

In more arid regions, the content of soil carbon in urban areas is often elevated compared to nearby natural landscapes. This difference is likely due to human importation of water to supplement the local climate and grow vegetation, which places more vegetative litter into the soil system (Trammel *et al.*, 2020). The greatest increases in urban soil carbon have been observed in the most highly managed soils, such as golf courses and lawns (Pataki *et al.*, 2006). Urban areas are also a source of organic matter that can be added to the soil or to containers. Urban sources include fresh and composted food waste, yard and park maintenance waste, and hay and manure from zoos and horse stables. Bio-solids and liquids are produced from disposal and recycling of human waste products. Organic matter in all forms may be used to increase the storage of carbon in the soil up to the point of equilibrium with microbial decomposition and harvesting removals. This point is reached in turfgrass after 25 to 40 years (Shi, Bowman and Ruffy, 2012). Tree density in a park in Almada, Portugal, was positively correlated with total carbon stocks. Grasslands with high tree density areas and the forest had 228 and 262 tC/ha of total carbon from soil to overstory. Soil carbon was highest under forest and lawns with low tree density (Mexia *et al.*, 2018).



**Table 77.** Changes in soil organic carbon stocks reported for management of parks, gardens and lawns

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Years)	Depth (cm)	More information	Reference
Nebraska City, NE, United States of America	udic, mesic = humid continental	Typic Argiudoll in Soil Taxonomy; Vertic Luvisols in World Resource Base		0.52 for unirrigated fine fescue 0.74 for irrigated fine fescue 0.32 for Kentucky bluegrass 0.78 for creeping bentgrass	4	20		Qian and Follet (2002)
East Lansing, MI, United States of America	udic, mesic = Humid Continental Mild Summer	container soil	8	0.5	2	6		Getter <i>et al.</i> (2009)
Seoul, Democratic People's Republic of Korea	Tropical	Clay loam	20–140	SOC concentrations in topsoil increased 256 percent	10	100	Seoul Forest park; SOC stock dependent on land use	Bae and Ryu (2015)
Helsinki and Lahti, Finland	Boreal	Spodosol	149	0.6	100	50	10 parks of contrasting age	Setälä <i>et al.</i> (2016)
Urumqui city, NW China	Arid	Solonetz, Castanozem	55	1.25	20	80	11 urban greenspaces; SOC storage due to irrigation and fertilisation	Yan <i>et al.</i> (2016)

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Adding organic matter to landscape and garden soils on a regular basis is a common practice to improve some of the physical and chemical properties of the soil. The additions should not exceed the long-term equilibrium organic carbon content of reference topsoil levels. Adding organic matter (which eventually decomposes into humus and other stable carbon compounds) to topsoil in sites that have lost topsoil to erosion or removal, should help in improving uptake of nutrients, providing N, P and S following mineralization, reducing the toxicity of certain metals such as Al at low pH, capturing pollutants such as Atrazine, increasing the cation exchange capacity of the soil, buffering against change in pH, lowering the density, improving soil structure, increasing infiltration and water-holding capacity, and decreasing surface evaporation losses and crusting. Adding mulch also helps to keep weeds from growing and provides long-term organic carbon and nutrient supply which eventually results in amended soil fertility and pH. Therefore, soil properties may be enhanced by adding organic residues following lawn mowing, shrub pruning, and tree-branch trimming; adding decaying perennial plant matter from the ground surface; and after composting (as practiced in Seoul Forest Park (Bae and Ryu, 2015)). Adequate garden management may decrease salinization and alkalinity.

### 4.2 Minimization of threats to soil functions

Properly designed and managed planting areas can increase infiltration and minimize soil exposure to wind, runoff, and erosion. For example, gardens on sloping ground should be terraced on the contour to decrease runoff and erosion. In parks and lawns, pavements can be made of pervious materials that create a firm surface for walking or driving but also allow infiltration and decrease runoff.

Table 78. Soil threats

Soil threats	
Soil erosion	Soil erosion can be prevented by adequate management; i.e. maintaining continuous soil cover.
Nutrient imbalance and cycles	Nutrient cycles may be fostered by careful fertilization practices and organic matter additions.
Soil salinization and alkalization	Soil salinization and alkalization may be amended through adequate garden management (see below).
Soil biodiversity loss	Soil biodiversity loss may be avoided by the establishment of gardens, which vary widely in structures that may promote plant biodiversity, such as ponds, moss, ground cover, and varied plant species (Tresch <i>et al.</i> , 2019).

Soil threats	
Soil sealing	Soil sealing will be avoided because parks and gardens generally contain few impervious surfaces.
Soil water management	Gardens, parks, and lawns allow for water infiltration, thus helping to prevent stormwater runoff.

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Gardens, parks, lawns, and school gardens produce significant amounts of vegetation and organic matter. Many of these are perennial plants with long-lived root systems that add organic matter beneath the soil surface. Production can be harvested and used for domestic consumption and commercial sales.

### 4.4 Mitigation of and adaptation to climate change

Parks, gardens, and lawns contribute to climate change mitigation through carbon sequestration in biomass as well as soil. This contribution, however, is mostly omitted in national and regional carbon stock estimates, especially in the global south, where it can be substantial. For example, in the metropolitan area of Kumasi, Ghana, a total of 1676 Gg or 66 t/ha of carbon is stored in above- and below-ground vegetation and soil of parks, gardens, and lawns (Nero *et al.*, 2017). Like other green infrastructures, parks that have high tree density are cooling islands. They thereby help to alleviate high temperatures and to increase moisture over a distance of up to 60 meters away (Grilo *et al.*, 2020).

### 4.5 Socio-economic benefits

Natural, semi-natural, and artificial networks of multifunctional ecological systems, gardens, parks, and lawns are extremely important for sustainable cities because they contribute to ecosystem health and human health (Tzoulas *et al.*, 2007). Pleasure, recreation, exercise, socialization, social inclusion, education, aesthetics, environmental improvement, and therapy are benefits of managed systems, such as gardens, parks, and lawns. The act of maintaining and recreating in these areas often involves groups of people that share common interests. Maintenance and management of gardens, parks, and lawns involve an ongoing learning and training process. Maintenance can be aesthetically pleasing and beneficial to the birds, insects, bees, earthworms, microbes, mammals, and other biology within an urban area. The process of designing, managing, and maintaining gardens, parks, and lawns can be a calming and peaceful way to experience outdoor areas in an otherwise stressful urban environment.

## 4.6 Additional benefits to the practice

The inclusion of communities is essential in decision making for the design and management of public land, ecological landscape types, and soil health that promotes carbon sequestration. By involving the public in the formation of adaptable and ecologically functional landscapes, greater education, local maintenance, and even use of these areas are enhanced by a sense of collective involvement, ownership, and responsibility.

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

Runoff, erosion, and deep percolation following the application of excess chemical fertilizers can cause both surface and groundwater pollution by nitrogen, phosphorous, and potassium. Certain herbicides, pesticides, and fungicides can build up in the soil, harm humans, or harm beneficial insects, birds, bees, and earthworms. Reading and following label directions are advised to minimize threats. Turfgrass management practices, such as frequent mowing, can cause soil compaction and may require significant amounts of supplemental irrigation in warm, semi-arid and arid regions.

**Table 79.** Soil threats

Soil threats	
<b>Soil erosion</b>	Uncovered soil in gardens, parks, and lawns can be subject to erosion.
<b>Soil salinization and alkalinization</b>	Soil salinization is increased on golf courses and lawns that are watered with saline groundwater, especially in soils that have surface compaction, restricted subsoil permeability, and higher clay contents (Miyamoto and Chacon, 2006).
<b>Soil contamination/pollution</b>	Private gardening activities may result in misuse of fertilizers and other agrochemicals. Heavy metal contamination may accumulate in close proximity to vehicle traffic or where the green infrastructure is located at an old building site.
<b>Soil acidification</b>	Fertilizing with sulfur or sulfate products can lead to acidification of the soil. Large additions of acidic organic matter or compost can also.

Soil threats	
Soil biodiversity loss	Excess additions of herbicides, pesticides, and fungicides can lead to loss of beneficial insects and biodiversity in soils.
Soil water management	Supplemental water may be needed to maintain vegetative cover in semi-arid and arid regions

## 5.2 Increases in greenhouse gas emissions

Management of gardens, parks, and lawns may include trade-offs of greenhouse gas emission following their maintenance. A recent study assessed SOC sequestration and greenhouse gas emissions from turfgrass on athletic fields and ornamental lawns of urban parks down to 50 cm (Townsend-Small and Czimzik, 2010). The results showed that greenhouse gas emission from fuel use, fertilization, and irrigation outweighed SOC sequestration and that turfgrass establishment is not a greenhouse gas mitigation option (Townsend-Small and Czimzik, 2010). Use of petrol-powered machinery for garden maintenance, e.g., mowing, may have an important carbon footprint.

## 5.3 Conflict with other practice(s)

In most gardens, parks, and lawns, the tree density is reduced as compared to woodlands. This practice may thus compete with urban forestry.

## 5.4 Other conflicts

Establishment of public gardens, parks, and lawns has been prevented in some places because of the need for land to build housing for city dwellers in highly populated urban areas. Space is needed for commercial buildings, transportation, and sealed surfaces, such as pavements.

## 6. Recommendations before implementation of the practice

Because of the wide range of potential carbon sequestration as well as many other performance factors, an integrative design approach should be used for the creation or renovation of any park, garden, or other public green space. Consideration should be given to all of the benefits and values these areas can provide if they are designed and managed as a system. It could be good to maintain areas of park landscapes as naturalized landscapes, either through preservation of existing natural conditions or through ecological restoration practices. Such practices increase biodiversity and lower the carbon and energy footprint, or the green infrastructure, because they can be maintained with minimal or no inputs of supplemental water, chemical fertilizers, herbicides, or pesticides. Semi-natural gardens and lawns, which have a reduced management intensity, should be preferred. Reduced management intensity has a positive effect on biodiversity and leads to reduction of external inputs, thereby increasing sustainable SOC sequestration. In most cities, plant litter is removed for cleanliness and to prevent inconvenience for users. Removal of grass clippings and plant and tree leaves in these areas was found to reduce the soil organic content (Yoon *et al.*, 2016). In such cases, organic waste treatment that includes transformation into organic amendments should be performed. For example, in the Seoul Forest Park, increased SOC storage could be achieved by returning composted litter layers to soil (Bae and Ryu, 2015).

Salinization can be remedied by decreasing soil compaction, choosing soil with a lower clay content and good subsoil permeability (Miyamoto and Chacon, 2006), decreasing surface evaporation, watering infrequently with salt-free water, and ensuring free drainage after watering. Alkalization is a common problem in urban areas because of dust derived from concrete and calcareous gravel road traffic. Garden management, therefore, must aim to achieve adequate soil pH. Soil reaction (pH) can be lowered by adding organic matter or sulfate compounds and can be raised by adding lime. Gypsum can be added to improve soil structure, and dense soil can be loosened by infrequent tillage or spading and by the addition of organic matter.

Because soil in humid and subhumid climates is prone to erosion, gardens, parks, and lawns should have minimal soil exposed to wind or water erosion, except briefly during establishment, construction, or reclamation.

Risks of nitrate and phosphate losses should be avoided through adequate fertilizer use. Heavy metal contamination should be avoided by testing amendments (such as composts, biosolids, and other waste products) for contaminants before use. Atmospheric dust and other potential current contamination sources (old buildings, construction activities, etc.) should be considered when establishing a proper management plan for a garden, park, or lawn.

Foot and vehicle traffic should be concentrated in a few areas rather than across the area to be managed for plant growth. When sidewalks and pavements must be added, pervious materials should be used if possible. Soils that have been compacted by foot and vehicle traffic should be ripped or loosened to lessen the density, and organic matter should be added to promote soil structure development. Walkways between garden beds and through parks can be covered with steppingstones or layers of wood chips to minimize compaction.

## 7. Potential barriers to adoption

Competition for land use in urban areas is high and is regulated through zoning, ordinances, and homeowners' associations. Ownership of gardens and lawns is private and public, while parks are most likely publicly owned. Biophysical, cultural, social, economic, institutional, legal, and knowledge barriers may exist for using land for gardens, parks, and lawns in urban areas.

**Table 80.** Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Urban soils may be impoverished and polluted, thus preventing establishment of vegetation.
Cultural	No	In general, people like to live close to green, nature-like environments. Likewise, studies have shown that creating neat edges on the perimeter of ecological landscapes can help to increase receptivity to these sites (Nassauer, 2007).
Economic	Yes	Land values are extremely high in urban areas, and conversion of privately-owned gardens and lawns into other land uses is economically advantageous. Parks require maintenance, which may be costly.
Institutional	Yes	Poor leadership, governance, and city planning may have reduced the number of gardens, parks, and lawn infrastructures. Miss-management may lead to their destruction.
Legal (Right to soil)	No	Zoning and ordinances protect gardens, parks, and lawns from alternative land uses.
Knowledge	Yes	There may be a misunderstanding or underappreciation of the importance of gardens, parks, and lawns.

## Photos of the practice



Photo 25. A flower garden in Chicago, Illinois, United States of America



Photo 26. Echo Urban Park in Los Angeles, California, United States of America





Photo 27. An urban park in Houston, Texas, United States of America




Photo 28. Two urban parks in San Francisco, California, United States of America

Table 81. Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Compost application to restore post-disturbance soil health in Montgomery county, Virginia, United States</i>	North America	4	6	28
<i>Management of ornamental lawns and athletic fields in California, United States</i>	North America	2, 10, 20 and 33	6	29
<i>Water and residues management on a golf course, Nebraska, United States</i>	North America	4	6	30

## References

- Bae, J. & Ryu, Y.** 2015. Land use and land cover changes explain spatial and temporal variations of the soil organic carbon stocks in a constructed urban park. *Landscape and Urban Planning*, 136: 57–67. <https://doi.org/10.1016/j.landurbplan.2014.11.015>
- Brown, S., Miltner, E. & Cogger, C.** 2012. Carbon Sequestration Potential in Urban Soils. In R. Lal & B. Augustin (Eds.) *Carbon Sequestration in Urban Ecosystems*, pp. 173–196. Dordrecht, Springer Netherlands. [https://doi.org/10.1007/978-94-007-2366-5\\_9](https://doi.org/10.1007/978-94-007-2366-5_9)
- Getter, K.L., Rowe, D.B., Robertson, G.P., Cregg, B.M. & Andresen, J.A.** 2009. Carbon sequestration potential of extensive green roofs. *Environ. Sci. Technol.*, 43(19): 7564–7570. <https://doi.org/10.1021/es901539x>.
- Grilo, F., Pinho, P., Aleixo, C., Catita, C., Silva, P., Lopes, N., Freitas, C., Santos-Reis, M., McPherson, T. & Branquinho, C.** 2020. Using green to cool the grey: Modelling the cooling effect of green spaces with a high spatial resolution. *Science of The Total Environment*, 724: 138182. <https://doi.org/10.1016/j.scitotenv.2020.138182>
- Jo, H.K. & Pherson, E.G.** 1995. Carbon storage and flux in urban residential greenspace. *Journal of Environmental Management*, 45(2): 109–133. <https://doi.org/10.1006/jema.1995.0062>
- Koerner, B.A. & Klopatek, J.M.** 2010. Carbon fluxes and nitrogen availability along an urban–rural gradient in a desert landscape. *Urban Ecosystems*, 13(1): 1–21. <https://doi.org/10.1007/s11252-009-0105-z>
- Logsdon, S.D., Sauer, P.A. & Cambardella, C.A.** 2017. Digging to the top(soil). *Canadian Journal of Soil Science*, 97(4): 793–795. <https://doi.org/10.1139/cjss-2017-0047>
- Mexia, T., Vieira, J.I., Silva, A.P., Anjos, A., Lopes, N., Freitas, C., Santos-Reis, M., Correia, O., Branquinho, C. & Pinho, P.** 2018. Ecosystem services: Urban parks under a magnifying glass. *Environmental Research*, 160: 469–478. <https://doi.org/10.1016/j.envres.2017.10.023>
- Miyamoto, S. & Chacon, A.** 2006. Soil salinity of urban turf areas irrigated with saline water: II. Soil factors. *Journal of Landscape and Urban Planning*, 77(1–2): 28–38. <https://doi.org/10.1016/j.landurbplan.2004.12.011>
- Nassauer, J.** 1995. Messy Ecosystems, Orderly Frames. *Landscape Journal*, 14(2): 161–169. <https://doi.org/10.3368/lj.14.2.161>
- Nero, B.F., Callo-Concha, D., Anning, A. & Denich, M.** 2017. Urban green spaces enhance climate change mitigation in cities of the global south: the case of Kumasi, Ghana. *Procedia Engineering*, 198: 69–83. <https://doi.org/10.1016/j.proeng.2017.07.074>
- Pataki D.E., Alig, R.J., Fung, A.S., Golubiewski, N.E., Kennedy, C.A., McPherson, E.G., Nowak, D.J., Pouyat, R.V., Romero-Lankao, P.** 2006. Urban ecosystems and the North American carbon cycle. *Global Change Biology*, 12: 2092–2102. <https://doi.org/10.1111/j.1365-2486.2006.01242.x>

- Pouyat, R., Groffman, P., Yesilonis, I. & Hernandez, L.** 2002. Soil carbon pools and fluxes in urban ecosystems. *Environmental Pollution*, 116: S107–S118. [https://doi.org/10.1016/S0269-7491\(01\)00263-9](https://doi.org/10.1016/S0269-7491(01)00263-9)
- Pouyat, R., Szlávecz, K., Yesilonis, I.D., Groffman, P. & Schwarz, K.** 2010. Chemical, physical, and biological characteristics of urban soils. *Urban ecosystem ecology*, 119–152.
- Qian, Y. & Follett, R.F.** 2002. Assessing Soil Carbon Sequestration in Turfgrass Systems Using Long-Term Soil Testing Data. *Agronomy Journal*, 94(4): 930–935. <https://doi.org/10.2134/agronj2002.9300>
- Setälä, H.M., Francini, G., Allen, J.A., Hui, N., Jumpponen, A. & Kotze, D.J.** 2016. Vegetation Type and Age Drive Changes in Soil Properties, Nitrogen, and Carbon Sequestration in Urban Parks under Cold Climate. *Frontiers in Ecology and Evolution*, 4: 93. <https://doi.org/10.3389/fevo.2016.00093>
- Shi, W., Bowman, D. & Rufty, T.** 2012. Microbial Control of Soil Carbon Accumulation in Turfgrass Systems. Ch. 11. pp. 215-231. In Lal, R. & Augustin, B. (Eds.) *Carbon Sequestration in Urban Ecosystems*. Springer Publ., New York, NY, USA.
- Townsend-Small, A. & Czimczik, C.I.** 2010. Carbon sequestration and greenhouse gas emissions in urban turf. *Geophysical Research Letters*, 37(2). <https://doi.org/10.1029/2009GL041675>
- Tresch, S., Frey, D., Bayon, R.-C.L., Mäder, P., Stehle, B., Fliessbach, A. & Moretti, M.** 2019. Direct and indirect effects of urban gardening on aboveground and belowground diversity influencing soil multifunctionality. *Scientific Reports*, 9(1): 9769. <https://doi.org/10.1038/s41598-019-46024-y>
- Tzoulas, K., Korpela, K., Venn, S., Yli-Pelkonen, V., Kazmierczak, A., Niemela, J. & James, P.** 2007. Promoting ecosystem and human health in urban areas using Green Infrastructure: A literature review. *Landscape and Urban Planning*, 81: 167–178. <https://doi.org/10.1016/j.landurbplan.2007.02.001>
- Yan, Y., Zhang, C., Hu, Y.F. & Kuang, W.H.** 2016. Urban land-cover change and its impact on the ecosystem carbon storage in a dryland city. *Remote Sensing*, 8: 6. <https://doi.org/10.3390/rs8010006>
- Yoon, T.K., Seo, K.W., Park, G.S., Son, Y.M. & Son, Y.** 2016. Surface Soil Carbon Storage in Urban Green Spaces in Three Major South Korean Cities. *Forests*, 7: 115. <https://doi.org/10.3390/f7060115>

# 21. Bioretention systems

Cornelia Rumpel

CNRS, Institute of Ecology and Environmental Sciences, Paris, France

## 1. Description of the practice

Bioretention or biofiltration systems (e.g. bioswales, constructed stormwater facilities, and raingardens) have multiple functions in cities. Their main role is in hydrology, as the systems are intended to channel, retain, and purify rainwater. They are an alternative to conventional drainage systems composed of networks of artificial pipes and pits. They consist of vegetated channels and are designed specifically to attenuate and treat stormwater runoff from large impermeable surfaces, such as roads and parking lots (Figure 14). Their size is variable but should be at least one percent of the area being drained (EPA, 2000). They are designed to have rapidly permeable, sand-based soil media that reduces stormwater volume. Commonly, they receive organic matter amendments. Addition of labile organic matter and a submerged zone at the bottom of the facility significantly affect the removal of nutrients and heavy metals from the stormwater. Vegetation in these infrastructures includes selected species and ranges from herbal species to woody plants. Trees planted in bioretention facilities can provide many co-benefits, including shade, carbon sequestration and storage, air pollution reduction, and aesthetic improvement. However, tree planting is a challenge because the trees generally cannot tolerate a soil substrate that has high hydraulic conductivity, low nutrient availability, and toxic concentrations of pollutants. The success of tree planting depends on the type and depth of the growing substrate (Tirpak *et al.*, 2018).




Figure 14. Schematic view of a bioretention system

Source: adapted from <https://www.idsnews.com/article/2017/10/bioswale-installation-will-help-slow-flooding>

## 2. Range of applicability

Bioretention systems were introduced in the 1990s as a best management practice for urban stormwater management (County, 1993). Traditionally, they were installed to reduce water volume, control erosion, and improve water quality. In recent years, their crucial role for human well-being and various other ecosystem services, such as air quality improvement, biodiversity, and carbon storage, has been documented (Prudencio and Null, 2018). These systems may be a sustainable practice to provide hydrologic ecological restoration of urban areas and may help to meet goals for downstream water quality (Liu *et al.*, 2014). Therefore, in the context of sustainability, the importance of bioretention systems is increasing in temperate regions as well as in tropical regions. The City of New York is an example for massive investment in bioretention systems, such as bioswales and raingardens (City of New York, 2010). Singapore’s Active Beautiful Clean (ABC) Waters Program was implemented in 2006 and includes a wide variety of urban bioretention systems (Lim and Lu, 2016).

## 3. Impact on soil organic carbon stocks

Bioswales may sequester 1.5 to 9 tons of soil organic carbon (SOC) per hectare per year in soil (Bouchard *et al.*, 2013). Because they are built from various materials, including soil, their baseline SOC stock can vary (

Table 82). Soil organic carbon accumulation rates are influenced by the type of bioswale. Such engineered bioretention systems have been found to support distinct microbial communities as compared to non-engineered urban soils, suggesting that management of these urban infrastructures may influence biogeochemical cycling as well as physical properties (Gill, Lee and McGuire, 2017)

**Table 82.** Changes in soil organic carbon stocks reported for bioretention systems

Location	Climate zone	Soil type	Baseline SOC stock (tC/ha)	Additional SOC storage (tC/ha/yr)	Duration (Years)	Depth (cm)	More information	Reference
Southeast Queensland, Australia	Subtropical	Constructed: 4 layers (drainage, transition, filter, mulch). Filter media of sandy loam or loamy sand with pH 5.5-7.5	1	3.1	13	20	Bioretention basin with <i>Carex</i> , <i>Ficinia</i> , and <i>Lomandra</i> vegetation	Kavehei <i>et al.</i> (2019)
United States of America	Subhumid, subtropical	Felsic and crystalline soils and lower coastal plain	0.05	1	21	20	Vegetated swale with grass vegetation	Bouchard <i>et al.</i> (2013)
		Topsoil material, acidic, 58% clay	2-5	0.8	15	10	Constructed stormwater wetland with grasses, sedges, and macrophytes	Moore and Hunt (2012)
Sweden	Subarctic	Podzol	NA	0.8	17	10	Constructed wet retention pond	Merriman <i>et al.</i> (2017)
	Humid continental	Entisol	NA	0.8	26	10		
Singapore	Humid tropical	Acid soil, 20% clay	NA	1.4	15	10		

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Biofiltration systems are a great improvement compared to conventional drainage systems, which consist of a network of pits and pipes and are mainly based on impervious materials. Biofiltration systems preserve soil from sealing and may counter impacts of urbanization (Prudencio and Null, 2018).

## 4.2 Minimization of threats to soil functions

Table 83. Soil threats

Soil threats	
<b>Soil erosion</b>	Due to the presence of vegetation, water infiltration is enhanced and soil loss is minimized.
<b>Nutrient imbalance and cycles</b>	The soils used for bioswale construction should be capable of retaining high amounts of phosphorus or nitrogen, which may be present in drainage water. For example, a high content of iron and aluminium in the soil may help retain phosphorus, and mulch may help retain nitrogen.
<b>Soil contamination / pollution</b>	The systems are designed to clean stormwater. They may, therefore, prevent downstream soil pollution.
<b>Soil sealing</b>	Bioretention systems prevent soil sealing and are, therefore, preferable to other stormwater management practices.
<b>Soil water management</b>	Bioretention systems have a positive effect on stormwater infiltration. Moreover, they improve water quality through retention of pollutants. They have higher removal efficiency for particulate pollutants than for dissolved pollutants (Boger <i>et al.</i> , 2018). Particulate pollutants, including Pb, Zn, and PAH, were very efficiently removed (around 90 percent) by biofiltration swales (Flanagan <i>et al.</i> , 2018). Removal efficiency was low in winter.

## 4.3 Increases in production (e.g. food/fuel/feed/timber)

Due to their limited size, biofiltration systems do not affect production.

## 4.4 Mitigation of and adaptation to climate change

Biofiltration systems may improve climate change adaptation of cities because the systems attenuate stormwater flow, which is an effect of extreme events (Prudencio and Null, 2018).



## 4.5 Socio-economic benefits

Biofiltration system installation is a low-cost multifunctional technology with relatively simple and quick implementation. Building a system using organic and mineral-waste material from within a city encourages recycling and a circular economy.

## 4.6 Additional benefits to the practice

A biofiltration system may provide cultural services, such as landscape improvement and education opportunities.

# 5. Potential drawbacks to the practice

## 5.1 Tradeoffs with other threats to soil functions

Table 84. Soil threats

Soil threats	
<b>Nutrient imbalance and cycles</b>	Bioswale establishment may lead to infiltration of high loads of nitrogen and phosphorus during stormwater events, especially if organic amendments were used.
<b>Soil contamination / pollution</b>	Pollutants may be released from artificial biofilter components if the components were not chosen adequately (Flanagan <i>et al.</i> , 2018).
<b>Soil biodiversity loss</b>	Constructed stormwater ponds may have reduced biodiversity as compared to natural ponds (Moore and Hunt, 2012).

## 5.2 Possible greenhouse gas (GHG) emissions

N<sub>2</sub>O and CH<sub>4</sub> emissions from bioretention basins were found to be low, while CO<sub>2</sub> emissions from bioretention basins may increase with increasing soil organic matter content (McPhillips, Goodale and Walter, 2018). Greenhouse gas emissions are driven by hydrologic conditions. Wet basins have higher CH<sub>4</sub> and N<sub>2</sub>O emissions than dry basins (McPhillips and Walter, 2015). Nitrification and denitrification processes can be promoted by introducing a low-permeability layer below a higher-permeability layer in the bioretention soil media (Hsieh, Davis and Needelman, 2007).

### 5.3 Conflict with other practice(s)

Bioretention facilities may conflict with conventional stormwater management strategies that are based on networks of impervious pits and pipes.

### 5.4 Decreases in production (e.g. food/fuel/feed/timber)

The material used for building bioretention facilities may have adverse effects on plant growth.

### 5.5 Other conflicts

In tropical and subtropical climates, bioretention facilities can favor mosquito proliferation and thus transmission of epidemic diseases, such as dengue-fever, zika virus disease, chikungunya, and malaria (Batalini de Macedo *et al.*, 2017).

## 6. Recommendations before implementation of the practice

The presence of vegetation was found to be more controlling than clay content for SOC accumulation in retention systems (Merriman *et al.*, 2017). The establishment of vegetation, therefore, is necessary in these systems. Littoral-shelf surface coverage around the perimeter of constructed wetland ponds should exceed the recommended 1- to 3-meter-wide swaths (Merriman *et al.*, 2017). Suitable shallow water levels can be achieved by an adjustable outlet structure and proper maintenance (Hunt *et al.*, 2011). Choice of vegetation is important. For example, plants that can use N and P effectively should be used to enhance water purification. In dry areas, local, drought-resistant species should be chosen. The tolerance of tree species to environmental conditions can differ widely; therefore, choice of trees should be adapted to soil type, sun exposure, soil moisture regime, soil pH, and climate (Tirpak *et al.*, 2018). Use of native plants should be encouraged. They were found to effectively remove nutrients from urban stormwater (Lim and Liu, 2016).

Organic matter additions in the form of compost should be limited to the amount necessary for plant growth (recommended <15 percent of volume) because a high content of organic matter leads to loss of excess P and N and thereby increases risks for water pollution and greenhouse gas emissions. If organic materials are added, materials that have C/N >20 and a low P content should be chosen. Such materials minimize excess nutrient availability and greenhouse gas production (McPhillips, Goodale and Walter, 2018). Using biochar in bioretention systems may greatly improve water purification through effective retention of organic carbon, total nitrogen, nitrates, and total dissolved phosphorus (>60 percent) and trace organic contaminants (>99 percent) (Ulrich, Loehnert and Higgins, 2017). Biodiversity in bioretention systems should be encouraged. It prevents soil erosion and is beneficial for water infiltration (Batalini de Macedo *et al.*, 2017). Biodiverse systems also increase below-ground root carbon input and SOC sequestration (Lange *et al.*, 2015). Inoculation of

bioretention systems with earthworms may benefit water infiltration and therefore enhance the function of the system in terms of stormwater attenuation.

Bioretention systems require regular maintenance to prevent surface clogging and to increase the amount and quality of stormwater runoff. The maintenance is mainly necessary to retain functioning and diverse vegetation cover (Batalini de Macedo *et al.*, 2017). To prevent the spread of disease in tropical and subtropical environments during the rainy season, weekly maintenance is required to deplete water in possible accumulation places (Erickson, Weiss and Gulliver, 2013).

## 7. Potential barriers to adoption

**Table 85.** Potential barriers to adoption

Barrier	Yes/No	
Biophysical	No	As these are artificial structures, biophysical barriers other than lack of space can be overcome.
Cultural	No	People generally favor vegetated spaces in cities (Kim and An, 2017).
Social	Yes	Public acceptance can be a barrier due to dirty appearance caused by plant litter and other organic residues (Dobbie, 2016).
Institutional	Yes	Some cities have strong stormwater management performance standards that may not be met by bioretention systems.
Legal (Right to soil)	Yes	Little regulatory control exists for design, construction, operation, and maintenance (Ashley <i>et al.</i> , 2015).
Knowledge	Yes	For optimal efficiency, technical knowledge is needed (Batalini de Macedo <i>et al.</i> , 2017).
Other	Yes	Perception that bioretention systems have higher maintenance costs than conventional stormwater retention systems can be a barrier; although this was found to be wrong (Houle <i>et al.</i> , 2013).

## References

- Ashley, R., Walker, L., D'Arcy, B., Wilson, S., Illman, S., Shaffer, P., Woods-Ballard, B. & Chatfield, P. 2015. UK sustainable drainage systems: past, present and future. *Proceedings of the Institution of Civil Engineers - Civil Engineering*, 168(3): 125–130. <https://doi.org/10.1680/cien.15.00011>
- Batalini de Macedo, M., Altair, R., Ferreira do Lago, C.A., Mendiondo, E.M. & Borges de Souza, V.C. 2017. Learning from the operation, pathology and maintenance of a bioretention system to optimize urban drainage practices. *Journal of Environmental Quality*, 15: 454–466. <https://doi.org/10.1016/j.jenvman.2017.08.023>
- Boger, A.R., Ahiablame, L., Mosase, E. & Beck, D. 2018. Effectiveness of roadside vegetated filter strips and swales at treating roadway runoff: a tutorial review. *Environmental Science: Water Research & Technology*, 4(4): 478–486. <https://doi.org/10.1039/C7EW00230K>
- Bouchard, N.R., Osmond, D.L., Winston, R.J. & Hunt, W.F. 2013. The capacity of roadside vegetated filter strips and swales to sequester carbon. *Ecological Engineering*, 54: 227–232. <https://doi.org/10.1016/j.ecoleng.2013.01.018>
- County, P.G.S. 1993. *Design Manual for Use of Bioretention in Stormwater Management*. Prince George's County (MD) Government, Department of Environmental Protection. Watershed Protection Branch, Landover, MD
- Dobbie, M. 2016. *Designing Raingardens for Community Acceptance*. Co-Operative Research Centre for Water Sensitive Cities: Melbourne, Australia.
- EPA. 2000. *Guiding Principles for Constructed Treatment Wetlands*. EPA 843-B-00003, US Environmental Protection Agency.
- Erickson, A.J., Weiss, P.T. & Gulliver, J.S. 2013. *Optimizing Stormwater Treatment Practices: a Handbook of Assessment and Maintenance*. Springer, New York.
- Flanagan, K., Branchu, P., Boudahmane, L., Caupos, E., Demare, D., Deshayes, S., Dubois, P., Meffray, L., Partibane, C., Saad, M. & Gromaire, M.-C. 2018. Field performance of two biofiltration systems treating micropollutants from road runoff. *Water Research*, 145: 562–578. <https://doi.org/10.1016/j.watres.2018.08.064>
- Gill, A.S., Lee, A. & McGuire, K.L. 2017. Phylogenetic and Functional Diversity of Total (DNA) and Expressed (RNA) Bacterial Communities in Urban Green Infrastructure Bioswale Soils. *Applied and Environmental Microbiology*, 83(16). <https://doi.org/10.1128/AEM.00287-17>
- Hunt, W.F., Greenway, M., Moore, T.C., Brown, R.A., Kennedy, S.G., Line, D.E. & Lord, W.G. 2011. Constructed Storm-Water Wetland Installation and Maintenance: Are We Getting It Right? *Journal of Irrigation and Drainage Engineering*, 137(8): 469–474. [https://doi.org/10.1061/\(ASCE\)IR.1943-4774.0000326](https://doi.org/10.1061/(ASCE)IR.1943-4774.0000326)

- Houle, J.J., Roseen, R.M., Ballestero, T.P., Puls, T.A. & Sherrard, J.** 2013. Comparison of Maintenance Cost, Labor Demands, and System Performance for LID and Conventional Stormwater Management. *Journal of Environmental Engineering*, 139(7): 932–938. [https://doi.org/10.1061/\(ASCE\)EE.1943-7870.0000698](https://doi.org/10.1061/(ASCE)EE.1943-7870.0000698)
- Hsieh, C.-H., Davis, A.P. & Needelman, B.A.** 2007. Nitrogen Removal from Urban Stormwater Runoff Through Layered Bioretention Columns. *Water Environment Research*, 79(12): 2404–2411. <https://doi.org/10.2175/106143007X183844>
- Kavehei, E., Jenkins, G. A., Lemckert, C. & Adame, M.F.** 2019. Carbon stocks and sequestration of stormwater bioretention/biofiltration basins. *Ecological Engineering*, 138: 227–236. <https://doi.org/10.1016/j.ecoleng.2019.07.006>
- Kim, S. & An, K.** 2017. Exploring psychological and aesthetic approaches of bio-retention facilities in the urban open space. *Sustainability*, 9: 2067. <https://doi.org/10.3390/su9112067>
- Lange, M., Eisenhauer, N., Sierra, C.A., Bessler, H., Engels, C., Griffiths, R.I., Mellado-Vázquez, P.G., Malik, A.A., Roy, J., Scheu, S., Steinbeiss, S., Thomson, B.C., Trumbore, S.E. & Gleixner, G.** 2015. Plant diversity increases soil microbial activity and soil carbon storage. *Nature Communications*, 6(1): 6707. <https://doi.org/10.1038/ncomms7707>
- Lim, H.S. & Lu, X.X.** 2016. Sustainable urban stormwater management in the tropics: An evaluation of Singapore’s ABC Waters Program. *Journal of Hydrology*, 538: 842–862. <https://doi.org/10.1016/j.jhydrol.2016.04.063>
- Liu, J., Sample, D.J., Bell, C. & Guan, Y.** 2014. Review and research needs of bioretention used for the treatment of urban stormwater. *Water*, 6(4): 1069–1099. <https://doi.org/10.3390/w6041069>
- McPhillips, L. & Walter, M.T.** 2015. Hydrologic conditions drive denitrification and greenhouse gas emissions in stormwater detention basins. *Ecological Engineering*, 85: 67–75. <https://doi.org/10.1016/j.ecoleng.2015.10.018>
- McPhillips, L., Goodale, C. & Walter, M.T.** 2018. Nutrient Leaching and Greenhouse Gas Emissions in Grassed Detention and Bioretention Stormwater Basins. *Journal of Sustainable Water in the Built Environment*, 4(1): 04017014. <https://doi.org/10.1061/JSWBAY.0000837>
- Merriman, L.S., Moore, T.L.C., Wang, J.W., Osmond, D.L., Al-Rubaei, A.M., Smolek, A.P., Blecken, G.T., Viklander, M. & Hunt, W.F.** 2017. Evaluation of factors affecting soil carbon sequestration services of stormwater wet retention ponds in varying climate zones. *Science of The Total Environment*, 583: 133–141. <https://doi.org/10.1016/j.scitotenv.2017.01.040>
- Moore, T.L.C. & Hunt, W.F.** 2012. Ecosystem service provision by stormwater wetlands and ponds – A means for evaluation? *Water Research*, 46(20): 6811–6823. <https://doi.org/10.1016/j.watres.2011.11.026>

**New York City.** 2010. Green Infrastructure Program. [Online] [Accessed 16 September 2020].  
<https://www1.nyc.gov/site/ddc/resources/features/2017/08/bioswales.page>

**Prudencio, L. & Null, S.E.** 2018. Stormwater management and ecosystem services: a review. *Environmental Research Letters*, 13(3): 033002. <https://doi.org/10.1088/1748-9326/aaa81a>

**Tirpak, R.A., Hathaway, J.M., Franklin, J.A. & Khojandi, A.** 2018. The Health of Trees in Bioretention: A Survey and Analysis of Influential Variables. *Journal of Sustainable Water in the Built Environment*, 4(4): 04018011. <https://doi.org/10.1061/JSWBAY.0000865>

**Ulrich, B.A., Loehnert, M. & Higgins, C.P.** 2017. Improved contaminant removal in vegetated stormwater biofilters amended with biochar. *Environmental Science: Water Research & Technology*, 3(4): 726–734.  
<https://doi.org/10.1039/C7EW00070G>

## 22. Green roofs

Cornelia Rumpel<sup>1</sup>, Jean Christophe Lata<sup>1</sup>, Claudia Marques-dos-Santos<sup>2</sup>

<sup>1</sup>CNRS, Sorbonne Université, Institute of Ecology and Environmental Sciences, Paris, France

<sup>2</sup>Instituto Superior de Agronomia de Universidade de Lisboa, Portugal

### 1. Description of the practice

A green roof is a roof that is partly or completely covered by plants and has an impermeable layer underneath. A green roof has an anti-root membrane and waterproofing, drainage, and water retention layers covered by growth substrate (Lata *et al.*, 2018). Mineral as well as organic materials are used as growth substrates for green roof construction. Green roofs may be managed intensively or extensively. Extensive green roofs are lightweight and support herbal vegetation on a shallow growing substrate that requires minimal maintenance. Intensive green roofs support heavier weight and thus a deeper layer of growing substrate and a wider variety of plant types, including herbaceous species and shrubs (Figure 15, Besir and Cuce, 2018).




Figure 15. Schematic view of green roof composition

Source: modified from Besir and Cuce (2018)

## 2. Range of applicability

Green roofs are used for limiting rainfall run-off, for aesthetics, and for thermal protection (Lata *et al.*, 2018). Green roofs may also be installed as a sustainable strategy to improve urban water availability through harvesting and purification of rainwater (Semeraro, Aretano and Pomes, 2019). In northern countries that have high rainfall, grass species may be used for green roof establishment. In drier areas (e.g. deserts), roofs are covered with biological soil crusts (biocrusts), which are composed of a complex mosaic of cyanobacteria, green algae, lichens, mosses, microfungi, and other bacteria/archaea. These biocrusts can photosynthesize when water is available, but their entire metabolism ceases in drought conditions (Paço *et al.*, 2014). Biocrusts can remain under these conditions for long periods and return to their normal functions after rain or dew events. They could be a solution for green roofs in these harsh conditions. Although governments all over the world have been promoting the establishment of extensive green roofs, the practice is poorly adopted in many cities (Paço *et al.*, 2014). In addition to improving water management, green roofs may enhance wildlife in urban areas. They were, for instance, found to support diverse wild bees (Tonietto *et al.*, 2011). Establishment of green roofs with vegetable crops has been suggested recently as a possibility to improve agricultural sustainability and food security in urban areas (Walters and Midden, 2018).

## 3. Impact on soil organic carbon stocks

Substrates used to establish green roofs may already contain organic carbon, especially if organic materials are included. Additional soil organic carbon (SOC) storage may depend on the vegetation, climate, and management practices. Intensively managed green roofs may store more SOC than extensively managed green roofs but may also require more external inputs in terms of irrigation, fertilisation, and other maintenance.

Results from studies of extensive green roofs in the United States of America indicated varying carbon storage in biomass depending on species. The average was 1.7 tC/ha. Substrate carbon storage averaged 9.1 tC/ha without species effect, representing a SOC sequestration 1.0 tC/ha/yr (Table 86, Getter *et al.*, 2009). Higher carbon storage can be achieved with the use of organic waste materials. A one-year study from China using sewage sludge for green roof construction indicated carbon storage of 18 tC/ha with an average carbon sequestration of 64 tC/ha/yr in the combined biomass and substrate organic matter (Luo *et al.*, 2015).



**Table 86.** Changes in soil organic carbon stocks reported for green roofs

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration	Depth (cm)	More information	Reference
East Lansing MI, United States of America	Temperate	XeroFolr XF-105 drainage layer and greenroof substrate (79% sand, pH 8.2)	8	1	2 years	6	<i>Sedum</i> vegetation ( <i>S. acre</i> , <i>S. album</i> , <i>S. kamtschaticum</i> , <i>S. spurium</i> )	Getter <i>et al.</i> (2009)
			31	15–208	3 years	10.2	3 types of vegetation with contrasting additional C storage: Sedum (15 t/ha) Native prairie mix (18 t/ha) Herbaceous perennials and grasses (208 t/ha)	Wittinghill and Rowe (2014)
Berlin, Germany	Temperate	-	-	0.8	1 year	9	Sedum	Heusinger and Weber (2017)
Dujiangyan, China	Subtropical, humid	Local natural soil (pH 6.3, 1.2%C)	38	3.89	1 year	25	3 plant species ( <i>Ligustrum vicaryi</i> , <i>Liriope spicata</i> , <i>Nephrolepis auriculata</i> )	Luo <i>et al.</i> (2015)
		Local natural soil amended with sewage sludge (1:1; v:v)	159	3.81				
Murcia, Spain	Semi-arid	Mixture compost/silica sand/crushed bricks (1:1:1.8; v:v:v)	3.9	0.8	10 months	10	SOC increase only under <i>Lotus creticus</i> L.; <i>Asteriscus maritimus</i> L. showed SOC loss	Ondoño, Martínez-Sánchez and Moreno (2016)
		Compost/crushed bricks (1:9; v:v)	10.9	-5.2			2 plant species ( <i>Lotus creticus</i> L. and <i>Asteriscus maritimus</i> L.)	
		Compost/Haplic calcisol/crushed bricks (1:1:1.8; v:v:v)	7.6	1.1				
		Compost/silica sand/Haplic calcisol (1:1:1.8; v:v:v)	9.7	2.3				

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Green roofs are constructed entities, and their properties are strongly dependent on the organic and mineral materials used (Dusza *et al.*, 2016). Soil properties are therefore not “improved” by this practice; rather, the establishment of green roofs adds soil functions to areas that previously supported impervious surfaces.

### 4.2 Minimization of threats to soil functions

Table 87. Soil threats

Soil threats	
Soil water management	Green roofs have positive effects on stormwater management and water purification (Shafique <i>et al.</i> , 2018) as well as on rainwater recovery, which is of particular importance in urban areas with impermeable pavements (Semeraro, Aretano and Pomes, 2019)

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Green roofs may have the potential to support urban agriculture. However, before the practice can be incorporated at scale, installation costs must be reduced, roof-weight limitations must be addressed, and appropriate management practices must be developed (Whittinghill and Rowe, 2012). Extensive green roof systems can support high productivity of lettuce, kale, and radish if sufficient moisture and nutrient inputs are available (Walters and Midden, 2018).

### 4.4 Mitigation of and adaptation to climate change

Green roofs alleviate the huge ecological footprint of cities (Getter *et al.*, 2009). This function is due to their isolating impact—reducing energy needed for cooling and heating—and to the creation of sustainable buildings. In tropical China, Hong Kong SAR, green roofs prevented 43.9 TJ of solar energy from penetrating buildings in summer (Tsang and Kim, 2011). In Italy, green roofs are used to isolate wine cellars (Conti, Barbari, and Monti, 2016). Moreover, green roofs may contribute to carbon removal from the atmosphere due to their great organic carbon sequestration potential in growth substrates. Nitrogen fixation from the atmosphere (both through free-living organisms in the substrate and symbiotic N-fixing organisms) must be ensured to avoid the need for fertilisation of the plants used.

## 4.5 Socio-economic benefits

Rainwater retention by green roofs reduces the urban heat-island effect in buildings (which results from an increase in temperature in urban environments with reduced vegetation cover); helps mitigate heat loss during the winter; increases urban biodiversity and carbon sequestration; improves air quality, soundproofing, and durability of buildings; and potentially slows the spread of fires. Green roofs therefore greatly reduce the footprint of cities through reduction of heating and cooling consumption while also contributing to rainwater retention (Karteris *et al.*, 2016).

## 4.6 Additional benefits to the practice

Green roofs provide cultural services by supplying or improving aesthetics and may have a positive effect on the psychology of urban dwellers (Lata *et al.*, 2018).

# 5. Potential drawbacks to the practice

## 5.1 Tradeoffs with other threats to soil functions

Table 88. Soil threats

Soil threats	
<b>Soil contamination / pollution</b>	Organic building materials in high concentrations (e.g. composts) could lead to water pollution due to P and N leaching (Wang, Tian and Zhao, 2017).

## 5.2 Increases in greenhouse gas emissions

The carbon footprint due to construction and maintenance has been estimated to be between 6.4 and 155.8 kg CO<sub>2</sub> eq/m<sup>2</sup>, depending on material used and management intensity. The lowest carbon footprint (6.4 kg CO<sub>2</sub> eq/m<sup>2</sup>) was found for a lightweight green roof (Chenani, Lehvävirta and Häkkinen, 2015). Green roofs may be effective CH<sub>4</sub> sinks and do not significantly affect N<sub>2</sub>O emissions (Teemusk *et al.*, 2019), but the number of studies concerning this issue is still very limited.

## 5.3 Conflict with other practice(s)

Green roofs replace traditional roofs and may thus create conflicts with other actors in the building industry.

## 5.4 Other conflicts

Green roofs have a carbon footprint due to the manufacturing process. The carbon “cost” of installing a root barrier, drainage layer, substrate, and plant material was assumed to be similar to the gravel ballast of a traditional roof (Kosareo and Ries, 2007). The use of permanent green roofs in the Mediterranean regions, which have hot and dry summers, requires the use of irrigation. The vegetation cannot survive summer without water because the installed vegetation is not native to the climatic region. Even species that are suited to the climate require irrigation to maintain growth and aesthetic appearance during the dry period. Green roofs in this context thus also have a water footprint.

## 6. Recommendations before implementation of the practice

Provision of ecosystem services of green roofs is strongly dependent on the growth substrates used (Lata *et al.*, 2018). Adequate substrate for green roof installation needs to find a compromise between an average pore size that allows for aeration of plant roots and a water holding capacity that minimizes the need for irrigation (Ondoño, Martínez-Sánchez and Moreno, 2016). Typical inorganic building materials are light, porous materials, such as volcanic rock (pumic or pozzolan), expanded clay, expanded shale, or a mixture of these elements (Lata *et al.*, 2018). However, artificial substrates are more vulnerable in terms of water loss than natural soil because they dry faster (Dusza *et al.*, 2016). An organic component of the growth substrate is needed to provide nutrients and improve physical conditions for plant growth. Although content of organic matter in green roofs may vary due to national regulations, an initial organic matter content between 10 and 20 percent seems to be optimal for plant growth (Ondoño, Martínez-Sánchez and Moreno, 2016). In the spirit of circular economy, local waste materials, such as compost and municipal sewage sludge, may be used for green roof construction. Biochar could be included in green roof substrate due to its beneficial effects on amending water quality and quantity of runoff (Qianqian *et al.*, 2019). Choice of plant species may be important as it can greatly influence the carbon sequestration potential of the roofs (Ondoño, Martínez-Sánchez and Moreno, 2016). While Sedum species are often used, SOC sequestration potential was found to be higher with thicker substrate layers when herbaceous species were included (Wittinghill and Rowe, 2012). However, this may also decrease water quality due to higher nitrogen and carbon leaching (Seidl *et al.*, 2013) and increase the maintenance effort and cost.

There are three different management systems for green roofs (Lata *et al.*, 2018):

- ◆ Extensive, without harvesting and other management activities (succulent plants, such as Sedum; light substrate; 4–15 cm thick; no irrigation)
- ◆ Semi-intensive (grasses or shrubs; light substrate; 12–30 cm thick; irrigation)
- ◆ Intensive (all plant types; natural soil; >30 cm thick; irrigation)

Moss- dominated biocrusts appear to be an innovative solution to urban landscape. These taxa do not have a root system, thereby reducing the thickness of the substrate, decreasing the installation costs, and decreasing the weight load on the building structure. The resulting product of this biological technology applied to urban landscape was designated by Paço *et al.* (2014), as MedMossRoofs (<https://www.facebook.com/medmossroofs>).

## 7. Potential barriers to adoption

**Table 89.** Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Climatic conditions in arid areas or boreal areas may impede green roof installation.
Social	Yes	Lack of awareness among public and private sectors on the benefits of green roofs (Hossain <i>et al.</i> , 2019); difficulty of acceptance from construction industry sector; negative visual appreciation of roofs when plants are too diverse or during winter.
Economic	Yes	Increased maintenance costs.
Institutional	Yes	Lack of promotion and incentives from the government; lack of guidelines (Hossain <i>et al.</i> , 2019).
Legal (Right to soil)	Yes	A planning permit may be necessary. Rooftop structures could be prohibited.
Knowledge	Yes	Many knowledge gaps exist concerning the effect of green roofs on ecosystem services related to biodiversity (e.g., pollination) and air quality (Lata <i>et al.</i> , 2018).

**Table 90.** Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Urban agriculture on rooftops in Paris, France - the T4P research project (Pilot Project of Parisian Productive Rooftops)</i>	Europe	5	6	23

## References

- Besir, A.B. & Cuce, E. 2018. Green roofs and facades: A comprehensive review. *Renewable and Sustainable Energy Reviews*, 82: 915–939. <https://doi.org/10.1016/j.rser.2017.09.106>
- Chenani, S.B., Lehvavirta, S. & Häkkinen, T. 2015. Life cycle assessment of layers of green roofs. *Journal of Cleaner Production*, 90: 153–162. <https://doi.org/10.1016/j.jclepro.2014.11.070>
- Conti, L., Barbari, M. & Monti, M. 2016. Design of Sustainable Agricultural Buildings. A Case Study of a Wine Cellar in Tuscany, Italy. *Buildings*, 6(2): 17. <https://doi.org/10.3390/buildings6020017>
- Dusza, Y., Barot, S., Kraepiel, Y., Lata, J.-C., Abbadie, L. & Raynaud, X. 2017. Multifunctionality is affected by interactions between green roof plant species, substrate depth, and substrate type. *Ecology and Evolution*, 7(7): 2357–2369. <https://doi.org/10.1002/ece3.2691>
- Getter, K.L., Rowe, D.B., Robertson, G.P., Cregg, B.M. & Andresen, J.A. 2009. Carbon sequestration potential of extensive green roofs. *Environmental Science and Technology*, 43(19): 7564–7570. <https://doi.org/10.1021/es901539x>
- Heusinger, J. & Weber, S. 2017. Extensive green roof CO<sub>2</sub> exchange and its seasonal variation quantified by eddy covariance measurements. *The Science of the Total Environment*, 607–608: 623–632. <https://doi.org/10.1016/j.scitotenv.2017.07.052>
- Hossain, M.A., Shams, S., Amin, M., Reza, M.S. & Chowdhury, T.U. 2019. Perception and Barriers to Implementation of Intensive and Extensive Green Roofs in Dhaka, Bangladesh. *Buildings*, 9(4): 79. <https://doi.org/10.3390/buildings9040079>
- Karteris, M., Theodoridou, I., Mallinis, G., Tsiros, E. & Karteris, A. 2016. Towards a green sustainable strategy for Mediterranean cities: Assessing the benefits of large-scale green roofs implementation in Thessaloniki, Northern Greece, using environmental modelling, GIS and very high spatial resolution remote sensing data. *Renewable and Sustainable Energy Reviews*, 58: 510–525. <https://doi.org/10.1016/j.rser.2015.11.098>
- Kosareo, L. & Ries, R. 2007. Comparative environmental life cycle assessment of green roofs. *Building and Environment*, 42(7): 2606–2613. <https://doi.org/10.1016/j.buildenv.2006.06.019>
- Lata, J.-C., Dusza, Y., Abbadie, L., Barot, S., Carmignac, D., Gendreau, E., Kraepiel, Y., Mériguet, J., Motard, E. & Raynaud, X. 2018. Role of substrate properties in the provision of multifunctional green roof ecosystem services. *Applied Soil Ecology*, 123: 464–468. <https://doi.org/10.1016/j.apsoil.2017.09.012>
- Luo, H., Liu, X., Anderson, B.C., Zhang, K., Li, X., Huang, B., Li, M., Mo, Y., Fan, L., Shen, Q., Chen, F. & Jiang, M. 2015. Carbon sequestration potential of green roofs using mixed-sewage-sludge substrate in Chengdu World Modern Garden City. *Ecological Indicators*, 49: 247–259. <https://doi.org/10.1016/j.ecolind.2014.10.016>
- Ondoño, S., Martínez-Sánchez, J.J. & Moreno, J.L. 2016. The composition and depth of green roof substrates affect the growth of *Silene vulgaris* and *Lagurus ovatus* species and the C and N sequestration under

two irrigation conditions. *Journal of Environmental Management*, 166: 330–340.

<https://doi.org/10.1016/j.jenvman.2015.08.045>

**Paço, T.A., Cameira, M.R., Branquinho, C., Cruz de Carvalho, R., Luís, L., Espírito-Santo, D., Valente, F., Brandão, C., Soares, A.L., Anico, A., Abreu, F. & Pereira, L.S.** 2014. Innovative green roofs for southern Europe: biocrusts and native species with low water use. In *Proceedings of the 40th IAHS World Congress on Housing–“Sustainable Housing Construction”*. Funchal, Portugal, December 16–19

**Qianqian, Z., Liping, M., Huiwei, W. & Long, W.** 2019. Analysis of the effect of green roof substrate amended with biochar on water quality and quantity of rainfall runoff. *Environmental Monitoring and Assessment*, 191(5): 304. <https://doi.org/10.1007/s10661-019-7466-4>

**Semeraro, T., Aretano, R. & Pomes, A.** 2019. Green Roof Technology as a Sustainable Strategy to Improve Water Urban Availability. *IOP Conference Series: Materials Science and Engineering*, 471: 092065. <https://doi.org/10.1088/1757-899X/471/9/092065>

**Seidl, M., Gromaire, M.C., Saad, M. & De Gouvello, B.** 2013. Effect of substrate depth and rain-event history on the pollutant abatement of green roofs. *Environ. Pollut.*, 183: 195–203. <https://doi.org/10.1016/j.envpol.2013.05.026>

**Shafique, M., Kim, R., Kyung-Ho, K., Shafique, M., Kim, R. & Kyung-Ho, K.** 2018. Green Roof for Stormwater Management in a Highly Urbanized Area: The Case of Seoul, Korea. *Sustainability*, 10(3): 584. <https://doi.org/10.3390/su10030584>

**Teemusk, A., Kull, A., Kanal, A. & Mander, Ü.** 2019. Environmental factors affecting greenhouse gas fluxes of green roofs in temperate zone. *Science of The Total Environment*, 694: 133699. <https://doi.org/10.1016/j.scitotenv.2019.133699>

**Tonietto, R., Fant, J., Ascher, J., Ellis, K. & Larkin, D.** 2011. A comparison of bee communities of Chicago green roofs, parks and prairies. *Landscape and Urban Planning*, 103(1): 102–108. <https://doi.org/10.1016/j.landurbplan.2011.07.004>

**Tsang, S.W. & Jim, C.Y.** 2011. Theoretical evaluation of thermal and energy performance of tropical green roofs. *Energy*, 36(5): 3590–3598. <https://doi.org/10.1016/j.energy.2011.03.072>

**Walters, S.A. & Stoelzle Midden, K.** 2018. Sustainability of Urban Agriculture: Vegetable Production on Green Roofs. *Agriculture*, 8(11): 168. <https://doi.org/10.3390/agriculture8110168>

**Wang, X., Tian, Y. & Zhao, X.** 2017. The influence of dual-substrate-layer extensive green roofs on rainwater runoff quantity and quality. *The Science of the Total Environment*, 592: 465–476. <https://doi.org/10.1016/j.scitotenv.2017.03.124>

**Whittinghill, L.J. & Rowe, D.B.** 2012. The role of green roof technology in urban agriculture. *Renewable Agriculture and Food Systems*, 27(4): 314–322.

## 23. Urban agriculture

John M. Galbraith<sup>1</sup>, Tatiana Morin<sup>2</sup>, Cláudia M.d.S. Cordovil<sup>3</sup>

<sup>1</sup>Virginia Tech, Blacksburg, VA, United States of America

<sup>2</sup>Urban Soil Institute, New York City, NY, United States of America

<sup>3</sup>University of Lisbon, School of Agriculture, CEF, Lisboa, Portugal

### 1. Description of the practice

Urban agriculture comprises all forms of agricultural production within or around (peri-urban) cities (Wagstaff and Wortman, 2013). It consists of growing, harvesting, processing, and distributing fresh vegetables and fruits in urban areas for personal consumption or for sale. Urban agriculture may involve urban dwellers who cultivate food in private yards, allotment gardens, common-area (community) gardens, containers, and raised beds or inside hoop houses and greenhouses. It also includes rooftop farming and urban farms run by private companies. According to FAO, urban agriculture refers to plant and animal production in an urban context, including not only small vegetable gardens but also farming activity on community land (FAO, 2010). Urban agriculture is substantial: 15 to 20 percent of the world's food is grown in urban areas (Gerster-Bentaya, 2013). In the United States, according to the 1990 Census, urban areas produced 40 percent of the dollar value of U.S. agricultural products (Deelstra and Girardet, 2000).

### 2. Range of applicability

The production of food in urban areas addresses food-insecurity issues of a growing population in global megacities. Urban agriculture is practiced by 800 million people and produces 15 to 20 percent of the world's food. The dependence on urban agriculture for food security is growing, especially in developing countries (Gallaher *et al.*, 2013; Karanja and Njenga, 2011). Supplementing food production by city farming is not new. It has been especially common during wars and depressions, when food scarcity was an issue (Burkhardt and Schneider, 2018). Nowadays, more city farming is practiced as a leisure time activity, but it is also practiced to improve personal health and economic situation. It improves personal and community access to food, creates jobs and higher incomes, improves the appearance and cohesiveness of the community, educates about



gardening and farming, and provides ecosystem services (Benth, *et al.* 2013). In some parts of Europe, the concepts of urban and peri-urban agriculture (UPA) are still a recent novelty in academic and political agendas. Regulated urban gardens, such as the regulated Municipal Urban Parks, started to emerge in 2008, largely due to a deep economic crisis. In a study conducted by Cordovil, Rodrigo and Gonçalves (2015), however, urban farming proved not to be related to unemployment. It was related to pleasure or passion for agriculture and working with the land. It was also related to a need to alter consumption habits towards the use of organic products and zero use of fertilizers. Most urban farmers have a previous link to rural areas, or have a history of farmers in their family, or both.

In many urban areas, soils and organic-growth media may be unfavourable for plant growth because of low physical quality and chemical fertility (Jim, 1998). Urban soils that were less than 20 years old in Pullman, Washington, United States of America, and Moscow, Idaho, United States of America, had lower soil quality and soil health than non-urban soils. These soils tended to be more compacted than natural soils, have a higher content of clay, and have less nitrogen mineralization, biological activity, and organic matter. However, urban soils that were 50 years old had reduced soil bulk densities, increased biological activity, and increased soil organic matter (Scharenbroch, Lloyd and Johnson-Maynard, 2005). Urban soils generally have a higher content of heavy metals (Burt *et al.*, 2014), pollutants, and artifacts than surrounding non-urban soils; however, their properties are quite variable (Pouyat *et al.*, 2002). Urban soils have higher pH and higher temperatures than comparable land uses in rural areas (Brown, Miltner and Cogger, 2012). Food production in urban agriculture often utilizes and recycles compost, bio-solids, and bio-liquids produced by human settlements (Lal, 2017). It also imports large quantities of topsoil materials from agricultural lands or forestlands. Urban farming often relies on specific engineered systems and techniques, such as “lasagne” gardening (Lanza, 1999) that consist of different layers of organic materials. Rooftop farming is practiced in many urban areas because of limited yard and garden space, better exposure to sunlight, and better ability to protect food from theft and predation.

### 3. Impact on soil organic carbon stocks

Agricultural soils have high plant production but are bare for large parts of the year, which could have contrasting effects on SOC sequestration. Many urban agriculture managers, such as gardeners, regularly add organic matter to improve their soil. Some of these soils could have elevated carbon storage compared to rural agricultural soils found in the same region (Edmondson *et al.*, 2014). Urban areas are a source of organic matter that can be added to the soil or containers to increase carbon storage over other land uses (Craul, 1999). The organic matter can be in the form of fresh and composted food waste, mulch from yard waste, tree trimmings, park maintenance waste, and hay and manure from zoos and horse stables. Brown, Miltner and Cogger (2012) reported that over 50 percent of yard wastes are land applied. Moreover, in urban areas, soil amendments may be produced by recycling of human waste products (Brown, Miltner and Cogger, 2012). Urban agriculture areas are hotspots for biodiversity of plant and animals, both below and above ground (Tresch *et al.*, 2019). This biodiversity may increase SOC sequestration in urban agricultural soils.

Table 91 reports on SOC stocks and their evolution in two trials in Europe and North America.

**Table 91.** Changes in soil organic carbon stocks reported for two urban agriculture trials

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Years)	Depth (cm)	More information	Reference
Bottrop, Germany	Temperate, oceanic	Hortic Gleysol	164	-	-	30	14 vegetable patches in allotments	Burghardt and Schneider (2018)
Youngtown, Ohio, United States of America	Temperate	Degraded, compacted urban soil from demolition site	22	14	2	10	Compost and biochar amendment on ground; tomato and sweet potato	Beniston, Lal and Mercer (2014)
			19	21			Raised bed with compost and biochar; tomato and sweet potato	

## 4. Other benefits of urban agriculture

### 4.1. Improvement of soil properties

In urban areas, soil is subjected to several deleterious effects that deplete its quality, such as sealing, compaction, erosion, and contamination by organic and inorganic pollutants (Ferreira, Walsh and Ferreira, 2018). Urban agricultural activity that includes high plant diversity may improve many physical, chemical, and biological soil properties, especially in urban gardens (Tresch *et al.*, 2019). By implementing urban agriculture, soil cover promotes restoration of soil organic matter and sequestration of soil organic carbon, reduces erosion, and improves water management. Adding organic matter to garden soils on a regular basis is a common practice to improve the physical and chemical properties of the soil. This practice results in better fertility, lower bulk density, improved soil structure, increased water-holding capacity, and decreased surface evaporation losses and crusting (Bretzel *et al.*, 2016).

## 4.2 Minimization of threats to soil functions

Table 92. Soil threats

Soil threats	
<b>Soil erosion</b>	Agricultural activities may limit soil erosion if they include maintenance of soil cover and adequate water management that prevents runoff from entering or leaving agricultural areas.
<b>Nutrient imbalance and cycles</b>	Appropriate additions of organic matter to mineral soils builds soil health by improving soil structure and promotes establishment of healthy and diverse microbial populations, allowing for better water and nutrient holding capacities (Salomon <i>et al.</i> , 2020). Cover crops and crop rotations keep nutrients and greenhouse gases from escaping. If fertilizer is applied, the use of recommended rates minimizes the threat to groundwater pollution from nitrate leaching and from nitrogen and phosphorus pollution in runoff.
<b>Soil salinization and alkalinization</b>	Some agricultural practices, such as mulching, can help to prevent surface evaporation and thus minimize salinization in arid regions.
<b>Soil contamination / pollution</b>	Increasing the content of soil organic matter attenuates contaminants through adsorption and microbial use, reducing plant uptake, leaching of pollutants, and human exposure from skin contact and ingestion (Lal, 2017; Lehmann and Kleber, 2015; Chaney, Sterrett and Mielke, 1984). Biosolids have been applied with great success to sequester carbon, increase fertility, and attenuate heavy metal contamination.
<b>Soil acidification</b>	Adding calcitic or dolomitic lime to raise pH in acidic soils offsets acidification.
<b>Soil biodiversity loss</b>	Urban agriculture minimizes soil biodiversity loss if it is based on sustainable practices (Scialabba <i>et al.</i> , 2003). Urban gardens are hotspots of biodiversity because they include many plant species (Tresch <i>et al.</i> , 2019).
<b>Soil sealing</b>	Pervious pavements and paving materials minimize soil functional losses, reduce runoff, and increase water infiltration.
<b>Soil compaction</b>	Mitigating concentrated vehicle and foot traffic when the soil is wet minimizes soil compaction. Compacted soil has high bulk density and loss of soil structure, gas exchange, water holding and movement, and root tip expansion.
<b>Soil water management</b>	Urban agricultural soils that have a high content of organic matter may improve soil structure, thus enhancing water infiltration.

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Urban agriculture brings many benefits and helps to build more resilient urban communities by ensuring stronger urban food systems (Dubelling, van Veenhuizen and Halliday, 2019). It can help to fulfil the food requirements of urban populations, especially in developing countries. Regardless of the objective of the urban agriculture (for personal use or for profit), it always contributes to a more cohesive community.

Many people plant fruit trees in their yards and share the excess. In humid areas, vegetables can be grown at a rate of over 20 000 pounds per acre (22 400 kg per ha). Community gardens and school gardens promote growing and cooking of fresh foods and sharing among community members (Cordovil, Rodrigo and Gonçalves, 2015). Education and training are provided through school programs, teachers, and trainers, such as extension agents and specialists. By using containers, rooftop gardens, and raised beds, impervious surfaces can be converted into areas for food production. Distribution of the food raised in urban areas is simple and inexpensive compared with shipping from remote sources (Lee, Lee and Lee, 2015). Non-profit organizations and charity groups help grow, glean, and distribute food in their communities.

### 4.4 Mitigation of and adaptation to climate change

Urban agriculture reduces greenhouse gas emissions by significantly reducing the amount fuel burned to transport foods from outside sources (Lee, G-G., Lee, H-W. and Lee, J-H., 2015; Kulak, Graves and Chatterton, 2013). Further reductions in greenhouse gas emissions are associated with rooftop gardens, which reduce heating and cooling costs (Kulak, Graves and Chatterton, 2013).

### 4.5 Socio-economic benefits

Urban agriculture contributes to creation of jobs, both voluntary and paid. It serves to educate young people, gives new meaning to life, and promotes respect towards the elderly. It also serves as a recycling pool for kitchen and other domestic wastes, thereby serving as a tool to clean urban waste sites (Turner, 2010). Pleasure, recreation, exercise, socialization, social inclusion, education, aesthetics, environmental improvement, and therapy are benefits of gardening and food production on larger scales (Leake, Adam-Bradford and Rigby, 2009). Growing food often involves groups of people that share common interests. Food production, preservation, storage, and use involve an ongoing learning and training process.

### 4.7 Additional benefits to the practice

In addition to being an economical way to produce food, urban agriculture can also be aesthetically pleasing and beneficial to the birds, insects, bees, earthworms, microbes, mammals, and other biology of an urban area (Benth *et al.*, 2013). The art of growing and harvesting food is a calming and peaceful way to experience outdoor areas in an otherwise stressful urban environment (Leake, Adam-Bradford and Rigby, 2009).

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

Runoff, erosion, and deep percolation following the application of excess chemical fertilizers or manure can cause both surface and groundwater pollution by nitrogen, phosphorous, and potassium. Certain herbicides, pesticides, and fungicides can build up in the soil, harm humans, or harm beneficial insects, birds, bees, and earthworms. Reading labels and following directions are advised to minimize threats. Successful food production, however, often requires a balance between controlling weeds, insects, and diseases and accepting damaged or minimal vegetables and fruits.

**Table 93.** Soil threats

Soil threats	
<b>Soil erosion</b>	Some erosion is to be expected following tillage and between growing seasons (unless cover crops are used) and in sloping land.
<b>Nutrient imbalance and cycles</b>	Excess fertilizer can drive fertilizer dependence and drive higher fluxes of CO <sub>2</sub> and other greenhouse gases (Lorenz and Lal, 2009; Oertel <i>et al.</i> , 2016). Nutrient loss results from exposed soils, tillage, overworking the soils, poor irrigation management, insufficient soil depth, and an insufficient mineral soil component (Oertel <i>et al.</i> , 2016)
<b>Soil salinization and alkalinization</b>	In areas where the only water source is saline, irrigation may lead to soil salinization—careful water management is required. Treated wastewater may also be a source of salinity.
<b>Soil contamination / pollution</b>	Excessive fertilizer use can result in nitrate and phosphate losses to waterways. Heavy metals may accumulate near vehicle traffic. In constructed growth substrates, organic matter may be lost due to leaching (Grard <i>et al.</i> , 2018). Amendments (such as composts, biosolids, and other waste products) and imported soils may be sources of contaminants (Kumar and Hundal, 2016; Haynes, Murtaza and Naidu, 2009).
<b>Soil acidification</b>	Fertilizing with sulfur or sulfate products can lead to acidification of the soil. Large additions of acidic organic matter or compost can also.
<b>Soil biodiversity loss</b>	Abandoning a farm or garden can lead to weeds taking over or to a change in land use that causes soil biodiversity loss. Dependence on inorganic fertilizers, pesticides, and herbicides; overwatering; bare soils; monoculture planting; or tilling, reworking, or replacing the soil every season leads to soil biodiversity loss.
<b>Soil sealing</b>	Soil sealing is a risk due to establishment of impervious foot- and vehicle-traffic paths or roads.
<b>Soil compaction</b>	Soils may be compacted by foot- or vehicle-traffic that are concentrated in specific areas.

## 5.2 Possible greenhouse gas (GHG) emissions

If rooftop agricultural space is installed, GHG emission may occur due to transportation of building materials. Use of high amounts of labile organic matter, such as compost, or high amounts of mineral fertilizer can lead to CO<sub>2</sub> and N<sub>2</sub>O emissions (Barthod *et al.*, 2018).

## 5.3 Conflict with other practice(s)

Urban agriculture may conflict with other urban planning priorities. Moreover, the correction of inequalities in the food system is neither inevitable nor guaranteed by urban agriculture. On the contrary, urban farms may even lead to displacement through eco-gentrification (Siegner, Sowerwine and Acey, 2018).

## 5.4 Other conflicts

Land tenure and the installation of security and anti-vandalism measures are possible sources of conflict arising from the implementation of urban farms (Turner, 2010). Conflicts between farming and other land uses, such as urbanization and municipal green spaces, are also possible.

# 6. Recommendations before implementation of the practice

Management of organic matter in soils or of engineered growth substrates used for food production is of primary importance to urban agriculture. Implementation of the practice must first consider the underlying (substrate) materials. The choice of materials depends on the site where production will take place (i.e. rooftop, garden, or urban farm). Organic materials used as growth substrates in urban agriculture include bark and composted materials, such as green (yard) wastes, municipal solid wastes, and even sewage sludge (Carlile, Cattivello and Zaccheo, 2015). “Lasagne” gardens with compost were shown to be a valuable technique for vegetable production on rooftops (Grard *et al.*, 2018).

Some urban soils have low fertility. Gardeners should avoid overusing chemical fertilizers and should instead use mature compost, which may greatly improve soil quality and allow for organic waste recycling (Bretzel *et al.*, 2016). Before compost application, soil properties should be evaluated to avoid detrimental effects, such as enhanced metal mobility (Murray, Pinchin and Macfie, 2011) and anti-germinative effects (Vidal *et al.*, 2020). As organic amendments and existing soils may be a source of contaminants, they should be tested and, if contaminated, amended to ensure safe gardening and edibles. Atmospheric dust and other potential current contamination sources (old buildings, construction activities, etc.) should be considered when establishing a proper garden or farm management plan.

The content of soil organic carbon (SOC) in soils that are used for urban agriculture in allotment gardens is strongly driven by the presence of trees (Edmondson *et al.*, 2014). The presence of trees and shrubs and the addition of carbon in the form of manure and compost are therefore recommended to maintain or increase SOC storage (Edmondson *et al.* 2014). Urban agriculture should increase soil biodiversity by using cover crops and companion plants, incorporating native crops and plants, integrating animals when possible, creating habitats, and minimizing monoculture farming, pesticides, herbicides, and commercial fertilizers (Scialabba *et al.*, 2003).

Erosion is a common threat in urban areas because of the high runoff from sealed and compacted surfaces. Poor soil structure and poor vegetative cover lead to exposure of soil particles and thereby erosion from wind and water. Managed food-plant growth and mulches can increase infiltration and minimize some of the excess exposure to wind, runoff, and erosion (Broz, Pfof and Thompson, 2017). Pavements can be made of pervious materials that still create a firm surface for walking or driving. Walking and traffic paths should be ripped or loosened to lessen their density; and organic matter should be added to promote development of soil structure. Walkways between garden beds can be covered with stepping stones or with layers of wood chips. These actions can minimize soil sealing, increase infiltration, and decrease runoff (Broz, Pfof and Thompson, 2017). Gardens on sloping ground should be terraced on the contour to decrease runoff and erosion. Water can be captured in cisterns or stormwater basins and used to water gardens. Most vegetable plants and fruits grow best in well drained soils that get regular, deep watering but do not stay saturated. Food plants, therefore, should not be grown in closed-bottom containers. Adequate water management is necessary in arid regions and in areas where irrigation water is saline. Cuevas *et al.* (2019) reviewed soil-improving cropping systems that minimize salinization and increase crop yields. Excess additions of pesticides, herbicides, and fungicides can lead to pollution and to loss of beneficial insects and biodiversity in soils (Scialabba *et al.*, 2003). Excess additions should thus be avoided; integrated pest management and use of organic gardening methods are advised whenever possible.

## 7. Potential barriers to adoption

Competition for land use in urban areas is high; and land use is regulated by zoning, ordinances, and homeowners' associations. Agriculture may be practiced on privately or publicly owned land. Biophysical, cultural, social, economic, institutional, legal, and knowledge barriers may exist for agricultural land use in urban areas.

**Table 94.** Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	Land for agriculture may not be available to urban dwellers. Fertility of urban soils may be low. Soils may contain a high concentration of toxic elements. Especially in the developing world, urban farmers may have limited access to such resources as organic matter, fertilizers, and water.
Cultural	No	Urban agriculture unites cultures and communities. It allows various cultures to share methods and to grow a variety of different crops, diversifying the palette and experiences.
Social	Yes/No	Gardens and farms nurture community and local pride and can become a social gathering place and outlet for neighborhoods. However, agricultural activities may not be accepted in some urban contexts due to the nuisance caused to neighbouring estates.
Economic	Yes	In cities, acquiring space for farming can be difficult. Significant funding may be required to secure a property and retain the right to farm year after year.
Institutional	Yes	Policy support and strategies for urban agricultural development are commonly missing in developing countries (Crush, Frayne and Pendleton, 2012).
Legal (Right to soil)	Yes	Inexperienced municipalities may enforce restrictions in soil testing or installation of structures such as high tunnels, resulting in a risk of liabilities.
Knowledge	Yes	Although gardening and farming are always a learning process best learned by experience, basic knowledge of potential crops and composting techniques requires capacity building.



## Photos of the practice



Photo 29. Urban-agriculture community gardens, Commerce, Texas, United States of America



Photo 30. Rooftop garden, Chicago Botanical Gardens, Chicago, Illinois, United States of America



Photo 31. Urban community garden, Los Angeles, California, United States of America



Photo 32. Community garden, West Hollywood, California, United States of America



Photo 33. Community garden, Governors Island, New York City, New York, United States of America

Table 95. Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n°
<i>Urban agriculture on rooftops in Paris, France - the T4P research project (Pilot Project of Parisian Productive Rooftops)</i>	Europe	5	6	23
<i>Urban Agriculture in Tacoma, Washington, United States of America</i>	North America	1	6	26

## References

- Beniston, J.W., Lal, R. & Mercer, K.L.** 2014. Assessing and managing soil quality for urban agriculture in a degraded vacant lot soil. *Land Degradation and Development*, 27: 4. <https://doi.org/10.1002/ldr.2342>
- Bretzel, F., Calderisi, M., Scatena, M. & Pini, R.** 2016. Soil quality is key for planning and managing urban allotments intended for the sustainable production of home-consumption vegetables. *Environmental Science and Pollution Research International*, 23(17): 17753–17760. <https://doi.org/10.1007/s11356-016-6819-6>
- Brown, S., Miltner, E. & Cogger, C.** 2012. Carbon sequestration potential in urban soils. Ch. 9. pp. 173–196. In Lal, R. & Augustin, B. (Eds.) *Carbon Sequestration in Urban Ecosystems*. Springer Publ., New York, NY, USA. <https://doi.org/10.1007/978-94-007-2366-5>.
- Broz, R., Pfost, D. & Thompson, A.** 2017. *Controlling runoff and erosion at urban construction sites*. MU Extension Publication, g01509, University of Missouri-Columbia. (also available at: <https://extensiondata.missouri.edu/pub/pdf/agguides/agengin/g01509.pdf>)
- Burghardt, W. & Schneider, T.** 2018. Bulk density and content, density and stock of carbon, nitrogen and heavy metals in vegetable patches and lawns of allotments gardens in the northwestern Ruhr area, Germany. *Journal of Soils and Sediments*, 18(2): 407–417. <https://doi.org/10.1007/s11368-016-1553-8>
- Burt, R., Hernandez, L., Shaw, R., Tunstead, R., Ferguson, R. & Peaslee, S.** 2014. Trace element concentration and speciation in selected urban soils in New York City. *Environmental Monitoring and Assessment*, 186(1): 195–215. <https://doi.org/10.1007/s10661-013-3366-1>
- Carlile, W.R., Cattivello, C. & Zaccheo, P.** 2015. Organic Growing Media: Constituents and Properties. *Vadose Zone Journal*, 14(6). <https://doi.org/10.2136/vzj2014.09.0125>
- Chaney, R., Sterrett, S. & Mielke, H.** 1984. The potential for heavy metal exposure from urban gardens and soils. In J.R. Preer (Ed.) *Proc. Symp. Heavy Metals in Urban Gardens*. Univ. Dist. Columbia Extension Service, 37–84. Washington, DC.
- Cordovil, C.M.d.S., Rodrigo, I. & Gonçalves, R.** 2015. Urban farming to grow a greener future. In *Proceedings of TE-O\_09. RAMIRAN 2015–16th International Conference. Rural-Urban Symbiosis*. September 8–10, 2015. Hamburg, Germany.
- Craul, P.J.** 1999. *Urban soils: applications and practices*. John Wiley & Sons Inc., New York, NY, USA. ISBN 10: 0471189030.
- Crush, J., Frayne, B. & Pendleton, W.** 2012. The Crisis of Food Insecurity in African Cities. *Journal of Hunger & Environmental Nutrition*, 7(2–3): 271–292. <https://doi.org/10.1080/19320248.2012.702448>
- Cuevas, J., Daliakopoulos, I.N., del Moral, F., Juan J., Hueso, J.J. & Tsanis, I.K.** 2019. A review of soil-improving cropping systems for soil salinization. *Agronomy*, 9(6): 295. <https://doi.org/10.3390/agronomy9060295>

- Deelstra, T. & Girardet, H.** 2000. *Urban agriculture and sustainable cities*. Pennsylvania State University. [online]. [cited 15 September 2020] <http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.168.4991&rep=rep1&type=pdf>
- FAO.** 2010. *Policy brief 10, Fighting Poverty and Hunger. What role for urban agriculture?* Department of Economics and Social Perspectives. Rome, Italy. <http://www.fao.org/docrep/012/al377e/al377e00.pdf>
- Ferreira, C.S.S., Walsh, R.P.D. & Ferreira, A.J.D.** 2018. Degradation in urban areas. *Current Opinion in Environmental Science & Health*, 5: 19–25. <https://doi.org/10.1016/j.coesh.2018.04.001>
- Gallaher, C.M., Kerr, J.M., Njenga, M., Karanja, N.K. & WinklerPrins, A.M.G.A.** 2013. Urban agriculture, social capital, and food security in the Kibera slums of Nairobi, Kenya. *Agriculture and Human Values*, 30(3): 389–404. <https://doi.org/10.1007/s10460-013-9425-y>
- Gerster-Bentaya, M.** 2013. Nutrition-sensitive urban agriculture. *Food Security*, 5(5): 723–737. <https://doi.org/10.1007/s12571-013-0295-3>
- Grard, B., Chenu, C., Manouchehri, N., Houot, S., Frascaria-Lacoste, N. & Aubry, C.** 2018. Rooftop farming on urban waste provides many ecosystem services. *Agronomy for Sustainable Development*, 38: 2. <https://doi.org/10.1007/s13593-017-0474-2>
- Haynes, R.J., Murtaza, G. & Naidu, R.** 2009. Inorganic and organic constituents and contaminants of biosolids: Implications for land application. *Advances in Agronomy*, 104(4): 165–267. [https://doi.org/10.1016/S0065-2113\(09\)04004-8](https://doi.org/10.1016/S0065-2113(09)04004-8)
- Jim, C.Y.** 1998. Urban soil characteristics and limitations for landscape planting in Hong Kong. *Landscape and Urban Planning*, 40: 235–249. [https://doi.org/10.1016/S0169-2046\(97\)00117-5](https://doi.org/10.1016/S0169-2046(97)00117-5)
- Karanja, N. & Njenga, M.** 2011. Feeding the cities. In Linda Starke (Ed.) *State of the world: Innovations that nourish the planet*. pp. 109–117. The World Watch Institute, Washington, D.C.
- Kulak, M., Graves, A. & Chatterton, J.** 2013. Reducing greenhouse gas emissions with urban agriculture: A life cycle assessment perspective. *Landscape and Urban Planning*, 111: 68–78. <https://doi.org/10.1016/j.landurbplan.2012.11.007>
- Kumar, K. & Hundal, L.S.** 2016. Soil in the city: sustainably improving urban soils. *Journal of Environmental Quality*, 45: 2–8. <https://doi.org/10.2134/jeq2015.11.0589>
- Lal, R.** 2017. Urban agriculture and food security. In Levin, M.J. *et al.* (eds.) *Soils within Cities*. Catena Soil Sciences, pp. 177.
- Lanza, P.** 1999. *Lasagna Gardening: A New Layering System for Bountiful Gardens: No Digging, No Tilling, No Weeding, No Kidding!* Rodale Press Inc., Emmaus. ISBN 10: 0875967957.
- Leake, J.R., Adam-Bradford, A. & Rigby, J.E.** 2009. Health benefits of ‘grow your own’ food in urban areas: implications for contaminated land risk assessment and risk management? *Environmental Health*, 8(1): S6. <https://doi.org/10.1186/1476-069X-8-S1-S6>

- Lee, G.-G., Lee, H.-W. & Lee, J.-H.** 2015. Greenhouse gas emission reduction effect in the transportation sector by urban agriculture in Seoul, Korea. *Landscape and Urban Planning*, 140: 1-7. <https://doi.org/10.1016/j.landurbplan.2015.03.012>
- Lehmann, J. & Kleber, M.** 2015. The contentious nature of soil organic matter. *Nature*, 528: 60-68. <https://doi.org/10.1038/nature16069>
- Lorenz, K. & Lal, R.,** 2009. Biogeochemical C and N cycles in urban soils. *Environment International*, 35: 1-8. <https://doi.org/10.1016/j.envint.2008.05.006>
- Murray, H., Pinchin, T.A. & Macfie, S.M.** 2011. Compost application affects metal uptake in plants grown in urban garden soils and potential human health risk. *Journal of Soils and Sediments*, 11: 815-829. <https://doi.org/10.1007/s11368-011-0359-y>
- Oertel, C., Matschullat, J., Zurba, K., Zimmermann, F. & Erasmi, S.** 2016. Greenhouse gas emissions from soils—A review. *Geochemistry*, 76(3): 327-352. <https://doi.org/10.1016/j.chemcr.2016.04.002>
- Pouyat, R., Groffman, P., Yesilonis, I. & Hernandez, L.** 2002. Soil carbon pools and fluxes in urban ecosystems. *Environmental Pollution*, 116: S107-S118. [https://doi.org/10.1016/S0269-7491\(01\)00263-9](https://doi.org/10.1016/S0269-7491(01)00263-9)
- Salomon, M.J., Watts-Williams, S.J., McLaughlin, M.J. & Cavagnaro, T.R.** 2020. Urban soil health: A city-wide survey of chemical and biological properties of urban agriculture soils. *Journal of Cleaner Production*, 275. <https://doi.org/10.1016/j.jclepro.2020.122900>.
- Siegner, A., Sowerwine, J. & Acey, C.** 2018. Does Urban Agriculture Improve Food Security? Examining the Nexus of Food Access and Distribution of Urban Produced Foods in the United States: A Systematic Review. *Sustainability*, 10(9): 2988. <https://doi.org/10.3390/su10092988>
- Scharenbroch, B.C., Lloyd, J.E. & Johnson-Maynard, J.L.** 2005. Distinguishing urban soils with physical, chemical, and biological properties. *Pedobiologia*, 49(4): 283-296. <https://doi.org/10.1016/j.pedobi.2004.12.002>
- Scialabba, N., El-Hage, Grandi, C. & Henatsch, C.** 2003. Case study no. 4: Organic agriculture and genetic resources for food and agriculture. In *Biodiversity and the Ecosystem Approach in Agriculture, Forestry and Fisheries. Proceedings of the Ninth Regular Session of the Commission on Genetic Resources for Food and Agriculture*. Rome, Italy. 12-13 October 2002. Food and Agriculture Organization. (also available at: <http://www.fao.org/3/y4586e/y4586e05.htm#>)
- Tresch, S., Frey, D., Bayon, R.-C.L., Mäder, P., Stehle, B., Fliessbach, A. & Moretti, M.** 2019. Direct and indirect effects of urban gardening on aboveground and belowground diversity influencing soil multifunctionality. *Scientific Reports*, 9(1): 9769. <https://doi.org/10.1038/s41598-019-46024-y>
- Turner, A.H.** 2010. *Establishing Urban Agriculture in Your Community: What You Need to Know Before You Get Your Hands Dirty*. Practice guide #27. University of Louisville, US. 29 p.
- Wagstaff, R.K. & Wortman, S.E.** 2015. Crop physiological response across the Chicago metropolitan region: Developing recommendations for urban and peri-urban farmers in the North Central US. *Renewable Agriculture and Food Systems*, 30(1): 8-14. <https://doi.org/10.1017/S174217051300046X>

## 24. Urban forestry

Jennifer Mason<sup>1</sup>, Thomas Meixner<sup>2</sup>, Cornelia Rumpel<sup>3</sup>, Jean Christophe Lata<sup>3</sup>, John M. Galbraith<sup>4</sup>

<sup>1</sup>USDA-NRCS Soil Survey Office Leader, Clinton, TN, the United States of America

<sup>2</sup>Department Head, Hydrology and Atmospheric Sciences, Univ. of Arizona, the United States of America

<sup>3</sup>CNRS, Sorbonne Université, Institute of Ecology and Environmental sciences, Paris, France

<sup>4</sup>Virginia Tech, Blacksburg, VA, the United States of America

### 1. Description of the practice

Urban forestry is the management of single trees and forest resources, such as woodlands, in and around urban settings for the benefits they provide society. Urban trees are part of the urban green infrastructure, and they provide physiological, sociological, economic, and aesthetic benefits (Konijnendijk *et al.*, 2006). Proper forest management that promotes stand structure and production and minimizes soil disturbance also promotes retention of soil organic carbon (SOC) (Jandl *et al.*, 2007). Urban areas of all sizes with human populations of 50 000 and higher have included trees in urban gardens, parks, and lawns, and many have preserved forests and woodlands. While trees cover 10–67 percent of urban areas in the United States of America (Edmondson *et al.*, 2014), most cities in Europe have a tree density of 50–80 street trees per 1000 inhabitants (Pauleit *et al.*, 2002). In a major African city like Addis Ababa, urban forest covers <10 percent of the surface and consists mainly of introduced species intended to satisfy local wood demand (Woldegerima, Yeshitela and Lindley, 2017).

### 2. Range of applicability

Trees in urban areas range in extent from natural forests, planted woodlots, and tree-lined avenues down to individual trees planted in soil containers. Containerized trees line city streets and are grown on buildings and rooftops. Trees in urban areas often are exposed to stressful growing conditions, such as excess heat, air pollution, inadequate water and air supply (Fite *et al.*, 2011), inadequate soil pH, poor soil quality, heavy metals

and other pollutants, and scarce availability of nutrients. Trees frequently are damaged by humans, animals, and cars. Root systems and trunks are constricted by pavements and sidewalks and are exposed to invasive diseases, insects, and other pests. This situation may be worsened by climate change because urban trees worldwide are composed of few species with low genetic variability of the species and varieties (Lohr, Kendal and Dobbs, 2016). Despite the stressful environment in which they grow, trees add substantial environmental functions and services to urban ecosystems.

### 3. Impact on soil organic carbon stocks

In the United States of America, the national average carbon storage density is 25.1 tC/ha for urban forests, compared with 53.5 tC/ha in forest stands (Nowak and Crane, 2002). Carbon storage in urban forests was found to vary between 15 and 160 tC/ha down to 1 m depth in cities in the United States of America and in China (Nowak *et al.*, 2013; Pouyat, Yesilonis and Novak, 2006). It was found to increase with increasing forest age (Table 96). Urban forest soils may contain greater proportions of recalcitrant carbon due to poorer quality leaf litter, enhanced mineralization of readily available carbon, and higher soil temperatures (Groffman *et al.*, 1995). Increases in urban soil carbon storage were observed by Pouyat *et al.* (2002) in New York City, where soil carbon density was about 30 percent higher in the urban forest than in suburban and rural forest soils. They also cited increased heavy metal and salt concentrations in urban forest soils. Higher soil carbon may be due to decreased respiration (Koerner and Klopatek, 2010) or reduced degradability of litter material (Pouyat *et al.*, 2002). However, the soil organic carbon storage under urban forests was much lower than under natural forests in the same region of Northeast China and increased with increasing age of the cities (Lv *et al.*, 2016). Studies have found that SOC is dependent on tree density (Mexia *et al.*, 2018) and on tree species or genus (Scharenbroch, 2012; Edmondson *et al.*, 2014). Urban trees that live longer and reach larger sizes have greater potential to sequester more carbon relative to shorter-lived, smaller trees (Nowak *et al.*, 2002).

Species mixture has been investigated as a strategy for conserving SOC in urban forests. This management technique is also associated with ecosystem resilience. However, the rate of carbon accumulation and its distribution within the soil profile differs between tree species. SOC enhancement was found to be highest under *Fraxinus* and *Acer spp.* and lower under *Quercus* and mixed woodland (Edmondson *et al.*, 2014).

For managed urban forests, Yoon *et al.* (2016) outline a strategy to enhance SOC by leaving litter and grass clippings in place to offset the loss of organic matter supplies. Application of compost, mulching, or both could increase urban soil carbon storage (Brown, Miltner and Cogger, 2012; Beesley *et al.*, 2012), as could application of biochar; however, it is not known how management practices might control inorganic carbon from carbonate reactions (Lorenz and Lal, 2015). Application of residual organic matter, such as yard and food waste and biosolids, to pervious surfaces in Tacoma, Washington, United States of America, would result in an annual carbon sequestration rate of 0.22 tC/ha/yr. This rate is similar to rates observed for no-till agriculture (Brown, Miltner and Cogger, 2012). Similarly, in arid settings, leaving plant material behind during maintenance has been shown to increase soil carbon storage (Rockhill, 2017; Swartz, 2019).

In arid regions, soils beneath the urban forest often have significantly higher soil carbon than the surrounding natural landscapes (Rockhill, 2017). This pattern of urban soil carbon content being higher in dry climates but lower in more humid settings has been documented as part of a broader convergence hypothesis for soil biogeochemistry in urban landscapes of North America (Trammell *et al.*, 2020).



**Table 96.** Changes in soil organic carbon stocks reported for urban forests from diverse regions

Location	Climate zone	Soil type	Baseline C stock (tC/ha)	Additional C storage (tC/ha/yr)	Duration (Years)	Depth (cm)	More information	Reference
Cities in United States of America and Canada	Temperate-subtropical	-	-	0.20-1.23	-	100	17 urban forests	Nowak <i>et al.</i> (2013) Pouyat, Yesilonis and Novak, (2006)
Shanghai, China	Subtropical	Alluvial	77.3	0.6	22		Young-near mature <i>Metasequoia glyptostroboides</i> stands	ChunBo <i>et al.</i> (2010)
Kumasi, Ghana	Tropical	Haplic Alisol and Lithic Leptosol	45.5	-	-	60	Plantations increase SOC storage (83.5 tC/ha)	Nero <i>et al.</i> (2017)
Harbin city, NW China	Temperate, cold	Luvic Phaeozem	50	0.15	100	20	219 plots in urban forests	Lv <i>et al.</i> (2016)

## 4. Other benefits of the practice

### 4.1. Improvement of soil properties

Edmondson *et al.* (2014) studied soils under three genera (*Fraxinus*, *Acer*, and *Quercus*) in Leicester, England, and found that tree cover type did not influence soil bulk density or C/N ratio. However, Livesley *et al.* (2016) studied trees, grass areas, and turfgrass in sandy urban soils in Melbourne, Australia, and found that the soils under tree canopy were less compacted ( $1.07 \text{ g/cm}^3$ ) than in grassy areas ( $1.32 \text{ g/cm}^3$ ) and had higher mean C/N ratio.

## 4.2 Minimization of threats to soil functions

Erosion is a common threat in urban areas because of the high runoff from sealed, compacted, or bare surfaces. Erosion can result from the loss of leaf litter in shady areas because of excessive earthworm activity (Pouyat, Yesilonis and Novak, 2006) and from tree or shrub density shading out understory perennial ground plane vegetation. Poor soil structure and poor vegetative cover expose soil particles to the hazard of erosion by wind and water. Properly designed and managed planting areas can increase infiltration and minimize soil exposure to wind, runoff, and erosion. Pavements can be made of pervious materials that create a firm surface for walking or parking under trees but still allow infiltration and decrease runoff. Supplemental water can maintain tree growth in semi-arid and arid regions and in humid areas where pavements limit infiltration of rainwater. Excess additions of herbicides, pesticides, and fungicides can lead to loss of beneficial insects and biodiversity in soils. Hydrophobicity of soil and litter under trees increase runoff from rainfall at the beginning of storm events. Removal of leaf litter soon after fall is recommended but must be compensated for by other runoff control measures.

**Table 97. Soil threats**

Soil threats	
<b>Soil erosion</b>	Depending on tree species, soil erosion can be reduced by tree planting in urban areas, especially in sloping areas (Woldegerima, Yeshitela and Lindley, 2017).
<b>Nutrient imbalance and cycles</b>	Fertilizing at recommended rates minimizes the threat of groundwater pollution from nitrate leaching and from nitrogen and phosphorus pollution in runoff. Net nitrogen mineralization rates in samples from the forest floor and A horizon vary by tree type (White and McDonnell, 1988). Trees can reduce oxidized nitrogen, reactive phosphorous, and stormwater nitrogen pollution (Denman, May and Moore, 2016).
<b>Soil contamination / pollution</b>	Planting trees may reduce the risk of soil contamination and in particular enhance the breakdown of organic contaminants (Dickson <i>et al.</i> , 2000).
<b>Soil biodiversity loss</b>	The importance of urban forest as a reserve of biodiversity will increase in the future (Alvey, 2006).
<b>Soil sealing</b>	Establishing urban woodland and trees prevents soil from sealing.
<b>Soil compaction</b>	Soil compaction may be reduced following tree planting.
<b>Soil water management</b>	Trees in flood plains slow down the velocity and increase turbulence of floodwaters, mitigating the harmful effects of flooding.

### 4.3 Increases in production (e.g. food/fuel/feed/timber)

Very little timber is harvested from individual urban trees for purposes other than firewood or making furniture because of the high cost to cut the tree safely and transport the pieces away using heavy machinery. Also, in places, production is low due to trace metal pollution. Most of the poor-quality material is used for firewood or made into mulch, especially in developing countries (Woldegerima, Yeshitela and Lindley, 2017). Woodlots or forests that have become assimilated into expanding urban areas may have timber values, especially in the southeastern United States where commercial stands of pine are common.

### 4.4 Mitigation of and adaptation to climate change

Urban forests also provide climate change mitigation services, including carbon sequestration and the reduction of the urban heat-island effect (Kleerekoper, van Esch and Salcedo, 2012). Temperature reductions to the heat island can be as much as 6 °C due to vegetation canopy (Shiflett *et al.*, 2017). Deciduous trees on southern and westerly aspects of urban buildings in the Northern hemisphere may potentially reduce energy emissions by shading in summer (McPherson, 1994).

### 4.5 Socioeconomic benefits

Urban forests provide a variety of essential ecosystem services, including decreasing air, water, and noise pollution, mitigating flood risk, and providing recreational areas (Escobedo, Kroeger and Wagner, 2011; Roy, Byrne and Pickering, 2012).

The economic benefits of urban forests have been quantified in a recent study. According to the authors, the economic benefit increases with tree cover and amounts to 0.93 million USD savings in air pollution health care costs, USD 20000 by capturing water runoff, and USD 478000 in building energy heating and cooling savings (Endreny *et al.*, 2017).

### 4.6 Additional benefits to the practice

Urban trees and woodlands can act as biological filters and remove airborne pollutants and thus improve air quality in urban areas (Beckett, Freer-Smith and Taylor, 1998).

## 5. Potential drawbacks to the practice

### 5.1 Tradeoffs with other threats to soil functions

Repeated timber harvesting and similar management practices can cause soil compaction and erosion. Following best management practices prescribed by conservation agencies and cooperative extension specialists is recommended.

**Table 98.** Soil threats

Soil threats	
<b>Soil erosion</b>	Commercial timber production or removal of stumps may cause soil erosion. Exposed and non-vegetated soil under urban forest and individual tree soils in semi-arid and non-arid climates are vulnerable to erosion. Excess recreation under tree canopies can result in exposure of the surface soil to erosion.
<b>Soil salinization and alkalinization</b>	Soil salinization of trees in golf courses is increased when saline groundwater is used to water soils that have surface compaction, restricted subsoil permeability, and higher clay contents (Miyomoto and Chacon, 2006).
<b>Soil contamination / pollution</b>	Tree canopies may capture atmospheric dust, which is a potential source of contamination.
<b>Soil acidification</b>	Urban trees that have highly acid needles (juniper, pines, spruce, larch, fir, other evergreens) may acidify the soil. Fertilizing with sulfur or sulfate products can lead to acidification of the soil, as can large additions of acid-producing leaf litter or compost from such matter.
<b>Soil compaction</b>	Excess recreational activities under tree canopies can result in compaction.
<b>Soil water management</b>	Trees are not suitable for growth within stormwater retention basins or on onsite waste disposal drain fields and must be removed if they grow there. All forest-floor and A-horizon samples from an urban forest study in New York City were extremely hydrophobic (White and McDonnell, 1988). Tree planting may lead to enhanced evaporation and thus less water in soil.

## 5.2 Increases in greenhouse gas emissions

Greenhouse gas emissions are associated with timber harvesting that uses mechanized equipment, such as chainsaws, skidders, and haul trucks. Cutting trees reduces the amount of carbon dioxide removal until the biomass regains its former level. Waste products and firewood that are burned release carbon dioxide into the atmosphere.

## 5.3 Conflict with other practice(s)

Forestry in urban areas can conflict with other land use choices that are higher producers of income or are deemed to be more valuable, such as housing and business developments, landfills, and transportation projects.

## 5.4 Decreases in production (e.g. food/fuel/feed/timber)

Soil compaction from recreational or sports activities may be harmful to nearby tree root systems. Construction of infrastructure and damage by impact from vehicles is harmful to tree growth. Air and water pollution reduce growth and make trees more vulnerable to diseases and pests. Invasive diseases and insects threaten native species that are not adapted to the threat.

# 6. Recommendations before implementation of the practice

Relatively few studies specifically address soil carbon management in urban forests. However, long-term studies of forest management strategies of rural forests recommend increasing productivity, e.g., through afforestation and planting of fast-growing tree species. Increased productivity has immediate effects on SOC by incorporating CO<sub>2</sub> in plant biomass and increasing carbon input to the soil (Jandl *et al.*, 2007). Urban sites are often characterized by limited rooting space and compacted or polluted soil, poor drainage, and high variability (Day and Bassuk, 1994), which need to be amended before trees can be planted. Species choice is important. Site-specific properties need to be matched to species performance. Eucalyptus species were reported to have high erosion incidence in arid sloping land, while mixed forests performed better and also increase biodiversity (Woldegerima, Yeshitela and Lindley, 2017). Foot and vehicle traffic should be concentrated in a few areas rather than underneath trees. When sidewalks and pavements must be added, pervious materials should be used if possible. Soils that have been compacted by foot and vehicle traffic should be ripped or loosened to lessen the density, and organic matter should be added to promote soil structure development. Walkways beneath trees can be covered with layers of wood chips to minimize compaction.

## 7. Potential barriers to adoption

Table 99. Potential barriers to adoption

Barrier	YES/NO	
Biophysical	Yes	In more arid regions, availability of water can be an impediment to the benefits of urban forests.
Economic	Yes	Excessive tree growth in humid areas may conflict with views from buildings and need costly thinning or removal. Cost of tree establishment and maintenance may be a barrier to planting, especially in arid and semi-arid areas where trees are not native.
Institutional	Yes	Inadequate funding and lack of political and public support despite community interest were identified as main barriers to urban forestry.
Legal (Right to soil)	Yes	Damage to infrastructure and personal property from falling trees may cause litigation.
Other	Yes	Inadequate care and maintenance of trees after planting can lead to high mortality rates.

## Photos of the practice



Photo 34. Tree growing in a containerized bunker in New York City. Beneath the pavement is a subway line



Photo 35. Trees surrounded by pervious brick pavement



Photo 36. Trees lining an urban park in New York City (Central Park)



Photo 37. Urban woodlots

Table 100. Related case studies available in volumes 4 and 6

Title	Region	Duration of study (Years)	Volume	Case-study n <sup>o</sup>
<i>Carbon storage in soils built from waste for tree plantation in Angers, France</i>	Europe	3	6	22
<i>Urban Forestry effects on soil carbon in Leicester, United Kingdom of Great Britain and Northern Ireland</i>	Europe	20 to 100	6	25
<i>Soil Organic Carbon in forested and non-forested urban plots in the Chicagoland Region, United States of America</i>	North America	Various	6	27
<i>Compost application to restore post-disturbance soil health in Montgomery county, Virginia, United States</i>	North America	4	6	28




## References

- Alvey, A.A. 2006. Promoting and preserving biodiversity in the urban forest. *Urban Forestry & Urban Greening*, 5(4): 195–201. <https://doi.org/10.1016/j.ufug.2006.09.003>
- Beckett, K.P., Freer-Smith, P.H. & Taylor, G. 1998. Urban woodlands: their role in reducing the effects of particulate pollution. *Environmental Pollution*, 99(3): 347–360. [https://doi.org/10.1016/S0269-7491\(98\)00016-5](https://doi.org/10.1016/S0269-7491(98)00016-5)
- Beesley, L. 2012. Carbon storage and fluxes in existing and newly created urban soils. *Journal of Environmental Management*, 104: 158–165. <https://doi.org/10.1016/j.jenvman.2012.03.024>
- Brown, S., Miltner, E. & Cogger, C. 2012. Carbon sequestration potential in urban soils. In Lal, R., Augustin, B. (Eds.) *Carbon Sequestration in Urban Ecosystems*. Springer: New York, NY. 47: 173–196.
- ChunBo, X., Hai, W., KaiFeng, F., Becuwe, X., YuJie, H., HongZhong, K. & ChunJiang, L. 2010. Carbon storage of Metasequoia glyptostroboides plantation ecosystems at different age stages in Chongming Island, East China. *Journal of Shanghai Jiaotong University - Agricultural Science*, 28(1): 30–34.
- Day, S.D. & Bassuk, N.L., 1994. A review of the effects of soil compaction and amelioration treatments on landscape trees. *Journal of Arboriculture*, 20: 9–16.
- Denman, E.C., May, P.B. & Moore, G.M. 2016. The Potential Role of Urban Forests in Removing Nutrients from Stormwater. *Journal of Environmental Quality*, 45(1): 207–214. <https://doi.org/10.2134/jeq2015.01.0047>
- Dickson, N.M., Mackay, J.M. & Goodman, A., Putwain, P. 2000. Planting trees on contaminated soils: issues and guidelines. *Land Contamination and Reclamation*, 8: 87–97. <https://doi.org/10.2462/09670513.561>
- Edmondson, J.L., O’Sullivan, O.S., Inger, R., Potter, J., McHugh, N., Gaston, K.J. & Leake, J.R. 2014. Urban Tree Effects on Soil Organic Carbon. *PLoS ONE*, 9(7): e101872. <https://doi.org/10.1371/journal.pone.0101872>
- Endreny, T., Santagata, R., Perna, A., Stefano, C., Rallo, R. & Ulgiati, S. 2017. Implementing and managing urban forests: A much needed conservation strategy to increase ecosystem services and urban wellbeing. *Ecological Modelling*, 360: 328–335. <https://doi.org/10.1016/j.ecolmodel.2017.07.016>
- Escobedo, F.J., Kroeger, T. & Wagner, J.E. 2011. Urban forests and pollution mitigation: analyzing ecosystem services and disservices. *Environmental Pollution*, 59(8-9): 2078–87. <https://doi.org/10.1016/j.envpol.2011.01.010>
- Fite, K., Smiley, E.T., McIntyre, J. & Wells, C.E. 2011. Evaluation of a Soil Decompaction and Amendment Process for Urban Trees. *Arboriculture & Urban Forestry*, 37(6): 293–300.
- Groffman, P.M., Pouyat, R.V., McDonnell, M.J., Pickett, S.T.A., Zipperer, W.C., Pouyat, R.V., McDonnell, M.J., Pickett, S.T.A. & Zipperer, W.C. 1995. Carbon pools and trace gas fluxes in urban forest soils. In Lal, R. et al. (Eds.) *Soil Management and the Greenhouse Effect*. CRC Press, Boca Raton, FL.

- Jandl, R., Lindner, M., Vesterdal, L., Bauwens, B., Baritz, R., Hagedorn, F., Johnson, D.W., Minkkinen, K. & Byrne, K.A. 2007. How strongly can forest management influence soil carbon sequestration? *Geoderma*, 137(3-4): 253–268. <https://doi.org/10.1016/j.geoderma.2006.09.003>
- Kleerekoper, L., van Esch, M. & Salcedo, T.B. 2012. How to make a city climate-proof, addressing the urban heat island effect. *Resources, Conservation and Recycling*, 64: 30–38. <https://doi.org/10.1016/j.resconrec.2011.06.004>
- Koerner, B.A. & Klopatek, J.M. 2010. Carbon fluxes and nitrogen availability along urban-rural gradient in a desert landscape. *Urban Ecosystems*, 13: 1–21. <https://doi.org/10.1007/s11252-009-0105-z>
- Konijnendijk, C., Ricard, R., Kenney, A. & Randrup, T. 2006. Defining urban forestry—A comparative perspective of North America and Europe. *Urban Forestry & Urban Greening*, 4(3-4): 93–103. <https://doi.org/10.1016/j.ufug.2005.11.003>
- Livesley, S.J., Ossola, A., Threlfall, C.G., Hahs, A.K. & Williams, N.S.G. 2016. Soil Carbon and Carbon/Nitrogen Ratio Change under Tree Canopy, Tall Grass, and Turf Grass Areas of Urban Green Space. *Journal of Environmental Quality*, 45(1): 215–223. <https://doi.org/10.2134/jeq2015.03.0121>
- Lohr, V.I., Kendal, D., Dobbs, C., 2016. Urban trees worldwide have low species and genetic diversity, posing high risks of tree loss as stresses from climate change increase. *Acta horticultrae*, 108: 263–270. <https://doi.org/10.17660/ActaHortic.2016.1108.34>
- Lorenz, K. & Lal, R. 2015. Managing soil carbon stocks to enhance the resilience of urban ecosystems. *Carbon Management*, 6(1–2): 35–50. <https://doi.org/10.1080/17583004.2015.1071182>
- Lv, H., Wang, W., He, X., Xiao, L., Zhou, W. & Zhang, B. 2016. Quantifying Tree and Soil Carbon Stocks in a Temperate Urban Forest in Northeast China. *Forests*, 7(9): 200. <https://doi.org/10.3390/f7090200>
- McPherson, E.G. 1994. Cooling urban heat islands with sustainable landscapes. In Platt, Rutherford, H., Rowntree, Rowan, A., Muick, Pamela, C. (Eds.) *The ecological city: preserving and restoring urban biodiversity*. Amherst, MA: University of Massachusetts Press: 151–171.
- Mexia, T., Vieira, J., Príncipe, A., Anjos, A., Silva, P., Lopes, N., Freitas, C., Santos-Reis, M., Correia, O., Branquinho, C. & Pinho, P. 2018. Ecosystem services: Urban parks under a magnifying glass. *Environmental Research*, 160: 469–478. <https://doi.org/10.1016/j.envres.2017.10.023>
- Nero, B.F., Callo-Concha, D., Anning, A. & Denich, M. 2017. Urban green spaces enhance climate change mitigation in cities of the global south: the case of Kumasi, Ghana. *Procedia Engineering*, 198: 69–83. <https://doi.org/10.1016/j.proeng.2017.07.074>
- Nowak, D.J., Stevens, J.C., Sisinni, S.M. & Luley, C.J. 2002. Effects of urban tree management and species selection on atmospheric carbon dioxide. *Journal of Arboriculture*. 28(3): 113–122.
- Nowak, D.J., Greenfield, E.J., Hoehn, R.E. & Lapoint, E. 2013. Carbon storage and sequestration by trees in urban and community areas of the United States. *Environmental Pollution*, 178: 229–236. <https://doi.org/10.1016/j.envpol.2013.03.019>

- Nowak, D. & Crane, D.** 2002. Carbon storage and sequestration by urban trees in the USA. *Environmental Pollution*, 116(3): 381–389. [https://doi.org/10.1016/S0269-7491\(01\)00214-7](https://doi.org/10.1016/S0269-7491(01)00214-7)
- Pauleit, S., Jones, N., Garcia-Martin, G., Garcia-Valdecantos, J.L., Rivière, L.M., Vidal-Beaudet, L., Bodson, M. & Randrup, T.B.** 2002. Tree establishment practice in towns and cities – Results from a European survey. *Urban Forestry & Urban Greening*, 1(2): 83–96. <https://doi.org/10.1078/1618-8667-00009>
- Pouyat, R.V., Yesilonis, I.D. & Novak, D.J.** 2006. Carbon storage by urban soils in the United States. *J. Environmental Quality* 35: 1566–1575. <https://doi.org/10.2134/jeq2005.0215>.
- Pouyat, R. Groffman, P., Yesilonis, I. & Hernandez, L.** 2002. Soil carbon pools and fluxes in urban ecosystems. *Environmental Pollution*, 116: 107–118. [https://doi.org/10.1016/S0269-7491\(01\)00263-9](https://doi.org/10.1016/S0269-7491(01)00263-9)
- Rockhill, T.** 2017. *Influence of Soil Physical and Chemical Properties on Soil CO<sub>2</sub> Flux in Semi-Arid Green Stormwater Infrastructure*. University of Arizona.
- Roy, S., Byrne, J. & Pickering, C.** 2012. A systematic quantitative review of urban tree benefits, costs, and assessment methods across cities in different climatic zones. *Urban Forestry and Urban Greening*, 11(4): 351–363. <https://doi.org/10.1016/j.ufug.2012.06.006>
- Scharenbroch, B.C.** 2012. Urban Trees for Carbon Sequestration. In Lal R., Augustin B. (Eds.) *Carbon Sequestration in Urban Ecosystems*, pp. 121-138. Springer, Dordrecht, Netherlands. [https://doi.org/10.1007/978-94-007-2366-5\\_6](https://doi.org/10.1007/978-94-007-2366-5_6)
- Shiflett, S.A., Liang, L.L., Crum, S.M., Feyisa, G.L., Wang, J. & Jenerette, G.D.** 2017. Variation in the urban vegetation, surface temperature, air temperature nexus. *Science of the Total Environment*, 579: 495–505. <https://doi.org/10.1016/j.scitotenv.2016.11.069>
- Swartz, S.** 2019. *Infiltration rates of green infrastructure curb-cut basins: finding balance between functions and aesthetic*. University of Arizona.
- Trammell, T.L.E., Pataki, D.E., Pouyat, R.V., Groffman, P.M., Rosier, C., Bettez, N., Cavender-Bares, J., Grove, M.J., Hall, S.J., Heffernan, J., Hobbie, S.E., Morse, J.L., Neill, C. & Steele, M.** 2020. Urban soil carbon and nitrogen converge at a continental scale. *Ecological Monographs*, 90(2): e01401. <https://doi.org/10.1002/ecm.1401>
- White, C.S. & McDonnell, M.J.** 1988. Nitrogen cycling processes and soil characteristics in an urban versus rural forest. *Biogeochemistry*, 5: 243–262. <https://doi.org/10.1007/BF02180230>
- Woldegerima, T., Yeshitela, K. & Lindley, S.** 2017. Ecosystem services assessment of the urban forests of Addis Ababa, Ethiopia. *Urban Ecosyst*, 20: 683–699. <https://doi.org/10.1007/s11252-016-0624-3>
- Yoon, T.K., Seo, K., Park, G., Son, Y.M. & Son, Y.** 2016. Surface Soil Carbon Storage in Urban Green Spaces in Three Major South Korean Cities. *Forests*, 7: 115. <https://doi.org/10.3390/f7060115>.












The Global Soil Partnership (GSP) is a globally recognized mechanism established in 2012. Our mission is to position soils in the Global Agenda through collective action. Our key objectives are to promote Sustainable Soil Management (SSM) and improve soil governance to guarantee healthy and productive soils, and support the provision of essential ecosystem services towards food security and improved nutrition, climate change adaptation and mitigation, and sustainable development.



Thanks to the financial support of



European  
Commission



Ministry of Finance of the  
Russian Federation

ISBN 978-92-5-134900-7



9 789251 349007

CB6606EN/1/09.21