

the corolla, (b) its excision, (c) loss of colour, (d) closing, (e) not opening, (f) absence of insects, (g) reduction of temperature, (h) transportation. 5. Highly self-fertile forms may arise under cultivation. 6. Special adaptations occur for self-fertilisation. 7. Inconspicuous flowers are highly self-fertile. 8. Cyclicotamous flowers are always self-fertilised. 9. Conservation of energy in reduction of pollen. 10. Relative fertility may equal or surpass that of crossed plants. 11. It does not decrease in successive generations. 12. It may increase. 13. Free from competition self-fertilised plants equal the intercrossed; (a) as seedlings, (b) planted in open ground. 14. They may gain no benefit from a cross from the same or a different stock. 15. They are as healthy as the intercrossed. 16. They may be much more productive than flowers dependent upon insects. 17. Naturalised abroad they gain great vigour; and (18) are the fittest to survive in the struggle for life.

**PHYSIOLOGICAL ACTION OF NICOTINE.**—About twenty years ago the Rev. Prof. Haughton called attention to the fact that there was an antagonism between the actions of nicotine and strychnia. His experiments were on frogs. About ten years afterwards Dr. Wernley experimented in the same direction with cats; and some five years ago Dr. Reese performed a series of experiments with these drugs on dogs. Not satisfied with the results of any of these experiments and recognising the great importance of the subject, Dr. Haynes has made a long series of experiments on dogs, cats, rabbits, and rats, and after some 143 experiments, has come to the following conclusions:—"The recorded cases of strychnia poisoning treated by tobacco are extremely unsatisfactory. If they prove anything it is merely that tobacco is a powerful emetic." "Haughton's experiments on this subject (really only two in number) were performed in such an unsatisfactory manner as to be utterly valueless." "Strychnia and nicotine are in no degree antagonistic poisons." "Strychnia increases the convulsive action and does not diminish the motor paralysis of nicotine." "Nicotine (even in paralyzing doses) increases the convulsive action of strychnia." "Both poisons cause death by paralyzing the respiratory organs. They may affect respiration in different ways, but the result is the same." "Animals may be killed by injecting together doses of the two drugs which, singly, are not fatal [*Proceedings of the American Philosophical Society, vol. xvi., No. 99.*]

**GLASSY SPONGES.**—Dr. W. Marshall and A. H. Meyer have published a memoir, as one of a series of contributions to the Zoological Museum at Dresden, "on some new or little-known sponges belonging to the Hexactinellida found in the Philippines." It seems but the other day since one could have numbered on the fingers of one hand all the known species of this family, so well known to many by that beautiful typical form, the Venus flower-basket (*Euplectella*), and now the number of described species is very large. In 1872 one of the authors (Dr. Meyer) was staying at Cebu one of the Philippine group, where *Euplectella aspergillum* is a regular article of trade, quoted at so much a dozen, and where it is not surprising that he should discover a number of other lovely forms in this memoir described and figured. Among the most interesting forms are the following:—*Hyalosarcus simplex*, *Mysium nitidum*, and two species of *Asiodictyon*, all of these found living attached to the basal portion of *Euplectella*. *Semperella schultzei* is figured of a natural size from a specimen twenty-one inches in length, and figures of the spicules of the various new species are also given.

**A MALE NURSE.**—The interest of the reproduction of Batrachians is by no means yet exhausted. A Spanish naturalist, Jimenez de la Espada, has recently discovered additional facts respecting *Rhombodroma dorsalis* (cf. ChIII), which was first made known by Mr. Darwin,

He finds that the supposed viviparous birth of the young from the female is a very different phenomenon. It is the males which are the nurses, and they have an extraordinary brood-sac, developed as a pouch from the throat, and extending over a great portion of the vertical surface of the animal. In this cavity a number of living tadpoles were found, in number of individuals, and the length of the tadpoles was about 14 mm. How these are first developed and nourished is not yet known. Dr. J. W. Spengel illustrates a portion of the Spanish paper in the current number of the *Zeitschrift für wissenschaftliche Zoologie*, vol. xxix. part 4.

**STRUCTURE OF CYCADLES.**—E. Warming, of Copenhagen, publishes (in Danish with French abstract) some fresh researches on this subject ("Recherches et Remarques sur les Cycadées," Copenhagen, 1877). He confirms in general the results previously arrived at by A. Braun and others, from the structure of the ovule and seed, the pro-embryonic characters, the mode of formation of the pollen and pollen-pipe, and of the growth of the stem and roots, &c., that the Cycadées are very nearly allied to the Conifères; and in particular he places them near to the Ginkgo (*Salsolaria alba*). Among Cryptogams he considers them to come nearest to Marattiaceæ and Ophioglossaceæ among Filicinae. He proceeds then to discuss the homology of the ovule of Phanerogams and Angiosperms—he thinks the structure of that of the Cycadées—intermediate between Vascular Cryptogams and Angiosperms—throws much light. The phanerogamic ovule he considers to be composed of two parts, of different morphological origin, viz., a nucleus which is homologous with the macrosporangium; and a lobe of the leaf which bears the nucleus, consisting partly of the funiculus and partly of the integumentum. In Angiosperms the nucleus rests on the surface of the leaf; in Gymnosperms it is partly imbedded in it. No part of the ovule is of axial origin (*Journal*).

**THE BRAIN OF A FOSSIL MAMMAL.**—Prof. Cope has been able to take a cast of the cranial cavity of a species of the Tapirid genus *Coryphodon*, from the Wahsatch beds of New Mexico. This has revealed remarkable primitive characters: (1) the small size of the cerebellum; (2) the large size of the region of the corpora quadrigemina; (3) the cerebral hemispheres were small, and (4) the olfactory lobes were very large. The medulla oblongata is wider than the cerebral hemispheres. In profile the brain closely resembles that of a lizard. These characters are so extraordinary that Prof. Cope considers them sufficient to mark a primary division of mammalia, which he, following Owen, calls Protencephalia. Prof. Cope describes and gives figures of a cast, the skull cavity, in the *Proceedings of the American Philosophical Society*, vol. xvi., No. 99.

#### INSECTIVOROUS PLANTS! [See Over]

SINCE the appearance of Mr. Darwin's work on "Insectivorous Plants" the want of direct proof that the plants profit by their carnivorous habits has been somewhat widely felt. Thus we find expressions to this effect by MM. Cassinir de Cardelle, Cramer, Duchateau, Daxal-Jouve, Faivre, Göppert, E. Moerer, Munk, Naudin, W. Pfeffer, Schenk, &c., &c.

The assent which many naturalists have given to Mr. Darwin's explanation of the meaning of the structure and physiological properties of carnivorous plants rests on a sound basis, namely, the impossibility of believing that highly specialised organs are unimportant to their possessor, and the difficulty of giving any rational explanation except the one proposed in "Insectivorous Plants." Mr. Darwin himself felt the desirableness of direct evidence on this head, and the experiments intended to

<sup>1</sup> From a paper "On the Nutrition of *Drosera rotundifolia*," by Francis Darwin, M.D., read before the Linnean Society, January 17, 1861.

I must put an appendix -

decide the question only settled through an accident. The present research by Dr. F. Darwin is practically a repetition of the same experiment.

The widely-spread belief that insectivorous plants thrive equally well when deprived of animal food rests on very insufficient grounds. Many observers have based their opinions on the general appearance of the plants, and in no case has observation been sufficiently extended in point of time or details of comparison. The plan of the present research was therefore (1) To cultivate a large number of plants. (2) To continue observation for a considerable space of time, during which artificial starving and feeding of two sets of plants was to be kept up. (3) To compare the starved and fed plants in a variety of ways and especially as to the production of seed.

With this object about six plants of *Drosera rotundifolia* were transplanted (June 19, 1875), and cultivated in compartments filled with moss during the rest of the summer.

Each plant was divided into halves by a low wooden partition, one side being destined to be fed with meat, while the plants in the opposite half were to be left starved. The plants were placed altogether under a glass case, so that the "starved" plants might be prevented from obtaining food by the capture of insects. The method of feeding consisted in supplying each half (on the fed side) of the six plants with one or two small bits of oxen meat, each weighing about one-fifth of a grain. This operation was repeated every two days from the beginning of July to the first days of September, when the final comparison of the two sets of plants was made, but long before this it was quite clear that the "fed" plants were growing by their roots &c. Thus, on July 27 it was evident that the leaves on the "fed" side were of a distinctly brighter green, showing that the increased supply of nitrogen had allowed a more active formation of chlorophyll grains to take place. It may be inferred, partly from microscopical examination of the starch in the leaves, but more certainly from the final comparison of dry weights, that the increase of chlorophyll was accompanied by an increased formation of cellulose. From this time forward the "fed" sides of the plants were clearly distinguishable by their shining appearance and their somewhat tall and stout flower-stems.

The advantage gained by the fed plants was estimated in many ways. First, on August 7 the ratio between the number of "starved" and "fed" flower stalks was 100 : 109.1. And by comparing the number of seeds actually in flower it was clear that the starved plants were losing the power of throwing up new flower stems at an earlier date than their rivals. In the middle of August the leaves were counted in three plants, and were found to be 117 on the starved, and 156 on the fed side—in the ratio of 100 : 133.5.

At the beginning of September the seeds being ripe, all the flower-stems were gathered, and the plants of these plants were picked out of the moss and carefully washed. As it seemed probable that one advantage of the fed over the starved plants would be the power of laying by a larger store of reserve-material, three plants were allowed to remain undisturbed after the flower-stems had been gathered. The relative number of plants which will appear in the spring on the "fed" and "starved" sides will be a means of estimating the relative quantities of reserve-material.

The following table gives the result of counting, measuring, and weighing the various parts of the two sets of plants. It will be seen the number of plants (judging from the three plants mentioned) were fairly equal on the starved and fed sides of the partitions so that a direct comparison of their produce is allowable—

Ratio between the number of starved and fed plants	100 : 101.1*
--	--------------

Ratio between weights of the plants on either of three dates	100 : 100.7
Total number of flower stems	100 : 104.9
Sum of the heights of the flower stems	100 : 103.4*
Total weight of flower stems	100 : 107.9
Total number of capsules	100 : 108.4
Average number of seeds per capsule	100 : 100.7
Average weight per seed	100 : 117.1
Total calculated number of seeds produced	100 : 111.7
Total calculated weight of seeds produced	100 : 133.9

The most important feature in the general result is that the advantage gained by the fed plants is far more conspicuously shown in all that relates to the seeds and flower-stems than in any other part. Thus the ratio between the weights of the plants, exclusive of flower-stems, were as 100 to 111.7; while the weights of the flower-stems, including seeds and capsules, were as 100 to 107.9. The highest ratio in seeds is between the total weight of seed produced, namely 100 : 133.9; and this is intelligible, because a store of nitrogen is laid by in the albuminous seeds.

Another point is that the difference between the starved and fed plants is more clearly shown in the comparison of weights than of numbers or heights. It is clear that increase of weight is a better proof of increased assimilation than any other character.

It may fairly be said that the above experiments prove beyond a doubt that insectivorous plants are largely benefited by a supply of animal food, and it can no longer be doubted that a similar benefit is gained in a more or less degree by the capture of insects.

ALBERT FOW HALLER

ON December 21 last the republic and city of Geneva celebrated the centenary of the death of one who is universally recognized as their greatest citizen. The important part played in science by Albert von Haller has already in a sufficient degree for us, pointing by the example of the recent celebration, to enable our readers to appreciate the marvellous aptitude of this eminent man for every kind of work, theoretical and practical, be it in any a department, theological, and pure, as well as a physiological, anatomical, and botanic.

Albert Haller was born at Bormio in October, 1707, of a family originally of St. Gall, one of whose members fell by the side of Zwargli in 1522. Very weak in body, like Isaac Newton, in his infancy he exhibited, like him, an extraordinary precocity, and his studies for books was something insatiable. Having finished his classical studies brilliantly and rapidly, he went to Tübingen at the age of fifteen years to study medicine, then came also to Leyden to follow the clinic of the illustrious Boerhaave, on whose works he at a later date published a commentary which greatly commended it to his country. Haller taught law anatomy and I Geneva history. At eighteen and a half years he obtained the degree of doctor, and afterwards attended, in London, the teaching of Dr. Winslow. After a sojourn at Paris he returned to Switzerland and studied mathematics with Jean Bernoulli, and that with such ardour that his friends were constrained to look after him.

In 1738 he made, with Gessner, his first great Alpine excursion, which, many times repeated, made him, in an eminent degree, master of the Swiss Alps. His most celebrated poem, entitled "Die Alpen," was another result of his mountain journey, which contributed to diffuse among those far away the magic charms of that magnificent scenery.

\* Showing the average height of the fed stems is slightly less than 100 of that of all the fed, but the average number of plants and seeds, the total weight of flower stems, and the total weight of seeds.

\*\* These statistical observations of the Fuchs's *Utricularia*, kindly at a very private lecture accompanied with the paper, will be found in a subsequent issue of this journal. It is further to be mentioned, that it was the intention of the author to publish this paper in the *Journal de Pharmacie*.