

Chapter 18. Detection and Attribution of Observed Impacts

Coordinating Lead Authors

Wolfgang Cramer (Germany), Gary Yohe (USA)

Lead Authors

Maximilian Auffhammer (USA), Christian Huggel (Switzerland), Ulf Molau (Sweden), Maria Assunção Faus da Silva Dias (Brazil), Andrew Solow (USA), Dáithí Stone (Canada / South Africa), Lourdes Tibig (Philippines)

Contributing Authors

Laurens Bouwer (Netherlands), Mark Carey (USA), Dim Coumou (Germany), Gerrit Hansen (Germany), Ove Hoegh-Guldberg (Australia), Solomon Hsiang (USA), Sari Kovats (UK), Paul Leadley (France), David Lobell (USA), Camille Parmesan (USA), Elvira Poloczanska (Australia), Hans Otto Pörtner (Germany)

Review Editors

Rik Leemans (Netherlands), Bernard Seguin (France), Neville Smith (Australia)

Volunteer Chapter Scientist

Gerrit Hansen (Germany)

Contents

Executive Summary

18.1. Introduction

18.1.1. Scope and Goals of the Chapter

18.1.2. Summary of Findings from the AR4

18.2. Methodological Concepts for Detection and Attribution of Impacts of Climate Change

18.2.1. Concepts and Approaches

18.2.1.1. Detecting and Attributing Change in the Earth System

18.2.1.2. Concepts of Detection and Attribution of Climate Change Impacts

18.2.1.3. Approaches to Attribution

18.2.2. Challenges to Detection and Attribution

18.2.2.1. Types and Quality of Observations

18.2.2.2. Spatial and Temporal Characteristics of Change

18.2.2.3. Publication Bias

18.3. Detection and Attribution of Observed Climate Change Impacts in Natural Systems

18.3.1. Freshwater Resources

18.3.1.1. The Regional Water Balance

18.3.1.2. Floods and Droughts

18.3.1.3. The Cryosphere

18.3.1.4. Erosion, Landslides, and Avalanches

18.3.2. Terrestrial and Inland Water Systems

18.3.2.1. Phenology

18.3.2.2. Productivity and Biomass

18.3.2.3. Biodiversity

18.3.2.4. Impacts on Major Systems

18.3.3. Coastal Systems and Low Lying Areas

18.3.3.1. Erosion, Shoreline Processes, and Coastal Aquifers

18.3.3.2. Coastal Ecosystems

18.3.3.3. Coastal Settlements, Infrastructure, and Economic Activities

- 1 18.3.4. Oceans
2 18.3.4.1. Impacts on Ocean System Properties and Marine Organisms and Ecosystems
3 18.3.4.2. Observed Climate Change Effects across Ocean Regions
4
- 5 18.4. Detection and Attribution of Observed Climate Change Impacts in Human and Managed Systems
6 18.4.1. Food Production Systems
7 18.4.1.1. Agricultural Crops
8 18.4.1.2. Fisheries
9 18.4.1.3. Food Security
10 18.4.2. Cities and Urbanization
11 18.4.3. Economic Impacts, Key Economic Sectors and Services
12 18.4.3.1. Economic Growth
13 18.4.3.2. Energy Systems
14 18.4.3.3. Tourism
15 18.4.4. Impacts of Extreme Weather Events
16 18.4.4.1. Economic Losses due to Extreme Weather Events
17 18.4.4.2. Detection and Attribution of the Impacts of Single Extreme Weather Events to Climate
18 Change
19 18.4.5. Human Health
20 18.4.6. Human Security
21 18.4.6.1. Violent Conflicts and Social Disruptions
22 18.4.6.2. Migration
23 18.4.7. Rural Areas, Livelihoods, and Poverty
24
- 25 18.5. Detection and Attribution of Observed Climate Change Impacts across Regions
26 18.5.1. Africa
27 18.5.2. Europe
28 18.5.3. Asia
29 18.5.4. Australasia
30 18.5.5. North America
31 18.5.6. Central and South America
32 18.5.7. Polar Regions
33 18.5.8. Small Islands
34 18.5.9. Impacts across Regions
35
- 36 18.6. Synthesis
37 18.6.1. Approach and History
38 18.6.2. Five Reasons for Concern
39 18.6.2.1. Risks to Unique and Threatened Systems
40 18.6.2.2. Risk of Extreme Weather Events
41 18.6.2.3. Distribution of Impacts
42 18.6.2.4. Aggregate Impacts
43 18.6.2.5. Risks of Large Scale Discontinuities
44 18.6.3. Conclusion
45
- 46 18.7. Gaps, Research Needs, Emerging Issues
47
- 48 Frequently Asked Questions
49 18.1: What are the main challenges in detecting climate change impacts?
50 18.2: What is the main challenge in attributing changes in a system to climate change?
51 18.3: Why are detection and attribution of climate impacts important?
52 18.4: Is it possible to attribute a single event, like a disease outbreak or the extinction of a species, to
53 climate change?
54

1 References

2
3
4 **Executive Summary**

5
6 **Impacts of recent observed climate change on physical, biological, and human systems have been detected on**
7 **all continents and in most oceans. This conclusion is strengthened by observations since the AR4 as well as**
8 **through more extensive analyses of earlier observations.** Most reported impacts of climate change are attributed
9 to regional warming of the atmosphere and the ocean. The level of confidence in attribution of observed impacts to
10 shifts in rainfall patterns is lower. There is emerging evidence of impacts of ocean acidification. [18.3-18.6]

11
12 **For many natural systems, new or stronger evidence for substantial and wide-ranging impacts of climate**
13 **change exists, including the cryosphere, water resources, coastal systems and ecosystems on land and in the**
14 **ocean.**

15
16 **Cryosphere and water resources**

17
18 **Glaciers worldwide continue to shrink;** new glacier lakes have formed and existing ones have changed;
19 seasonal ice in many lakes and rivers forms later and breaks up earlier. A major part of these changes can
20 be attributed with *high confidence* to climate change. [18.3.1.3, 18.5; Figure 18-3]

21
22 **Widespread changes and degradation of permafrost of both high-latitude and high-elevation**
23 **mountain regions have been observed over the past years and decades (*high confidence*).** The
24 permafrost boundary has been moving polewards and to higher elevations, and the active layer thickness
25 has increased at many sites (*medium confidence* in attribution to climate change). [18.3.1.3, 18.5]

26
27 **Hydrological systems have changed in many regions,** due to changing rainfall or melting glaciers,
28 affecting water resources, water quality, and sediment transport (*medium confidence*). In many river
29 systems, the frequency of floods has been altered by climate change (*low to medium confidence*). The
30 duration of droughts in some regions has been altered by climate change (*medium confidence*). [18.3.1.1,
31 18.5]

32
33 **Terrestrial and freshwater ecosystems**

34
35 **Across all climate zones and continents, an increasing range of species and communities in terrestrial**
36 **ecosystems have been impacted by recent climate change and increasing atmospheric CO₂.** Many
37 plants and animals show changes in phenology (*high confidence*), productivity and / or geographic range
38 (*medium confidence*). Elevated rates of extinction cannot be attributed to climate change. [18.3.2, 18.5]

39
40 **In freshwater ecosystems of most continents and climate zones, rising temperatures have been linked**
41 **to shifts in animal community composition, especially in headwater streams.** While confidence in
42 detection of change is *high*, there is *low confidence* in attribution to climate change due to many
43 confounding factors. [18.3.2.4, 18.5]

44
45 **Several major terrestrial ecosystems are undergoing broad-scale changes that can be characterized**
46 **as early warnings for coming regime shifts, in part due to climate change.** For wide-spread shrub
47 encroachment in the Arctic tundra there is *high confidence*, for boreal forest tree mortality there is *low*
48 *confidence* of climate change as driver. The recession and degradation of the Amazon forest cannot be
49 attributed to climate change. [18.3.2.4, 18.5.6, 18.5.7]

50
51 **Oceans and coastal systems**

52
53 **The physical and chemical properties of the ocean have changed significantly over the past 60 years,**
54 **due to anthropogenic climate change. As a result, marine organisms have moved to higher latitudes,**

1 **changed their depth distribution or their phenology (*high confidence*).** Facilitated by changes in the
2 distribution of sea ice and changes in ocean currents, organisms such as crustaceans, zooplankton and fish
3 have migrated toward higher latitudes. These changes also affect coastal ecosystems including kelp forests
4 and seagrass meadows [18.3.3, 18.3.4.1]
5

6 **Climate change has influenced ocean primary productivity, with positive consequences for some**
7 **fisheries and negative ones for others (*medium confidence*).** Together with non-climate influences such
8 as excess nutrient input from human activities, climate change has contributed to an increase in the
9 frequency, geographical distribution, and severity of hypoxic areas in the ocean. [18.3.4.1, 18.3.4.2]
10

11 **Due to the increased frequency of stress events arising from elevated sea temperatures, coral reefs**
12 **have experienced increased mass bleaching and mortality (*very high confidence*).** These events have
13 contributed to the loss of reef building corals in many parts of the world since the early 1980s. [18.3.3,
14 18.3.4, Box 18-3, 18.5, Table 18-8, Box CC-CR]
15

16 **Arctic sea ice has been shrinking in extent, thickness, composition, with observed impacts on marine**
17 **biology and the livelihoods of indigenous people (*medium to high confidence*).** [18.3.1.3, 18.3.4, 18.4.7,
18 18.5.7]
19

20 **Despite the known sensitivity of coastal systems to sea-level rise, local perturbations from regional**
21 **variability in the ocean and human activities preclude the confident detection of sea level-related**
22 **impacts attributable to climate change outside of the Arctic.** [18.3.3]
23

24 **For managed ecosystems and human systems, the effects of changing social and economic factors often**
25 **dominate over any direct impact of climate change. Despite this, numerous impacts of climate change have**
26 **been detected.**
27

28 **Even accounting for changes in technology and other non-climate factors, agricultural crop yields**
29 **have changed in many regions in response to climate.** Yields have increased in some (mid to high
30 latitude) regions, due to warming and higher CO₂ (*low confidence*), and decreased in other (mainly low
31 latitude) regions due to water shortages and higher temperatures (*medium confidence*). Despite the high
32 sensitivity of crop yields to temperature extremes, observed trends in the agricultural markets cannot
33 presently be attributed to climate change, due presence of other drivers. [18.4.1, Table 18-9]
34

35 **The catch potential of fisheries has increased in some regions and decreased in others with**
36 **consequences for the food and livelihood of involved human communities (*high confidence*).** Fisheries
37 at high latitudes are showing increased productivity due to sea ice retreats and increases in net primary
38 productivity. In other regions, stratification of the water column driven by warming has reduced net
39 primary productivity of the ocean. [18.3.4, 18.4.1.2, 18.5.7]
40

41 **In some regions, increases in the prevalence of vector-borne diseases have been detected and**
42 **attributed to warming.** Dengue fever and malaria have increased in several regions of the world over the
43 past few decades, but there is *very low confidence* in attribution of these trends to climate change. [18.4.5]
44

45 **Climate impacts on Arctic indigenous groups have been detected and attributed to climate change.**
46 These include changes in seasonal migration and hunting patterns, health, and cultural identity (*medium*
47 *confidence*). [18.4.7, Box 18-5, 18.5.7, Table 18-9]
48

49 **Extreme climate events have impacted natural and physical livelihood assets, incomes, public health,**
50 **and social institutions.** Economic losses due to extreme weather events have increased globally, mostly
51 due to increase in wealth and exposure, but with a documented contribution of climate change and
52 variability in some cases. [18.4.4, 18.4.7]
53

1 **Despite known vulnerabilities and increasing exposure to climatic stressors, impacts of climate**
2 **change on human livelihoods have rarely been detected with confidence.** Such detection is complicated
3 by the effects of other economic and social factors. There is emerging literature on the impact of climate on
4 poverty, working conditions, violent conflict, migration, and economic growth, but evidence for detection
5 or attribution remains limited. [18.4.3, 18.4.6, 18.4.7]
6

7 **Detection and attribution of observed impacts of climate change supports assessments of current conditions**
8 **with respect to “Reasons for Concern” formulated by earlier IPCC assessments.** The degree to which projected
9 damages are now manifest, or the detection of stronger early warning signals for expected impacts, can contribute to
10 a more comprehensive risk assessment for dangerous anthropogenic interference with the climate system.
11

12 **Increases in “risks to unique and threatened systems” are now documented with higher confidence**
13 **by the observed impacts on Arctic marine and terrestrial ecosystems and indigenous livelihoods**
14 **(medium to high confidence), tropical coral reefs (high confidence) and glaciers in most mountain**
15 **regions (high confidence). Observed impacts thus confirm this reason for concern.** [18.6.2.1]
16

17 **“Risks of extreme weather events” – extreme warming events continue to increase for warm water**
18 **coral reefs (high confidence), confirming this reason for climate-related concern. Risks have also**
19 **increased in some other systems, causing economic losses, however this is predominantly due to the**
20 **increases in wealth and exposure.** [18.6.2.2]
21

22 **Impacts of climate change have now been documented globally, covering all continents and the ocean**
23 **(high confidence). However, research coverage is still insufficient and too heterogeneous to effectively**
24 **address spatial or social disparities concerning the “distribution of impacts” beyond local case**
25 **studies.** [18.6.2.3]
26

27 **“Aggregate impacts” have been documented with measures of glacier volume and permafrost extent**
28 **(medium to high confidence). There is limited evidence of a climate contribution to aggregate impacts**
29 **on biospheric carbon fluxes, agricultural yields, fishery production and economic losses due to**
30 **extreme weather, due to the presence of dominant non-climatic drivers. For the cryospheric impacts,**
31 **observations clearly confirm the reason for concern.** [18.6.2.4]
32

33 **“Risks from large-scale singularities” relating to irreversible shifts with significant feedback**
34 **potential in the Earth system have yet to be observed. However, there is now robust evidence of**
35 **significant early warning signals in observed impacts of climate change that indicate large-scale**
36 **regime shifts for the Arctic region and the tropical coral reef systems. This evidence confirms reasons**
37 **for concern.** [18.6.2.5]
38

39 **Though evidence is improving, there is a persistent gap of knowledge regarding how large parts of the world**
40 **are being affected by observed climate change.** The number of attribution studies linking observed impacts to the
41 anthropogenic component of climate change remains limited. Research to improve the timeliness and knowledge
42 about the detection and attribution is needed in particular for the risk of extreme events. [18.7]
43

44 **In most studies, the attribution of observed impacts and vulnerabilities is related to all changes in climate that**
45 **represent deviations from historical means and / or historic variability. Only a smaller number of robust**
46 **attribution studies link responses in physical and biological systems to anthropogenic climate change.** [18.1]
47

48 **Methods for rigorous assessment of observed impacts of climate change are evolving.** Evidence for attribution
49 comes from assessment of the relative contribution by all known drivers affecting the dynamics of a system to its
50 behavior, using scientific methods, involving an assessment of confidence in both detection and attribution. Formal
51 meta-analysis or aggregated assessments of many observations or studies can help to improve confidence. [18.2.1,
52 Box 18-1]
53
54

18.1. Introduction

This chapter synthesizes the scientific literature on the detection and attribution of observed changes in physical, biological and human systems in response to the climate change that has occurred during the recent few decades. It assesses the degree to which detected changes in such systems can be attributed to observed climate change and where possible, separates out the relative importance of anthropogenic drivers of climate change. For most natural and essentially all human systems, climate is only one of many drivers that cause systems to change, which therefore requires a careful accounting of the importance of other confounding factors on the overall change in these systems.

18.1.1. Scope and Goals of the Chapter

Previous assessments, notably Rosenzweig *et al.* (2007), in the IPCC Fourth Assessment Report (AR4), and the increasing body of literature published since, indicated that numerous physical and biological systems are affected by recent climate change. Human systems have received comparatively little attention in this literature, with the exception of the food system, which is a coupled human/natural system. Rigorous formal assessment of the literature going beyond just detection of change across a variety of regions and sectors, to scientifically robust attribution to climate change and its anthropogenic drivers is critical for several purposes (Brander *et al.*, 2011; Hoegh-Guldberg, 2011; Stocker *et al.*, 2011). Only formal detection studies provide robust evidence of where climate change impacts already being observed and where they are not, supporting near-term planned adaptation if and where necessary.

For policy makers and the public, detection and attribution of observed impacts will be a key element to determine the necessity and degree of mitigation and adaptation efforts. Detection and attribution are vital parts of the evidence base requested of the IPCC by signatories to the United Nations Framework Convention on Climate change to judge policies aiming to stabilize “greenhouse gas concentrations in the atmosphere at a level that would prevent dangerous atmospheric interference with the climate system”.

Full attribution of changes in these systems to anthropogenic climate is extremely hard to accomplish for many complex systems with diverse responses and multiple significant confounding drivers (Parmesan *et al.*, 2011). There has, however, been a significant increase in studies attributing changes in some systems to local climate change without direct linkage to anthropogenic drivers. This chapter assesses the studies that exist for both full and partial attribution, and the methodologies that can be brought to bear on attribution and the uncertainties inherent in doing so.

18.1.2. Summary of Findings from the AR4

Rosenzweig *et al.* (2007) reported that “observational evidence from all continents and most oceans shows that many natural systems are being affected by regional climate changes, particularly temperature increases.” In particular, they highlighted several areas where this general conclusion could be supported by specific conclusions that were reported with *high confidence*:

- Changes in snow, ice and frozen ground had been seen to increase ground instability in mountain and other permafrost regions; these changes had led to changes in some Arctic and Antarctic ecosystems and produced increases in the number and size of glacial lakes.
- Some hydrological systems had been affected by increased runoff and earlier spring peak discharges; in particular many glacier- and snow-fed rivers and lakes had warmed, producing changes in their thermal structures and water quality.
- Spring events had appeared earlier in the year so that some terrestrial ecosystems had moved poleward and upward; these shifts in plant and animal ranges were attributed to recent warming.
- Shifts in ranges and changes in algal, plankton and fish abundance as well as changes in ice cover, salinity, oxygen levels and circulation had been associated with rising water temperatures in some marine and freshwater systems.

1 In terms of a global synthesis, this assessment noted “that it is likely that anthropogenic warming over the last three
2 decades has had a discernible influence on many physical and biological systems” (Rosenzweig *et al.*, 2007).
3 Though it was based on analyses of a very large number of observational datasets, the assessment noted a lack of
4 geographic balance in data and literature on observed changes, with marked scarcity in low and middle income
5 countries.

6
7 Observed impacts to human systems were less obviously attributed to anthropogenic climate change. Rosenzweig *et al.*
8 *al.* (2007) concluded with *medium confidence* only that, “other effects of regional climate change on natural and
9 human environments are emerging, although many are difficult to discern due to adaptation and non-climatic
10 drivers”. They especially noted effects of temperature increases on:

- 11 • Some agricultural and forestry management practices in the higher latitudes of the Northern Hemisphere;
12 these included earlier spring planting as well as changes in the disturbance regimes of fires and pests.
- 13 • Some aspects of human health, including heat-related mortality in Europe, changes in some vectors of
14 infectious diseases across the world, and phenological changes in allergenic pollen in the mid to high
15 latitudes of the Northern Hemisphere.
- 16 • Some human activities in the Arctic (such as hunting and travel) and in lower-elevation alpine areas (such
17 as mountain sports).

18.2. Methodological Concepts for Detection and Attribution of Impacts of Climate Change

21
22 There are three substantial challenges to the detection and attribution of change in environmental systems to a
23 changing climate. First, all systems are affected also by environmental factors other than climate; to detect a climate
24 change impact, it is therefore necessary to control for the effects of such confounding factors. Second, the ability of
25 many systems to adapt to change complicates the detection and attribution of any impacts; that is, adaptation can
26 mask an impact. Detection (and attribution) of adaptation to climate change is in this chapter considered to be
27 detection (and attribution) of a climate change impact (Box 18-1). Third, because systems are typically affected by
28 local or regional climate change, attribution of a detected climate change impact to anthropogenic climate change
29 can be difficult. To overcome these difficulties and to best account for all available knowledge on observed impacts,
30 a range of methods is employed here. They are summarized in this section.

18.2.1 Concepts and Approaches

18.2.1.1. Detecting and Attributing Change in the Earth System

36
37 From an analysis perspective, the Earth system can be separated into three connected subsystems; we name them
38 here as the climate system, the natural system, and the human system (Figure 18-1). Many external drivers may
39 influence any system, including the changing climate and other confounding factors (Hegerl *et al.*, 2010). Each
40 climate, natural or human subsystem affects the other two directly or indirectly. For example, the human system
41 may directly affect the natural system through deforestation, which in turn affects the climate system through
42 changes in albedo; this can alter surface temperatures which in turn feed back on natural and human systems.

43
44 [INSERT FIGURE 18-1 HERE

45 Figure 18-1: Schematic of the subject covered in this chapter. The Earth system can be divided into three broad
46 interacting systems. Direct drivers of the human system on the climate system are denoted with a red arrow; some of
47 these drivers may also directly affect natural systems. These effects can in turn influence other systems (dashed red
48 arrows). Further influences of each of the systems on each other (confounding factors) that do not involve climate
49 drivers are represented by blue arrows. Examples of drivers and their impacts are given in the table. Adapted from
50 Stone *et al.* (2013).]

51
52 If an observed change in the human system impacts the climate system, we call this an anthropogenic driver of
53 climate change. In order to highlight potential reasons for concern (including the uncertainties in detection and

1 attribution), impacts of any observed climate change are covered in this chapter, irrespective of whether the
2 particular aspect of climate change has been identified as anthropogenic.

3 4 5 *18.2.1.2. Concepts of Detection and Attribution of Climate Change Impacts*

6
7 Concepts and approaches to detection and attribution have been evolving throughout the work of the IPCC. In 2010,
8 an IPCC Expert Meeting on Detection and Attribution was held to reconcile methods and terminology across IPCC
9 working groups. The report from this meeting (Hegerl *et al.*, 2010), along with the discussion of Stone *et al.* (2013),
10 is an important conceptual basis for this chapter.

11
12 ***Detection*** addresses the question of whether a system is changing beyond what might be considered normal
13 behaviour in the absence of climate change. While Hegerl *et al.* (2010) proposed a broad definition that could be
14 used across IPCC Working Groups, this definition, used in this chapter and in other Working Group II chapters, is
15 more specific in not just considering any observed changes (Stone *et al.*, 2013). The appropriate reference normal
16 behaviour may be stationary (e.g. glaciers) or non-stationary (e.g. economic activity), and the nature of that
17 reference needs to be spelled out clearly. Typically, detection studies involve the initial assumption of climate being
18 at least one of the drivers of that change, but other drivers (such as changes in land use) may be recognized to play a
19 significant role too. For many systems, the role of these confounding factors may exceed that of climate change.

20
21 ***Attribution*** addresses the question of whether climate change has contributed substantially to the detected change in
22 a system. In practice, attribution studies ask how much of the observed change is due to climate change. Attribution
23 requires the evaluation of the contributions of all external drivers to the system change.

24
25 Failure to include relevant factors in an attribution study can lead to erroneous conclusions. This is referred to as the
26 omitted variable problem (Greene, 2003). It is a particular problem in time series regressions when both climate and
27 non-climate factors may vary roughly monotonically over time. It is also important not to equate empirical
28 correlation with causality (Holland, 1986). This underscores the crucial role that system understanding must play in
29 high-quality detection and attribution studies.

30
31 Detection and attribution studies rely on quantitative or qualitative analysis. Quantitative studies adhere to basic
32 principles of statistical inference. One is that effects of formal or informal variable selection – in which a range of
33 statistical models is screened to identify the best-fitting model – must be taken into account to avoid spurious
34 attribution (Chatfield, 1995).

35
36 Qualitative attribution studies typically involve the identification of multiple, intersecting “impact chains” that
37 recognize confounding variables acting alongside climate change. These studies usually do not specifically isolate
38 certain social or environmental changes to climate change alone but rather, they aim to identify dynamic interactions
39 among an assemblage of intersecting forces in human-environment systems.

40
41 Evidence for testing the hypothesis that some observed change can be attributed to climate change may come from
42 different sources. The nature of the tools used for testing hypotheses varies depending on the details of the study
43 (Stone *et al.*, 2013). Mechanistic models can be used, for example representing a system through a chain of explicit
44 functions based on understanding of the individual processes that together comprise the mechanics of the system.
45 These mechanistic models can sometimes be formulated numerically. In other cases, empirical models may be
46 useful, relating the response of a system to external drivers according to mathematical relationships estimated
47 through observation, experimentation, or survey. In practice, though, many modeling setups will contain both
48 mechanistic and empirical components.

1 ____ START BOX 18-1 HERE ____

2
3 **Box 18-1. The Role of Climate Sensitivity and Adaptation for Impact Models in Human Systems**

4
5 Impacts of climate change on a measurable attribute of a human system only occur if a) the attribute is sensitive to
6 climate and b) a change in climate has occurred. A large literature has developed attempting to quantify both climate
7 sensitivity of various systems and observed changes in climate.

8
9 The literature estimating the climate sensitivity of an outcome such as crop yields, heat related mortality or
10 migration, has relied on observed climate variability either across space (e.g., Schlenker *et al.*, 2005), time (e.g.,
11 Mann and Emanuel, 2012), or space and time (Hsiang *et al.*, 2011; e.g., Dell *et al.*, 2012). While there is a rich
12 literature using climate variability across space, the long observational weather time series required for exploring
13 climate variability across space and time have limited the opportunities for study. A number of studies have instead
14 estimated the weather sensitivity of outcomes in order to project the future impacts of weather under climate change
15 (Deschênes and Greenstone, 2007; Deschênes and Greenstone, 2011), or attribute impacts for the past (Auffhammer
16 *et al.*, 2006). The issue with impact studies using a weather based sensitivity measure is that they cannot provide
17 estimates of impacts based on the climate sensitivity. For example, farmers may respond to an unusually hot
18 summer, which is a weather event, by applying more irrigation water. However, in the long run farmers may
19 respond to a warmer climate by switching crops, changing irrigation technology or abandoning farming altogether.
20 The two sensitivities and resulting magnitudes of attributable impacts due to a change in weather versus a change in
21 climate are therefore different.

22
23 In order to detect and attribute a change in a system to climate change, one needs to combine a measure of
24 sensitivity of the outcome to climate with an observed measure of climate under climate change. This can be done
25 via a multi-step approach (Hegerl *et al.*, 2010), whereby one estimates the outcome of interest using measures of
26 climate sensitivity combined with measures of observed climate and compares these to a simulated outcome using
27 the same sensitivity but removing the climate change from the climate input.

28
29 ____ END BOX 18-1 HERE ____

30
31
32 *18.2.1.3. Approaches to Attribution*

33
34 There are many levels at which drivers and responses can be defined for observed impacts of climate change (Table
35 18-1). In practice, two different levels of external drivers are used:

- 36 • Attribution to climate change, where impacts are related to identified long-term trends in climate (including
37 changes in variability patterns). Studies examining this relationship do not formally identify the role of
38 anthropogenic emissions in the observed climate change or, by extension, in the observed impact, but they
39 do indicate the degree to which the system is sensitive to long term climate change.
- 40 • Attribution to anthropogenic climate change, where impacts are related, via the climate, to anthropogenic
41 emissions of greenhouse gases and other human activities that are affecting the climate. Because of the
42 complexity of the causal chain, investigation of this relationship is extremely challenging (Parmesan *et al.*,
43 2011) and only a limited number of these studies have been performed to date.

44
45 Attribution studies use two main approaches (see Figure 18-2). Single-step methods consider all relevant systems in
46 a single setup. The assessment tool can comprise a single model or a sequence of models, provided the output of
47 each model is directly employed as an input for the next model in a logical sequence. The attribution analysis is
48 performed through a comparison of the change in the final output from the model(s) against the observed change in
49 the system of interest.

50
51 [INSERT FIGURE 18-2 HERE

52 Figure 18-2: A schematic diagram comparing approaches to attribution for an ecological system. The multi-step
53 approach differs from the single-step approach in having a discontinuity between the attributed climate change and
54 the observed weather driving the ecological model. Adapted from Stone *et al.* (2013).]

1
2 Multi-step attribution approaches comprise multiple attribution analyses, with an overall conclusion deduced from
3 the collection of analyses. A multi-step approach could include an analysis of the attribution of changes in a measure
4 of interest to changing climatic conditions and a separate analysis of how external climate drivers affected relevant
5 climatic conditions. Typically, the separation of these analyses creates gaps or inconsistencies between the outputs
6 from one step and the inputs to the next. These gaps must be assessed themselves with respect to their impact on the
7 confidence in the overall attribution.

8
9 In areas where extensive monitoring records are available, such as for phenological measures in ecological systems,
10 studies may be combined into a formal quantitative synthesis assessment. For instance, Rosenzweig *et al.* (2008)
11 noted that the spatial pattern of observed changes in a large number of natural systems around the globe matched the
12 pattern expected based on the observed warming pattern, and argued that other global factors such as land use
13 change could not account for this pattern (see also Box 18-2).

14 _____ START BOX 18-2 HERE _____

15 16 17 **Box 18-2. Quantitative Synthesis Assessment of Detection and Attribution Studies**

18
19 There are a number of powerful tools for quantitative synthesis assessment of detection and attribution across
20 multiple studies. These tools include associative pattern analyses (e.g., Rosenzweig *et al.*, 2008) and regression
21 analyses (Chen *et al.*, 2011) which compare expected changes due to anthropogenic climate change across multiple
22 studies against observed changes.

23
24 Quantitative synthesis assessments have been particularly prominent in ecology, where measures of phenology
25 (timing of seasonal events) and geographical range can be assembled across species into standardized indices
26 (Parmesan and Yohe, 2003; Rosenzweig *et al.*, 2008; Chen *et al.*, 2011). Synthesizing across multiple species can
27 increase our confidence in detection of general patterns of change in the biological indices. This confidence
28 increases with the number of species / ecosystems observed, the number of independent studies, the geographical
29 distribution of these observations, the temporal depth and resolution of the data, and the representativeness of
30 species/ecosystems and locations studied.

31
32 Synthesis assessments examining climate change as the only driver can provide some insight into the attribution of
33 biological responses to climate change because they can indicate whether plausible hypotheses are supported by
34 data. However, increasing spatial coverage, numbers of species, etc. does not a priori increase confidence that
35 climate change is a more credible explanation for biological change than alternative hypotheses. Additional data can
36 contribute to confidence in causal relationships, i.e. attribution, in a synthesis assessment when it provides new
37 evidence for explicit testing against a credible range of alternative hypotheses.

38
39 _____ END BOX 18-2 HERE _____

40 41 42 **18.2.2. Challenges to Detection and Attribution**

43 44 **18.2.2.1. Types and Quality of Observations**

45
46 The nature of indicators can take many forms, some of which are more amenable for precise long-term monitoring
47 than others. Many changing phenomena can be measured directly, e.g., the temperature of the surface waters of a
48 lake. Often, however, important indicators are less specific and thus difficult to pinpoint, for instance the excess
49 mortality and increased human vulnerability associated with heat waves, or the percentage of species migrating
50 polewards or upwards.

51
52 The quality of observations of these indicators varies in space and time. Monitoring of ecosystems, for example, is
53 often not designed with the intention of measuring incremental long-term changes and are not optimized for the
54 detection of climate-related changes (Midgley *et al.*, 2007). Consequently, the record length may be too short, too

1 heterogeneous or discontinuous to be useful, or measuring standards may have been improved through time
2 resulting in measurement artifacts. This latter factor is most visible in the extensive and long-running networks
3 designed to monitor human health, where the priority is an accurate timely assessment of current health status and
4 risk rather than the determination of long-term trends (see also 18.4.5).

7 *18.2.2.2. Spatial and Temporal Characteristics of Change*

9 Detection studies require observational data spanning a time period longer than the typical time scale of natural
10 variability in the system in the absence of external drivers. A particular challenge is that this required time scale is
11 usually difficult to discern from available observational records and thus other sources of information must be used
12 to determine them (Hegerl *et al.*, 2010). Lags between changes in external drivers and responses to those drivers
13 vary among different impact systems and within them, from seconds or minutes to centuries or millennia, and thus
14 need to be recognised (Stone *et al.*, 2013).

16 Non-linear system response to change is a fundamental issue for detection studies. Some systems may respond by
17 step-wise changes, other systems may have tipping points, showing little or no change until a certain threshold
18 where they suddenly start to respond vigorously and often in a chaotic way (e.g., De Young and Jarre, 2009;
19 Wassmann and Lenton, 2012, 18.6.2.5).

21 Change in environmental and human systems occurs on multiple spatial scales from local to global. Attribution of
22 change to climate change (e.g., over the last several decades) faces different challenges at different spatial scales. At
23 the local scale, detection may be straightforward due to the availability of long-term observations, but attribution
24 may be difficult because many local factors have been changing as well (e.g., Parmesan *et al.*, 2013). Furthermore,
25 local climate may be affected by localised drivers such as land cover change (De Noblet-Ducoudré *et al.*, 2012),
26 which complicates the evaluation of the role of global anthropogenic emissions.

29 *18.2.2.3. Publication Bias*

31 Conclusions about the effect of climate change on natural and human systems in this report are based on a synthesis
32 of findings in the scientific literature. A potential problem with this approach is publication bias, specifically the
33 preferential publication of papers reporting statistically significant findings (Parmesan and Yohe, 2003). The effect
34 of publication bias could be a false impression of the strength of the evidence in favor of a hypothesized effect.
35 Methods exist for detecting and correcting for publication bias in formal quantitative synthesis analysis (Rothstein *et al.*,
36 2005). For instance, the availability of a large sample of observations from an existing phenological monitoring
37 network has permitted the assessment that publication bias has not been an important factor in conclusions about the
38 role of climate change in studies of the timing of leafing, flowering, and fruiting in Europe (Menzel *et al.*, 2006), but
39 the lack of such long-term monitoring networks preclude such a conclusion for measures related to vector-borne
40 diseases (Kovats *et al.*, 2001).

43 **18.3. Detection and Attribution of Observed Climate Change Impacts in Natural Systems**

45 The IPCC AR4 provided extensive reporting on observed impacts of climate change (in Rosenzweig *et al.*, 2007, but
46 also other chapters of the WGII report). The scientific literature on this topic is growing quickly. Rather than a full
47 analysis of this literature, the following section provides a synthetic overview of the state of knowledge across major
48 sectors of natural systems, based largely on the respective sectoral chapters (3, 4, 5, 6) and chapter 30 of this report,
49 and a methodological framework with these chapters.

18.3.1 Freshwater Resources

The availability of freshwater resources, and essentially all aspects of the hydrological cycle, are affected by climate change on all continents and probably most islands, with different characteristics of change in different regions. Observed changes in each of the components of the water system are assessed by the IPCC WGI in their chapters 2 and 10, and WGII in their chapters 3, 4 and the regional chapters.

Figure 18-3 presents a synthesis of confidence in detection of changes in freshwater resources and related systems (notably erosion and slope stability), and their attribution to climate change. Frozen components of freshwater systems tend to show higher confidence in detection and attribution, while components that are strongly influenced by non-climatic drivers, such as groundwater or river flow, have lower confidence.

[INSERT FIGURE 18-3 HERE]

Figure 18-3: Levels of confidence in detection and attribution of observed climate change impacts for freshwater systems over the past several decades, based on expert assessment contained in this section 18.3.1 and augmented by subsections of chapter 3 as indicated. Numbered symbols refer to: Freshwater systems (18.3.1.1): 1 groundwater depletion (Ch.3.2.4), 2 changing river flow (Ch. 3.2.3), 3 changing flood frequency or intensity (Ch. 3.2.3), 4 reduction in lake and river ice duration or thickness (Ch. 18.3.1.2); Cryosphere: 5 shrinking glaciers (Ch. 3.2.2; 18.3.1.2), 6 changes in glacier lakes (Ch. 18.3.1.1), 7 erosion and degradation of arctic coastal permafrost (Ch. 18.3.1.2), 8 degradation and thaw of lowland and mountain permafrost (Ch. 18.3.1.2), Soils and rock (18.3.1.3): 9 increasing erosion (Ch.3.2.6), 10 changes in shallow landslides (Ch.3.2.6), 11 increasing frequency of alpine rock failures.]

18.3.1.1. The Regional Water Balance

The regional water balance is the net result of gains (precipitation, ice and snow melt, river and groundwater inflow) and losses (evapotranspiration, water use and river and groundwater outflow). Impacts of climate change include reduced availability of freshwater for use (one of the variables defining drought) or excess water (floods). Evapotranspiration, being a function of surface temperature, vegetation cover, soil moisture and wind, is affected by the changing climate, but also by changing vegetation processes and land cover.

At the global scale, human influence has contributed to large scale changes in precipitation patterns over land and, since the mid 20th century, in extreme precipitation (medium confidence, WGI AR5 Chapter10, Min *et al.*, 2011). More locations worldwide have experienced an increase than a decrease in heavy rainfall events, yet with significant regional and seasonal variations (Seneviratne *et al.*, 2012). While runoff has not changed in the majority of rivers, year-to-year variability has increased (WGI AR5 Chapter 2). At the regional scale, however, human influence has affected streamflow and also evapotranspiration (WGI AR5 Chapter 10).

Change in river flow is a direct indicator of a changing regional water balance. Globally, one-third of the top 200 rivers (ranked by river flow) show statistically significant trends during 1948–2004, with the rivers having downward trends (45) outnumbering those with upward trends (19) (Dai *et al.*, 2009). In the western United States, observed changes in the hydrological cycle (river flow, snow pack) during the second half of the 20th century have been linked to recent climate change, and 60% thereof were due to anthropogenic influences (Barnett *et al.*, 2008). Likewise, the Uruguay River in South America experienced a positive trend in average streamflow from 1960 to 2000, due to increased rainfall, with shorter-term peaks in runoff caused by changes in land cover (Saurral *et al.*, 2008). In the Yellow River in China, however, long-term change in streamflow are mostly attributed to anthropogenically enhanced soil erosion rather than to climatic changes (Zhang *et al.*, 2007). Changes in rainfall seasonality in monsoon systems also affect river flows. In South America, for example, wet seasons have increased in duration from an average of 170 days before 1972, to 195 days after (Carvalho *et al.*, 2010), with significant impacts on the Amazon and the La Plata basins.

18.3.1.2. Floods and Droughts

Floods, defined as impacts caused by the overtopping of river banks and levées, have increased in magnitude and frequency over recent decades in many river systems. There are limited instrumental records for these, and the existing gauging stations do not necessarily report whether impact-relevant overtopping has occurred. In addition, the highest annual flood will not necessarily imply the same overtopping each year. Other confounding factors, such as human alteration of river channels and land use also play a role, hence there is only *low to medium confidence* in global detection of a change in floods.

In regions with detected increase in heavy rainfall events and supposable consequences for pluvial floods (North America, Central Europe), both increases and decreases in floods have been found (Petrow and Merz, 2009; Villarini *et al.*, 2009). An attribution study using a multi-step modeling framework of floods suggests that an anthropogenic signal is detectable for a 20% increase in flood risk for the autumn 2000 floods in England and Wales (Kay *et al.*, 2011; Pall *et al.*, 2011, see also 18.4.4.2).

In mountain areas, glacial lake outburst floods (GLOFs) are characterized by their low frequency and high magnitude with most devastating impacts on downstream areas. While there is no evidence for a change in frequency or magnitude of GLOFs anywhere in the world (Seneviratne *et al.*, 2012), changes in the number and area of glacial lakes have been observed, with varying degrees of increasing trends in several regions over the Hindu Kush Himalayan mountain arc in the past two decades (Gardelle *et al.*, 2011), and a similarly strong increase in lake numbers in the Andes of Peru in the second half of the 20th century (Carey, 2005), and in northern Patagonia from 1945 to 2011 (Loriaux and Casassa, 2013). The growing number of these lakes is of concern since it suggests an increased likelihood of GLOFs.

Since the 1950s some regions of the world have experienced more intense and longer droughts (*medium confidence*), although a global trend cannot currently be established (Seneviratne *et al.*, 2012). Drought conditions have increased (with *medium confidence*) in Southern Europe and the Mediterranean, West Africa, East Asia, Southern Australia and New Zealand, and decreased in most of North America and Northern Australia (the rest of the world had either no change or insufficient data).

Groundwater storage has been reduced in large parts of the world, and this has been primarily attributed to human activities, such as in northeastern India where groundwater depletion, as detected by satellite data for the 21st century, was largely attributed to groundwater withdrawal for irrigation and other human use (Rodell *et al.*, 2009). Attribution of groundwater change to climatic drivers is more rare (Taylor *et al.*, 2012). For Kashmir (India), Jeelani (2008) suggests that the observed decline in groundwater recharge between 1981 and 2005 can be attributed to decreasing precipitation and glacier retreat, while a modeling study for southeast Spain indicates an effect of temperature related changes in evapotranspiration on groundwater (Aguilera and Murillo, 2009).

Water quality in watersheds and lakes is expected to change with increasing temperature through an increase in eutrophication. It is difficult, however, to link observed changes in water quality to climate change due to the confounding factors. Eutrophication is mostly driven by other causes, such as untreated sewage inflows in urban and industrial areas and surface runoff delivering residues of fertilizers used in agriculture, and vegetation decay and manure inputs in flooding events (Kundzewicz and Krysanova, 2010). There is emerging evidence for downstream impacts on water quality due to upstream climate impacts, such as high sulfide content in rivers of Peru's Cordillera Blanca due to sulfide-rich rocks that became exposed as glaciers retreated (Fortner *et al.*, 2011).

18.3.1.3. The Cryosphere

There is extensive evidence of significant recent changes in various components of the cryosphere, including glaciers, ice sheets and floating ice shelves, sea, lake and river ice, subsurface ice (permafrost) and snow (WGI AR5 Chapter 4). It is *likely* that there is an anthropogenic component in the changes observed in Arctic sea ice, Greenland surface melt, glaciers, permafrost and snow cover (WGI AR5 Chapter 10).

1 Changes in glaciers continue to provide a globally largely homogeneous but regionally variable signal of retreat.
2 There is *high confidence* that glacier changes over the past 2-3 decades exceed internal variability, but only few
3 studies are available that attribute these glacier changes to anthropogenic forcing (WGI AR5 Chapter 10). The
4 absolute contribution of glaciers and ice caps to sea level rise has increased since the early 20th century and has been
5 close to 1 mm yr⁻¹ for the past two decades (WGI AR5 Chapter 4), around a third of total observed sea level rise.
6 The decadal-scale mass loss of ice sheets and glaciers causes accelerated uplift of underlying land in the North
7 Atlantic Region (Jiang *et al.*, 2010). There is *medium confidence* regarding the effects of decadal ice loss on
8 seismicity due to unloading of the lithosphere beneath ice sheets (Hempel *et al.*, 2010) and Alaskan glaciers (Sauber
9 and Ruppert, 2008), and on volcanic activity such as enhanced magma generation (Sigmundsson *et al.*, 2010). The
10 strong and rapid downwasting observed on alpine glaciers has prompted a number of impacts. Expanding or new
11 lakes at the margin of many retreating glaciers in the Alps of Europe, Himalayas, Andes and other mountain regions
12 have altered the risk of outburst floods. In the Swiss Alps and the Peruvian Andes, outburst floods from several
13 lakes in the 21st century have caused damages, and required risk reduction measures on the order of tens of millions
14 USD (Huggel *et al.*, 2011; Carey *et al.*, 2012b). New glacier lakes have also become a tourist attraction, led to
15 additional infrastructure, and stimulated assessment of potential for hydropower generation (Terrier *et al.*, 2011).
16 There is also evidence of slope instabilities as a consequence of recent decadal scale glacier downwasting (Haerberli
17 and Hohmann, 2008; Huggel *et al.*, 2011).

18
19 Depending on local conditions, variations in runoff in high-mountain regions are often attributed to glacier and
20 climate change (Casassa *et al.*, 2009). Current understanding suggests that during the continuous shrinkage of
21 glaciers a ‘peak meltwater’ exists, with increasing runoff trends before, and decreasing trend after this threshold
22 (3.4.4). An increase in runoff from glacier areas has been documented for catchments in western and south-central
23 China over the past several decades, and for western Canada and Europe (Zhang *et al.*, 2008; Moore *et al.*, 2009; Li
24 *et al.*, 2010; Stahl *et al.*, 2010). In the Peruvian Andes there is evidence that ‘peak meltwater’ has recently been
25 passed, based on runoff decrease during the dry season in seven out of nine glacier-fed catchments in the Cordillera
26 Blanca (Baraer *et al.*, 2012), also confirmed by qualitative observations made by local people (Bury *et al.*, 2010;
27 Carey *et al.*, 2012a). In the Swiss Alps, positive variations in river runoff have occurred primarily in highly glaciated
28 catchments during warm periods of the 20th century (1940’s, 1990’s), while for less glaciated catchments a runoff
29 decrease was detected for the warm and dry 1990s (Collins, 2006; Pellicciotti *et al.*, 2010). For large catchments (Po
30 and Rhône catchments with <1% basin glacier cover), the contribution of glacier melt to total runoff in August was
31 significantly lower for 2004-2008 than for the previous twenty years (Huss, 2011).

32
33 Earlier reported trends of Arctic sea ice decline in terms of extent and thickness have continued, with a significant
34 increase of the rate of sea ice decline in the first decade of the 21st century (WGI AR5 Chapter 4). It is *likely* that at
35 least some of the decline in Arctic sea ice extent can be attributed to anthropogenic climate forcing (WGI AR5
36 Chapter 10). Observations by Inuit people in the Canadian Arctic confirm with *high confidence* the instrumental
37 observations on the various changes of sea ice (see Box 18-5). Antarctic sea ice has slightly increased over the past
38 30 years, yet with strong regional differences (WGI AR5 Chapter 4).

39
40 For lake and river ice, there is generally *high confidence* of later freeze-up and earlier break-up over the past 100+
41 years, yet with regional differences (WGI AR5 Chapter 4). Changes in lake and river ice can have effects on
42 freshwater ecosystems, transport and traffic over frozen lakes and rivers, and ice induced floods during freeze-up
43 and break-up events (Voigt *et al.*, 2011). Some evidence exists in Europe that ice-jam floods were reduced during
44 the last century due to reduced freshwater freezing (Svensson *et al.*, 2006).

45
46 Combined in-situ and satellite observations indicate a decline in snow cover extent in most months of the period
47 1922-2010, with the largest decline in spring (8%) (WGI AR5 Chapter 4). Only few formal detection and attribution
48 studies exist but they consistently indicate an anthropogenic influence on snow cover reduction (WGI AR5 Chapter
49 10), including an up to 60% contribution of anthropogenic climate forcing on changes in snow pack and runoff
50 timing between 1950 and 1999 in the Western United States (Barnett *et al.*, 2008). Impacts on winter tourism have
51 been observed (18.4.3.3).

52
53 Widespread changes and degradation of permafrost of both, high-latitude/low-land and high-elevation mountain
54 regions, have been observed over the past years and decades (WGI AR5 Chapter 4). Generally, the permafrost

1 boundary has been moving polewards and to higher elevations, and the active layer thickness has increased at many
2 sites. While formal attribution studies are hardly available for any of the above mentioned permafrost attributes,
3 several impacts have been related to permafrost changes, including an increase of flow speed of Alpine rock glaciers
4 in particular in the 21st century, resulting in rock fall and debris flows (Kääb *et al.*, 2007; Delaloye *et al.*, 2010),
5 expansion and deepening of thermokarst lakes, erosion at the Arctic coast, resulting in a doubling of the erosion rate
6 at Alaska's northern coastline over the past 50 years (Karl *et al.*, 2009). Furthermore, expansion of channel networks
7 (Toniolo *et al.*, 2009), increased river bank erosion (Costard *et al.*, 2007) and higher dynamics in shrinkage and
8 expansion of lakes and ponds have been observed in the Arctic (Rowland *et al.*, 2010), as well as an increase in
9 hillslope erosion and landsliding in Northern Alaska since the 1980's (Gooseff *et al.*, 2009). Complex feedbacks and
10 interactions across surface systems, and spatial and temporal scales complicate detection of drivers and effects. For
11 example, drying of land surface due to permafrost degradation may cause an increase in wildfires, in turn resulting
12 in a loss of ground surface insulation and change in surface albedo that accelerates permafrost thawing (Rowland *et*
13 *al.*, 2010; Forkel *et al.*, 2012).

16 18.3.1.4. Erosion, Landslides, and Avalanches

17
18 Erosion and landsliding typically increases during phases of deglaciation in mountain areas (Ballantyne, 2002;
19 Korup *et al.*, 2012) and there is emerging evidence for this to occur during contemporary deglaciation (Schneider *et*
20 *al.*, 2011; Uhlmann *et al.*, 2012). Erosion related to sediment flux changes from mountain areas has been observed in
21 the Western Himalaya in relation with hydrologic extreme events (medium confidence, Wulf *et al.*, 2012),
22 increasing over the past 60 years (Malik *et al.*, 2011), with important impacts on hydropower schemes. For southern
23 China, there is robust evidence of decline in sediment load in some rivers since the 1980s and 1990s (Zhang *et al.*,
24 2008a). Dam construction is an important driver of the recent decline in sediment load, leaving only *low confidence*
25 in attributing the change to any climate impacts for the Yangtze catchment in China (Xu *et al.*, 2008).

26
27 Changes in sediment yield, e.g. from rockfall or disintegration of rock glaciers, related or unrelated to climate
28 impacts, can significantly influence frequency and magnitude of alpine shallow landslides and debris flows (Lugon
29 and Stoffel, 2010), but no clear evidence exists so far for a change in frequency of shallow landslides and debris
30 flows from recently deglaciated mountain areas in the European Alps (Jomelli *et al.*, 2004; Stoffel and Huggel,
31 2012).

32
33 Rock slope failures in mountain areas with permafrost occurrence have increased since the 1990s (*high confidence*
34 in the Western European Alps, *medium confidence* for New Zealand Alps, and *low confidence* globally). There also
35 is *high confidence* that glacier retreat and downwasting, permafrost degradation and high-temperature events have
36 contributed to many high-mountain rock slope failures over the past 20 years (Allen *et al.*, 2010; Raveland and
37 Deline, 2011; Schneider *et al.*, 2011; Fischer *et al.*, 2012; Huggel *et al.*, 2012a). Damages and costs for risk
38 reduction measures in Alpine areas have increased due to rock fall and debris flows, with climate change playing a
39 major role in triggering complex downstream impact chains and feedback (*medium to high confidence* of influence
40 of anthropogenic climate forcing), such as recently documented in the Swiss Alps with costs on the order of tens of
41 millions USD (Huggel *et al.*, 2012b). Rock and ice avalanches, and other landslides from destabilized slopes have
42 also impacted glacier lakes and caused downstream damage in several high-mountain regions (e.g., Xin *et al.*, 2008;
43 Bajracharya and Mool, 2009; Künzler *et al.*, 2010; Carey *et al.*, 2012a; Huggel *et al.*, 2012b).

44
45 Other than for the above mentioned types of landslides, there is no clear evidence that their frequency or magnitude
46 has changed over the past decades (Huggel *et al.*, 2012b). This is true for shallow landslides and also for regions
47 with a relatively complete event record (e.g. Switzerland, see Hilker *et al.*, 2009). In general, detection of changes in
48 the occurrence of landslides is complicated by incomplete inventories, both in time and space, and inconsistency in
49 terminology. In line with loss from other extreme events (see 18.4.4) an increase in terms of casualties, or loss due
50 to landslides has been documented in South, East and South-East Asia over the past years, but are largely attributed
51 to changes in exposure, i.e. population growth (Petley, 2010).

1 No change in snow avalanche activity has been detected so far in Europe (Latenser and Schneebeli, 2002; Voigt *et*
2 *al.*, 2011). However, the detection of changes in snow avalanche impacts, such as fatalities and property loss, is
3 difficult over the past decades due to changes in snow sport activities and avalanche defense measures.
4
5

6 **18.3.2. Terrestrial and Inland Water Systems**

7

8 As documented by previous IPCC reports (notably Rosenzweig *et al.*, 2007), numerous changes in terrestrial and
9 inland water systems have been attributed to recent climate change. Confidence in such detection of change is often
10 *very high*, reflecting high agreement among many independent sources of evidence of change, and robust evidence
11 that changes in ecosystems or species are outside of their natural variation. Confidence in attribution to climate
12 change is also often *high*, due to process understanding of responses to climate change, or strong correlations with
13 climate trends and where confounding factors are understood to have limited importance. The scientific literature in
14 this field is growing quickly, precluding a full review in this chapter. Therefore, statements of confidence for
15 detection and attribution are given without references, as detailed traceability is provided in chapter 4.
16

17 Organisms respond to changing climate in a multitude of ways, including through their phenology (the timing of key
18 life history events such as flowering in plants or migration of birds), productivity (the assimilation of carbon and
19 nutrients in biomass), spatial distribution, mortality / extinction, or by invading new territory. Noticeable changes
20 may occur at the level of individual organisms, ecosystems, landscapes, or by modification of entire biomes.

21 Organisms and ecosystems are adapted to a variable environment, and they are capable to adapt to gradual change to
22 some degree – for the scope of this chapter, available knowledge will be reviewed concerning changes in terrestrial
23 and freshwater ecosystems that occur beyond natural variability and which can be assumed to be due to recent
24 climate change and/or increased in atmospheric CO₂. Confidence in the detection of such change involves therefore
25 assumptions about natural variability in these ecosystems, and confidence in the attribution of detected change to
26 climate drivers (or CO₂) implies the assessment of confounding drivers such as land use change.
27

28 **18.3.2.1. Phenology**

29

30
31 Since the AR4 there has been a further significant increase in observations of phenology of plants and animals,
32 showing that many, but not all species have changed functioning to some degree over the last decades to centuries
33 on all continents (*high confidence* due to robust evidence but only moderate agreement across all species). New
34 satellite-based analyses confirm earlier trends, showing, for example, that the onset of the growing season in the
35 northern hemisphere has advanced by 5.4 days from 1982 to 2008 and its end has been delayed by 6.6 days (Jeong *et*
36 *al.*, 2011). Significant changes have been detected, by direct observation, for many different species, for example,
37 for amphibians, birds (breeding, migration), mammals, and plants; a number of meta-analyses have been carried out
38 summarizing this literature (e.g., Cook *et al.*, 2012). Attribution of these changes to climate change is supported by
39 more refined analyses that consider also the regional changes in several variables such as temperature, growing
40 season length, precipitation, snow cover duration and others, as well as experimental evidence. The *high confidence*
41 in attributing many observed changes in phenology to changing climate is a result of these analyses, as well as of
42 improved knowledge of confounding factors such as land use and land management (for more references and details
43 see 4.3.2.1).
44

45 **18.3.2.2. Productivity and Biomass**

46

47
48 Many terrestrial ecosystems are now net sinks for carbon over much of the Northern hemisphere and also in parts of
49 the Southern hemisphere (*high confidence*). This is shown, for example, by inference from atmospheric chemistry,
50 but also by direct observations of increased tree growth in many regions including Europe, the United States,
51 tropical Africa and the Amazon. During the decade 2000 to 2009, global land net primary productivity was approx.
52 5% above the preindustrial level, contributing to a net carbon sink on land of 2.6 ± 0.7 Pg C yr⁻¹ (Raupach *et al.*,
53 2008; Le Quéré *et al.*, 2009, WGI AR5 Chapter 6), despite ongoing deforestation. These trends are in part due to
54 nitrogen deposition, afforestation and altered land management which makes direct attribution of the increase to

1 climate change difficult (*low confidence* in attribution). The degree to which rising atmospheric CO₂ concentrations
2 contribute to this trend remains a particularly important source of uncertainty (for more references and details, see
3 4.3.2.2 and 4.3.2.3).

6 18.3.2.3. Biodiversity

8 Each species responds differently to a changing environment, therefore the composition of species, genotypes,
9 communities and even ecosystems varies in different ways from place to place, in response to climate change. The
10 consequences are changing ranges of species, changing composition of the local species pool, invasions, mortality
11 and ultimately extinctions. For different species and species groups, detected range shifts vary, and so do the
12 confidence of detection and the degree of attribution to climate change. The number of species studied has
13 considerably increased since the AR4. Overall, many terrestrial species have recently moved, on a global average,
14 17 km poleward and 11 m up in altitude per decade (e.g., Europe, North America, Chile, Malaysia), which
15 corresponds to predicted range shifts due to warming (Chen *et al.*, 2011) and is 2 to 3 times faster than previous
16 estimates (Parmesan and Yohe, 2003; Fischlin *et al.*, 2007), *high confidence* in detection. For example, over the last
17 decades, arthropods have moved large and statistically significant distances towards the poles (many 10s of km).
18 Species with short life cycles and high dispersal capacity – such as butterflies (*high confidence* in attribution) – are
19 generally tracking climate more closely than longer-lived species or those with more limited dispersal such as trees
20 (*medium confidence* in attribution). There are many less well-studied species for which detection of change and its
21 attribution to climate change are more uncertain.

23 Across the world, species extinctions are at or above the highest rates of species extinction in the fossil record (*high*
24 *confidence* in detection). However, only a small fraction of observed species extinctions have been attributed to
25 climate change — most have been ascribed to non-climatic factors such as invasive species, overexploitation or
26 habitat loss (*very low confidence* in attribution to climate change). For those species where climate change has been
27 invoked as a causal factor in extinction (such as for the case of Central American amphibians), there is *low*
28 *agreement* among investigators concerning the importance of climate variation in driving extinction and even less
29 agreement that extinctions were caused by global warming. *Confidence* in the attribution of extinctions across all
30 species to climate change is *very low*.

32 Species invasions have been increasing over the last several decades world-wide, notably in freshwater ecosystems
33 (*very high confidence*), often causing biodiversity loss or other negative impacts. While there is a documented
34 contribution of recent climate trends to establishment, growth, spread and survival of some invasive species
35 populations, there is only *low confidence* that the species invasions have generally been assisted by recent climatic
36 trends because of the overwhelming importance of human facilitated dispersal in mediating invasions (for more
37 references and details see 4.3.2.5 and 4.3.2.6).

40 18.3.2.4. Impacts on Major Systems

42 The extent of recent change in major ecosystems such as the boreal forest, the Arctic tundra, or the Amazon forest
43 has been characterized as a “regime shift” by some authors (e.g., Biggs *et al.*, 2009), implying significant and broad-
44 scale changes in both distribution and functioning of the ecosystem.

46 Field and satellite measurements indicate a substantial increase in shrub growth (often linked to permafrost thawing)
47 in many areas of the Arctic tundra (*high confidence* in detection). This change corresponds to expectations, based on
48 experiments, models and paleoecological responses to past warming, of broad-scale boreal forest encroachment into
49 tundra, a process which takes decades and which would have very large impacts on ecosystem structure and
50 function. The particular strength of warming over the last 50 years for most of the Arctic further facilitates
51 attribution (*high confidence*). The change affects a significant area of the tundra biome and can be considered an
52 early warning for an upcoming regime shift.

1 For the boreal forest, increases in tree mortality are observed in many regions, including wide-spread dieback related
2 to insect infestations in North America, but there is *low confidence* in detection of a global trend. Local and regional
3 mortality has in some cases been linked to climate fluctuations, but there is *low confidence* in attributing any
4 perceived overall dieback to climate change. At the Southern “trailing end” of the shifting boreal zone, there are
5 indications of enhanced mortality of plant species (notably trees), but there is *low confidence* in overall detection
6 due to a lack of temporal and spatial coverage of observations. There is consequently also *low confidence* in
7 attribution of the changes to climatic change.

8
9 In the humid tropical forests of the Amazon basin, increases in tree turnover (increased mortality and growth) have
10 been detected with *medium confidence* for recent decades. These changes have been explained by a number of
11 factors, including the direct effect of rising CO₂ on lianas, recovery from past disturbance and changing climate.
12 Overall, there is *very low confidence* in attribution of these observations to climate change.

13
14 In freshwater ecosystems of most continents and climate zones, rising temperatures have been linked to shifts in
15 invertebrate and fish community composition, especially in headwater streams where species are more sensitive to
16 warming (*high confidence* in detection, *low confidence* in attribution due to numerous confounding factors; 4.3.3.1,
17 4.3.3.3).

18 19 20 **18.3.3. Coastal Systems and Low Lying Areas**

21
22 In coastal waters, both average warming rate and changes in seasonal timing are larger than in the open oceans (see
23 5.3.3). Sea surface temperatures have warmed significantly during the past 30 years along more than 70% of the
24 world’s coastlines, albeit with large spatial and seasonal variation in the rates of change (*very high confidence*). The
25 frequency of extreme temperature events has changed in many areas (Lima and Wethey, 2012). Seawater pH also
26 spans larger ranges and exhibits higher variability near coastlines, and anthropogenic ocean acidification can be
27 enhanced or lessened by coastal geochemical processes (Borges and Gypens, 2010; Feely *et al.*, 2010; Duarte *et al.*,
28 2013, see also Box CC-OA), and extreme winds, changes in wave regime and sea level play important roles in
29 coastal processes (see Table 5-1).

30
31 While it is *extremely likely* that observed global sea level rise can at least in part be attributed to anthropogenic
32 emissions (WGI AR5 Chapter 10.4.3), the evaluation of local sea level trends must consider important local
33 confounding factors, such as regional variability in ocean and atmospheric circulation, subsidence, coastal erosion,
34 and coastal modification (see also 5.3.2). Thus far, it has not been possible to isolate an anthropogenic climate signal
35 in local sea level changes from the contributions of these confounding factors. A possible emerging exception is
36 along the coasts of regions with melting glaciers and ice sheets, where local gravitational effects leading to a
37 lowering of sea level may dominate other factors (Kopp *et al.*, 2010; Tamisiea and Mitrovica, 2011). As a
38 consequence, attribution of observed effects of relative sea level change to climate change is hardly possible (see
39 Nicholls *et al.*, 2007; Nicholls *et al.*, 2009), despite attempts for very few locations, e.g. for flood damages in Venice
40 (Carbognin *et al.*, 2010).

41 42 43 **18.3.3.1. Erosion, Shoreline Processes, and Coastal Aquifers**

44
45 Throughout the world, beaches and dunes, as well as bluffs and cliffs, are eroding due to a variety of climate related
46 processes, such as rising mean sea levels (Leatherman *et al.*, 2000; Ranasinghe and Stive, 2009), more frequent
47 extreme sea levels (Woodworth *et al.*, 2011), changes in wave regimes (Tamura *et al.*, 2010; Reguero *et al.*, 2013),
48 the loss of natural protective structures such as coral reefs (Grevelle and Mimura, 2008) or mangrove forests due to
49 increased ocean temperatures or ocean acidification (Bongaerts *et al.*, 2010), or permafrost degradation and sea ice
50 retreat (Manson and Solomon, 2007). However, there are multiple non-climate related drivers involved in shoreline
51 erosion, including dams capturing fluvial sands, subsidence due to resource extraction, mining and coastal
52 engineering and development. Due to the fragmentary nature of the information available and to the multiple natural
53 and anthropic stressors contributing to coastal erosion, confidence in attribution of shoreline changes to climate
54 change is *very low*, with the exception of polar regions (Forbes, 2011, see also 18.5).

1
2 Coastal lagoons and estuaries, as well as deltas are highly susceptible to alterations of sediment input and
3 accumulation (Syvitski *et al.*, 2005; Ravens *et al.*, 2009), processes that can be influenced by climate change via
4 changes in sea level, storminess, and precipitation. However, the primary drivers of widespread observed changes in
5 those systems are human drivers other than climate change (thereby *very low confidence* in attribution to climate
6 change, see 5.4.2.6, 5.4.2.7).

7
8 Coastal aquifers are crucial for the water supply of densely populated coastal areas, in particular in Small Island
9 environments. Aquifer recharge is sensitive to changes in temperature and precipitation; and rising sea levels and sea
10 water overwash from storm surges can contribute to saline intrusion into groundwater (Post and Abarca, 2010; Terry
11 and Falkland, 2010; White and Falkland, 2010, see also 29.3, Table 18-8). However, excessive groundwater
12 extraction for coastal settlements and agriculture is the main cause for widely observed groundwater degradation in
13 coastal aquifers (e.g., White *et al.*, 2007a; Barlow and Reichard, 2010). Attribution to climate change, in particular
14 incremental sea level rise, is not supported in the literature (Rozell and Wong, 2010; White and Falkland, 2010).

15 16 17 *18.3.3.2. Coastal Ecosystems*

18
19 Coastal habitats and ecosystems experience cumulative impacts of land- and ocean-based anthropogenic stressors
20 (Halpern *et al.*, 2008). Most coral reefs, seagrass beds, mangroves, rocky reefs and shelves have undergone
21 substantial changes. Coral reefs have been degraded due to both local anthropogenic factors such as unsustainable
22 fishing and pollution, and global change factors such as increased heat waves, ocean warming and acidification (see
23 also Box CC_CR). Coral bleaching is being detected with *high confidence* on all coasts, and warming is a major
24 contributor (*high confidence*, for further discussion see Box 18-3, and Box CC_OA). Overexploitation and habitat
25 destruction have been responsible for a large fraction of historical changes observed in coastal ecosystems (Lotze *et al.*, 2006).

26
27
28 Hypoxia poses a serious threat to marine life, which is exacerbated when combined with elevated temperature
29 (Vaquer-Sunyer and Duarte, 2011). Increased loads of nutrients from anthropogenic sources generate coastal
30 eutrophication and constitute the primary cause of increasing hypoxia, while upwelling of low oxygen waters and
31 ocean warming constitute secondary drivers (see Zhang *et al.*, 2010). Persisting hypoxia can result in so-called
32 “dead zones”, which have approximately doubled each decade since 1960 (Diaz and Rosenberg, 2008).

33
34 Changes in abundance and distribution of rocky shore species have been observed since the late 1940s in the North
35 East Atlantic (Hawkins *et al.*, 2008), and the role of temperature has been demonstrated by experiments (e.g., Peck
36 *et al.*, 2009; Somero, 2012, see also 5.4.2.2). Distinguishing the response to climatic stressors from changes due to
37 hydrology, or natural temporal and spatial fluctuations, is nevertheless challenging.

38
39 Globally, the range limits of many intertidal species have shifted up to 50 km per decade, much faster than most
40 recorded shifts of terrestrial species (Helmuth *et al.*, 2006, see also Box 18-4). However, the geographical
41 distribution of some species did not change in the past decades. This may be due to weak local warming
42 (Rivadeneira and Fernández, 2005), or overriding effects of variables such as timing of low tide, hydrographic
43 features, lack of suitable bottom types, larval dispersal, food supply, predation and competition (Helmuth *et al.*,
44 2002; Helmuth *et al.*, 2006; Poloczanska *et al.*, 2011). Changes in current patterns and increased storminess can
45 dislodge benthic invertebrates and affect the distribution of propagules and recruitment.

46
47 Changes in musselbeds in response to higher temperatures induced by climatic change have been observed along the
48 West coasts of the United States (Smith *et al.*, 2006a; Menge *et al.*, 2008; Harley, 2011). On Tatoosh Island
49 (Washington, USA), a shift in community structure from a mussel to an algal-barnacle-dominated community has
50 been attributed to rapidly declining pH (Wootton *et al.*, 2008; Wootton and Pfister, 2012).

51
52 Ocean warming is also leading to range shifts in vegetated coastal habitats such as coastal wetlands, mangrove
53 forests and seagrass meadows (*medium confidence*, see 5.4.2.3). Extreme temperature events can alter marine and

1 coastal communities, as shown for the European heatwave 2003 (Garrahou *et al.*, 2009), and the early 2011 heat
2 wave off the Australian West Coast (Wernberg *et al.*, 2012).

3
4 Poleward expansion of mangrove forests, consistent with expected behavior under climate change, has been
5 observed in the Gulf of Mexico (Perry and Mendelssohn, 2009; Comeaux *et al.*, 2011; Raabe *et al.*, 2012), and New
6 Zealand (Stokes *et al.*, 2010). High temperatures have impacted seagrass biomass in the Atlantic Ocean (Reusch *et*
7 *al.*, 2005; Díez *et al.*, 2012; Lamela-Silvarrey *et al.*, 2012), the Mediterranean Sea (Marbà and Duarte, 2010) and
8 Australian waters (Rasheed and Unsworth, 2011). Extreme weather events contributed to the overall degradation of
9 seagrass meadows in a Portuguese estuary (Cardoso *et al.*, 2008).

10
11 Decline in kelp populations attributed to ocean warming has occurred off the North coast of Spain (Fernández,
12 2011), as well as in southern Australia, where the poleward range expansion of some herbivores have also
13 contributed to observed kelp decline (Johnson *et al.*, 2011; Wernberg *et al.*, 2011; Wernberg *et al.*, 2011). The
14 spread of subtropical invasive macroalgal species (e.g., Lima *et al.*, 2007) may be adding to the stresses temperate
15 seagrass meadows experience from ocean warming.

16
17 Overall, there is *high confidence* in detection of range shifts and biodiversity changes in intertidal and other coastal
18 species, as well as in a decline in kelp forests, and increased mortality of seagrasses, and *high confidence* that
19 climate change has contributed to these effects (5.4.4, Figure 5-5). For the widespread decline observed in salt
20 marshes and mangroves, there is *very low confidence* in a role of climate change due to overriding effects of other
21 human drivers.

22 23 24 18.3.3.3. Coastal Settlements, Infrastructure, and Economic Activities

25
26 Recent global (e.g., Menéndez and Woodworth, 2010; Woodworth *et al.*, 2011) and regional studies (e.g., Marcos *et*
27 *al.*, 2009; Haigh *et al.*, 2010; Haigh *et al.*, 2011) have found that observed trends in extreme sea levels are mainly
28 consistent with mean sea level trends (Woodworth *et al.*, 2011), indicating that the increasing frequency of extreme
29 events affecting coastal infrastructures observed so far is related to rising mean sea level rather than to changes in
30 the behaviour of severe storms. Increased damages from coastal flooding have been observed with *high confidence*,
31 however, with exposure and subsidence constituting the major drivers, confidence in attribution to climate change is
32 *very low* (Seneviratne *et al.*, 2012, see also 5.4.3, 5.4.4).

33
34 Increases in saltwater intrusion and flooding have been observed in low-lying agricultural areas of deltaic regions
35 and Small Islands, but the contribution of climate change to this is not clear (e.g., Rahman *et al.*, 2011, see also
36 18.5.9). Both climate variability and change impact fishermen livelihoods (Badjeck *et al.*, 2010) and physiological
37 and ecological properties of fish (e.g., Barange and Perry, 2009, see also 18.3.4, 18.4.1.2); however local and
38 regional observations of climate impacts on output of coastal fisheries are scarce.

39
40 While vulnerability of coastal settlements and infrastructure to future climate change, in particular sea level rise and
41 coastal flooding, is widely accepted and well-documented (see Table 5-5), there is a shortage of studies on the role of
42 climate change in observed impacts on coastal systems.

43
44 _____ START BOX 18-3 HERE _____

45 46 **Box 18-3. Detection and Attribution of Mass Coral Bleaching and Mortality to Climate Change**

47
48 Declining water quality as well as increasing fishing pressure and coastal development have been implicated in the
49 rapid decline in the abundance of corals and coral reefs over the past 50 years (Bryant *et al.*, 1998; Gardner *et al.*,
50 2003; Bruno and Selig, 2007; Sheppard *et al.*, 2010; Burke *et al.*, 2011; De'ath *et al.*, 2012). Since 1980, mass coral
51 bleaching and mortality events began to occur on reefs throughout the tropics and subtropics with no precedent in
52 the scientific literature (Box CC-CR). While bleaching of individual coral colonies has been reported prior to 1980
53 (Yonge and Nichols, 1931), mass coral bleaching events involving hundreds and thousands of coral colonies across
54 entire reef and coastal regions have not (Hoegh-Guldberg, 1999; Baker *et al.*, 2008). These novel events are often

1 followed by the mass mortality of coral communities, especially if conditions remain anomalously warm for long
2 periods (ibid). In the very warm year of 1998, for example, mass coral bleaching affected almost every part of the
3 seas associated with the tropics and subtropics resulting in the loss of 16% of the world's reef-building coral
4 (Wilkinson and Hodgson, 1999).
5

6 There is broad agreement that mass coral bleaching can be triggered by small increases in sea temperature ($> 1^{\circ}\text{C}$)
7 above the summer maxima for a region over several weeks (very high confidence, Hoegh-Guldberg, 1999; Baker *et al.*,
8 2008; Strong *et al.*, 2011). The impact of temperature is also exacerbated by strong solar irradiance (Hoegh-
9 Guldberg, 1999). There is also broad agreement that thermal stress impacts the ability of *Symbiodinium* to capture
10 and process light, leading to the production of damaging reactive oxygen species which precedes the loss of
11 symbionts (Jones *et al.*, 1998). As the symbiosis breaks down, the brown dinoflagellates leave their host corals,
12 turning their tissues white and depriving corals of an important source of energy (Muscatine, 1986; Hoegh-Guldberg
13 and Smith, 1989).
14

15 There is little or no evidence that reef-building corals and their *Symbiodinium* have, or evolved substantially in terms
16 of the thermal tolerance over yearly or decadal time frames, or can be expected to do so (Hoegh-Guldberg, 2012).
17 The relationship between elevated sea temperatures and mass coral bleaching remains robust enough to be used
18 within algorithms that reliably detect and project the incidence of mass coral bleaching from satellites (Strong *et al.*,
19 2004; Strong *et al.*, 2011). Depending on the size of the thermal anomaly and exposure time, communities of reef-
20 building corals have either recovered, or have experienced large-scale mortalities such as those seen worldwide in
21 1998 (Hoegh-Guldberg, 1999; Wilkinson and Hodgson, 1999; Baker *et al.*, 2008) and 2005 in the Caribbean (Eakin
22 *et al.*, 2010). The detection of mass coral bleaching has been improved using satellite algorithms such as Degree
23 Heating Weeks (DHW) (ibid).
24

25 Of the many variables investigated, only elevated sea temperature co-occurs consistently with mass coral bleaching
26 and mortality events, and has a well worked physiological model associated with it (Hoegh-Guldberg, 1999).
27 Consequently, given the relationship between global climate change and increased ocean temperatures, and that
28 between coral bleaching and elevated temperature, the detected increase in the frequency of mass coral bleaching
29 and mortality events can be attributed to anthropogenic climate change as the dominant driver with *very high*
30 *confidence*.
31

32 _____ END BOX 18-3 HERE _____
33
34

35 **18.3.4. Oceans**

36
37 Since 1970, ocean temperatures have increased by around $0.1^{\circ}\text{C decade}^{-1}$ in the upper 75 m and approximately
38 $0.02^{\circ}\text{C decade}^{-1}$ in the upper 500m, due to anthropogenic warming.
39

40 The increased flux of CO_2 from the atmosphere to the ocean has reduced the average pH of seawater by 0.1 pH units
41 over the past century, with the greatest reduction occurring at high latitudes (Box CC-OA). These changes have been
42 attributed to increases in the atmospheric concentration of greenhouse gases as result of human activities (*high to*
43 *very high confidence*, WGI AR5 Chapter 10.4.1-10.4.4). Changes in wind speed, upwelling, water column
44 stratification, surface salinity, ocean currents, solar irradiance, cloud distribution, and oxygen depth profile have also
45 occurred (WGI AR5 Chapter 3.2 – 3.8; Figure 30-5, 30-6, 30-7).
46

47 Changes in the physical and chemical nature of ocean environments are predicted to have impacts on marine
48 organisms and ecosystems, with many already having been observed across most ocean regions (see 6.2, 6.3, 6.5,
49 30.4, 30.5). However, the detection and attribution of recent changes in ocean systems is complicated by the
50 influence of long-term variability such as the Pacific Decadal Oscillation (PDO), El Niño-Southern Oscillation
51 (ENSO), and the Atlantic Multidecadal Oscillation (AMO). The fragmentary nature of ocean observations and the
52 influence of confounding factors such as fishing, habitat alteration, and pollution also represent significant
53 challenges to detection and attribution (Hoegh-Guldberg *et al.*, 2011; Parmesan *et al.*, 2011).
54

1
2 *18.3.4.1. Impacts on Ocean System Properties and Marine Organisms and Ecosystems*
3

4 Greater thermal stratification in many regions has reduced ocean ventilation and mixing depth, thereby reducing the
5 availability of inorganic nutrients and hence primary productivity in surface layers, with controversial trends
6 depending on methodology (see 6.1.3, 6.2.3, 30.5.6). However, upwelling has increased in some regions bringing
7 greater concentrations of nutrients to surface waters, boosting productivity (see 30.5.5) and enhancing fisheries
8 output (see also 18.4.1.2 for fisheries). Increases in productivity also occurred with warming and sea ice loss at high
9 latitude. At a global scale a small increase in net primary production has been detected (Table 18-1, *medium*
10 *confidence*).

11
12 Poleward shifts in the distributions of zooplankton, fish, seabirds and benthic invertebrate have been observed,
13 particularly from the well-studied NE Atlantic, with a clear attribution to warming (*high confidence*, 6.3.2, 30.5.1).
14 In many regions, Temperature exerts the strongest influence on ecosystems and the responses of ecological systems
15 to changing temperature are well-studied. However, it is often difficult to clearly identify the interaction of
16 temperature with other factors (6.3.5). Some studies have found changes in the abundance of fish species that are
17 consistent with regional warming, with differences in response between species, in line with differential
18 specializations of coexisting species (6.2, 6.3.2, Pörtner, 2012). Anthropogenic influences modulate responses to
19 climate, e.g., due to exploitation status (Tasker, 2008; Belkin, 2009; Overland *et al.*, 2010; Schwing *et al.*, 2010),
20 with more heavily exploited species being more sensitive to environmental variability in general, including
21 temperature trends and extremes (Hsieh *et al.*, 2005; Stige *et al.*, 2006; Hsieh *et al.*, 2008).
22

23 Changes to water column mixing have combined with other factors such as nutrient loading to drive down oxygen
24 concentrations and increase the number and extent of hypoxic zones. These zones are characterized by very low
25 oxygen and high CO₂ levels and have been detected with *high confidence*, and attributed in part to enhanced
26 stratification and microbial respiration caused by warming (*medium confidence*, see Table 18-1). In some cases,
27 expanding or shifted hypoxia exerts strong local and regional effects on marine biota such as distribution shifts,
28 habitat contraction or loss, and fish kills.
29

30 Laboratory experiments have shown that a broad range of marine organisms (e.g. corals, fish, pteropods
31 coccolithophores, and macroalgae), physiological processes (e.g., skeleton formation, gas exchange, reproduction,
32 growth and neural function), and ecosystems processes (e.g., productivity, reef building and erosion) are sensitive to
33 changes in pH and carbonate chemistry of seawater (*high confidence*, 6.2.2-5, 6.3.4, Box CC-OA). However, few
34 field studies have been able to detect and attribute specific changes in marine ecosystems to anthropogenic ocean
35 acidification due to the inability to identify the effect of ocean acidification from ocean warming or local factors
36 (Wootton *et al.*, 2008; De Moel *et al.*, 2009; Moy *et al.*, 2009; Bednaršek *et al.*, 2012, see also 6.3.4).
37

38 [INSERT TABLE 18-1 HERE

39 Table 18-1: Observed changes in ocean system properties and their effects, with confidence levels for detection and
40 attribution to climate change, based on assessment in Chapter 6.2, 6.3, and summarized in 6.6.1 and Figure 6-16.
41 Observed impacts related to temperature effects have been attributed to warming, although the relative contributions
42 of regional climate variation and long-term global trends have not been quantified.]
43

44 There has been a substantial increase in the number of studies documenting significant changes in marine species
45 and processes since the AR4 (Hoegh-Guldberg and Bruno, 2010). A new meta-analysis using a database of long-
46 term observations from peer-reviewed studies of biological systems, with nearly half of the time series extending
47 prior to 1960, shows that a high proportion (81%) of observed responses are consistent with regional climate change
48 (see 30.4). Poloczanska *et al.* (2013) argue that the high consistency of marine species' responses across geographic
49 regions (coastal to open ocean, polar to tropical), taxonomic groups (phytoplankton to top predators), and types of
50 responses (distribution, phenology, abundance) reported in their analysis support the detection of a widespread
51 impact of climate change on marine populations and ecosystems (see 30.4 and 30.5 for more detail).
52

1 The impacts of climate change on marine organisms such as fish, invertebrates and marine algae include changes in
2 abundance, distribution, and community structure and shifts in phenology and migration patterns. Table 18-2 gives
3 examples of the manifestation of climate change on marine species.

4
5 [INSERT TABLE 18-2 HERE

6 Table 18-2: Observed changes on marine species and ecosystems, with confidence levels for assessment of detection
7 of changes and their attribution to climate change, based on assessment in Chapter 6.2, 6.3, summarized in 6.6.1 and
8 Figure 6-16, and 30.4. Observed impacts related to temperature effects have been attributed to warming, although
9 the relative contributions of regional climate variation and long-term global trends have not been quantified.]

10 11 12 *18.3.4.2. Observed Climate Change Effects across Ocean Regions*

13
14 While climate change is evident across the Ocean, its impacts vary between ocean regions (Table 30.2; Figures 30-2,
15 30.3 and 30.5; WGI AR5 Chapter 3). Considerable differences in system understanding, data availability, and the
16 potential contribution of climate change in relation to other factors add to the heterogeneity of the assessment.

17 Attribution of regional heat content changes are less certain than on a global level, but warming patterns have been
18 detected in all basins (Table 30-2, Figure 30-2) and attributed to anthropogenic influence with *high confidence* (WGI
19 AR5 Chapters 3, 10.3.1, and 10.4), with the important exception of Eastern Boundary Upwelling Ecosystems
20 (EBUE), where two of the four major upwelling systems show no change in sea surface temperatures over the last
21 60 years (table 30-2). These differences may relate to geographical differences that contribute to different levels of
22 change in wind stress and hence upwelling, which influences the amount of cooler water flooding the surface layers
23 of the water column in these regions (see Box 30.8.2). Recent research shows deep penetration of warming in some
24 regions, and declining oxygen levels (*low to medium confidence*, see Table 18-3). Regional estimates of CO₂ uptake
25 are in line with global estimates, and ocean acidification has been detected and attributed with *high confidence* in
26 most regions (WGI AR5 Chapter 3.8.1, 30.5, Box CC-CAO). Table 18-3 shows confidence in detection and
27 attribution of observed climate change impacts across the world's major open ocean regions, with the exception of
28 the Deep Sea.

29
30 [INSERT TABLE 18-3 HERE

31 Table 18-3: Confidence in detection and attribution of observed climate change effects across ocean regions, based
32 on expert assessment in Chapter 30.5 (respective subsections are given below). Confidence levels assigned are for
33 detection / attribution, respectively.]

34
35 The high latitude spring bloom systems of the Northern Hemisphere show strong warming and associated effects
36 (see above). In the North Pacific, the Bering Sea is one of the productive sub-regions and has undergone major
37 changes in recent decades as a result of climate variability, climate change and fishing impacts (Litzow *et al.*, 2008;
38 Mueter and Litzow, 2008; Jin *et al.*, 2009; Hunt, 2010). Loss of sea ice has led to the retreat of the cold pool in the
39 Bering Sea, and northward expansion of productivity.

40
41 Marginal seas such as the East China Sea are also warming rapidly. There is *high confidence* that this has
42 contributed to declining primary productivity and fisheries yields as well as other ecological changes (30.5.4.1).
43 However, other human pressures including over-fishing, habitat alteration, and nutrient loading are important
44 contributing factors and it is difficult to disentangle these from the impacts of climate change.

45
46 Semi-enclosed seas like the Black and Baltic Seas and the Arabian / Persian Gulf have their own specific
47 environmental conditions and responses (30.5.3.1). The Baltic and Black Seas show an expansion of hypoxic zones
48 attributed in part to climate change (*medium confidence*). Coral reefs in the Arabian / Persian Gulf and Red Seas
49 appear to have experienced widespread bleaching in 1996 and 1998 associated with elevated temperature. It is
50 highly likely that these impacts are associated with long-term variability that has, combined with climate change,
51 resulted in warmer than normal summer sea temperatures that now periodically exceed the thermal tolerance of reef-
52 building corals (*high confidence*).

1 Warming of the Mediterranean has been associated with mass mortality events as well as with invasions and
2 subsequent spread of new warm water species, which has resulted in the 'tropicalisation' of fauna (30.5.3.1.3). In
3 many tropical regions and the subtropical gyres of the Pacific, Indian, and Atlantic, periodic heat stress combined
4 with other local stresses has driven mass coral bleaching and mortality (see also Box CC-CR, 30.5).

5
6 In other regions, such as the California Current upwelling system, high-quality databases support *very high*
7 *confidence* in the detection of ecological changes, but attribution of these to climate change can be made with *very*
8 *low confidence* due to large-scale environmental variability associated with ENSO and the PDO.

9
10 In overall terms, attributing observed local and regional changes in marine species and ecosystems to climate change
11 remains an important question for on-going research (Stock *et al.*, 2010).

12
13 _____ START BOX 18-4 HERE _____

14 15 **Box 18-4. Differences in Detection and Attribution of Ecosystem Change on Land and in the Ocean**

16
17 Marine and terrestrial ecosystems differ in fundamental ways. Gradients in turbulence, light, pressure and nutrients
18 uniquely drive fundamental characteristics of organisms and ecosystems in the ocean. While the critical factor for
19 transporting nutrients to marine primary producers ocean mixing driven by wind, water is the primary mode for
20 transporting nutrients to land plants. In addition to these characteristics, marine ecosystems are often more
21 technically difficult and costly to explore than terrestrial equivalents, which explains the low number and shorter
22 scientific studies of marine ecosystems (Hoegh-Guldberg and Bruno, 2010). The latter has restricted the extent to
23 which we can accurately detect and attribute of changes within the Ocean.

24
25 Impacts of climate change in terrestrial and marine systems differ significantly for the same types of measures, e.g.,
26 species phenology and range shifts, leading to differences in expert's interpretations of the data and possibly
27 divergent levels of confidence in detection and attribution. There are also fundamental differences in exposure of
28 organisms to recent warming, their biological responses and our ability to detect change through observations.
29 Changes in temperature of ocean systems have generally been less than those of terrestrial ecosystems over the last
30 four decades (Burrows *et al.*, 2011). Furthermore, despite higher variability the horizontal spatial gradient of
31 temperature change ($^{\circ}\text{C km}^{-1}$) is generally much higher in terrestrial ecosystems than in marine ecosystems. All else
32 being equal, the net result is that species have generally needed to move much shorter distances in terrestrial
33 ecosystems to stay within their preferred climates, also due to the influence of the topography such as mountain
34 ranges (Burrows *et al.*, 2011), although many marine species can potentially exploit strong vertical thermal gradients
35 to attenuate the need for range shifts in response to warming.

36
37 Species and ecosystems may respond very differently to these climate signals in ways that influence the ability to
38 detect change. For example, a comparison of ectotherm species (i.e., species that do not actively regulate their body
39 temperatures such as reptiles and fish) indicates that marine species' ranges have tracked recent warming at both
40 their poleward and equatorial range limits, while many other terrestrial species' ranges have only tracked warming at
41 their poleward range limits (Sunday *et al.*, 2012). Biological processes influencing phenological shifts may also
42 differ substantially between systems. For example, the effect of climate on the timing of flowering of terrestrial
43 plants at high latitudes is only moderately influenced by confounding effects, whereas the timing of phytoplankton
44 blooms in high latitude marine systems is highly dependant on ocean temperature and associated stratification and
45 changes in nutrient availability.

46
47 _____ End BOX 18-4 HERE _____

48 49 50 **18.4. Detection and Attribution of Observed Climate Change Impacts in Human and Managed Systems**

51
52 Observed impacts on human systems have received considerably less attention in previous IPCC reports and the
53 scientific literature, compared to observed impacts on natural systems. Human systems' "normal state in the absence
54 of climate change" is almost never stationary. Confounders other than climate change have been and continue to

1 drive the normal evolution of these systems with climate often playing a relatively minor role. It is therefore difficult
2 to detect and attribute the signal of climate change in the majority of human systems. The food system is one
3 noteworthy exception. There is emerging literature estimating the climate *sensitivity* of many sectors within the
4 human system, yet climate impacts are often not detectable over the impacts from non-climate confounders.
5

6 For some human systems, the only observed situations where a climate signal had a detectable and sometimes
7 attributable impact are during extreme weather events. Extreme events for a variety of sectors are therefore
8 discussed in a single section below. Overall, the literature has made significant progress for certain sectors, such as
9 food systems, since AR4. The following sections provide a synthesis of findings with regard to food systems, cities,
10 economic systems, human health, human security and human livelihoods and poverty, which are documented in
11 greater detail in chapters 7, 8, 9, 10, 11, 12 and 13. We have also incorporated evidence from regional chapters and
12 further available literature, especially for the discussion of extreme events, human security, and observed changes in
13 indigenous communities.
14

16 **18.4.1. Food Production Systems**

17

18 Over the past several decades food production systems across the globe have changed significantly. Factors other
19 than atmospheric CO₂ or climate, such as cultivar improvement and increased use of synthetic fertilizers, herbicides,
20 and irrigation were primarily responsible for these changes. In a number of settings the effects of past changes in
21 weather or CO₂ have been regarded as noise that obscures other effects of interest (Bell and Fischer, 1994). Due to
22 the large number and relative importance of non-climate drivers in food systems and food security, formal detection
23 and attribution of impacts is extremely difficult for this sector. The majority of confounders, such as fertilizer
24 application or adoption of modern crop varieties, are not well measured in terms of their distribution across space
25 and time. Further, it is difficult to quantify the exact relationship between these confounders and outcomes of
26 interest (e.g., crop yields). The identification of a unique fingerprint from greenhouse gas emissions in these systems
27 is therefore impractical. There are no known studies simulating historical trends in food-related outcomes with and
28 without changes in anthropogenic emissions of greenhouse gases. A possible exception is the study by Auffhammer
29 et al. (2006) who compared predicted rice yields in India using climate model simulations of temperature in the late
30 20th century with yields estimated from observed temperatures for 1930-1960, using the latter period as a surrogate
31 for climate without recent changes in greenhouse gases. They find that rice yields in a world without greenhouse gas
32 emissions would have been significantly higher, thus attributing negative impacts to emissions.
33

35 **18.4.1.1. Agricultural Crops**

36

37 In order to make attribution statements regarding crop changes to climate change, one needs to make assumptions
38 about how farmers adapt. Some studies assume that farmers do not change their practices or technology as a
39 response to changes in climate during the period of study. This may be a valid assumption in some cases (Schlenker
40 and Roberts, 2009) yet there is evidence of significant adaptation to climate via technology for a variety of crops and
41 locations (Zhang *et al.*, 2008b; Liu *et al.*, 2010).
42

43 A significant number of studies have provided impact estimates of observed changes in climate on cropping systems
44 over the past few decades (see chapter 7.2). Based on this literature, there is *medium confidence* that over the past
45 several decades observed climate trends have adversely affected wheat and maize production for many regions.
46 There is *medium confidence* that observed climate trends since 1980 have had adverse impacts on global total
47 production of these crops (Figure 7-2). There is *medium confidence* that climate change impacts on rice and soybean
48 yields over this time period have been small in major production regions and globally (Figure 7-2). There is *high*
49 *confidence* that in some cold regions warming has benefitted crop production in recent decades (Jaggard *et al.*, 2007;
50 Chen *et al.*, 2011).
51

52 Many crop modeling studies focus on production for single sites or provinces and/or short time-series, rendering
53 attribution of observed yields to climate change problematic. Some recent studies, however, examine outcomes at
54 the continental or global scale (Lobell and Field, 2007; You *et al.*, 2009; Lobell *et al.*, 2011). At this scale, observed

1 trends in some climatic variables, including mean summer temperatures, can be attributed to anthropogenic activity
2 (e.g., Jones *et al.*, 2008). These studies indicate that this observed warming has had significant negative impacts on
3 trends in crop yields for certain crops. Figure 7-3 presents a summary of the detectability of changes in growing
4 season climate and crop yield changes, as well as the ability to attribute changes to climate and CO₂ trends (in the
5 case of yield changes) or anthropogenic emissions (in the case of growing season climate changes).
6

7 The recent literature has increasingly documented attributable trends not only in the seasonal averages of climate
8 variables, but also for extremes. Extreme rainfall events are widely recognized as important to cropping systems
9 (Rosenzweig *et al.*, 2002). Changes in the patterns of rainfall extremes have been attributed to anthropogenic
10 activity (Min *et al.*, 2011). A similar observation has been made for frost patterns in nearly every region of the world
11 (Alexander *et al.*, 2006; Zwiers *et al.*, 2011), as well as for the occurrence of very hot nights (WGI AR5 Chapter
12 10.6.1). High nighttime temperatures are harmful to most crops, but this effect has been observed most frequently
13 for rice yield (Peng *et al.*, 2004; Wassmann *et al.*, 2009; Welch *et al.*, 2010) and quality (Okada *et al.*, 2009).
14 Daytime extreme heat is also damaging and sometimes lethal to crops (Porter and Gawith, 1999; Schlenker and
15 Roberts, 2009). At the global scale, trends in annual maximum daytime temperatures have been attributed to
16 greenhouse gas emissions (Zwiers *et al.*, 2011).
17

18 Further, changing atmospheric conditions are affecting crops both positively and negatively. It is *virtually certain*
19 that the dramatic increase in atmospheric CO₂ concentrations since preindustrial times has improved water use
20 efficiency and yields most notably in C₃ crops. It is important to note that these effects are of relatively minor
21 importance when explaining total yield trends (Amthor, 2001; Long *et al.*, 2006; McGrath and Lobell, 2011).
22

23 Emissions of CO₂ have been associated with ozone (O₃) precursors (Morgan *et al.*, 2006; Mills *et al.*, 2007). There is
24 *high confidence* that elevated O₃ currently suppresses global output of major crops, with reductions estimated at
25 roughly 10% for wheat and soy and 3-5% for maize and rice (Van Dingenen *et al.*, 2009). Detected impacts are most
26 significant for India and China, but can also be found for soybean production in the United States in recent decades
27 (Fishman *et al.*, 2010).
28
29

30 18.4.1.2. Fisheries

31

32 There is a large literature focusing on the relationship between the dynamics of marine fish stocks and climate
33 variability, suggesting that climate change has impacts on these stocks and on the fisheries that exploit them
34 (Hollowed *et al.*, 2001; Roessig *et al.*, 2004; Shriver *et al.*, 2006; Brander, 2007). Some fisheries and aquaculture do
35 not show evidence of climate change impacts (e.g. aquaculture in the UK and Ireland, Callaway *et al.*, 2012), while
36 many others do with both positive and negative changes (30.6.2.1; Figure 30-15B, see also 18.3.4).
37

38 There is *high confidence* in the detection and attribution of shifts in the spatial distributions of marine fishes (Perry
39 *et al.*, 2005, 30.6.2.1) and in the timing of events like spawning and migration (Sydeman and Bograd, 2009, 30.4).
40 The challenges produced by ocean warming and acidification vary from region to region, however, with decreases in
41 many regions and increases (probably short-term) in some regions, especially at high latitudes (30.6.2.1; Figure 30-
42 15B). The ability to attribute detected changes to climate change is confounded by the influence of other factors
43 such as harvesting, habitat modification, technological development, pollution, and interannual to decadal climate
44 variability (Brander, 2010).
45

46 Ecosystems such as kelp forests, mangroves and coral reefs provide habitat to fisheries which provide food and
47 income to hundreds of millions of people worldwide. The strong linkage between increasing sea temperatures, the
48 decline of carbonate reef frameworks (see Box 18-3), and habitat for key fisheries resources underpins a strong
49 traceable account from global climate change to impacts on coastal fisheries (*high confidence*; see Box CC-CR,
50 Figure 30-12, 30.5.2 - 30.5.4, 30.5.6).
51

52 Similar linkages between climate change and fisheries can be made for pelagic fisheries such as tuna. Shifts in tuna
53 fisheries have occurred in the Indian and Pacific oceans, driven by climate variability (ENSO). The detected changes

1 are consistent with the thermal biology of these important species and projections of change under further changes to
2 ocean temperature (*high confidence*, 30.5.1.1.1, 30.5.6.1, 30.6.2.1).

5 *18.4.1.3. Food Security*

7 Food security depends crucially on the production and storage of food. The evidence on observed climate change
8 impacts on food production therefore has implications for food security. Further, the term food security needs to be
9 discussed at different spatial scales, as there is enough food available per capita globally, yet a significant share of
10 humankind is affected by permanent or periodic food shortages. Quantifying the impact of climate change on food
11 security is extremely challenging at any scale as there are significant numbers of non-climate confounders which
12 affect food security at various spatial scales (7.1).

14 One measurable indicator of food security is the global price of food. Food prices have dropped slowly throughout
15 the last century, yet even since AR4 there have been several periods of food price increase and periodic spikes in
16 food prices (figure 7-4). Increased demand for crops recently has been partially driven by biofuel productions, which
17 is both mandated by policy and market driven by oil price spikes (Roberts and Schlenker, 2010; Mueller *et al.*, 2011;
18 Wright, 2011). There is some evidence that supply side fluctuations also impacted food prices (figure 7.4). While
19 not all significant weather events result in food price spikes, high price episodes are more probable during periods
20 with low stored stocks. This can be due to more slowly growing supply relative to demand. Government
21 interventions, such as export bans, can exacerbate price responses to weather events (Dawe, 2008). There is some
22 recent evidence that climate trends have had some influence on global supply. Lobell *et al.* (2011) estimated a price
23 increase of 19% due to the impacts of temperature and precipitation trends on supply, or an increase of 6% once the
24 beneficial yield effects of increased CO₂ over the study period were considered.

27 *18.4.2. Cities and Urbanization*

29 The world continues to urbanize rapidly. Both new and existing urban systems are evolving due to rapidly changing
30 incomes, institutions and population. There is robust evidence across a set of case studies that climate has changed
31 in many urban areas and that variability has increased, consistent with climate change projections. The most robust
32 evidence has emerged from observational data for mean annual temperature and precipitation rates, days of extreme
33 temperature, number of extreme rainfall events, and rate of sea level rise (8.2.2.1, 8.3). These shifts are associated
34 with an increased probability of flooding, droughts, inland flooding, coastal flooding, storm surges, heat waves, and
35 declines in the number of extreme cold days (see Hunt and Watkiss, 2011; Romero-Lankao and Dodman, 2011;
36 Rosenzweig *et al.*, 2011 for recent reviews).

38 Detection and attribution of climate change impacts in cities is extremely difficult due to the large role of other
39 confounding factors. Opportunities to discern climate change signals in cities are complicated by the pattern and
40 pace of urbanization which have consequences for local environmental conditions such as intensification of urban
41 heat islands, land subsidence associated with groundwater withdrawal, and heightened flooding probability resulting
42 from increase of impervious surfaces. These conditions interact with ongoing climate change and as a result make it
43 difficult to provide evidence and attribution agreement of climate change signals in cities given the current state of
44 observed data and models.

47 *18.4.3. Economic Impacts, Key Economic Sectors and Services*

49 *18.4.3.1. Economic Growth*

51 A negative cross sectional correlation between per capita income and temperature has been observed both across
52 countries (Nordhaus, 2006) and across regions within countries (Dell *et al.*, 2009a; Dell *et al.*, 2009b). Such
53 correlations should never be interpreted as causal, because the underlying mechanisms are usually complex. In low
54 income countries, careful tracking of incomes and temperatures over an extended period, taking into account

1 important confounders, shows that higher annual temperatures as well as higher temperatures averaged over 15 year
2 periods result in substantially lower economic growth (Dell *et al.*, 2012). This effect is not limited to the level of per
3 capita income, but also to its rate of growth.

4
5 Broadly, a 1 degree Celsius increase in annual average temperature has been found to lower economic growth in the
6 same year by 1.3%, which is both statistically and economically significant (Dell *et al.*, 2012). Based on 15 year
7 averages of weather, which is a measure of climate, the impacts become larger (1.9% for low income countries). The
8 same relationships do not hold for high income countries. Generally, higher temperatures affect economic growth
9 through impacts on the agricultural and industrial sectors (Dell *et al.*, 2012). One proposed mechanism for this is the
10 impact of heat stress on workers in the workplace (Dash and Kjellstrom, 2011). Temperature shocks negatively
11 affect the growth of developing countries' exports, for which 1 degree Celsius of warming in a given year reduces
12 the growth rate of its exports by 2.0-5.7% (Jones and Olken, 2010). The export sectors most affected are agricultural
13 and light manufacturing exports. There is no detectable effect for higher income countries (Jones and Olken, 2010).

14 15 16 *18.4.3.2. Energy Systems*

17
18 Energy production and consumption is growing rapidly globally, with much of the growth taking place in low
19 income and emerging economies. Due to the large number of non-climate confounders, the literature on detection
20 and attribution for the energy sector is sparse. There is a significant literature identifying the climate sensitivity of
21 various parts of the energy sector. Higher temperatures have been shown to raise the demand for cooling and lower
22 the demand for heating. Cooling demand is largest in the summer and it has been shown for some areas that peak
23 loads during the summer months have increased and that this peak is highly correlated with summer maximum
24 temperatures (Franco and Sanstad, 2008). The literature showing the opposing effects of warmer winters and
25 summers on electricity and gas demand using statistical methods have confirmed this U-shaped relationship of
26 energy and electricity demand in temperature for the United States and elsewhere (Isaac and van Vuuren, 2009;
27 Akpinar-Ferrand and Singh, 2010; Deschênes and Greenstone, 2011).

28
29 Production losses from thermal power plants increase when temperatures exceed standard design criteria (e.g.,
30 Erdem and Sevilgen, 2006), as would be expected to occur more frequently under climate change. Power generation
31 facilities may also experience performance losses and other impacts related to changes in access to and temperature
32 of cooling water, as well as sea level rise and extreme weather events (Durmazyan and Sogut, 2006; CCSP, 2007;
33 Kopytko and Perkins, 2011, see also 10.2). Further, solar photovoltaic cells become less efficient during hot days
34 (Skoplaki and Palyvos, 2009).

35
36 The impacts of higher temperatures and extreme weather events on energy delivery, transmission and distribution
37 vary across different empirical studies, facility characteristics, geographic regions, and other factors. Barges and
38 ocean vessels that transport energy resources have been shown to be particularly vulnerable to hurricanes, storms,
39 and flooding; pipeline performance can be affected by increasing ambient and soil temperatures, as well as extreme
40 events (CCSP, 2007, see also 10.2). Some studies have quantified the general relationship between temperature and
41 electricity transmission and distribution infrastructure, finding that increased temperatures can accelerate the aging
42 of transformer insulation, lead to efficiency losses, and create power system reliability issues (Swift *et al.*, 2001; Li
43 *et al.*, 2005; Askari *et al.*, 2009).

44 45 46 *18.4.3.3. Tourism*

47
48 Tourism is a climate sensitive economic sector and ample research has been performed to understand its sensitivity
49 to climate change and impacts of (future) climate change on tourism, yet little to no literature has focused on
50 detection and attribution of observed impacts (cf. Scott *et al.*, 2008, see also 10.6). A comparatively well-studied
51 area is the climate sensitivity of wintersports in lower lying areas. For example, the increase in investment in
52 artificial snow machines in the European Alps can be attributed with *high confidence* to a general decrease of snow
53 depth, snow cover duration and snow fall days since the end of the 1980's for low-elevation mountain stations
54 (Durand *et al.*, 2009; Valt and Cianfarra, 2010; Voigt *et al.*, 2011), which in turn has been attributed to anomalous

1 warm winter temperatures over the past 20 years (Marty, 2008). Increased variability in precipitation, shrinking
2 glaciers and milder winters have been shown to negatively affect visitor numbers in winter sports areas in Europe
3 and North America (Becken and Hay, 2007). Eijgelaar *et al.* (2010) argue that so-called “last chance tourism” is a
4 strong pull for tourists to visit Antarctica to admire the glaciers while they still can. Farbotko (2010) uses a similar
5 mechanism to explain the rise in popularity of Tuvalu as a destination choice. In contrast, Zeppel (2012) states a low
6 level of concern for coral bleaching by tourists visiting the Great Barrier Reef.

9 **18.4.4. Impacts of Extreme Weather Events**

10
11 The impacts of extreme weather events depend on the frequency and intensity of the events, as well exposure and
12 vulnerability. Climate change is expected to affect both the frequency and intensity of extreme weather events
13 including extreme temperature, droughts, heavy rainfall, and tropical and extratropical cyclones (IPCC, 2012). There
14 is *low to high* confidence that such changes have already occurred, depending on the type of extreme (WGI AR5
15 Chapter 10.6). However, the impacts of extreme weather events also depend on the vulnerability and exposure of
16 systems. It is possible that climate change can affect vulnerability and exposure, but both can also be affected by
17 non-climate confounders, most notably economic development.

20 **18.4.4.1. Economic Losses due to Extreme Weather Events**

21
22 Extreme weather events can have economic impacts arising from damage to private and public assets as well as the
23 temporary disruption of economic activities. Although immediate economic impacts are most noticeable, long-term
24 impacts are also possible, as are impacts beyond the area directly affected by the event. Some economic impacts are
25 not readily monetizable and are thus excluded from most economic assessments (Handmer *et al.*, 2012, their chapter
26 4.5.1, 4.5.3).

27
28 There is *high confidence* that economic costs of extreme weather events have increased over the period 1960-2000,
29 with insured losses increasing more rapidly than overall losses (Handmer *et al.*, 2012, their chapter 4.5.3.3, 4.5.4.1).
30 This is also reflected by an increase in the frequency of extreme weather-related disasters over the same period
31 (Neumayer and Barthel, 2011). Recent studies from Latin America (Mexico, Colombia, Peru) highlight both
32 variability and positive trends in disaster frequency, (unadjusted) losses and other damage metrics (Saldaña-Zorrilla
33 and Sandberg, 2009; Marulanda *et al.*, 2010; Rodriguez-Oreggia *et al.*, 2012; Huggel *et al.*). However, there is *high*
34 *confidence* that the greatest contributor to increased cost is rising exposure associated with population growth and
35 growing value of assets (Bouwer *et al.*, 2007; Bouwer, 2011; Barthel and Neumayer, 2012; Handmer *et al.*, 2012,
36 their Chapter 4.2.2, Box 4.2 and 4.5.3.3).

37
38 To account for changes over time in the value of exposed assets, many studies attempt to normalize monetary losses
39 by an overall measure of changes in asset value. Although there is considerable year-to-year variability associated
40 with the occurrence of extreme weather events, most studies have found no detectable trend in normalized losses
41 consistent with anthropogenic climate change. Studies of normalized insured losses, that in general meet higher data
42 quality standards than data on overall losses due to thoroughly monitored payouts, have focused on developed
43 countries including Australia, Germany, Spain, the US, Spain (Changnon, 2007; Changnon, 2008; Changnon,
44 2009a; Changnon, 2009b; Barredo *et al.*, 2012; Barthel and Neumayer, 2012, see also 10.7.3; Sander *et al.*, 2013),
45 Studies of normalized losses from extreme winds associated with hurricanes in the US (Miller *et al.*, 2008; Pielke Jr
46 *et al.*, 2008; Schmidt *et al.*, 2010; Bouwer and Botzen, 2011) and the Caribbean (Pielke Jr *et al.*, 2003), tornadoes in
47 the US (Brooks and Doswell, 2002; Boruff *et al.*, 2003; Simmons *et al.*, 2012) and wind storms in Europe (Barredo,
48 2010) have failed to detect trends consistent with anthropogenic climate change, although some studies were able to
49 find signals in loss records related to climate variability, such as century-scale damage and loss of life due to
50 wildfires in Australia related to El Niño Southern Oscillation and Indian Ocean dipole phenomena (Crompton *et al.*,
51 2010), or decadal-scale typhoon loss variability in the Western North Pacific (Welker and Faust, 2013).

52
53 Although there are a number of data issues that may limit the reliability of assessments of the economic impacts of
54 extreme weather events (Crompton and McAneney, 2008; Bouwer and Botzen, 2011; Nicholls, 2011; Handmer *et*

1 *al.*, 2012, their Chapter 4.5.3.3, see also 10.7.3), there is *limited evidence* of a trend in the economic impacts of
2 extreme weather events that is consistent with a change driven by observed anthropogenic climate change.

3 4 5 18.4.4.2. *Detection and Attribution of the Impacts of Single Extreme Weather Events to Climate Change* 6

7 Although most studies of the relationship between climate change and extreme weather events have focused on
8 changes over time in their frequency and intensity, a few studies have focused on the contribution of climate change
9 to specific events (WGI AR5 Chapter 10.6.2). Assessing the contribution of climate change to a specific event poses
10 particular challenges, both in terms of methodology and communication of results (Allen, 2011; Curry, 2011; Hulme
11 *et al.*, 2011; Trenberth, 2011). If climate change was among the drivers for a specific extreme weather event, there
12 remains the question how this contribution translates into associated impacts and damages. Only a few studies have
13 attempted to evaluate the role of climate change in the impacts of individual extreme weather events. Pall *et al.*
14 (2011) and Kay *et al.* (2011), using observational constraints on climate and hydrologic model simulations,
15 concluded that greenhouse gas emissions have increased the probability of occurrence of a comparable flooding
16 event in autumn 2000 over the UK. A similar study for recent high flood years in the Okavango Delta, Botswana,
17 found a decreased probability of high floods (Wolski *et al.*, 2013). The autumn 2000 UK studies did not evaluate the
18 relative role of other factors, however, while the Okavango study argued that the effects of land use, land cover, and
19 water management change in the Okavango Basin were negligible.

20
21 In highly temperature-sensitive regions such as high mountains there is *high confidence* that several extreme impact
22 events of the 20th and 21st century can be qualitatively attributed to effects of climate change, namely glacier lake
23 outburst floods due to glacier recession and subsequent formation of unstable lakes (Evans and Clague, 1994; Carey,
24 2005; Bajracharya and Mool, 2009), debris flows from recently deglaciated areas, and rock fall and avalanches
25 following the loss of mechanical support accompanying glacier retreat (Haeberli and Beniston, 1998; Oppikofer *et al.*
26 *et al.*, 2008; Huggel *et al.*, 2012b; Stoffel and Huggel, 2012, see also 18.3.1.3). Similar multiple-step approaches can
27 be used to evaluate the contributions of anthropogenic emissions to recent damaging extreme events (see Table 18-
28 4).

29
30 Irrespective of whether a specific event can be attributed in part to climate change, there is ample evidence of the
31 severity of related impacts on people and various assets. Both low- and high-income countries have been strongly
32 impacted by extreme weather events in recent years, but the impacts relative to economic strength have been higher
33 in low-income countries (Handmer *et al.*, 2012). Similarly, at the national scale, poor or elderly people have been
34 disproportionately affected, as documented for Hurricane Katrina in the US in 2005 (Elliott and Pais, 2006; Bullard
35 and Wright, 2010) or the 2003 European heat wave (Fouillet *et al.*, 2008). Exacerbating effects of extreme weather
36 events are mostly of non-climatic nature, including increasing exposure and urbanization, land-use changes
37 including deforestation, or increasingly vulnerable infrastructure.

38
39 An illustrative case is the flood damage in New York City during the landfall of Hurricane Sandy in 2012. The
40 storm surge hazard is expected to increase with rising local sea level as a result of anthropogenic emissions,
41 notwithstanding any changes in associated meteorology (Lin *et al.*, 2012). However, increased exposure of
42 buildings, people, and infrastructure in New York City has probably contributed to a much larger increase in risk
43 over the past century (Aerts *et al.*, 2013). Against that increase, though, there has been a major investment in
44 reducing vulnerability and increasing resilience, partly in response to past extreme events (Tollefson, 2013).
45 Determining a contribution from anthropogenic climate change amongst such large competing factors, including
46 diagnosing the forces behind adaptation, remains a major challenge.

47
48 [INSERT TABLE 18-4 HERE

49 Table 18-4: Illustrative selection of some recent extreme impact events for which the role of climate has been
50 assessed in the literature. The table shows confidence assessments as to whether the associated meteorological
51 events made a substantial contribution to the impact event, as well as confidence assessments of a contribution of
52 anthropogenic emissions to the meteorological event. The assessment of confidence in the findings is not necessarily
53 a conclusion of the listed literature but rather results from assessment of the literature. Assessment of the role of

1 anthropogenic emissions in the impact event requires a multi-step evaluation. Partly based on Coumou and
2 Rahmstorf (2012).]

5 **18.4.5. Human Health**

7 IPCC AR4 (Confalonieri *et al.*, 2007) concluded that there was *weak to moderate evidence* (with *low to medium*
8 *confidence*) of climate change effects on three main categories of health exposures: vectors of human infectious
9 diseases (changes in distribution), allergenic pollen (changes in phenology), and extreme heat exposures (trend in
10 increased frequency of very hot days and heat wave events).

12 For pollen production, changes in phenology have been consistently observed in mid to high latitudes with, for
13 example, earlier onset in Finland (e.g., Yli-Panula *et al.*, 2009) and Spain (D'Amato *et al.*, 2007; García-Mozo *et al.*,
14 2010, see also 4.3). In North America, the pollen season of ragweed (*Ambrosia* spp.) has been extended by 13-27
15 days since 1995 at latitudes above 44°N (Ziska *et al.*, 2011). Allergic sensitization of humans has changed over a 25
16 year period in Italy, but the attribution to observed warming remains unclear (Ariano *et al.*, 2010).

18 The most direct potential health impact of climate change is through exposure to higher temperatures. The
19 association between very hot days and increases in mortality in temperate populations is very robust but depends on
20 factors such as geographic location, the age structure of the population, and the prevalence of air-conditioning
21 (11.2.2). Seneviratne *et al.* (2012) concluded that there is *medium to high confidence* in detection of non-zero trends
22 in increasing frequency of hot days and nights, decreasing frequency of cold days and nights, and in the duration of
23 heat waves and warm spells over temperate subcontinental regions, and that on the global scale it is *likely* that
24 anthropogenic emissions have had at least a minor role in these changes. However, the translation of this trend in
25 hazard to a trend in exposure is complicated by evidence of changes in social, environmental, and behavioral factors
26 (e.g., Carson *et al.*, 2006).

28 Overall, Confalonieri *et al.* (2007) concluded that there was a lack of evidence for observed effects in human health
29 outcomes, and this remains the case. Disease patterns change considerably over time due to changes in exposures
30 (e.g. smoking patterns), control measures (vaccination, drug resistance), and population structures (population
31 aging). Therefore, observations of trends in health outcomes provide little insight into a role of climate change. The
32 detection of a climate-change-induced trend for any health outcome requires that changes in reporting over time
33 need to be taken into account. Adaptation is an added confounding factor, as health agencies are mandated to
34 intervene, and people are liable to alter exposures, as soon as risks are identified. With regard to temperature-related
35 mortality, Christidis *et al.* (2010) examined the roles of changing exposure and health care and of anthropogenic
36 climate change in observed trends in excess mortality during cool and warm days. They concluded that
37 anthropogenic climate change had at most a minor role in trends in temperature-related mortality in England and
38 Wales over the three decades, relative to a major role for the combined trends of exposure and health care; however,
39 assessment of the existence of a minor role for anthropogenic climate change may have been sensitive to the lack of
40 interseasonal mortality relationships in their modeling setup (Rocklöv *et al.*, 2009; Ha *et al.*, 2011).

42 There is *limited evidence* regarding the role of observed warming in changes in tick-borne disease in mid to high
43 latitudes. The upsurge of tick borne encephalitis (TBE) in the 1980-90s in central and Eastern Europe has been
44 attributed to socio-economic factors (human behavior) rather than temperature (Šumilo *et al.*, 2008; Šumilo *et al.*,
45 2009). Changes in the observed incidence of TBE in central Sweden remain unexplained however (Randolph, 2010).
46 Changes in the latitudinal and altitudinal distribution of ticks in Europe are consistent with observed warming trends
47 (e.g., Gray *et al.*, 2009), but there is no evidence so far of any associated changes in the distribution of human cases
48 of tick-borne diseases. In North America, a northward expansion of the distribution of the tick vector (*Ixodes*
49 *scapularis*) was observed in the period 1996 to 2004 (Ogden *et al.*, 2010).

51 There is *limited evidence* of a change in the distribution of rodent-borne infections in the US (plague and tularaemia)
52 consistent with observed warming (Nakazawa *et al.*, 2007). Specifically, a northward shift of the southern edge of
53 the distributions of the diseases (based on human case data for period 1965-2003) was observed. There was no
54 change detected in the northern edge of the distributions though.

1
2 Malaria incidence has been monitored in the Kericho region of Kenya for over three decades, with a trend toward
3 increasing incidence in the late 1990s. A local warming trend also occurred during the end of the observation period
4 (Omumbo *et al.*, 2011). Other studies have confirmed that malaria incidence is sensitive to temperature and rainfall
5 effects in nonlinear ways, but it is a complex ecological system, with changes in vector, human and parasite
6 behavior that need to be considered. A mosquito-human model, however, has shown that predicted malaria cases
7 exhibit a strongly non-linear response to observed warming (Alonso *et al.*, 2011). A detailed review by Chaves and
8 Koenraadt (2010) concludes that there is *robust evidence* that decadal temperature changes have played at least a
9 minor role in the malarial trends of the East African highlands, arguing that it may be the combination of many factors
10 that has been crucial. However, temperature trends should nonetheless not be considered the main or sole cause of
11 such changes in malaria in the east African highland region, and globally the dominant trend has been a contraction
12 of the geographical range of malaria and a decrease in malaria endemicity over the past century (Gething *et al.*,
13 2010).

14 15 16 **18.4.6. Human Security**

17 18 *18.4.6.1. Violent Conflicts and Social Disruptions*

19
20 A small number of studies have examined the connection between the collapse of civilizations and large-scale
21 climate disruptions like severe or prolonged drought. DeMenocal (2001) summarized the evidence for a number of
22 cases. However, attribution to climate change can be made with only *low confidence* due to limitations on both
23 historical understanding and data. Similarly, a small number of studies have suggested that levels of warfare in
24 Europe and Asia were relatively high during the Little Ice Age (Parker, 2008; Brook, 2010; Tol and Wagner, 2010;
25 White, 2011; Zhang *et al.*, 2011), but again attribution to climate can only be made with *low confidence*. There is no
26 evidence of a climate effect on inter-state conflict in the post-WWII period.

27
28 Most of the recent research in this area has focused on the relationship between climate variability – as opposed to
29 climate change – and civil conflict, with most studies focusing on Africa. Although some studies have established
30 statistical relationships (Miguel *et al.*, 2004; Hendrix and Glaser, 2007; Burke *et al.*, 2009; Hsiang *et al.*, 2011), their
31 results have been questioned (Buhaug *et al.*, 2010; Buhaug and Theisen, 2012; Slettebak, 2012; Theisen *et al.*, 2012)
32 and no consensus has been reached. Both the detection of a climate effect on civil conflict and its attribution to
33 climate change can be made with only *low confidence*.

34
35 Very recent work has begun to examine links between climate variability and small-scale communal violence
36 (Adano *et al.*, 2012; Butler and Gates, 2012; Hendrix and Salehyan, 2012; Raleigh and Kniveton, 2012; Theisen,
37 2012), but it is too early to attach anything but *low confidence* to the detection of a climate effect and its attribution to
38 climate change.

39
40 Finally, a number of efforts have been made to establish a link between violent crime and high temperatures
41 (Anderson, 1987; Field, 1992; Anderson, 2001; Rotton and Cohn, 2001; Butke and Sheridan, 2010; Breetzke and
42 Cohn, 2012; Gamble and Hess, 2012). However, the findings remain controversial with most studies identifying
43 other factors as explaining variations in rates of violent crime (Kawachi *et al.*, 1999; Fajnzylber *et al.*, 2002;
44 Neumayer, 2003; Cole and Gramajo, 2009) and the detection of a temperature effect on violent crime and its
45 attribution to climate change can be made only with *low confidence*.

46 47 48 *18.4.6.2. Migration*

49
50 Large population movements, in response to climatic events, are sometimes considered a human security issue.
51 Empirical detection of such relationships has been slow because data sets on population movements are not yet well
52 developed. Moreover, the attribution of migration to climate change is difficult because economic, political, social,
53 demographic, and other environmental drivers interact with climatic drivers to influence migration (Black *et al.*,
54 2011). Few studies measure or empirically demonstrate how rainfall or temperature changes cause a strengthening

1 or weakening of the various forces driving migration, especially income levels and income variability (Lilleør and
2 van den Broeck, 2011).

3
4 Some large sample studies have been able to detect population movements in response to natural disasters (Smith *et al.*,
5 2006b; Boustan *et al.*, 2012) and climate-induced agricultural losses (Feng *et al.*, 2012) in the United States,
6 where data quality is high. In both the United States and African contexts, crop losses have also been associated with
7 rural to urban population movements within a country (Barrios *et al.*, 2006; Feng *et al.*, 2012). By statistical
8 attribution, Marchiori *et al.* (2012) estimate that anomalous temperature and rainfall have displaced roughly 128,000
9 people per year in Sub-Saharan Africa during 1960–2000.

10
11 Drought has prompted both short-distance (Tacoli, 2009) and long distance international migration, with the
12 Mexican drought of the 1990s providing an example of the latter (Saldaña-Zorrilla and Sandberg, 2009; Feng *et al.*,
13 2010). In Burkina Faso, temporary moves to other rural areas have increased as a result of a reduction in rainfall
14 (Henry *et al.*, 2004). Even though there is a statistically significant relationship between migration outcomes and
15 rainfall variability, Kniveton *et al.* (2011) report from own fieldwork that only 27 of 3,517 households identified
16 rainfall as a driver of migration.

17 18 19 **18.4.7. Rural Areas, Livelihoods and Poverty**

20
21 Much like for the other sectors discussed above, available research about climate related impacts on livelihood and
22 poverty distinguishes between climate sensitivity and impacts driven by climate change (e.g. gradual temperature
23 changes, and changes in climate variability). A climate sensitivity of livelihoods has been observed with *high*
24 *confidence*, while there is a paucity of evidence about impacts of gradual climate change on livelihoods and poverty.
25 Detection of changes in livelihood aspects is often difficult due to a lack of data. If trends are actually detected,
26 many confounding factors contribute to detected changes. Because of the important confounders and the current
27 limitations in attributing climate extremes and changes in variability to climate change (see 18.4.4.2), confidence in
28 attribution of changes in livelihoods and poverty to climate change is typically *low* or *very low* (Nielsen and
29 Reenberg, 2010).

30
31 Primarily negative impacts from historical climate variability and extremes, and to a much lesser degree from
32 climate change, have been observed on people's natural, physical, economic, social and cultural assets (see Table
33 18-5). Impacted natural assets include land, water, fish stocks and livestock (Osborne *et al.*, 2008; Bunce *et al.*,
34 2010). There is growing concern about negative effects of climate change and ocean acidification on marine and
35 coastal fisheries, and the livelihoods of fisherfolk (Cooley and Doney, 2009; Badjeck *et al.*, 2010), however there is
36 no literature available discussing observed impacts.

37
38 [INSERT TABLE 18-5 HERE

39 Table 18-5: Cases of regional livelihood impacts attributable with varying degree to weather- and climate related
40 events, climate change or climate variability.]

41
42 Natural and physical assets (e.g. settlements) of many poor people have been affected by weather- and climate-
43 related extreme events. Poor people living in hazard exposed areas in Africa and Latin America were affected by
44 floods and landslides in the 1990's and 2000's with some upward trend mainly due to increased urbanization (*low to*
45 *medium confidence* for increasing floods and landslides, *high confidence* for increasing number of people affected)
46 (Douglas *et al.*, 2008; Hardoy and Pandiella, 2009). Although there is evidence of a decline in average precipitation
47 in West Africa since 1960 (Lacombe *et al.*, 2012) including repeated droughts (Dietz *et al.*, 2004; Armah *et al.*,
48 2011), which in some cases has been partly attributed to anthropogenic climate forcing (Jenkins *et al.*, 2005; Biasutti
49 and Giannini, 2006), there is *limited evidence* of changes in poverty among affected small-holder and subsistence
50 farmers that can be attributed to climate drivers such as rainfall decline and droughts.

51
52 In rural areas correlations have been found between climatic variability and extreme events and livelihood
53 diversification in West Africa (Nielsen and Reenberg, 2010) and Mexico (Eakin, 2006) and in Mexico frequent

1 impacts of natural disasters correlate with substantial income drops, marginalized, grain-intensive farming and low
2 access to credit (Saldaña-Zorrilla and Sandberg, 2009).

3
4 Climate impacts disproportionately affect poor populations, thus increasing social and economic inequalities, both in
5 urban and rural areas, and in low-, middle- and high-income countries. Climatic factors thereby interact with aspects
6 of race, class, gender, ethnicity or age (Nightingale, 2011). Evidence for poor people in high income nations being
7 disproportionately affected by weather- and climate related extreme events comes, for instance, from 2005 US
8 Hurricane Katrina (Elliott and Pais, 2006; Bullard and Wright, 2010) or drought events in Australia (Alston, 2011).
9 Glacial lake outburst floods in the Peruvian Andes also affect different populations depending on their degree of
10 exposure, level of vulnerability, race, ethnicity, and socio-economic class (Carey, 2010; Carey *et al.*, 2012b). There
11 has been an observed gender bias in impacts from climate variability and extremes, due to gender specific roles
12 within the household, communities, and wider socio-political and institutional networks (Carr, 2008; Arora-Jonsson,
13 2011, see also Box 13-1).

14
15 For indigenous peoples, specific rights, including the right to life, adequate food, water, health, adequate housing,
16 and the right to self-determination, are directly implicated by the impacts of climate change and variability (Ford,
17 2009b, see also Box 18-5). Livelihoods of indigenous people in the Arctic have been identified as among the most
18 severely affected by both climate change and variability, including food security aspects, traditional travel and
19 hunting, and cultural values and references (Beaumier and Ford, 2010; Pearce *et al.*, 2010). Negative effects on
20 social and cultural assets by climatic and non-climatic stressors have also been identified in Africa with respect to
21 social networks of the poorest, elderly and female-headed households (Osbahr *et al.*, 2008). Impacts of rising
22 temperatures and increased variability of weather extremes on crops and livestock of indigenous people in highlands
23 were furthermore reported from Tibet (Byg and Salick, 2009) and the Andes of Bolivia (McDowell and Hess, 2012).
24 Shifts from transient to chronic poverty due to climate change and variability are suggested for livelihoods and
25 households that, unlike more affluent ones, lack appropriate response to, and ability to cope with altered
26 seasonalities, unpredictable seasons, and extreme events such as floods and droughts (Hardoy and Pandiella, 2009).

27
28 _____ START BOX 18-5 HERE _____

29 30 **Box 18-5. Detection, Attribution, and Traditional Ecological Knowledge (TEK)**

31
32 Indigenous and local peoples often possess detailed knowledge of climate change that is derived from observations
33 of environmental conditions over many generations. Consequently, there is increasing interest in merging this
34 traditional ecological knowledge (TEK)—also referred to as indigenous ecological knowledge (IEK) or simply
35 indigenous knowledge (IK)—with the natural and social sciences in order to better understand and detect climate
36 change impacts (Huntington *et al.*, 2004; Parry *et al.*, 2007; Salick and Ross, 2009; Green and Raygorodetsky, 2010;
37 Ford *et al.*, 2011; Diemberger *et al.*, 2012). TEK, however, does not simply augment the sciences, but rather stands
38 on its own as a valued knowledge system that can, together with or independently of the natural sciences, produce
39 useful knowledge for climate change detection or adaptation (Agrawal, 1995; Cruikshank, 2001; Hulme, 2008;
40 Berkes, 2009; Byg and Salick, 2009; Maclean and Cullen, 2009; Wohling, 2009; Ford *et al.*, 2011; Herman-Mercer
41 *et al.*, 2011).

42
43 Cases in which TEK and scientific studies both detect the same phenomenon offer a higher level of confidence
44 about climate change impacts and environmental change and the value of indigenous knowledge (Huntington *et al.*,
45 2004; Laidler, 2006; Krupnik and Ray, 2007; Salick and Ross, 2009; Gamble *et al.*, 2010; Green and
46 Raygorodetsky, 2010; Alexander *et al.*, 2011; Cullen-Unsworth *et al.*, 2011). For example, in Peru's Cordillera
47 Blanca mountains, local residents and instrument-based scientific analysis both report increasingly rapid glacial
48 recession, less snow in the upper watershed, decreased water supplies in glacier-fed basins, and an increase of
49 falling glacier “blocks” since the latter half of the 20th century (Bury *et al.*, 2010; Carey, 2010; Baraer *et al.*, 2012;
50 Carey *et al.*, 2012b). For another, in Tibet, many, but certainly not all, local residents observed warming
51 temperatures, less snow, and shrinking glaciers, which are consistent with scientific interpretations (Byg and Salick,
52 2009). At Clyde River, Nunavut, Canada, Inuit and scientific observations have detected that wind speed has
53 increased in recent years and that wind direction changes more often over short periods (within a day) than it did
54 during past decades (Gearheard *et al.*, 2010). Finally, in the Canadian Arctic, Inuit sea ice experts and scientists have

1 both observed the thinning of multiyear sea ice, the shortening of the sea ice season, and the declining extent of sea
2 ice cover, with Inuit observers reporting less predictability in the sea ice and more hazardous travel and hunting at
3 ice edges (Nichols *et al.*, 2004; Laidler, 2006; Krupnik and Ray, 2007; Ford *et al.*, 2009; Aporta *et al.*, 2011).

4
5 TEK can also inspire scientists to study new issues in the detection of climate change impacts. In one case,
6 experienced Inuit weather forecasters in Baker Lake, Nunavut, Canada, reported that it had become increasingly
7 difficult for them to predict weather, suggesting an increase of weather variability and anomalies in recent years. To
8 test Inuit observations, scientists analyzing hourly temperature data over a 50 year period confirmed that afternoon
9 temperatures fluctuated much more during springtime during the last 20 years—precisely when Inuit forecasters
10 noted unpredictability—than they had during the previous 30 years (Weatherhead *et al.*, 2010).

11
12 Despite frequent confluence between TEK and scientific observations, there are sometimes discrepancies between
13 them. These discrepancies indicate uncertainty in the identification of climate change impacts. Attribution of
14 impacts to anthropogenic climate change, for example, tends to have much less convergence between TEK and the
15 natural sciences. While community members in Canada’s Northwest Territories report that less ice cracking during
16 the last decade was a result of winter warming caused by climate change, scientists have concluded that the
17 relationship between ice cracking and air temperature is much more complex and requires more research on water
18 temperature, ice thickness, snow cover, and ice properties in order to attribute reduced ice cracking to global climate
19 change (Woo *et al.*, 2007).

20
21 Scale is another problem in the detection of climate change: TEK and scientific studies frequently focus on different
22 and distinct scales that make comparison difficult. Local knowledge may fail to detect regional environmental
23 changes while scientific regional or global scale analyses may miss local variation (Wohling, 2009; Gamble *et al.*,
24 2010). In some cases TEK and scientific studies measure or note distinct phenomenon that cannot be compared or
25 have inaccuracies (Gearheard *et al.*, 2010). Furthermore, TEK based observations and related interpretations
26 necessarily need to be viewed within the context of the respective cultural, social, and political backgrounds
27 (Agrawal, 1995). Therefore, a direct translation of TEK into a western science perspective is often not feasible.

28
29 _____ END BOX 18-5 HERE _____
30
31

32 **18.5. Detection and Attribution of Observed Climate Impacts across Regions**

33
34 Since the AR4, significant new knowledge about detected impacts of recent climate change has been gained from all
35 continents of the world, assessed in the regional chapters 22-29 of this report (for relevant information on Ocean
36 regions from Chapter 30, see 18.3.2). The following sections provide a short overview about key findings for each
37 of the eight world regions discussed in Chapters 22-29, along with a summary of relevant recent observed climate
38 change. Further details of regional detection and attribution findings are summarized in 18.5.9 for various
39 collections of natural, managed, and human systems.

40 41 42 **18.5.1. Africa**

43
44 For much of Africa, knowledge about recent climate change is limited, due to weak climate monitoring, and gaps in
45 coverage continue to exist. On the other hand, the low natural temperature variability over the continent allows
46 earlier detection of warming signals. Thus there is *medium to high confidence* in regional warming, with *low to high*
47 *confidence* in attribution to anthropogenic emissions. A main regional feature has been the drying of the Sahel
48 during the decades following 1970, but that trend has halted during the most recent decade (see also 22.2).

49
50 African environmental systems present many strong challenges for the potential detection and attribution of
51 responses to climate change. Given the weak spatial and temporal variations in temperature, there is smaller scope
52 for migrational and phenological responses to anthropogenic climate change than in other parts of the world.
53 Furthermore, high quality monitoring is relatively sparse in time and space, and is often unsuitable for detecting
54 changes across margins and borders where responses to climate change are most expected. The dearth of studies

1 examining attribution questions means it is currently difficult to estimate the degree to which studies are selectively
2 published based on results, and thus to determine whether each attribution study is only indicative of local reasons
3 for concern or if it is more generally representative of a broader domain.

4
5 Since the AR4 there has been a particular research focus on three geographic domains: the effects of dryness in the
6 Sahel since 1970 on tree density; the effects of surface warming and resulting increased stratification on the Great
7 Lakes on the lake ecology; and the effect of warming on species ranges in southern Africa, where spatial
8 temperature gradients are larger and there is more scope for range shifts as a measureable response.

11 **18.5.2. Europe**

12
13 Amongst all continents, Europe has the longest tradition in climate monitoring. Warming has been occurring across
14 the continent in all seasons, with an associated decreasing frequency of cold extremes and increasing frequency of
15 hot extremes (Seneviratne *et al.*, 2012). The Southern parts of the continent (the Mediterranean) have been getting
16 drier, while northern areas have been getting wetter (23.2.2.1), with a general increase in the frequency of extreme
17 wet events everywhere (Seneviratne *et al.*, 2012).

18
19 Significant warming-related impacts have long been observed in Northern and Central Europe, with substantial loss
20 of Alpine glaciers, longer growing seasons, associated productivity increases and changed phenology of many plant
21 and animal species on land as well as in the Atlantic Ocean. Higher temperatures and generally drier conditions have
22 likely contributed to the stagnation of agricultural yields that would otherwise have been expected to increase as a
23 consequence of improved agricultural practices and higher atmospheric CO₂. In the Mediterranean drier and hotter
24 conditions have contributed to increased wildfires.

27 **18.5.3. Asia**

28
29 Asia spans a wide range of climate types and expected climate changes, but warming has been observed throughout
30 the continent with northern areas amongst the fastest warming on the planet. Precipitation trends vary
31 geographically, with a more frequent but weaker Indian monsoon (WGI AR5 Chapter 14.7.10), and contrasting
32 increasing and drying trends over coastal and inland China (24.3.1).

33
34 Among the most wide-spread observed impacts of recent climate change in Asia are the degradation of permafrost
35 occurring throughout its current distribution in Siberia, Central Asia and on the Tibetan Plateau, matching
36 observations from elsewhere in the world and to a very high degree explained by warming. Substantial new
37 evidence has been collected since AR4 on glaciers in Asia. Across most of Asia glaciers have been shrinking, except
38 from some areas in the Karakorum and Pamir. In some rivers (e.g. in China) an increase in runoff was observed in
39 glaciated catchments.

40
41 Plants and animals are changing their phenology and growth in many parts of Asia, largely due to climate change,
42 many species are also shifting their distribution northwards or upwards in elevation. In the oceans of tropical Asia
43 and around Japan, coral reefs and large seaweeds are in decline.

46 **18.5.4. Australasia**

47
48 There is *very high confidence* of warming over Australian and New Zealand during the past century, and *high*
49 *confidence* in hot extremes becoming more frequent and cold extremes becoming less frequent (25.2, Table 25-1).
50 Winters in southern areas of Australia have become drier in the past few decades and the northwest has become
51 wetter, and precipitation increased over the south and west of both islands of New Zealand (*high confidence*). While
52 there have been no significant trends in drought frequency over Australia, regional warming may have increased
53 their hydrological intensity (*low confidence*, Table 25-1).

1 There is *high confidence* of a significant decline in late season snow depth several sites in Australia's Snowy
2 Mountains, and a decline of glacier volume by 25% in New Zealand since the middle of the 20th century (table 25-
3 1). In many Australian terrestrial ecosystems, some observed changes in the distributions, genetics, phenology of
4 individual species, and in the structure and composition of some ecological communities can be attributed to recent
5 climatic and atmospheric trends, but non-climatic drivers, such as changes in fire management, grazing and land-
6 use, also play a significant role. The 1997-2009 drought has affected freshwater systems in the eastern states and the
7 Murray Darling Basin, but for many freshwater systems, direct climate impacts are difficult to detect. In New
8 Zealand, few changes in ecosystems have been attributed to climate change.

9
10 Climate change impacts have been profound in marine systems around Australia and New Zealand, involving large
11 latitudinal shifts of climatic zones and associated biota. Many of these changes have been associated with ocean
12 warming, although for others the impacts of interacting non-climate stresses, including habitat degradation, coastal
13 pollution and fisheries are also significant. About 10% of the observed 50% decline in coral cover on the Great
14 Barrier Reef has been attributed to bleaching, the remainder to cyclones and predators.

15
16 Exceptional heatwave conditions in Australia have been associated with substantial increases in excess deaths and
17 heat-related hospital admissions (Khalaj *et al.*, 2010; Loughnan *et al.*, 2010).

18 19 20 **18.5.5. North America**

21
22 North America spans a wide range of climate types and observed climate changes. While the northwest of the
23 continent has been amongst the fastest warming on the planet, the southeast of the USA has experienced slight
24 cooling (26.2.2). Hot extremes have been becoming more frequent while cold extremes and frost days have been
25 becoming less frequent over the past several decades. Trends in precipitation over western parts of the continent are
26 strongly influenced by the variability of the El Niño/Southern Oscillation, with a matching drying and decreasing
27 snowpack. The intensity of precipitation events has been increasing over most of the continent, but trends in dryness
28 are spatially heterogeneous (26.2.2). There is *robust evidence* of an increase in intense tropical storms in the North
29 Atlantic over the past several decades (WGI AR5 Chapter 2.6.3).

30
31 There is evidence for hydrological change in many parts of North America, particularly in catchments that are
32 dominated by snowpack and ice. Terrestrial ecosystems have responded to warming with range shifts (upwards and
33 polewards), changed phenology, productivity and mortality (insect outbreaks). In some regions, wildfires have
34 become more frequent. Marine ecosystems significantly shift ranges northwards. Agricultural production is affected
35 by increased drought conditions in the Southern US and Mexico. There is some evidence for infrastructural damage
36 from more frequent climatic extremes.

37 38 39 **18.5.6. Central and South America**

40
41 Most of South America has been observed to be warming over the past half century, except for an observed cooling
42 over a western coastal strip (27.2.1.1). Precipitation over much of Central and South America is strongly influenced
43 by the El Niño/Southern Oscillation, with accompanying long-term variability. There has been in reduction in the
44 number of dry summer months in the southern half of the continent, while observed trends over the Amazon are
45 sensitive to the selection of time period (27.2.1.1).

46
47 Among the observed impacts of climate change in South and Central America there are some that stand with *high* or
48 *very high confidence* in detection and attribution. These are: the retreat of tropical glaciers and ice-fields in the
49 Andes, decreasing river flows in the western tropical Andes, increased streamflow in the La Plata basin and
50 bleaching of coral reefs in the western Caribbean. Several impacts are associated to many confounding causes and
51 although detection of change may be found with *high* levels of *confidence*, the confidence in attribution is *low* or
52 *very low*. In this category are included the degrading and receding of the Amazon rainforest, mangrove degradation
53 on the Northern coast of South America and the expansion of agricultural areas in climatically marginal regions of
54 Argentina. With *high confidence* in detection and *medium confidence* in attribution, denoting perhaps an impact

1 more sensitive to climate than in the previous case, are included the increase in agricultural yields in Southeastern
2 South America and the increase in frequency and extension of dengue fever, yellow fever and malaria.

3
4 Although the number of articles in the literature and of consolidated datasets has both increased in Central and South
5 America during the past decade, still there are many gaps to be filled in mainly in remote regions and regarding
6 historic temporal coverage.

9 **18.5.7. Polar Regions**

10
11 The areas of largest observed warming are all polar: the northwest of North America, northern Asia, and the
12 Antarctic Peninsula. While this occurs against a large background of natural variability, WGI AR5 Chapter 10
13 conclude that it is *likely* that there has been an anthropogenic contribution to Arctic warming; for the Antarctic,
14 however, there is *low confidence* in attribution due a sparse monitoring network, an apparent lack of warming in the
15 continental interior, and the contrasting effects of greenhouse gas emissions and stratospheric ozone depletion
16 (10.3.1.1.4). The nature of polar regions means that warming can lead to large changes in other aspects of the
17 climate system, in particular the observed decrease in summer sea ice cover, earlier thaw, earlier spring runoff, and
18 melting of permafrost (28.2).

19
20 Recent changes in the Arctic tundra hydrology are attributed to permafrost degradation caused by increasing air
21 temperature and reduced albedo. Swamp formation, lake drainage, and ecosystem shifts are among the observed
22 consequences. Permafrost degradation also affects traditional living for local and indigenous communities.

23
24 The ongoing reduction of sea ice cover and duration has profound impacts on the subsistence of many species in the
25 Arctic as well as the Antarctic. Polar bears are suffering from reduced vigor and reproduction, and arctic seabird
26 populations are facing longer flight distances for provisioning of their offspring due to phenological mismatch,
27 resulting in decreasing reproductive success. Antarctic penguin species show decreasing numbers, attributed to
28 shrinking stock of their main prey, the Antarctic krill, which in turn is affected by sea ice loss.

29
30 Changing snow conditions affect arctic plants and animals in various ways. Mild spells in the winter, with increasing
31 frequency of rain on snow events and the formation of ice crusts in the snow pack has affected the populations and
32 cyclicity of arctic grazers (voles, ptarmigans, reindeer and caribou) with cascading effects on their predators, e.g.,
33 the Arctic fox: Decreasing range of snow-beds in the Arctic summer have strong impacts on animal communities
34 and reindeer husbandry.

35
36 The most easily detected change in the physiognomy of the arctic tundra is the ongoing increase in shrub cover as
37 seen in many studies from the North American as well as the Eurasian Low Arctic and Subarctic, attributed to
38 increasing temperature. In the Antarctic Peninsula, the two resident species of vascular plants have passed a summer
39 temperature threshold, and the attribution of the observed sudden increase in production of mature seeds is not
40 confounded by other drivers.

43 **18.5.8. Small Islands**

44
45 Despite the widely accepted high vulnerability of many small islands to climate change, including their close
46 relationship with Coral Reefs, there are only few formal studies on observed impacts. Detection of climate change
47 impacts in small islands is challenging due to the strong presence of other anthropogenic drivers of local
48 environmental change. Attribution is further challenged by the strong influence of natural variability compared to
49 incremental changes of climate drivers, and the lack of long term monitoring, high quality data.

50
51 Observations of impacts of sea level rise on small islands are scarce. In many cases phenomena which are associated
52 with expected impacts of sea level rise such as erosion, inundation, wave overwash are also strongly linked with on-
53 going, known climate variability and extremes and cyclic processes such as ENSO, storms and extreme tidal

1 phenomena. It is therefore difficult to quantify the degree to which such impacts can be attributed to climate change
2 and sea level rise.

3
4 For example, there is evidence for higher than average sea level rise, with rates up to four times the global average
5 (approximately 12 mm yr⁻¹ between 1993 and 2009), in the tropical western Pacific region, where a large number of
6 small island and atoll communities are located (Becker *et al.*, 2012; Meyssignac and Cazenave, 2012). However,
7 these changes may be transient, caused by strong and persistent La Nina like conditions superimposed over sea-level
8 rise and are therefore currently not unambiguously attributed to climate change (Becker *et al.*, 2012).

11 **18.5.9. Impacts across Regions**

12
13 In order to allow for comparison between the various regions of the world, the following synopsis regroups the main
14 conclusions for various components of the physical, biological, and human systems. The regional synopsis is
15 divided into various systems, covering observed impacts of regional climate change for: snow, ice and mountains,
16 rivers and lakes (Table 18-6); terrestrial ecosystems, droughts and wildfires (Table 18-7); marine ecosystems and
17 coastal processes (Table 18-8); and human and managed systems such as food production, economic impacts, and
18 health (Table 18-9). These assessments follow from discussions in the various sectoral and regional chapters of this
19 report and in earlier sections of this chapter.

20
21 Broadly, there is now at least *robust evidence* for observed impacts in most of the categories of physical and
22 biological systems listed in most of the regions. There is also evidence of observed impacts on food production in
23 many of the regions, while detection of impacts in other human systems remains limited. Further interpretation of
24 these regional synopses within the “Reasons for Concern” framework forms the basis of the synthesis assessment of
25 18.6.

26
27 [INSERT TABLE 18-6 HERE

28 Table 18-6: Observed impacts of climate change on snow, ice and mountains, and rivers and lakes, across eight
29 major world regions, with confidence in detection/confidence in attribution to climate change stated for each impact.
30 References to related chapters are given as well as key references underlying the assessment.]

31
32 [INSERT TABLE 18-7 HERE

33 Table 18-7: Observed impacts of climate change on terrestrial ecosystems, and occurrence of Drought and Wildfire,
34 across eight major world regions, with confidence in detection/confidence in attribution to climate change stated for
35 each impact. References to related chapters are given as well as key references underlying the assessment.]

36
37 [INSERT TABLE 18-8 HERE

38 Table 18-8: Observed impacts of climate change on Marine and marine influenced Ecosystems, and Coastal
39 processes, across eight major world regions, with confidence in detection/confidence in attribution to climate change
40 stated for each impact. References to related chapters are given as well as key references underlying the assessment.]

41
42 [INSERT TABLE 18-9 HERE

43 Table 18-9: Observed impacts of climate change on Human Systems across eight major world regions, with
44 confidence in detection/confidence in attribution to climate change stated for each impact. References to related
45 chapters are given as well as key references underlying the assessment.]

48 **18.6. Synthesis: Detected Impacts of Climate Change and Reasons for Concern**

50 **18.6.1. Approach and History**

51
52 A key motivation for the effort in assessing observed changes is the possibility that observed impacts could
53 constitute indications of future expected changes. Observed losses in glacial volume, for example, lend additional
54 plausibility to model-based expectations that additional warming could result in additional ice loss. Due to the

1 complex nonlinear behavior of most environmental systems, it cannot always be assumed that past impacts scale
2 linearly to future impacts. Likewise, absence of past impacts cannot constitute evidence against the possibility of
3 future impacts. Nonetheless, detection and attribution of observed impacts may serve as part of the foundation for a
4 climatic risk analysis. In order to do so, the total body of observed impacts needs to undergo a synthetic analysis
5 pointing towards the conceivable risks.
6

7 The AR4 precursor of the current chapter (Rosenzweig *et al.*, 2007) organized its efforts around a geographically
8 distributed empirical analysis of correlation across numerous detailed and localized studies of changing systems as
9 described in Rosenzweig *et al.* (2008). Rather than expand that approach using the quickly growing scientific
10 literature, the goal for this synthesis is to organize findings on detection and attribution in a way that fully covers the
11 disciplinary, sectoral and geographic diversity of observed impacts. The approach aims to establish current
12 conditions concerning the risk analysis model formulated earlier by the IPCC through the establishment of a limited
13 number of “Reasons for Concern” (RFC). The RFC concept was developed in IPCC-TAR (Smith *et al.*, 2001),
14 adopted for a second time in IPCC-AR4 (IPCC, 2007a), and updated in Smith *et al.* (2009). The RFCs respond
15 directly to requests from countries that the IPCC assesses the science with respect to the United Nations Framework
16 Convention on Climate Change’s (UNFCCC) commitment to “stabilization of greenhouse gas concentrations in the
17 atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system” (UNFCCC,
18 1992), without being policy prescriptive in terms of the definition of “dangerous”. For the present synthesis, the goal
19 is not to determine the levels of warming that correspond to a particular level of impact (as done in, e.g., IPCC,
20 2007b, Figure SPM.2); instead this analysis seeks to establish, qualitatively, the importance of impacts already
21 observed in relation to the critical levels for concern indicated by the red colour of each bar portrayed in IPCC
22 (2007a), using the levels revised by Smith *et al.* (2009).
23

24 While the summary of observed impacts, taken for themselves, cannot constitute an assessment of a probability-
25 based measure of risk for dangerous anthropogenic interference, the degree to which the earlier projected impacts
26 now have occurred can be useful for policy-relevant conclusions about the risks (chapters 1, 19). The actual
27 quantification of risk for a given aspect of dangerous interference depends on the metrics chosen for the societal
28 importance of a given impact and is not part of this analysis. Instead, the degree of concern is expressed in
29 qualitative terms only, with an assessment of confidence in the assessment using standard IPCC terminology
30 (Mastrandrea *et al.*, 2010, 18.2). This is thought to directly support the development of policy-relevant conclusions
31 about the risks expressed by each RFC.
32

33 As described more fully in Chapter 1 and earlier documentation, the five RFCs are (1) “Risks to Unique and
34 Threatened Systems”, (2) “Risks of Extreme Weather Events”, (3) “Distribution of Impacts”, (4) “Aggregate
35 Impacts”, and (5) “Risks of Large-Scale Discontinuities”. Their broad definitions imply significant overlap, hence
36 some observed impacts are referred to under more than one RFC. For consistency with the earlier assessments, the
37 definitions and naming conventions in Smith *et al.* (2009) were nevertheless followed as rigorously as possible.
38
39

40 **18.6.2. Five Reasons for Concern**

41 *18.6.2.1. Risks to Unique and Threatened Systems*

42
43
44 The RFC of “Risks to Unique and Threatened Systems” is concerned with the potential for increased damage to, or
45 irreversible loss of, plant and animal species, physical systems, and human livelihoods which are known to be highly
46 sensitive to temporal and/or spatial variations in climate. Figure 18-4 displays current evidence derived from
47 detection and attribution studies concerning observed impacts to unique and threatened systems as a result of
48 observed changes in climate. Changes in the three indicated physical systems have at least *high confidence* in
49 detection and at least *medium confidence* in attribution, with regional assessments also tending to have *high*
50 *confidence*. There is at least *medium confidence* in both detection and attribution for at least one each of ecosystems,
51 physical systems, and human systems, indicating observational evidence for moderate to strong reasons for concern
52 for all three classes.
53
54

1 [INSERT FIGURE 18-4 HERE

2 Figure 18-4: Confidence in detection and attribution of observed impacts on unique and threatened systems as a
3 result of recent climate change. Assessments are for confidence in detection of a change and for confidence in
4 attribution of a major role of observed climate change. Large letters denote conclusions for the full-scale system,
5 while small letters denote conclusions for regional subsets of the systems. System changes include shrinking /
6 receding glaciers, mountain and lowland permafrost degradation, Arctic coastal permafrost degradation, increased
7 mortality and bleaching of warm water reef-building corals, increased shrub cover and permafrost decay in the
8 northern tundra, increased tree mortality in the Amazon, increased tree mortality in boreal forests, changes in Arctic
9 marine ecosystems, decline in extent of mangroves and coastal wetlands, and impacts on livelihoods of indigenous
10 Arctic peoples. See Table 18-10 for details and references.]

11
12 The systems with strongest detection and attribution evidence cover the Arctic, warm-water coral reefs, and
13 mountains. In the Arctic, there is at least *medium confidence* of impacts on glaciers, permafrost, the tundra, marine
14 ecosystems, and livelihoods of indigenous peoples, reflecting macroscopic changes across both natural and human
15 systems and across the physical and ecological subregions. Evidence for the detection and attribution of shrinkage
16 and recession of glaciers comes from all continents, while evidence for attribution of coral bleaching spans a
17 similarly broad area of the tropical oceans (see Table 18-10). These observed impacts confirm the reason for
18 concern about risks to unique and threatened systems.

19
20 [INSERT TABLE 18-10 HERE

21 Table 18-10: Reported confidence of detection of observed changes in “Unique and Threatened Systems” and
22 attribution of observed changes to observed climate change. Assessments are for global or the indicated regional
23 domain, with regional variations indicated in the final column.]

24 25 26 *18.6.2.2. Risk of Extreme Weather Events*

27
28 “This RFC tracks increases in extreme events with substantial consequences for societies and natural systems.
29 Examples include increase in the frequency, intensity, or consequences of heat waves, floods, droughts, wildfires, or
30 tropical cyclones” (Smith *et al.*, 2009). The risk of impacts by an extreme weather event is a combination of the
31 probability of an extreme weather hazard and the consequences of the realization of that hazard. Hence, a change in
32 the risk of impacts of extreme weather events could be caused by a change in the probability, intensity, or
33 sequencing of the weather event itself (which are manifestations of recent climate change), or by a change in
34 exposure, vulnerability, or the resilience of the impacted system. It follows that assessments of change in risk to
35 anthropogenic climate change are particularly influenced by new evidence concerning the dominant drivers of
36 change, whether climate change is one of those major drivers or not.

37
38 In the TAR, WGII found itself unable to detect a climatic component in the causal analysis of the increasing losses
39 from extreme weather events, but outlined instead the significant vulnerabilities that could be expected if extreme
40 weather events would increase with future climate change (IPCC, 2001, Technical Summary). In the AR4, detection
41 and attribution was reported in terms of likelihood and supported in terms of consequence, with the conclusion that
42 “It is now more likely than not that human activity has contributed to observed increases in heat waves, intense
43 precipitation events, and the intensity of tropical cyclones. There are, as well, more observations of climate change
44 impacts from extremes than in the TAR. Responses to some recent extreme climate events have also revealed higher
45 levels of vulnerability across the globe, producing significant loss of life and property damage in both developing
46 and developed countries” (IPCC, 2007b).

47
48 The current assessment confirms that, in many human systems, combined changes in exposure, vulnerability, and
49 resilience have dominated past trends in risk. For instance, while there is *high confidence* that monetary losses
50 related to extreme weather events have increased over the past 3–4 decades (ISDR, 2009, 10.7.3, 18.4.2), it is
51 generally accepted that climate change has not been a major driver of that change in risk: instead, trends in exposure,
52 wealth, and vulnerability of human systems have been the major factors for the increase in losses (IPCC, 2012,
53 10.7.3, 18.4.2). There has been considerable recent research and assessment into various components of the overall
54 chain involved in extreme weather risk. Table 18-11 summarizes some notable new evidence concerning changes in

1 hazard and damage associated with extreme weather events that has been reported in the SREX, the WGI
2 contribution to the AR5, and throughout this WGII contribution to the AR5.

3
4 [INSERT TABLES 18-11a AND 18-11b HERE

5 Table 18-11a: Confidence in detection of changes in extreme weather hazard and attribution of a major role of
6 changing greenhouse gas concentrations in the observed changes.

7 Table 18-11b: Confidence in detection of observed trends in damage in various systems and attribution of a major
8 role of observed changes in extreme weather to those observed trends.]

9
10 With regard to extreme weather hazard, temperature extremes have changed in many regions as a consequence of
11 climate change (*high confidence*), with lower confidence in weather extremes that are less directly related to
12 temperature or that occur on smaller spatial and time scales (where the connection with global mean temperature is
13 less secure) (Seneviratne *et al.*, 2012, WGI AR5 Chapter 10.6.1). Given the documented sensitivity of many systems
14 to hot temperatures, the increase in extreme heat hazard implies a significant reason for concern. Indeed, for some
15 systems there is evidence that local trends in temperature extremes have had a direct impact (Table 18-11b). There is
16 also evidence that other trends in extreme weather hazard which are closely tied to global or regional warming, such
17 as hazards related to sea level rise or sea ice retreat, have also been leading to increased damage.

18
19 Most of the evidence of attributable changes in impact risk due to weather extremes concerns corals or Arctic
20 systems (see Figure 18-5). Evidence in both cases is *robust*. Outside of these systems, however, evidence is *limited*
21 and highly localised. Recent detection and attribution literature has shed light on various components of the impact
22 risk chain, but is otherwise limited in its extent for evaluating changes in impact risk for systems other than warm-
23 water coral reefs and Arctic systems. Overall, observed impacts confirm reasons for concern about the risks of
24 extreme weather events.

25
26 [INSERT FIGURE 18-5 HERE

27 Figure 18-5: Left: confidence in detection of observed changes in extreme weather hazards and attribution of a
28 major role of changing greenhouse gas concentrations in the observed changes. Right: confidence in detection of
29 observed changes in damages related to extreme weather and attribution of major role to observed changes in
30 extreme weather.]

31 32 33 18.6.2.3. Distribution of Impacts

34
35 This RFC concerns the disparities of impacts between regions, countries, and populations. Evidence of differentiated
36 climate change impacts have been projected for all IPCC regions since the days of the TAR. However, because of
37 major geographical gaps in detection and attribution evidence the TAR did not report any observed regionally
38 differentiated impacts, and nor did the AR4. The “before present” segments of the RFC box were therefore left
39 uncolored in both the TAR and Smith *et al.* (2009). The survey of recent studies presented in 18.5 indicates that the
40 gap is being reduced. While evidence for detected impacts is still more exhaustive from Europe and North America,
41 considerable confidence in conclusions has been developed elsewhere since the AR4, particularly in Central and
42 South America and Australasia. Tables 18-6 through 18-9 summarize what we have learned from the more recent
43 evidence, as assessed in this chapter and other chapters throughout this report based on a consistent interpretation of
44 the underlying criteria. Figure 18-6 portrays the assessment visually.

45
46 In terms of level of confidence, it is no longer the case that observed evidence to support the regional RFC is
47 dominated by any particular region. The number of studies and coverage of systems may still be asymmetrically
48 distributed, but rigorous analyses of observed climate change effects and attribution (to a lesser but not insignificant
49 degree) have emerged from Central and South America, Australasia, Asia, Africa, and Small Islands. The synthesis
50 of assessments for natural systems in the middle-to-high range for detection and the middle-to-high range for
51 detection and attribution spans all major regions. The number of systems covered is only an indicative metric of
52 coverage, because many options exist for aggregation and disaggregation of evidence. Thus detection and attribution
53 analyses currently do not provide evidence of differing severity of impacts between continents. There exists,
54 however, a particular emphasis on evidence concerning the Arctic where systems exhibit a high sensitivity to

1 climate change. Given the strong levels of confidence in changes across a broad range of Arctic systems, this
2 emphasis indicates a stronger severity of impacts on the Arctic generally.

3
4 Coverage of human and managed systems is noticeably less extensive and more concentrated in the traditionally
5 well-studied regions. Notwithstanding these differences, the qualitative conclusion that observed impacts have now
6 been detected and attributed with *medium* to *high confidence* across the various IPCC-defined regions of the world is
7 new and noteworthy. These higher confidence impacts generally concern the UNFCCC's metric that "food
8 production is not threatened"; confidence in detection and attribution of observed impacts in other human systems
9 generally remains much lower.

10
11 Throughout its assessments, the IPCC has repeatedly noted that there is significant disparity between the
12 vulnerability of countries, regions, and social groups, due to differences in adaptive capacity. The AR4 cited
13 examples that indicate how impacts of extreme events differ between social groups (Wilbanks *et al.*, 2007). The
14 current coverage of detection and attribution studies is insufficient for broad evaluation of social disparities in
15 impacts. Additional research effort is required to more fully address the nature of differences in impacts on various
16 groups (18.7).

17
18 [INSERT FIGURE 18-6 HERE

19 Figure 18-6: Confidence in detection of observed changes in natural systems (panel a) and human and managed
20 systems (panel b) across regions, and confidence in attribution of such trends to observed climate change. Based on
21 assessments developed in Tables 18-6 through 18-9.]

22 23 24 *18.6.2.4. Aggregate Impacts*

25
26 The original intent of the Aggregate Impacts (or Aggregate Damages) RFC was to assess economic impacts,
27 damages, and economic risk driven by climate change at a globally aggregated level. However, while some observed
28 impacts are calibrated using a common monetary currency, others are now directly measured in a common
29 calibrated metric or are otherwise qualitatively comparable across the globe and thus amenable to aggregation. In
30 recognition of this, the scope of the RFC has been expanded over time (Smith *et al.*, 2009). Consequently, a variety
31 of globally comparable impacts are considered here.

32
33 Table 18-2 lists various aggregate systems for which: the extent is global or near-global; there is a quantitatively or
34 qualitatively calibrated measure for comparison across space and subsystems; the detection and attribution evidence
35 has sufficient geographical distribution for a spatially representative sample. Confidence assessments for both
36 detection and attribution span a wide range. Confidence is highest in cryospheric systems (glaciers and permafrost),
37 with confidence in detection at least *medium* for various ecosystem measures but confidence in attribution lower
38 (see Figure 18-7). Within human and managed systems, confidence in attribution to observed climate change is *low*
39 *or very low* because of the dominating contribution of other drivers of change, including increased wealth, changes
40 in exposure, new crop varieties, and new technologies. Overall, detection and attribution analysis reveals evidence
41 for concern that risk of changes in aggregate measures has already changed, but also reveals *limited evidence* of a
42 climate change influence on aggregate measures which do not concern the cryosphere.

43
44 Overall, the synthesis of aggregated impact measures confirms the concern for globally aggregated impacts of recent
45 climate change being detected for a variety of systems and metrics.

46
47 [INSERT TABLE 18-12 HERE

48 Table 18-12: Confidence in detection of impacts on aggregate impact measures and confidence in attribution of at
49 least a minor role of climate change in those observed changes.]

50
51 [INSERT FIGURE 18-7 HERE

52 Figure 18-7: Confidence in detection of changes in aggregate impact measures and attribution of at least a minor
53 role of climate change in those trends]

1
2 18.6.2.5. Risks of Large-Scale Discontinuities
3

4 The RFC of “Risks of Large-Scale Discontinuities” “represents the likelihood that certain phenomena (sometimes
5 called singularities or tipping points) would occur, any of which may be accompanied by very large impacts” (Smith
6 *et al.*, 2009). Such discontinuities are hence a source of future risk, as they refer to thresholds in the Earth system
7 that, once passed, will alter the nature of the system itself. While several of the identified “tipping elements” form
8 part of the physical climate-ocean-cryosphere system (and are discussed in WGI), there are also elements in the
9 biosphere that exhibit non-linear behavior with potentially strong feedbacks on the Earth system (Lenton *et al.*,
10 2008; Leadley *et al.*, 2010).

11
12 For observed impacts, the concern translates into a question of the possible presence of “early warning signals” for
13 discontinuities that may be derived from monitoring changes in some climate or natural systems (Collie *et al.*, 2004;
14 deYoung *et al.*, 2008; Andersen *et al.*, 2009; Lenton, 2011). To this effect, conceptual, observational and modelling
15 studies of climate-change-induced biome or large-scale ecological regime shifts have intensified over the last decade
16 (deYoung *et al.*, 2008; Marengo *et al.*, 2011; e.g., Duarte *et al.*, 2012b; Mann *et al.*, 2012).

17
18 Evidence from detection and attribution analysis supports concerns that both the Arctic and the global warm-water
19 coral reef system are experiencing irreversible regime shifts, attributable to recent climate change, while
20 observations from the boreal and Amazon forest are less conclusive.

- 21 • For the Arctic region, new evidence indicates a biophysical regime shift is taking place, with social and
22 economic consequences (Post *et al.*, 2009; CAFF, 2010; Callaghan *et al.*, 2010; AMAP, 2011; Duarte *et al.*,
23 *et al.*, 2012b, see also 18.3.1, 24.4.2.2, 28.2). For the marine biota in the Arctic Ocean, the rapid reduction of
24 summer ice cover is now severely affecting pelagic ecosystems as well as ice-dependent mammals such as
25 seals and polar bears (Duarte *et al.*, 2012a). Thawing of Arctic permafrost and shrub encroachment on the
26 tundra have both been detected with *high confidence* (18.3.2.4, 18.5.8, 24.4.2.2), driven by warming and an
27 associated prolongation of the growing season. Wide-spread hydrological changes have been observed in
28 the tundra region, as a result of increasing winter rains and the degradation of permafrost, leading to lake
29 formation or disappearance within a few years’ time (CAFF, 2010; Callaghan *et al.*, 2013) and cascading
30 consequences for the tundra food webs (Post *et al.*, 2009; Callaghan *et al.*, 2013; Hansen *et al.*, 2013).
- 31 • Warm-water coral reefs have been lost at a large scale due to thermal stress and other impacts, as was noted
32 by AR4 already. Increased mass bleaching and mortality of corals has been detected and attributed globally
33 with *very high confidence* (see Box 18-3), with at least *high confidence* regionally in tropical Asia, Japan,
34 the Australian Great Barrier Reef, the western Caribbean and coast of Central America, and the waters
35 around other small tropical islands (6.3.2, 24.4.3.2, 27.3.3.1, 29.3.1.2). This irreversible loss of biodiversity
36 has significant feedbacks within the marine biosphere, and cascading consequences for regional marine
37 ecosystems as well as the human livelihoods which depend on them. The additional evidence for ongoing
38 change and its attribution to warming gained since the AR4 strengthens the conclusion that coral reef loss
39 constitutes a strong warning signal for the irreversible loss of an entire biome.
- 40 • Dieback and degradation in the Boreal forests and the Amazonian rainforest have been identified as
41 potential tipping elements, due to their large extent and the possible feedbacks with the carbon cycle
42 (Lenton *et al.*, 2008; Malhi *et al.*, 2009; Leadley *et al.*, 2010; Marengo *et al.*, 2011, see also 4.3.3.1,
43 4.3.3.4). For the boreal forest, increases in tree mortality are observed in many regions (4.3.3), including
44 wide-spread dieback related to insect infestations in North America (26.4.1), but there is *low confidence* in
45 detection of this as a global trend attributable to climate change. In the humid tropical forests of the
46 Amazon basin, tree turnover (both mortality and growth) has increased during recent decades. A number of
47 factors have likely played a role, including the direct effect of rising CO₂ on lianas, recovery from past
48 disturbance, and changing climate. The reason for concern about the Amazon forest is the interaction
49 between global climate change, regional climate change related to deforestation, and the high susceptibility
50 of forests to fire, which together could lead to degradation of forests in large areas of the Amazon above
51 deforestation itself (Malhi *et al.*, 2009). There are indications of droughts in the Amazon and evidence that
52 dry years and deforestation increase the vulnerability of forests to fire, but there is only *very low confidence*
53 in attribution of observed changes in tree turnover to climate change. In conclusion, there is currently

1 insufficient evidence from observed climate change impacts to support a climate-related warning sign of
2 possible large-scale discontinuities in the boreal and Amazonian forest.
3

4 In the TAR, the risk of large-scale discontinuities was evaluated to be a concern only after a few degrees warming.
5 At the time of the AR4, new understanding of physical climate processes led to a major revision of that assessment
6 by Smith *et al.* (2009), bringing concerns forward that discontinuities could occur with much less warming. While
7 the present analysis does not indicate the manifestation of a tipping point being reached, it does indicate significant
8 early warning indicators for possible regime shifts in the Arctic region and for the world's warm-water coral
9 ecosystems. Observations therefore confirm the reasons for concern about risks from large-scale singularities.
10

11 12 **18.6.3. Conclusion** 13

14 The body of scientific evidence on observed impacts of recent climate change, after rigorous assessment of scientific
15 confidence in its attribution to recent trends, lends new qualitative support to four out of five reasons for concern
16 established by earlier IPCC assessments. Specifically, concerns are confirmed for risks to unique and threatened
17 systems, risks stemming from extreme weather events, globally aggregated impacts and – in terms of early warnings
18 – risks of large-scale discontinuities. Only the spatial or social disparities covered under “distribution of impacts”
19 are still insufficiently studied to permit a synthesis of available observations for the characterization of a global
20 concern. While the Arctic stands out as a region with robust evidence of impacts across numerous systems, current
21 detection and attribution literature does not address whether the severity of those impacts differs from other regions.
22 Across RFCs, the critical evidence often comes from the same environmental systems, notably the Arctic region,
23 warm-water coral reef systems and mountain glaciers, but there are also important observations from impacted
24 hydrological systems and human systems including agriculture.
25

26 Detection and attribution studies evaluate the agreement between observations and process understanding, with the
27 important requirement for direct observational evidence of the impacts. This sets a higher bar for establishing
28 confidence in past changes than is generally used for assessing confidence in projected future changes, because
29 observational evidence has important gaps, while the plausibility of future changes can be established on the basis of
30 process knowledge only (WGI AR5 Chapter 10.2.5).
31

32 Despite this constraint, the evidence gathered since the AR4 on detection and attribution of observed impacts from
33 climate change has reached a level at which it can inform evaluation of many of the aspects of present-day climate
34 change risk as described by the RFCs. In particular, the geographic distribution of studies is reaching the point
35 where assessment of the global nature of impacts is possible:

- 36 • There is now *robust evidence* of observed changes in natural systems in all of the regional groupings used
37 in this report. There is at least *medium confidence* in attribution of observed changes in various components
38 of the cryosphere to observed climate change for each of the inhabited continents. There is also at least
39 *medium confidence* in attribution of observed changes in terrestrial ecosystems to observed climate change
40 for six continents (exception being South America) and for some small islands, and for marine ecosystems
41 surrounding six continents (exception being Africa) and for some small islands.
- 42 • There is *good evidence* of the detection of impacts in human systems on the inhabited continents. There is
43 at least *medium confidence* in detection of impacts on food production in all the inhabited continents.
- 44 • While the current detection and attribution literature does not reveal observational evidence of geographical
45 differences in the severity of climate change impacts between continents, it does indicate that the unique
46 systems of the Arctic region and warm water coral reefs are undergoing rapid changes in response to
47 observed warming in ways that are potentially irreversible.
48

49 50 **18.7. Gaps, Research Needs, Emerging Issues** 51

52 While the literature formalizing accepted approaches and methods for detection and attribution studies on human
53 and natural systems is relatively recent, the underlying question is one with a long tradition: how to detect a
54 relationship between cause (climate change) and effect (impacts)? The literature on climate change impacts has

1 largely focused on providing estimates of *future* impacts covering the globe and almost any possible affected human
2 and natural system, *observed* impacts have received comparably little attention. In order to provide better guidance
3 for the policy-relevant risk assessment of future impacts, the coverage of impacts that have already occurred needs
4 to be strengthened
5

6 With the exception of the food system, detection and attribution studies are most scarce in human systems. While
7 sensitivity to climate variability is clear for many important economic sectors and infrastructures, this is not
8 mirrored in the literature available on observed impacts of climate change.
9

10 In part, this is due to the fact that human systems are extremely complex and evolving at a very rapid pace, with
11 many other drivers and their interactions challenging assessments of the comparatively weak recent climate signals.
12 Human systems are also capable of autonomous and planned adaptation, including adaptation to predicted climate
13 change, complicating the diagnosis of impacts of observed climate change. The literature on observed climate
14 adaptation response is sparse. More fundamentally, much of the research on the relation between past climate
15 change and impacts in human systems has been of a nature that has not fit into the deterministic, quantitative
16 detection and attribution framework adopted in earlier Assessment Reports (Stone *et al.*, 2013). Fully including
17 human systems into a complete understanding of observed impacts should be a priority for assessing future
18 vulnerability to climate change.
19

20 The current detection and attribution literature does not provide a sufficiently comprehensive sampling of the
21 distribution of impacts around the world, for instance between poor and wealthy populations. This dearth of studies
22 reflects inadequate or a lack of long term monitoring or observational systems, an insufficient network density of
23 meteorological observations, and/or lack of research funding, together leading to limited research capability. One
24 other important reason is that in some sub-regions, particularly in the tropics, temperature sensitivities are poorly
25 understood because rainfall is the primary climate driver, and precipitation trends under anthropogenic climate
26 change are not as clearly understood as temperature trends.
27

28 Another challenge lies in addressing changes in impacts of extreme weather events and their relation to climate
29 change, as well as determining the role of climate change in special events in impact systems. In some cases
30 difficulty arises from dominating influences of non-climate factors on the extent of actual impacts of extreme
31 weather events and the often highly nonlinear interactions of those factors, but a general characteristic lies in the
32 difficulty of characterizing changes in the frequency or intensity of events that occur only very rarely.
33
34

35 **Frequently Asked Questions**

36 ***FAQ 18.1: What are the main challenges in detecting climate change impacts?***

37 Detection addresses the question of whether a system has changed beyond its expected behavior in the absence of
38 climate change. One challenge in detection is distinguishing between a change in a system and natural variability in
39 the system. Most natural and human systems will exhibit variability over time even in the absence of a change in
40 external factors. Particularly over short periods of time, this kind of natural variability can give the appearance of a
41 systematic change. For example, even in the absence of changes in external factors, some wild populations can
42 undergo prolonged periods of boom or bust. Distinguishing natural variability from systematic change usually
43 requires an understanding of the characteristics of the former. This can be based on an analysis of historical data or an
44 understanding of the dynamics of the system. A second challenge to detection is distinguishing between between a
45 change in the system that is potentially attributable to climate change and one that is due to a change in non-climatic
46 factors. For example, wild populations may change as a result of harvesting or habitat loss unconnected to climate
47 change. Distinguishing such changes from those potentially attributable to climate change requires an understanding
48 of the non-climatic factors that can affect the system and how these factors have changed.
49
50

51 ***FAQ 18.2: What is the main challenge in attributing changes in a system to climate change?***

52 Attribution addresses the question of whether changes detected in a system can be attributed to climate change.
53 Once a climate-related effect on a system has been detected, the main challenge in attribution is determining
54 whether this effect is due to climate change rather than natural climate variability. For example, widespread flooding

1 in Australia and other parts of the western Pacific during 2010 and 2011 was caused by unusually heavy rainfall.
2 However, this heavy rainfall was found to be related to the occurrence of La Niña, part of the naturally occurring
3 ENSO variation, and therefore the flooding is not attributable to climate change.

4
5 **FAQ 18.3: Why are detection and attribution of climate impacts important?**

6 In deciding how to respond to climate change, policy-makers and others need to understand what the future impacts
7 of climate change will be. One way to gain understanding about the future impacts of climate is to identify impacts
8 of climate change that has already occurred. For example, it has been predicted that climate change will cause a
9 poleward shift in the geographical ranges of species. Detecting such a shift in response to climate change that has
10 already occurred would tend to validate this prediction.

11
12 **FAQ 18.4: Is it possible to attribute a single event, like a disease outbreak
13 or the extinction of a species, to climate change?**

14 Scientists are usually reluctant to attribute a single event to climate change and instead tend to focus on the
15 frequency or severity of classes of events. One reason for this is that the scientific knowledge needed to attribute a
16 single event to climate change is much greater than that needed to attribute a change in the frequency or severity of
17 classes of events. For example, there is good evidence that there has been an increase in the frequency of heat waves
18 in many parts of the world and that this increase can be attributed to climate change. However, claiming that, in the
19 absence of climate change, a particular heat wave would not have occurred is largely beyond present scientific
20 capabilities.

21
22
23 **References**

- 24
25 ACIA, 2005: *Arctic Climate Impact Assessment, Scientific Report*. In: Cambridge University Press, Cambridge,
26 U.K.
- 27 Adano, W.R., T. Dietz, K. Witsenburg, and F. Zaal, 2012: Climate change, violent conflict and local institutions in
28 Kenya's drylands. *Journal of Peace Research*, **49**, 65-80.
- 29 Aerts, J.C.J.H., N. Lin, W.J.W. Botzen, K. Emanuel, and H. De Moel, 2013: Low-probability flood risk modeling
30 for New York City. *Risk Analysis*, **in press**.
- 31 Agrawal, A., 1995: Dismantling the divide between indigenous and scientific knowledge. *Development and Change*,
32 **26(3)**, 413-439.
- 33 Aguilera, H. and J. Murillo, 2009: The effect of possible climate change on natural groundwater recharge based on a
34 simple model: a study of four karstic aquifers in SE Spain. *Environmental Geology*, **57(5)**, 963-974.
- 35 Akpınar-Ferrand, E. and A. Singh, 2010: Modeling increased demand of energy for air conditioners and consequent
36 CO₂ emissions to minimize health risks due to climate change in India. *Environmental Science & Policy*,
37 **13(8)**, 702-712.
- 38 Alderman, H., 2010: Safety nets can help address the risks to nutrition from increasing climate variability. *The*
39 *Journal of Nutrition*, **140(1)**, 148-152.
- 40 Alexander, C., N. Bynum, E. Johnson, U. King, T. Mustonen, P. Neofotis, N. Oettle, C. Rosenzweig, C. Sakakibara,
41 V. Shadrin, M. Vicarelli, J. Waterhouse, and B. Weeks, 2011: Linking indigenous and scientific knowledge of
42 climate change. *Bioscience*, **61**, 477-484.
- 43 Alexander, L.V., X. Zhang, T.C. Peterson, J. Caesar, B. Gleason, A. Klein Tank, M. Haylock, D. Collins, B. Trewin,
44 and F. Rahimzadeh, 2006: Global observed changes in daily climate extremes of temperature and precipitation.
45 *Journal of Geophysical Research*, **111**.
- 46 Alheit, J., T. Pohlmann, M. Casini, W. Greve, R. Hinrichs, M. Mathis, K. O'Driscoll, R. Vorberg, and C. Wagner,
47 2012: Climate variability drives anchovies and sardines into the North and Baltic Seas. *Progress in*
48 *Oceanography*, **96(1)**, 128-139.
- 49 Allen, M., 2011: In defense of the traditional null hypothesis: remarks on the Trenberth and Curry WIREs opinion
50 articles. *Wiley Interdisciplinary Reviews: Climate Change*, **2**, 931-934.
- 51 Allen, S.K., S.C. Cox, and I.F. Owens, 2010: Rock-avalanches and other landslides in the central Southern Alps of
52 New Zealand: a regional assessment of possible climate change impacts. *Landslides*, **8(1)**, 33-48.

- 1 Alling, A., O. Doherty, H. Logan, L. Feldman, and P. Dustan, 2007: *Catastrophic coral mortality in the remote*
2 *central Pacific Ocean: Kiribati Phoenix Islands*. In: Atoll Research Bulletin. National Museum of Natural
3 History Smithsonian Institution, Washington DC, USA.
- 4 Allison, E.H., A.L. Perry, M.C. Badjeck, W.N. Adger, K. Brown, D. Conway, A.S. Halls, G.M. Pilling, J.D.
5 Reynolds, N.L. Andrew, and N.K. Dulvy, 2009: Vulnerability of national economies to the impacts of climate
6 change on fisheries. *Fish and Fisheries*, **10(2)**, 173-196.
- 7 Alongi, D.M., 2008: Mangrove forests: Resilience, protection from tsunamis, and responses to global climate
8 change. *Estuarine Coastal and Shelf Science*, **76(1)**, 1-13.
- 9 Alonso, D., M.J. Bouma, and M. Pascual, 2011: Epidemic malaria and warmer temperatures in recent decades in an
10 East African highland. *Proceedings of the Royal Society B: Biological Sciences*, **278**, 1661-1669.
- 11 Alston, M., 2011: Gender and climate change in Australia. *Journal of Sociology*, **47(1)**, 53-70.
- 12 AMAP, 2011: *Snow, water, ice and permafrost in the Arctic (SWIPA): Climate change and the cryosphere*. Arctic
13 *monitoring and assessment programme (AMAP)*. Oslo, Norway, pp. 538.
- 14 Amstrup, S.C., E.T. DeWeaver, D.C. Douglas, B.G. Marcot, G.M. Durner, C.M. Bitz, and D.A. Bailey, 2010:
15 Greenhouse gas mitigation can reduce sea-ice loss and increase polar bear persistence. *Nature*, **468(7326)**,
16 955-958.
- 17 Amthor, J.S., 2001: Effects of atmospheric CO₂ concentration on wheat yield. Review of results from experiments
18 using various approaches to control CO₂ concentration. *Field Crops Research*, **73(1)**, 1-34.
- 19 Andersen, T., J. Carstensen, E. Hernández-García, and C.M. Duarte, 2009: Ecological thresholds and regime shifts:
20 approaches to identification. *Trends in Ecology & Evolution*, **24(1)**, 49-57.
- 21 Anderson, C.A., 1987: Temperature and aggression: effects on quarterly, yearly, and city rates of violent and
22 nonviolent crime. *Journal of Personality and Social Psychology*, **52(6)**, 1161.
- 23 Anderson, C.A., 2001: Heat and violence. *Current Directions in Psychological Sciences*, **10(1)**, 33-38.
- 24 Aporta, C., D.R.F. Taylor, and G.J. Laidler, 2011: Geographies of Inuit sea ice use. Introduction. *Canadian*
25 *Geographer / Le Géographe Canadien*, **55**, 1-5.
- 26 Ariano, R., G.W. Canonica, and G. Passalacqua, 2010: The possible role of climate changes in variations of pollen
27 seasons and allergic sensitizations over 27 years. *The Journal of Allergy and Clinical Immunology*, **125(2)**,
28 AB192.
- 29 Armah, F.A., J.O. Odoi, G.T. Yengoh, S. Obiri, D.O. Yawson, and E.K.A. Afrifa, 2011: Food security and climate
30 change in drought-sensitive savanna zones of Ghana. *Mitigation and Adaptation Strategies for Global Change*,
31 **16(3)**, 291-306.
- 32 Arora-Jonsson, S., 2011: Virtue and vulnerability: Discourses on women, gender and climate change. *Global*
33 *Environmental Change*, **21(2)**, 744-751.
- 34 Arrigo, K.R. and G.L. Van Dijken, 2011: Secular trends in Arctic Ocean net primary production. *Journal of*
35 *Geophysical Research*, **116(C9)**, C09011.
- 36 Askari, M.T., M.Z.A. Ab Kadir, W.F.W. Ahmad, and M. Izadi, 2009: Investigate the effect of variations of ambient
37 temperature on HST of transformer. Proceedings of 2009 IEEE Student Conference on Research and
38 Development (SCORED), 16.11.2009, pp. 363-367.
- 39 Atkinson, A., V. Siegel, E. Pakhomov, and P. Rothery, 2004: Long-term decline in krill stock and increase in salps
40 within the Southern Ocean. *Nature*, **432**, 100-103.
- 41 Auffhammer, M., V. Ramanathan, and J.R. Vincent, 2006: Integrated model shows that atmospheric brown clouds
42 and greenhouse gases have reduced rice harvests in India. *Proceedings of the National Academy of Sciences of*
43 *the United States of America*, **103(52)**, 19668-19672.
- 44 Ayyappan, R., S. Sankar, P. Rajkumar, and K. Balakrishnan, 2009: Work-related heat stress concerns in automotive
45 industries. A case study from Chennai, India. *Global Health Action*, **2**.
- 46 Badjeck, M.C., E.H. Allison, A.S. Halls, and N.K. Dulvy, 2010: Impacts of climate variability and change on
47 fishery-based livelihoods. *Marine Policy*, **34(3)**, 375-383.
- 48 Bajracharya, S.R. and P. Mool, 2009: Glaciers, glacial lakes and glacial lake outburst floods in the Mount Everest
49 region, Nepal. *Annals of Glaciology*, **50(53)**, 81-86.
- 50 Baker, A.C., P.W. Glynn, and B. Riegl, 2008: Climate change and coral reef bleaching: An ecological assessment of
51 long-term impacts, recovery trends and future outlook. *Estuarine, Coastal and Shelf Science*, **80(4)**, 435-471.
- 52 Balakrishnan, K., A. Ramalingam, V. Dasu, J.C. Stephen, M.R. Sivaperumal, D. Kumarasamy, K. Mukhopadhyay,
53 S. Ghosh, and S. Sambandam, 2010: Case studies on heat stress related perceptions in different industrial
54 sectors in southern India. *Global Health Action*, **3**.

- 1 Ballantyne, C.K., 2002: Paraglacial geomorphology. *Quaternary Science Reviews*, **21(18-19)**, 1935-2017.
- 2 Ballu, V., M.N. Bouin, P. Siméoni, W.C. Crawford, S. Calmant, J.M. Boré, T. Kanas, and B. Pelletier, 2011:
3 Comparing the role of absolute sea-level rise and vertical tectonic motions in coastal flooding, Torres Islands
4 (Vanuatu). *Proceedings of the National Academy of Sciences of the United States of America*, **108(32)**, 13019-
5 13022.
- 6 Banfai, D.S. and D. Bowman, 2007: Drivers of rain-forest boundary dynamics in Kakadu National Park, Northern
7 Australia. A field assessment. *Journal of Tropical Ecology*, **23**, 73-86.
- 8 Baraer, M., B.G. Mark, J.M. McKenzie, T. Condom, J. Bury, K.I. Huh, C. Portocarrero, J. Gómez, and S. Rathay,
9 2012: Glacier recession and water resources in Peru's Cordillera Blanca. *Journal of Glaciology*, **58(207)**, 134-
10 150.
- 11 Barange, M. and R.I. Perry, 2009: *Physical and ecological impacts of climate change relevant to marine and inland*
12 *capture fisheries and aquaculture*. In: Climate change implications for fisheries and aquaculture: Overview
13 over current scientific knowlegde [Cochrane, K., de Young, C., Soto, D. and Bahri, T. (eds.)]. FAO, Rome, pp.
14 7-212.
- 15 Barlow, P.M. and E.G. Reichard, 2010: Saltwater intrusion in coastal regions of North America. *Hydrogeology*
16 *Journal*, **18(1)**, 247-260.
- 17 Barnett, T.P., D.W. Pierce, H.G. Hidalgo, C. Bonfils, B.D. Santer, T. Das, G. Bala, A.W. Wood, T. Nozawa, A.A.
18 Mirin, D.R. Cayan, and M.D. Dettinger, 2008: Human-induced changes in the hydrology of the western United
19 States. *Science*, **319(5866)**, 1080-1080.
- 20 Barredo, J.I., D. Saurí, and M.C. Llasat, 2012: Assessing trends in insured losses from floods in Spain 1971–2008.
21 *Natural Hazards Earth System Sciences*, **12**, 1723-1729.
- 22 Barredo, J.I., 2010: No upward trend in normalised windstorm losses in Europe: 1970-2008. *Natural Hazards and*
23 *Earth System Sciences*, **10**, 97-104.
- 24 Barriopedro, D., E.M. Fischer, J. Luterbacher, R.M. Trigo, and R. García-Herrera, 2011: The hot summer of 2010:
25 redrawing the temperature record map of Europe. *Science*, **332(6026)**, 220-224.
- 26 Barrios, S., L. Bertinelli, and E. Strobl, 2006: Climatic change and rural-urban migration. The case of sub-Saharan
27 Africa. *Journal of Urban Economics*, **60(3)**, 357-371.
- 28 Barros, V.R., 2010: Agro y ambiente: Una agenda compartida para el desarrollo sustentable. El cambio climático en
29 Argentina (Chapter 3). In: Foro de la Cadena Agroindustrial Argentina, Buenos Aires, Argentina, .
- 30 Barthel, F. and E. Neumayer, 2012: A trend analysis of normalized insured damage from natural disasters. *Climatic*
31 *Change*, **113(2)**, 215-237.
- 32 Bauder, A., M. Funk, and M. Huss, 2007: Ice-volume changes of selected glaciers in the Swiss Alps since the end of
33 the 19th century. *Annals of Glaciology*, **46(1)**, 145-149.
- 34 Beaugrand, G., M. Edwards, and L. Legendre, 2010: Marine biodiversity, ecosystem functioning, and carbon cycles.
35 *Proceedings of the National Academy of Sciences of the United States of America*, **107(22)**, 10120-10124.
- 36 Beaugrand, G., C. Luczak, and M. Edwards, 2009: Rapid biogeographical plankton shifts in the North Atlantic
37 Ocean. *Global Change Biology*, **15(7)**, 1790-1803.
- 38 Beaugrand, G., P.C. Reid, F. Ibanez, J.A. Lindley, and M. Edwards, 2002: Reorganization of North Atlantic marine
39 copepod biodiversity and climate. *Science*, **296(5573)**, 1692-1694.
- 40 Beaumier, M. and J.D. Ford, 2010: Food insecurity among Inuit women exacerbated by socio-economic stresses and
41 climate change. *Canadian Journal of Public Health*, **101(3)**, 196-201.
- 42 Becken, S. and J. Hay, 2007: *Tourism and climate change: risks and opportunities*. Channel View Publications,
43 Clevedon, pp. 350.
- 44 Becker, M., B. Meyssignac, C. Letetrel, W. Llovel, A. Cazenave, and T. Delcroix, 2012: Sea level variations at
45 tropical Pacific islands since 1950. *Global and Planetary Change*, **80**, 85-98.
- 46 Bednaršek, N., G.A. Tarling, D.C.E. Bakker, S. Fielding, E.M. Jones, H.J. Venables, P. Ward, A. Kuzirian, B. Lézé,
47 R.A. Feely, and E.J. Murphy, 2012: Extensive dissolution of live pteropods in the Southern Ocean. *Nature*
48 *Geoscience*, **5**, 881-885.
- 49 Behrenfeld, M.J., R.T. O'Malley, D.A. Siegel, C.R. McClain, J.L. Sarmiento, G.C. Feldman, A.J. Milligan, P.G.
50 Falkowski, R.M. Letelier, and E.S. Boss, 2006: Climate-driven trends in contemporary ocean productivity.
51 *Nature*, **444(7120)**, 752-755.
- 52 Belkin, I.M., 2009: Rapid warming of large marine ecosystems. *Progress in Oceanography*, **81(1-4)**, 207-213.
- 53 Bell, M.A. and R.A. Fischer, 1994: Using yield prediction models to assess yield gains. A case study for wheat.
54 *Field Crops Research*, **36(2)**, 161-166.

- 1 Beniston, M., M. Stoffel, and M. Hill, 2011: Impacts of climatic change on water and natural hazards in the Alps:
2 Can current water governance cope with future challenges? Examples from the European “ACQWA” project.
3 *Environmental Science & Policy*, **14(7)**, 734-743.
- 4 Benning, T.L., D. LaPointe, C.T. Atkinson, and P.M. Vitousek, 2002: Interactions of climate change with biological
5 invasions and land use in the Hawaiian Islands: Modeling the fate of endemic birds using a geographic
6 information system. *Proceedings of the National Academy of Sciences of the United States of America*, **99(22)**,
7 14246-14249.
- 8 Berkes, F., 2009: Indigenous ways of knowing and the study of environmental change. *Journal of the Royal Society*
9 *of New Zealand*, **39**, 151-156.
- 10 Beven, J.L., L.A. Avila, E.S. Blake, D.P. Brown, J.L. Franklin, R.D. Knabb, R.J. Pasch, J.R. Rhome, and S.R.
11 Stewart, 2008: Atlantic hurricane season of 2005. *Monthly Weather Review*, **136(3)**, 1109-1173.
- 12 Biasutti, M. and A. Giannini, 2006: Robust Sahel drying in response to late 20th century forcings. *Geophysical*
13 *Research Letters*, **33(11)**, L11706.
- 14 Biggs, R., S.R. Carpenter, and W.A. Brock, 2009: Turning back from the brink: Detecting an impending regime
15 shift in time to avert it. *Proceedings of the National Academy of Sciences of the United States of America*,
16 **106(3)**, 826-831.
- 17 Björk, R.G. and U. Molau, 2007: Ecology of alpine snow beds and the impact of global change. *Arctic, Antarctic*
18 *and Alpine Research*, **39(1)**, 34-43.
- 19 Black, R., W.N. Adger, N.W. Arnell, S. Dercon, A. Geddes, and D. Thomas, 2011: The effect of environmental
20 change on human migration. *Global Environmental Change*, **21(Supplement 1)**, 3-11.
- 21 Blok, D., U. Sass-Klaassen, G. Schaepman-Strub, M. Heijmans, P. Sauren, and F. Berendse, 2011: What are the
22 main climate drivers for shrub growth in Northeastern Siberian tundra? *Biogeosciences*, **8(5)**, 1169-1179.
- 23 Bolch, T., A. Kulkarni, A. Kääb, C. Huggel, F. Paul, G. Cogley, H. Frey, J.S. Kargel, K. Fujita, M. Scheel, S.
24 Bajracharya, and M. Stoffel, 2012: The state and fate of Himalayan glaciers. *Science*, **336(6079)**, 310-314.
- 25 Bongaerts, P., T. Ridgway, E. Sampayo, and O. Hoegh-Guldberg, 2010: Assessing the ‘deep reef refugia’
26 hypothesis. Focus on Caribbean reefs. *Coral Reefs*, **29(2)**, 309-327.
- 27 Borges, A. and N. Gypens, 2010: Carbonate chemistry in the coastal zone responds more strongly to eutrophication
28 than to ocean acidification. *Limnology & Oceanography*, **55(1)**.
- 29 Boruff, B.J., J.A. Easoz, S.D. Jones, H.R. Landry, J.D. Mitchem, and S.L. Cutter, 2003: Tornado hazards in the
30 United States. *Climate Research*, **24(2)**, 103-117.
- 31 Boustan, L.P., M.E. Kahn, and P.W. Rhode, 2012: Moving to higher ground. Migration response to natural disasters
32 in the early twentieth century. *The American Economic Review*, **102(3)**, 238-244.
- 33 Bouwer, L.M., R.P. Crompton, E. Faust, P. Hoppe, and R.A. Pielke Jr, 2007: Confronting disaster losses. *Science*,
34 **318(11)**, 753-753.
- 35 Bouwer, L.M., 2011: Have disaster losses increased due to anthropogenic climate change? *Bulletin of the American*
36 *Meteorological Society*, **92(1)**, 39-46.
- 37 Bouwer, L.M. and W.J.W. Botzen, 2011: How sensitive are US hurricane damages to climate? Comment on a paper
38 by W.D. Nordhaus. *Climate Change Economics*, **02(01)**, 1-7.
- 39 Bowman, D., B.P. Murphy, and D.S. Banfai, 2010: Has global environmental change caused monsoon rainforests to
40 expand in the Australian monsoon tropics? *Landscape Ecology*, **25(8)**, 1247-1260.
- 41 Bradley, R.S., F.T. Keimig, H.F. Diaz, and D.R. Hardy, 2009: Recent changes in freezing level heights in the tropics
42 with implications for the deglaciation of high mountain regions. *Geophysical Research Letters*, **36(17)**,
43 L17701.
- 44 Brander, K.M., 2007: Global fish production and climate change. *Proceedings of the National Academy of Sciences*
45 *of the United States of America*, **104**, 19709-19714.
- 46 Brander, K., 2010: Impacts of climate change on fisheries. *Journal of Marine Systems*, **79(3-4)**, 389-402.
- 47 Brander, K., J. Bruno, A. Hobday, and D. Schoeman, 2011: The value of attribution. *Nature Climate Change*, **1(2)**,
48 70-71.
- 49 Breetzke, G.D. and E.G. Cohn, 2012: Seasonal assault and neighborhood deprivation in South Africa - some
50 preliminar findings. *Environment and Behavior*, **44(5)**, 641-667.
- 51 Brisson, N., P. Gate, D. Gouache, G. Charmet, F. Oury, and F. Huard, 2010: Why are wheat yields stagnating in
52 Europe? A comprehensive data analysis for France. *Field Crops Research*, **119(1)**, 201-212.
- 53 Brook, T., 2010: *The troubled empire: China in the Yuan and Ming Dynasties*. Belknap Press, Cambridge, pp. 329.

- 1 Brooks, H.E. and C.A. Doswell, 2002: Deaths in the 3 may 1999 Oklahoma city tornado from a historical
2 perspective. *Weather and Forecasting*, **17(3)**, 354-361.
- 3 Bruno, J.F. and E.R. Selig, 2007: Regional decline of coral cover in the Indo-Pacific: timing, extent, and subregional
4 comparisons. *PLoS ONE*, **2(8)**, e711.
- 5 Bryant, D., L. Burke, J. McManus, and M. Spalding, 1998: *Reefs at risk: a map-based indicator of threats to the*
6 *world's coral reefs*. World Resources Institute, Washington, DC, .
- 7 Buhaug, H. and O.M. Theisen, 2012: On environmental change and armed conflict. In: *Climate change, human*
8 *security and violent conflict*. [Scheffran, J. (ed.)]. Springer-Verlag, Berlin Heidelberg, pp. 43-55.
- 9 Buhaug, H., H. Hegre, and H. Strand, 2010: *Sensitivity analysis of climate variability and civil war*. PRIO Paper.
10 Peace Research Institute Oslo, PRIO, Oslo.
- 11 Bullard, R. and B. Wright, 2010: *Race, place, and environmental justice after Hurricane Katrina: Struggles to*
12 *reclaim, rebuild, and revitalize New Orleans and the Gulf Coast*. Westview Press, Boulder, CO, pp. 312.
- 13 Bunce, M., S. Rosendo, and K. Brown, 2010: Perceptions of climate change, multiple stressors and livelihoods on
14 marginal African coasts. *Environment, Development and Sustainability*, **12(3)**, 407-440.
- 15 Burke, M.B., E. Miguel, S. Satyanath, J.A. Dykema, and D.B. Lobell, 2009: Warming increases the risk of civil war
16 in Africa. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 20670-
17 20674.
- 18 Burke, L., K. Reyntar, M. Spalding, and A. Perry, 2011: *Reefs at risk revisited*. World Resources Institute,
19 Washington, DC, .
- 20 Burrows, M.T., D.S. Schoeman, L.B. Buckley, P. Moore, E.S. Poloczanska, K.M. Brander, C. Brown, J.F. Bruno,
21 C.M. Duarte, B.S. Halpern, J. Holding, C.V. Kappel, W. Kiessling, M.I. O'Connor, J.M. Pandolfi, C.
22 Parmesan, F.B. Schwing, W.J. Sydeman, and A.J. Richardson, 2011: The pace of shifting climate in marine
23 and terrestrial ecosystems. *Science*, **334(6056)**, 652-655.
- 24 Bury, J., B.G. Mark, J. McKenzie, A. French, M. Baraer, K. In Huh, M. Zapata Luyo, and R.J. Gómez López, 2010:
25 Glacier recession and human vulnerability in the Yanamarey Watershed of the Cordillera Blanca, Peru.
26 *Climatic Change*, **105**, 179-206.
- 27 Butke, P. and S.C. Sheridan, 2010: An analysis of the relationship between weather and aggressive crime in
28 Cleveland, Ohio. *Weather, Climate, and Society*, **2(2)**, 127-139.
- 29 Butler, C. and S. Gates, 2012: African range wars: Climate, conflict, and property rights. *Journal of Peace Research*,
30 **49(1)**, 23-34.
- 31 Butt, N., P.A. De Oliveira, and M.H. Costa, 2011: Evidence that deforestation affects the onset of the rainy season in
32 Rondonia, Brazil. *Journal of Geophysical Research-Atmospheres*, **116**, D11120.
- 33 Byg, A. and J. Salick, 2009: Local perspectives on a global phenomenon-Climate change in Eastern Tibetan
34 villages. *Global Environmental Change*, **19**, 156-166.
- 35 CAFF, 2010: *Arctic biodiversity trends 2010 – Selected indicators of change*. Arctic Council, Conservation of
36 Arctic Flora and Fauna Working Group, International Secretariat, Aujureyri, Iceland, pp. 124.
- 37 Cai, W., T. Cowan, P. Briggs, and M. Raupach, 2009: Rising temperature depletes soil moisture and exacerbates
38 severe drought conditions across southeast Australia. *Geophysical Research Letters*, **36(21)**, L21709.
- 39 Cai, W. and P. Van Rensch, 2012: The 2011 southeast Queensland extreme summer rainfall: A confirmation of a
40 negative Pacific Decadal Oscillation phase? *Geophysical Research Letters*, **39(8)**, L08702.
- 41 Callaghan, T.V., F. Bergholm, T.R. Christensen, C. Jonasson, U. Kokfelt, and M. Johansson, 2010: A new climate
42 era in the sub-Arctic. Accelerating climate changes and multiple impacts. *Geophysical Research Letters*,
43 **37(14)**, L14705.
- 44 Callaghan, T.V., M. Johansson, R.D. Brown, P.Y. Groisman, N. Labba, V. Radionov, R.G. Barry, R.S. Bradley, S.
45 Blangy, and O.N. Bulygina, 2011: Changing snow cover and its impacts. In: *Snow, water, ice and permafrost*
46 *in the arctic (SWIPA)*. Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway, pp. 1-58.
- 47 Callaghan, T.V., C. Jonasson, T. Thierfelder, Y. Zhenlin, H. Hedenås, M. Johansson, U. Molau, R. Van Bogaert, J.
48 Olofsson, D. Jones, S. Bokhorst, G. Phoenix, J. Bjerke, H. Tømmervik, T. Christensen, E. Hanna, E. Koller,
49 and V. Sloan, 2013: Ecosystem change and stability over multiple decades in the Swedish sub-Arctic: complex
50 processes and multiple drivers. *Philosophical Transactions of the Royal Society, B, Biological Sciences*, (in
51 press).
- 52 Callaway, R., A.P. Shinn, S.E. Grenfell, J.E. Bron, G. Burnell, E.J. Cook, M. Crumlish, S. Culloty, K. Davidson,
53 and R.P. Ellis, 2012: Review of climate change impacts on marine aquaculture in the UK and Ireland. *Aquatic*
54 *Conservation: Marine and Freshwater Ecosystems*, **22(3)**, 389-421.

- 1 Cambers, G., 2009: Caribbean beach changes and climate change adaptation. *Aquatic Ecosystem Health and*
2 *Management*, **12(2)**, 168-176.
- 3 Carbognin, L., P. Teatini, A. Tomasin, and L. Tosi, 2010: Global change and relative sea level rise at Venice: what
4 impact in term of flooding. *Climate Dynamics*, **35(6)**, 1039-1047.
- 5 Cardoso, P.G., D. Raffaelli, and M.A. Pardal, 2008: The impact of extreme weather events on the seagrass *Zostera*
6 *noltii* and related *Hydrobia ulvae* population. *Marine Pollution Bulletin*, **56(3)**, 483-492.
- 7 Carey, M., 2005: Living and dying with glaciers. People's historical vulnerability to avalanches and outburst floods
8 in Peru. *Global and Planetary Change*, **47(2-4)**, 122-134.
- 9 Carey, M., 2010: *In the shadow of melting glaciers. Climate change and Andean society*. Oxford University Press,
10 New York, .
- 11 Carey, M., A. French, and E. O'Brien, 2012a: Unintended effects of technology on climate change adaptation. An
12 historical analysis of water conflicts below Andean glaciers. *Journal of Historical Geography*, **38**, 181-191.
- 13 Carey, M., C. Huggel, J. Bury, C. Portocarrero, and W. Haeberli, 2012b: An integrated socio-environmental
14 framework for glacier hazard management and climate change adaptation. Lessons from Lake 513, Cordillera
15 Blanca, Peru. *Climatic Change*, **112(3-4)**, 733-767.
- 16 Carilli, J.E., R.D. Norris, B.A. Black, S.M. Walsh, and M. McField, 2009: Local stressors reduce coral resilience to
17 bleaching. *PLoS ONE*, **4(7)**, e6324.
- 18 Carr, E.R., 2008: Between structure and agency. Livelihoods and adaptation in Ghana's central region. *Global*
19 *Environmental Change*, **18(4)**, 689-699.
- 20 Carson, C., S. Hajat, B. Armstrong, and P. Wilkinson, 2006: Declining vulnerability to temperature-related mortality
21 in London over the 20th century. *American Journal of Epidemiology*, **164**, 77-84.
- 22 Carvalho, L.M.V., C. Jones, A.E. Silva, B. Liebmann, and P.L. Silva Dias, 2010: The South American monsoon
23 system and the 1970s climate transition. *International Journal of Climatology*, **31(8)**, 1248-1256.
- 24 Casassa, G.P., B. Pouyaud, and F. Escobar, 2009: Detection of changes in glacial run-off in alpine basins. Examples
25 from North America, the Alps, central Asia and the Andes. *Hydrological Processes*, **23(1)**, 31-41.
- 26 Cashman, A., L. Nurse, and C. John, 2010: Climate change in the Caribbean: The water management implications.
27 *The Journal of Environment & Development*, **19(1)**, 42-67.
- 28 Cattiaux, J., R. Vautard, and P. Yiou, 2009: Origins of the extremely warm European fall of 2006. *Geophysical*
29 *Research Letters*, **36(6)**, L06713.
- 30 CCSP, 2007: *Effects of climate change on energy production and use in the United States*. U.S. Climate Change
31 Science Program and the subcommittee on Global Change Research, pp. 160.
- 32 Chambers, L.E., C.A. Devney, B.C. Congdon, N. Dunlop, E.J. Woehler, and P. Dann, 2011: Observed and predicted
33 effects of climate on Australian seabirds. *Emu*, **111(3)**, 235-251.
- 34 Chambers, L.E., 2008: Trends in timing of migration of south-western Australian birds and their relationship to
35 climate. *Emu*, **108(1)**, 1-14.
- 36 Changnon, S.A., 2007: Catastrophic winter storms. An escalating problem. *Climatic Change*, **84(2)**, 131-139.
- 37 Changnon, S.A., 2008: Assessment of flood losses in the United States. *Journal of Contemporary Water Research &*
38 *Education*, **138(1)**, 38-44.
- 39 Changnon, S.A., 2009a: Characteristics of severe Atlantic hurricanes in the United States: 1949-2006. *Natural*
40 *Hazards*, **48(3)**, 329-337.
- 41 Changnon, S.A., 2009b: Increasing major hail losses in the U.S. *Climatic Change*, **96(1)**, 161-166.
- 42 Chatfield, C., 1995: Model uncertainty, data-mining, and statistical inference. *Journal of the Royal Statistical*
43 *Society*, **A158**, 419-466.
- 44 Chaves, L.F. and C.J.M. Koenraadt, 2010: Climate change and highland malaria. Fresh air for a hot debate. *The*
45 *Quarterly Review of Biology*, **85(1)**, 27-55.
- 46 Chen, C., C. Lei, A. Deng, C. Qian, W. Hoogmoed, and W. Zhang, 2011: Will higher minimum temperatures
47 increase corn production in Northeast China? An analysis of historical data over 1965–2008. *Agricultural and*
48 *Forest Meteorology*, **151(12)**, 1580-1588.
- 49 Chen, C.C. and B.A. McCarl, 2009: Hurricanes and possible intensity increases: Effects on and reactions from U.S.
50 agriculture. *Journal of Agricultural and Applied Economics*, **41(1)**, 125-144.
- 51 Chen, I.C., J.K. Hill, R. Ohlemüller, D.B. Roy, and C.D. Thomas, 2011: Rapid range shifts of species associated
52 with high levels of climate warming. *Science*, **333(6045)**, 1024-1026.
- 53 Chinn, T.B., B. Fitzharris, A. Willsman, and M.J. Salinger, 2012: Annual ice volume changes (1976-2008) for the
54 New Zealand Southern Alps. *Global and Planetary Change*, **92**.

- 1 Christidis, N., G.C. Donaldson, and P.A. Stott, 2010: Causes for the recent changes in cold- and heat-related
2 mortality in England and Wales. *Climatic Change*, **102(3-4)**, 539-553.
- 3 Christie, D.A., J.A. Boninsegna, M.K. Cleaveland, A. Lara, C.L. Quesne, M.S. Morales, M. Mudelsee, D.W. Stahle,
4 and R. Villalba, 2011: Aridity changes in the Temperate-Mediterranean transition of the Andes since ad 1346
5 reconstructed from tree-rings. *Climate Dynamics*, **36(7-8)**, 1505-1521.
- 6 Cogley, J.G., 2012: Climate science: Himalayan glaciers in the balance *Nature*, **488(7412)**, 7412-7469.
- 7 Cole, J.H. and A.M. Gramajo, 2009: Homicide rates in a cross-section of countries: Evidence and interpretations.
8 *Population and Development Review*, **35(4)**, 749-776.
- 9 Coles, S.L. and B.M. Riegl, 2012: Thermal tolerances of reef corals in the Gulf: A review of the potential for
10 increasing coral survival and adaptation to climate change through assisted translocation. *Marine Pollution*
11 *Bulletin*, (in press).
- 12 Collie, J.S., K. Richardson, and J.H. Steele, 2004: Regime shifts: can ecological theory illuminate the mechanisms?
13 *Progress in Oceanography*, **60(2)**, 281-302.
- 14 Collins, D.N., 2006: Climatic variation and runoff in mountain basins with differing proportions of glacier cover.
15 *Nordic Hydrology*, **37(4-5)**, 315-326.
- 16 Comeaux, R.S., M.A. Allison, and T.S. Bianchi, 2011: Mangrove expansion in the Gulf of Mexico with climate
17 change: Implications for wetland health and resistance to rising sea levels. *Estuarine, Coastal and Shelf*
18 *Science*, .
- 19 Confalonieri, U., B. Menne, R. Akhtar, K.L. Ebi, M. Hauengue, R.S. Kovats, B. Revich, and A. Woodward, 2007:
20 *Human health. Climate change 2007. Impacts, adaptation and vulnerability. Contribution of Working Group II*
21 *to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. [Parry, M.L., Canziani,
22 O.F., Palutikof, J.P., van der Linden, P.J. and Hanson, C.E. (eds.)]. Cambridge University Press, Cambridge,
23 pp. 391-431.
- 24 Conway, D. and G. Mahé, 2009: River flow modelling in two large river basins with non-stationary behaviour: the
25 Paraná and the Niger. *Hydrological Processes*, **23(22)**, 3186-3192.
- 26 Cook, B.I., E.M. Wolkovich, and C. Parmesan, 2012: Divergent responses to spring and winter warming drive
27 community level flowering trends. *Proceedings of the National Academy of Sciences of the United States of*
28 *America*, **109(23)**, 9000-9005.
- 29 Cooley, S.R. and S.C. Doney, 2009: Anticipating ocean acidification's economic consequences for commercial
30 fisheries. *Environmental Research Letters*, **4(2)**, 024007.
- 31 Cooper, T.F., G. De'ath, K.E. Fabricius, and J.M. Lough, 2008: Declining coral calcification in massive Porites in
32 two nearshore regions of the northern Great Barrier Reef. *Global Change Biology*, **14(3)**, 529-538.
- 33 Costard, F., E. Gautier, D. Brunstein, J. Hammadi, A. Fedorov, D. Yang, and L. Dupeyrat, 2007: Impact of the
34 global warming on the fluvial thermal erosion over the Lena River in Central Siberia. *Geophysical Research*
35 *Letters*, **34(14)**, L14501.
- 36 Coumou, D. and S. Rahmstorf, 2012: A decade of weather extremes. *Nature Climate Change*, **2(7)**, 491-496.
- 37 Craine, J.M., A.J. Elmore, K.C. Olson, and D. Tolleson, 2010: Climate change and cattle nutritional stress. *Global*
38 *Change Biology*, **16(10)**, 2901-2911.
- 39 Crompton, R.P. and K.J. McAneney, 2008: Normalised Australian insured losses from meteorological hazards:
40 1967-2006. *Environmental Science and Policy*, **11(5)**, 371-378.
- 41 Crompton, R.P., K.J. McAneney, K. Chen, R.A. Pielke Jr, and K. Haynes, 2010: Influence of location, population,
42 and climate on building damage and fatalities due to Australian Bushfire: 1925-2009. *Weather, Climate, and*
43 *Society*, **2(4)**, 300-310.
- 44 Croxall, J., P. Trathan, and E. Murphy, 2002: Environmental change and Antarctic seabird populations. *Science*,
45 **297(5586)**, 1510-1514.
- 46 Cruikshank, J., 2001: Glaciers and climate change. Perspectives from oral tradition. *Arctic*, **54**, 377-393.
- 47 Cullen-Unsworth, L.C., R. Hill, J.R.A. Butler, and M. Wallace, 2011: A research process for integrating indigenous
48 and scientific knowledge in cultural landscapes. Principles and determinants of success in the wet tropics
49 world heritage area, Australia. *The Geographical Journal*, .
- 50 Curry, J., 2011: Nullifying the climate null hypothesis. *Wiley Interdisciplinary Reviews: Climate Change*, **2**, 919-
51 924.
- 52 Cutter, S., B. Osman-Elasha, J. Campbell, S.M. Cheong, S. McCormick, R. Pulwarty, S. Supratid, and G. Ziervogel,
53 2012: Managing the risks from climate extremes at the local level. In: *Managing the Risks of Extreme Events*
54 *and Disasters to Advance Climate Change Adaptation. A Special Report of Working Groups I and II of the*

- 1 *Intergovernmental Panel on Climate Change (IPCC)*. [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J.
2 Dokken, K.L. Ebi *et al.*(eds.)]. Cambridge University Press, Cambridge, UK and New York, USA, pp. 291-
3 338.
- 4 D'Amato, G., L. Cecchi, S. Bonini, C. Nunes, I. Annesi-Maesano, H. Behrendt, G. Liccardi, T. Popov, and P. Van
5 Cauwenberge, 2007: Allergenic pollen and pollen allergy in Europe. *Allergy*, **62(9)**, 976-990.
- 6 Dai, A., K.E. Trenberth, and J.D. Milliman, 2009: Changes in continental freshwater discharge from 1948 to 2004.
7 *Journal of Climate*, **22(10)**, 2773-2792.
- 8 Dai, A., 2011: Drought under global warming: a review. *Wiley Interdisciplinary Reviews-Climate Change*, **2(1)**, 45-
9 65.
- 10 Dash, S. and T. Kjellstrom, 2011: Workplace heat stress in the context of rising temperature in India. *Current*
11 *Science*, **101(4)**, 496.
- 12 Dawe, D., 2008: *Have recent increases in international cereal prices been transmitted to domestic economies? The*
13 *experience in seven large Asian countries* FAO, Rome, .
- 14 De Moel, H., G.M. Ganssen, F.J.C. Peeters, S.J.A. Jung, D. Kroon, G.J.A. Brummer, and R.E. Zeebe, 2009: Planktic
15 foraminiferal shell thinning in the Arabian Sea due to anthropogenic ocean acidification? *Biogeosciences*, **6(9)**,
16 1917-1925.
- 17 De Noblet-Ducoudré, N., J.P. Boisier, A. Pitman, G.B. Bonan, V. Brovkin, F. Cruz, C. Delire, V. Gayler, B.J.J.M.
18 Van den Hurk, P.J. Lawrence, M.K. van der Molen, C. Müller, C.H. Reick, B.J. Strengers, and A. Voldoire,
19 2012: Determining robust impacts of land-use-induced land cover changes on surface climate over North
20 America and Eurasia: results from the first set of LUCID experiments. *Journal of Climate*, **25(9)**, 3261-3281.
- 21 De Young, B. and A. Jarre, 2009: Regime shifts: Methods of analysis. In: *Encyclopedia of Ocean Sciences (Second*
22 *Edition)*. [Steele, J.H., K.K. Turekian, and S.A. Thorpe(eds.)]. Academic Press, Oxford, pp. 717-721.
- 23 De'ath, G., K. Fabricius, H. Sweatman, and M. Puotinen, 2012: The 27 Year Decline of Coral Cover on the Great
24 Barrier Reef and its causes. *Proceedings of the National Academy of Sciences of the United States of America*,
25 **109(44)**, 17995-17999.
- 26 Delaloye, R., C. Lambiel, and I. Gärtner-Roer, 2010: Overview of rock glacier kinematics research in the Swiss
27 Alps. Seasonal rhythm, interannual variations and trends over several decades. *Geographica Helvetica*, **62(2)**,
28 135-145.
- 29 Dell, M., B.F. Jones, and B.A. Olken, 2012: Temperature shocks and economic growth: Evidence from the last half
30 century. *American Economic Journal: Macroeconomics*, **4(3)**, 66-95.
- 31 Dell, M., B.F. Jones, and B.A. Olken, 2009a: Climate Change and Economic Growth: Evidence from the Last Half
32 Century. *American Economic Journal: Macroeconomics*, **2(99)**.
- 33 Dell, M., B.F. Jones, and B.A. Olken, 2009b: Temperature and Income: Reconciling New Cross-Sectional and Panel
34 Estimates. *American Economic Review*, **99(2)**, 198-204.
- 35 Delpla, I., A.V. Jung, E. Baures, M. Clement, and O. Thomas, 2009: Impacts of climate change on surface water
36 quality in relation to drinking water production. *Environment International*, **35(8)**, 1225-1233.
- 37 DeMenocal, P.B., 2001: Cultural responses to climate change during the late Holocene. *Science*, **292(5517)**, 667-
38 673.
- 39 Deschênes, O. and M. Greenstone, 2007: The economic impacts of climate change: Evidence from agricultural
40 output and random fluctuations in weather. *The American Economic Review*, **97(1)**, 354-385.
- 41 Deschênes, O. and M. Greenstone, 2011: Climate change, mortality, and adaptation: Evidence from annual
42 fluctuations in weather in the US. *American Economic Journal: Applied Economics*, **3(4)**, 152-185.
- 43 Descy, J.P. and H. Sarmento, 2008: Microorganisms of the East African Great Lakes and their response to
44 environmental changes. *Freshwater Reviews*, **1**, 59-73.
- 45 deYoung, B., M. Barange, G. Beaugrand, R. Harris, R.I. Perry, M. Scheffer, and F. Werner, 2008: Regime shifts in
46 marine ecosystems: detection, prediction and management. *Trends in Ecology & Evolution*, **23(7)**, 402-409.
- 47 Di Baldassarre, G., A. Montanari, H. Lins, D. Koutsoyiannis, L. Brandimarte, and G. Blöschl, 2010: Flood fatalities
48 in Africa: From diagnosis to mitigation. *Geophysical Research Letters*, **37(22)**, L22402.
- 49 Diaz, R.J. and R. Rosenberg, 2008: Spreading dead zones and consequences for marine ecosystems. *Science*,
50 **321(5891)**, 926-929.
- 51 Diemberger, H., K. Hastrup, S. Schaffer, C.F. Kennel, D. Sneath, M. Bravo, H.F. Graf, J. Hobbs, J. Davis, and M.L.
52 Nodari, 2012: Communicating climate knowledge: Proxies, processes, politics. Commentaries. *Current*
53 *Anthropology*, **53(2)**, 226-244.

- 1 Dietz, A.J., R. Ruben, and A. Verhagen, 2004: *The impact of climate change on drylands: With a focus on West*
2 *Africa*. Springer, .
- 3 Díez, I., N. Muguerza, A. Santolaria, U. Ganzedo, and J. Gorostiaga, 2012: Seaweed assemblage changes in the
4 eastern Cantabrian Sea and their potential relationship to climate change. *Estuarine, Coastal and Shelf*
5 *Science.*, **99**, 108-120.
- 6 Dole, R., M. Hoerling, J. Perlwitz, J. Eischeid, P. Pegion, T. Zhang, X.W. Quan, T. Xu, and D. Murray, 2011: Was
7 there a basis for anticipating the 2010 Russian heat wave? *Geophysical Research Letters*, **38(6)**, L06702.
- 8 Donat, M.G., L.V. Alexander, H. Yang, I. Durre, R. Vose, R.J.H. Dunn, K.M. Willett, E. Aguilar, M. Brunet, J.
9 Caesar, B. Hewitson, C. Jack, A.M.G. Klein Tank, A.C. Kruger, J. Marengo, T.C. Peterson, M. Renom, C.
10 Oria Rojas, M. Rusticucci, J. Salinger, A. Sanhoury Elayah, S.S. Sekele, A.K. Srivastava, B. Trewin, C.
11 Villarroel, L.A. Vincent, P. Zhai, X. Zhang, and S. Kitching, 2013: Updated analyses of temperature and
12 precipitation extreme indices since the beginning of the twentieth century: The HadEX2 dataset. *Journal of*
13 *Geophysical Research: Atmospheres*, **in press**.
- 14 d'Orgeval, T. and J. Polcher, 2008: Impacts of precipitation events and land-use changes on West African river
15 discharges during the years 1951-2000. *Climate Dynamics*, **31(2-3)**, 249-262.
- 16 Douglas, I., K. Alam, M.A. Maghenda, Y. McDonnell, L. McLean, and J. Campbell, 2008: Unjust waters. Climate
17 change, flooding and the urban poor in Africa. *Environment and Urbanization*, **20(1)**, 187-205.
- 18 Doyle, M.E. and V.R. Barros, 2011: Attribution of the river flow growth in the Plata basin. *International Journal of*
19 *Climatology*, **31(15)**, 2234-2248.
- 20 Duarte, C., I. Hendriks, T. Moore, Y. Olsen, A. Steckenbauer, L. Ramajo, J. Carstensen, J. Trotter, and M.
21 McCulloch, 2013: Is ocean acidification an open-ocean syndrome? Understanding anthropogenic impacts on
22 marine pH. *Estuaries and Coasts*, (in press).
- 23 Duarte, C.M., S. Agustí, P. Wassmann, J.M. Arrieta, M. Alcaraz, A. Coello, N. Marbà, I.E. Hendriks, J. Holding,
24 and I. García-Zarandona, 2012a: Tipping elements in the Arctic marine ecosystem. *AMBIO: A Journal of the*
25 *Human Environment*, **41(1)**, 44-55.
- 26 Duarte, C.M., T.M. Lenton, P. Wadhams, and P. Wassmann, 2012b: Abrupt climate change in the Arctic. *Nature*
27 *Climate Change*, **2(2)**, 60-62.
- 28 Dulamsuren, C., M. Hauck, H. Leuschner, and C. Leuschner, 2011: Climate response of tree-ring width in *Larix*
29 *sibirica* growing in the drought-stressed forest-steppe ecotone of northern Mongolia. *Annals of Forest Science*,
30 **68(2)**, 275-282.
- 31 Durand, Y., G. Giraud, M. Laternser, P. Etchevers, L. Mérindol, and B. Lesaffre, 2009: Reanalysis of 47 years of
32 climate in the French Alps (1958-2005): Climatology and trends for snow cover. *Journal of Applied*
33 *Meteorology and Climatology*, **48(12)**, 2487-2512.
- 34 Durmayaz, A. and O.S. Sogut, 2006: Influence of cooling water temperature on the efficiency of a pressurized-water
35 reactor nuclear-power plant. *International Journal of Energy Research*, **30(10)**, 799-810.
- 36 Eakin, C.M., J.A. Morgan, S.F. Heron, T.B. Smith, G. Liu, L. Alvarez-Filip, B. Baca, E. Bartels, C. Bastidas, C.
37 Bouchon, M. Brandt, A.W. Bruckner, L. Bunkley-Williams, A. Cameron, B.D. Causey, M. Chiappone, T.R.L.
38 Christensen, M.J.C. Crabbe, O. Day, E. de la Guardia, G. Diaz-Pulido, D. DiResta, D.L. Gil-Agudelo, D.S.
39 Gilliam, R.N. Ginsburg, S. Gore, H.M. Guzman, J.C. Hendee, E.A. Hernandez-Delgado, E. Husain, C.F.G.
40 Jeffrey, R.J. Jones, E. Jordan-Dahlgren, L.S. Kaufman, D.I. Kline, P.A. Kramer, J.C. Lang, D. Lirman, J.
41 Mallela, C. Manfrino, J. Marechal, K. Marks, J. Mihaly, W.J. Miller, E.M. Mueller, E.M. Muller, C.A. Orozco
42 Toro, H.A. Oxenford, D. Ponce-Taylor, N. Quinn, K.B. Ritchie, S. Rodriguez, A. Rodriguez Ramirez, S.
43 Romano, J.F. Samhoury, J.A. Sanchez, G.P. Schmahl, B.V. Shank, W.J. Skirving, S.C.C. Steiner, E.
44 Villamizar, S.M. Walsh, C. Walter, E. Weil, E.H. Williams, K.W. Roberson, and Y. Yusuf, 2010: Caribbean
45 corals in crisis. Record thermal stress, bleaching, and mortality in 2005. *PLoS ONE*, **5(11)**, e13969.
- 46 Eakin, H., 2006: *Weathering risk in rural Mexico: climatic, institutional, and economic change*. University of
47 Arizona Press, Tucson, .
- 48 Edwards, M. and A.J. Richardson, 2004: Impact of climate change on marine pelagic phenology and trophic
49 mismatch. *Nature*, **430(7002)**, 881-884.
- 50 Eijgelaar, E., C. Thaper, and P. Peeters, 2010: Antarctic cruise tourism: the paradoxes of ambassadorship, 'last
51 chance tourism' and greenhouse gas emissions. *Journal of Sustainable Tourism*, **18(3)**, 337-354.
- 52 Ekau, W., H. Auel, H.O. Pörtner, W. Gilbert, and D. Gilbert, 2010: Impacts of hypoxia on the structure and
53 processes in pelagic communities (zooplankton, macro-invertebrates and fish). *Biogeosciences*, **7(5)**, 1669-
54 1699.

- 1 Eliason, E.J., T.D. Clark, M.J. Hague, L.M. Hanson, Z.S. Gallagher, K.M. Jeffries, M.K. Gale, D.A. Patterson, S.G.
2 Hinch, and A.P. Farrell, 2011: Differences in thermal tolerance among sockeye salmon populations. *Science*,
3 **332(6025)**, 109-112.
- 4 Elliott, J.R. and J. Pais, 2006: Race, class, and Hurricane Katrina: Social differences in human responses to disaster.
5 *Social Science Research*, **35(2)**, 295-321.
- 6 Elmendorf, S.C., G.H.R. Henry, R.D. Hollister, R.G. Björk, N. Boulanger-Lapointe, E.J. Cooper, J.H.C.
7 Cornelissen, T.A. Day, E. Dorrepaal, T.G. Elumeeva, M. Gill, W.A. Gould, J. Harte, D.S. Hik, A. Hofgaard,
8 D.R. Johnson, J.F. Johnstone, I.S. Jónsdóttir, J.C. Jorgenson, K. Klanderud, J.A. Klein, S. Koh, G. Kudo, M.
9 Lara, E. Lévesque, B. Magnússon, J.L. May, J.A. Mercado-Díaz, A. Michelsen, U. Molau, I.H. Myers-Smith,
10 S.F. Oberbauer, V.G. Onipchenko, C. Rixen, N.M. Schmidt, G.R. Shaver, M.J. Spasojevic, T.E.
11 Thórhallsdóttir, A. Tolvanen, T. Troxler, C.E. Tweedie, S. Villareal, C.H. Wahren, X. Walker, P.J. Webber,
12 J.M. Welker, and S. Wipf, 2012: Plot-scale evidence of tundra vegetation change and links to recent summer
13 warming. *Nature Climate Change*, **2(6)**, 453-457.
- 14 Emanuel, K., 2005: Increasing destructiveness of tropical cyclones over the past 30 years. *Nature*, **436**, 686-688.
- 15 Erdem, H.H. and S.H. Sevilgen, 2006: Case study: Effect of ambient temperature on the electricity production and
16 fuel consumption of a simple cycle gas turbine in Turkey. *Applied Thermal Engineering*, **26(2-3)**, 320-326.
- 17 Espinoza, J.C., J. Ronchail, F. Frappart, W. Lavado, W. Santini, and J.L. Guyot, 2013: The 2012 flood in the
18 Amazonas River and tributaries (Western Amazon basin). *Journal of Hydrometeorology*, (in press).
- 19 Etter, A., C. McAlpine, S. Phinn, D. Pullar, and H. Possingham, 2006: Unplanned land clearing of Colombian
20 rainforests: Spreading like disease? *Landscape and Urban Planning*, **77(3)**, 240-254.
- 21 Evans, S.G. and J.J. Clague, 1994: Recent climatic change and catastrophic geomorphic processes in mountain
22 environments. *Geomorphology*, **10(1-4)**, 107-128.
- 23 Fajnzylber, P., D. Lederman, and N. Loayza, 2002: What causes violent crime? *European Economic Review*, **46(7)**,
24 1323-1357.
- 25 Farbotko, C., 2010: 'The global warming clock is ticking so see these places while you can': Voyeuristic tourism
26 and model environmental citizens on Tuvalu's disappearing islands. *Singapore Journal of Tropical Geography*,
27 **31(2)**, 224-238.
- 28 Feely, R.A., S.R. Alin, J. Newton, C.L. Sabine, M. Warner, A. Devol, C. Krembs, and C. Maloy, 2010: The
29 combined effects of ocean acidification, mixing, and respiration on pH and carbonate saturation in an
30 urbanized estuary. *Estuarine, Coastal and Shelf Science*, **88(4)**, 442-449.
- 31 Feng, S., A.B. Krueger, and M. Oppenheimer, 2010: Linkages among climate change, crop yields and Mexico-US
32 cross-border migration. *Proceedings of the National Academy of Sciences of the United States of America*,
33 **107(32)**, 14257-14262.
- 34 Feng, S., M. Oppenheimer, and W. Schlenker, 2012: *Climate change, crop yields, and internal migration in the*
35 *United States*. National Bureau of Economic Research.
- 36 Fernández, C., 2011: The retreat of large brown seaweeds on the north coast of Spain: the case of *Saccorhiza*
37 *polyschides*. *European Journal of Phycology*, **46(4)**, 352-360.
- 38 Field, S., 1992: The effect of temperature on crime. *British Journal of Criminology*, **32(3)**, 340-351.
- 39 Figueira, W.F., P. Biro, D.J. Booth, and V.C. Valenzuela, 2009: Performance of tropical fish recruiting to temperate
40 habitats: role of ambient temperature and implications of climate change. *Marine Ecology-Progress Series*,
41 **384**, 231-239.
- 42 Fischer, L., R.S. Purves, C. Huggel, J. Noetzli, and W. Haeberli, 2012: On the influence of geological, topographic
43 and glaciological factors on slope instabilities. Analyses of recent Alpine rock avalanches. *Natural Hazards*
44 *and Earth System Sciences*, **12**, 241-254.
- 45 Fischlin, A., G.F. Midgley, J.T. Price, R. Leemans, B. Gopal, C. Turley, M.D.A. Rounsevell, O.P. Dube, J.
46 Tarazona, and A.A. Velichko, 2007: Ecosystems, their properties, goods, and services. Climate Change 2007:
47 Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of
48 the Intergovernmental Panel on Climate Change. In: *Fourth Assessment Report of the Intergovernmental Panel*
49 *on Climate Change*. [Parry, M.L., O.F. Canziani, J.P. Palutikof, P.J. van der Linden, and C.E. Hanson(eds.)].
50 Cambridge University Press, Cambridge, pp. 211-272.
- 51 Fishman, J., J.K. Creilson, P.A. Parker, E.A. Ainsworth, G.G. Vining, J. Szarka, L. Booker, and X. Xu, 2010: An
52 investigation of widespread ozone damage to the soybean crop in the upper Midwest determined from ground-
53 based and satellite measurements. *Atmospheric Environment*, **44(18)**, 2248-2256.

- 1 Foden, W., G.F. Midgley, G. Hughes, W.J. Bond, W. Thuiller, M.T. Hoffmann, P. Kaleme, L.G. Underhill, A.
2 Rebelo, and L. Hannah, 2007: A changing climate is eroding the geographical range of the Namib Desert tree
3 Aloe through population declines and dispersal lags. *Diversity and Distributions*, **13**, 645-653.
- 4 Forbes, D.L. (ed.), 2011: *State of the Arctic Coast 2010 – Scientific Review and Outlook*. International Arctic
5 Science Committee, Land-Ocean Interactions in the Coastal Zone, Arctic Monitoring and Assessment
6 Programme, International Permafrost Association [IASC, LOICZ, AMAP, IPA], Helmholtz-Zentrum,
7 Geesthacht, Germany, pp. 178.
- 8 Forcada, J., P.N. Trathan, and E.J. Murphy, 2008: Life history buffering in Antarctic mammals and birds against
9 changing patterns of climate and environmental variation. *Global Change Biology*, **14(11)**, 2473-2488.
- 10 Ford, J.D., 2009a: Vulnerability of Inuit food systems to food insecurity as a consequence of climate change: a case
11 study from Igloolik, Nunavut. *Regional Environmental Change*, **9(2)**, 83-100.
- 12 Ford, J.D., 2009b: Dangerous climate change and the importance of adaptation for the Arctic's Inuit population.
13 *Environmental Research Letters*, **4(2)**, 024006.
- 14 Ford, J.D., W.A. Gough, G.J. Laidler, J. MacDonald, C. Irngaut, and K. Qrunnut, 2009: Sea ice, climate change, and
15 community vulnerability in northern Foxe Basin, Canada. *Climate Research*, **38**, 137-154.
- 16 Ford, J.D., W. Vanderbilt, and L. Berrang-Ford, 2011: Authorship in IPCC AR5 and its implications for content.
17 Climate change and Indigenous populations in WGII. *Climatic Change*, **113(2)**, 201-213.
- 18 Ford, M., 2012: Shoreline changes on an urban atoll in the central Pacific Ocean. Majuro atoll, Marshall Islands.
19 *Journal of Coastal Research*, **28(1)**, 11-22.
- 20 Forkel, M., K. Thonicke, C. Beer, W. Cramer, S. Bartalev, and C. Schmullius, 2012: Extreme fire events are related
21 to previous-year surface moisture conditions in permafrost-underlain larch forests of Siberia. *Environmental*
22 *Research Letters*, **7(4)**, 044021.
- 23 Fortner, S.K., B.G. Mark, J.M. McKenzie, J. Bury, A. Trierweiler, M. Baraer, P.J. Burns, and L. Munk, 2011:
24 Elevated stream trace and minor element concentrations in the foreland of receding tropical glaciers. *Applied*
25 *Geochemistry*, **26**, 1792-1801.
- 26 Fouillet, A., G. Rey, V. Wagner, K. Laaidi, P. Empereur-Bissonnet, A. le Tertre, P. Frayssinet, P. Bessemoulin, F.
27 Laurent, P. Crouy-Chanel, E. Jouglu, and D. Hémon, 2008: Has the impact of heat waves on mortality changed
28 in France since the European heat wave of summer 2003? A study of the 2006 heat wave. *International*
29 *Journal of Epidemiology*, **37(2)**, 309-317.
- 30 Fowbert, J.A. and R.I.L. Smith, 1994: Rapid population increases in native vascular plants in the Argentine Islands,
31 Antarctic Peninsula. *Arctic and Alpine Research*, **26**, 290-296.
- 32 Franco, G. and A.H. Sanstad, 2008: Climate change and electricity demand in California. *Climatic Change*, **87(1)**,
33 139-151.
- 34 Freire, K.M.F. and D. Pauly, 2010: Fishing down Brazilian marine food webs, with emphasis on the east Brazil large
35 marine ecosystem. *Fisheries Research*, **105(1)**, 57-62.
- 36 Galarneau, T.J., T.M. Hamill, R.M. Dole, and J. Perlwitz, 2012: A multiscale analysis of the extreme weather events
37 over western Russia and northern Pakistan during July 2010. *Monthly Weather Review*, **140(5)**, 1639-1664.
- 38 Gamble, D.W., D. Campbell, T.L. Allen, D. Barker, S. Curtis, D. McGregor, and J. Popke, 2010: Climate change,
39 drought, and Jamaican agriculture: local knowledge and the climate record. *Annals of the Association of*
40 *American Geographers*, **100(4)**, 880-893.
- 41 Gamble, J.L. and J.J. Hess, 2012: Temperature and violent crime in Dallas, Texas: relationships and implications of
42 climate change. *Western Journal of Emergency Medicine*, **13(3)**, 239-246.
- 43 García-Mozo, H., C. Galán, P. Alcázar, C. de la Guardia, D. Nieto-Lugilde, M. Recio, P. Hidalgo, F. González-
44 Minero, L. Ruiz, and E. Domínguez-Vilches, 2010: Trends in grass pollen season in southern Spain.
45 *Aerobiologia*, **26(2)**, 157-169.
- 46 Gardelle, J., Y. Arnaud, and E. Berthier, 2011: Contrasted evolution of glacial lakes along the Hindu Kush Himalaya
47 mountain range between 1990 and 2009. *Global and Planetary Change*, **75**, 47-55.
- 48 Gardner, T.A., I.M. Côté, J.A. Gill, A. Grant, and A.R. Watkinson, 2003: Long-term region-wide declines in
49 Caribbean corals. *Science*, **301(5635)**, 958-960.
- 50 Gardner, A.S., G. Moholdt, J.G. Cogley, A.A. Arendt, B. Wouters, J. Wahr, W.T. Pfeffer, G. Kaser, S.R.M.
51 Ligtenberg, T. Bolch, R. Hock, E. Berthier, M.J. Sharp, J.O. Hagen, M. van den Broeke, F. Paul, and V. Radic,
52 2013: A consensus estimate of glacier contributions to sea level rise: 2003 to 2009. *Science*, (in press).
- 53 Garrabou, J., R. Coma, N. Bensoussan, M. Bally, P. Chevaldonné, M. Cigliano, D. Diaz, J.G. Harmelin, M.C.
54 Gambi, D.K. Kersting, J.B. Ledoux, C. Lejesune, C. Linares, C. Marschal, T. Pérez, M. Ribes, J.C. Romano,

- 1 E. Serrano, N. Teixido, O. Torrents, M. Zabala, F. Zuberer, and C. Cerrano, 2009: Mass mortality in
2 Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave. *Global Change*
3 *Biology*, **15(5)**, 1090-1103.
- 4 Gaston, A.J., H.G. Gilchrist, M.L. Mallory, and P.A. Smith, 2009: Changes in seasonal events, peak food
5 availability, and consequent breeding adjustment in a marine bird: a case of progressive mismatching. *The*
6 *Condor*, **111(1)**, 111-119.
- 7 Gearheard, S., M. Pocernich, R. Stewart, J. Sanguya, and H.P. Huntington, 2010: Linking Inuit knowledge and
8 meteorological station observations to understand changing wind patterns at Clyde River, Nunavut. *Climatic*
9 *Change*, **100**, 267-294.
- 10 Gehrig-Fasel, J., A. Guisan, and N.E. Zimmermann, 2007: Tree line shifts in the Swiss Alps: Climate change or land
11 abandonment? *Journal of Vegetation Science*, **18(4)**, 571-582.
- 12 Georgakakos, A., P. Fleming, M. Dettinger, C. Peters-Lidgard, T.C. Richmond, K. Reckhow, K. White, and D.
13 Yates, 2013: *Federal Advisory Committee Draft Climate Assessment Report Released for Public Review.*
14 *Chapter 3: Water Resources. Available: <http://ncadac.globalchange.gov/>.*
- 15 Gething, P.W., D.L. Smith, A.P. Patil, A.J. Tatem, R.W. Snow, and S.I. Hay, 2010: Climate change and the global
16 malaria recession. *Nature*, **465**, 342-345.
- 17 Giannini, A., M. Biasutti, and M.M. Verstraete, 2008: A climate model-based review of drought in the Sahel.
18 Desertification, the re-greening and climate change. *Global and Planetary Change*, **64**, 119-128.
- 19 Gilman, E.L., J. Ellison, N.C. Duke, and C. Field, 2008: Threats to mangroves from climate change and adaptation
20 options: a review. *Aquatic Botany*, **89(2)**, 237-250.
- 21 Giri, C., E. Ochieng, L.L. Tieszen, Z. Zhu, A. Singh, T. Loveland, J. Masek, and N. Duke, 2011: Status and
22 distribution of mangrove forests of the world using earth observation satellite data. *Global Ecology and*
23 *Biogeography*, **20(1)**, 154-159.
- 24 Gonzalez, P., C.J. Tucker, and H. Sy, 2012: Tree density and species decline in the African Sahel attributable to
25 climate. *Journal of Arid Environments*, **78**, 55-64.
- 26 Goodman, J., J. Maschinski, P. Hughes, J. Mcauliffe, J. Roncal, D. Powell, and L.O. Sternberg, 2012: Differential
27 response to soil salinity in endangered key tree cactus. Implications for survival in a changing climate. *PLoS*
28 *ONE*, **7(3)**, e32528.
- 29 Gooseff, M.N., A. Balsler, W.B. Bowden, and J.B. Jones, 2009: Spatio- temporal evolution of a thermokarst in
30 Interior Alaska. *EOS Transactions of the American Geophysical Union*, **90(4)**, 29-30.
- 31 Gray, J.S., H. Dautel, A. Estrada-Peña, O. Kahl, and E. Lindgren, 2009: Effects of climate change on ticks and tick-
32 borne diseases in Europe. *Interdisciplinary Perspectives on Infectious Diseases*, **2009**.
- 33 Green, D. and G. Raygorodetsky, 2010: Indigenous knowledge of a changing climate. *Climatic Change*, **100**, 239-
34 242.
- 35 Green, K., 2010: Alpine taxa exhibit differing responses to climate warming in the Snowy Mountains of Australia.
36 *Journal of Mountain Science*, **7**, 167-175.
- 37 Greene, W.H., 2003: *Econometric analysis*. Prentice Hall, Upper Saddle River, NJ, USA, pp. 1026.
- 38 Greene, A.M., A. Giannini, and S.E. Zebiak, 2009: Drought return times in the Sahel: a question of attribution.
39 *Geophysical Research Letters*, **36(12)**, L12701.
- 40 Gregory, P.J. and B. Marshall, 2012: Attribution of climate change: a methodology to estimate the potential
41 contribution to increases in potato yield in Scotland since 1960. *Global Change Biology*, **18(4)**, 1372-1388.
- 42 Grémillet, D. and T. Boulinier, 2009: Spatial ecology and conservation of seabirds facing global climate change. A
43 review. *Marine Ecology Progress Series*, **391**, 121-137.
- 44 Grevelle, G. and N. Mimura, 2008: Vulnerability assessment of sea-level rise in Viti Levu, Fiji Islands.
45 *Sustainability Science*, **3**, 171-180.
- 46 Guzman, H.M., S. Benfield, O. Breedy, and J.M. Mair, 2008: Broadening reef protection across the Marine
47 Conservation Corridor of the Eastern Tropical Pacific: Distribution and diversity of reefs in Las Perlas
48 Archipelago, Panama. *Environmental Conservation*, **35(1)**, 46-54.
- 49 Ha, J., H. Kim, and S. Hajat, 2011: Effect of previous-winter mortality on the association between summer
50 temperature and mortality in South Korea. *Environmental Health Perspectives*, **119**, 542-546.
- 51 Haeberli, W. and R. Hohmann, 2008: Climate, glaciers and permafrost in the Swiss Alps 2050: scenarios,
52 consequences and recommendations. In: *Proceedings Ninth International Conference on Permafrost* pp. 607-
53 612.

- 1 Haeberli, W. and M. Beniston, 1998: Climate change and its impacts on glaciers and permafrost in the Alps. *Ambio*,
2 **27(4)**, 258-265.
- 3 Haigh, I., M. Eliot, C. Pattiaratchi, and T. Wahl, 2011: Regional changes in mean sea level around Western
4 Australia between 1897 and 2008. Proceedings of Coasts and Ports 2011: Diverse and Developing: Proceedings
5 of the 20th Australasian Coastal and Ocean Engineering Conference and the 13th Australasian Port and
6 Harbour Conference, pp. 280.
- 7 Haigh, I., R. Nicholls, and N. Wells, 2010: Assessing changes in extreme sea levels: application to the English
8 Channel, 1900–2006. *Continental Shelf Research*, **30(9)**, 1042-1055.
- 9 Halpern, B.S., S. Walbridge, K.A. Selkoe, C.V. Kappel, F. Micheli, C. D'agrosa, J.F. Bruno, K.S. Casey, C. Ebert,
10 H.E. Fox, R. Fujita, D. Heinemann, H.S. Lenihan, E.M.P. Madin, M.T. Perry, E.R. Selig, M. Spalding, R.
11 Steneck, and R. Watson, 2008: A global map of human impact on marine ecosystems. *Science*, **319(5865)**,
12 948-952.
- 13 Hampel, A., R. Hetzel, and G. Maniatis, 2010: Response of faults to climate-driven changes in ice and water
14 volumes on Earth's surface. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and*
15 *Engineering Sciences*, **368(1919)**, 2501-2517.
- 16 Handmer, J., Y. Honda, Z.W. Kundzewicz, N. Arnell, G. Benito, J. Hatfield, I.F. Mohamed, P. Peduzzi, S. Wu, B.
17 Sherstyukov, K. Takahashi, and Z. Yan, 2012: *Changes in impacts of climate extremes. Human systems and*
18 *ecosystems. In: Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation.*
19 In: A Special Report of Working Groups I and II of the Intergovernmental Panel on Climate Change (IPCC)
20 [Field, C.B., Barros, V., Stocker, T.F., Qin, D., Dokken, D.J., Ebi, K.L., Mastrandrea, M.D., Mach, K.J.,
21 Plattner, G.K., Allen, S.K., Tignor, M. and Midgley, P.M. (eds.)]. Cambridge University Press, Cambridge and
22 New York, pp. 231-290.
- 23 Hansen, B.B., V. Grøtán, R. Aanes, B.E. Sæther, A. Stien, E. Fuglei, R.A. Ims, N.G. Yoccoz, and Å.Ø. Pedersen,
24 2013: Climate events synchronize the dynamics of a resident vertebrate community in the high Arctic. *Science*,
25 **339(6117)**, 313-315.
- 26 Hardoy, J. and G. Pandiella, 2009: Urban poverty and vulnerability to climate change in Latin America.
27 *Environment and Urbanization*, **21(1)**, 203-224.
- 28 Harley, C.D., 2011: Climate change, keystone predation, and biodiversity loss. *Science*, **334(6059)**, 1124-1127.
- 29 Hatfield, J.K., K.J. Boote, B.A. Kimball, D.W. Wolfe, D.R. Ort, R.C. Izaurralde, and G.L. Hahn, 2008: The effects
30 of climate change on agriculture, land resources, water resources, and biodiversity in the United States. In:
31 *Synthesis and Assessment Product 4.3. Agriculture, US Climate Change Science Program, Washington DC,*
32 *USA, pp. 21-74.*
- 33 Hawkins, S.J., P.J. Moore, M.T. Burrows, E. Poloczanska, N. Mieszkowska, R.J.H. Herbert, S.R. Jenkins, R.C.
34 Thompson, M.J. Genner, and A.J. Southward, 2008: Complex interactions in a rapidly changing world:
35 responses of rocky shore communities to recent climate change. *Climate Research*, **37**, 123-133.
- 36 Hayes, J. and A. Goonetilleke, 2012: Building community resilience - learning from the 2011 floods in Southeast
37 Queensland, Australia. In: *8th Annual Conference of International Institute for Infrastructure, Renewal and*
38 *Reconstruction: International Conference on Disaster Management (IIIRR 2012)* [Kakimoto, R. and F.
39 Yamada(eds.)]. Kumamoto University, Kumamoto, Japan, pp. 51-60.
- 40 Hayhoe, K., D. Cayan, C.B. Field, P.C. Frumhoff, E.P. Maurer, N.L. Miller, S.C. Moser, S.H. Schneider, K.N.
41 Cahill, E.E. Cleland, L. Dale, R. Drapek, R.M. Hanemann, L.S. Kalkstein, J. Lenihan, C.K. Lunch, R.P.
42 Neilson, S.C. Sheridan, and J.H. Verville, 2004: Emissions pathways, climate change, and impacts on
43 California. *Proceedings of the National Academy of Sciences of the United States of America*, **101(34)**, 12422-
44 12427.
- 45 Hecky, R.E., R. Mugidde, P.S. Ramlal, M.R. Talbot, and G.W. Kling, 2010: Multiple stressors cause rapid
46 ecosystem change in Lake Victoria. *Freshwater Reviews*, **55**, 19-42.
- 47 Hedenås, H., H. Olsson, C. Jonasson, J. Bergstedt, U. Dahlberg, and T.V. Callaghan, 2011: Changes in tree growth,
48 biomass and vegetation over a 13-year period in the Swedish sub-Arctic. *AMBIO: A Journal of the Human*
49 *Environment*, **40(6)**, 672-682.
- 50 Hegerl, G.C., O. Hoegh-Guldberg, G. Casassa, M.P. Hoerling, R.S. Kovats, C. Parmesan, D.W. Pierce, and P.A.
51 Stott, 2010: Good practice guidance paper on detection and attribution related to anthropogenic climate
52 change. In: *Meeting Report of the Intergovernmental Panel on Climate Change Expert Meeting on Detection*
53 *and Attribution of Anthropogenic Climate Change.* [Stocker, T.F., C.B. Field, D. Qin, V. Barros, G.K. Plattner,

- 1 M. Tignor *et al.*(eds.)]. IPCC Working Group I Technical Support Unit, University of Bern, Bern, Switzerland,
2 .
- 3 Helmuth, B., C.D. Harley, P.M. Halpin, M. O'Donnell, G.E. Hofmann, and C.A. Blanchette, 2002: Climate change
4 and latitudinal patterns of intertidal thermal stress. *Science*, **298(5595)**, 1015-1017.
- 5 Helmuth, B., N. Mieszkowska, P. Moore, and S.J. Hawkins, 2006: Living on the edge of two changing worlds:
6 forecasting the responses of rocky intertidal ecosystems to climate change. *Annual Review of Ecology,
7 Evolution, and Systematics*, **37**, 373-404.
- 8 Hemp, A., 2005: Climate change-driven forest fires marginalize the impact of ice cap wasting on Kilimanjaro.
9 *Global Change Biology*, **11**, 1013-1023.
- 10 Hendrix, C.S. and I. Salehyan, 2012: Climate change, rainfall, and social conflict in Africa. *Journal of Peace
11 Research*, **49**, 35-50.
- 12 Hendrix, C.S. and S.M. Glaser, 2007: Trends and triggers: Climate, climate change and civil conflict in Sub-Saharan
13 Africa. *Political Geography*, **26(6)**, 695-715.
- 14 Hennessy, K.J., P.H. Whetton, K. Walsh, I.N. Smith, J.M. Bathols, M. Hutchinson, and J. Sharples, 2008: Climate
15 change effects on snow conditions in mainland Australia and adaptation at ski resorts through snowmaking.
16 *Climate Research*, **35(3)**, 255-270.
- 17 Henry, G.H.R. and S. Elmendorf, 2010: *Greening of the Arctic*. In: Arctic Biodiversity Trends 2010 - Selected
18 Indicators of Change. CAFF International Secretariat, Akureyri, Iceland, pp. 62-64.
- 19 Henry, S., B. Schoumaker, and C. Beauchemin, 2004: The impact of rainfall on the first out-migration. A multi-
20 level event history analysis on Burkina Faso. *Population and Environment*, **25**, 423-460.
- 21 Herman-Mercer, N., P.F. Schuster, and K.B. Maracle, 2011: Indigenous observations of climate change in the
22 Lower Yukon River Basin, Alaska. *Human Organization*, **70**, 244-252.
- 23 Hiddink, J.G. and R. ter Hofstede, 2008: Climate induced increases in species richness of marine fishes. *Global
24 Change Biology*, **14(3)**, 453-460.
- 25 Hilker, N., A. Badoux, and C. Hegg, 2009: The Swiss flood and landslide damage database 1972-2007. *Natural
26 Hazards and Earth System Sciences*, **9(3)**, 913-925.
- 27 Hockey, P.A.R. and G.F. Midgley, 2009: Avian range changes and climate change. A cautionary tale from the Cape
28 Peninsula. *Ostrich*, **80**, 29-34.
- 29 Hockey, P.A.R., C. Sirami, A.R. Ridley, G.F. Midgley, and H.A. Babiker, 2011: Interrogating recent range changes
30 in South African birds. Confounding signals from land use and climate change present a challenge for
31 attribution. *Diversity and Distributions*, **17**, 254-261.
- 32 Hoegh-Guldberg, O., 1999: Climate change, coral bleaching and the future of the world's coral reefs. *Marine and
33 Freshwater Research*, **50(8)**, 839-866.
- 34 Hoegh-Guldberg, O., 2011: Coral reef ecosystems and anthropogenic climate change. *Regional Environmental
35 Change*, **11(Suppl.1)**, 215-227.
- 36 Hoegh-Guldberg, O., G. Hegerl, T. Root, F. Zwiers, P. Stott, D. Pierce, and M. Allen, 2011: Difficult but not
37 impossible. *Nature Climate Change*, **1(2)**, 72-72.
- 38 Hoegh-Guldberg, O. and G.J. Smith, 1989: The effect of sudden changes in temperature, light and salinity on the
39 population-density and export of zooxanthellae from the reef corals *stylophora-pistillata* esper and *seriatopora-
40 hystrix dana*. *Journal of Experimental Marine Biology and Ecology*, **129(3)**, 279-303.
- 41 Hoegh-Guldberg, O., P.J. Mumby, A.J. Hooten, R.S. Steneck, P. Greenfield, E. Gomez, C.D. Harvell, P.F. Sale, A.J.
42 Edwards, K. Caldeira, N. Knowlton, C.M. Eakin, R. Iglesias-Prieto, N. Muthiga, R.H. Bradbury, A. Dubi, and
43 M.E. Hatziolos, 2007: Coral reefs under rapid climate change and ocean acidification. *Science*, **318(5857)**,
44 1737-1742.
- 45 Hoegh-Guldberg, O. and J.F. Bruno, 2010: The impact of climate change on the world's marine ecosystems.
46 *Science*, **328(5985)**, 1523-1528.
- 47 Hoegh-Guldberg, O., 2012: The adaptation of coral reefs to climate change: Is the Red Queen being outpaced?
48 *Scientia Marina*, **76(2)**, 403-408.
- 49 Hoerling, M., J. Hurrell, J. Eischeid, and A. Phillips, 2006: Detection and attribution of twentieth-century northern
50 and southern African rainfall change. *Journal of Climate*, **19(16)**, 3989-4008.
- 51 Hoinka, K.P., A. Carvalho, and A.I. Miranda, 2009: Regional-scale weather patterns and wildland fires in central
52 Portugal. *International Journal of Wildland Fire*, **18(1)**, 36-49.
- 53 Holland, P.W., 1986: Statistics and causal inference. *Journal of the American Statistical Association*, **81**, 945-960.

- 1 Hollowed, A.B., S.R. Hare, and W.S. Wooster, 2001: Pacific Basin climate variability and patterns of Northeast
2 Pacific marine fish production. *Progress in Oceanography*, **49**, 257-282.
- 3 Hong, C., H. Hsu, N. Lin, and H. Chiu, 2011: Roles of European blocking and tropical-extratropical interaction in
4 the 2010 Pakistan flooding. *Geophysical Research Letters*, **38(13)**, L13806.
- 5 Houze Jr, R.A., K.L. Rasmussen, S. Medina, S.R. Brodzik, and U. Romatschke, 2011: Anomalous atmospheric
6 events leading to the Summer 2010 floods in Pakistan. *Bulletin of the American Meteorological Society*, **92(3)**,
7 291-298.
- 8 Hoyos, L.E., A.M. Cingolani, M.R. Zak, M.V. Vaieretti, D.E. Gorla, and M.R. Cabido, 2012: Deforestation and
9 precipitation patterns in the arid Chaco forests of central Argentina. *Applied Vegetation Science*, **2012**.
- 10 Hsiang, S.M., K.C. Meng, and M.A. Cane, 2011: Civil conflicts are associated with the global climate. *Nature*,
11 **476(7361)**, 438-441.
- 12 Hsieh, C., C.S. Reiss, R.P. Hewitt, and G. Sugihara, 2008: Spatial analysis shows that fishing enhances the climatic
13 sensitivity of marine fishes. *Canadian Journal of Fisheries and Aquatic Sciences*, **65**, 947-961.
- 14 Hsieh, C.H., C. Reiss, W. Watson, M.J. Allen, J.R. Hunter, R.N. Lea, R.H. Rosenblatt, P.E. Smith, and G. Sugihara,
15 2005: A comparison of long-term trends and variability in populations of larvae of exploited and unexploited
16 fishes in the Southern California region. A community approach. *Progress in Oceanography*, **67(1)**, 160-185.
- 17 Huang, X., M. Sillanpaa, E.T. Gjeesing, and P.R. Vogt, 2009: Water quality in the Tibetan Plateau: Major ions and
18 trace elements in the headwaters of four major Asian rivers. *Science of the Total Environment*, **407(24)**, 6242-
19 6254.
- 20 Huggel, C., A. Raissig, M. Rohrer, G. Romero, A. Díaz, and N. Salzmann, Spatio-temporal changes and impacts of
21 weather- and climate related disasters in the Andes of southern Peru. (submitted manuscript).
- 22 Huggel, C., L. Blaškovičová, H. Breien, P. Dobesberger, R. Frauenfelder, B.G. Kalsnes, A. Solheim, M.
23 Stankoviansky, K. Hagen, and K. Kronholm, 2011: Landslide and rock slope failures. In: *Impacts of climate
24 change on snow, ice, and permafrost in Europe: Observed trends, future projections, and socio-economic
25 relevance*. [Voigt, T., H.M. Füssel, I. Gärtner-Roer, C. Huggel, C. Marty, and M. Zemp(eds.)]. European
26 Environment Agency, Copenhagen, pp. 95-101.
- 27 Huggel, C., S. Allen, P. Deline, L. Fischer, J. Noetzli, and L. Raveland, 2012a: Ice thawing, mountains falling - Are
28 alpine rock slope failures increasing? *Geology Today*, **28(3)**, 102-108.
- 29 Huggel, C., J.J. Clague, and O. Korup, 2012b: Is climate change responsible for changing landslide activity in high
30 mountains? *Earth Surface Processes and Landforms*, **37**, 77-91.
- 31 Hulme, M., 2008: Geographical work at the boundaries of climate change. *Transactions of the Institute of British
32 Geographers*, **33(1)**, 5-11.
- 33 Hulme, M., S.J. O'Neill, and S. Dessai, 2011: Is weather event attribution necessary for adaptation funding? *Science*,
34 **334**, 764-765.
- 35 Hunt, A. and P. Watkiss, 2011: Climate change impacts and adaptation in cities: a review of the literature *Climatic
36 Change*, **104(1)**, 13-49.
- 37 Hunt, G.L., 2010: Bering Sea. In: *Marine Ecosystems of the North Pacific Ocean, 2003-2008. PICES Special
38 Publication 4*. [S. M. McKinnell and M. J. Dagg(eds.)]. North Pacific Marine Science Organization, Sidney,
39 B.C., Canada, pp. 196-267.
- 40 Huntington, H., T. Callaghan, S. Fox, and I. Krupnik, 2004: Matching traditional and scientific observations to
41 detect environmental change: A discussion on Arctic terrestrial ecosystems. *Ambio*, **13**, 18-23.
- 42 Huss, M., 2011: Present and future contribution of glacier storage change to runoff from macroscale drainage basins
43 in Europe. *Water Resources Research*, **47**, W07511.
- 44 IPCC, 2001: *Climate Change 2001: Impacts, Adaptation, and Vulnerability: Contribution of Working Group II to
45 the Third Assessment Report to the International Panel on Climate Change*. [McCarthy, J.J., Canziani, O.F.,
46 Leary, N.A., Dokken, D.J. and White, K.S. (eds.)]. Cambridge University Press, Cambridge, pp. 1032.
- 47 IPCC, 2007a: Summary for Policymakers. In: *Climate Change 2007: Synthesis Report. Contribution of Working
48 Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Core
49 Writing Team, R.K. Pachauri, and A. Reisinger(eds.)]. IPCC, Geneva, Switzerland, pp. 104.
- 50 IPCC, 2007b: Summary for Policymakers. In: *Climate Change 2007: Impacts, Adaptation and Vulnerability.
51 Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate
52 Change*. [Parry, M.L., O.F. Canziani, J.P. Palutikof, P.J. Van der Linden, and C.E. Hanson(eds.)]. Cambridge
53 University Press, Cambridge, UK, pp. 7-22.

- 1 IPCC, 2012: Managing the risk of extreme events and disasters to advance climate change adaptation. In: *A special*
2 *report of Working Groups I and II of the Intergovernmental Panel on Climate Change*. [Field, C.B., V. Barros,
3 T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi, M.D. Mastrandrea, K.J. Mach, G.K. Plattner, S.K. Allen, M.
4 Tignor, and P.M. Midgley (ed.)]. Cambridge University Press, Cambridge, UK, and New York, USA, pp. 582.
5 Isaac, M. and D.P. van Vuuren, 2009: Modeling global residential sector energy demand for heating and air
6 conditioning in the context of climate change. *Energy Policy*, **37(2)**, 507-521.
7 ISDR, 2009: null. In: *Risk and poverty in a changing climate. Global Assessment Report on Disaster Risk Reduction*
8 UNISDR, pp. 207.
9 Jaggard, K., A. Qi, and M. Semenov, 2007: The impact of climate change on sugarbeet yield in the UK: 1976–2004.
10 *The Journal of Agricultural Science*, **145(4)**, 367-375.
11 Jeelani, G., 2008: Aquifer response to regional climate variability in a part of Kashmir Himalaya in India.
12 *Hydrogeology Journal*, **16(8)**, 1625-1633.
13 Jenkins, G.S., A.T. Gaye, and B. Sylla, 2005: Late 20th century attribution of drying trends in the Sahel from the
14 Regional Climate Model (RegCM3). *Geophysical Research Letters*, **32(22)**, L22705.
15 Jenouvrier, S., H. Weimerskirch, C. Barbraud, and Y.H. Park, 2005: Evidence of a shift in the cyclicity of Antarctic
16 seabird dynamics linked to climate. *Proceedings of the Royal Society B: Biological Sciences*, **272(1566)**, 887-
17 895.
18 Jentes, E.S., G. Poumerol, M.D. Gershman, D.R. Hill, J. Lemarchand, R.F. Lewis, J.E. Staples, O. Tomori, A.
19 Wilder-Smith, and T.P. Monath, 2011: The revised global yellow fever risk map and recommendations for
20 vaccination, 2010: consensus of the Informal WHO Working Group on Geographic Risk for Yellow Fever.
21 *The Lancet Infectious Diseases*, **11(8)**, 622-632.
22 Jeong, S., C.H. Ho, H.J. Gim, and M. Brown, 2011: Phenology shifts at start vs. end of growing season in temperate
23 vegetation over the Northern Hemisphere for the period 1982-2008. *Global Change Biology*, **17**, 2385-2399.
24 Jiang, Y., T.H. Dixon, and S. Wdowinski, 2010: Accelerating uplift in the North Atlantic region as an indicator of
25 ice loss. *Nature Geoscience*, **3(6)**, 404-407.
26 Jin, M., C. Deal, J. Wang, and C.P. McRoy, 2009: Response of lower trophic level production to long-term climate
27 change in the southeastern Bering Sea. *Journal of Geophysical Research*, **114(C4)**, C04010.
28 Johnson, C.R., S.C. Banks, N.S. Barrett, F. Cazassus, P.K. Dunstan, G.J. Edgar, S.D. Frusher, C. Gardner, M.
29 Haddon, F. Helidoniotis, K.L. Hill, N.J. Holbrook, G.W. Hosie, P.R. Last, S.D. Ling, J. Melbourne-Thomas,
30 K. Miller, G.T. Pecl, A.J. Richardson, K.R. Ridgway, S.R. Rintoul, D.A. Ritz, D.J. Ross, J.C. Sanderson, S.A.
31 Shepherd, A. Slotvinski, K.M. Swadling, and N. Taw, 2011: Climate change cascades: Shifts in
32 oceanography, species' ranges and subtidal marine community dynamics in eastern Tasmania. *Journal of*
33 *Experimental Marine Biology and Ecology*, **400(1-2)**, 17-32.
34 Jomelli, V., V.P. Pech, C. Chochillon, and D. Brunstein, 2004: Geomorphic variations of debris flows and recent
35 climatic change in the French Alps. *Climatic Change*, **64(1)**, 77-102.
36 Jomelli, V., V. Favier, A. Rabatel, D. Brunstein, G. Hoffmann, and B. Francou, 2009: Fluctuations of glaciers in the
37 tropical Andes over the last millennium and palaeoclimatic implications: A review. *Palaeogeography,*
38 *Palaeoclimatology, Palaeoecology*, **281(3)**, 269-282.
39 Jones, G.S., P.A. Stott, and N. Christidis, 2008: Human contribution to rapidly increasing frequency of very warm
40 Northern Hemisphere summers. *Journal of Geophysical Research*, **113(D2)**, D02109.
41 Jones, B.F. and B.A. Olken, 2010: Climate shocks and exports. *American Economic Review*, **100(2)**, 454-459.
42 Jones, R., O. Hoegh-Guldberg, A. Larkum, and U. Schreiber, 1998: Temperature induced bleaching of corals begins
43 with impairment to the carbon dioxide fixation mechanism of zooxanthellae. *Plant, Cell and Environment*, **21**,
44 1219-1230.
45 Jump, A., J. Hunt, and J. Peñuelas, 2006: Rapid climate change-related growth decline at the southern range-edge of
46 *Fagus sylvatica*. *Global Change Biology*, **12(11)**, 2163-2174.
47 Jump, A.S., T. Huang, and C. Chou, 2012: Rapid altitudinal migration of mountain plants in Taiwan and its
48 implications for high altitude biodiversity. *Ecography*, **35(3)**, 204-210.
49 Käab, A., R. Frauenfelder, and I. Roer, 2007: On the response of rockglacier creep to surface temperature increase.
50 *Global and Planetary Change*, **56(1)**, 172-187.
51 Käab, A., E. Berthier, C. Nuth, J. Gardelle, and Y. Arnaud, 2012: Contrasting patterns of early twenty-first-century
52 glacier mass change in the Himalayas. *Nature*, **488(7412)**, 495-498.
53 Karl, T.R., J.M. Melillo, and T.C. Peterson, 2009: *Global Climate Change Impacts in the United States*. Cambridge
54 University Press, Cambridge and New York, .

- 1 Kawachi, I., B.P. Kennedy, and R.G. Wilkinson, 1999: Crime: social disorganization and relative deprivation. *Social*
2 *Science and Medicine*, **48(6)**, 719-731.
- 3 Kay, A.L., S.M. Crooks, P. Pall, and D.A. Stone, 2011: Attribution of autumn/winter 2000 flood risk in England to
4 anthropogenic climate change: A catchment-based study. *Journal of Hydrology*, **406(1)**, 97-112.
- 5 Kearney, M.R., N.J. Briscoe, D.J. Karoly, W.P. Porter, M. Norgate, and P. Sunnucks, 2010: Early emergence in a
6 butterfly causally linked to anthropogenic warming. *Biology Letters*, **6(5)**, 674-677.
- 7 Keatley, M.R. and I.L. Hudson, 2012: Detecting change in an Australian flowering record: Comparisons of linear
8 regression and cumulative sum analysis change point analysis. *Austral Ecology*, **37(7)**, 825-835.
- 9 Keith, D.A., S. Rodoreda, and M. Bedward, 2010: Decadal change in wetland-woodland boundaries during the late
10 20th century reflects climatic trends. *Global Change Biology*, **16(8)**, 2300-2306.
- 11 Kelly, A.E. and M.L. Goulden, 2008: Rapid shifts in plant distribution with recent climate change. *Proceedings of*
12 *the National Academy of Sciences of the United States of America*, **105(33)**, 11823-11826.
- 13 Khalaj, B., G. Lloyd, V. Sheppard, and K. Dear, 2010: The health impacts of heat waves in five regions of New
14 South Wales, Australia: a case-only analysis. *International Archives of Occupational and Environmental*
15 *Health*, **83(7)**, 833-842.
- 16 Kharuk, V.I., S.T. Im, M.L. Dvinskaya, and K.I. Ranson, 2010: Climate-induced mountain tree-line evolution in
17 southern Siberia. *Scandinavian Journal of Forest Research*, **25**, 446-454.
- 18 Killeen, T.J., A. Guerra, M. Calzada, L. Correa, V. Calderon, L. Soria, B. Quezada, and M.K. Steininger, 2008:
19 Total historical land-use change in Eastern Bolivia: Who, where, when, and how much? *Ecology and Society*,
20 **13(1)**, 36.
- 21 Kingsford, R.T. and J.E.M. Watson, 2011: Climate change in Oceania - A synthesis of biodiversity impacts and
22 adaptations. *Pacific Conservation Biology*, **17**, 270-284.
- 23 Kniveton, D., C. Smith, and S. Wood, 2011: Agent-based model simulations of future changes in migration flows
24 for Burkina Faso. *Global Environmental Change*, **21(1)**, 34-40.
- 25 Knutson, T.R., J.L. McBride, J. Chan, K. Emanuel, G. Holland, C. Landsea, I. Held, J.P. Kossin, A. Srivastava, and
26 M. Sugi, 2010: Tropical cyclones and climate change. *Nature Geoscience*, **3(3)**, 157-163.
- 27 Kopp, R.E., J.X. Mitrovica, S.M. Griffies, J. Yin, C.C. Hay, and R.J. Stouffer, 2010: The impact of Greenland melt
28 on local sea levels: a partially coupled analysis of dynamic and static equilibrium effects in idealized water-
29 hosing experiments. *Climatic Change*, **103**, 619-625.
- 30 Kopytko, N. and J. Perkins, 2011: Climate change, nuclear power, and the adaptation-mitigation dilemma. *Energy*
31 *Policy*, **39(1)**, 318-333.
- 32 Korup, O., T. Görüm, and Y. Hayakawa, 2012: Without power? Landslide inventories in the face of climate change.
33 *Earth Surface Processes and Landforms*, **37(1)**, 92-99.
- 34 Koutsias, N., M. Arianoutsou, A.S. Kallimanis, G. Mallinis, J.M. Halley, and P. Dimopoulos, 2012: Where did the
35 fires burn in Peloponnisos, Greece the summer of 2007? Evidence for a synergy of fuel and weather.
36 *Agricultural and Forest Meteorology*, **156(0)**, 41-53.
- 37 Kovach, R.P., A.J. Gharrett, and D.A. Tallmon, 2012: Genetic change for earlier migration timing in a pink salmon
38 population. *Proceedings of the Royal Society B: Biological Sciences*, **279(1743)**, 3870-3878.
- 39 Kovats, R.S., D.H. Campbell-Lendrum, A.J. McMichael, A. Woodward, and J.S. Cox, 2001: Early effects of climate
40 change: do they include changes in vector-borne disease? *Philosophical Transactions of the Royal Society of*
41 *London B: Biological Sciences*, **356**, 1057-1068.
- 42 Krauss, K.W., D.R. Cahoon, J.A. Allen, K.C. Ewel, J.C. Lynch, and N. Cormier, 2010: Surface elevation change
43 and susceptibility of different mangrove zones to sea-level rise on pacific high islands of Micronesia.
44 *Ecosystems*, **13(1)**, 129-143.
- 45 Krepper, C.M., N.O. Garcia, and P.D. Jones, 2008: Low-frequency response of the upper Parana basin. *International*
46 *Journal of Climatology*, **28(3)**, 351-360.
- 47 Krepper, C.M. and G.V. Zucarelli, 2010: Climatology of water excesses and shortages in the La Plata Basin.
48 *Theoretical and Applied Climatology*, **102(1-2)**, 13-27.
- 49 Krishnan, P., S. Dam Roy, G. George, R.C. Srivastava, A. Anand, S. Murugesan, M. Kaliyamoorthy, N. Vikas, and
50 R. Soundararajan, 2011: Elevated sea surface temperature during May 2010 induces mass bleaching of corals
51 in the Andaman. *Current Science*, **100(1)**, 111-117.
- 52 Kristensen, K., K. Schelde, and J.E. Olesen, 2011: Winter wheat yield response to climate variability in Denmark.
53 *The Journal of Agricultural Science*, **149(1)**, 33-47.

- 1 Krupnik, I. and G.C. Ray, 2007: Pacific walruses, indigenous hunters, and climate change: Bridging scientific and
2 indigenous knowledge. *Deep Sea Research Part II: Topical Studies in Oceanography*, **54**, 2946-2957.
- 3 Kundzewicz, Z.W. and V. Krysanova, 2010: Climate change and stream water quality in the multi-factor context.
4 *Climatic Change*, **103(3-4)**, 353-362.
- 5 Kundzewicz, Z.W., I. Pińskwar, and G.R. Brakenridge, 2013: Large floods in Europe, 1985-2009. *Hydrological
6 Sciences Journal*, **58(1)**, 1-7.
- 7 Künzler, M., C. Huggel, A. Linsbauer, and W. Haeberli, 2010: Emerging risks related to new lakes in deglaciating
8 areas of the Alps. In: *Mountain Risks: Bringing Science to Society. Proceedings of the "Mountain Risk"
9 International Conference 24-26 November 2010, Firenze, Italy*, pp. 453-458.
- 10 Kurz, W.A., C.C. Dymond, G. Stinson, G.J. Rampley, E.T. Neilson, A.L. Carroll, T. Ebata, and L. Safranyik, 2008:
11 Mountain pine beetle and forest carbon feedback to climate change. *Nature*, **452(7190)**, 987-990.
- 12 Lacombe, G., M. McCartney, and G. Forkuor, 2012: Drying climate in Ghana over the period 1960-2005: evidence
13 from the resampling-based Mann-Kendall test at local and regional levels. *Hydrological Sciences Journal*,
14 **57(8)**, 1594-1609.
- 15 Laidler, G.J., 2006: Inuit and scientific perspectives on the relationship between sea ice and climate change: The
16 ideal complement? *Climatic Change*, **78**, 407-444.
- 17 Laidre, K.L., I. Stirling, L.F. Lowry, O. Wiig, M.O. Heide-Jørgensen, and S.H. Ferguson, 2008: Quantifying the
18 sensitivity of arctic marine mammals to climate-induced habitat change. *Ecological Applications*, **18(2)**, 97-
19 125.
- 20 Lamela-Silvarrey, C., C. Fernández, R. Anadón, and J. Arrontes, 2012: Furoid assemblages on the north coast of
21 Spain: past and present (1977–2007). *Botanica Marina*, **55(3)**, 199-207.
- 22 Lampis, A., 2010: *Challenges to adaptation for risk-prone coastal livelihoods in Tumaco, Pacific Coast (Colombia)*.
23 Urbanization and Global Environmental Change (UGEC), pp. 18-22.
- 24 Landsea, C., G.A. Vecchi, L. Bengtsson, and T.R. Knutson, 2009: Impact of duration thresholds on Atlantic tropical
25 cyclone counts. *Journal of Climate*, **23**, 2508-2519.
- 26 Lantuit, H., P.P. Overduin, N. Couture, S. Wetterich, F. Aré, D. Atkinson, J. Brown, G. Cherkashov, D. Drozdov, D.
27 Forbes, A. Graves-Gaylord, M. Grigoriev, H.W. Hubberten, J. Jordan, T. Jorgenson, R. Ødegård, S. Ogorodov,
28 W. Pollard, V. Rachold, S. Sedenko, S. Solomon, F. Steenhuisen, I. Streletskaaya, and A. Vasiliev, 2011: The
29 Arctic coastal dynamics database: A new classification scheme and statistics on Arctic permafrost coastlines.
30 *Estuaries and Coasts*, **35(2)**, 383-400.
- 31 Laternser, M. and M. Schneebeli, 2002: Temporal trend and spatial distribution of avalanche activity during the last
32 50 years in Switzerland. *Natural Hazards*, **27(3)**, 201-230.
- 33 Laybourn-Parry, J., 2003: Polar limnology, the past, the present and the future. In: *Antarctic biology in a global
34 context*. [Huiskes, A.H.L., W.W.C. Gieskes, J. Rozema, R.M.L. Schorno, S.M. van der Vies, and W.J.
35 Wolff(eds.)]. Backhuys Publishers, Leiden, pp. 321-329.
- 36 Lazo, J.K., M. Lawson, P.H. Larsen, and D.M. Waldman, 2011: U.S. economic sensitivity to weather variability.
37 *Bulletin of the American Meteorological Society*, **92(6)**, 709-720.
- 38 Le Polain de Waroux, Y. and E.F. Lambin, 2012: Monitoring degradation in arid and semi-arid forests and
39 woodlands: The case of the Argan woodlands (Morocco). *Applied Geography*, **32**, 777-786.
- 40 Le Quééré, C., M.R. Raupach, J.G. Canadell, and G. Marland, 2009: Trends in the sources and sinks of carbon
41 dioxide. *Nature Geoscience*, **2**, 831-836.
- 42 Leadley, P., H.M. Pereira, R. Alkemade, J.F. Fernandez-Manjarrés, V. Proença, J.P.W. Scharlemann, and M.J.
43 Walpole, 2010: *Biodiversity Scenarios: Projections of 21st century change in biodiversity and associated
44 ecosystem services*. Secretariat of the Convention on Biological Diversity, pp. 132.
- 45 Leatherman, S.P., K. Zhang, and B.C. Douglas, 2000: Sea level rise shown to drive coastal erosion. *Eos*,
46 *Transactions American Geophysical Union*, **81(6)**, 55-57.
- 47 Lenoir, J., J.C. Gegout, P.A. Marquet, P. de Ruffray, and H. Brisse, 2008: A significant upward shift in plant species
48 optimum elevation during the 20th century. *Science*, **320(5884)**, 1768-1771.
- 49 Lenton, T.M., 2011: Early warning of climate tipping points. *Nature Climate Change*, **1(4)**, 201-209.
- 50 Lenton, T.M., H. Held, E. Kriegler, J.W. Hall, W. Lucht, S. Rahmstorf, and H.J. Schellnhuber, 2008: Tipping
51 elements in the Earth's climate system. *Proceedings of the National Academy of Sciences of the United States
52 of America*, **105(6)**, 1786-1793.

- 1 Levin, L.A., W. Ekau, A.J. Gooday, F. Jorissen, J.J. Middelburg, S.W.A. Naqvi, C. Neira, N.N. Rabalais, and J.
2 Zhang, 2009: Effects of natural and human-induced hypoxia on coastal benthos. *Biogeosciences*, **6(10)**, 2063-
3 2098.
- 4 Li, X., R.W. Mazur, D.R. Allen, and D.R. Swatek, 2005: Specifying transformer winter and summer peak-load
5 limits. *IEEE Transactions on Power Delivery*, **20**, 185-190.
- 6 Li, Z.X., Y.Q. He, T. Pu, W.X. Jia, X.Z. He, H.X. Pang, N.N. Zhang, Q. Liu, S.J. Wang, G.F. Zhu, S.X. Wang, L.
7 Chang, J.K. Du, and H.J. Xin, 2010: Changes of climate, glaciers and runoff in China's monsoonal temperate
8 glacier region during the last several decades. *Quaternary International*, **218(1-2)**, 13-28.
- 9 Lilleør, H.B. and K. van den Broeck, 2011: Economic drivers of migration and climate change in LDCs. *Global
10 Environmental Change*, **21**, 70-81.
- 11 Lima, F.P., P.A. Ribeiro, Queiroz, S.J. Hawkins, and A.M. Santos, 2007: Do distributional shifts of northern and
12 southern species of algae match the warming pattern? *Global Change Biology*, **13**, 2592-2604.
- 13 Lima, F.P. and D.S. Wetthey, 2012: Three decades of high-resolution coastal sea surface temperatures reveal more
14 than warming. *Nature Communications*, **3**.
- 15 Lin, B.B., 2007: Agroforestry management as an adaptive strategy against potential microclimate extremes in coffee
16 agriculture. *Agricultural and Forest Meteorology*, **144(1-2)**, 85-94.
- 17 Lin, N., K. Emanuel, M. Oppenheimer, and E. Vanmarcke, 2012: Physically based assessment of hurricane surge
18 threat under climate change. *Nature Climate Change*, **2(6)**, 462-467.
- 19 Ling, S.D., C.R. Johnson, K. Ridgway, A.J. Hobday, and M. Haddon, 2009: Climate-driven range extension of a sea
20 urchin: inferring future trends by analysis of recent population dynamics. *Global Change Biology*, **15(3)**, 719-
21 731.
- 22 Litzow, M.A., J.D. Urban, and B.J. Laurel, 2008: Increased spatial variance accompanies reorganization of two
23 continental shelf ecosystems. *Ecological Applications*, **18(6)**, 1331-1337.
- 24 Liu, Y., E. Wang, X. Yang, and G. Wang, 2010: Contributions of climatic and crop varietal changes to crop
25 production in the North China Plain since 1980s. *Global Change Biology*, **16**, 2287-2299.
- 26 Lloyd, A.H., A.G. Bunn, and I. Berner, 2011: A latitudinal gradient in tree growth response to climate warming in
27 the Siberian taiga. *Global Change Ecology*, **17**, 1935-1945.
- 28 Lobell, D.B., W. Schlenker, and J. Costa-Roberts, 2011: Climate trends and global crop production since 1980.
29 *Science*, **333(6042)**, 616.
- 30 Lobell, D.B. and C.B. Field, 2007: Global scale climate - crop yield relationships and the impacts of recent
31 warming. *Environmental Research Letters*, **2(1)**, 014002.
- 32 Long, S.P., E.A. Ainsworth, J. Nösberger, and D.R. Ort, 2006: Food for thought: Lower-than-expected crop yield
33 stimulation with rising CO₂ concentrations. *Science*, **312(5782)**, 1918-1921.
- 34 Loriaux, T. and G. Casassa, 2013: Evolution of glacial lakes from the Northern Patagonia Icefield and terrestrial
35 water storage in a sea-level rise context. *Global and Planetary Change*, (in press).
- 36 Lotze, H.K., H.S. Lenihan, B.J. Bourque, R.H. Bradbury, R.G. Cooke, M.C. Kay, S.M. Kidwell, M.X. Kirby, C.H.
37 Peterson, and J.B.C. Jackson, 2006: Depletion, degradation, and recovery potential of estuaries and coastal
38 seas. *Science*, **312(5781)**, 1806-1809.
- 39 Loughnan, M.E., N. Nicholls, and N.J. Tapper, 2010: The effects of summer temperature, age and socioeconomic
40 circumstance on Acute Myocardial Infarction admissions in Melbourne, Australia. *International Journal of
41 Health Geographics*, **210(9)**, 41.
- 42 Lucey, S.M. and J.A. Nye, 2010: Shifting species assemblages in the Northeast US continental shelf large marine
43 ecosystem. *Marine Ecology Progress Series*, **415**, 23-33.
- 44 Lugon, R. and M. Stoffel, 2010: Rock-glacier dynamics and magnitude-frequency relations of debris flows in a
45 high-elevation watershed: Ritigraben, Swiss Alps. *Global and Planetary Change*, **73**, 202-210.
- 46 Luterbacher, J., M.A. Liniger, A. Menzel, N. Estrella, P.M. Della-Marta, C. Pfister, and E. Xoplaki, 2007:
47 Exceptional European warmth of autumn 2006 and winter 2007: Historical context, the underlying dynamics,
48 and its phenological impacts. *Geophysical Research Letters*, **34(12)**, L12704.
- 49 Luterbacher, J., D. Dietrich, E. Xoplaki, M. Grosjean, and H. Wanner, 2004: European seasonal and annual
50 temperature variability, trends, and extremes since 1500. *Science*, **303**, 1499-1503.
- 51 Ma, T. and C. Zhou, 2012: Climate-associated changes in spring plant phenology in China. *International Journal of
52 Biometeorology*, **56(2)**, 269-275.
- 53 MacDonald, G.M., 2010: Water, climate change, and sustainability in the southwest. *Proceedings of the National
54 Academy of Sciences of the United States of America*, **107(50)**, 21256-21262.

- 1 Mack, M.C., M.S. Bret-Harte, T.N. Hollingsworth, R.R. Jandt, E.A.G. Schuur, G.R. Shaver, and D.L. Verbyla,
2 2011: Carbon loss from an unprecedented Arctic tundra wildfire. *Nature*, **475(7357)**, 489-492.
- 3 Maclean, K. and L. Cullen, 2009: Research methodologies for the co-production of knowledge for environmental
4 management in Australia. *Journal of the Royal Society of New Zealand*, **39**, 205-208.
- 5 Magrin, G.O., M.I. Travasso, W.E. Baethgen, M.O. Grondona, A. Giménez, G. Cunha, J.P. Castaño, and G.R.
6 Rodriguez, 2007: *Past and Future Changes in Climate and their Impacts on Annual Crops Yield in South East*
7 *South America*. In: IPCC TGICA Expert Meeting Integrating Analysis of Regional Climate Change and
8 Response Options. Nadi, Fiji [Intergovernmental Panel on Climate Change (IPCC) (ed.)]. Meeting Report, pp.
9 121-124.
- 10 Malhi, Y., L. Aragao, D. Galbraith, C. Huntingford, R. Fisher, P. Zelazowski, S. Sitch, C. McSweeney, and P. Meir,
11 2009: Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rainforest.
12 *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 20610-20615.
- 13 Malik, N., B. Bookhagen, N. Marwan, and J. Kurths, 2011: Analysis of spatial and temporal extreme monsoonal
14 rainfall over South Asia using complex networks. *Climate Dynamics*, **39(3-4)**, 971-987.
- 15 Mann, D.H., T.S. Rupp, M.A. Olson, and P.A. Duffy, 2012: Is Alaska's boreal forest now crossing a major
16 ecological threshold? *Arctic, Antarctic, and Alpine Research*, **44(3)**, 319-331.
- 17 Mann, M.E. and K.A. Emanuel, 2012: Atlantic hurricane trends linked to climate change. **87**, 233-241.
- 18 Manson, G.K. and S.M. Solomon, 2007: Past and future forcing of Beaufort Sea coastal change. *Atmosphere-Ocean*,
19 **45(2)**, 107-122.
- 20 Manzello, D.P., J.A. Kleypas, D.A. Budd, C.M. Eakin, P.W. Glynn, and C. Langdon, 2008: Poorly cemented coral
21 reefs of the eastern tropical Pacific: Possible insights into reef development in a high-CO₂ world. *Proceedings*
22 *of the National Academy of Sciences of the United States of America*, **105(30)**, 10450-10455.
- 23 Maranz, S., 2009: Tree mortality in the African Sahel indicates an anthropogenic ecosystem displaced by climate
24 change. *Journal of Biogeography*, **36**, 1181-1193.
- 25 Marbà, N. and C.M. Duarte, 2010: Mediterranean warming triggers seagrass (*Posidonia oceanica*) shoot mortality.
26 *Global Change Biology*, **16**, 2366-2375.
- 27 Marchiori, L., J.F. Maystadt, and I. Schumacher, 2012: The impact of weather anomalies on migration in sub-
28 Saharan Africa. *Journal of Environmental Economics and Management*, **63(3)**, 335-374.
- 29 Marcos, M., M.N. Tsimplis, and A.G.P. Shaw, 2009: Sea level extremes in southern Europe. *Journal of Geophysical*
30 *Research*, **114**, C01007.
- 31 Marengo, J.A., C.A. Nobre, G. Sampaio, L.F. Salazar, and L.S. Borma, 2011: *Climate change in the Amazon Basin:*
32 *Tipping points, changes in extremes, and impacts on natural and human systems*. Springer, New York, pp.
33 259-283.
- 34 Marengo, J.A., L.M. Alves, W.R. Soares, D.A. Rodriguez, H. Camargo, M. Paredes, and A.D. Pablo, 2013: Two
35 contrasting seasonal extremes in tropical South America in 2012: Flood in Amazonia and drought in Northeast
36 Brazil. *Journal of Climate*, **(submitted)**.
- 37 Marsh, P., M. Russell, S. Pohl, H. Haywood, and C. Onclin, 2008: Changes in thaw lake drainage in the Western
38 Canadian Arctic from 1950 to 2000. *Hydrological Processes*, **23(1)**, 145-158.
- 39 Marshall, B.E., 2012: Does climate change really explain changes in the fisheries productivity of Lake Kariba
40 (Zambia-Zimbabwe)? *Transactions of the Royal Society of South Africa*, **67**, 45-51.
- 41 Marty, C., 2008: Regime shift of snow days in Switzerland. *Geophysical Research Letters*, **35**, L12501-L12501.
- 42 Marulanda, M.C., O.D. Cardona, and A.H. Barbat, 2010: Revealing the socioeconomic impact of small disasters in
43 Colombia using the DesInventar database. *Disasters*, **34(2)**, 552-570.
- 44 Mastrandrea, M.D., C.B. Field, T.F. Stocker, O. Edenhofer, K.L. Ebi, D.J. Frame, H. Held, E. Kriegler, K.J. Mach,
45 P.R. Matschoss, G.K. Plattner, G.W. Yohe, and F.W. Zwiers, 2010: *Guidance notes for lead authors of the*
46 *IPCC Fifth Assessment Report on consistent treatment of uncertainties*. Intergovernmental Panel on Climate
47 Change (IPCC), .
- 48 McDowell, J.Z. and J.J. Hess, 2012: Accessing adaptation: Multiple stressors on livelihoods in the Bolivian
49 highlands under a changing climate. *Global Environmental Change*, **22(2)**, 342-352.
- 50 McGrath, J. and D.B. Lobell, 2011: An independent method of deriving the carbon dioxide fertilization effect in dry
51 conditions using historical yield data from wet and dry years. *Global Change Biology*, **17(8)**, 2689-2696.
- 52 McIntyre, T., I.J. Ansorge, H. Bornemann, J. Plötz, C.A. Tosh, and M.N. Bester, 2011: Elephant seal dive behaviour
53 is influenced by ocean temperature. Implications for climate change impacts on an ocean predator. *Marine*
54 *Ecology Progress Series*, **441**, 257-272.

- 1 McKee, K.L., D.R. Cahoon, and I.C. Feller, 2007: Caribbean mangroves adjust to rising sea level through biotic
2 controls on change in soil elevation. *Global Ecology and Biogeography*, **16(5)**, 545-556.
- 3 Menéndez, M. and P.L. Woodworth, 2010: Changes in extreme high water levels based on a quasi-global tide-gauge
4 data set. *Journal of Geophysical Research*, **115(C10)**, C10011.
- 5 Menge, B.A., F. Chan, and J. Lubchenco, 2008: Response of a rocky intertidal ecosystem engineer and community
6 dominant to climate change. *Ecology Letters*, **11(2)**, 151-162.
- 7 Menzel, A., T.H. Sparks, N. Estrella, E. Koch, A. Aasa, R. Ahas, K. Alm-Kübler, P. Bissolli, O. Braslavskaja, A.
8 Briede, F.M. Chmielewski, Z. Crepinsek, Y. Curnel, A. Dahl, C. Defila, A. Donnelly, Y. Yolanda Filella, K.
9 Jatzczak, F. Måge, A. Mestre, Ø. Nordli, J. Peñuelas, P. Pirinen, V. Remišová, H. Scheifinger, M. Striz, A.
10 Susnik, A.J.H. Van Vliet, F. Wielgolaski, S. Zach, and A. Züst, 2006: European phenological response to
11 climate change matches the warming pattern. *Global Change Biology*, **12**, 1969-1976.
- 12 Merico, A., T. Tyrrell, E.J. Lessard, T. Oguz, P.J. Stabeno, S.I. Zeeman, and T.E. Whitledge, 2004: Modelling
13 phytoplankton succession on the Bering Sea shelf: role of climate influences and trophic interactions in
14 generating *Emiliania huxleyi* blooms 1997-2000. *Deep-Sea Research Part I*, **51(12)**, 1803-1826.
- 15 Meyssignac, B. and A. Cazenave, 2012: Sea level: A review of present-day and recent-past changes and variability.
16 *Journal of Geodynamics*, **58**, 96-109.
- 17 Midgley, G.F., S.L. Chown, and B.S. Kgope, 2007: Monitoring effects of anthropogenic climate change on
18 ecosystems: A role for systematic ecological observation? *South African Journal of Science*, **103(7-8)**, 282-
19 286.
- 20 Miguel, E., S. Satyanath, and E. Sergenti, 2004: Economic shocks and civil conflict: An instrumental variables
21 approach. *Journal of Political Economy*, **112(4)**, 725-753.
- 22 Miller, S., R. Muir-Wood, and A. Boissonade, 2008: An exploration of trends in normalized weather-related
23 catastrophe losses. In: *Climate extremes and society*. [Diaz, H.F. and R.J. Murnane(eds.)]. Cambridge
24 University Press, Cambridge, UK, pp. 225-247.
- 25 Mills, G., A. Buse, B. Gimeno, V. Bermejo, M. Holland, L. Emberson, and H. Pleijel, 2007: A synthesis of AOT40-
26 based response functions and critical levels of ozone for agricultural and horticultural crops. *Atmospheric*
27 *Environment*, **41(12)**, 2630-2643.
- 28 Min, S.K., X. Zhang, F.W. Zwiers, and G.C. Hegerl, 2011: Human contribution to more-intense precipitation
29 extremes. *Nature*, **470(7334)**, 378-381.
- 30 Moiseev, P.A., A.A. Bartysh, and Z.Y. Nagimov, 2010: Climate changes and tree stand dynamics at the upper limit
31 of their growth in the North Ural mountains. *Russian Journal of Ecology*, **41(6)**, 486-497.
- 32 Molau, U., 2010a: Long-term impacts of observed and induced climate change on tussock tundra near its southern
33 limit in northern Sweden. *Plant Ecology and Diversity*, **3(1)**, 29-34.
- 34 Molau, U., 2010b: *Recent changes of vegetation patterns in the mountains of northern Sweden. Europe's ecological*
35 *backbone: recognising the true value of mountains*. EEA Report, No. 6/2010. European Environment Agency,
36 pp. 159-160.
- 37 Mölg, T., N.J. Cullen, D.R. Hardy, G. Kaser, and L. Klok, 2008: Mass balance of a slope glacier on Kilimanjaro and
38 its sensitivity to climate. *International Journal of Climatology*, **28**, 881-892.
- 39 Mölg, T., M. Großhauser, A. Hemp, M. Hofer, and B. Marzeion, 2012: Limited forcing of glacier loss through land-
40 cover change on Kilimanjaro. *Nature Climate Change*, **2**, 254-258.
- 41 Møller, A.P., D. Rubolini, and E. Lehikoinen, 2008: Populations of migratory bird species that did not show a
42 phenological response to climate change are declining. *Proceedings of the National Academy of Sciences of the*
43 *United States of America*, **105(42)**, 16195-16200.
- 44 Moore, R.D., S.W. Fleming, B. Menounos, R. Wheate, A. Fountain, K. Stahl, K. Holm, and M. Jakob, 2009: Glacier
45 change in western North America: influences on hydrology, geomorphic hazards and water quality.
46 *Hydrological Processes*, **23(1)**, 42-61.
- 47 Morgan, P.B., T.A. Mies, G.A. Bollero, R.L. Nelson, and S.P. Long, 2006: Season-long elevation of ozone
48 concentration to projected 2050 levels under fully open-air conditions substantially decreases the growth and
49 production of soybean. *New Phytologist*, **170(2)**, 333-343.
- 50 Morton, T.A., P. Bretschneider, D. Coley, and T. Kershaw, 2011: Building a better future: An exploration of beliefs
51 about climate change and perceived need for adaptation within the building industry. *Building and*
52 *Environment*, **46(5)**, 1151-1158.
- 53 Moser, S. and M. Davidson, 2013: Coastal zone development and ecosystems. Chapter 25. In: *National climate*
54 *assessment. Draft for public comment*. US Global Change Research Program, .

- 1 Mote, P.W., 2006: Climate-driven variability and trends in mountain snowpack in western north America. *Journal*
2 *of Climate*, **19**, 6209-6220.
- 3 Moy, A.D., W.R. Howard, S.G. Bray, and T.W. Trull, 2009: Reduced calcification in modern Southern Ocean
4 planktonic foraminifera. *Nature Geoscience*, **2(4)**, 276-280.
- 5 Mueller, S.A., J.E. Anderson, and T.J. Wallington, 2011: Impact of biofuel production and other supply and demand
6 factors on food price increases in 2008. *Biomass and Bioenergy*, **35(5)**, 1623-1632.
- 7 Mueter, F.J. and M.A. Litzow, 2008: Sea ice retreat alters the biogeography of the Bering Sea continental shelf.
8 *Ecological Applications*, **18(2)**, 309-320.
- 9 Müller, P., 2011: Summer 2010 – Wildfires in Russia. In: *Topics Geo. Natural catastrophes 2010. Analyses,*
10 *assessments, positions.* Munich Reinsurance Company, Munich, pp. 26-28.
- 11 Müller, R., T. Laepple, I. Bartsch, and C. Wiencke, 2009: Impact of oceanic warming on the distribution of
12 seaweeds in polar and cold-temperate waters. *Botanica Marina*, **52(6)**, 617-638.
- 13 Muscatine, L., 1986: Bioenergetics of reef-building corals. *Indian Edition Series*, **12**, 297-306.
- 14 Nagai, S., G. Yoshida, and K. Tarutani, 2011: Change in species composition and distribution of algae in the coastal
15 waters of western Japan. In: *Global warming impacts - case studies on the economy, human health, and on*
16 *urban and natural environments.* [Casalegno, S. (ed.)]. InTech, .
- 17 Nakazawa, Y., R. Williams, A.T. Peterson, P. Mead, E. Staples, and K.L. Gage, 2007: Climate change effects on
18 plague and tularemia in the United States *Vector-Borne and Zoonotic Diseases*, **7**, 529-540.
- 19 Ndebele-Murisa, M.R., E. Mashonjowa, and T. Hill, 2011: The implications of a changing climate on the Kapenta
20 fish stocks of Lake Kariba. *Transactions of the Royal Society of South Africa*, **66**, 105-119.
- 21 Nepstad, D.C., C.M. Stickler, and O.T. Almeida, 2006: Globalization of the Amazon soy and beef industries:
22 Opportunities for conservation. *Conservation Biology*, **20(6)**, 1595-1603.
- 23 Nepstad, D.C. and C.M. Stickler, 2008: Managing the tropical agriculture revolution. *Journal of Sustainable*
24 *Forestry*, **27(1-2)**, 43-56.
- 25 Neuheimer, A.B., R.E. Thresher, J.M. Lyle, and J.M. Semmens, 2011: Tolerance limit for fish growth exceeded by
26 warming waters. *Nature Climate Change*, **1(2)**, 110-113.
- 27 Neumayer, E., 2003: Good policy can lower violent crime: Evidence from a cross-national panel of homicide rates,
28 1980-97. *Journal of Peace Research*, **40(6)**, 619-640.
- 29 Neumayer, E. and F. Barthel, 2011: Normalizing economic loss from natural disasters: a global analysis. *Global*
30 *Environmental Change*, **21(1)**, 13-24.
- 31 Nicholls, N., 2011: Comments on “ Have disaster losses increased due to anthropogenic climate change?”. *Bulletin*
32 *of the American Meteorological Society*, **92(6)**, 791-791.
- 33 Nicholls, R.J., P.P. Wong, V.R. Burkett, J.O. Codignotto, J.E. Hay, R.F. McLean, S. Ragoonaden, and C.D.
34 Woodroffe, 2007: Coastal systems and low-lying areas. *Climate Change 2007: Impacts, Adaptation and*
35 *Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental*
36 *Panel on Climate Change.* In: [Parry, M.L., O.F. Canziani, J.P. Palutikof, P.J. van der Linden, and C.E.
37 Hanson(eds.)]. Cambridge University Press, Cambridge, UK, pp. 315-356.
- 38 Nicholls, R.J., C. Woodroffe, and V. Burkett, 2009: Coastline degradation as an indicator of global change. In:
39 *Climate Change: Observed Impacts on Planet Earth.* [Letcher, T.M. (ed.)]. Elsevier, pp. 409-424.
- 40 Nicholls, N., 2006: Detecting and attributing Australian climate change: a review. *Australian Meteorological*
41 *Magazine*, **55(3)**, 199-211.
- 42 Nichols, T., F. Berkes, D. Jolly, N.B. Snow, and The Community of Sachs Harbour, 2004: Climate change and sea
43 ice: Local observations from the Canadian Western Arctic. *Arctic*, **57(1)**, 68-79.
- 44 Nielsen, J.Ø. and A. Reenberg, 2010: Temporality and the problem with singling out climate as a current driver of
45 change in a small West African village. *Journal of Arid Environments*, **74(4)**, 464-474.
- 46 Nightingale, A.J., 2011: Bounding difference: Intersectionality and the material production of gender, caste, class
47 and environment in Nepal. *Geoforum*, **42(2)**, 153-162.
- 48 Nordhaus, W., 2006: Geography and macroeconomics: New data and new findings. *Proceedings of the National*
49 *Academy of Sciences of the United States of America*, **103(10)**, 3510-3517.
- 50 Novelo-Casanova, D.A. and G. Suarez, 2010: Natural and man-made hazards in the Cayman Islands. *Natural*
51 *Hazards*, **55(2)**, 441-466.
- 52 Nye, J.A., J.S. Link, J.A. Hare, and W.J. Overholtz, 2009: Changing spatial distribution of fish stocks in relation to
53 climate and population size on the Northeast United States continental shelf. *Marine Ecology Progress Series*,
54 **393**, 111-129.

- 1 Nye, J.A., T.M. Joyce, Y. Kwon, and J.S. Link, 2011: Silver hake tracks changes in Northwest Atlantic circulation.
2 *Nature Communications*, **2**, 412.
- 3 Ogawa-Onishi, Y. and P.M. Berry, 2013: Ecological impacts of climate change in Japan: The importance of
4 integrating local and international publications. *Biological Conservation*, **157(0)**, 361-371.
- 5 Ogden, N.H., C. Bouchard, K. Kurtenbach, G. Margos, L.R. Lindsay, L. Trudel, S. Nguon, and F. Milord, 2010:
6 Active and passive surveillance and phylogenetic analysis of borrelia burgdorferi elucidate the process of lyme
7 disease risk emergence in Canada. *Environmental Health Perspectives*, **118(7)**, 909-914.
- 8 Okada, M., T. Iizumi, Y. Hayashi, and M. Yokozawa, 2009: A climatological analysis on the recent declining trend
9 of rice quality in Japan. *Journal of Agricultural Meteorology*, **65(4)**, 327-337.
- 10 Oliveira, P.J.C., G.P. Asner, D.E. Knapp, A. Almeyda, R. Galvan-Gildemeister, S. Keene, R.F. Raybin, and R.C.
11 Smith, 2007: Land-use allocation protects the Peruvian Amazon. *Science*, **317(5842)**, 1233-1236.
- 12 Olsen, M.S., T.V. Callaghan, J.D. Reist, L.O. Reiersen, D. Dahl-Jensen, M.A. Granskog, B. Goodison, G.K.
13 Hovelsrud, M. Johansson, and R. Kallenborn, 2011: The changing arctic cryosphere and likely consequences:
14 an overview. *AMBIO: A Journal of the Human Environment*, **40(1)**, 111-118.
- 15 Omumbo, J.A., B. Lyon, S.M. Waweru, S.J. Connor, and M.C. Thomson, 2011: Raised temperatures over the
16 Kericho tea estates: revising the climate in the East African highlands malaria debate. *Malaria Journal*, **10(1)**,
17 12.
- 18 Oppikofer, T., M. Jaboyedoff, and H.R. Keusen, 2008: Collapse at the eastern Eiger flank in the Swiss Alps. *Nature*
19 *Geoscience*, **1(8)**, 531-535.
- 20 Osbahr, H., C. Twyman, W. Neil Adger, and D.S.G. Thomas, 2008: Effective livelihood adaptation to climate
21 change disturbance: scale dimensions of practice in Mozambique. *Geoforum*, **39(6)**, 1951-1964.
- 22 Otto, F.E.L., N. Massey, G.J. Van Oldenborgh, R.G. Jones, and M.R. Allen, 2012: Reconciling two approaches to
23 attribution of the 2010 Russian heat wave. *Geophysical Research Letters*, **39(4)**, L04702.
- 24 Overeem, I. and J.P.M. Syvitsky, 2010: Shifting discharge peaks in Arctic rivers, 1977–2007. *Geografiska Annaler:*
25 *Series A, Physical Geography*, **92(2)**, 285-296.
- 26 Overland, J.E., J. Alheit, A. Bakun, J.W. Hurrell, D.L. Mackas, and A.J. Miller, 2010: Climate controls on marine
27 ecosystems and fish populations. *Journal of Marine Systems*, **79(3)**, 305-315.
- 28 Oxenford, H., R. Roach, A. Brathwaite, L. Nurse, R. Goodridge, F. Hinds, K. Baldwin, and C. Finney, 2008:
29 Quantitative observations of a major coral bleaching event in Barbados, Southeastern Caribbean. *Climatic*
30 *Change*, **87(3-4)**, 435-449.
- 31 Pall, P., T. Aina, D.A. Stone, P.A. Stott, T. Nozawa, A.G.J. Hilberts, D. Lohmann, and M.R. Allen, 2011:
32 Anthropogenic greenhouse gas contribution to flood risk in England and Wales in autumn 2000. *Nature*,
33 **470(7334)**, 382-385.
- 34 Panday, P.K. and B. Ghimire, 2012: Time-series analysis of NDVI from AVHRR data over the Hindu Kush–
35 Himalayan region for the period 1982–2006. *International Journal of Remote Sensing*, **33(21)**, 6710-6721.
- 36 Parker, G., 2008: Crisis and catastrophe: The global crisis of the seventeenth century reconsidered. *American*
37 *Historical Review*, **113(4)**, 1053-1079.
- 38 Parmesan, C. and G. Yohe, 2003: A globally coherent fingerprint of climate change impacts across natural systems.
39 *Nature*, **421**, 37-42.
- 40 Parmesan, C., 2006: Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology,*
41 *Evolution, and Systematics*, **37**, 637-669.
- 42 Parmesan, C., C. Duarte, E. Poloczanska, A.J. Richardson, and M.C. Singer, 2011: Overstretching attribution.
43 *Nature Climate Change*, **1**, 2-4.
- 44 Parmesan, C., M.T. Burrows, C.M. Duarte, E.S. Poloczanska, A.J. Richardson, D.S. Schoeman, and M.C. Singer,
45 2013: Beyond climate change attribution in conservation and ecological research. (in press).
- 46 Parnikoza, I., P. Convey, I. Dykyy, V. Trokhymets, G. Milinevsky, O. Tyschenko, D. Inozemtseva, and I.
47 Kozeretska, 2009: Current status of the Antarctic herb tundra formation in the Central Argentine Islands.
48 *Global Change Biology*, **15(7)**, 1685-1693.
- 49 Parry, M.L., O.F. Canziani, J.P. Palutikof, van der Linden, P J, and C.E. Hanson, 2007: Cross-chapter case study. In:
50 *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth*
51 *Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press,
52 Cambridge, UK, pp. 864-868.
- 53 Pasquini, A.I. and P.J. Depetris, 2007: Discharge trends and flow dynamics of South American rivers draining the
54 southern Atlantic seaboard: An overview. *Journal of Hydrology*, **333(2)**, 385-399.

- 1 Patterson, D.L., A. Easter-Pilcher, and W.R. Fraser, 2003: The effects of human activity and environmental
2 variability on long-term changes in Adélie penguin populations at Palmer Station, Antarctica. In: *Antarctic*
3 *biology in a global context*. [Huiskes, A., W. Gieskes, J. Rozema, R. Schorno, S. van der Vies, and W.
4 Wolff(eds.)]. Backhuys Publishers, Leiden, The Netherlands, pp. 301-307.
- 5 Paul, F. and W. Haeberli, 2008: Spatial variability of glacier elevation changes in the Swiss Alps obtained from two
6 digital elevation models. *Geophysical Research Letters*, **35(21)**, L21502-L21502.
- 7 Payet, R. and W. Agricole, 2006: Climate change in the seychelles: implications for water and coral reefs. *AMBIO:*
8 *A Journal of the Human Environment*, **35(4)**, 182-189.
- 9 Paz, S., N. Bisharat, E. Paz, O. Kidar, and D. Cohen, 2007: Climate change and the emergence of *Vibrio vulnificus*
10 disease in Israel. *Environmental Research*, **103(3)**, 390-396.
- 11 Pearce, T., B. Smit, F. Duerden, J.D. Ford, A. Goose, and F. Kataoyak, 2010: Inuit vulnerability and adaptive
12 capacity to climate change in Ulukhaktok, Northwest Territories, Canada. *Polar Record*, **46(237)**, 157-177.
- 13 Peck, L.S., M.S. Clark, S.A. Morley, A. Massey, and H. Rossetti, 2009: Animal temperature limits and ecological
14 relevance: effects of size, activity and rates of change. *Functional Ecology*, **23(2)**, 248-256.
- 15 Pellicciotti, F., A. Bauder, and M. Parola, 2010: Effect of glaciers on streamflow trends in the Swiss Alps. *Water*
16 *Resources Research*, **46(10)**, W10522.
- 17 Peng, S., J. Huang, J.E. Sheehy, R.C. Laza, R.M. Visperas, X. Zhong, G.S. Centeno, G.S. Khush, and K.G.
18 Cassman, 2004: Rice yields decline with higher night temperature from global warming. *Proceedings of the*
19 *National Academy of Sciences of the United States of America*, **101(27)**, 9971-9975.
- 20 Peng, C., Z. Ma, X. Lei, Q. Zhu, H. Chen, W. Wang, S. Liu, W. Li, X. Fang, and X. Zhou, 2011: A drought-induced
21 pervasive increase in tree mortality across Canada's boreal forests. *Nature Climate Change*, **1(9)**, 467-471.
- 22 Perry, C.L. and I.A. Mendelsohn, 2009: Ecosystem effects of expanding populations of *Avicennia germinans* in a
23 Louisiana salt marsh. *Wetlands*, **29(1)**, 396-406.
- 24 Perry, A.L., P.J. Low, J.R. Ellis, and J.D. Reynolds, 2005: Climate change and distribution shifts in marine fishes.
25 *Science*, **308(5730)**, 1912-1915.
- 26 Petley, D.N., 2010: On the impact of climate change and population growth on the occurrence of fatal landslides in
27 South, East and SE Asia. *Quarterly Journal of Engineering Geology & Hydrogeology*, **43(4)**, 487-496.
- 28 Petrow, T. and B. Merz, 2009: Trends in flood magnitude, frequency and seasonality in Germany in the period
29 1951-2002. *Journal of Hydrology*, **371(1-4)**, 129-141.
- 30 Philippart, C.J.M., R. Anadón, R. Danovaro, J.W. Dippner, K.F. Drinkwater, S.J. Hawkins, T. Oguz, G. O'Sullivan,
31 and P.C. Reid, 2011: Impacts of climate change on European marine ecosystems: Observations, expectations
32 and indicators. *Journal of Experimental Marine Biology and Ecology*, **400(1-2)**, 52-69.
- 33 Pielke Jr, R.A., J. Gratz, C.W. Landsea, D. Collins, M.A. Saunders, and R. Musulin, 2008: Normalised hurricane
34 damage in the United States: 1900-2005. *Natural Hazards Review*, **9**, 29-42.
- 35 Pielke Jr, R.A., J. Rubiera, C. Landsea, M.L. Fernandez, and R. Klein, 2003: Hurricane vulnerability in Latin
36 America and the Caribbean: normalized damage and loss potentials. *Natural Hazards Review*, **4(3)**, 101-114.
- 37 Pitt, N.R., E.S. Poloczanska, and A.J. Hobday, 2010: Climate-driven range changes in Tasmanian intertidal fauna.
38 *Marine and Freshwater Research*, **61(9)**, 963-970.
- 39 Polidoro, B.A., K.E. Carpenter, L. Collins, N.C. Duke, A.M. Ellison, J.C. Ellison, E.J. Farnsworth, E.S. Fernando,
40 K. Kathiresan, N.E. Koedam, S.R. Livingstone, T. Miyagi, G.E. Moore, V.N. Nam, J.E. Ong, J.H. Primavera,
41 S.G. Salmo III, J.C. Sanciangco, S. Sukardjo, Y. Wang, and J.W.H. Yong, 2010: The loss of species:
42 Mangrove extinction risk and geographic areas of global concern. *PLoS ONE*, **5(4)**, e10095.
- 43 Poloczanska, E.S., C.J. Brown, W.J. Sydeman, W. Kiessling, D.S. Schoeman, P.J. Moore, K. Brander, J.F. Bruno, L.
44 Buckley, M.T. Burrows, C.M. Duarte, B.S. Halpern, J. Holding, C.V. Kappel, M.I. O'Connor, J.M. Pandolfi,
45 C. Parmesan, F. Schwing, S.A. Thompson, and A.J. Richardson, 2013: Global imprint of climate change on
46 marine life. *Nature Climate Change*, (submitted).
- 47 Poloczanska, E.S., S. Smith, L. Fauconnet, J. Healy, I.R. Tibbetts, M.T. Burrows, and A.J. Richardson, 2011: Little
48 change in the distribution of rocky shore faunal communities on the Australian east coast after 50 years of
49 rapid warming. *Journal of Experimental Marine Biology and Ecology*, **400(1)**, 145-154.
- 50 Porter, J.R. and M. Gawith, 1999: Temperatures and the growth and development of wheat. A review. *European*
51 *Journal of Agronomy*, **10(1)**, 23-36.
- 52 Pörtner, H.O. and R. Knust, 2007: Climate change affects marine fishes through the oxygen limitation of thermal
53 tolerance. *Science*, **315(5808)**, 95-97.
- 54 Pörtner, H.O. and A.P. Farrell, 2008: Ecology. Physiology and climate change. *Science*, **322(5902)**, 690-692.

- 1 Pörtner, H.O., C. Bock, R. Knust, G. Lannig, M. Lucassen, F.C. Mark, and F.J. Sartoris, 2008: Cod and climate in a
2 latitudinal cline: physiological analyses of climate effects in marine fishes. *Climate Research*, **37(2-3)**, 253-
3 270.
- 4 Pörtner, H.O., 2012: Integrating climate-related stressor effects on marine organisms: unifying principles linking
5 molecule to ecosystem-level changes. *Marine Ecology Progress Series*, **470**, 273-290.
- 6 Post, V. and E. Abarca, 2010: Preface: Saltwater and freshwater interactions in coastal aquifers. *Hydrogeology*
7 *Journal*, **18(1)**, 1-4.
- 8 Post, E., M.C. Forchhammer, M.S. Bret-Harte, T.V. Callaghan, T.R. Christensen, B. Elberling, A.D. Fox, O. Gilg,
9 D.S. Hik, T.T. Høye, R.A. Ims, E. Jeppesen, D.R. Klein, J. Madsen, A.D. McGuire, S. Rysgaard, D.E.
10 Schindler, I. Stirling, M.P. Tamstorf, N.J.C. Tyler, R. van der Wal, J. Welker, P.A. Wookey, N.M. Schmidt,
11 and P. Aastrup, 2009: Ecological dynamics across the Arctic associated with recent climate change. *Science*,
12 **325(5946)**, 1355-1358.
- 13 Potter, N.J., F. Chiew, and A. Frost, 2010: An assessment of the severity of recent reductions in rainfall and runoff
14 in the Murray-Darling Basin. *Journal of Hydrology*, **381(1)**, 52-64.
- 15 Pouliotte, J., B. Smit, and L. Westerhoff, 2009: Adaptation and development: Livelihoods and climate change in
16 Subarnabad, Bangladesh. *Climate and Development*, **1(1)**, 31-46.
- 17 Poveda, G., Ó.A. Estrada-Restrepo, J.E. Morales, Ó.O. Hernández, A. Galeano, and S. Osorio, 2011: Integrating
18 knowledge and management regarding the climate-malaria linkages in Colombia. *Current Opinion in*
19 *Environmental Sustainability*, **3(6)**, 449-460.
- 20 Poveda, G. and K. Pineda, 2009: Reassessment of Colombia's tropical glaciers retreat rates: are they bound to
21 disappear during the 2010-2020 decade? *Advances in Geosciences*, **22(2008)**, 107-116.
- 22 Powers, L.A., T.C. Johnson, J.P. Werne, I.S. Castañeda, E.C. Hopmans, J.S. Sinninghe Damsté, and S. Schouten,
23 2011: Organic geochemical records of environmental variability in Lake Malawi during the last 700 years, Part
24 I: the TEX86 temperature record. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **303(1)**, 133-139.
- 25 Prathumratana, I., S. Sthiannopkao, and K. Woong Kim, 2008: The relationship of climatic and hydrological
26 parameters to surface water quality in the lower Mekong River. *Environment International*, **34**, 860-866.
- 27 Prowse, T.D. and K. Brown, 2010: *Appearing and disappearing of lakes in the Arctic and their impacts on*
28 *biodiversity*. In: Arctic Biodiversity Trends 2010 - Selected Indicators of Change. CAFF International
29 Secretariat, Akureyri, Iceland, pp. 68-70.
- 30 Prudhomme O'Meara, W., J. Nekesa Mangeni, R. Steketee, and B. Greenwood, 2010: Changes in the burden of
31 malaria in sub-Saharan Africa. *The Lancet Infectious Diseases*, **10**, 545-555.
- 32 Quayle, W.C., L.S. Peck, H. Peat, J. Ellis-Evans, and P.R. Harrigan, 2002: Extreme responses to climate change in
33 Antarctic lakes. *Science*, **295(5555)**, 645-645.
- 34 Raabe, E.A., L.C. Roy, and C.C. McIvor, 2012: Tampa Bay coastal wetlands: Nineteenth to twentieth century tidal
35 marsh-to-mangrove conversion. *Estuaries and Coasts*, **35(5)**, 1145-1162.
- 36 Rabatel, A., B. Francou, A. Soruco, J. Gomez, B. Cáceres, J.L. Ceballos, R. Basantes, M. Vuille, J.E. Sicart, C.
37 Huggel, M. Scheel, Y. Lejeune, Y. Arnaud, M. Collet, T. Condom, G. Consoli, V. Favier, V. Jomelli, R.
38 Galarraga, P. Ginot, L. Maisincho, J. Mendoza, M. Ménégoz, E. Ramirez, P. Ribstein, W. Suarez, M. Villacis,
39 and P. Wagnon, 2013: Current state of glaciers in the tropical Andes: a multi-century perspective on glacier
40 evolution and climate change. *The Cryosphere*, **7**, 81-102.
- 41 Rahman, M., T. Lund, and I. Bryceson, 2011: Salinity impacts on agro-biodiversity in three coastal, rural villages of
42 Bangladesh. *Ocean & Coastal Management*, **54(6)**, 455-468.
- 43 Rahmstorf, S. and D. Coumou, 2011: Increase of extreme events in a warming world. *Proceedings of the National*
44 *Academy of Sciences of the United States of America*, **108(44)**, 17905-17909.
- 45 Raleigh, C. and D. Kniveton, 2012: Come rain or shine: an analysis of conflict and climate variability in East Africa.
46 *Journal of Peace Research*, **49**, 51-64.
- 47 Ranasinghe, R. and M.J.F. Stive, 2009: Rising seas and retreating coastlines. *Climatic Change*, **97(3)**, 465-468.
- 48 Randolph, S.E., 2010: To what extent has climate change contributed to the recent epidemiology of tick-borne
49 diseases? *Veterinary Parasitology*, **167(2)**, 92-94.
- 50 Rankey, E.C., 2011: Nature and stability of atoll island shorelines: Gilbert Island chain, Kiribati, equatorial Pacific.
51 *Sedimentology*, **58(7)**, 1831-1859.
- 52 Rasheed, M.A. and R.K.F. Unsworth, 2011: Long-term climate-associated dynamics of a tropical seagrass meadow.
53 Implications for the future *Marine Ecology Progress Series*, **422**, 93-103.

- 1 Raupach, M.R., J.G. Canadell, and C. Le Quéré, 2008: Anthropogenic and biophysical contributions to increasing
2 atmospheric CO₂ growth rate and airborne fraction. *Biogeosciences*, **5**, 2867-2896.
- 3 Ravanel, L. and P. Deline, 2011: Climate influence on rockfalls in high-Alpine steep rockwalls. The north side of
4 the Aiguilles de Chamonix (Mont Blanc massif) since the end of the 'Little Ice Age'. *The Holocene*, **21(2)**, 357-
5 365.
- 6 Ravens, T.M., R.C. Thomas, K.A. Roberts, and P.H. Santschi, 2009: Causes of salt marsh erosion in Galveston Bay,
7 Texas. *Journal of Coastal Research*, **25(2)**, 265-272.
- 8 Raxworthy, C.J., R.G. Pearson, N. Rabibisoa, A.M. Rakotondrazafy, J.B. Ramanamanjato, A.P. Raselimanana, S.
9 Wu, R.A. Nussbaum, and D.A. Stone, 2008: Extinction vulnerability of tropical montane endemism from
10 warming and upslope displacement: a preliminary appraisal for the highest massif in Madagascar. *Global
11 Change Biology*, **14(8)**, 1703-1720.
- 12 Razumov, S.O., 2010: Permafrost as a factor of the dynamics of the coastal zone of the Russian East Arctic Seas.
13 *Oceanology*, **50(2)**, 262-267.
- 14 Reguero, B., F.J. Mendez, and I.J. Losada, 2013: Variability of multivariate wave climate in Latin America and the
15 Caribbean. *Global and Planetary Change*, **100**, 70-84.
- 16 Resurreccion, B.P., 2011: *The gender and climate debate. More of the same or new pathways of thinking and
17 doing?*. Asia Security Initiative Policy Series No.10. RSIS Center for non-traditional security studies,
18 Singapore.
- 19 Reusch, T.B., A. Ehlers, A. Hammerli, and B. Worm, 2005: Ecosystem recovery after climatic extremes enhanced
20 by genotypic diversity. *Proceedings of the National Academy of Sciences of the United States of America*,
21 **102(8)**, 2826-2831.
- 22 Riordan, B., D. Verbyla, and D.A. McGuire, 2006: Shrinking ponds in subarctic Alaska based on 1950–2002
23 remotely sensed images. *Journal of Geophysical Research*, **111(G4)**, G02004.
- 24 Rivadeneira, M.M. and M. Fernández, 2005: Shifts in southern endpoints of distribution in rocky intertidal species
25 along the south-eastern Pacific coast. *Journal of Biogeography*, **32(2)**, 203-209.
- 26 Roberts, M.J. and W. Schlenker, 2010: Identifying supply and demand elasticities of agricultural commodities:
27 Implications for the US ethanol mandate. *National Bureau of Economic Research Working Paper Series*,
28 **15921**.
- 29 Robine, J.M., S.L.K. Cheung, S. Le Roy, H. Van Oyen, C. Griffiths, J.P. Michel, and F.R. Herrmann, 2008: Death
30 toll exceeded 70,000 in Europe during the summer of 2003. *Comptes Rendus Biologies*, **331(2)**, 171-178.
- 31 Rocklöv, J., B. Forsberg, and K. Meister, 2009: Winter mortality modifies the heat-mortality association the
32 following summer. *European Respiratory Journal*, **33(2)**, 245-251.
- 33 Rodell, M., I. Velicogna, and J.S. Famiglietti, 2009: Satellite-based estimates of groundwater depletion in India.
34 *Nature*, **460(7258)**, 999-1002.
- 35 Rodríguez-Morales, A.J., A. Risquez, and L. Echezuria, 2010: Impact of climate change on health and disease in
36 Latin America. In: *Climate change and variability*. [Simard, S. (ed.)]. Sciyo, pp. 463-486.
- 37 Rodríguez-Oreggia, E., A. De La Fuente, R. De La Torre, and H.A. Moreno, 2012: Natural Disasters, Human
38 Development and Poverty at the Municipal Level in Mexico. *The Journal of Development Studies*, (**ahead-of-
39 print**), 1-14.
- 40 Roessig, J.M., C.M. Woodley, J.J. Cech, and L.J. Hansen, 2004: Effects of global climate change on marine and
41 estuarine fishes and fisheries. *Reviews in Fish Biology and Fisheries*, **14**, 251-275.
- 42 Romanovsky, V.E., D.S. Drozdov, N.G. Oberman, G.V. Malkova, A.L. Kholodov, S.S. Marchenko, N.G.
43 Moskalenko, D.O. Sergeev, N.G. Ukraintseva, A.A. Abramov, and others, 2010: Thermal state of permafrost
44 in Russia. *Permafrost and Periglacial Processes*, **21(2)**, 136-155.
- 45 Romero-Lankao, P. and D. Dodman, 2011: Cities in transition: transforming urban centers from hotbeds of GHG
46 emissions and vulnerability to seedbeds of sustainability and resilience: Introduction and editorial overview.
47 *Current Opinion in Environmental Sustainability*, **3(3)**, 113-120.
- 48 Rosenzweig, C., W.D. Solecki, S. Hammer, and S. Mehrotra (eds.), 2011: *Climate change and cities: First
49 assessment report of the Urban Climate Change Research Network*. Cambridge University Press, Cambridge,
50 UK, pp. 312.
- 51 Rosenzweig, C., F.N. Tubiello, R. Goldberg, E. Mills, and J. Bloomfield, 2002: Increased crop damage in the US
52 from excess precipitation under climate change. *Global Environmental Change*, **12(3)**, 197-202.
- 53 Rosenzweig, C., G. Casassa, D.J. Karoly, A. Imeson, C. Liu, A. Menzel, S. Rawlins, T.L. Root, B. Seguin, and P.
54 Tryjanowski, 2007: Assessment of observed changes and responses in natural and managed systems. In:

- 1 *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth*
2 *Assessment Report of the Intergovernmental Panel on Climate Change.* [Parry, M.L., O.F. Canziani, J.P.
- 3 Palutikof, van der Linden, P. J., and C.E. Hanson(eds.)]. Cambridge University Press, Cambridge, U.K, pp. 79-
4 131.
- 5 Rosenzweig, C., D. Karoly, M. Vicarelli, P. Neofotis, Q. Wu, G. Casassa, A. Menzel, T.L. Root, N. Estrella, B.
6 Seguin, P. Tryjanowski, C. Liu, S. Rawlins, and A. Imeson, 2008: Attributing physical and biological impacts
7 to anthropogenic climate change *Nature*, **453(7193)**, 353-357.
- 8 Ross, M.S., J.J. O'Brien, R.G. Ford, K. Zhang, and A. Morkill, 2009: Disturbance and the rising tide: The challenge
9 of biodiversity management on low-island ecosystems. *Frontiers in Ecology and the Environment*, **7(9)**, 471-
10 478.
- 11 Rothstein, H.R., A.J. Sutton, and M. Borenstein (eds.), 2005: *Publication bias in meta-analysis: prevention,*
12 *assessment, and adjustments.* Wiley, Chichester, UK, pp. 374.
- 13 Rotton, J. and E.G. Cohn, 2001: Temperature, routine activities, and domestic violence: A reanalysis. *Violence and*
14 *Victims*, **16(2)**, 203-215.
- 15 Rowland, J.C., C.E. Jones, G. Altmann, R. Bryan, B.T. Crosby, G.L. Geernaert, L.D. Hinzman, L.D. Kane, D.M.
16 Lawrence, A. Mancino, P. Marsh, J.P. McNamara, V.E. Romanovsky, H. Toniolo, B.J. Travis, E. Trochim,
17 and C.J. Wilson, 2010: Arctic landscapes in transition: Responses to thawing permafrost. *Eos*, **91(26)**, 229-
18 236.
- 19 Rozell, D.J. and T.F. Wong, 2010: Effects of climate change on groundwater resources at Shelter Island, New York
20 State, USA. *Hydrogeology Journal*, **18(7)**, 1657-1665.
- 21 Saba, V.S., M.A.M. Friedrichs, M.E. Carr, D. Antoine, R.A. Armstrong, I. Asanuma, O. Aumont, N.R. Bates, M.J.
22 Behrenfeld, V. Bennington, L. Bopp, J. Bruggeman, E.T. Buitenhuis, M.J. Church, A.M. Ciotti, S.C. Doney,
23 M. Dowell, J. Dunne, S. Dutkiewicz, W. Gregg, N. Hoepffner, K.J.W. Hyde, J. Ishizaka, T. Kameda, D.M.
24 Karl, I. Lima, M.W. Lomas, J. Marra, G.A. McKinley, F. Melin, J.K. Moore, A. Morel, J. O'Reilly, B.
25 Salihoglu, M. Scardi, T.J. Smyth, S.L. Tang, J. Tjiputra, J. Uitz, M. Vichi, K. Waters, T.K. Westberry, and A.
26 Yool, 2010: Challenges of modeling depth-integrated marine primary productivity over multiple decades: a
27 case study at BATS and HOT. *Global Biogeochemical Cycles*, **24(3)**.
- 28 Saldaña-Zorrilla, S. and K. Sandberg, 2009: Impact of climate-related disasters on human migration in Mexico: a
29 spatial model. *Climatic Change*, **96(1-2)**, 97-118.
- 30 Salick, J. and N. Ross, 2009: Traditional peoples and climate change: Introduction. *Global Environmental Change*,
31 **19**, 137-139.
- 32 Sander, J., J.F. Eichner, E. Faust, and M. Steuer, 2013: Rising variability in thunderstorm-related U.S. losses as a
33 reflection of changes in large-scale thunderstorm forcing. *Journal of the American Meteorological Society*,
34 (acc).
- 35 Sauber, J.M. and N.A. Ruppert, 2008: Rapid ice mass loss: Does it have an influence on earthquake occurrence in
36 southern Alaska? *Geophysical Monograph*, **179**, 369-384.
- 37 Saurral, R.I., V.R. Barros, and D.P. Lettenmaier, 2008: Land use impact on the Uruguay river discharge.
38 *Geophysical Research Letters*, **35(12)**, L12401.
- 39 Schär, C. and G. Jendritzky, 2004: Hot news from summer 2003. *Nature*, **432(7017)**, 559-560.
- 40 Schlenker, W., W.M. Hanemann, and A.C. Fisher, 2005: Will U.S. agriculture really benefit from global warming?
41 Accounting for irrigation in the hedonic approach. *The American Economic Review*, **95(1)**, 395-406.
- 42 Schlenker, W. and M.J. Roberts, 2009: Nonlinear temperature effects indicate severe damages to U.S. crop yields
43 under climate change. *Proceedings of the National Academy of Sciences of the United States of America*,
44 **106(37)**, 15594-15598.
- 45 Schlepner, C., 2008: Evaluation of coastal squeeze and its consequences for the Caribbean island Martinique.
46 *Ocean & Coastal Management*, **51(5)**, 383-390.
- 47 Schmidt, S., C. Kemfert, and P. Höpfe, 2010: The impact of socio-economics and climate change on tropical
48 cyclone losses in the USA. *Regional Environmental Change*, **10(1)**, 13-26.
- 49 Schmocker-Fackel, P. and F. Naef, 2010: More frequent flooding? Changes in flood frequency in Switzerland since
50 1850. *Journal of Hydrology*, **381(1)**, 1-8.
- 51 Schneider, D., C. Huggel, W. Haeberli, and R. Kaitna, 2011: Unraveling driving factors for large rock-ice avalanche
52 mobility. *Earth Surface Processes and Landforms*, **36(14)**, 1948-1966.
- 53 Schneider, P. and S.J. Hook, 2010: Space observations of inland water bodies show rapid surface warming since
54 1985. *Geophysical Research Letters*, **37(22)**, L22405.

- 1 Schwing, F.B., R. Mendelssohn, S.J. Bograd, J.E. Overland, M. Wang, and S. Ito, 2010: Climate change,
2 teleconnection patterns, and regional processes forcing marine populations in the Pacific. *Journal of Marine*
3 *Systems*, **79(3)**, 245-257.
- 4 Scott, D., B. Amelung, S. Becken, J.P. Ceron, G. Dubois, S. Gössling, P. Peeters, and M.C. Simpson, 2008: *Climate*
5 *change and tourism: Responding to global challenges*. Report for UNWTO, UNEP and WMO. World Tourism
6 Organization; United Nations Environment Programme, Madrid; Paris.
- 7 Senapathi, D., M.A.C. Nicoll, C. Teplitsky, C.G. Jones, and K. Norris, 2011: Climate change and the risks
8 associated with delayed breeding in a tropical wild bird population. *Proceedings of the Royal Society B:*
9 *Biological Sciences*, **278(1722)**, 3184-3190.
- 10 Seneviratne, S.I., N. Nicholls, D. Easterling, C.M. Goodess, S. Kanae, J. Kossin, Y. Luo, J. Marengo, K. McInnes,
11 M. Rahimi, M. Reichstein, A. Sorteberg, C. Vera, and X. Zhang, 2012: Changes in climate extremes and their
12 impacts on the natural physical environment. In: *Managing the Risks of Extreme Events and Disasters to*
13 *Advance Climate Change Adaptation. A Special Report of Working Groups I and II of the Intergovernmental*
14 *Panel on Climate Change (IPCC)*. [Field, C.B., V. Barros, T.F. Stocker, D. Qin, D.J. Dokken, K.L. Ebi *et*
15 *al.*(eds.)]. Cambridge University Press, Cambridge, UK and New York, NY, USA, pp. 109-230.
- 16 Sheppard, C., M. Al-Husiani, F. Al-Jamali, F. Al-Yamani, R. Baldwin, J. Bishop, F. Benzoni, E. Dutrieux, N.K.
17 Dulvy, S.R. Durvasula, D.A. Jones, R. Loughland, D. Medio, M. Nithyanandan, G.M. Pilling, I. Polikarpov,
18 A.R. Price, S. Purkis, B. Riegl, M. Saburova, K.S. Namin, O. Taylor, S. Wilson, and K. Zainal, 2010: The
19 Gulf: a young sea in decline. *Marine Pollution Bulletin*, **60(1)**, 13-38.
- 20 Shiklomanov, A.I., R.B. Lammers, M.A. Rawlins, L.C. Smith, and T.M. Pavelsky, 2007: Temporal and spatial
21 variations in maximum river discharge from a new Russian data set. *Journal of Geophysical Research*, **112**,
22 G04S53.
- 23 Shrestha, A.B. and R. Aryal, 2011: Climate change in Nepal and its impact on Himalayan glaciers. *Regional*
24 *Environmental Change*, **11(1)**, 65-77.
- 25 Shrestha, U.B., S. Gautam, and K.S. Bawa, 2012: Widespread climate change in the Himalayas and associated
26 changes in local ecosystems. *PLoS ONE*, **7(5)**, e36741.
- 27 Shriver, A.L., B.H. Yeo, K.O. Ting, M. Garcia, and M. Ahmed, 2006: Annotated bibliography on the economic
28 effects of global climate change on fisheries. *World Fish Bibliography*, **14**, 48.
- 29 Sigmundsson, F., V. Pinel, B. Lund, F. Albino, C. Pagli, H. Geirsson, and E. Sturkell, 2010: Climate effects on
30 volcanism: influence on magmatic systems of loading and unloading from ice mass variations, with examples
31 from Iceland. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering*
32 *Sciences*, **368(1919)**, 2519-2534.
- 33 Silva Dias, M.A.F., J. Dias, L. Carvalho, E. Freitas, and P.L. Silva Dias, 2013: Changes in extreme daily rainfall for
34 São Paulo, Brazil. *Climatic Change*, **116(3-4)**, 705-722.
- 35 Simmons, K., D. Sutter, and R.A. Pielke Jr, 2012: Normalized tornado damage in the United States: 1950-2011.
36 *Environmental Hazards*, , 1-16 (in press).
- 37 Skoplaki, E. and J.A. Palyvos, 2009: On the temperature dependence of photovoltaic module electrical performance:
38 A review of efficiency/power correlations. *Solar Energy*, **83(5)**, 614-624.
- 39 Slettebak, R.T., 2012: Don't blame the weather! Climate-related natural disasters and civil conflict. *Journal of Peace*
40 *Research*, **49(1)**, 163-176.
- 41 Smith, J., J. Schellnhuber, M. Mirza, S. Fankhauser, R. Leemans, L. Erda, L. Ogallo, B. Pittock, R. Richels, C.
42 Rosenzweig, U. Safriel, R.S.J. Tol, J. Weyant, and G. Yohe, 2001: Vulnerability to climate change and reasons
43 for concern: A synthesis. Chapter 19. In: *Climate change 2001: Impacts, adaptation and vulnerability*,
44 Cambridge University Press, Cambridge, UK, pp. 913-967.
- 45 Smith, J.B., S.H. Schneider, M. Oppenheimer, G.W. Yohe, W. Hare, M.D. Mastrandrea, A. Patwardhan, I. Burton,
46 J. Corfee-morlot, C.H.D. Magadza, H.M. Fussler, A.B. Pittock, A. Rahman, A. Suarez, and J.P. Van Ypersele,
47 2009: Assessing dangerous climate change through an update of the Intergovernmental Panel on Climate
48 Change (IPCC) 'Reasons for concern'. *Proceedings of the National Academy of Sciences of the United States*
49 *of America*, **106(11)**, 4133-4137.
- 50 Smith, J.R., P. Fong, and R.F. Ambrose, 2006a: Dramatic declines in mussel bed community diversity: response to
51 climate change? *Ecology*, **87(5)**, 1153-1161.
- 52 Smith, V.K., J.C. Carbon, J.C. Pope, D.G. Hallstrom, and M.E. Darden, 2006b: Adjusting to natural disasters.
53 *Journal of Risk and Uncertainty*, **33**, 37-54.

- 1 Smol, J.P. and M.S.V. Douglas, 2007: Crossing the final ecological threshold in high Arctic ponds. *Proceedings of*
2 *the National Academy of Sciences of the United States of America*, **104(30)**, 12395-12397.
- 3 Somero, G.N., 2012: The physiology of global change: Linking patterns to mechanisms. *Annual Review of Marine*
4 *Science*, **4**, 39-61.
- 5 Stahl, K., H. Hisdal, J. Hannaford, L. Tallaksen, H. Van Lanen, E. Sauquet, S. Demuth, M. Fendekova, and J.
6 Jordar, 2010: Streamflow trends in Europe: evidence from a dataset of near-natural catchments. *Hydrology and*
7 *Earth System Sciences Discussions*, **14**, 2367-2382.
- 8 Stern, D.I., P.W. Gething, C.W. Kabaria, W.H. Temperley, A.M. Noor, E.A. Okiro, G.D. Shanks, R.W. Snow, and
9 S.I. Hay, 2011: Temperature and malaria trends in highland East Africa. *PLoS ONE*, **6(9)**, e24524.
- 10 Stewart, I.T., D.R. Cayan, and M.D. Dettinger, 2005: Changes toward earlier streamflow timing across western
11 North America. *Journal of Climate*, **18**, 1136-1151.
- 12 Stige, L.C., G. Ottersen, K. Brander, K.S. Chan, and N.C. Stenseth, 2006: Cod and climate. Effect of the North
13 Atlantic Oscillation on recruitment in the North Atlantic. *Marine Ecology Progress Series*, **325**, 227-241.
- 14 Stige, L.C., G. Ottersen, P. Dalpadado, K.S. Chan, D. Hjermann, D.L. Lajus, N.A. Yaragina, and N.C. Stenseth,
15 2010: Direct and indirect climate forcing in a multi-species marine system. *Proceedings of the Royal Society*
16 *B: Biological Sciences*, **277(1699)**, 3411-3420.
- 17 Stock, C.A., M.A. Alexander, N.A. Bond, K.M. Brander, W.W.L. Cheung, E.N. Curchitser, T.L. Delworth, J.P.
18 Dunne, S.M. Griffies, M.A. Haltuch, J.A. Hare, A.B. Hollowed, P. Lehodey, S.A. Levin, J.S. Link, K.A. Rose,
19 R.R. Rykaczewski, J.L. Sarmiento, R.J. Stouffer, F.B. Schwing, G.A. Vecchi, and F.E. Werner, 2010: On the
20 use of IPCC-class models to assess the impact of climate on Living Marine Resources. *Progress in*
21 *Oceanography*, **88(1)**, 1-27.
- 22 Stocker, T.F., Q. Dahe, G.K. Plattner, P. Midgley, and M. Tignor, 2011: Tried and tested. *Nature Climate Change*,
23 **1(2)**, 71.
- 24 Stoffel, M. and C. Huggel, 2012: Effects of climate change on mass movements in mountain environments.
25 *Progress in Physical Geography*, **36(6)**, 421-439.
- 26 Stokes, D.J., T.R. Healy, and P.J. Cooke, 2010: Expansion dynamics of monospecific, temperate mangroves and
27 sedimentation in two embayments of a barrier-enclosed lagoon, Tauranga Harbour, New Zealand. *Journal of*
28 *Coastal Research*, **26(1)**, 113-122.
- 29 Stokes, C.R., M. Shahgedanova, I.S. Evans, and V.V. Popovnin, 2013: Accelerated loss of alpine glaciers in the
30 Kodar Mountains, south-eastern Siberia. *Global and Planetary Change*, **101(0)**, 82-96.
- 31 Stolper, D.A., N.P. Revsbech, and D.E. Canfield, 2010: Aerobic growth at nanomolar oxygen concentrations.
32 *Proceedings of the National Academy of Sciences of the United States of America*, **107(44)**, 18755-18760.
- 33 Stone, D., M. Auffhammer, M. Carey, G. Hansen, C. Huggel, W. Cramer, D. Lobell, U. Molau, A. Solow, L. Tibig,
34 and G. Yohe, 2013: The challenge to detect and attribute effects of climate change on human and natural
35 systems. (submitted).
- 36 Storey, D. and S. Hunter, 2010: Kiribati: an environmental 'perfect storm'. *Australian Geographer*, **41(2)**, 167-181.
- 37 Stott, P.A., D.A. Stone, and M.R. Allen, 2004: Human contribution to the European heatwave of 2003. *Nature*,
38 **432(7017)**, 610-614.
- 39 Stramma, L., E.D. Prince, S. Schmidtko, J. Luo, J.P. Hoolihan, M. Visbeck, D.W.R. Wallace, P. Brandt, and A.
40 Kortzinger, 2012: Expansion of oxygen minimum zones may reduce available habitat for tropical pelagic
41 fishes. *Nature Climate Change*, **2(1)**, 33-37.
- 42 Stramma, L., G.C. Johnson, J. Sprintall, and V. Mohrholz, 2008: Expanding oxygen-minimum zones in the tropical
43 oceans. *Science*, **320(5876)**, 655-658.
- 44 Stramma, L., S. Schmidtko, L.A. Levin, and G.C. Johnson, 2010: Ocean oxygen minima expansions and their
45 biological impacts. *Deep-Sea Research Part I: Oceanographic Research Papers*, **57(4)**, 587-595.
- 46 Strong, A.E., G. Liu, W. Skirving, and C.M. Eakin, 2011: NOAA's Coral Reef Watch program from satellite
47 observations. *Annals of GIS*, **17(2)**, 83-92.
- 48 Strong, A.E., G. Liu, J. Meyer, J.C. Hendee, and D. Sasko, 2004: Coral Reef Watch 2002. *Bulletin of Marine*
49 *Science*, **75(2)**, 259-268.
- 50 Šumilo, D., A. Bormane, L. Asokliene, V. Vasilenko, I. Golovljova, T. Avsic-Zupanc, Z. Hubalek, and S.E.
51 Randolph, 2008: Socio-economic factors in the differential upsurge of tick-borne encephalitis in Central and
52 Eastern Europe. *Reviews in Medical Virology*, **18(2)**, 81-95.

- 1 Šumilo, D., A. Bormane, V. Vasilenko, I. Golovljova, L. Asokliene, M. Žygtiene, and S. Randolph, 2009: Upsurge
2 of tick-borne encephalitis in the Baltic States at the time of political transition, independent of changes in
3 public health practices. *Clinical Microbiology and Infection*, **15(1)**, 75-80.
- 4 Sunday, J.M., A.E. Bates, and N.K. Dulvy, 2012: Thermal tolerance and the global redistribution of animals. *Nature*
5 *Climate Change*, **2(9)**, 686-690.
- 6 Svensson, C., J. Hannaford, Z.W. Kundzewicz, and T.J. Marsh, 2006: Trends in river floods: why is there no clear
7 signal in observations? *IAHS Publications-Series of Proceedings and Reports*, **305**, 1-18.
- 8 Swanson, D.A., J.C. Hiley, H.D. Venema, and R. Grosshans, 2007: *Indicators of Adaptive Capacity to Climate*
9 *Change for Agriculture in the Prairie Region of Canada: An analysis based on Statistics Canada's Census of*
10 *Agriculture*. In: Working Paper for the Prairie Climate Resilience Project [International Institute for
11 Sustainable Development (ed.)], Winnipeg, Canada.
- 12 Swift, G.W., Zocholl, E.S., Bajpai, M., J.F. Burger, C.H. Castro, S.R. Chano, F. Cobelo, P. de Sa, E.C. Fennell, J.G.
13 Gilbert, S.E. Grier, R.W. Haas, W.G. Hartmann, R.A. Hedding, P. Kerrigan, S. Mazumdar, D.H. Miller, P.G.
14 Mysore, M. Nagpal, R.V. Rebbapragada, M.V. Thaden, J.T. Uchiyama, S.M. Usman, J.D. Wardlow, and M.
15 Yalla, 2001: Adaptive transformer thermal overload protection. *IEEE Transactions on Power Delivery*, **16**,
16 516-521.
- 17 SwissRe, 2011: *Swiss Re provides estimate of its claims costs from Thailand flood* Swiss Re news release, 6
18 December 2011. [online] Available from: [www.swissre.com/media/news_releases/
19 Swiss_Re_provides_estimate_of_its_claims_costs_from_Thailand_flood.html](http://www.swissre.com/media/news_releases/Swiss_Re_provides_estimate_of_its_claims_costs_from_Thailand_flood.html).
- 20 Sydeman, W.J. and S.J. Bograd, 2009: Marine ecosystems, climate, and phenology: introduction. *Marine Ecology*
21 *Progress Series*, **393**, 185-188.
- 22 Syvitski, J.P., C.J. Vörösmarty, A.J. Kettner, and P. Green, 2005: Impact of humans on the flux of terrestrial
23 sediment to the global coastal ocean. *Science*, **308(5720)**, 376-380.
- 24 Tacoli, C., 2009: Crisis or adaptation? Migration and climate change in a context of high mobility. *Environment and*
25 *Urbanization*, **21(2)**, 513-525.
- 26 Takasuka, A., Y. Oozeki, and I. Aoki, 2007: Optimal growth temperature hypothesis. Why do anchovy flourish and
27 sardine collapse or vice versa under the same ocean regime? *Canadian Journal of Fisheries and Aquatic*
28 *Sciences*, **64(5)**, 768-776.
- 29 Takasuka, A., Y. Oozeki, and H. Kubota, 2008: Multi-species regime shifts reflected in spawning temperature
30 optima of small pelagic fish in the western North Pacific. *Marine Ecology Progress Series*, **360**, 211-217.
- 31 Tamisiea, M.E. and J.X. Mitrovica, 2011: The moving boundaries of sea level change: Understanding the origins of
32 geographic variability. *Oceanography*, **24**, 24-39.
- 33 Tamura, T., K. Horaguchi, Y. Saito, L.N. Van, M. Tateishi, K.O.T. Thi, F. Nanayama, and K. Watanabe, 2010:
34 Monsoon-influenced variations in morphology and sediment of a mesotidal beach on the Mekong River delta
35 coast. *Geomorphology*, **116(1-2)**, 11-23.
- 36 Tan, A., J.C. Adam, and D.P. Lettenmaier, 2011: Change in spring snowmelt timing in Eurasian Arctic rivers.
37 *Journal of Geophysical Research*, **116(D3)**, D03101.
- 38 Tape, K.D., M. Sturm, and C. Racine, 2006: The evidence for shrub expansion in northern Alaska and the
39 Pan-Arctic. *Global Change Biology*, **12**, 686-702.
- 40 Tape, K.D., M. Hallinger, J.M. Welker, and R.W. Ruess, 2012: Landscape heterogeneity of shrub expansion in
41 arctic Alaska. *Ecosystems*, **15(5)**, 1-14.
- 42 Tasker, M.L., 2008: *The effect of climate change on the distribution and abundance of marine species in the OSPAR*
43 *Maritime Area*. In: ICES Cooperative Research Report No. 293. ICES, Copenhagen, Denmark, pp. 45.
- 44 Taylor, R.G., L. Mileham, C. Tindimugaya, and L. Mwebembezi, 2009: Recent glacial recession and its impact on
45 alpine riverflow in the Rwenzori Mountains of Uganda. *Journal of African Earth Sciences*, **55(3-4)**, 205-213.
- 46 Taylor, R.G., B. Scanlon, P. Doll, M. Rodell, R. Van Beek, Y. Wada, L. Longuevergne, M. Leblanc, J.S.
47 Famiglietti, M. Edmunds, L. Konikow, T.R. Green, J. Chen, M. Taniguchi, M.F.P. Bierkens, A. MacDonald,
48 Y. Fan, R.M. Maxwell, Y. Yechieli, J.J. Gurdak, D.M. Allen, M. Shamsudduha, K. Hiscock, P.J.F. Yeh, I.
49 Holman, and H. Treidel, 2012: Ground water and climate change. *Nature Climate Change*, (advance online
50 publication).
- 51 Teixeira, M.G., M.C.N. Costa, F. Barreto, and M.L. Barreto, 2009: Dengue: twenty-five years since reemergence in
52 Brazil. *Cadernos De Saúde Pública*, **25**, 7-18.
- 53 Terrier, S., F. Jordan, A. Schleiss, W. Haeberli, C. Huggel, and M. Künzler, 2011: Optimized and adapted
54 hydropower management considering glacier shrinkage in the Swiss Alps.[Schleiss, A. and R.M. Boes(eds.)].

- 1 Proceedings of the International Symposium on Dams and Reservoirs under Changing Challenges–79th
2 Annual Meeting of ICOLD, Swiss Committee on Dams, Lucerne, Switzerland, Lucerne, Switzerland, pp. 291-
3 298.
- 4 Terry, J.P. and A.C. Falkland, 2010: Responses of atoll freshwater lenses to storm-surge overwash in the Northern
5 Cook Islands. *Hydrogeology Journal*, **18(3)**, 749-759.
- 6 Theisen, O.M., H. Holtermann, and H. Buhaug, 2012: Climate wars? Assessing the claim that drought breeds
7 conflict. *International Security*, **36(3)**, 79-106.
- 8 Theisen, O.M., 2012: Climate clashes? Weather variability, land pressure, and organized violence in Kenya, 1989-
9 2004. *Journal of Peace Research*, **49(1)**, 81-96.
- 10 Thresher, R.E., J.A. Koslow, A.K. Morison, and D.C. Smith, 2007: Depth-mediated reversal of the effects of climate
11 change on long-term growth rates of exploited marine fish. *Proceedings of the National Academy of Sciences
12 of the United States of America*, **104(18)**, 7461-7465.
- 13 Tierney, J.E., M.T. Mayes, N. Meyer, C. Johnson, P.W. Swarzenski, A.S. Cohen, and J.M. Russell, 2010: Late-
14 twentieth-century warming in Lake Tanganyika unprecedented since AD 500. *Nature Geoscience*, **3**, 422-425.
- 15 Tol, R.S.J. and S. Wagner, 2010: Climate change and violent conflict in Europe over the last millennium. *Climatic
16 Change*, **99(1-2)**, 65-79.
- 17 Tollefson, J., 2013: Natural hazards: New York vs the sea. *Nature*, **494(7436)**, 162-164.
- 18 Toniolo, H., P. Kodial, L.D. Hinzman, and K. Yoshikawa, 2009: Spatio-temporal evolution of a thermokarst in
19 Interior Alaska. *Cold Regions Science and Technology*, **56(1)**, 39-49.
- 20 Trathan, P.N., P.T. Fretwell, and B. Stonehouse, 2011: First recorded loss of an emperor penguin colony in the
21 recent period of Antarctic regional warming: implications for other colonies. *PLoS ONE*, **6(2)**, e14738.
- 22 Trenberth, K.E. and D.J. Shea., 2006: Atlantic hurricanes and natural variability in 2005. *Geophysical Research
23 Letters*, **33**, 1-4.
- 24 Trenberth, K.E. and J.T. Fasullo, 2012: Climate extremes and climate change: the Russian heat wave and other
25 climate extremes of 2010. *Journal of Geophysical Research*, **117(D17)**, D17103.
- 26 Trenberth, K.E., 2011: Attribution of climate variations and trends to human influences and natural variability.
27 *Wiley Interdisciplinary Reviews: Climate Change*, **2(6)**, 925-930.
- 28 Trivelpiece, W.Z., J.T. Hinke, A.K. Miller, C.S. Reiss, S.G. Trivelpiece, and G.M. Watters, 2011: Variability in krill
29 biomass links harvesting and climate warming to penguin population changes in Antarctica. *Proceedings of the
30 National Academy of Sciences of the United States of America*, **108(18)**, 7625-7628.
- 31 Tschakert, P., R. Tutu, and A. Alcaro, 2011: Embodied experiences of environmental and climatic changes in
32 landscapes of everyday life in Ghana. *Emotion, Space and Society*, (in press).
- 33 Uhlmann, M., O. Korup, C. Huggel, L. Fischer, and J.S. Kargel, 2012: Supra-glacial deposition and flux of
34 catastrophic rock-slope failure debris, south-central Alaska. *Earth Surface Processes and Landforms*, .
- 35 UNFCCC, 1992: *The United Nations Framework Convention on Climate Change*
36 <http://unfccc.int/resource/docs/convkp/conveng.pdf>, accessed 8.3.2013.
- 37 Valt, M. and P. Cianfarra, 2010: Recent snow cover variability in the Italian Alps. *Cold Regions Science and
38 Technology*, **64(2)**, 146-157.
- 39 Van Bogaert, R., K. Haneca, J. Hoogesteger, C. Jonasson, M. De Dapper, and T.V. Callaghan, 2011: A century of
40 tree line changes in sub-Arctic Sweden shows local and regional variability and only a minor influence of 20th
41 century climate warming. *Journal of Biogeography*, **38(5)**, 907-921.
- 42 Van den Honert, R.C. and J. McAneney, 2011: The 2011 Brisbane floods: Causes, impacts and implications. *Water*,
43 **3(4)**, 1149-1173.
- 44 Van Dingenen, R., F.J. Dentener, F. Raes, M.C. Krol, L. Emberson, and J. Cofala, 2009: The global impact of ozone
45 on agricultural crop yields under current and future air quality legislation. *Atmospheric Environment*, **43(3)**,
46 604-2310.
- 47 Van Oldenborgh, G.J., 2007: How unusual was autumn 2006 in Europe? *Climate of the Past*, **3**, 659-669.
- 48 Van Oldenborgh, G.J., A. Van Urk, and M. Allen, 2012: The absence of a role of climate change in the 2011
49 Thailand floods. *Bulletin of the American Meteorological Society*, **93**, 1047-1049.
- 50 Vaquer-Sunyer, R. and C.M. Duarte, 2011: Temperature effects on oxygen thresholds for hypoxia in marine benthic
51 organisms. *Global Change Biology*, **17(5)**, 1788-1797.
- 52 Vecchi, G.A., K.L. Swanson, and B.J. Soden, 2008: Whither hurricane activity? *Science*, **322(5902)**, 687-689.
- 53 Véran, S., O. Gimenez, E. Flint, W.L. Kendall, P.F.J. Doherty, and J.D. Lebreton, 2007: Quantifying the impact of
54 longline fisheries on adult survival in the black-footed albatross. *Journal of Applied Ecology*, **44(5)**, 942-952.

- 1 Veron, J.E., O. Hoegh-Guldberg, T.M. Lenton, J.M. Lough, D.O. Obura, P. Pearce-Kelly, C.R. Sheppard, M.
2 Spalding, M.G. Stafford-Smith, and A.D. Rogers, 2009: The coral reef crisis: The critical importance of <350
3 ppm CO₂. *Marine Pollution Bulletin*, **58(10)**, 1428-1436.
- 4 Villarini, G., F. Serinaldi, J.A. Smith, and W.F. Krajewski, 2009: On the stationarity of annual flood peaks in the
5 continental United States during the 20th century. *Water Resources Research*, **45(8)**, W08417.
- 6 Voigt, T., H.M. Füssel, I. Gärtner-Roer, C. Huggel, C. Marty, and M. Zemp (eds.), 2011: *Impacts of climate change
7 on snow, ice, and permafrost in Europe: Observed trends, future projections, and socio-economic relevance*.
8 European Environment Agency, Copenhagen, Denmark, pp. 117.
- 9 Vorogushyn, S. and B. Merz, 2012: What drives flood trends along the Rhine River: climate or river training?
10 *Hydrology and Earth System Sciences Discussions*, **9**, 13537-13567.
- 11 Vuille, M., B. Francou, P. Wagnon, I. Juen, G. Kaser, B.G. Mark, and R.S. Bradley, 2008: Climate change and
12 tropical Andean glaciers: Past, present and future. *Earth Science Reviews*, **89(3-4)**, 79-96.
- 13 Walker, M.D., C.H. Wahren, R.D. Hollister, G.H.R. Henry, L.E. Ahlquist, J.M. Alatalo, M.S. Bret-Harte, M.P.
14 Calef, T.V. Callaghan, A.B. Carroll, H.E. Epstein, I.S. Jónsdóttir, J.A. Klein, B. Magnússon, U. Molau, S.F.
15 Oberbauer, S.P. Rewa, C.H. Robinson, G.R. Shaver, K.N. Suding, C.C. Thompson, A. Tolvanen, Ø. Totland,
16 P.L. Turner, C.E. Tweedie, P.J. Webber, and P.A. Wookey, 2006: Plant community responses to experimental
17 warming across the tundra biome. *Proceedings of the National Academy of Sciences of the United States of
18 America*, **103(5)**, 1342-1346.
- 19 Walther, G., A. Roques, P.E. Hulme, M.T. Sykes, P. Pyšek, I. Kühn, M. Zobel, S. Bacher, Z. Botta-Dukát, H.
20 Bugmann, B. Czúcz, J. Dauber, T. Hickler, V. Jarošík, M. Kenis, S. Klotz, D. Minchin, M. Moora, W.
21 Nentwig, J. Ott, V.E. Panov, B. Reineking, C. Robinet, V. Semenchenko, W. Solarz, W. Thuiller, M. Vilà, K.
22 Vohland, and J. Settele, 2009: Alien species in a warmer world: risks and opportunities. *Trends in Ecology &
23 Evolution*, **24(12)**, 686-693.
- 24 Wang, G., S. Sun, and R. Mei, 2011: Vegetation dynamics contributes to the multi-decadal variability of
25 precipitation in the amazon region. *Geophysical Research Letters*, **38**, L19703.
- 26 Wassenaar, T., P. Gerber, P.H. Verburg, M. Rosales, M. Ibrahim, and H. Steinfeld, 2007: Projecting land use
27 changes in the Neotropics: The geography of pasture expansion into forest. *Global Environmental Change-
28 Human and Policy Dimensions*, **17(1)**, 86-104.
- 29 Wassmann, P. and T.M. Lenton, 2012: Arctic tipping points in a Earth system perspective. *Ambio*, **41**, 1-9.
- 30 Wassmann, R., S.V.K. Jagadish, S. Heuer, A. Ismail, E. Redona, R. Serraj, R.K. Singh, G. Howell, H. Pathak, and
31 K. Sumfleth, 2009: Climate change affecting rice production. The physiological and agronomic basis for
32 possible adaptation strategies. *Advances in Agronomy*, **101**, 59-122.
- 33 Weatherhead, E., S. Gearheard, and R.G. Barry, 2010: Changes in weather persistence: Insight from Inuit
34 knowledge. *Global Environmental Change*, **20**, 523-528.
- 35 Webb, A.P., 2006: *Analysis of Coastal Change and Erosion - Tebunginako Village, Abaiang, Kiribati*. SOPAC
36 Technical Paper ER53. SOPAC, Suva Fiji.
- 37 Webb, A.P., 2007: *Assessment of Salinity of Groundwater in Swamp Taro (Cyrtosperma Chamissonis) Pulaka Pits
38 in Tuvalu*. In: SOPAC Technical Paper ER75. SOPAC Technical Paper ER75. SOPAC, Suva, Fiji.
- 39 Webb, L.B., P.H. Whetton, J. Bhend, R. Darbyshire, P.R. Briggs, and E.W.R. Barlow, 2012: Earlier wine-grape
40 ripening driven by climatic warming and drying and management practices. *Nature Climate Change*, **2(4)**,
41 259-264.
- 42 Webster, P.J., G.J. Holland, J.A. Curry, and H.R. Chang, 2005: Changes in tropical cyclone number, duration, and
43 intensity in a warming environment. *Science*, **309**, 1844-1846.
- 44 Webster, P.J., V.E. Toma, and H.M. Kim, 2011: Were the 2010 Pakistan floods predictable? *Geophysical Research
45 Letters*, **38(4)**, L04806.
- 46 Welch, J.R., J.R. Vincent, M. Auffhammer, P.F. Moya, A. Dobermann, and D. Dawe, 2010: Rice yields in
47 tropical/subtropical Asia exhibit large but opposing sensitivities to minimum and maximum temperatures.
48 *Proceedings of the National Academy of Sciences of the United States of America*, **107(33)**, 14562-14567.
- 49 Welker, C. and E. Faust, 2013: Tropical cyclone-related socio-economic losses in the western North Pacific region.
50 *Natural Hazards and Earth System Sciences*, **13**, 115-124.
- 51 Wernberg, T., B.D. Russell, P.J. Moore, S.D. Ling, D.A. Smale, A. Campbell, M.A. Coleman, P.D. Steinberg, G.A.
52 Kendrick, and S.D. Connell, 2011: Impacts of climate change in a global hotspot for temperate marine
53 biodiversity and ocean warming. *Journal of Experimental Marine Biology and Ecology*, **400(1)**, 7-16.

- 1 Wernberg, T., D.A. Smale, F. Tuya, M.S. Thomsen, T.J. Langlois, T. de Bettignies, S. Bennett, and C.S. Rousseaux,
2 2012: An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot. *Nature*
3 *Climate Change*, **3**, 78-82.
- 4 Wernberg, T., B.D. Russell, M.S. Thomsen, C.F.D. Gurgel, C.J.A. Bradshaw, E.S. Poloczanska, and S.D. Connell,
5 2011: Seaweed communities in retreat from ocean warming. *Current Biology*, **21(21)**, 1828-1832.
- 6 Westerling, A.L., H.G. Hidalgo, D.R. Cayan, and T.W. Swetnam, 2006: Warming and earlier spring increases
7 Western U.S. forest fire activity. *Science*, **313(5789)**, 940-943.
- 8 Wezel, A. and A.M. Lykke, 2006: Woody vegetation change in Sahelian West Africa: Evidence from local
9 knowledge. *Environment, Development, and Sustainability*, **8**, 553-567.
- 10 White, I., T. Falkland, T. Metutera, E. Metai, M. Overmars, P. Perez, and A. Dray, 2007a: Climatic and human
11 influences on groundwater in low atolls. *Vadose Zone Journal*, **6(3)**, 581-590.
- 12 White, I., T. Falkland, P. Perez, A. Dray, T. Metutera, E. Metai, and M. Overmars, 2007b: Challenges in freshwater
13 management in low coral atolls. *Journal of Cleaner Production*, **15(16)**, 1522-1528.
- 14 White, I. and T. Falkland, 2010: Management of freshwater lenses on small Pacific islands. *Hydrogeology Journal*,
15 **18(1)**, 227-246.
- 16 White, S., 2011: *The climate of rebellion in the early modern Ottoman Empire*. Cambridge University Press, New
17 York, USA, pp. 354.
- 18 Wilbanks, T.J., P. Romero Lankao, M. Bao, F. Berkhout, S. Cairncross, J.-. Ceron, M. Kapshe, R. Muir-Wood, and
19 R. Zapata-Marti, 2007: Industry, settlement and society. In: *Climate Change 2007: Impacts, Adaptation and*
20 *Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental*
21 *Panel on Climate Change*. [Parry, M.L., Canziani, O.F., Palutikof, J.P., P. Van der Linden, and C.E.
22 Hanson (eds.)]. Cambridge University Press, Cambridge, UK, pp. 357-390.
- 23 Wilkinson, C. and G. Hodgson, 1999: Coral reefs and the 1997-1998 mass bleaching and mortality. *Nature and*
24 *Resources*, **35(2)**, 16-25.
- 25 Wohling, M., 2009: The problem of scale in indigenous knowledge: a perspective from Northern Australia. *Ecology*
26 *and Society*, **14(1)**, 1-14.
- 27 Wolfe, D., L. Ziska, C. Petzoldt, A. Seaman, L. Chase, and K. Hayhoe, 2008: Projected change in climate thresholds
28 in the Northeastern U.S.: implications for crops, pests, livestock, and farmers. *Mitigation and Adaptation*
29 *Strategies for Global Change*, **13(5)**, 555-575.
- 30 Wolski, P., D. Stone, M. Tadross, M. Wehner, and B. Hewitson, 2013: Attribution of floods in the Okavango Basin,
31 Southern Africa. *Journal of Hydrology*, (submitted).
- 32 Woo, M.K., P. Modeste, L. Martz, J. Blondin, B. Kochtubajda, D. Tutcho, J. Gyakum, A. Takazo, C. Spence, J.
33 Tutcho, P. Di Cenzo, G. Kenny, J. Stone, I. Neyelle, G. Baptiste, M. Modeste, B. Kenny, and W. Modeste,
34 2007: Science meets traditional knowledge: Water and climate in the Sahtu (Great Bear Lake) region,
35 Northwest Territories, Canada. *Arctic*, **60**, 37-46.
- 36 Woodworth, P.L., M. Menéndez, and W. Roland Gehrels, 2011: Evidence for century-timescale acceleration in
37 mean sea levels and for recent changes in extreme sea levels. *Surveys in Geophysics*, **32(4)**, 603-618.
- 38 Wootton, J.T. and C.A. Pfister, 2012: Carbon System Measurements and Potential Climatic Drivers at a Site of
39 Rapidly Declining Ocean pH. *PLoS ONE*, **7(12)**, e53396.
- 40 Wootton, J.T., C.A. Pfister, and J.D. Forester, 2008: Dynamic patterns and ecological impacts of declining ocean pH
41 in a high-resolution multi-year dataset. *Proceedings of the National Academy of Sciences of the United States*
42 *of America*, **105(48)**, 18848-18853.
- 43 WorldBank, 2011: *The World Bank supports Thailand's post-floods recovery effort*. [online] Available from:
44 <http://go.worldbank.org/TCFEHXJML0>, The World Bank, Washington.
- 45 Wright, B.D., 2011: The economics of grain price volatility. *Applied Economic Perspectives and Policy*, **33(1)**, 32-
46 58.
- 47 Wulf, H., B. Bookhagen, and D. Scherler, 2012: Climatic and geologic controls on suspended sediment flux in the
48 Sutlej River Valley, western Himalaya. *Hydrology and Earth System Sciences*, **16(7)**, 2193-2217.
- 49 Xin, W., L. Shiyin, G. Wanqin, and X. Junli, 2008: Assessment and simulation of glacier lake outburst floods for
50 Longbasaba and Pida Lakes, China. *Mountain Research and Development*, **28(3)**, 310-317.
- 51 Xu, K., J.D. Milliman, Z. Yang, and H. Xu, 2008: Climatic and anthropogenic impacts on water and sediment
52 discharges from the Yangtze River (Changjiang), 1950-2005. In: *Large rivers: Geomorphology and*
53 *management*. [Gupta, A. (ed.)]. John Wiley & Sons, Chichester, UK, pp. 609-626.

- 1 Yamano, H., H. Kayanne, T. Yamaguchi, Y. Kuwhara, H. Yokoki, H. Shimazaki, and M. Chicamori, 2007: Atoll
2 island vulnerability to flooding and inundation revealed by historical reconstruction. Fongafale island, funafuti
3 atoll, tuvalu. *Global and Planetary Change*, **57(3-4)**, 407-416.
- 4 Yang, Z., J. Gao, L. Zhao, X. Xu, and H. Ouyang, 2012: Linking thaw depth with soil moisture and plant
5 community composition: effects of permafrost degradation on alpine ecosystems on the Qinghai-Tibet Plateau.
6 *Plant and Soil*, , 1-14.
- 7 Yao, T., L. Thompson, W. Yang, W. Yu, Y. Gao, X. Guo, X. Yang, K. Duan, H. Zhao, B. Xu, J. Pu, A. Lu, Y.
8 Xiang, D.B. Kattel, and D. Joswiak, 2012: Different glacier status with atmospheric circulations in Tibetan
9 Plateau and surroundings *Nature Climate Change*, **2**, 663-667.
- 10 Yiou, P., R. Vautard, P. Naveau, and C. Cassou, 2007: Inconsistency between atmospheric dynamics and
11 temperatures during the exceptional 2006/2007 fall/winter and recent warming in Europe. *Geophysical*
12 *Research Letters*, **34**, L21808.
- 13 Yli-Panula, E., D.B. Fekedulegn, B.J. Green, and H. Ranta, 2009: Analysis of airborne betula pollen in Finland; a
14 31-year perspective. *International Journal of Environmental Research and Public Health*, **6(6)**, 1706-1723.
- 15 Yonge, C.M. and A.G. Nichols, 1931: Studies on the physiology of corals: V. The effect of starvation in light and in
16 darkness on the relationship between corals and zooxanthellae. *Scientific Report of the Great Barrier Reef*
17 *Expedition*, **1**, 177-211.
- 18 You, L., M.W. Rosegrant, S. Wood, and D. Sun, 2009: Impact of growing season temperature on wheat productivity
19 in China. *Agricultural and Forest Meteorology*, **149(6)**, 1009-1014.
- 20 Zemp, M., M. Hoelzle, and W. Haeberli, 2009: Six decades of glacier mass-balance observations: a review of the
21 worldwide monitoring network. *Annals of Glaciology*, **50(50)**, 101-111.
- 22 Zemp, M., I. Roer, A. Käab, M. Hoelzle, F. Paul, and W. Haeberli, 2008: *Global glacier changes. Facts and figures*.
23 WGMS, Zurich, Switzerland, pp. 88.
- 24 Zeppel, H., 2012: Climate change and tourism in the Great Barrier Reef Marine Park. *Current Issues in Tourism*,
25 **15(3)**, 287-292.
- 26 Zhang, G.H., S.H. Fu, W.H. Fang, H. Imura, and X.C. Zhang, 2007: Potential effects of climate change on runoff in
27 the yellow river basin of China. *Transactions of the American Society of Agricultural and Biological*
28 *Engineers (ABASE)*, **50(3)**, 911-918.
- 29 Zhang, J., D. Gilbert, A. Gooday, L. Levin, S. Naqvi, J. Middelburg, M. Scranton, W. Ekau, A. Peña, and B.
30 Dewitte, 2010: Natural and human-induced hypoxia and consequences for coastal areas: synthesis and future
31 development. *Biogeosciences*, **7(5)**, 1443-1467.
- 32 Zhang, J.G., Y.L. Wang, Y.S. Ji, and D.Z. Yan, 2011: Melting and shrinkage of cryosphere in Tibet and its impact
33 on the ecological environment. *Journal of Arid Land*, **3(4)**, 292-299.
- 34 Zhang, S., X.X. Lu, D.L. Higgitt, C.T.A. Chen, J. Han, and H. Sun, 2008a: Recent changes of water discharge and
35 sediment load in the Zhujiang (Pearl River) Basin, China. *Global and Planetary Change*, **60(3)**, 365-380.
- 36 Zhang, T., J. Zhu, X. Yang, and X. Zhang, 2008b: Correlation changes between rice yields in North and Northwest
37 China and ENSO from 1960 to 2004. *Agricultural and Forest Meteorology*, **148(6)**, 1021-1033.
- 38 Zhang, D.D., H.F. Lee, C. Wang, B. Li, Q. Pei, J. Zhang, and Y. An, 2011: The causality analysis of climate change
39 and large-scale human crisis. *Proceedings of the National Academy of Sciences of the United States of*
40 *America*, **108(42)**, 17296-17301.
- 41 Zhang, Y., S. Liu, J. Xu, and D. Shangguan, 2008: Glacier change and glacier runoff variation in the Tuotuo River
42 basin, the source region of Yangtze River in western China. *Environmental Geology*, **56(1)**, 59-68.
- 43 Ziska, L., K. Knowlton, C. Rogers, D. Dalan, N. Tierney, M.A. Elder, W. Filley, J. Shropshire, L.B. Ford, C.
44 Hedberg, P. Fleetwood, K.T. Hovanky, T. Kavanaugh, G. Fulford, R.F. Vrtis, J.A. Patz, J. Portnoy, F. Coates,
45 L. Bielory, and D. Frenz, 2011: Recent warming by latitude associated with increased length of ragweed pollen
46 season in central North America. *Proceedings of the National Academy of Sciences of the United States of*
47 *America*, **108(10)**, 4248-4251.
- 48 Zongxing, L., H. Yuanqing, P. Tao, J. Wenxiong, H. Xianzhong, P. Hongxi, Z. Ningning, L. Qiao, W. Shijing, Z.
49 Guofeng, W. Shuxin, C. Li, D. Jiankuo, and X. Huijuan, 2010: Changes of climate, glaciers and runoff in
50 China's monsoonal temperate glacier region during the last several decades. *Quaternary International*, **218(1-**
51 **2)**, 13-28.
- 52 Zwiers, F.W., X. Zhang, and Y. Feng, 2011: Anthropogenic influence on long return period daily temperature
53 extremes at regional scales. *Journal of Climate*, **24(3)**, 881-892.
- 54

Table 18-1: Observed changes in ocean system properties and their effects, with confidence levels for detection and attribution to climate change, based on assessment in Chapter 6.2, 6.3, and summarized in 6.6.1 and Figure 6-16. Observed impacts related to temperature effects have been attributed to warming, although the relative contributions of regional climate variation and long-term global trends have not been quantified.

Process	confidence in		Context	Ref
	detection	attribution		
Impacts of ocean acidification on pelagic marine biota	<i>low</i>	<i>low</i>	e.g., reduction in foraminiferan, coccolithophores and pteropod shell weight. Attribution supported by experimental evidence and physiological knowledge	[1]
Expansion of midwater hypoxic zones	<i>high</i>	<i>med</i>	oxygen minimum zones (OMZs) caused by enhanced stratification and bacterial respiration due to effects of warming	[2]
Regional and local impacts of expanding OMZs	<i>medium</i>	<i>low - medium</i>	reduction of biodiversity, compression of oxygenated habitat for intolerant species, range expansion for tolerant taxa	[3]
Direct temperature effects on marine biota related to limited physiological tolerance ranges	<i>very high</i>	<i>high</i>	e.g., large scale latitudinal shifts of species distribution, changes in community composition; attribution supported by experimental and statistical evidence as well as physiological knowledge	[4]
Increase in global net primary production	<i>medium</i>	<i>low</i>	discrepancy between satellite observations and open ocean time-series sites; in higher latitudes, NPP is increasing due to sea ice decline and warming	[5]
Changes in microbial processes	<i>low</i>	<i>low</i>	<i>low confidence</i> and limited understanding of microbial processes, drivers and interactions	[6]
Large scale shifts in biogeochemical pathways	<i>low</i>	<i>low</i>	<i>low confidence</i> for large scale shifts in biogeochemical pathways such as oxygen production, carbon sequestration and export production, nitrogen fixation, climate-feedback by DMS production, nutrient recycling	[7]

Key references and traceable account for the statements above:[1] (Wootton *et al.*, 2008; De Moel *et al.*, 2009; Moy *et al.*, 2009; Bednaršek *et al.*, 2012); 6.2.2, 6.3.4; [2] (Stramma *et al.*, 2008; Stolper *et al.*, 2010); 6.2.2; [3] (Levin *et al.*, 2009; Ekau *et al.*, 2010; Stramma *et al.*, 2010; Stramma *et al.*, 2012), 6.3.3; [4] (Merico *et al.*, 2004; Perry *et al.*, 2005; Pörtner and Farrell, 2008; Beaugrand *et al.*, 2010; Alheit *et al.*, 2012); 6.2.2, 6.3.2; [5] (Behrenfeld *et al.*, 2006; Saba *et al.*, 2010; Arrigo and Van Dijken, 2011); 6.3.1; [6] 6.2.2.4; [7] 6.3.3;6.3.4;6.3.5

Table 18-2: Observed changes on marine species and ecosystems, with confidence levels for assessment of detection of changes and their attribution to climate change, based on assessment in Chapter 6.2, 6.3, summarized in 6.6.1 and Figure 6-16, and 30.4. Observed impacts related to temperature effects have been attributed to warming, although the relative contributions of regional climate variation and long-term global trends have not been quantified.

Process	confidence in		Context	Ref
	detection	attribution		
Range shifts of fish and macroalgae	<i>high</i>	<i>high</i>	changes in species biogeographical ranges to higher latitudes or greater depths	[1a]
Changes in community composition	<i>high</i>	<i>high</i>	due to effects of warming, hypoxia, and sea ice retreat	[1b]
Changes in abundance	<i>high</i>	<i>high</i>	observed in fish, corals and intertidal species	[1c]
Impacts on marine air breathers, e.g. walruses, penguins, and other sea-birds	<i>high</i>	<i>high</i>	observed effects include changing abundance, phenology, species distribution and turtle sex ratios, and are mostly mediated through changes in resource availability including prey	[2]
Impacts on warm-water reef-building corals	<i>very high</i>	<i>very high</i>	effects mostly attributed to warming and rising extreme temperatures, though ocean acidification may contribute	[3]
Increments in fish species richness in temperate and high latitude zones	<i>high</i>	<i>medium</i>	effect associated with loss of sea ice and latitudinal species shifts due to warming trends	[4]
Change in regional fishery catch potential due to species shifts	<i>high</i>	<i>high</i>	partly attributable to climate change, and to fishing pressure	[5]
Change in fishery catch potential following changes in net primary production	<i>low</i>	<i>low</i>	effect associated with warming	[6]

Key references and traceable account for the statements above: [1a,b,c] (Müller *et al.*, 2009; Stige *et al.*, 2010); 6.3.2, 30.4; [2] (Grémillet and Boulinier, 2009; McIntyre *et al.*, 2011); [3] (Hoegh-Guldberg, 1999; Hoegh-Guldberg *et al.*, 2007; Baker *et al.*, 2008; Veron *et al.*, 2009); [4] (Hiddink and ter Hofstede, 2008; Beaugrand *et al.*, 2010); 6.3.7, 6.5.2; [5] Figure 6-16.5a; [6] Figure 6-16 5b

Table 18-3: Confidence in detection and attribution of observed climate change effects across ocean regions, based on expert assessment in Chapter 30.5 (respective subsections are given below). Confidence levels assigned are for detection / attribution, respectively.

	high latitude spring bloom systems	equatorial upwelling	semi-enclosed seas	coastal boundary systems	Eastern boundary upwelling ecosystems	Subtropical gyres
	30.5.1	30.5.4	30.5.6	30.5.3	30.5.2	30.5.5
Regional warming	<i>very high-high / high</i>	<i>very high-high / very high-high</i>	<i>very high-high / high</i>	<i>very high / high</i>	<i>medium/medium- low*</i>	<i>very high-high / very high-high</i>
Declining oxygen	<i>medium / medium-low</i>	<i>medium / medium-low</i>	<i>medium / medium-low</i>	<i>high / medium</i>	<i>medium / medium</i>	<i>medium / low-medium</i>
Declining primary productivity	n/s	n/s	n/s	<i>medium / medium-low</i>	<i>medium-low / medium-low</i>	<i>medium / medium</i>
Shifts in phenology	<i>high / high-medium</i>	n/s	<i>medium / medium-low</i>	<i>medium / medium-low</i>	<i>medium / medium-low</i>	n/s
Changes in abundance and species distribution	<i>high / high</i>	n/s	<i>high / high</i>	<i>medium / medium-low</i>	<i>medium / medium-low</i>	n/s

* California and Canary systems only

Table 18-4: Illustrative selection of some recent extreme impact events for which the role of climate has been assessed in the literature. The table shows confidence assessments as to whether the associated meteorological events made a substantial contribution to the impact event, as well as confidence assessments of a contribution of anthropogenic emissions to the meteorological event. The assessment of confidence in the findings is not necessarily a conclusion of the listed literature but rather results from assessment of the literature. Assessment of the role of anthropogenic emissions in the impact event requires a multi-step evaluation. Partly based on Coumou and Rahmstorf (2012).

Year	Region	Extreme impact event		Meteorological Event	
		Impact / damage	Confidence in contribution of extreme weather event to observed damage	Meteorological event	Confidence in contribution of anthropogenic emissions to extreme weather event
2003	Europe	excess death toll exceeding 70,000 (Robine <i>et al.</i> , 2008)	<i>very high</i>	hottest summer in at least 500 years (Luterbacher <i>et al.</i> , 2004; Schär and Jendritzky, 2004; Stott <i>et al.</i> , 2004; Christidis <i>et al.</i> , 2010)	<i>high</i>
2005	North Atlantic / USA	1,700 deaths and over US\$100 Bn in damage (Beven <i>et al.</i> , 2008)	<i>very high</i>	record number of tropical storms, hurricanes and category 5 hurricanes since 1970 (Emanuel, 2005; Webster <i>et al.</i> , 2005; Trenberth and Shea., 2006; Pielke Jr <i>et al.</i> , 2008; Vecchi <i>et al.</i> , 2008; Landsea <i>et al.</i> , 2009; Knutson <i>et al.</i> , 2010)	<i>very low</i>
2006-2007	Europe	partial second flowering or extended flowering in 2006, early flowering in 2007 (Luterbacher <i>et al.</i> , 2007)	<i>high</i>	hottest record fall and winter in at least 500 years (Luterbacher <i>et al.</i> , 2007; Van Oldenborgh, 2007; Yiou <i>et al.</i> , 2007; Cattiaux <i>et al.</i> , 2009)	<i>medium</i>
2010	Western Russia	burned area > 12,500km (Müller, 2011)	<i>low</i>	hottest summer since 1500 (Barriopedro <i>et al.</i> , 2011; Dole <i>et al.</i> , 2011; Rahmstorf and Coumou, 2011; Otto <i>et al.</i> , 2012)	<i>medium</i>
2011	Thailand	prolonged (up to 2 month) inundation of urban and industrialized areas, insured loss US\$ 8-11 B, total loss ca. US\$45 (SwissRe, 2011; WorldBank, 2011)	<i>very high</i>	wettest monsoon on record in middle and upper Chao Phraya Basin (Van Oldenborgh <i>et al.</i> , 2012)	<i>very low</i>
2010	Colombia	exceptionally heavy rainfall and floods, 4 M people affected, US\$ 7.8 Bn total damage (Hoyos <i>et al.</i> , 2013)	<i>very high</i>	ENSO-related second and third highest SST in Caribbean on record in late 2010; second most active storm and hurricane season (Trenberth and Fasullo, 2012)	<i>low</i>
2010	Pakistan	worst ever known floods in the region, 2000 people killed, 20 M affected, total loss US\$ 40Bn (Hong <i>et al.</i> , 2011)	<i>very high</i>	Exceptionally high rainfall amounts over northern Pakistan with unusual atmospheric circulation patterns (Houze Jr <i>et al.</i> , 2011; Webster <i>et al.</i> , 2011; Galarnau <i>et al.</i> , 2012)	<i>very low</i>
2011	Queensland, Australia	>200,000 people affected, >30,000 homes flooded, damages and cost to economy US\$ 2.5 – 10 B (Van den Honert and McAneney, 2011; Hayes and Goonetilleke, 2012)	<i>very high</i>	2010 wettest year on record for Queensland, with extreme precipitation in January 2011 on saturated ground; record high Southern Oscillation Index in 2010 (Van den Honert and McAneney, 2011; Cai and Van Rensch, 2012; Hayes and Goonetilleke, 2012)	<i>low</i>

Table 18-5: Cases of regional livelihood impacts attributable with varying degree to weather- and climate related events, climate change or climate variability.

Impacted Population	Climate-related driver	Impact on livelihood	Reference
Small-scale farmers, Ghana	Drought	Landscape transformation, poverty	(Tschakert <i>et al.</i> , 2011)
Middle-class farmers, Australia	Drought	Landscape transformation, income loss from agriculture, social conflict, poverty	(Alston, 2011)
High Arctic native people	Warming	Changing ice and snow conditions, dwindling access to hunting grounds	(Ford, 2009a; Ford, 2009b, see also Box 18-5)
Urban populations in Maputo, Accra, Nairobi, Lagos, Kampala	Flood frequency and severity increase	Direct impacts on people and loss of physical assets (e.g. housing)	(Douglas <i>et al.</i> , 2008)
Industry workers in India	Temperature variability and heat waves	Limited ability to carry out physical work, health impacts	(Ayyappan <i>et al.</i> , 2009; Balakrishnan <i>et al.</i> , 2010; Dash and Kjellstrom, 2011)
Farmers in Subarnabad, Bangladesh	Sea-level rise	Salt water intrusion, shift from agriculture to shrimp farming, loss of agricultural livelihoods	(Pouliotte <i>et al.</i> , 2009)
Women farmers, Ghana	Rainfall-related climate variability	Pressure from husbands limiting involvement in agriculture, poverty	(Carr, 2008)
Cambodian rice farmers	Warming, rainfall-related climate variability	Shift in income generation patterns between men and women	(Resurreccion, 2011)
Poor children in Africa and Latin America	Weather and climate-related events	Food price shocks, reduced caloric intake, physical stunting, long-term effects such as reduced lifetime earnings	(Alderman, 2010)
Smallholder farmers in highlands of Bolivia	Rising temperatures and higher variability in weather and climate-related extremes	Stress on household resources due to need to respond to increasing plant pests, switching to other crop types or livestock	(McDowell and Hess, 2012)

Table 18-6: Observed impacts of climate change on snow, ice and mountains, and rivers and lakes, across eight major world regions, with confidence in detection/confidence in attribution to climate change stated for each impact. References to related chapters are given as well as key references underlying the assessment.

	Snow, Ice and Mountains	Rivers and Lakes
Africa	<p>Retreat of tropical highland glaciers in East Africa <i>very high / medium confidence</i> [22.5.1, (Mölg <i>et al.</i>, 2008; Taylor <i>et al.</i>, 2009; Mölg <i>et al.</i>, 2012)]</p>	<p>Reduced discharge in West African rivers <i>medium / low confidence</i> (d'Orgeval and Polcher, 2008; Dai <i>et al.</i>, 2009; Di Baldassarre <i>et al.</i>, 2010)</p> <p>Decreased flood frequency in Okavango delta <i>low / low confidence</i> (Wolski <i>et al.</i>, 2013)</p> <p>Lake surface warming and water column stratification increases in the Great Lakes and Lake Kariba <i>high / high confidence</i> [22.3.2.2, (Tierney <i>et al.</i>, 2010; Ndebele-Murisa <i>et al.</i>, 2011; Powers <i>et al.</i>, 2011)]</p>
Europe	<p>Retreating glaciers in the Alps <i>very high / high confidence</i>, (Bauder <i>et al.</i>, 2007; Paul and Haeberli, 2008; Zemp <i>et al.</i>, 2008; Zemp <i>et al.</i>, 2009)</p> <p>Increase in rock slope failures in Western Alps <i>high / medium confidence</i> [18.3.1.4; (Fischer <i>et al.</i>, 2012; Huggel <i>et al.</i>, 2012a)]</p>	<p>Changes in the occurrence of extreme river discharges and floods <i>low / very low confidence</i> (Schmocker-Fackel and Naef, 2010; Beniston <i>et al.</i>, 2011; Cutter <i>et al.</i>, 2012; Vorogushyn and Merz, 2012; Kundzewicz <i>et al.</i>, 2013)</p>
Asia	<p>Permafrost degradation in Siberia, Central Asia, and the Tibetan Plateau <i>high / high confidence</i> [Box 3-2, WGI AR5 Chapter 10.5.3, (Romanovsky <i>et al.</i>, 2010; Yang <i>et al.</i>, 2012)]</p> <p>Mountain glaciers across Asia are shrinking <i>high / medium confidence</i> [WGI AR5 Chapter 4.3.2-4.3.3; Box 3-1; (Bolch <i>et al.</i>, 2012; Cogley, 2012; Kääb <i>et al.</i>, 2012; Yao <i>et al.</i>, 2012; Gardner <i>et al.</i>, 2013; Stokes <i>et al.</i>, 2013)]</p>	<p>Changes in water availability in many Chinese rivers <i>high / low confidence</i> [24.4.1.2; (Casassa <i>et al.</i>, 2009; Zongxing <i>et al.</i>, 2010)]</p> <p>Increased runoff in many rivers due to shrinking glaciers in the Himalayas and Central Asia <i>high / high confidence</i> [24.4.1.2; Box 3-1; (Casassa <i>et al.</i>, 2009; Zongxing <i>et al.</i>, 2010; Shrestha and Aryal, 2011; Zhang <i>et al.</i>, 2011)]</p> <p>Surface water degradation in various parts of Asia partially related to climate change <i>medium / medium confidence</i> [24.4.1.2; (Prathumratana <i>et al.</i>, 2008; Delpla <i>et al.</i>, 2009; Huang <i>et al.</i>, 2009)]</p> <p>Earlier timing of maximum spring flood in Russian rivers <i>high / high confidence</i> [28.2.1.1; (Shiklomanov <i>et al.</i>, 2007; Tan <i>et al.</i>, 2011)]</p>
Australasia	<p>Reduction in glacier ice volume in New Zealand <i>high / low confidence</i> [WGI AR5 Chapter 4.3.3; Table 25-1; (Nicholls, 2006; Chinn <i>et al.</i>, 2012)]</p> <p>Significant decline in late-season snow depth at four alpine sites in Australia (1957-2002) <i>high / medium confidence</i> (Hennessy <i>et al.</i>, 2008)</p>	n/s

North America	<p>Observed snowpack show primarily decreasing trends in the amount of water stored in spring snowpack from 1960-2002 <i>high / high confidence</i> (Stewart <i>et al.</i>, 2005; Mote, 2006)</p>	<p>Observed shift to earlier peak flow in snow dominated rivers in Western North America <i>high / high confidence</i> [WGI AR5 Chapter 2.6.2; (Barnett <i>et al.</i>, 2008)]</p> <p>Runoff increases in the Midwestern and Northwestern US, decreases in Southern states <i>high / medium confidence</i> (Georgakakos <i>et al.</i>, 2013)</p>
South and Central America	<p>Retreat of tropical Andean glaciers in Venezuela, Colombia, Ecuador, Peru and Bolivia (second half of the 20th century) and glaciers and ice-fields in the extra tropical Andes (Central-South Chile and Argentina) <i>high / high confidence</i> [27.3.1.1; (Vuille <i>et al.</i>, 2008; Bradley <i>et al.</i>, 2009; Jomelli <i>et al.</i>, 2009; Poveda and Pineda, 2009; Rabatel <i>et al.</i>, 2013)]</p> <p>Increased landslide frequency due to heavy precipitation in SE South America <i>medium / low confidence</i> [27.2.1.1; (Donat <i>et al.</i>, 2013; Marengo <i>et al.</i>, 2013; Silva Dias <i>et al.</i>, 2013)]</p>	<p>Changes in extreme flows in Amazon River <i>high / medium confidence</i> [27.2.1.1; (Rodríguez-Morales <i>et al.</i>, 2010; Butt <i>et al.</i>, 2011; Wang <i>et al.</i>, 2011; Espinoza <i>et al.</i>, 2013)]</p> <p>Changes in discharge patterns in rivers in the Western Andes due to retreating glaciers and reduced snowpack; for major river basins in Colombia discharge has decreased during the last 30-40 years <i>high / high confidence</i> [27.3.1.1; (Vuille <i>et al.</i>, 2008; Bradley <i>et al.</i>, 2009; Jomelli <i>et al.</i>, 2009; Poveda and Pineda, 2009; Rabatel <i>et al.</i>, 2013)]</p> <p>Increased stream flow in Sub-basins of the La Plata River, attributed to increasing precipitation, but also to trends in land use changes that have reduced evapotranspiration <i>very high / high confidence</i> [27.3.1.1; (Pasquini and Depetris, 2007; Krepper <i>et al.</i>, 2008; Conway and Mahé, 2009; Krepper and Zucarelli, 2010; Doyle and Barros, 2011)]</p>
Polar Regions	<p>Decreasing Arctic sea ice cover in summer, reduction in glacier ice volume, due to warming <i>high / high confidence</i> [WGI AR5 Chapter 10.5.1.1; (ACIA, 2005; AMAP, 2011)]</p> <p>Decreasing snow cover duration across the entire Arctic <i>high / medium confidence</i> [28.2.3.4, 5; (AMAP, 2011; Callaghan <i>et al.</i>, 2011)]</p> <p>Widespread permafrost degradation, especially in the southern Arctic <i>high / high confidence</i> (AMAP, 2011; Olsen <i>et al.</i>, 2011)</p> <p>Increased ice mass loss along coastal Antarctica <i>medium / very low</i> [WGI AR5 Chapter 4.2.3, 4.4, 4.6, 10.5.2.1]</p>	<p>Increased river discharge for large circumpolar rivers (1997–2007) <i>high / low confidence</i> [28.2.1.1; (Overeem and Syvitsky, 2010)]</p> <p>Winter minimum flows have risen in most sectors of the Arctic due to enhanced groundwater input due to permafrost thawing <i>high / medium</i> [28.2.1.1; (Tan <i>et al.</i>, 2011)]</p> <p>Increasing surface lake water temperatures 1985–2009 and prolonged ice-free seasons, due to warming <i>medium / medium confidence</i> [28.2.1.1; (Callaghan <i>et al.</i>, 2010; Schneider and Hook, 2010)]</p> <p>Thermokarst lakes disappear due to permafrost degradation in the low Arctic, while new ones are being created in areas of formerly frozen peat <i>high / high confidence</i> (Riordan <i>et al.</i>, 2006; Marsh <i>et al.</i>, 2008; Prowse and Brown, 2010)</p>

Table 18-7: Observed impacts of climate change on terrestrial ecosystems, and occurrence of drought and wildfires, across eight major world regions, with confidence in detection/confidence in attribution to climate change stated for each impact. References to related chapters are given as well as key references underlying the assessment.

	Terrestrial Ecosystems	Drought and Wildfire
Africa	<p>Tree density decreases in Sahel and semi-arid Morocco <i>high / medium confidence</i> [22.3.2.1.2; (Gonzalez <i>et al.</i>, 2012; Le Polain de Waroux and Lambin, 2012)]</p> <p>Climate-driven range shifts of several southern plants and animals: South African bird species polewards; Madagascar reptiles and amphibians upwards; Namib aloe contracting ranges. <i>high / medium confidence</i> [22.3.2.1; (Foden <i>et al.</i>, 2007; Raxworthy <i>et al.</i>, 2008; Hockey and Midgley, 2009; Hockey <i>et al.</i>, 2011)]</p>	<p>Increased drought in the Sahel since 1970, partially wetter conditions since 1990 <i>medium / medium confidence</i> [22.2.2.1; (Hoerling <i>et al.</i>, 2006; Giannini <i>et al.</i>, 2008; Greene <i>et al.</i>, 2009; Seneviratne <i>et al.</i>, 2012)]</p> <p>Wildfires increase on Mt. Kilimanjaro due to warming and drying trends <i>medium / low confidence</i> [22.5.1; (Hemp, 2005)]</p>
Europe	<p>Earlier greening, earlier leaf emergence and fruiting in temperate and boreal trees <i>high / high confidence</i> [4.4.1.1; (Menzel <i>et al.</i>, 2006)]</p> <p>Increased colonization of alien plant species in Europe <i>medium / medium confidence</i> [4.2.4.7; (Walther <i>et al.</i>, 2009)]</p> <p>Earlier arrival of migratory birds in Europe since 1970 <i>medium / medium confidence</i> [4.4.1.1; (Møller <i>et al.</i>, 2008)]</p> <p>Upward shift in tree-line in Europe <i>medium / low confidence</i> [18.3.2.1; (Gehrig-Fasel <i>et al.</i>, 2007; Lenoir <i>et al.</i>, 2008)]</p>	<p>Increasing burnt forest areas during recent decades <i>high / high confidence</i> (Hoinka <i>et al.</i>, 2009; Koutsias <i>et al.</i>, 2012)</p>
Asia	<p>Changes in plant phenology and growth occur in many parts of Asia, observed from ground observations as well as from satellites (earlier greening), particularly in the North and the East <i>high / medium confidence</i> [24.4.2.2; 4.2.1, Box 4-1; (Ma and Zhou, 2012; Panday and Ghimire, 2012; Shrestha <i>et al.</i>, 2012; Ogawa-Onishi and Berry, 2013)]</p> <p>Many plant and animal species have shifted their distribution, particularly in the North of Asia, generally upwards in elevation or polewards <i>high / medium confidence</i> [24.4.2.2; (Moiseev <i>et al.</i>, 2010; Chen <i>et al.</i>, 2011; Jump <i>et al.</i>, 2012; Ogawa-Onishi and Berry, 2013)]</p> <p>Siberian larch forests are being invaded by pine and spruce during recent decades, Mongolian larches show decreasing growth <i>medium / low confidence</i> (Kharuk <i>et al.</i>, 2010; Dulamsuren <i>et al.</i>, 2011; Lloyd <i>et al.</i>, 2011)</p> <p>Advance of shrubs into the Siberian tundra <i>high / medium confidence</i> [28.2.3.1; (Blok <i>et al.</i>, 2011)]</p>	

Australasia	<p>Climate-related changes in genetics, growth distribution and phenology of many species (earlier emergence of butterflies, change in plant flowering dates, breeding times of bird, decline in body size of passerine birds etc.) <i>medium / medium confidence</i> [table 25-3; (Green, 2010; Kearney <i>et al.</i>, 2010; Keatley and Hudson, 2012)]</p> <p>Change in timing of migration of water and land birds in Australia possibly due to changes in precipitation rather than changes in temperature <i>low / low confidence</i> [25.6.1, Table 25-3; (Chambers, 2008)]</p> <p>Expansion of some wetlands and corresponding contraction of adjacent woodlands in SE Australia <i>medium / low confidence</i> [table 25-3; (Banfai and Bowman, 2007; Bowman <i>et al.</i>, 2010; Keith <i>et al.</i>, 2010)]</p>	<p>No significant change in drought occurrence in Australia (over 20th century, using rainfall only) or New Zealand (since 1972, using a soil water balance model) <i>medium-low / n.a.</i> [table 25-1. 3.5.1; (Cai <i>et al.</i>, 2009; Potter <i>et al.</i>, 2010; Kingsford and Watson, 2011; IPCC, 2012)]</p>
North America	<p>Species distribution shifts upward in elevation and northward in latitude across multiple taxa <i>high / medium confidence</i> [26.4.1, 2; (Kelly and Goulden, 2008)]</p> <p>Phenology changes <i>high / medium confidence</i> [26.4.2; (Parmesan, 2006)]</p> <p>Increases in tree mortality at regional scales and insect infestations in forests <i>medium / low confidence</i> [26.4.1; (Peng <i>et al.</i>, 2011)]</p>	<p>Increases in wildfire activity, including fire frequency and duration, length of fire season, and area burned <i>high / medium confidence</i> [Box 26.2; (Westerling <i>et al.</i>, 2006)]</p>
South and Central America	<p>Degrading and receding rainforest in the Amazon <i>high / very low confidence</i> [27.2.2.1; (Etter <i>et al.</i>, 2006; Nepstad <i>et al.</i>, 2006; Oliveira <i>et al.</i>, 2007; Wassenaar <i>et al.</i>, 2007; Killeen <i>et al.</i>, 2008; Nepstad and Stickler, 2008)]</p>	<p>Increase dryness for most of the west coast of South America and in the Andes between 35.6°S and 39.9°S since 1950 <i>medium / low confidence</i> [table 27-1; (Christie <i>et al.</i>, 2011; Dai, 2011)]</p>
Polar Regions	<p>Increase in shrub cover in tundra in North America and Eurasia <i>high / high confidence</i> [28.2.3.2; (Tape <i>et al.</i>, 2006; Walker <i>et al.</i>, 2006; Henry and Elmendorf, 2010; Blok <i>et al.</i>, 2011; Elmendorf <i>et al.</i>, 2012; Tape <i>et al.</i>, 2012)]</p> <p>Significant advance of Arctic tree-line in latitude and altitude, due to warming, although the pace is lower than expected due insect outbreaks, land use history <i>high / medium confidence</i> [28.2.3.3; (AMAP, 2011; Hedenås <i>et al.</i>, 2011; Van Bogaert <i>et al.</i>, 2011)]</p> <p>Snow-bed ecosystems and tussock tundra are retreating, due to prolonged thawing season and less precipitation in the form of snow <i>high / high confidence</i> [28.2.3.2; (Björk and Molau, 2007; Molau, 2010a; Hedenås <i>et al.</i>, 2011; Callaghan <i>et al.</i>, 2013)]</p> <p>Animal populations in the tundra being affected by increasing occurrence of ice layers in the annual snow pack due to rain-on-snow events <i>medium / medium confidence</i> (Callaghan <i>et al.</i>, 2013; Hansen <i>et al.</i>, 2013)</p> <p>Breeding area and population size of subarctic birds has changed, due to warming and shrub encroachment in the tundra <i>high / medium confidence</i> (Molau, 2010b; Callaghan <i>et al.</i>, 2013)</p>	<p>Increasing drought in high Arctic polar deserts <i>high / high confidence</i> [28.2.1.1; (Smol and Douglas, 2007)]</p> <p>Increased frequency of wildfires in the conifer forest at the southern fringe of the Arctic, due to increasing summer temperature <i>high / medium confidence</i> (Mann <i>et al.</i>, 2012)</p> <p>Tundra wildfires are increasing in frequency in the Low Arctic, due to increasing summer air temperature and subsequent surface drought <i>high / medium confidence</i> [28.2.3.6; (Mack <i>et al.</i>, 2011)]</p>

	<p>Plant species in the West Antarctic Peninsula and nearby islands have increased over the past 50 years <i>high / high confidence</i> [28.2.3.2; (Fowbert and Smith, 1994; Parnikoza <i>et al.</i>, 2009)]</p> <p>Increasing phytoplankton productivity in Signy Island lake waters <i>high / high confidence</i> [28.2.1.2; (Quayle <i>et al.</i>, 2002; Laybourn-Parry, 2003)]</p>	
Small Islands	<p>Upward trend in tree-lines and associated fauna on high elevation islands <i>low / low confidence</i> [29.3.2.1; (Benning <i>et al.</i>, 2002; Jump <i>et al.</i>, 2006)]</p> <p>Changes in tropical bird populations in Mauritius, due to changes in rainfall <i>medium / medium confidence</i> [29.3.2.1; (Senapathi <i>et al.</i>, 2011)]</p>	<p>Increased drought frequency in Seychelles and Southern Jamaica over past 30 years, decrease in rainfall over past 100 years in the Caribbean islands may have contributed to multiple water stress <i>low / very low confidence</i> [29.3.3.2; (Payet and Agricole, 2006; Cashman <i>et al.</i>, 2010; Gamble <i>et al.</i>, 2010)]</p>

Table 18-8: Observed impacts of climate change on marine ecosystems, and coastal processes, across eight major world regions, with confidence in detection / confidence in attribution to climate change stated for each impact. References to related chapters are given as well as key references underlying the assessment.

	Marine Ecosystems	Coastal Processes
Europe	<p>Poleward shifts in the distributions of zooplankton, fish, seabirds and benthic invertebrate, and conversion of polar into more temperate and temperate into more subtropical system Characteristics in NE Atlantic <i>high / high confidence</i> [Table 6-8, 6.3.2, 30.5.1, 18.3.3.1, (Beaugrand <i>et al.</i>, 2009; Philippart <i>et al.</i>, 2011)]</p> <p>Phenology changes and retreat of colder water plankton to the north in the Northeast Atlantic, with mean poleward movement of plankton reaching up to 200–250 km per decade between 1958–2005 <i>medium / medium confidence</i> [6.3.2, Table 6-8, Fig 6-16; (Beaugrand <i>et al.</i>, 2002; Edwards and Richardson, 2004; Beaugrand <i>et al.</i>, 2009; Philippart <i>et al.</i>, 2011)]</p> <p>Shift in distribution of Atlantic cod due to warming, interacting with regime shift and regional changes in Plankton Phenology in North Sea <i>high / medium confidence</i> [6.3.2, Fig 6-16; (Perry <i>et al.</i>, 2005; Pörtner <i>et al.</i>, 2008; Beaugrand <i>et al.</i>, 2009; Beaugrand <i>et al.</i>, 2010)]</p> <p>Decreasing abundance of eelpout (Helcom indicator species) in Wadden Sea <i>medium / high confidence</i> [6.3.2, Fig 6-16; (Pörtner and Knust, 2007)]</p>	

Asia	<p>Coral reefs and large seaweeds decline in tropical Asian and Japanese waters <i>high / high confidence</i> [24.4.3.2; (Krishnan <i>et al.</i>, 2011; Nagai <i>et al.</i>, 2011; Coles and Riegl, 2012)]</p> <p>Shift from sardines to anchovies in Japanese Sea <i>medium / medium confidence</i> [6.3.22, Fig 16-6; (Takasuka <i>et al.</i>, 2007; Takasuka <i>et al.</i>, 2008)]</p>	<p>Coastal erosion is accelerating in Arctic Asia, due to changes in permafrost, storm wave energy and sea-ice retreat <i>medium / low confidence</i> (Razumov, 2010; Lantuit <i>et al.</i>, 2011; Handmer <i>et al.</i>, 2012)</p>
Australasia	<p>Mass bleaching of corals in the Great Barrier Reef, changes in coral calcification rates and changes in coral disease dynamics (e.g., black band disease, white syndrome) <i>high / high confidence</i> [6.3.2, 25.6.2, Box 18-3; (Cooper <i>et al.</i>, 2008; De'ath <i>et al.</i>, 2012)]</p> <p>Multiple impacts of climate change on marine ecosystems from warming oceans, although other environmental changes may play a role. Examples are increase of growth rates in fish, range shifts of intertidal invertebrates, retreat of seaweeds, range shift in near-shore fish related to kelp decline, increasing abundance of northern marine species in Tasmania, declines in recruitment of rock lobster and of abalone, decline in growth rate and biomass of phytoplankton, retreat of macroalgae, southward expansion of some tropical seabirds in Australia <i>high / high confidence</i> (Thresher <i>et al.</i>, 2007; Figueira <i>et al.</i>, 2009; Ling <i>et al.</i>, 2009; Pitt <i>et al.</i>, 2010; Chambers <i>et al.</i>, 2011; Neuheimer <i>et al.</i>, 2011; Wernberg <i>et al.</i>, 2011; Wernberg <i>et al.</i>, 2011, see also Table 25-3)</p>	
North America	<p>Northwest Atlantic fish show northward range shift in response to warming since the 1960s, with some of the shifts being correlated with the Atlantic multidecadal Oscillation <i>high / medium confidence</i> [Table 6-8, Fig 6-16 (Nye <i>et al.</i>, 2009; Lucey and Nye, 2010; Nye <i>et al.</i>, 2011)]</p> <p>Earlier onset of migration of Pink Salmon in Alaska, collapse of spawning migration of Sockeye Salmon in Fraser River, BC, due to warming <i>high / high confidence</i> [(Eliason <i>et al.</i>, 2011; Kovach <i>et al.</i>, 2012)]</p> <p>Loss of biomass of midwater fish off California Coast <i>high / high confidence</i> [6.3.3, 6.6.3]</p>	<p>Coastal roads affected by flooding during high tides in the US, though contribution of climate change to relative sea-level rise uncertain <i>low / very low confidence</i> (Moser and Davidson, 2013)</p>
South and Central America	<p>Bleaching of coral reefs in the western Caribbean near the coast of Central America <i>very high / high confidence</i> [27.3.3.1, (Guzman <i>et al.</i>, 2008; Manzello <i>et al.</i>, 2008; Carilli <i>et al.</i>, 2009; Eakin <i>et al.</i>, 2010)]</p>	<p>Mangrove degradation on Northern South American coast <i>high / low confidence</i> (Alongi, 2008; Lampis, 2010; Polidoro <i>et al.</i>, 2010; Giri <i>et al.</i>, 2011)</p>
Polar Regions	<p>Many arctic and subarctic marine non-migratory mammals (walrus, seals, whales) are negatively affected by sea ice loss <i>high / high confidence</i> [28.2.2.1.3, (Laidre <i>et al.</i>, 2008; McIntyre <i>et al.</i>, 2011)]</p> <p>Reduced growth rate and body mass, lower survival and reproductive capacity of polar bears, linked to reduced off-shore range and sea-ice loss due to warming <i>high / high confidence</i> [28.2.2.1.2; (Amstrup <i>et al.</i>, 2010)]</p> <p>Arctic seabirds experience reduced reproductive success, due to earlier sea-ice break-up <i>medium / medium confidence</i> [28.2.2.1.1; (Gaston <i>et al.</i>, 2009; Grémillet and Boulinier, 2009)]</p>	<p>Increased coastal erosion in Arctic, due to prolonged ice-free season at shore, increased exposure to wave activity, and degrading permafrost <i>high / high confidence</i> [28.2.4, 5; 28.3.4, (Forbes, 2011)]</p>

	<p>Acidification of Southern Ocean waters has resulted in reduced thickness of foraminifera shells <i>medium / medium confidence</i> [6.3.4, 28.2.2.2, (Moy <i>et al.</i>, 2009)]</p> <p>Antarctic krill density in the Scotia Sea has declined by ca. 30 % since the 1980s, due to reduced winter sea ice extent and duration <i>medium / medium confidence</i> (Atkinson <i>et al.</i>, 2004; Trivelpiece <i>et al.</i>, 2011)</p> <p>Many Southern Ocean species of seals and seabirds, e.g. penguins and Albatross, show negative responses to warmer conditions <i>high / medium confidence</i> [28.2.2.2; (Croxall <i>et al.</i>, 2002; Patterson <i>et al.</i>, 2003; Jenouvrier <i>et al.</i>, 2005; Véran <i>et al.</i>, 2007; Forcada <i>et al.</i>, 2008; Trathan <i>et al.</i>, 2011)]</p>	
<p>Small Islands</p>	<p>Coral bleaching near many tropical small islands <i>high / high confidence</i> [29.3.1.2,3; (Alling <i>et al.</i>, 2007; Bruno and Selig, 2007; Oxenford <i>et al.</i>, 2008)]</p> <p>Degradation of mangroves, wetlands and seagrass in small islands, mostly due to disturbances and only a lesser extent possibly due to sea-level rise <i>high / low confidence</i> [29.3.1.4; (McKee <i>et al.</i>, 2007; Gilman <i>et al.</i>, 2008; Schlepner, 2008; Krauss <i>et al.</i>, 2010; Marbà and Duarte, 2010; Rankey, 2011)]</p> <p>Degradation of freshwater dependent ecosystems, due to saline intrusion following sea level rise and more frequent and intense hurricanes in the Florida Keys, USA <i>low / low confidence</i> [29.3.2.1; (Ross <i>et al.</i>, 2009; Goodman <i>et al.</i>, 2012)]</p>	<p>Shoreline erosion is widespread and increasing on many small islands, though impact of sea-level rise can presently not be discriminated from climate variability and local disturbance <i>high / low confidence</i> [29.3.1.1, 2; (Yamano <i>et al.</i>, 2007; Cambers, 2009; Novelo-Casanova and Suarez, 2010; Storey and Hunter, 2010; Ford, 2012)]</p> <p>More frequent inundation in low-lying flood prone areas, increasing appearance of freshwater lenses at surface <i>medium / low confidence</i> [29.3.1.1, 2;(Webb, 2006; Webb, 2007; Yamano <i>et al.</i>, 2007; Ballu <i>et al.</i>, 2011)]</p> <p>Increases in groundwater degradation, mostly driven by over-pumping or pollution. Limited evidence for saline intrusion due to sea level rise or overtopping <i>high / low confidence</i> [29.3.2.2; (White <i>et al.</i>, 2007a; White <i>et al.</i>, 2007b; Terry and Falkland, 2010; White and Falkland, 2010)]</p>

Table 18-9: Observed impacts of climate change on human systems across eight major world regions, with confidence in detection / confidence in attribution to climate change stated for each impact. References to related chapters are given as well as key references underlying the assessment.

	Impacts on Human Systems: Health, Food production, Infrastructure and Livelihoods
Africa	Malaria increases in Kenyan highlands , partly due to warming <i>low / low confidence</i> [22.3.5.2, (Prudhomme O'Meara <i>et al.</i> , 2010; Alonso <i>et al.</i> , 2011; Stern <i>et al.</i> , 2011)]
	Reduced productivity of Great Lakes and Lake Kariba , partly due to warming <i>high / low confidence</i> [23.3.2.2, 23.3.4.4; (Descy and Sarmento, 2008; Hecky <i>et al.</i> , 2010; Ndebele-Murisa <i>et al.</i> , 2011; Marshall, 2012)]
	Fruit-bearing trees in Sahel decline <i>medium / medium confidence</i> (Wezel and Lykke, 2006; Maranz, 2009)
Europe	Increased allergic sensitization to pollen in Northern Italy <i>very low / very low confidence</i> [11.3; (Ariano <i>et al.</i> , 2010)]
	Shift from cold-related mortality to heat-related mortality in England and Wales <i>medium / low confidence</i> [18.4.5; (Christidis <i>et al.</i> , 2010)]
	Stagnation of wheat yields in some countries in recent decades, due to warming and/or drought <i>high / medium confidence</i> (Brisson <i>et al.</i> , 2010; Kristensen <i>et al.</i> , 2011) Positive yield impacts for wheat, sugar beet and potato in the UK since 1980 <i>low / low confidence</i> [Fig 7-3; (Gregory and Marshall, 2012)]
Asia	Negative impacts on aggregate wheat yields in South Asia <i>medium/low confidence</i> [Fig.7-3, 7.2.1.1]
	Negative impacts on aggregate wheat and maize yields in China <i>low / low confidence</i> [Fig.7-3, 7.2.1.1]
	Increases in water-borne diseases have been linked to warming in Israel <i>low / low confidence</i> (Paz <i>et al.</i> , 2007)
Australasia	Wine-grape maturation has advanced in recent decades, partly due to warming <i>high / medium confidence</i> (Webb <i>et al.</i> , 2012)
North America	Yields of grains, forage, livestock and dairy have declined due to increasing temperatures unless where accompanied by increased precipitation; soil organic content has declined; salinity has increased; climate change has affected product quality <i>high / low confidence</i> [26.5.1; (Hayhoe <i>et al.</i> , 2004; Lin, 2007; Hatfield <i>et al.</i> , 2008; Wolfe <i>et al.</i> , 2008; Schlenker and Roberts, 2009; Craine <i>et al.</i> , 2010)]
	Reduced economic returns from agriculture following extreme heat and storms, particularly in Mexico <i>high / low confidence</i> [Boxes 26.1, 26.2, 26.4; (Swanson <i>et al.</i> , 2007; Chen and McCarl, 2009)]
	Direct and indirect economic impacts of climate extremes on industry through reduced supply of raw material, the production process, the transportation of goods, and the demand for certain products <i>high / medium confidence</i> [26.8; (Lazo <i>et al.</i> , 2011)]
	Damages from climate extremes on infrastructure, especially in the transport and housing sectors <i>medium / low confidence</i> [26.8; (Morton <i>et al.</i> , 2011)] Impacts of extreme events on economy, lives and livelihoods in urban and rural settlements <i>medium / low confidence</i> [26.7.1, 26.7.2; (Kurz <i>et al.</i> , 2008; MacDonald, 2010)]
Central and South America	Increase in frequency and extension of dengue fever and yellow fever <i>high / low confidence</i> [27.3.7.1; (Teixeira <i>et al.</i> , 2009; Rodríguez-Morales <i>et al.</i> , 2010; Jentes <i>et al.</i> , 2011)]
	Increase in frequency and extension of malaria <i>high / medium confidence</i> [27.3.7.1; (Rodríguez-Morales <i>et al.</i> , 2010; Poveda <i>et al.</i> , 2011)]
	Increase in agricultural yields in Southeastern South America <i>high / medium confidence</i> [27.3.4.1; (Magrin <i>et al.</i> , 2007; Barros, 2010)]
	Expansion of agricultural areas in climatically marginal regions of Argentina <i>high / low confidence</i> [27.3.4; (Barros, 2010; Hoyos <i>et al.</i> , 2012)] Reduction in fish stocks (Peru, Colombia, Brazil) <i>low / very low confidence</i> [27.3.7.1; (Allison <i>et al.</i> , 2009; Freire and Pauly, 2010)]
Polar Regions	Impact on livelihoods of high Arctic indigenous peoples <i>medium / medium confidence</i> [18.4.5, Box 18-5; (Ford <i>et al.</i> , 2009; Beaumier and Ford, 2010; Pearce <i>et al.</i> , 2010)]

Small Islands	Degradation of coastal fisheries <i>high / low confidence</i> [29.3.3; 18.4.5; 30.4, 6.3.2] Casualties and damage from extreme weather events <i>very high / very low confidence</i> [29.3.3]
----------------------	--

Table 18-10: Reported confidence of detection of observed changes in “Unique and Threatened Systems” and attribution of observed changes to observed climate change. Assessments are for global or the indicated regional domain, with regional variations indicated in the final column.

Impact on unique or threatened system	Source for overarching assessment	Confidence in		Regional details, with corresponding subsections and confidence in detection/confidence in attribution
		detection	attribution	
Shrinking / receding glaciers	18.3.1	<i>very high</i>	<i>high</i>	Tropical Highland Glaciers, East Africa (22.5.1): <i>very high / medium</i> European Alps: <i>very high / high</i> Asia (24.4.2.2): <i>high / medium</i> New Zealand (Table 25-1): <i>high / low</i> North America: <i>high / high</i> Andean tropical glaciers and extra tropical icefields (27.3.1.1): <i>high / high</i> Polar glaciers: <i>high / high</i>
Mountain and lowland permafrost degradation	18.3.1.2	<i>high</i>	<i>medium</i>	Europe: <i>medium / medium</i> Siberia, Central Asia and Tibetan Plateau (24.4.2.2): <i>high / high</i> North America: <i>medium / medium</i> Southern Arctic (28.2.1.1): <i>high / high</i>
Arctic coastal permafrost degradation	18.3.1.2, 24.4.3.2, 28.2.1.1	<i>high</i>	<i>high</i>	
Warm water reef building corals – increased mortality and bleaching	5.3.1.6, 6.3.2.1.2, 30.3.1.1, 30.5, 30.8.2, Box 18-3	<i>very high</i>	<i>high</i>	Asia (24.4.3.2): <i>high / high</i> Australian Great Barrier Reef: <i>high / high</i> Western Caribbean (27.3.3.1): <i>very high / high</i> Coastal reefs surrounding small islands (29.3.1): <i>high / high</i>
Shrub increase and permafrost decay in Arctic tundra, onset of biome shift	18.3.2.4, 28.2.3.2	<i>high</i>	<i>high</i>	
Increased tree mortality in the Amazon; onset of biome shift	4.3.3	<i>low</i>	<i>very low</i>	
Increased tree mortality in boreal forests; onset of biome shift	4.3.3	<i>low</i>	<i>low</i>	
Changes in Arctic marine ecosystems	18.5.7, 28.2.2.1	<i>high</i>	<i>medium</i>	
Decline in extent of mangroves and coastal wetlands	27.3.3.1, 29.3.1.4	<i>very high</i>	<i>very low</i>	Mangrove degradation northern South American coast (27.3.3.1): <i>high / low</i> Impacts on mangroves and sea grasses in coastal areas of small islands (29.3.1.4): <i>high / low</i>
Livelihood impacts on indigenous Arctic peoples	18.4.7, Box 18-5	<i>medium</i>	<i>medium</i>	

Table 18-11a: Confidence in detection of changes in extreme weather hazard and attribution of a major role of changing greenhouse gas concentrations in the observed changes.

Observed changes in extreme weather or hydrologic hazard	Confidence in		Assessment*
	detection	attribution	
Increasing frequency and intensity of extreme hot events	<i>high</i>	<i>high</i>	WGI AR5 Chapter 10.6.1
Decreasing frequency and intensity of extreme cold events	<i>high</i>	<i>high</i>	WGI AR5 Chapter 10.6.1
Increases in the number and intensity of heavy precipitation events	<i>medium</i>	<i>medium</i>	WGI AR5 Chapter 10.6.1
Increases in tropical cyclone activity	<i>low</i>	<i>low</i>	WGI AR5 Chapter 10.6.1
Changes in tornadoes or hail	<i>very low</i>	<i>very low</i>	Chapter 10.6.1
More intense and longer dry spells in some low and mid latitude regions	<i>low</i>	<i>low</i>	WGI AR5 Chapter 10.6.1
Changes in inland flood magnitude and frequency	<i>low</i>	<i>low</i>	3.2.3, 18.3.1.1
Increasingly frequent coastal flooding	<i>medium</i>	<i>low</i>	29.3.1.1, 29.3.2.2, 5.3.3.3

*more detailed regional information consistent with AR5 WGI Chapter 10 assessment can be found in Seneviratne *et al.* (2012)


Table 18-11b: Confidence in detection of observed trends in damage in various systems and attribution of a major role of observed changes in extreme weather to those observed trends.

Observed changes in impacts of extreme weather	Relevant weather extremes	Confidence in		Assessment
		detection	attribution	
Increased coral bleaching	Hot surface waters	<i>very high</i>	<i>high</i>	5.3.1.6, 6.3.2, Box 18-3, 24.4.3.2, 27.3.3.1, 29.3.1, 30.3.1.1, 30.5. 30.8.2
Increased beach erosion in low and mid latitudes	High storm waves and surges	<i>high</i>	<i>very low</i>	5.3.1.2
Increased erosion of Arctic coastal bluffs	Lack of sea ice protection from wind storms	<i>high</i>	<i>high</i>	24.4.3.2, 28.2, 28.3.4
Increased South American landslide frequency	Heavy precipitation	<i>medium</i>	<i>low</i>	27.2.1.1
Cascading food web effects on Arctic grazers and predators	Higher frequency of rain-on-snow events	<i>high</i>	<i>high</i>	28.2.1.1
Increased drought in high Arctic deserts	Warm spells	<i>high</i>	<i>high</i>	28.2.1.1
Increased monetary losses due to extreme weather events	Storms, floods	<i>very high</i>	<i>very low</i>	10.7, 18.4.4.2

Table 18-12: Confidence in detection of impacts on aggregate impact measures and confidence in attribution of at least a minor role of climate change in those observed changes.

Global aggregated impact measure	Reference chapters	Confidence in	
		detection	attribution
Glacier ice volume reduction	18.3.1.3; 3.2.2	<i>very high</i>	<i>high</i>
Shift of permafrost boundary to higher altitudes and latitudes, and increase of active layer thickness	18.3.1.3	<i>high</i>	<i>medium</i>
Increase in terrestrial net primary production and C stocks	18.3.2.2	<i>high</i>	<i>low</i>
Negative yield impacts on global wheat against trends in technology, practice, and coverage	18.4.1.1, Fig 7-3	<i>very low</i>	<i>very low</i>
Negative yield impacts on global maize against trends in technology, practice, and coverage	18.4.1.1, Fig 7-3	<i>low</i>	<i>low</i>
Small increase in marine net primary production	Table 18-1	<i>medium</i>	<i>low</i>
Change in fishery productivity	Table 18-2; Fig 30.14, 6.6.1	<i>high</i>	<i>low</i>
Increase in monetary losses due to extreme weather	18.4.4.2 and 10.7	<i>very high</i>	<i>very low</i>

Figure 18-1: Schematic of the subject covered in this chapter. The Earth system can be divided into three broad interacting systems. Direct drivers of the human system on the climate system are denoted with a red arrow; some of these drivers may also directly affect natural systems. These effects can in turn influence other systems (dashed red arrows). Further influences of each of the systems on each other (confounding factors) that do not involve climate drivers are represented by blue arrows. Examples of drivers and their impacts are given in the table. Adapted from Stone *et al.* (2013).



	Example of Drivers	Example of Impacts	
		Direct impacts	Subsequent impacts
1	Emission of CO ₂	Warming	
1a			Altered crop yield
1b			Shift in species phenology
2	Emission of CO ₂	Carbon fertilization of plants	
2a			Increase in forestry yield
2b			Change in humidity
3	Pollution of river catchment	Decrease in fisheries	
4	Plague of crop pests	Decrease in crop yield	
5	Forest fire	Increased windiness	
6	El Nino event	More wildfires	
7	El Nino event	Fisheries	

Figure 18-2: A schematic diagram comparing approaches to attribution for an ecological system. The multi-step approach differs from the single-step approach in having a discontinuity between the attributed climate change and the observed weather driving the ecological model. Adapted from Stone *et al.* (2013).




Figure 18-3: Levels of confidence in detection and attribution of observed climate change impacts for freshwater systems over the past several decades, based on expert assessment contained in this section 18.3.1 and augmented by subsections of chapter 3 as indicated. Numbered symbols refer to: Freshwater systems (18.3.1.1): 1 groundwater depletion (Ch.3.2.4), 2 changing river flow (Ch. 3.2.3), 3 changing flood frequency or intensity (Ch. 3.2.3), 4 reduction in lake and river ice duration or thickness (Ch. 18.3.1.2); Cryosphere: 5 shrinking glaciers (Ch. 3.2.2, 18.3.1.2), 6 changes in glacier lakes (Ch. 18.3.1.1), 7 erosion and degradation of arctic coastal permafrost (Ch. 18.3.1.2), 8 degradation and thaw of lowland and mountain permafrost (Ch. 18.3.1.2), Soils and rock (18.3.1.3): 9 Increasing erosion (Ch.3.2.6), 10 changes in shallow landslides (Ch.3.2.6), 11 increasing frequency of Alpine rock failures




Figure 18-4: Confidence in detection and attribution of observed impacts on unique and threatened systems as a result of recent climate change. Assessments are for confidence in detection of a change and for confidence in attribution of a major role of observed climate change. Large letters denote conclusions for the full-scale system, while small letters denote conclusions for regional subsets of the systems. System changes include shrinking / receding glaciers, mountain and lowland permafrost degradation, Arctic coastal permafrost degradation, increased mortality and bleaching of warm water reef-building corals, increased shrub cover and permafrost decay in the northern tundra, increased tree mortality in the Amazon, increased tree mortality in boreal forests, changes in Arctic marine ecosystems, decline in extent of mangroves and coastal wetlands, and impacts on livelihoods of indigenous Arctic peoples. See Table 18-10 for details and references.




Figure 18-5: Left: confidence in detection of observed changes in extreme weather hazards and attribution of a major role of changing greenhouse gas concentrations in the observed changes. Right: confidence in detection of observed changes in damages related to extreme weather and attribution of major role to observed changes in extreme weather.






Figure 18-6: Confidence in detection of observed changes in natural systems (panel a) and human and managed systems (panel b) across regions, and confidence in attribution of such trends to observed climate change. Based on assessments developed in Tables 18-6 through 18-9.



Panel a: Regional Impacts on Natural Systems



Panel b: Regional Impacts on Human Systems

Figure 18-7: Confidence in detection of changes in aggregate impact measures and attribution of at least a minor role of climate change in those trends

