



Washington
State Department of
Agriculture

The Grayland Ditch

An Evaluation of the Effectiveness of Best Management Practices in Preventing Pesticides from Entering Cranberry Bog Drainage Ditches



March, 2014

Publication No. *AGR PUB 102-401*

Publication and Contact Information

This report is available on the Department of Agriculture's website at

<http://agr.wa.gov/FP/Pubs/NaturalResourcesAssessmentPubs.aspx>

Contact Information

Author: Rodney Baker
Natural Resource Assessment Section
P.O. Box 42560
Olympia, WA 98504-2560

Communication Officer
Hector Castro
Phone: 360.902.1815

Any use of product or firm names in this publication is for descriptive purposes only and does not imply endorsement by the author of the Department of Agriculture.

The Grayland Ditch

By

Rodney Baker,

Washington State Department of Agriculture
Natural Resource Assessment Section
Olympia, Washington 98504-2560

Acknowledgements

Kirk Cook, NRAS

Perry Beale, NRAS

Kelly McLain, NRAS

George Tuttle, NRAS

Desirée R. Whittenberg, WSDA Pesticide Management

This page is left blank intentionally

Table of Contents

List of Figures	6
List of Tables	6
Summary	8
Purpose and Goals	10
Study Area	10
Background	10
Methods	15
Sampling Design	15
Quality Objectives and Data Criteria	16
Field Procedures	16
Lab Procedures	17
Sample Results	19
Ambient Data	22
Description and Findings	30
References	33
Appendix A	36
Glossary	36
Acronyms and Abbreviations	37
Units of Measurement	38
Appendix B	39
Appendix C	42

List of Figures

Figure 1: Grays Harbor Ditch study area 12
Figure 2. 2012 Diazinon Concentrations 19
Figure 3. 2013 Diazinon Concentrations. 20
Figure 4. 2012 Chlorpyrifos Concentrations. 21
Figure 5. 2013 Chlorpyrifos Concentrations. 21
Figure 6. Discharge (Flow)..... 23
Figure 7. Water Temperature. 24
Chart 7. pH. 25
Figure 9. Turbidity..... 26
Figure 10. Conductivity. 27
Figure 11. Dissolved Oxygen. 28
Figure 12. Total Suspended Solids. 29

List of Tables

Table 1 . Sampling locations and descriptions for GHCCD-1 and PCDD-1..... 15
Table 2. Measurement quality objectives for conventional parameters measured by field meters or determined by a standard method..... 17
Table 3. Laboratory measurement quality objectives. 17
Table 4. Available water quality criteria. 18
Table 5. Summarized study results from 1996, 1998, and 2009 sampling sites..... 39
Table 6. Ambient Measurements (2013). 42
Table 6. Ambient Measurements (2013). 43
Table 6. Ambient Measurements (2013). 44

This page left blank intentionally

Summary

During 1996, 1998, 2002, and 2009, the Washington State Department of Ecology (Ecology) sampled for pesticides in surface water draining cranberry growing areas near Grayland on the Washington coast. Ecology found concentrations of three organophosphate pesticides in excess of water quality criteria: azinphos-methyl (Guthion™), chlorpyrifos (Lorsban™), and diazinon.

For the past few years, cranberry farmers have been implementing best management practices (BMPs) to reduce concentrations of organophosphate pesticides in adjacent surface waters. The 2009 Ecology study presented an evaluation of BMP effectiveness in reducing pesticide levels during peak application periods. Three sites were sampled in both Grays Harbor County Drainage Ditch No. 1 and Pacific County Drainage Ditch No. 1. Samples were collected one week prior to pesticide application, during the week of peak application, and two weeks following application.

Although it appeared some improvements had been made, concentrations of chlorpyrifos and diazinon continued to periodically exceed state water quality standards. A majority of the cranberry growers in the Grayland area are employing BMPs, but a small number of growers have yet to implement them. Lack of detectable concentrations of azinphos-methyl in the 2009 study was likely due to the cancellation of registered uses on cranberries. Increased usage of chlorpyrifos and diazinon, due to the loss of azinphos-methyl, did not consistently show corresponding increases in detectable concentrations in the current sampling project.

Chlorpyrifos and diazinon have been continually detected above state water quality standards and federal National Recommended Water Quality Criteria (NRWQC) over the last 17 years. This requires the Washington State Department of Agriculture (WSDA) to follow the guidance described in the *Washington State Pesticide Management Strategy for Water Quality Protection* (Cook and Cowles, 2009). WSDA is the State Lead Agency for pesticide registration in Washington as delegated by EPA under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA).

WSDA began sampling the water in the Grayland Ditch in 2012, continued to sample in 2013 and will continue to sample in 2014 and 2015.

Purpose and Goals

Study Area

A major cranberry growing area in Washington State is located between Grayland (Grays Harbor County) and North Cove (Pacific County) on the Washington Coast. Grays Harbor County and Pacific County each manage a ditch system, collectively known as the Grayland Ditch. The Grayland Ditch drains these cranberry growing areas as well as residential properties.

Precipitation runoff from woodland areas east and upslope of the cranberry bogs also feeds into the ditches. These ditches originate near the Grays Harbor/Pacific County line, west of Highway 105 (Figure 1).

Grays Harbor County Drainage Ditch No. 1 (GHCDD-1) flows north for approximately 2.8 miles, draining water from around the county line through the Grayland area, and discharges to the South Bay of Grays Harbor. Pacific County Drainage Ditch No. 1 (PCDD-1) flows south for approximately 5 miles, from the county line, and discharges to the North Cove of Willapa Bay. Figure 1 shows the locations of GHCDD-1 and PCDD-1.

Background

In Washington State, drainage ditches such as the Grayland Ditch are designated as surface waters of the state. As such, state water quality standards apply (Chapter 173-201A WAC). Washington State water quality standards are the basis for protecting and regulating the quality of surface waters. Section 303(d) of the federal Clean Water Act requires Washington State to every three years prepare a list of all surface waters in the state for which beneficial uses – such as for drinking, recreation, aquatic habitat, or industrial use – are impaired by pollutants. Ecology’s assessment of waters to be placed on the 303(d) list is guided by federal laws, state water quality standards, and state policy.

In 1994 and 1995, Ecology identified several pesticides frequently detected at concentrations exceeding Washington State or federal water quality standards in the Grayland Ditch (Davis et

al., 1997). The resulting 303(d) listings include chlorpyrifos and diazinon, as well as DDT and its metabolites (4,4'-DDD, 4,4'-DDE).¹

Azinphos-methyl was found above federal water quality criteria, but there is not a corresponding state water quality criterion. As a result, azinphos-methyl was not put on the Washington State 303(d) list. However, due to the number of detections above the federal NRWQC, azinphos-methyl was still considered a chemical in need of further investigation.

Local cranberry growers responded to these listings by sponsoring research and development of BMPs for their growing operations. Use of BMPs to reduce pesticide levels in Grayland Ditch began in 1994 (Pacific Conservation District and the Pacific Coast Cranberry Research Foundation, 1999).


A 1996 study conducted by Ocean Spray (Frantz et al., 1997) investigated the effectiveness of several different BMPs for use in the Grayland area cranberry bogs using rubidium chloride (RbCl) as a surrogate for pesticides. The study was able to show cribbing (lining) and covering of ditches within and bordering cranberry bogs was the most effective BMP. Cribbing and covering reduced the interior and perimeter ditch concentrations of RbCl to non-detection levels at the study's detection limit of 100 µg/L.

At the time of the BMP effectiveness study, approximately one-half of the growers were implementing BMPs. By 2000, approximately 95% of the Grayland area cranberry growers were using at least one BMP for reducing pesticide pollution (Rountry, 2008).

To track the progress of the implemented BMPs, Ecology conducted studies to evaluate the reduction in pesticide concentrations (Anderson and Davis, 2000; Coots, 2003, Anderson 2009). Results of the studies showed reductions in pesticide levels present in the ditches. However, the concentrations of pesticides continued to exceed water quality standards.

¹ DDT registration was cancelled in the United States in 1972 and it is considered a legacy pesticide.

Figure 1: Grays Harbor Ditch study area



In 2010, WSDA staff were contacted by the cranberry growers on the Grayland Ditch system and asked to provide technical expertise to address the ongoing water quality concerns. In 2011, WSDA staff gathered information from the growers on pesticide application timing, application rates, and rate of BMP implementation across farms. Using this information, WSDA staff conducted baseline sampling during the spring 2012 application period (April 15 – May 31). The results of this initial sampling showed most portions of the ditch exceeded water quality standards for chlorpyrifos and diazinon. Based on these test results, WSDA followed up with a pesticide application records request to all cranberry growers in the Grayland Ditch area for the 2012 pesticide application season.

It was determined the most effective way to make reductions was to continue supporting development and implementation of BMPs and to re-evaluate pesticide concentrations with future sampling. To help with implementation of BMPs, the Natural Resource Conservation Service, Pacific and Grays Harbor Conservation Districts, and other organizations have provided growers with technical assistance and grants.

The re-evaluation of pesticide concentrations in the Grayland Ditch is now being conducted by WSDA. Under the guidance of the *Washington State Pesticide Management Strategy for Water Quality Protection* and the authority of WAC 16-232-400/450, WSDA began a three year water sampling project in 2013. The project will continue through the 2015 pesticide application season.

As a State Lead Agency, WSDA reviews pesticides occurring in surface and groundwater at concentrations approaching or exceeding human health or ecological reference points. If the pesticide(s) are determined to pose a risk to surface water and/or groundwater quality, the development and implementation of management measures is required. A response matrix detailing WSDA's approach to assessing pesticide detections in surface and groundwater is provided in Appendix One of the *Washington State Pesticide Management Strategy for Water Quality Protection*.

The water quality results within cranberry growing area of Grayland have already passed response levels 1 and 2 of the response matrix contained within the “*Strategy*”. This places the response at level 3 and may result in implementation of mandatory BMPs. As stated in Bicki et al. (2003), ditch lining and covering BMPs are effective when properly implemented.

Under the *Washington State Pesticide Management Strategy for Water Quality Protection*, WSDA will coordinate with Ecology and, if needed, Washington State Department of Health to implement the management strategies.

Methods

Sampling Design

The sampling was conducted by WSDA’s Natural Resource Assessment Section (NRAS) staff.

WSDA sampling sites differ somewhat from previous sites sampled by Ecology. A new site, Tide Gate North, was added to the monitoring program to better capture runoff from all cranberry bogs draining into GHCDD. In addition the Grange Road and Jacobson Road sites were removed from the study. The Schmid Road, County Line Road, Heather Road, and Tide Gate South sites remained unchanged for a total of five sampling locations (see Figure 1 and Table 1).

Table 1 . Sampling locations and descriptions for GHCDD-1 and PCDD-1.

Station Name	Latitude	Longitude	Description
GHCDD-1			
Tide Gate North	46.83611	-124.07705	Upstream side of bridge on GHCDD-1
Schmid Road	46.8161	-124.0916	Upstream side of bridge on GHCDD-1
County Line Road	46.7938	-124.0866	Upstream side of culvert on GHCDD-1
PCDD-1			
Heather Road	46.7758	-124.0777	Upstream side of bridge on PCDD-1
Tide Gate South	46.7372	-124.0688	Upstream of tide gate on PCDD-1

Datum NAD 1983

For 2012, WSDA obtained only water samples. Beginning in 2013, WSDA began measuring ambient surface water quality for temperature, pH, conductivity, dissolved oxygen (DO), and turbidity. Stream flow measurements were also taken beginning in 2013. Water samples were collected for laboratory analysis of organophosphate pesticides, total suspended solids (TSS) and biological oxygen demand (BOD).

The timing of the sampling focused on the most intensive application periods for insecticides in the cranberry bogs. Intensive application of insecticides typically occurs during the spring and again in the middle of July. Many factors influence the start of pesticide applications. To achieve the target sample collection windows, WSDA staff queried the cranberry growers

regarding pesticide application timing, application rates, and rate of BMP implementation across farms. WSDA also conducted a pesticide application record request for the 2012 pesticide application season. In typical years insecticide applications occur twice, once in late spring (April-May) and again immediately after removal of the honey bees used for pollination. Removal of the bees is weather dependent, but in the past has occurred around the middle of July.

The sampling schedule was determined using prior studies and information gathered by WSDA. Samples were obtained beginning the week of April 22 and sampling continued through the week of May 20. Samples were also obtained July 8 through July 25. This sampling regime is similar to what has been used in previous studies. The laboratory analysis was completed by ALS Environmental (ALS) under contract.

Quality Objectives and Data Criteria

Quality objectives for this project are to obtain data of sufficient quality and quantity so it can be used to assess the current concentrations of chlorpyrifos and diazinon in the Grayland Ditch system; and subjectively evaluate the effectiveness and extent of implemented BMPs. This will be accomplished by comparing data from this study to data collected from previous studies. These objectives will be achieved through careful planning and adherence to procedures described in the project Quality Assurance Project Plan (QAPP).

Field Procedures

Field measurements of temperature, pH, conductivity, DO, and turbidity were adapted from Ecology Standard Operating Procedures (SOP) (Nipp, 2006; Swanson, 2007; Ward, 2007a, 2007b) and the manufactures instruction manual. All field parameters were measured at the sampling site by field staff using a Hydrolab Quanta® multi-parameter probe (table 2).

Table 2. Measurement quality objectives for conventional parameters measured by field meters or determined by a standard method.

Parameter	Method/Equipment	Field Replicate MQO	Reporting Limits
Water Temperature	Hydrolab Quanta™	±0.2°C	-5.0°C
Conductivity	Hydrolab Quanta™	±1% of reading ±1 count	0.1 mS/cm
pH	Hydrolab Quanta™	±0.2 s.u.	2 s.u.
Dissolved Oxygen	Hydrolab Quanta™	±0.2 mg/L	0.1 mg/L
Turbidity	Hydrolab Quanta™	±0.5% of reading ± NTU	0 NTU

MQO – measurement quality objectives, s.u. – standard unit, Nephelometric Turbidity Units (NTU)

The Hydrolab Quanta™ was calibrated before each field use by using commercially available reagents and in accordance with the manufacturer’s instructions.

Stream flow measurements were obtained using a Marsh-McBirney flow meter, an OTT MF Pro electromagnetic flow sensor and a top-setting rod. Flow SOP was adapted from Ecology’s *Measuring and Calculating Stream Discharge v1 IEAP056*.

Lab Procedures

All surface water samples were collected as grab samples using a pole sampler. Surface water sampling SOPs were adapted from Ecology SOPs described in EAP003 *Sampling of Pesticides in Surface Waters* (Anderson, 2006) and EAP015 *Manually Obtaining Surface Water Samples* (Joy, 2006) (table 3). The water samples were sent to ALS in Kelso WA for analysis.

Table 3. Laboratory measurement quality objectives.

Parameter	Laboratory Control Samples	Duplicate Lab Control Samples	Matrix Spikes Samples	Duplicate Samples	Surrogate Standards
	% recovery	RPD	% Recovery	RPD	% Recovery
Diazinon	82	30	N/A	N/A	N/A
Chlorpyrifos	71	30	N/A	N/A	N/A

N/A – not applicable RPD – relative percent difference

Results from ALS included case narratives describing QA/QC procedures used during analysis. These QA/QC results included holding times, instrument calibrations, method blanks, matrix spikes, laboratory duplicates, laboratory control samples, and surrogate spikes. Case narratives describing the quality of the data are available upon request. ND and U were data qualifiers used to indicate non detection of the target analytes.

Transfer blanks were analyzed to evaluate the potential for contamination. Transfer blanks were prepared using blank water supplied by ALS. Laboratory water was transferred from its container at the sampling site to a new sample container. No target analytes were detected in transfer blanks.

The sample results were compared to available water criteria in Table 4.

Table 4. Available water quality criteria.

Chemical	Type	Common Name	WAC		NRWQC	
			Acute	Chronic	CMC	CCC
Chlorpyrifos	Organophosphate	Lorsban	0.083 µg/L	0.041 µg/L	0.083 µg/L	0.041 µg/L
Diazinon	Organophosphate	(several)	--	--	0.17 µg/L	0.17 µg/L

WAC – Washington Administrative Code (Chapter 172-201A)


CMC – criteria maximum concentration

CCC – criteria continuous concentration

Sample Results


In 2012 only water samples were obtained. In 2013 both water samples and ambient data were collected. The water samples were analyzed for the presence of diazinon and chlorpyrifos. The results are shown in Figures 2 through 5

Figure 2. 2012 Diazinon Concentrations




In 2012, diazinon was detected at each site at least twice at each site. All detections were above the NRWQC maximum/continuous concentration of 0.17 µg/L.

Figure 3. 2013 Diazinon Concentrations.




In 2013, diazinon was detected seven times at four sites. All detections were above the NRWQC maximum/continuous concentration of 0.17 µg/L. Only one detection (Heather on 7/18) was substantially above the NRWQC at levels similar to those observed in 2012.

Figure 4. 2012 Chlorpyrifos Concentrations.



Chlorpyrifos was detected at three sites in 2012. All detections were well above the established acute and chronic water quality standards.

Figure 5. 2013 Chlorpyrifos Concentrations.




Chlorpyrifos was detected at one site once in 2013. This detection was well above the State Acute and Chronic concentrations.

Ambient Data

Washington State water quality standards for conventional water quality parameters are set forth in Chapter 173-201A of the WAC. Water bodies are required to meet water quality standards based on the designated uses of the water body. When designated uses are not specified for a particular water body, then default designated uses are used to identify appropriate water quality criteria. The Grayland Ditch does not have specified designated uses, so the default becomes (1) salmonid spawning, rearing, and migration, and (2) primary contact recreation. There is no documented use by, or presence of, any salmon species in Grayland Ditch (WDFW, 2010). The numeric pH criteria for the above designated uses should be within the range of 6.5 to 8.5, with a human-caused variation of less than 0.5 standard units.


The ambient data collected is shown in Figures 6 through 12. Ambient data was not available for all sites for all sampling events. The tabular data for all measurements is available in Appendix C.

Figure 6. Discharge (Flow).



Stream discharge (flow) was measured at three sites in 2013. Discharge peaked on 5/8 and declined as the year progressed. The low flow in this system indicates a chance for increased pesticide exposure to aquatic invertebrates and non-anadromous fish.


Figure 7. Water Temperature.



Ambient data was not available for all sites for all sampling events.

Water temperature varied from just above eight degrees (C) in April to approximately fourteen degrees in July. All of these temperatures are within the range to support aquatic life. This ditch system has an abundance of riparian plantings and shade to mediate water temperature.


Chart 7. pH.



Ambient data was not available for all sites for all sampling events.

The pH as measured was most often within the aquatic criteria. The County Line reading on 5/1 was the single exception. The pH generally tended towards slightly acidic. The water in the Grayland Ditch primarily comes from wetlands and bog drainage which are typically acidic.


Figure 9. Turbidity.



Ambient data was not available for all sites for all sampling events.

Turbidity was measured using the Hach Hydrolab Quanta[®] beginning with the July site visits. The Schmid and County Line sites showed considerable variability between visits. Turbidity will be measured in the field during all sampling events in 2014 and 2015.


Figure 10. Conductivity.



Ambient data was not available for all sites for all sampling events.

Conductivity measurements showed considerable variability between the sampling sites. The measurements indicate the presence of salt water, at least in small amounts, throughout the Grayland Ditch system. This is expected and observed in other tidegate protected ditch systems.


Figure 11. Dissolved Oxygen.



Ambient data was not available for all sites for all sampling events.

The dissolved oxygen levels were generally low and did not meet any of the criteria for salmonid habitat.

Figure 12. Total Suspended Solids.



Ambient data was not available for all sites for all sampling events.

Water samples were obtained from most site visits and tested for total suspended solids. While there was a certain amount of variability between the samples, the Schmid sample on 7/18 was an extremely high. Turbidity was also measured in the field on 7/18/2013. This measurement was also very high. This anomalous result may be a natural occurrence or may be due to errors in the sampling technique.

Description and Findings

Ambient water quality information indicates the Grayland ditch system has relatively consistent water temperatures, dissolved oxygen and pH levels during the spring and summer. All areas of the system show some salt water intrusion, which is expected in a tidegate managed ditch.

Although this system is high in silt and other organic material, all but one turbidity sample were below 20 mg/L, indicating relatively good water clarity. Evaluation of ambient criteria will continue during the 2014 and 2015 sampling seasons.

The number of detections of both diazinon and chlorpyrifos declined during the first two years of WSDA's sampling study. The concentrations detected have also generally declined, although all detections are still above State and Federal water quality levels. The number of detections and concentrations have also declined from previous studies conducted by Ecology.

There are three factors that may be contributing to the decline in chlorpyrifos and diazinon detections:

1. The BMPs (cribbing and covering the ditches) is slowly increasing on all or most of the bog ditches and may be functioning properly. This would indicate the BMPs are successful in decreasing pesticide concentrations in the ditch system.
2. Diazinon and chlorpyrifos may not be used as much as they were in the past. This could indicate a decline in the number of applications and also the applied rate of these pesticides.
3. A combination of one and two.

A visual inspection of the bogs near the Grayland Ditch would help determine if the BMPs are in place and functioning properly. Discussions with the growers, WSU Extension and Ocean Spray would also help ascertain the status of the BMPs.

The purpose of this study is to evaluate the effectiveness of the "cribbing and covering" BMP in the Grayland Ditch. Implementation of the required BMPs should keep the majority of applied pesticide residues from entering the waters of the state. If diazinon and chlorpyrifos are being phased out, then continued testing for only diazinon and chlorpyrifos will not be a good indicator

to determine the effectiveness of the BMPs. WSDA should make an effort to identify all pesticides used in the production of cranberries and expand the water sampling testing in the 2014 and 2015 application seasons to include all pesticides applied.

This page left blank intentionally

References

- Anderson, P., 2006. Standard Operating Procedure (SOP) for Sampling of Pesticides in Surface Waters, Version 1.0. Washington State Department of Ecology, Olympia, WA. SOP Number EAP003. www.ecy.wa.gov/programs/eap/quality.html.
- Anderson, P., 2009. Quality Assurance Project Plan: Grayland Ditch: An Evaluation of Organophosphate Pesticides and Pesticide Test Kits. Washington State Department of Ecology, Olympia, WA. Publication No. 09-03-115. www.ecy.wa.gov/biblio/0903115.html.
- Anderson, P. and D. Davis, 2000. Evaluation of Efforts to Reduce Pesticide Contamination in Cranberry Bog Drainage. Washington State Department of Ecology, Olympia, WA. Publication No. 00-03-041. www.ecy.wa.gov/biblio/0003041.html.
- APHA, 2005. Standard Methods for the Analysis of Water and Wastewater, 21st Edition. Joint publication of the American Public Health Association, American Water Works Association, and Water Environment Federation. www.standardmethods.org/.
- Cook, K.V. and J. Cowles, 2009. Washington State Pesticide Management Strategy: Water Quality Protection, Version 2.22. Washington State Department of Agriculture, Olympia, WA. <http://agr.wa.gov/pestfert/natresources/docs/comprehensivepesticidemanagementstrategy.pdf>
- Coots, R., 2003. Pesticide Reduction Evaluation for Cranberry Bog Drainage in the Grayland Area. Washington State Department of Ecology, Olympia, WA. Publication No. 03-03-034. www.ecy.wa.gov/biblio/0303034.html.
- Davis, D., D. Serdar, and A. Johnson, 1997. Assessment of Cranberry Bog Drainage Pesticide Contamination. Washington State Department of Ecology, Olympia, WA. Publication No. 97-329. www.ecy.wa.gov/biblio/97329.html.
- Embrey, S.S. and L.M. Frans, 2003. Surface-Water Quality of the Skokomish, Nooksack, and Green-Duwamish Rivers and Thornton Creek, Puget Sound Basin, Washington, 1995-98. U.S. Geological Survey, Water-Resources Investigation Report 02-4190. <http://pubs.usgs.gov/wri/wri024190/pdf/wri024190.pdf>.
- EPA, 1991. Guidance for Data Usability in Risk Assessment. (Part A). EPA/540/R-92/003. U.S. Environmental Protection Agency, Washington, D.C.
- EPA, 1994. Using Qualified Data to Document an Observed Release. EPA/540/F-91/028. U.S. Environmental Protection Agency, Washington, D.C.
- EPA, 1998. Method 8270. Semivolatile Organic Compounds by GC/MS, Revision 4.0. U.S. Environmental Protection Agency, Office of Solid Waste. www.epa.gov/epawaste/hazard/testmethods/sw846/pdfs/8270d.pdf.

EPA, 2004. Method 3535. Solid-Phase Extraction for Organic Analytes, Revision 6.0. U.S. Environmental Protection Agency, Office of Solid Waste.
www.epa.gov/epawaste/hazard/testmethods/sw846/pdfs/3535a.pdf.

EPA, 2006. National Recommended Water Quality Criteria. U.S. Environmental Protection Agency, Office of Water and Office of Science and Technology.
www.epa.gov/waterscience/criteria/wqctable/.

Federal Register, 2006a. Environmental Protection Agency: Azinphos-methyl; Order to Amend Registrations to Terminate Certain Uses. March 29, 2006. Volume 71, Number 60: 15731-15732. Document ID: fr29mr06-77.

Federal Register, 2006b. Environmental Protection Agency: Azinphos-methyl; Amending Existing Stocks Provisions of Distribution for Cranberry Growers. July 5, 2006. Volume 71, Number 128: 38148-38149. Document ID: fr05jy06-46.

Frantz, W. T., T. Bicki, K. Talbot, and E. Unruh, 1997. Determining the Effectiveness of Surface Water Best Management Practices (BMPs) for Grayland, WA Cranberry Bogs. Cranberry Institute, Wareham, Massachusetts.

Grayland Cranberry Growers Board, 2010. Personal communication at the January 2010 board meeting regarding the use of other pesticides in place of azinphos-methyl. Markham, WA. Last communication January 2010.

Joy, J., 2006. Standard Operating Procedure (SOP) for Manually Obtaining Surface Water Samples, Version 1.0. Washington State Department of Ecology, Olympia, WA. SOP Number EAP015.
www.ecy.wa.gov/programs/eap/quality.html.

MEL, 2008. Manchester Environmental Laboratory Lab Users Manual, Ninth Edition. Manchester Environmental Laboratory, Washington State Department of Ecology, Manchester, WA.

Menconi, M. and C. Cox, 1994. Hazard Assessment of the Insecticide Diazinon to Aquatic Organisms in the Sacramento-Jan Joaquin River System. State of California Department of Fish and Game, Environmental Services Division, Pesticide Investigations Unit, Rancho Cordova, CA. Administrative Report 94-2. www.cdpr.ca.gov/docs/emon/surfwttr/hazasm/hazasm94_2.pdf.

Mitsch, W.J. and J.G. Gosselink, 2007. Wetlands, 4th edition. John Wiley & Sons, Inc., Hoboken, NJ.

Nipp, B., 2006. Standard Operating Procedure (SOP) for Instantaneous Measurements of Temperature in Water, Version 1.0. Washington State Department of Ecology, Olympia, WA. SOP Number EAP011. www.ecy.wa.gov/programs/eap/quality.html.

Pacific Conservation District and Pacific Coast Cranberry Research Foundation, 1999. BMPs for Cranberry Agriculture. Washington State Conservation Commission.
Rounry, D., 2008. Grayland Cranberry Growers Get Pesticides Out of Water and Reduce Costs.

Washington State Department of Ecology, Olympia, WA. Publication No. 08-10-087.
www.ecy.wa.gov/biblio/0810087.html.

Swanson, T., 2007. Standard Operating Procedure (SOP) for Hydrolab® DataSonde® and MiniSonde® Multiprobes, Version 1.0. Washington State Department of Ecology, Olympia, WA. SOP Number EAP033. www.ecy.wa.gov/programs/eap/quality.html.

Talbot, Kevin, 2010. Personal communication regarding types and amounts of pesticides used in Grayland cranberry bogs. Ocean Spray, Markham, WA. Last communication January 2010. WAC 173-201A. Water Quality Standards for Surface Waters in the State of Washington

Washington State Department of Ecology, Olympia, WA. www.ecy.wa.gov/laws-rules/ecywac.html.
Ward, W., 2007a. Standard Operating Procedure for the Collection and Analysis of pH Samples, Version 1.3. Washington State Department of Ecology, Olympia, WA. SOP Number EAP031. www.ecy.wa.gov/programs/eap/quality.html.

Ward, W., 2007b. Standard Operating Procedure for the Collection and Analysis of Conductivity Samples, Version 1.3. Washington State Department of Ecology, Olympia, WA. SOP Number EAP032. www.ecy.wa.gov/programs/eap/quality.html.

WDFW, 2010. SalmonScape interactive mapping application. Washington Department of Fish and Wildlife, Olympia, WA. <http://wdfw.wa.gov/mapping/salmonscape/index.html>.

Appendix A

Glossary, Acronyms, and Abbreviations

Glossary

303(d) list: Section 303(d) of the federal Clean Water Act requires Washington State to periodically prepare a list of all surface waters in the state for which beneficial uses of the water – such as for drinking, recreation, aquatic habitat, and industrial use – are impaired by pollutants. These are water quality limited estuaries, lakes, and streams that fall short of state surface water quality standards.

Acetylcholinesterase: An enzyme that hydrolyzes the neurotransmitter acetylcholine: its action is blocked by nerve gases and certain drugs.

Carbamate insecticides: N-methyl carbamate insecticides are similar to organophosphate insecticides in that they are nerve agents that inhibit cholinesterase enzymes. However they differ in action from the organophosphate compounds in that the inhibitory effect on cholinesterase is brief. Carbamates degrade rapidly in the environment.

Clean Water Act: A federal act passed in 1972 that contains provisions to restore and maintain the quality of the nation's waters. Section 303(d) of the Clean Water Act establishes the TMDL program.

Conductivity: A measure of water's ability to conduct an electrical current. Conductivity is related to the concentration and charge of dissolved ions in water.

Exceeds criteria: Fails to meet criteria.

Grab sample: A discrete sample from a single point in the water column or sediment surface.

Organophosphate pesticides: Organophosphate pesticides are all derived from phosphoric acid. They are nerve poisons which kill the target pest (usually insects). The mechanism of action is similar to carbamate insecticides, both are neurotoxins, inhibiting the enzyme acetylcholinesterase by inhibiting cholinesterase. They break down relatively quickly in the environment.

Parameter: Water quality constituent being measured (analyte). A physical, chemical, or biological property whose values determine environmental characteristics or behavior.

Pesticide: A pesticide is any substance or mixture of substance intended for preventing, destroying, repelling or mitigating any pest. Pests include nuisance microbes, plants, fungus, and animals.

pH: A measure of the acidity or alkalinity of water. A low pH value (0 to 7) indicates that an acidic condition is present, while a high pH (7 to 14) indicates a basic or alkaline condition. A pH of 7 is

considered to be neutral. Since the pH scale is logarithmic, a water sample with a pH of 8 is ten times more basic than one with a pH of 7.

Surface waters of the state: Lakes, rivers, ponds, streams, inland waters, salt waters, wetlands and all other surface waters and watercourses within the jurisdiction of Washington State.

Total Maximum Daily Load (TMDL): Water cleanup plan. A calculated or managed distribution of a substance in a waterbody designed to protect it from exceeding water quality standards. A TMDL is equal to the sum of all of the following: (1) individual wasteload allocations for point sources, (2) the load allocations for nonpoint sources, (3) the contribution of natural sources, and (4) a Margin of Safety to allow for uncertainty in the wasteload determination. A reserve for future growth is also generally provided.

Total suspended solids: The suspended particulate matter in a water sample as retained by a filter.

Acronyms and Abbreviations

Following are acronyms and abbreviations used frequently in this report.

BMP Best management practices
BOD Biological Oxygen Demand
CCC Criteria Continuous Concentrations
CFU Cubic Feet per Second
CMC Criteria Maximum Concentration
DO Dissolved Oxygen
Ecology Washington State Department of Ecology
EPA U.S. Environmental Protection Agency
FIFRA Federal Insecticide Fungicide Rodenticide Act
GHCCDD-1 Grays Harbor County Drainage Ditch No. 1
MEL Manchester Environmental Laboratory
MQO Measurement Quality Objectives
NAD North American Datum
NOAA National Oceanic and Atmospheric Administration
NRAS Natural Resource Assessment Section
NRWQC National Recommended Water Quality Criteria
NTU Nephelometric Turbidity Units
OP Organophosphate
PCDD-1 Pacific County Drainage Ditch No. 1
QAPP Quality Assurance Project Plan
QA/QC Quality Assurance / Quality Control
RPD Relative percent difference
SOP Standard operating procedures
SU Standard Units
TSS Total Suspended Solids
WAC Washington Administrative Code
WSDA Washington State Department of Agriculture

Units of Measurement

°C degrees centigrade

mg/L milligrams per liter

s.u. standard unit

µg/L micrograms per liter (parts per billion)

µS/cm microSeimans per centimeter

Appendix B.

Summarized Study Results from Common Sampling Sites, 1996-2009 (Ecology).

Table 5. Summarized study results from 1996, 1998, and 2009 sampling sites.

Location			1996	1998	2002	2009
GHCDD-1						
Schmid Road	Azinphos-methyl	number	26	5	3	3
		detections	16	3	2	0
		mean	0.13	0.25	0.11	nd
		range	0.010-0.73	0.004-1.2	0.033-0.20	nd
	Chlorpyrifos	n	26	5	3	3
		detections	7	5	2	3
		mean	0.021	0.38	0.008	0.09
		range	0.003-0.016	0.0095-1.8	0.0050-0.010	0.051-0.13
	Diazinon	n	26	5	3	3
detections		26	5	3	2	
mean		0.86	1.1	0.17	0.76	
range		0.026-5.42	0.033-4.4	0.018-0.35	0.034-2.2	
Grange Road	Azinphos-methyl	n	-	-	3	3
		detections	-	-	3	0
		mean	-	-	0.13	nd
		range	-	-	0.019-0.020	nd
	Chlorpyrifos	n	-	-	3	3
		detections	-	-	3	3
		mean	-	-	0.013	0.14
		range	-	-	0.0065-0.023	0.089-0.21
	Diazinon	n	-	-	3	3
detections		-	-	3	2	
mean		-	-	1.9	0.054	
range		-	-	0.033-5.7	0.034-0.094	
County Line Road	Azinphos-methyl	n	-	-	3	3
		detections	-	-	3	0
		mean	-	-	0.18	nd
		range	-	-	0.020-0.30	nd
	Chlorpyrifos	n	-	-	3	3
		detections	-	-	2	3
		mean	-	-	0.012	0.16
		range	-	-	0.0067-0.020	0.1-0.22
	Diazinon	n	-	-	3	3
detections		-	-	3	1	
mean		-	-	0.80	0.029	
range		-	-	0.036-2.0	0.032-0.055	

Location			1996	1998	2002	2009
PCDD-1						
Heather Road	Azinphos-methyl	number	-	-	3	3
		detections	-	-	3	0
		mean	-	-	0.11	nd
		range	-	-	0.048-0.22	nd
	Chlorpyrifos	number	-	-	3	3
		detections	-	-	3	1
		mean	-	-	0.027	0.027
		range	-	-	0.019-0.039	0.022-0.052
	Diazinon	number	-	-	3	3
detections		-	-	3	3	
mean		-	-	0.16	0.19	
range		-	-	0.079-0.31	0.089-0.36	
Jacobson Road	Azinphos-methyl	number	-	-	3	3
		detections	-	-	3	0
		mean	-	-	0.08	nd
		range	-	-	0.0079-0.22	nd
	Chlorpyrifos	number	-	-	3	3
		detections	-	-	3	2
		mean	-	-	0.39	0.046
		range	-	-	0.034-0.59	0.035-0.082
	Diazinon	number	-	-	3	3
detections		-	-	3	3	
mean		-	-	0.31	0.21	
range		-	-	0.11-0.69	0.096-0.42	
Larkin Road	Azinphos-methyl	number	26	5	3	-
		detections	23	5	3	-
		mean	0.17	0.33	0.050	-
		range	0.006-0.74	0.012-1.4	0.0061-0.13	-
	Chlorpyrifos	number	26	5	3	-
		detections	25	5	3	-
		mean	0.44	0.59	0.028	-
		range	0.003-3.7	0.019-1.3	0.015-0.036	-
	Diazinon	number	26	5	3	-
detections		25	5	3	-	
mean		0.28	2.4	0.48	-	
range		0.008-1.7	0.033-7.0	0.20-0.64	-	
Tidegate	Azinphos-methyl	number	-	-	3	3
		detections	-	-	3	0
		mean	-	-	0.06	nd
		range	-	-	0.0067-0.15	nd
	Chlorpyrifos	number	-	-	3	3
		detections	-	-	3	2
		mean	-	-	0.025	0.08
		range	-	-	0.014-0.030	0.027-0.18
	Diazinon	number	-	-	3	3
detections		-	-	3	3	
mean		-	-	0.53	0.13	
range		-	-	0.18-0.71	0.085-0.22	

Appendix C

Environmental Measurements

Table 6. Ambient Measurements (2013).

Location	Date	Discharge	Water Temp(°C)	pH	Turb (NTU)	Cond (µS/cm)	Cond (MS/cm)	DO(%)	DO(mg/L)	TSS (mg/L)
Tidegate North	4/25/13	NA	NA	NA	NA	NA	NA	NA	NA	8.5
Tidegate North	5/1/13	NA	8.30	7.52	NA	175	0.175	62.7	7.45	8
Tidegate North	5/8/13	NA	11.61	7.24	NA	174	0.174	66.6	7.26	11
Tidegate North	5/15/13	NA	11.43	6.95	NA	171	0.171	67.1	7.31	18.5
Tidegate North	5/22/13	NA	NA	NA	NA	NA	NA	NA	NA	NA
Tidegate North	7/9/13	NA	13.76	7.08	19.2	179	0.179	68.7	7.11	6.5
Tidegate North	7/18/13	NA	13.30	7.08	18.6	181	0.181	63.5	6.63	6.5
Tidegate North	7/25/13	NA	12.82	7.04	17.5	176	0.176	64.6	6.88	6
Schmid	4/25/13	NA	NA	NA	NA	NA	NA	NA	NA	11
Schmid	5/1/13	6.36678	11.73	6.92	NA	233	0.233	40.1	4.41	11.5
Schmid	5/8/13	7.6644	12.85	6.92	NA	197	0.197	55.7	5.84	10.5
Schmid	5/15/13	4.8186	12.32	6.82	NA	195	0.195	54.4	5.85	19
Schmid	5/22/13	4.6652	11.43	6.96	NA	193	0.193	55.7	6.11	12
Schmid	7/9/13	5.1496	13.43	6.96	26.2	233	0.233	47.2	4.95	10
Schmid	7/18/13	2.53052	13.92	6.88	100	229	0.229	37	3.96	83.3
Schmid	7/25/13	2.02868	14.11	6.82	23.1	213	0.213	56	5.77	12

Table 6. Ambient Measurements (2013).

Location	Date	Discharge	Water Temp(°C)	pH	Turb (NTU)	Cond (µS/cm)	Cond (MS/cm)	DO(%)	DO(mg/L)	TSS (mg/L)
County Line	4/25/13	NA	NA	NA	NA	NA	NA	NA	NA	9
County Line	5/1/13	2.7264	11.55	6.00	NA	216	0.216	35.5	3.9	11.5
County Line	5/8/13	2.606	12.70	6.92	NA	187	0.187	52.8	5.62	8
County Line	5/15/13	1.8956	12.06	6.76	NA	185	0.185	48.2	5.18	10
County Line	5/22/13	2.5832	11.26	6.83	NA	194	0.194	47.7	5.25	13
County Line	7/9/13	0.889339	14.00	6.8	24.6	208	0.208	34.1	3.52	11
County Line	7/18/13	0.51191	14.68	6.7	68.5	217	0.217	28.8	2.74	24.5
County Line	7/25/13	0.713712	14.28	6.75	40.1	200	0.2	48.5	4.99	12
Heather	4/25/13	NA	NA	NA	NA	NA	NA	NA	NA	ND
Heather	5/1/13	4.33136	9.88	6.9	NA	157	0.157	50.2	5.76	15.5
Heather	5/8/13	5.98	11.92	6.96	NA	134	0.134	62.4	6.77	5
Heather	5/15/13	3.97925	11.70	6.925	NA	145	0.145	58.2	6.38	5.5
Heather	5/22/13	4.17075	11.07	NA	NA	150	0.15	56.7	6.37	ND
Heather	7/9/13	2.52888	14.13	6.88	NA	134	0.134	57.7	5.91	5.5
Heather	7/18/13	2.337	13.40	6.97	11.2	135	0.135	62.6	6.22	ND
Heather	7/25/13	1.75972	12.24	6.92	5.9	158	0.158	52.4	5.54	ND

Table 6. Ambient Measurements (2013).

Location	Date	Discharge	Water Temp(°C)	pH	Turb (NTU)	Cond (µS/cm)	Cond (MS/cm)	DO(%)	DO(mg/L)	TSS (mg/L)
Tidegate South	4/25/13	NA	NA	NA	NA	NA	NA	NA	NA	10
Tidegate South	5/1/13	NA	9.61	7.3	NA	264	0.264	45.4	5.22	7
Tidegate South	5/8/13	NA	12.11	7.13	NA	226	0.226	54.8	5.94	5.5
Tidegate South	5/15/13	NA	11.94	6.96	NA	226	0.226	56.4	6.07	7.5
Tidegate South	5/22/13	NA	11.11	7.04	NA	209	0.209	59.1	6.51	8
Tidegate South	7/9/13	NA	14.09	7.11	12.9	232	0.232	55.5	5.72	5.5
Tidegate South	7/18/13	NA	13.98	7.16	12.1	227	0.227	57.8	5.97	ND
Tidegate South	7/25/13	NA	13.40	7.08	10.3	232	0.232	56.8	5.97	ND

This page is left blank intentionally