

ISAAC HALL #2238
 2087 Wells Street
 Wailuku, Maui, Hawaii 96793
 Telephone: (808) 244-9017
 Attorney for Maui Tomorrow Foundation, Inc.,
 and its supporters

COMMISSION ON WATER RESOURCE MANAGEMENT

STATE OF HAWAII

In re Petitions to Amend Interim
 Instream Flow Standards for
 Honopou, Huelo (Puolua), Hanehoi,
 Waikamoi, Alo, Wahinepe'e,
 Puohokamoa, Haipua'ena,
 Punalau/Kōlea, Honomanu, Nu'ailua,
 Pi`ina`au, Palauhulu, Ohia (Waianu),
 Waiokamilo, Kualani, Wailuanui, West
 Wailuaiki, East Wailuaiki, Kopili`ula,
 Puaka`a, Waiohue, Pa`akea, Waiaka`a,
 Kapa`ula, Hanawī and Makapipi
 streams.

Case No. CCH-MA13-01

EXHIBIT LIST OF MAUI TOMORROW
 FOUNDATION, INC. AND ITS
 SUPPORTERS FOR RE-OPENED HEARING;
 CERTIFICATE OF SERVICE


mt/exlist

**EXHIBIT LIST OF MAUI TOMORROW
 FOUNDATION, INC. AND ITS SUPPORTERS
 FOR RE-OPENED HEARING**

EXHIBIT NUMBER	DESCRIPTION	REFERENCES	ADM
E-1 - E-158	Exhibits, incorporated by reference, as admitted during original contested case hearing		
E-159	Curriculum Vitae for Albert Perez		
E-160	Mālama `Āina: A Conversation About Maui's Farming Future		
E-161	Maui News; August 28, 2016		
E-162	Second Amended Chart, Hanehoi Watershed		
E-163	Hanehoi Stream Diversions; Photos		
E-164	EMI Map of Honopou and Hanehoi Diversions		
E-165	Letter dated September 16, 2016 from Schulmeister to Case		

DATED: Wailuku, Maui, Hawaii

10.17.16

A handwritten signature in black ink, appearing to be 'I-h', written over a horizontal line.

Isaac Hall
Attorney for Maui Tomorrow Foundation,
Inc., and its Supporters

CERTIFICATE OF SERVICE

I hereby certify that one copy of the foregoing document was duly served upon the parties listed below by email, on October 17, 2016.

Commission on Water Resource Management
(via U.S. Mail and email
c/o kathy.s.yoda@hawaii.gov)
c/o Kathy S. Yoda
P.O. Box 621
Honolulu, HI 96809

Camille K. Kalama, Esq.
(via email: camille.kalama@nhlchi.org)
Summer L.H. Sylva, Esq.
(via email: summer.sylva@nhlchi.org)
Native Hawaiian Legal Corporation
1164 Bishop Street, Suite 1205
Honolulu, HI 96813
Attorneys for Na Moku Aupuni O Koolau Hui

Robert H. Thomas, Esq.
(via email: rht@hawaiilawyer.com)
Damon Key Leong Kupchak Hastert
1003 Bishop Street
Pauahi Tower, Suite 1600
Honolulu, HI 96813
Attorneys for Hawaii Farm Bureau Federation

William J. Wynhoff, Esq.
(via email: bill.j.wynhoff@hawaii.gov)
Linda L.W. Chow, Esq.
(via email: linda.l.chow@hawaii.gov)
Department of the Attorney General
465 South King Street, Room 300
Honolulu, HI 96813

Lawrence Miike, Hearings Officer
(via email: lmiike@hawaii.rr.com)
c/o Commission on Water Resource Management
P.O. Box 621
Honolulu, HI 96809

Elijah Yip, Esq.
(via email: eyip@cades.com)
David Schulmeister, Esq.
(via email: dschulmeister@cades.com)
Cades Schutte, LLP
1000 Bishop Street, 10th Floor
Honolulu, HI 96813
Attorneys for Alexander & Baldwin, Inc. and East Maui
Irrigation Co., Ltd.

Patrick K. Wong, Esq.
(via email: pat.wong@co.maui.hi.us)
Caleb Rowe, Esq.
(via email: Caleb.Rowe@co.maui.hi.us)
Kristin Tarnstrom, Esq.
(via email: Kristin.Tarnstrom@co.maui.hi.us)
Department of the Corporation Counsel
County of Maui,
200 S. High Street
Wailuku, HI 96793
Attorneys for County Dept of Water Supply

Jeffrey C. Paisner
(via email: jeffreypaisner@mac.com)
403 West 49th Street #2
New York, New York 10019
Pro Se

DATED: Wailuku, Maui, Hawaii

10.17.16


Isaac Hall
Attorney for Maui Tomorrow Foundation, Inc., and its Supporters

David Schulmeister
Direct Line: (808) 521-9392
Direct Fax: (808) 540-5022
E-mail: dschulmeister@ca des.com

September 16, 2016

VIA E-MAIL AND MAIL

Ms. Suzanne D. Case
Chairperson
State of Hawaii
Department of Land and Natural Resources
Commission on Water Resource Management
P.O. Box 621
Honolulu, Hawaii 96809

Re: *In the Matter of Petitions to Amend Interim Instream Flow Standards for East Maui Streams, Case No. CCH-MA13-01*

Dear Ms. Case:

The purpose of this letter is to provide, on behalf of East Maui Irrigation Co., Ltd. ("EMI"), a further update to the Commission regarding the status of EMI's compliance with the Commission's July 18, 2016 Order Re Interim Restoration of Stream Flow with regard to the streams that are to be permanently restored.

Please find enclosed an update of the plan previously submitted that includes the following additional information:

1. A listing of all of EMI's registered major and minor stream diversions on the taro streams cross referenced to the numbers used on the EMI map of the Ditch System so as to show their precise location;
2. An updated timeline.

The stream diversion work completed and to be completed is broken out on the updated plan both by stream and by phase of work for ease of reference.

I have been further advised that EMI has completed assembling the information required to be submitted with EMI's applications for stream diversion abandonment permits, and those

C S

Cades Schutte Building
1000 Bishop Street, Suite 1200
Honolulu, Hawaii 96813
Tel: 808.521-9200
Fax: 808.521-9210
www.cades.com

Kona Office
75-170 Hualalai Road, Suite B-303
Kailua Kona, Hawaii 96740
Tel: 808.329-5817
Fax: 808.326-1125

Maui Office
444 Hana Highway, Suite 204
Kahului, Hawaii 96732
Tel: 808.871.6016
Fax: 808.871.6017


Exhibit "E-165"

Ms. Suzanne D. Case
September 16, 2016
Page 2

applications are expected to be filed with the Commission today. Work on the other regulatory submissions and required approvals is continuing.

EMI proposes to provide another update in thirty days, or by October 17, 2016.

Very truly yours,



David Schulmeister

for

CADES SCHUTTE

A Limited Liability Law Partnership

Enc.

Cc (w/enc.):

Dr. Lawrence H. Miike
Linda L.W. Chow, Esq.
Summer L.H. Sylva, Esq.
Camille K. Kalama, Esq.
Issac Hall, Esq.
Patrick K. Wong, Esq.
Caleb P. Rowe, Esq.
Kristin K. Tarnstrom, Esq.
Robert H. Thomas, Esq.
Jeffrey C. Paisner
John Blumer-Buell
Nikhilananda

EMI Taro Stream Diversions

DIVERSIONS BY STREAM

	EMI Map #	General Description of Work
Honopou		
Honopou	W-22	Bolt steel plates or concrete over diversion intake grate.
Honopou long pipe intake	W-22a	Bolt steel plates or concrete over diversion intake grate.
Wailole	W-22b	Bolt steel plates or concrete over diversion intake grate.
Honopou	NH-22	Bolt steel plates or concrete over diversion intake grate.
Wailole	NH-23	Bolt steel plates or concrete over diversion intake grate.
Honopou long strainer	L-15	Seal intake opening with rocks and concrete.
Honopou siphon	L-16	Construct stream overpass over ditch.
Honopou side ditch	L-17	Construct stream overpass over ditch.
Honopou	H-8	Close existing control gate.
		Bolt steel plates or concrete over diversion intake grate and seal opening below grate with rock and concrete.
Hanehoi (Puolua)		
Hanehoi (Huelo intake)	W-18	Bolt steel plates or concrete over diversion intake grate.
Hanehoi (Huelo intake)	NH-17	Concrete over diversion intake grate.
West Hanehoi intake (Puolua)	NH-17a	Construct stream overpass over ditch.
Hanehoi Huelo # 1	L-5	Bolt steel plates or concrete over diversion intake grate.
Hanehoi Huelo # 2	L-6	Bolt steel plates or concrete over diversion intake grate.
Hanehoi Huelo # 3	L-7	Bolt steel plates or concrete over diversion intake grate.
Hanehoi	L-5a	Construct stream overpass over ditch.
Hanehoi	L-5b	Construct stream overpass over ditch.
Hanehoi	L-5c	Construct stream overpass over ditch.
Hanehoi Roseapple (Puolua)	L-7a	Install pipe or box culvert with wing walls through which ditch can pass beneath stream or construct stream overpass over ditch.
West Hanehoi	L-7b	Construct stream overpass over ditch.
East Hanehoi (Pancho)	H-3	Bolt steel plates or concrete over diversion intake grate.
West Hanehoi (School)	H-4	Seal intake opening with rock and concrete.
Pfina'au		
Pfina'au	K-31	Seal intake opening with rocks and concrete.
Hanehoi Wailole small intake controlled by gate	K-30d	Construct overpass over ditch.
Pfina'au 6" steel and pvc pipe intake	K-31a	Remove steel and pvc pipes.
Palauhulu		
Kano	K-26	Most flow will be restored by removal of sluice gate. Scope of work for full restoration is to be determined. All work is anticipated to be restricted to tunnel.

Lalahai (# 3 intake - Hauolo Wahine Ditch)	K-27	Scope of work for full restoration is to be determined. All work is anticipated to be restricted to tunnel.
Lalapipi (# 2 intake - Hauolo Wahine Ditch)	K-28	Scope of work for full restoration is to be determined. All work is anticipated to be restricted to tunnel.
Kaauau (# 1 intake - Hauolo Wahine Ditch)	K-29	Scope of work for full restoration is to be determined. All work is anticipated to be restricted to tunnel.
Hauolo Wahine	K-30	Bolt steel plates or concrete over diversion intake grate.
Kaauau diversion tunnel to # 1 intake	K-29a	Seal diversion ditch with rock and concrete.
Hauolo Wahine small diversion	K-30a	Remove pipe.
Hauolo Wahine small intake	K-30b	Remove pipe.
Hauolo Wahine small intake	K-30c	Remove pipe.
Waiokamilo		
Kualani (East Waiokamilo)	K-22	Diversion closed in 2007. Concreted over diversion intake.
Waiokamilo # 11 intake	K-23	Diversion closed in 2007. Blocked water from entering ditch with boards located in tunnel.
Waiokamilo # 12 intake	K-24	Diversion closed in 2007. Concreted over diversion intake.
Waiokamilo Kikokiko intake	K-25	Diversion closed in 2007. Concreted over diversion intake.
Lalapipi Ditch Diversion	K-21b	No Diversion.
Kualani alamahuna pipe	K-22a	Diversion closed in 2007. Removed pipe.
Koolau ditch # 10 crosscut intake # 1	K-22b	Diversion closed in 2007. Removed pipe.
Koolau ditch # 10 crosscut intake # 2	K-22c	Diversion closed in 2007. Removed pipe.
Koolau ditch # 10 crosscut intake # 3	K-22d	Diversion closed in 2007. Removed pipe.
Koolau ditch # 10 crosscut intake # 4	K-22e	Diversion closed in 2007. Removed pipe.
Koolau ditch # 10 crosscut intake # 5	K-22f	Diversion closed in 2007. Removed pipe.
Koolau ditch # 10 crosscut intake # 6	K-22g	Diversion closed in 2007. Removed pipe.
5' PVC pipe intake East of # 11 intake	K-23a	Diversion closed in 2007. Removed pipe.
Koolau Ditch # 12 crosscut intake # 1	K-24a	Diversion closed in 2007. Removed pipe.
Koolau Ditch # 12 crosscut intake # 2	K-24b	Diversion closed in 2007. Removed pipe.
Koolau Ditch # 12 crosscut intake # 3	K-24c	Diversion closed in 2007. Removed pipe.
Koolau Ditch # 12 crosscut intake # 4	K-24d	Diversion closed in 2007. Removed pipe.
Koolau Ditch # 12 crosscut intake # 5	K-24e	Diversion closed in 2007. Removed pipe.
Small intake west of main # 12 crosscut	K-24f	Diversion closed in 2007. Removed pipe.
Small intake west of main # 12 crosscut	K-24g	Diversion closed in 2007. Removed pipe.
Small intake west of main # 12 crosscut	K-24h	Diversion closed in 2007. Removed pipe.
Small intake west of main # 17 crosscut	K-24i	Diversion closed in 2007. Removed pipe.
Small intake west of main # 12 crosscut	K-24j	Diversion closed in 2007. Removed pipe.
East Kikokiko 7" pipe intake	K-25a	Diversion closed in 2007. Removed pipe.
Kikokiko small intake	K-25b	Diversion closed in 2007. Removed pipe.

Kikohiko #1 pipe intake manuka of bridge	K-25c	Diversion closed in 2007. Removed pipe.
West Kikohiko #4 pipe intake	K-25d	Diversion closed in 2007. Removed pipe.
West Kikohiko #3 pipe intake	K-25e	Diversion closed in 2007. Removed pipe.
Kikohiko #2 PVC pipe intake water bridge	K-25f	Diversion closed in 2007. Removed pipe.
Wailuanui (East and West)		
East Wailuanui # 6 intake and sluice basin	K-18	Seal intake opening with rocks and concrete.
East Wailuanui # 6 control house intake	K-19	Bolt steel plates or concrete over diversion intake grate.
Wailuanui # 7 intake	K-20	Seal intake opening with rocks and concrete.
West Wailuanui # 9 intake	K-21	Seal intake opening with rocks and concrete.
3" aluminum pipe intake by # 9 control house intake		
Wailuanui stream intake # 8 intake pipe	K-19a	Remove pipe.
8" steel pipe intake East of # 9 intake	K-20a	Remove pipe
	K-21a	Remove pipe

Total diversions = 69 (Not including Filipino Ditch Diversion) - 28 Waioakamilo diversions that were closed in 2007 = 41
Major diversions registered with CWRM in 1989
Minor diversions registered with CWRM in 1989

Phase I Projects			Timetable to Complete Project
Stream Diversion	EMI Map #	General Description of Work Plan	
Lowrie Honopou side ditch	L-17	Close existing control gate.	5 to 7 months after obtaining all required approvals and completing any required consultations
Haiku East Hanehoi (Pancho)	H-3	Bolt steel plates or concrete over diversion intake grate.	
Haiku West Hanehoi (School)	H-4	Seal intake opening with rock and concrete.	
Lowrie Hanehoi Huelo # 1	L-5	Bolt steel plates or concrete over diversion intake grate.	
Koolau Kano	K-26	Most flow will be restored by removal of sluice gate. Scope of work for full restoration is to be determined. All work is anticipated to be restricted to tunnel.	
Lowrie Hanehoi Huelo # 2	L-6	Bolt steel plates or concrete over diversion intake grate.	
Lowrie Hanehoi Huelo # 3	L-7	Bolt steel plates or concrete over diversion intake grate.	
Koolau Lalaha (# 3 intake - Hauolo Wahine Ditch)	K-27	Scope of work for full restoration is to be determined. All work is anticipated to be restricted to tunnel.	
Koolau Lalapipi (# 2 intake - Hauolo Wahine Ditch)	K-28	Scope of work for full restoration is to be determined. All work is anticipated to be restricted to tunnel.	
Kaaau (# 1 intake - Hauolo Wahine Ditch)	K-29	Scope of work for full restoration is to be determined. All work is anticipated to be restricted to tunnel.	
Hauolo Wahine	K-30	Bolt steel plates or concrete over diversion intake grate.	
	K-29a	Seal diversion ditch with rock and concrete.	
	K-30a	Remove pipe.	
	K-30b	Remove pipe.	
	K-30c	Remove pipe.	

Phase II Projects			Timetable to Complete Project
Stream Diversion	EMI Map #	General Description of Work Plan	
New Hamakua Hanehoi (Huelo intake)	NH-17	Bolt steel plates or concrete over diversion intake grate.	2 to 3 months after obtaining all required approvals and completing any required consultations
Koolau East Waiuanui # 6 intake and sluice basin	K-18	Seal intake opening with rocks and concrete.	
Koolau West Waiuanui # 9 intake	K-21	Seal intake opening with rocks and concrete.	
Koolau East Waiuanui # 6 control house intake	K-19	Bolt steel plates or concrete over diversion intake grate.	
Koolau Waiuanui # 7 intake	K-20	Seal intake opening with rocks and concrete.	
	K-19a	Remove pipe.	
	K-20a	Remove pipe.	
	K-21a	Remove pipe.	

Phase III Projects			Timetable to Complete Project
Stream Diversion	EMI Map #	General Description of Work Plan	
Wailoa Honopou	W-22	Bolt steel plates or concrete over diversion intake grate.	4 to 5 months after obtaining all required approvals and completing any required consultations
New Hamakua Honopou	NH-22	Bolt steel plates or concrete over diversion intake grate.	
Haiku Honopou	H-8	Bolt steel plates or concrete over diversion intake grate and seal opening below grate with rock and concrete.	
Wailoa Hanehoi (Huelo intake)	W-18	Bolt steel plates or concrete over diversion intake grate.	
Koolau Pinaau	K-31	Seal intake opening with rocks and concrete.	
	W-22a	Bolt steel plates or concrete over diversion intake grate.	
	W-22b	Bolt steel plates or concrete over diversion intake grate.	
	K-30d	Construct overpass over ditch.	
	K-31a	Remove steel and pvc pipes.	
	NH-23	Seal intake opening with rocks and concrete.	

Phase IV Projects			
Stream Diversion	EMI Map #	General Description of Work Plan	Timetable to Complete Project
Lowrie Honopou long strainer	NH-17a	Construct stream overpass over ditch.	17 to 23 months after obtaining all required approvals and completing any required consultations
Lowrie Honopou long siphon	L-7a	Install pipe or box culvert with wing walls through which ditch can pass beneath stream or construct stream overpass over ditch.	
	L-7b	Construct stream overpass over ditch.	
	L-5a	Construct stream overpass over ditch.	
	L-5b	Construct stream overpass over ditch.	
	L-5c	Construct stream overpass over ditch.	
	L-15	Construct stream overpass over ditch.	
	L-16	Construct stream overpass over ditch.	

Major diversions registered with CWRM in 1989

Minor diversions registered with CWRM in 1989

Honopou / Hanehoi




Exhibit "E-164"

Hanehoi & Puolua Streams and
their tributaries
Major Diversions to be restored

EXHIBIT E-163

HANEHOI STREAM & TRIBUTARIES

Hanehoi stream is diverted by:

Wailoa & New Hamakua Ditches

E. & W. Hanehoi streams and their tributary, Huelo Stream
are all separately diverted by Lowrie Ditch

Hanehoi Stream, after its confluence with Huelo and W.
Hanehoi stream, is diverted by Haiku Ditch

Wailoa Ditch Diversion- Hanehoi Stream (W-18) 191.6



Water collected thru diversion grate (57" wide x 78" long)
Grate proposed to be sealed w/ bolted iron plate or concrete.

- no plan given for maintenance of modified diversion structure
 - no plan for any adaptation needed for streamlife passage
- Phase III project: no status report on needed permits

view: downstream

MT EXH E-163; Photo #1
5/2016

Wailoa Ditch Diversion (W-18) 191.6)



Diversion structure W-18 spans Hanehoi stream channel. Minimal flow in stream- May 2016 below ditch structure. flow collects in stagnant pond in foreground of picture

MT EXH E-163; Photo #2
5/2016

Wailoa Ditch Diversion-Hanehoi Stream (W-18) 191.6



Diversion structure W-18 Hanehoi Stream

Sluice gate on stream channel.

Gate open 1 in. or less allows minimal flow in stream past ditch structure.

No adjustment to gate noted in A&B/EMI reports as of October 2016

MT EXH E-163; Photo #3
5/2016

Wailoa Ditch Diversion- Hanehoi Stream (W-18) 191.6



Diversion structure
W-18- showing East
Hanehoi natural
stream flow.

View is to upstream of
diversion structure

MT EXH E-163; Photo #4
5/2016

Wailoa Ditch Diversion-Hanehoi Stream (W-18) 191.6



E. Hanehoi stream immediately below Diversion structure W-18- showing stream bed with little flow except wet rocks downstream of diversion structure and stagnant pond

MT EXH E-163; Photo #5
5/2016

New Hamakua Ditch Diversion- Hanehoi Stream (NH-17) intake 264.6



MT EXH E-163; Photo #6
5/2016

Hanehoi Diversion structure NH-17/ 264.6

Dam & grate on East Hanehoi stream channel, several hundred ft. makai (downstream) of Wailoa intake 191.6.

- proposed to fill grate area with concrete
- appears this intake has not had regular flows to divert.
 - grate covered in debris (right side behind dam wall- see next slide)

New Hamakua Ditch Diversion-Hanehoi Stream (NH-17) intake 264.6 Diversion structure NH-17/ 264.6



Closeup of existing grate on East Hanehoi stream channel. Intake grate is 88" across at dam end and much narrower (48" wide) at upstream end. Grate is 24 ft long and covered in debris (May 2016)

- Portion of dam should be removed to facilitate flows and streamlife migration- not discussed in HC&S/EMI plan
- no plans for maintenance of structure discussed

Phase III project: no status report on needed permits.

MT EXH E-163; Photo #7
5/2016

New Hamakua Ditch Diversion-Hanehoi Stream (NH-17) intake 264.6



Hanehoi stream,
immediately
upstream of
Diversion structure
NH-17/ 264.6 on
East Hanehoi stream
channel. Stream
bed overgrown w/
invasive plants w/
little flow

MT EXH E-163; Photo #8
5/2016

New Hamakua Ditch Diversion-Hanehoi Stream (NH-17) intake 264.6



East Hanehoi stream immediately downstream of Diversion structure NH-17/ 264.6 on East Hanehoi stream channel.

- Stream bed overgrown w/ invasive trees and plants.
- No plan for cleaning or maintenance

MT EXH E-163; Photo #9
5/2016

Lowrie Ditch Diversion- E. Hanehoi Stream (L-5) intake 240.6



MT EXH E-163; Photo #10

5/1989

East Hanehoi stream at Lowrie Ditch

Diversion structure
L-5/ 240.6 on East Hanehoi stream
channel. Huelo Community pipe
(Intake 538.6) located upstream a
few hundred yards.

- Intake grate proposed to be covered with iron plate or cement.
- no plan given for maintenance of modified diversion structure
- no plan for any adaptation needed for streamlife passage

Phase I project: no status report on
needed permits

Lowrie Ditch Diversion- E. Hanehoi Stream (L-5) intake 240.6



East Hanehoi stream Lowrie Ditch intake

- bypass pipes in diversion structure are only current source of East Hanehoi flow above Hana Hwy
 - Pipes are subject to blockage on upstream side.

HC&S was asked to modify this diversion to allow native streamlife to travel.

MT EXH E-163; Photo #11

8/2013

Lowrie Ditch Diversion- E. Hanehoi Stream (L-5) intake 240.6
Close up of bypass pipes shown in previous slide.
Pipes commonly get clogged.



MT EXH E-163; Photo #12
8/2013

Lowrie Ditch Diversion- E. Hanehoi Stream (L-5) intake 240.6



East Hanehoi stream at Lowrie Diversion structure

L-5/ 240.6 on East Hanehoi
stream channel. View
upstream.

Two 4" pipes (left side of
diversion) carry water into
a stagnant pond below
diversion. This is only
water that bypasses Lowrie
diversion except during
storms.

MT EXH E-163; Photo #13

7/2012

Lowrie Ditch Diversion- E. Hanehoi Stream (L-5) intake 240.6



East Hanehoi stream Intake Lowrie Ditch

Far view, immediately downstream of Diversion structure L-5/ 240.6 on East Hanehoi stream channel. Stream bed overgrown w/ invasive plants.

EMI service road crosses stream bed (cement area)

MT EXH E-163; Photo #14
7/2012

Pool Above Lowrie Ditch Diversion- E. Hanehoi Stream intake 538.6



East Hanehoi stream pool
above Diversion structure
L-5/ 240.6 on East
Hanehoi stream channel.

Huelo Community pipe
(Intake 538.6) located
upstream a few hundred
yards in a small pool.

MT EXH E-163; Photo #15
8/2013

Lowrie Ditch Diversion- W. Hanehoi Stream (L-6) intake 242.6



MT EXH E-163; Photo #16
10/2009

West Hanehoi stream intake at Lowrie ditch

Diversion structure
L-6/ 242.6 on West
Hanehoi stream channel.

- intake not included in May restoration plan, but now is
- Intake grate proposed to be covered with iron plate or cement.
- no plan given for maintenance of modified diversion structure
 - no plan for any adaptation needed for streamlife passage

Phase I project: no status report on needed permits

Lowrie Ditch Diversion- W. Hanehoi Stream (L-6) intake 242.6



**West Hanehoi stream
intake at Lowrie ditch**

Far View of Diversion
structure
L-6/ 242.6 on West
Hanehoi stream
channel.

View is downstream of
diversion intake.

MT EXH E-163; Photo #17
10/2009


Lowrie Ditch Diversion- W. Hanehoi Stream (L-6) intake 242.6



Spillway crossing at
W. Hanehoi Stream
over Lowrie Ditch
makai view

MT EXH E-163; Photo #18

10/2009



A&B EXH C-01 Map

Red circle

shows where E. & W. Hanehoi Streams meet, just mauka of Haiku ditch.

Blue circle

is where W. Hanehoi and Huelo streams meet, just makai of Lowrie ditch

Shown on EXH C-01-A&B Map of streams and ditch system

E. & W. Hanehoi, Huelo and Puolua streams each have a diversion intake @ Lowrie ditch (red dots).

CRWM Maps should indicate all diverted stream branches

MT EXH E-163 Map #1

Lowrie Ditch Diversion- Huelo Stream (L-7) intake 155.6



Huelo stream intake at Lowrie ditch

Diversion structure L-7/ 155.6 on Huelo stream channel.

- intake not included in May restoration plan, but now is
- Intake grate proposed to be covered with iron plate or cement.
- no plan given for maintenance of modified diversion structure
 - no plan for any adaptation needed for streamlife passage

Phase I project: no status report on needed permits

MT EXH E-163; Photo #19
5/1989

Lowrie Ditch Diversion- Huelo Stream (L-7) intake 155.6



**Huelo stream intake at
Lowrie ditch**

View down Lowrie ditch
of diversion structure
L-7/ 155.6 on Huelo
stream channel.

MT EXH E-163; Photo #20

10/2009

Lowrie Ditch Diversion- Huelo Stream (L-7) intake 155.6



**Huelo stream intake at
Lowrie Ditch**

**Spillway over
Lowrie Ditch
mauka view**

MT EXH E-163; Photo #21

10/2009

Lowrie Ditch Diversion- Huelo Stream (L-7) intake 155.6



Huelo stream intake at Lowrie ditch

Close View of Lowrie
ditch intake dam and
grate L-7/ 155.6 on Huelo
stream channel.

This stream should be
added to CWRM maps


MT EXH E-163; Photo #22

10/2009

gistered diversions and EMI minor diversions identified in the Hanehoi hydrologic unit (Source: East Maui
, 1970; State of Hawaii, Commission on Water Resource Management, 2008e).

156°14'W

156°13'W



Huelo stream & intake actually exists, but is mislocated on CWRM's IFSTAR Map for Hanehoi Unit (2008).

Puolua stream is incorrectly labelled "Huelo Stream" on this map. They are not the "same stream"

See A&B's EXH C-01 map in Slide 20 for correct locations.

Lowrie Intake 155.6 (L-7) is shown on this CWRM IFSTAR map on what is really Puolua stream intake (L-7a). this stream is mislabelled "Huelo" stream. Huelo stream is added in (green line) on this map. This is where intake 155.6 is located. Only E. & W. Hanehoi and Puolua streams were shown on original map

**Maui County Plat Map
2-9-09**

Clearly shows three
streams crossing Lowrie
ditch

Lowrie Ditch (Blue) & three
streams that are diverted
by the ditch.


East Hanehoi stream
(brite green) crosses
Lowrie ditch at intake L-5.

West Hanehoi stream
crosses Lowrie ditch at
intake L-6.


Huelo stream (dark green)
crosses Lowrie Ditch at
intake L-7.

**Puolua stream is on Plat
map 06 so it does not
show here.**

Hana Hwy is red



MT EHB E-163 Map # 3



**County Plat Map
2-9-06**

Huelo Stream Lowrie Intake 155.6 (L-7) (blue circle) is shown.

Puolua stream intake on Lowrie Ditch is circled in red.

They are two completely different streams with separate diversions

Huelo stream is not shown on CWRM maps but is on Plat maps and EMI maps

MT EXH E-163 Map #4

ADVANCE SHEET
SUBJECT TO CHANGE

SECOND DIVISION	
ZONE	SEC. PLAT
2	9 06
CONTAINING PARCELS	
SCALE: 1 in. = 200 ft.	

PRINTED.....

New Haiku Ditch Diversion- Hanehoi Stream (H-3) intake 217.6



Hanehoi stream immediately up stream of Diversion structure H-3/ 217.6 on Hanehoi stream channel. Grate on left transports stream water to New Haiku Ditch tunnel.

Concrete dam blocks downstream flow and becomes filled up with large rocks.

West and East branches of Hanehoi stream join just mauka (upstream) of the Haiku Ditch diversion & become one stream (Hanehoi)

MT EX #23

5/2016

New Haiku Ditch Diversion- E. Hanehoi Stream (H-3) intake 217.6



MT EXH E-163; Photo #24
5/2016

Close view of Haiku intake grate on Hanehoi stream. Part of Diversion intake structure H-3/ 217.6 on Hanehoi stream channel.

- grate proposed to be sealed w/steel plate or concrete
- dam needs modification to allow stream life migration
- modification needed to allow larger rocks to move naturally downstream.
 - stagnant water collects in lower diversion trough (Phot 24-a)

New Haiku Ditch Diversion- E. Hanehoi Stream (H-3) intake 217.6



Hanehoi stream intake at New Haiku ditch

New Haiku ditch intake dam has a lower level that has been filled with stagnant water since the sluice grate has been open at H-3 diversion (July 2016 view)

- stagnant water can breed large mosquito population
- this area should be modified to allow water to drain and streamlife to migrate

MT EXH E-163; Photo #24-a
7/2016

New Haiku Ditch Diversion- E. Hanehoi Stream (H-3) intake 217.6

New Haiku Ditch



Hanehoi stream
Diversion structure
H-3/ 217.6 on Hanehoi
stream channel. Sluice
gate completely closed.
(May 2016) Only minimal
flows bypassed H-3
diversion structure.

Gate opened June 2016,
now large stones piling
up along dam and
opening

MT EXH E-163; Photo #25
5/2016

New Haiku Ditch Diversion- Hanehoi Stream (H-3) intake 217.6



New Haiku Ditch Hanehoi stream

Makai view of mostly dry Hanehoi stream immediately downstream of New Haiku diversion intake 217.6 (H-3) in May before sluice gates opened .

EMI owns overgrown stream bed and does not maintain.

Some flow enters the Hanehoi stream since the sluice gate was opened in June 2016 but it is beginning to erode the left bank, since the gate concentrates all flow on that side of the stream

MT EXH E-163; Photo #26
5/2016

Puolua Stream Diversions

Puolua stream is diverted by:
New Hamakua
Lowrie &
Haiku Ditches

New Hamakua Ditch Diversion- Puolua Stream (NH-17a) intake



MT EXH E-163; Photo #27
5/2016

New Hamakua Ditch Puolua stream

Puolua flow is captured directly into the New Hamakua Ditch.

New Hamakua Ditch and EMI service road would need modification to allow flow to continue makai

Stream “over pass” is proposed for Puolua intake NH-17-a

- no plan to allow any migrating streamlife
- Phase IV project. 1 to 2 years in future. No update on permit status

Lowrie Ditch Diversion- Puolua Stream (L-7a) intake



MT EXH E-163; Photo #28

3/2015

Lowrie Ditch Puolua stream

Puolua flow is captured directly into the Lowrie Ditch with no grate structure. Stream is essential to Huelo community

Lowrie Ditch and EMI service road would need modification to allow flow to continue makai

- Stream “over pass” is proposed for Puolua intake L-7-a
- no plan to allow any migrating streamlife
- Phase IV project. 1 to 2 years in future. No update on permit status



Lowrie Ditch near Puolua stream

Lowrie Ditch has open sections and tunnel sections.

A dirt EMI service road runs just makai (to right) of the ditch

MT EXH E-163; Photo #29
3/2015

Lowrie Ditch Diversion- Puolua Stream (L-7a) intake (Oct 2009)



Lowrie Ditch Puolua stream

For decades these two leaky 4" pipes ran under EMI service road to a 8" PVC pipe and provided the only downstream flow into Puolua stream past Lowrie Ditch intake. Pipes were often clogged.

EMI replaced pipes with 8 inch pipe in 2015

MT EXH E-163; Photo #30

10/2009



**Video of Lowrie Ditch
at Puolua stream
intake**

Video shows how a portion of Puolua flow is channeled towards new 8 inch wide bypass pipe, while the remaining flow is captured directly into the Lowrie Ditch.
Video 3/2015

**MT EXH E-163 Vid #1
3/2013**

Lowrie Ditch Diversion- Puolua Stream (L-7a) intake (March 2015)



Lowrie Ditch at Puolua stream

Lowrie Ditch flows under new 8" bypass pipe and into a tunnel. A dirt EMI service road runs along ditch, and Puolua stream continues makai of the road

MT EXH E-163; Photo #3 | 3/2015



Channel
to bypass
pipe



Lowrie Ditch at Puolua stream

Puolua stream was modified to create a channel for a portion of stream water to flow into new 8" bypass pipe that passes under the EMI service road and empties into Puolua stream makai of the road

- after months of storm surges the bypass pipe entrance and channel leading to pipe are clogged w/ debris 32-a
- No access keys are offered residents to facilitate maintenance and they must hike in

MT EXH E-163; Photo #32
3/2015

Lowrie Ditch Diversion- Puolua Stream (L-7a) intake (October 2016)



Narrow channel leading to Puolua bypass pipe can
become clogged w/ debris
View after debris recently cleaned

MT EXH E-163; Photo #32-a



Lowrie Ditch Puolua stream bypass pipe

Makai end of new 8" bypass pipe that currently passes under EMI dirt service road and conveys stream water into Puolua stream.

Puolua stream has a bend where dirt road crosses and restoring the stream across the service road will require a concrete area for stream flow and native stream life

MT EXH E-163; Photo #33
3/2015

Lowrie Ditch Puolua stream intake L-7a

Far view of Lowrie Ditch intake at Puolua stream.

Overgrown Puolua stream bed (owned by EMI) is in foreground. Bypass pipe passes under service road & empties into stream (red circle.)

Road would need to be modified if stream flowed. Blue line shows natural curve in stream channel



MT EXH E-163; Photo #34
3/2015



Puolua stream just upstream of Lowrie Ditch Diversion L-7a

Puolua stream bed and surrounding land are all “owned” by EMI/A&B, who do not maintain the stream channel. It is choked with alien species and fallen trees, impeding stream flows.

MT EXH E-163; Photo #35

3/2015

New Haiku Ditch Diversion- Puolua (H-4) intake (2/2009)



MT EXH E-163; Photo #35
2/2009

Haiku Ditch Intake at Puolua stream

Puolua stream channel is dammed and passed through a grate into New Haiku Ditch tunnel.

Bypass pipes (red circle) at this diversion allow some water to return to stream as does an open sluice gate



New Haiku Ditch Diversion- Puolua (H-4) intake

EMI Sluice Gate on Puolua Stream Feb 2009

- gate has been opened completely as of June 2016
- grate behind dam proposed to be sealed with plate or concrete
- other areas of diversion need to be modified, but there has been no consultation with community
- Phase I project. No update on permit status

Area appears to have been
cleared and cleaned by EMI for 2009 CWRM visit.
Usually, area is very overgrown
(next slide)

MT EXH E-163; Photo #36
2/2009

New Haiku Ditch Diversion- Puolua (H-4) intake (4/2011)



Haiku Ditch Intake gate at Puolua stream

Same sluce gate as last slide, two years later- April 2011- on Haiku Ditch/Puolua intake.(same white bucket on diversion gate handle)

Passing of two years has brought no regular maintenance by the landowner EMI/A&B.

Visits by CWRM to Puolua stream to monitor IIFS compliance ended in 2009.

- Intake currently (10/2016) in this overgrown condition due to no EMI maintenance

MT EXH E-163; Photo #36
4/2011

New Haiku Ditch Diversion- Puolua (H-4) intake (4/2011)



Haiku Ditch Intake gate at Puolua stream

View downstream to Haiku Ditch/Puolua intake wall (arrow) buried under fallen Roseapple trees.

No regular maintenance of stream channel by the landowner EMI/A&B. This must be part of stream restoration plan.

Visits by CWRM to Puolua stream to monitor IIFS compliance ended in 2009.

MT EXH E-163; Photo #39
4/2011



Feb 2009
Puolua Stream
Waterfall and pool just
upstream of Haiku Ditch on
EMI/A&B land.

This pool once had
recreational use by local
families, but the stream bed
is now overgrown

MT EXH E-163; Photo #40
2/2009

Conclusions:

- W. Hanehoi and Huelo Lowrie Ditch stream diversions need to be clearly noted on CWRM Maps. They are tributaries of Hanehoi stream and have CWRM registered Major Diversions (L-6 and L-7 on EMI map)
- Maintenance of abandoned stream diversion structures and stream channels and needs to be addressed in the restoration plan.
 - Puolua/New Hamakua and Puolua/Lowrie diversions will need modifications of the EMI service roads. Possibly other diversions as well.
- The dam structure on Hanehoi and Puolua streams @New Haiku ditch will likely need to be modified to let large stones travel along stream.

EXHIBIT "A"
SECOND AMENDED CHART
HANEHOI WATERSHED

Diversions	East Hanehoi	West Hanehoi	Huelo (East and West Branches)	Puolua (East and West Branches)
<p align="center">Wailoa</p>	<p>Div. #1 W-18 Reg. Intake 191.6 Wailoa/ Hanehoi (III) MT Ph. #4, 2, 5</p> <p>Grate diversion MT Ph. #1</p> <p>sluice gate open only 1" MT Ph. #3</p> <p>Dam can be removed (Ditch in tunnel many feet below diversion) MT Ph. #4</p> <p>EMI: Intent is to seal 3 ft x 5 ft intake grate by bolting steel plate. Haiku, Lowrie and New Hamakua ditch project needs to be completed prior to work on this diversion.</p> <p>(Phase III 4-5 mos) Blue/#3: 3 to 4 months after obtaining all required approvals and completing any required</p>	<p align="center">N/A</p>	<p align="center">N/A</p>	<p align="center">N/A</p>

	consultations AC, 1, 3			
New Hamakua	<p>Div. #2 NH-17 New Hamakua/ Hanehoi (IV) Reg. Intake 264.6 MT Ph. #8, 9</p> <p>Grate diversion (covered by debris) MT Ph. #6, 7</p> <p>Dam could be removed (grate to side of dam) MT Ph. #6, 7</p> <p>EMI: Sealing of stream intake is by bolting plates over intake grate. Haiku and Lowrie ditch project work needs to be completed prior to work on this diversion.</p> <p>Option of passing stream in pipe/culvert over ditch, but this may complicate CWA Section 404 exemption. Haiku and Lowrie ditch projects needs to be completed prior to work on this diversion.</p> <p>(Phase II- 2-3 mos)</p>	<p>Div. #1 NH-17a (EMI) New Hamakua/ West Hanehoi(IV) MT Ph. #27</p> <p>EMI:</p> <p>Ditch is cut into stream bed, so would need to install a pipe or box culvert with wing walls in the stream bed through which the ditch can pass beneath the stream. Anticipate up to 42-inch diameter pipe to handle 60 MGD. Haiku ditch project needs to be completed prior to work on this diversion.</p> <p>Pink/ #4 12 to 15 months after obtaining all required approvals and completing any required consultations Option of passing stream in pipe/culvert over ditch, but this may complicate CWA Section 404 exemption. Haiku and Lowrie ditch projects needs to be completed prior to work on this diversion.</p>		<p>Div. #1 NH-17a (NH-18a) (EMI) (incorrect) New Hamakua/ Puolua (IV) MT Ph. #29, 30</p> <p>Stream crosses road above ditch, cars cross stream</p> <p>EMI: Ditch is cut into stream bed, so would need to install a pipe or box culvert with wing walls in the stream bed through which the ditch can pass beneath the stream. Anticipate large diameter pipe to handle up to 100 MGD. (Construct stream overpass over ditch)</p> <p>Option of passing stream in pipe/culvert over ditch, but this may complicate CWA Section 404 exemption. Haiku and Lowrie ditch projects needs to be completed prior to work on this diversion.</p> <p>(Phase IV 17-23 mos) Pink #4 1,2,3,4</p>

	Green/#2 1 to 2 months after obtaining all required approvals and completing any required consultations AC,1,2,3			5 to 6 months after obtaining all required approvals and completing any required consultations
Lowrie	Div. #3 L-5a EMI: Ditch is cut into stream bed, so would need to install a pipe or box culvert with wing walls in the stream bed through which the ditch can pass beneath the stream. Anticipate up to 42-inch diameter pipe to handle 60 MGD. Haiku ditch project needs to be completed prior to work on this diversion. (Construct stream overpass over ditch) (Phase IV 17-23 mos) 12 to 15 months after obtaining all required approvals and completing any required consultations Div. # 4 L-5 Reg. Intake 240.6 Lowrie-Hanehoi (I)	Div. #2 L-5b Lowrie-West Hanehoi #1 (IV) EMI: Ditch is cut into stream bed, so would need to install a pipe or box culvert with wing walls in the stream bed through which the ditch can pass beneath the stream. Anticipate up to 42-inch diameter pipe to handle 60 MGD. Haiku ditch project needs to be completed prior to work on this diversion. (Construct stream overpass over ditch) (Phase IV 17-23 mos) 12 to 15 months after obtaining all required approvals and completing any required consultations Div. #3 L-5c Lowrie-Hanehoi small	Div. #1 L-7 Reg. Intake 155.6 Lowrie-Hanehoi Huelo #3 (I) MT Ph. #19, 20, 21, 22 Grate diversion MT Ph. #22 Dam could be removed (Grate is to one side of ditch) EMI: Sealing of stream intake is by bolting plates over intake grate. Need to repair leakage into ditch along edge if grate first. Haiku ditch project work needs to be completed prior to work on this diversion. (Phase I – 5-7 mos) Yellow/#1 4 to 6 months after obtaining all required approvals and completing any required consultations	Div. #2 L-7a Reg. Intake 225.6 Lowrie-Puolua (IV) (Roseapple) MT Ph. #28, 29 Deteriorating 4” pipes thereafter joined to 8” pipe; 4” pipes regularly blocked with debris; Just above Schupp kuleana MT Ph. #30, 31, 32, 33, 34, 35, MT Vid. #1 Stream crosses road above ditch, cars cross stream EMI: Ditch is cut into stream bed, so would need to install a pipe or box culvert with wing walls in the stream bed through which the ditch can pass beneath the stream. Anticipate up to 42-inch diameter pipe to handle 60 MGD. Haiku ditch project needs to be completed

	<p>Grate diversion MT Ph. #10</p> <p>2 4" bypass pipes totally blocked MT Ph. #11, 12, 13, 14</p> <p>Huelo Community pipe is above this diversion East Hanehoi Stream Intake 538.6 MT Ph. #15</p> <p>EMI: Sealing of stream intake is by bolting plates over intake grate. Need to repair leakage into ditch along edge if grate first. Haiku ditch project work needs to be completed prior to work on this diversion.</p> <p>Yellow/#1 4 to 6 months after obtaining all required approvals and completing any required consultations 1,3</p>	<p>EMI: Ditch is cut into stream bed, so would need to install a pipe or box culvert with wing walls in the stream bed through which the ditch can pass beneath the stream. Anticipate up to 42-inch diameter pipe to handle 60 MGD. (Construct stream overpass over ditch)</p> <p>Haiku ditch project needs to be completed prior to work on this diversion.</p> <p>(Phase IV 17-23 mos) Pink/ #4 12 to 15 months after obtaining all required approvals and completing any required consultations</p> <p>Div. #4 L-6 Reg. Intake 242.6 Lowrie-Hanehoi Huelo #2 (I) MT Ph. #16, 17, 18</p> <p>EMI: Sealing of stream intake is by bolting plates over intake grate. Need to</p>	<p>Div. #2 L-7b</p> <p>Lowrie-West Hanehoi #2 (IV) (incorrect)</p> <p>EMI: Ditch is cut into stream bed, so would need to install a pipe or box culvert with wing walls in the stream bed through which the ditch can pass beneath the stream. Anticipate up to 42-inch diameter pipe to handle 60 MGD. Haiku ditch project needs to be completed prior to work on this diversion. (Construct stream overpass over ditch)</p> <p>(Phase IV 17-23 mos) Pink/ #4 12 to 15 months after obtaining all required approvals and completing any required consultations</p>	<p>prior to work on this diversion. (Install pipe or box culvert with wing walls through which ditch can pass beneath stream or construct overpass over ditch)</p> <p>(Phase IV 17-23 mos) Pink/#4 1,2,3,4 12 to 15 months after obtaining all required approvals and completing any required consultations</p>
--	--	---	---	---


		<p>repair leakage into ditch along edge if grate first. Haiku ditch project work needs to be completed prior to work on this diversion.</p> <p>(Phase I – 5-7 mos) Yellow/ #1 4 to 6 months after obtaining all required approvals and completing any required consultations</p>		
Haiku	<p>Div. #5 H-3 (Pancho) Reg. Intake 217.6 Haiku-Hanehoi (I) MT Ph. #26</p> <p>Grate diversion MT Ph. #23, 24</p> <p>Sluice completely Closed 4/16 MT Ph. #25</p> <p>Pipes removed in last 10-15 yrs Holes left were filled with cement</p> <p>Dam can be removed (Grate is to side of ditch) MT Ph. #24</p> <p>EMI: Sealing of stream intake opening is by bolting plate over intake grate openings in intake</p>			<p>Div. #3 H-4 (School) Haiku-Puolua (I)</p> <p>Waterfall and pool just above Haiku diversion MT Ph. #40</p> <p>Grate diversion MT Ph. #39</p> <p>Bypass pipes MT Ph. #36</p> <p>Sluice gate should be opened MT Ph. #37, 38,</p> <p>Dam can be removed (Ditch is in tunnel many feet below dam)</p> <p>EMI: Sealing of stream intake is by bolting small plate over opening into</p>

	<p>grate.</p> <p>(Phase I – 5-7 mos) Yellow/#1 4 to 6 months after obtaining all required approvals and completing any required consultations 1, 3</p>			<p>ditch (not in stream). (Seal intake opening with rock and concrete)</p> <p>(Phase I – 5-7 mos) Yellow/#1 4 to 6 months after obtaining all required approvals and completing any required consultations</p>
--	--	--	--	--

- 1 CWRM approval – diversion abandonment permit
- 2 OCCL (conservation) site plan approval
- 3 USACE consultation/approval
- 4 Extensive work required
- 5 DoH permit

Mobile: mobile.mauinews.com

Subscribe | Submit News | Vac Rental | E-Edition | Home



ORDER LIVE UH FOOTBALL NOW!
 Order on Digital Channel 254.1254 or call 643-2100 [Learn More](#)

Oceanic REALTOR ASSOCIATION OF MAUI, INC.

Local News Opinion Blogs Sports Scene Vac Rental Ads Jobs Classifieds Contacts Extras TV

BREAKING NEWS» Baldwin edges Kamehameha Maui

Local News

- Local News
- Breaking News (free)
- Hawaii News
- Business
- Obituaries -- see also Online Newspaper
- Ads/In Memoriam
- Entertainment
- On The Campaign Trail
- Community
- Religion
- Opinion
- National News
- International News
- Weather
- SUBMIT NEWS: Virtual Newsroom

Biodiversity fest honors ind...

Maui REALTOR ASSOCIATION OF MAUI, INC.

Click here for local business and residential phone listings.

News Blogs & Events

SEARCH

News, Blogs & Events Web EZToUse.com

EZToUse! SLASH PAGES.com

Click here for local business and residential phone listings.

HC&S testing crops, like sorghum, that can be used for feed and fuel

Waving fields of . . . sorghum?

By MELISSA TANII - Staff Writer (mtanii@mauinews.com) , The Maui News

Save | Post a comment

PUNENE - Sorghum is a top contender to replace sugar cane in some of Hawaiian Commercial & Sugar Co.'s 36,000 acres after the plantation closes at the end of this year.

"We feel the most comfortable with it," HC&S General Manager Rick Volner said of sorghum, the grass crop that can be used for cattle feed and biofuel.

Now, several different varieties of sorghum are being grown on 182 acres of HC&S land. Trials began around five years ago, said Mae Nakahata, a 30-year HC&S veteran and director of agricultural research and crop control.

Article Photos



Sorghum, which has similar properties as sugar cane, L.



Silversea® Senior Cruise x

More Choices Than Any Luxury Line. Best Single Fares: Get a Quote Now!

Catch Your Cheater Spous x

Enter Their Email Address. See Hard To Find Pics & Social Profiles Now!

Orchid Growing Secrets+ x

The Most Important Things You Need To Know All About Growing Orchids!

Outstanding


Get great performance that can't be beat.

From \$149.99

HP ENVY 5660 e-All-in-One Printer

Shop now at HP

Free



Sorghum "seems to have a lot less pest problems than the other crops," she said.

And, after it's harvested, sorghum will regrow on its own, she said.

For now, the sorghum trial plantings are being given away for free as cattle feed to island ranches. Nakahata and Volner said sorghum could be sold to businesses and ranches. Ranchers have already offered to pay for sorghum to have an ample supply of feed, even after the sugar operations have ended, they said.

On Wednesday, a forage harvester was being driven through a field of sorghum - harvesting it with mechanical blades in a flat field off of Mokulele Highway near the Puunene Mill. Volner said sorghum is planted in flat, relatively rock-free fields to allow for equipment to easily harvest the crop. However, not all portions of the plantation are conducive to mechanical harvesting.

White cattle egrets flew by, landing and poking around in the green field in a search for bugs, a familiar sight after cane is burned and cane spiders and bugs scatter from the fields.

While birds can be seen as pests, especially when some species have been eating HC&S' trial crops, Nakahata said the egrets are a good sign because they eat the pests.

"They are the good birds," she said smiling.

Volner said another plus about sorghum is that it needs less irrigation than sugar cane.

Experts have said sorghum uses only about one-quarter as much water as cane. From afar, the sorghum plants look like sugar cane with their green, long, skinny shape and slender leaves. In the United States, some varieties are used as livestock feed or turned into ethanol.

Like the sorghum plantings, HC&S has a variety of other trial crops and diversified agriculture models under experimentation as the plantation's closure approaches.

Other experiments include energy crops, such as a sorghum plants and corn - both of which can be used for anaerobic conversion to biogas. HC&S is working with oil seed crops, such as soybeans and sunflowers for biodiesel production by crushing the seeds to extract oil. And trials are being conducted with grass-finished livestock on old sugar cane fields in Hamakuapoko.

Other planting trials are being done on plots smaller than an acre. Hemp could be a crop as well, company officials said.

Shyloh Stafford-Jones, a project manager for diversified crop operations, said that in addition to the sorghum crops, HC&S' larger experiments include different varieties of sunflowers on 2 acres, soybeans on 3 acres, corn on 3 acres and purple bana grasses on 2 acres.

He added that along with those crops, other types of crop cover and plants that help condition the soil are being planted in between crop rotations. These include large radishes, known to longtime residents as "daikon," which have roots that dig into the ground to help with soil compaction. A crop of legumes helps fertilize the soil with nitrogen.

Nakahata said the plantation is having trouble with birds eating soybean and sunflower plants and seeds.

She picked up a sunflower in a dusty field, which had nearly a quarter of its seeds eaten by birds. The seeds are what would be

used to provide oil for biodiesel.

The sunflowers are not typical ornamental sunflowers one would see sold in flower shops, Volner said. The ones the plantation is growing are for oil seed crops and are close to 3 feet high, with a smaller flower diameter than the larger ornamental ones used in bouquets.

Also nearby are soybeans, which could also be used for biodiesel.

Nakahata said a big problem with the birds was when plantings were going on for the oil seed crops, and the birds knew when seeds were being planted.

"The birds were literally lined up on the field waiting to go," she said.

Also, another test crop is the purple bama grasses, Volner said. The grasses from afar look like sugar cane, and they could be used for biofuel, he said.

While most of the test crops are being grown on flat parcels on the plantation near the mill, the grass-fed beef production project is ongoing at the higher and cooler elevations of the plantation on 30 acres of irrigated land in Hamakuapoko.

Work began about a year ago to get a variety of plants to grow in fields for cattle grazing. The crops included a variety of tropical grasses and legumes, which are high in protein and good for the cattle to eat, said Jacob Tavares, who does double duty as the livestock project manager and works in human resources at HC&S.

Another 150 acres is dedicated to grazing. But those lands consist of naturally grown vegetation, Tavares said.

Now, there are 104 head of adult cattle on the properties, with 24 steer on the 30 acres of irrigated fields; along with 50 steer, 30 cows and two calves on the 150 acres, HC&S said.

HC&S is working on the beef production project with Maui Cattle Co., which is made up of a partnership of Maui ranches. Cattle company officials have said the ranches produce 1,000 head of cattle annually, most of which stays on Maui. Some cattle are sent to Oahu.

Normally, it would take about a year for the cattle to get to an optimum weight of 1,200 pounds, Tavares said. The cattle now on the experimental HC&S properties were placed there in February. But so far they are gaining weight quick enough that it may take less than year for them to reach the optimum weight, he said.

The project is going "very, very well . . . exceeding our expectations," said Tavares, who has his own small ranch in Huelo.


The Hamakuapoko site is a good area for cattle-grazing crops because they grow best in areas that receive about 55 to 60 inches of rain a year.

Company officials estimated that perhaps 10,000 acres on the plantation could be used for cattle grazing.

* Melissa Tanji can be reached at mtanji@mauinews.com.

Copyright © 2016 The Maui News. All rights reserved. This material may not be published, broadcast, rewritten or redistributed.

Save | Post a comment

 Be the first of your friends to recommend this.



[Subscribe to The Maui News](#)

MĀLAMA `ĀINA: A CONVERSATION ABOUT MAUI'S FARMING FUTURE

A PROJECT OF THE MAUI TOMORROW FOUNDATION

Revised October 2016



Looking towards Iao Valley
Illustration by Silvia Yordanova

Report by Permaculture Design International LLC

EXHIBIT E-160



MAUI TOMORROW

Copyright 2016 by Maui Tomorrow Foundation, Inc.

Acknowledgements:

This report was funded by MAUI TOMORROW FOUNDATION, Inc. as part of its on-going commitment to state of the art sustainable development and the maintenance of Maui's rural lifestyle.

The content of this report was prepared by Permaculture Design International, LLC. for Maui Tomorrow.

Report Authors:

Jenny Pell, PDI
Loren Luyendyk, PDI

Contributors:

Neil Bertrando, PDI
Rico Zook, PDI
Koreen Brennan, PDI
Darren Doherty, Regrarians
Bill Reed, Regenesiis
Joel Glanzberg, Regenesiis
Alika Atay, Maui Farmer
Michael Howden, Maui Permaculture
Lucienne DeNaie, Maui Tomorrow
Albert Perez, Maui Tomorrow

Illustrations and Graphic Layout:

Silvia Yordanova
Loren Luyendyk
Jenny Pell

Disclaimer:

The material contained herein is based upon work supported by Maui Tomorrow. Any opinions, findings, conclusions, or recommendations are those of the authors and do not necessarily reflect the views of Maui Tomorrow, its employees or its administration.

Copyright:

No content may be reproduced without express written permission from Maui Tomorrow and Permaculture Design International.

Table of Contents

INTRODUCTION	1
A Brief Overview of Maui's "Central Valley" and Sugarcane	2
REGENERATIVE AGRICULTURE	3
Climate Change and Regenerative Agriculture	5
Regenerative Agricultural Land Use Potential and Transition Strategy	6
Retrofitting Sugarcane Operations and the Transition to Regenerative Agriculture	9
Mainframe Design	13
Methods to Reduce Overhead	13
Livestock and Holistic Management	14
Ancillary Agricultural Enterprises	16
Case Studies and Precedents	17
Diversified Operations	19
Biofuels	20
WATER AND SOIL	22
Water	22
Soil	24
Soil Building Strategies and Bioremediation	25
FARM ENTERPRISE OPPORTUNITIES	28
CONCLUSION	35
APPENDICES	36

PREFACE

In 2015, it became apparent that sugar cultivation on Maui was likely to end soon. It also appeared at that time that - other than raising cattle and leasing land to seed growers – landowner Alexander and Baldwin, Inc. (A&B) did not have much planned in the way of agriculture in central Maui that would increase the likelihood of long term success. The existing model of importing expensive chemical fertilizers and pesticides to grow a low-priced commodity crop for export across the Pacific Ocean to compete in world markets was clearly no longer viable. Diversified agriculture had been discussed as an alternative, but that had already been attempted for decades, and had yielded only mixed results throughout Hawai`i. Given that much of A&B's revenue is derived from real estate development, Maui Tomorrow's Board of Directors was concerned that urbanization of central Maui might soon follow. The Board discussed the importance of expressing a positive, sustainable vision, and considered hiring an agronomist to find crops that could be grown profitably here on Maui while minimizing the need for toxic chemical inputs. As a result of this inquiry, we discovered much more.

Properly designed and implemented, it turns out that this large, consolidated 35,000-acre block of central Maui farmland can be used to generate multiple income streams while growing food and fuel profitably for local consumption and value-added export. Regenerative agricultural methods can rebuild our soil, while using far less water. Agricultural jobs can be saved, along with Maui's open space and rural character. There is also an exciting possibility that we can create a new, regenerative agriculture education industry. Maui can share this important knowledge – which will help sequester carbon, and thus battle climate change - with people from around the world, and thus have an impact far beyond our shores.

The first printing of Mālama `Āina: A Conversation About Maui's Farming Future was released in March 2016. Simultaneously, it was released online at www.futureofmaui.org. As intended, it has generated a wide variety of comments, many of which are incorporated in this revised edition. We invite people to continue to share their mana`o, so that this living document can serve as a timely and valuable resource for prospective agricultural entrepreneurs, government decision makers, and the wider community.

As stated in this report, there needs to be a more detailed examination of the costs and methods necessary to shift agricultural operations on the former sugar lands toward profitable and sustainable regenerative agriculture. To address that need, Maui Tomorrow is in the process of preparing a follow up report that will include business plans and detailed data, both of which will be essential as we examine these issues in more depth.

We invite you to join us on this vital journey.

INTRODUCTION

Aloha`āina - love and respect the land, make it yours and claim stewardship for it

Mālama`āina - care for and nurture the land so it can give back all we need to sustain life for ourselves and our future generations

-Puanani Rogers, Ho`okipa Network

Beloved Maui is at a crossroads. The January 2016 announcement by Alexander and Baldwin (A&B) that Hawaiian Commercial & Sugar (HC&S) will be ending their 35,000 acre sugarcane operation has flung the door wide open to a much-needed conversation regarding what the future of farming on Maui can truly be. It's an opportunity to invite all the stakeholders into this discussion with the spirit of aloha, and draw on our collective mana`o to consider how we will plan ahead to mālama`āina - care for and nurture the land so it can sustain life for ourselves and future generations.

Maui now imports upwards of 90% of its food and energy, most of our building materials, and all of our textiles - a precarious reality for a remote island. We need many more living wage jobs, ample affordable housing, abundant and affordable local food, and clean water to provide for our 145,000 citizens and 2.6 million tourists annually.

Facing the future, we have choices - what will farming look like on Maui from this moment forward? This question is intimately tied to the wai, our precious water resources, and any answers must offer solutions that care for and restore watersheds.



Laguna Blanca, Argentina. Twelve years after transitioning to diversified regenerative farming.
http://www.tompkinsconservation.org/farm_laguna_blanca.htm



MA'O Organic Farms Hawai'i Investment Ready Program
<http://social-impact.org/regional-programs/investment-ready-hawaii>

The closure of the HC&S sugarcane enterprise is an opening to the next generation of diversified farm businesses. 35,000 acres of sugarcane plantation land farmed by HC&S are in question, of which 27,000 acres are designated Important Agricultural Land, and


receive tax and water benefits intended to help keep large tracts of contiguous farmland intact, and make farming more affordable. **Maui's farming future is tied to this land.**

What kinds of agriculture will benefit Maui's people moving forward? For 150 years Maui agriculture has been large-scale, mono-crop, chemical dependent, and export oriented. Can a new farming model bring both economic and biological benefits? The sugarcane era is officially ending this year - citizens of Maui are concerned about the loss of jobs for so many families, and want to see Maui's agricultural legacy continue.

This report is the start of a community conversation - bringing our diverse people and businesses together to find long-term solutions that are pono. With that in mind, there are many stories that need to be told, and discussions that need to take place. This report offers a window into abundant, resilient **regenerative agriculture** - a way forward that prioritizes food crops, livestock, diverse and profitable enterprises, and can build a whole farming economy that is just and environmentally sound.

The cessation of sugar cultivation raises important questions:

- What kinds of crops will grow here well and profitably?
- Should we prioritize food crops over commodity crops, and why? Can we have both?
- How do we remediate contaminated soils and aquifers?



Portuguese sugar workers - 6,000 in 1913
https://en.wikipedia.org/wiki/Portuguese_immigration_to_

- Will A&B sell the 27,000 acres now designated as Important Agricultural Lands? If so, at what price?
- What would be the implications of alternative ownership scenarios, such as non-profit land trusts, state ownership, or some combination thereof?

This report is a snapshot of what is possible, profitable, and pono. Looking at case-studies from similar climates, there are compelling precedents and sound science the world over that support making the transition from conventional to regenerative agriculture. Each specific area will need further research and detail before proceeding.



Japanese sugarcane workers 1885
<https://s-media-cache-ak0.pinimg.com/736x/84/42/80/844280dac7ef0154bde250816a634ba2.jpg>

Cultivating beneficial relationships between stakeholders is the foundation for the success of any project. We look forward to hearing your stories, addressing your questions and concerns, and incorporating new ideas.

A Brief Overview of Maui's "Central Valley" and Sugarcane

The central valley of Maui, once a native dryland forest, is now a wind-swept arid landscape with intermittent streams that were once perennial. Streams such as Pulehunui, Kaliainui, Kailua, Pohakukea, Waikapu and others were free flowing until the advent of large scale grazing and logging on upper slopes. This area previously provided nēnē habitat, as evidenced by the name Pu`unēnē; nēnē are now returning to areas such as Waikapu. Most native Hawaiians traditionally lived around flowing water, where the sophisticated ahupua`a-based farming and aquaculture systems were developed. Master farmers and master fishermen grew and raised plenty of food to feed upwards of a million people across the islands.

Before the sugarcane era, the central valley was sparsely populated, and was not intensively farmed. The birth of the sugarcane industry changed the face of the central valley and all of Maui forever. Water from the northeast coast of Maui was diverted to irrigate the sugar crop, impoverishing stream ecosystems and negatively impacting communities by restricting their access to water, and hence their ability to grow traditional foods. The once-parched land flourished with this abundant water, and the sugarcane industry has dominated the local economy for almost one hundred and fifty years. Waves of immigrant workers came to seek their fortunes, worked the cane fields, and settled down to raise families.


- Can we have diverse livestock operations? What would that look like?
- How many more jobs can we create with diversified regenerative agriculture?
- Does Maui have enough farmers for all these businesses?
- How can we assure long-term access to land for farmers?
- Can affordable housing be integrated into the design?
- Does regenerative agriculture use less water, and can some of that water be returned to East Maui watersheds?
- Can we and should we expand agritourism?
- What type of infrastructure do we have for "food hubs" - processing, creating value-added products, improving local distribution, and providing education for farmers?
- How can we bring all the stakeholders to the table, including A&B, in a transparent and meaningful way?

INTRODUCTION

The sugarcane industry the world over has shifted to chemical farming and systematic mechanization of jobs. On Maui, as in other places, not only has the number of industry jobs decreased steadily, but the use of more and different agricultural chemicals has contaminated the soil and the main Pa'ia aquifer, which is now polluted to the point where the water is considered unsuitable for drinking.

Pesticides, herbicides, fungicides, synthetic fertilizers, and ripening agents such as glyphosate have been applied to the soil for years, with very little independent research available to evaluate any deleterious effects on farm workers, neighbors, the groundwater, the ocean, and endemic wildlife. There are no legal obligations for HC&S to remediate the soils or the aquifers.

This style of agriculture, a monoculture crop with substantial chemical inputs, has been found to be a significant contributor to global warming - rather than sequestering carbon in healthy soil, repeated tillage and the application of nitrogen-based synthetic fertilizers release large amounts of CO₂ into the atmosphere. Records show significant amounts of pesticides have been applied to central Maui soils, and these chemicals or their derivatives have now shown up in the soil. Mechanization and global transportation of commodity crops only adds more CO₂ to the atmosphere, an externality that industrial agriculture has passed on to the citizens of the world.



Herb Kane's painting of Ka'anapali pre-contact
<http://herbkanehawaii.com>



Ka'anapali today
www.islandbreath.blogspot.com

HC&S has been unable to compete on the world market in recent years, and after reporting \$30 million in losses in 2015, they recently announced they will be shuttering the last remaining plantation, not only on Maui, but in the State of Hawai'i. 675 planters and skilled mill workers will be laid off this year.

Regenerative Agriculture

Regenerative agriculture integrates specific farm management practices and site design to build ecological and economic resiliency at every opportunity. Regenerative agricultural practices significantly improve upon the USDA National Organic Standards, and are applicable at any scale.


Systematically increasing soil health is the foundation of regenerative agriculture. Building healthy soil improves crop yields and resistance to pests, and makes crops more profitable. It decreases the need for external inputs such as fertilizers and pesticides, improves the water holding capacity by adding organic matter to the soil, and dramatically increases carbon sequestration as a byproduct of the above functions. Regenerative agricultural systems are based on perennial crops with sustained yields using resources generated on-site, as compared to annual agriculture which is often dependent on tillage and external inputs.

Regenerative Agriculture Characteristics:

- Improves water and mineral cycles on agricultural lands through contour farming and soil conservation methods
- Increases effective precipitation (the percentage of rainfall which becomes available to plants and crops) by improving soil structure and proper grading of land
- Reduces water use by selecting crops that are adapted to the local climate
- Preserves and creates soil through sound soil management practices
- Reduces or eliminates soil degradation and erosion caused by tillage through the use of perennial crops
- Sequesters carbon in the soil through organic production methods, thus counteracting climate change
- Is based on increasing diversity of both agricultural crops and native species to achieve Integrated Pest Management (organic farming techniques for controlling pest predation)
- Decreases reliance on agricultural chemicals such as fertilizers and biocides
- Integrates humanely raised livestock into crop production
- Improves economic resiliency of farming operations through diversified production
- Prioritizes local distribution and value-added products to improve profit margins
- Produces nutrient-dense food products that are healthier
- Improves natural capital and ecosystem services on agricultural lands
- Uses socially just business models like cooperatives, profit sharing, and nested enterprises
- Demands more skilled labor through the diversification of farming enterprises
- Embeds a full complement of necessary

Copyrighted Material
**THE
 CARBON FARMING
 SOLUTION**

A Global Toolkit of Perennial Crops and Regenerative Agriculture
 Practices for Climate Change Mitigation and Food Security



Cover of The Carbon Farming Solution, by Eric Toensmeier

- agricultural and business skills in the community for generations
- Generates a significant economic multiplier effect in the community, creating real wealth well beyond the agricultural businesses

Regenerative agriculture is based on sound design of the **mainframe** of farming operations. **This means that the site is designed to reduce or completely eliminate soil erosion from wind and rain through contour farming, proper grading of roads, and covering the soil with plants and/or mulches - the goal is to build soil, not lose it.** Infrastructural elements such as processing centers and agricultural facilities are

located near each other to increase efficiency of farming operations, and systems are integrated so that there is synergy and economy of management.

The reduction of water use through the selection of appropriate crops and the increase of effective precipitation through various soil management practices is another very important facet of regenerative agricultural systems. There is a global water shortage, but this shortage is really an issue of management of our water resources as opposed to a lack of water in the environment. Common agricultural practices like tillage, which leaves soil bare, reduce soil organic matter and therefore the capacity of soil to hold and store water. One percent of organic matter added to the soil enables it to hold 8 times more water, allowing the soil to act as a sponge. Shifting to ecologically sound management practices can conservatively increase water-holding capacity of soils by up to 15% or more.



Side by side comparison of conventional grazing (left) and regenerative agriculture (right) grazing lands. <http://www.abc.net.au/news/2015-09-16/6780458>

REGENERATIVE AGRICULTURE

Regenerative agriculture is rooted in ancient techniques and wedded to the best of modern agro-ecological technologies. Techniques such as composting and cover cropping for soil building are enhanced with our modern understanding of soil microbiology. Sophisticated rotational grazing of livestock to improve agricultural lands is now more effective with new and evolving practices such as Holistic Management and Management Intensive Grazing.


The business models and corollary social systems - the "invisible structures" - are the foundation for any successful agricultural operation that has the interests of its local community at heart. Regenerative agriculture systematically improves on select business models to build successful and profitable farming enterprises, taking into account the short and long-term health effects on land, water, agricultural workers and surrounding communities.

Maui residents have access to more information than ever before through the worldwide web. The ability to share successes and failures across the globe in real time is perhaps the most important advancement of our culture. This allows the future of Maui agriculture to incorporate improved agricultural systems from other similar climate zones as appropriate to our local culture, thus saving time and investment. A speedy transition to a diversified, sustainable farming model for the lands formerly in sugar cultivation will benefit local workers, enable local food production and enhance Maui's overall economy.



Regenerative Agriculture results:
Loess Plateau, China 1995 (top) & 2005 (bottom)


<http://permaculturenews.org/2012/06/28/hope-for-a-new-era-before-after-examples-of-permaculture-earth-restoration-solving-our-problems-from-the-ground-up/>



Climate Change and Regenerative Agriculture

Climate change and agriculture are strange bedfellows. Agriculture is responsible for 50% of global greenhouse gas emissions, and is one of the main contributors to climate change. Conversely, climate change negatively affects agriculture. Droughts, floods, and heat waves all have profound impacts on our food production systems leading to crop losses and food supply shortages. Resource scarcity is a leading cause of conflicts globally, as the surge of climate and economic refugees is being linked directly to competition for these resources.

Yet the problem reveals the solution. As agriculture is a major contributor to climate change, shifting our production methods to regenerative agriculture as described above can also reverse this trend. By some estimates, if all the arable land in the world increased the soil organic matter by 2%, we could reduce atmospheric carbon to pre-industrial levels.



<http://ecowatch.com/2015/01/06/regenerative-organic-agriculture/>

To quote a white paper by The Rodale Institute ¹:

- If management of all current cropland shifted to reflect the regenerative model, we could potentially sequester more than 40% of annual carbon emissions
- If all global pasture was managed using a regenerative model, 71% of carbon emissions could be sequestered
- Even if modest assumptions about soil's carbon sequestration potential are made, regenerative agriculture can easily keep annual emissions to within a desirable range


Andre Leu, President of the International Federation of Organic Agricultural movements (IFOAM), provides a thorough review on carbon sequestration in organic soils from diverse sources and ecosystems. These findings are corroborated by international agronomists and climate scientists; quickly converting farms to healthy soils is now a leading topic at climate change solutions forums.

(Endnotes)

1) <http://rodaleinstitute.org/assets/>



Carbon Farming Educational Series, Permaculture and Holistic Management for Carbon Negative Agriculture



<http://beyondpesticides.org/dailynewsblog/2016/02/organic-agriculture-essential-to-a-sustainable-future/>

Regenerative Agricultural Land Use Potential and Transition Strategy

From the 15th century on, Native Hawaiian chiefs governed and managed land based upon a system involving mauka-makai land divisions known as ahupua`a. This sophisticated agricultural and ecological management system included complete watersheds from mountain peak (mauka) to reef (makai), with several distinct sub-systems for food production, aquaculture, and communal land use.¹


The ahupua`a-based management system is a regenerative system; the local environment is continuously improved, and water cycles are maintained, all the while producing abundant foods. Food production and ecosystem functionality go hand in hand, and the system is resilient in the face of drought, flood, fire, and

hurricane. Skills, knowledge, and cultural identity grow from generation to generation. Today, upper portions of the watersheds are managed through collaborative watershed partnerships, while most farming land is outside that management system. The Maui Island Plan includes a policy that requires application of ahupua`a-based management to all lands within the watersheds:

All present and future watershed management plans shall incorporate concepts of ahupua`a management based on the interconnectedness of upland and coastal ecosystems/species.

Maui farmers have the opportunity to incorporate the principles of the ahupua`a system into overall land management to meet this goal and reconnect to our agricultural legacy.

REGENERATIVE AGRICULTURE



Ahupua'a Agricultural Association of Hawaii (AAA)
Kauai University Web log

Retrofitting the current mono-crop system to one that borrows from the more crop-diverse ahupua'a-based management system is an excellent way forward. Agricultural land uses would change according to elevation and soil type, exposure, and proximity to infrastructure. Maui has opportunities to implement solutions that require the least change for the greatest effect; this may begin with adding tree crops and livestock to the current sugarcane operation, as outlined below.

Despite the fact that Hawai'i's climate is ideal for the production of many types of crops, there is less than a ten-day supply of food on the islands at any given time.² The public is rightfully concerned about our food future, and people are demanding real solutions.

The rationales for changing the HC&S operation to regenerative agriculture are many:

- Increase the number of skilled jobs in the agricultural sector
- Satisfy the demand for locally produced food and renewable energy
- Create an economically resilient agricultural operation based on diversified products
- Recharge groundwater and restore hydrological cycles on the land
- Eliminate storm water discharge of agricultural chemicals and come into compliance with the Clean Water Act
- Eliminate air pollution from cane processing and burning, spray drift of agricultural chemicals, and airborne dust
- Address public outcry against industrial farming and Genetically Modified Organisms (GMOs)
- Restore water and stream flows to native habitat and farms in East Maui
- Restore native habitats and increase biodiversity on site
- Make it pono by providing access to land for farming through Cooperatives and provide farmworker housing for local workers
- Reduce / eliminate the use of chemical fertilizers and biocides, thus allowing cleansing / remediation of our soils, aquifers, and coastal ecosystems to begin

Regenerative agriculture pays dividends

Regenerative agricultural systems can be more profitable than conventional agriculture, offering better risk-reward scenarios for agri-business and farmers.

According to a recent report by Paul McMahon of SLM partners (an asset manager that acquires and manages rural land on behalf of institutional investors) there are a number of reasons why

these types of systems can deliver superior risk-adjusted returns:⁴


- **Comparable or better yields in most cases**
- **Lower operating costs because of less reliance on external inputs**
- **Enhanced natural capital, with the opportunity to increase asset values by regenerating degraded land**
- **Climatic resilience because healthy soils cope better with droughts and floods**
- **Positive environmental externalities and the chance to be paid for them, for example through carbon credits**
- **The ability to sell to higher value markets such as those for organic or grass-fed meats**
- **Higher profitability with less volatility**

Converting 35,000 acres of industrial sugarcane to diversified organic regenerative agriculture will take many years and the final outcome will likely be very different than what we now see as most practical. We must chart our course, and be prepared to adjust as the winds and tides dictate.


Designing the 'visible structures' - the plants and animals, windbreaks and terracing - is the easy part. Designing new 'invisible structures' - the contracts and agreements, business models, fair and inclusive governance, and land access - is much more challenging; however, if done properly, the result can be transparency, inclusivity, and shared prosperity.

This transition will require an adaptive management strategy. The system must be able to respond to cultural patterns, biological indicators, and economic pressures - it must be resilient. Nature is our model, as resilient systems have evolved over eons. By mimicking natural systems we create agricultural and cultural systems that are dynamic and respond to change and external pressures without wholesale collapse.


10 year regenerative agriculture transition to tree crops using sugarcane for fodder, swales on contour for water infiltration, and aquaculture



Year 1




Year 3



Year 10

Illustration by Silvia Yordanova



Keyline Design, photo by Mark Shepard, New Forest Farm

Retrofitting Sugarcane Operations and the Transition to Regenerative Agriculture

The most sensible way to switch the HC&S plantation to regenerative agriculture would be to retrofit the current sugarcane operation. A fresh look at the intrinsic characteristics and benefits of sugarcane will help us understand how management practices could be shifted slightly to take advantage of existing infrastructure and resources currently available.

While beyond the scope of this report, a detailed Business Plan could be prepared to examine what percentage of current HC&S field and processing employees would be needed in the early years of a regenerative agriculture program to harvest and process higher value-added cane products. Further, this Business Plan could review whether HC&S's current mill on Maui could be retrofitted, and at what cost, to process these new products.

The Business Plan would analyze whether the sale of additional sugar related products could generate enough funding over some time period to cover additional employees and infrastructure costs related to future regenerative agriculture operations.

Sugarcane, or *kō*, is a traditional Polynesian canoe crop that has been transplanted the world over because of its very desirable characteristics and adaptation to varied soils and regions.

Sugarcane is *the* champion crop with respect to carbon sequestration and soil building.⁵ It is one of the best tropical fodder crops for livestock, including cattle and swine. In its raw state, sugarcane juice is actually very nutritious. Raw cane juice is high in polyphenols, vitamins and minerals such as calcium, potassium, magnesium, manganese, and iron, along with a complete profile of essential amino-acids. It has even been found to lower cholesterol--both LDL and triglycerides, and is high in antioxidants.⁶

Sugarcane's outstanding characteristics are:

- Its perennial growth habit
- The quantity and nutritional quality of sugarcane increase with harvest interval, with optimum values being reached at a harvest interval of between 12 and 18 months. This is in marked contrast with almost all other tropical forage crops, which deteriorate in yield and quality as the interval between successive cuts is increased. For this reason, sugarcane has been called "ensilaje vivo", or living silage in many Central American countries
- The dry matter content of mature sugarcane averages 30 percent, which exceeds that of most other forage grasses (the average for Elephant and King grasses is closer to 17 percent). Thus harvest, transport and processing costs per unit dry matter are less for sugarcane than for most other forages
- There is a long tradition in sugarcane agronomy, especially in breeding, pest control and cultural practices. Admittedly this has been mainly

directed to enhanced production of sucrose rather than total sugars, which is the important criterion for animal feed. However, the implication of this practice in terms of the loss of potentially promising varieties is one of degree rather than direction, as there is a direct correlation between sucrose yield and feed value.

- Sugarcane is widely tolerant of soil and climatic characteristics. Maintaining a canopy of green leaves (or a mulch of dead ones) throughout the year helps to combat erosion, giving it a distinct advantage over competitive forage crops such as cassava and maize.⁷

Sugar production on Maui has centered around the crop as a source of sucrose: refined sugar; raw sugar and the value-added products of molasses and rum. There is considerable opportunity for additional value-added products from sugarcane processing. There is a market for boutique products such as organic raw sugar and juice. The antioxidant levels in sugarcane juice are thousands of times higher than the next best vegetable sources, and can sell for as much as \$60/kg.⁸ There are many uses for the bagasse including alcohol fuels, fiber for disposable plates, and other products. There are many cultivars of sugarcane possessing very different characteristics. Deeper analyses of the conditions unique to Maui can be done to indicate whether ***sugarcane may even be much more valuable converted to soil carbon and animal protein, as opposed to sucrose.*** It could be used as the fuel to feed the transition to regenerative agriculture.

Integrating Trees

The simple act of planting trees has so many beneficial effects. By integrating tree crops into the existing sugarcane fields as contour orchards and windbreaks, the current system may be retrofitted relatively easily and economically.³⁰

Trees serve multiple functions:

- Diversifying production
 - Fruits, nuts, and fungi
 - Medicinals and herbs
 - Spices and oils
 - Timber
 - Biofuels
- Providing fodder for livestock
- Decreasing overall water usage and improving water cycling
- Windbreaks for protection
- Moderate temperatures
- Sequester carbon while producing oxygen
- Biomass for soil remediation
- Increasing the overall diversity of the system

Contour Orchards

Contour orchards may be established by planting tree rows directly into the sugarcane fields on contour at 120 foot centers, creating alley cropping and silvopasture systems. Trees can be planted mechanically at the rate of several thousand per day per planting team. Additional herbaceous cover crops can be undersown to sugarcane fields to manage and prevent weeds, and improve the physical, chemical, and biological characteristics of the soil.

Integration of trees will reduce the total acreage of sugarcane significantly. **If the total acreage devoted to sugarcane was reduced by 30-50% by replacing with tree crops with less water demand, then the cumulative water use would conceivably be reduced by that fraction.** In addition, contour farming and rainwater harvesting earthworks combined with organic soil building strategies will offset supplemental irrigation even more.

Multi-Function Windbreaks

Winds have many harmful effects upon soil, plants, and animals, that are exacerbated as wind speeds increase. These impacts include drying of the soil and resultant loss of nutrients and biota, and increased plant and livestock stress resulting in reduced production and performance.

To minimize and even eliminate the impacts of wind stress, windbreaks of diverse tree, shrub, and deep-rooted herbaceous species are sited across the landscape. Generally, windbreaks are oriented perpendicular to the wind's dominant direction; however, while there is commonly a dominant wind direction, winds can and do occur from any direction at any time of the year. A 'net pattern' of windbreaks both down and across slope would significantly buffer winds in the central valley while supporting bio-remediation and providing important ecosystem services. In certain areas the windbreaks may be oriented on contour as part of Agroforestry, Holistic Range Management, and other production systems.

Windbreaks are a significant feature of the mainframe design for the central valley's regenerative agricultural system, providing many key services while increasing the aesthetic value of the landscape.



Multi-Function Windbreak

<https://www.agric.wa.gov.au/land-use/establishing-effective-windbreaks-swan-coastal-plain>




Illustration by Silvia Yordanova

Mainframe Design

The future of any successful regenerative agriculture venture hinges on excellent **Mainframe Design and Implementation**, as well as good management. The following design features must be prioritized as part of the transition strategy, as the success of the whole system is predicated on proper implementation of these elements.

In the case of the HC&S plantation, many existing elements may be retrofitted only slightly to maximize efficiency and economy. Below is a sequence of what that process may look like. Since the site is relatively dry, particular emphasis must be placed on harvesting the rain.

Conceptualized Implementation Sequence of Retrofitting Mainframe Design Elements:

1. Integrating tree crops into sugarcane alley cropping systems on contour while maintaining equal distance as much as possible to facilitate harvesting
2. As much as possible, shifting to contour farming for sugarcane fields for soil preservation
3. Planting multi-function windbreaks
4. Adjusting the shape and orientation of fields and grading the site to maximize rainwater harvesting, increase groundwater recharge, build soil, and eliminate erosion and storm-water runoff
5. Improving access, roads, ponds, and ditch systems to assist the above functions
6. Planting forestry blocks on slopes too steep to farm on contour
7. Implementing diverse soil remediation strategies
8. Restoring habitat to field borders, roads, ditches, and drainage gullies by planting native trees and shrubs
9. Researching and developing value-added products for local distribution and export

10. Creating cooperative business models to allow access to land for local farmers
11. Locating and building Farm Hubs; clustering facilities for processing, fertility management (composting and fertilizer production areas), plant nurseries, and agricultural sales facilities in order to increase efficiency of operations
12. Locating farm dwelling villages and affordable housing for farmers and farm labor
13. Integrating alternative energy production into operations, like solar and wind power, hydroelectric, biodiesel, and methane/biogas



Managed timber interplanted with commodity crops
<http://www.aftaweb.org/latest-newsletter/tempo-rate-agroforester/94-2006-vol-14/april-no-2/86-summary-of-the-silvoarable-agroforestry-for-europe-safe-project.html>

As the Mainframe elements are designed, planted, and built, a thorough study of appropriate crops should be conducted. These include:

- Annual crops - vegetables and row crops
- Herbaceous perennial crops - perennial vegetables and fodder crops
- Tree crops - fruits, nuts, timber, and fodder crops

- Cereals and grain crops - including pasture cropping⁹
- Livestock systems - ruminants, swine, fowl, invertebrates, and aquaculture
- Biofuels - both annual and perennial
- Textiles - including fiber and dye crops
- Specialty crops - including spices, medicinal herbs, cut flowers, cosmetic and essential oil crops

The selection criteria for crops may include:

- Low supplemental water needs
- Suitability to soils and climate
- Nutritive value
- Marketability and consumer demand
- Potential for job creation and value adding

Certain systems will work better together and should be appropriately linked. ***Isolating any one of these cropping systems is like taking one note out of a song - there is no harmony.***

Methods to Reduce Overhead

In addition to the production of crops, designing for energy independence of farming operations is important. There are many types of energy that farms rely on, from petrochemical to biological. Fuels and nutrients run the farm.

Production of Compost

The value of adding biologically active compost to the soil is measured in the reduction of operational expenses for materials such as fertilizers and biocides, as well as electricity and fuels costs for pumping irrigation water. As a plant relies on the macro-nutrients to produce sturdy roots, stems, and fruits, it also relies on macro and micro-nutrients to build up immunity to pathogens.

The production of compost is important as a source

of humus and for the recycling of agricultural wastes. Compost is actually the final stage of the energy cascade of agricultural products. Below is a conceptual sequence of the “energy cascade” of an agricultural product in regenerative agriculture (in this case, fruit), from source to sink;

Agricultural Product Energy Cascade:


- Agricultural product: fruit
- Process: pulp for juice product
- Mash from juicing (adding value to processing “waste” feeds livestock such as for swine and layer chickens)
- Manure and feed scraps are used in a variety of composting operations
 - Anaerobic composting/methanogenesis generates biogas to run boilers or produce electricity
 - Thermophillic compost can generate hot water
 - Vermicompost (worm compost)
- End uses for compost
 - Apply to fields
 - compost tea injection
 - soil medium for nursery plants

Biogas: Methane Production

George Chan of ZERI (Zero Emissions Research and Initiative) developed sophisticated systems for maximizing returns and reducing overhead for agricultural operations by integrating anaerobic methane digesters to process agricultural wastes. The production of biogas for power produces sludge that can be used to produce algae for animal feed and feedstock for vermi-compost.^{10, 11}

Vermicompost

Vermicompost is an excellent solution to the accumulation of organic wastes on farms. It produces high-quality compost using worms with the added benefit of reproducing the worms



Pigs grazing in the understory
<http://silvopasture.ning.com/photo/pigs-at-hopkins-woods-low-res?context=user>



Holistic management vs. desertification in South Africa
<http://sheldonfrith.com/2015/11/23/the-future-of-agriculture-is-regenerative/>

themselves, a high protein animal feed for chickens, fish, and pigs. The value and potential for worm composting on Maui is reported but certainly understated.¹² Extensive research has been done in Mexico using worms to compost “cachaza”, (the residue of sugarcane juice filtration)¹³; this would be a natural choice for diverse fertility and soil-building strategies on Maui.

Pruning as a Management Practice

Pruning and coppicing are important management practices to increase soil organic matter and manage shade in agroforestry systems. Plants are specifically cultivated for biomass to provide feedstock for livestock and mulch. There is further opportunity to integrate biomass energy (syngas) from coppice wood.

Fodder trees can be mechanically pruned and the forage dropped in the field, and livestock allowed to graze on the wilted foliage, leaving manure residue to enrich the soil. Better yet, the animals can browse shrubs for late season stockpiled fodder.

Integrated Pest Management

The species diversity of regenerative agricultural systems provides a built-in Integrated Pest Management (IPM) function, further reducing reliance on agricultural chemicals. Insects and birds reduce pest populations, and maybe take a little fruit or nectar for their services. Yield is not only measured in production volume, but reduced expenses of operations.

Mechanization

Because of the scale of this operation, some degree of mechanization will be necessary as a transitional strategy and long-term reality. Therefore the layout of the plantings should reflect this, and have appropriate access for harvesting and pruning equipment.

Livestock and Holistic Management

Maui’s cattle industry dates back to 1793. The Maui Cattle Company, six independent family-owned businesses with over 60,000 acres of prime grazing land, states as their mission “the re-ignition of an agricultural lifestyle through the establishment of a sustainable ranching industry”.

REGENERATIVE AGRICULTURE

The demand for local beef far exceeds the island's production capacity. Most meats are currently exported off-island, and even if all the meat raised here were consumed locally it would only supply 20 – 25% of the Maui market.

Livestock production doesn't just feed people, it creates a wealth of skilled jobs in and around the industry - diversified production of multiple species, research, industry support, processing, value-added products, distribution and marketing, pasture consulting, business management, and more. Prioritizing meats for local markets requires a local slaughterhouse. Expanding and improving the livestock industry into Maui's central valley will require further research and analysis.

Holistic Management

Holistic Management is a systems approach to decision making that includes land planning, grazing planning, financial planning, and biological monitoring as they all relate to one another in the context of an agricultural operation. Holistic Planned Grazing is one aspect of the Holistic Management process, and is a revolutionary livestock management practice that mimics natural herbivore behavior with domesticated livestock. This method has been proven to be one of the best and most expedient ways to repair damaged ecosystems and reverse desertification.

Holistic Planned Grazing uses cattle and other ruminants that are moved frequently so their impact does not harm the land, but rather benefits it. Much like a herd of wild herbivores responding to predator pressure, herds are constantly on the move, only grazing the tops of plants. Pasture is allowed to rest so that overgrazing does not occur, and plants are allowed to recover and release carbohydrates into the soil, feeding soil microbes. With proper management, weedy species are

Copyright 2016 by Maui Tomorrow Foundation, Inc.



Sunflowers and lavender

<http://imgc.allpostersimages.com/images/P-473-488-90/26/2659/D8DUD00Z/posters/stevevidler-sunflower-fields-provence-france.jpg>



Harvesting sunflowers

<https://i.ytimg.com/vi/YpzTinl65iM/maxresdefault.jpg>

replaced by more beneficial pasture species, and soil organic matter builds rapidly.

Livestock are bunched together in small groups called mobs, and their grazing patterns shift in response to increased competition for pasture; basically they eat weeds and other plants they might pass up if they were in a set stocked system

Fire & Drought Resistant Farms & Landscapes

with Darren J. Doherty

September 12th -14th, 2014 9 am - 5 pm daily
@ The Ojai Foundation Ojai, California



REGRIAN PLATFORM

1. Climate
2. Geography
3. Water
4. Access
5. Forestry
6. Buildings
7. Fencing
8. Soils
9. Marketing
10. Energy

with little competition for feed. Stocking rates can be doubled or tripled, with up to one million pounds of livestock per acre possible, in rapid rotation. Livestock may be moved daily and even hourly, depending on the forage quality and quantity, and paddocks may be rested for up to one year depending on conditions and management, to allow recovery.

Multiple species can be integrated into one management system for diversified production. For example, cattle can be followed by pigs, which can be followed by chickens. The cattle browse grasses and forbs, pigs root for insects and tubers, and chickens eat fly larvae in the manure of the previous species, reducing the vectors for disease. These animals all work in synergy and complement each other, much like the wild populations of animals that are diverse and occupy various niches. This is an example of stacking systems in time. Economic opportunities and a cascade of skilled jobs are the result of diversifying operations.

Ancillary Agricultural Enterprises

There is considerable opportunity for embedding ancillary agricultural enterprises within the Holistic Management system early in the transition, such as:

- Breeding and sales of organic open-pollinated heirloom seeds
- Nursery plants - including native, edible, and ornamental plants
- Organic fertilizer production - including compost and microbial inoculants
- Value added products - including preserves and fermented foods
- Construction products - including timber and bamboo
- Agri-tourism - including farm tours, fresh farm lunches and dinners, direct sales of value-added farm products, and educational workshops
- Regenerative agriculture training and implementation programs


Plant Breeding and Seed Production

Seed saving, genetic selection, and animal breeding techniques have traditionally been passed down through intact farming lineages. Over time “landrace” cultivars and breeds develop that are hardy, resilient, and perfectly adapted to local conditions.

Modern plant breeding has been relegated to seed companies and research institutions. Hybrids and genetically modified organisms are usually patented and their genetic material owned by the companies who produce them. It is technically illegal to save these seeds, or propagate them without paying royalties. Farmers now need to buy their seed every year, and sadly, saving seed has become an antiquated, and sometimes illegal practice.



Silvopasture grazing with sheep
<https://www.flickr.com/photos/baalands/2216308884/>



Cattle grazing on leucaena, a perennial nitrogen-fixing fodder crop
<http://portfolios.pratt.edu/gallery/3880677/Rethinking-food-production-in-the-tropics-research>

Plants reproduced through biotech engineering, hybridization, or clonal propagation are genetically identical. Mono-crop plantings of clones are more susceptible to pests and diseases, as pathogens need only crack the code of one genetic makeup to infect the whole field. In nature all plants are sexually propagated through pollination, therefore their genetic makeup is varied. This means pests and pathogens must crack the code of many genetic expressions, conferring natural disease resistance to native populations.³¹

Globally there is an enormous need for open-pollinated seed varieties of both annual and perennial crops. While many sources for open-pollinated and heirloom annual vegetable seeds exist, there are very few seed companies that develop fruit and nut tree seed with stabilized traits – qualities that come true to type when planted. Changes in climate are creating the need for new provenances that are adapted to increasingly variable climatic conditions.

Hawai'i can grow certain seed crops that yield three harvests each year. GMO seed corporations have successfully and profitably capitalized on the unique attributes of Hawai'i's favorable climate, year-round sun, and available water. HC&S lands have locations suited to lease to organic open-pollinated seed companies, if they were recruited to relocate here to Maui. This nationally expanding agriculture sector provides good, highly skilled jobs. Local workers could be trained in these breeding and harvesting protocols.

Developing a breeding program for fruit and nut trees, much like was done by the legendary plant breeder Luther Burbank - who developed the russet potato and other improved fruit tree varieties - is much needed in the world, especially Hawai'i. The islands are well suited to testing new and improved seed varieties.


REGENERATIVE AGRICULTURE

Hawai'i needs to develop new landrace seeds that have desirable characteristics, are adapted to the local climate, that are true to type, and make this seed available to farmers and gardeners to propagate at will. This is the real foundation of food sovereignty and security.

The University of Hawai'i College of Tropical Agriculture and Human Resources (CTAHR) recently began work on The Hawai'i Public Seed Initiative, which emphasizes the importance and value of local seed systems. Working with the Kohala Center, they have gathered baseline data, taught seed-saving workshops, created seed networks on all the islands, and are establishing statewide and regional variety testing trials.³²

Educational Opportunities

One important limiting factor inhibiting the proliferation of regenerative agriculture in Hawai'i is the lack of trained farmers. This is certainly not exclusive to Hawai'i; it is a global issue. Again, the problem reveals the solution. The sugarcane



Regenerative agriculture practices build deep rich soil (left), compared side-by-side with conventional grazing soil (right) at Winona Farm in Australia.

fields can be used as a living classroom to train farmers, and the initial curriculum would parallel the transition phase plan.

The potential for creating a world-class educational curriculum on this site should not be understated. The demand for this type of learning experience is as great as the demand for the food these systems produce. There is a serious lack of hands-on training for aspiring farmers, and for seasoned farmers wanting to move out of conventional farming. An ongoing educational series can be developed in partnership with the many existing Maui institutions that will attract professionals from around the world.

As a venue, Hawai'i and Maui are in high demand, and this project would expand on the ever-growing sector of agricultural and educational tourism. Recruiting the world's leading regenerative designers as instructors and hands-on trainers could put Maui on the map for excellence in regenerative agricultural education.


Examples of workshops and regenerative agriculture trainings include:

- Stakeholder process and community engagement
- Whole Farm Design
- Regenerative Agriculture Design and Management
- Holistic Management and Livestock Operations
- Tropical Agroforestry Design and Management
- Seed Breeding and Plant Propagation
- Earthworks Design and Implementation

There is significant income potential for educational offerings. Students regularly pay \$200/day for specialized trainings from 4 days – 2 weeks, and attend longer in-depth courses for certification-level trainings.

Case Studies and Precedents

There are a growing number of successful



<http://www.glenwoodgarden.com/wp-content/uploads/2014/06/SAGE-Food-farms-future-image-logo.jpg>

regenerative farms sprouting up in Hawai'i and across the world. It is important to look to existing success stories to glean information applicable to central Maui lands. The challenges that these enterprises have faced during their development provide the most valuable information we can seek in order to avoid similar mistakes and move forward swiftly. The following case studies were chosen as they are particularly relevant to Maui, showcasing projects from similar climates, farm businesses that integrate sugarcane, profitable livestock operations that regenerate farmlands, and demonstrate how farmers have diversified income from multiple products within a mainframe design.

Organic Sugarcane Production

Sugarcane Best Cultivation Practices, Brazil

Brazil has done pioneering work in developing and studying no-till management of sugarcane production, particularly for ethanol production. In a 15 year study, they found that no-till sugarcane operations provide many benefits to both the farms, the soils, and the environment. Through the combination of no-burn and mechanical harvesting while leaving the dry matter on the field, no-till sugarcane operations eliminate the air pollution caused by burning and at the same time create a net increase in soil carbon stocks of 9.7

Mg/hectare. Rather than coming at the expense of yields, no-till sugarcane operations yield 10 tons more per hectare than conventional plantings. Fertility is cycled by returning the sugarcane byproducts from the processing mills to the field and growing nitrogen-fixing cover crops such as sunn hemp (*Crotalaria juncea*) as a green manure. Advantages to no-till management of sugarcane include reduced costs in soil preparation, increased organic matter in the soil, improved fertility, reduced fertilizer applications, reduced compaction, reduced erosion, and lower emissions of greenhouse gases.^{14, 15}

No-till Systems

The Brazilian sugarcane industry employs modern agronomic management practices to enhance productivity and protect the environment. Key features of Brazil’s sustainable approach to cultivation and processing include:

Low Soil Erosion

Brazilian sugarcane fields have relatively low levels of soil loss, due in part to the semi-perennial nature of sugarcane. The same plant will grow back many times after it is cut, and its cane juice is extracted. In fact, sugarcane is typically only replanted every six or seven years. The Brazilian industry also emphasizes farming techniques that preserve soil stability while yielding approximately 34 tons of sugarcane per acre, as compared to an average of 11.5 tons per acre at HC&S.³⁰

Strategies include:

- No-till production systems
- Crop rotation with soybeans or peanuts
- Green fertilization by planting cover crops such as *Crotalaria juncea* or using leftover sugarcane straw after mechanized harvesting as ground cover



Silvopasture at Pongamia, Australia

Thanks to these responsible agricultural practices, soil erosion in sugarcane fields is minimal when compared to many other crops such as rice and soybeans (for more information, read the “Environmental sustainability of sugarcane ethanol in Brazil” study). In some regions of the country, sugarcane has been produced on the same soil for more than 200 years with continuous yield and soil carbon increases.¹⁶

Organic Sugarcane Production, Ingenio El Mante, México

Nacho Simon recognized that the “cachaza”, a by-product of sugarcane processing, was a problem for the majority of operations, so he decided to transform it into compost and return it to the fields. The addition of micro-organisms to the soil was very beneficial, as his soils had been depleted from repeated harvests.

The organic matter is the food and home of the life in the soil, and maintains the equilibrium and also transforms the dry waste in the fields to nutrients. The organic matter also retains humidity, reduces erosion, and improves the structure and texture of the soil, allowing roots to grow rapidly. The results of applying compost to the fields were a 100% increase in production and also the reduction of operating costs, allowing sugarcane operations to remain profitable.¹⁷

Cattle: Holistic Management

Florida, USA: Jim Elizondo, RegenGraz

Jaime (Jim) Elizondo of RegenGraz manages Mashona Cattle in an intensive silvopasture system with *Leucaena leucophylla* in North-central Florida. He also integrates high density grazing on mixed cover crops grown as stockpiled forages, which can be rapidly implemented on conventionally managed farmland under annual cropping and tillage practices.

Understory forages he uses in his intensive silvopasture system include bahiagrass, bermuda grass, torpedograss and a wide variety of herbaceous legumes. He also uses free-choice mineral supplementation based on the work of Mark Bader to improve the overall balance of soil minerals and forage quality for his livestock. Through these integrated systems, Elizondo has achieved cost savings “from \$100-200 per cow per year compared to normal practice in the area, plus [he] carries double the normal stocking rate while improving the soil and respecting wildlife.”

Starting with low fertility soils, they integrated compost extract liquid fertilizers at one gallon per acre to stimulate the soil microbes for the first two years. Once 100% soil cover was achieved, they stopped applying compost extracts and relied upon crop and litter management to feed soil microbes. Pasture cropping was also practiced, planting into winter perennial pastures with a mixture of summer annuals including lab-lab bean, sunflower, cowpea, hairy indigo, soybean, sudangrass, millet, sunn hemp, and clover.

Cover crops are trampled and harvested as forage using ultra-high density strip grazing (~1,000,000 lb -cow/acre moved 4-6x per day during this treatment). Based on his current success, he

REGENERATIVE AGRICULTURE

plans to develop shelterbelts and fodder tree rows of mimosa and mulberry plantings.^{18, 19}

Commodity crops: Textiles and fiber

Winona NSW, Australia: Colin Seis

2,075 acres pasture cropping: cereals, sheep, native grass seed

One of the most powerful innovations in large scale regenerative agriculture is pasture cropping, a system pioneered by Colin Seis at his home and farm Winona in New South Wales, Australia. Colin has been practicing and refining the pasture cropping system on his 840 acre farm since 1992, when a wildfire wiped out his operation. By integrating native warm season perennial pastures with no-till cool season cereal grain production, Colin produces three 'stacked' enterprise product lines from each pasture: sheep wool and meat, cereal grain, and native grass seed. The balance between pasture grazing and cereal cropping is achieved through carefully timed grazing rotations. At the same time, he has decreased annual production costs by \$120,000 AUS and increased soil carbon by over 200% in 10 years. While he has decreased costs and improved soil fertility and water holding capacity, his wool quality has also increased. All this is done on 650 mm (26") of avg. rainfall per year. Colin now works to expand his system to other farms and reports "over 2,000 farmers pasture cropping" cereal crops into summer (C4) and winter (C3) perennial native grass in NSW, South Australia, Victoria Queensland, West Australia and Tasmania as well as other areas around the world.²⁰

Agroforestry, Contour Orchards, and Alleycropping

Agenda Gøtsch, Brazil: Ernst Gøtsch

Another example of regenerative agriculture

transforming a landscape is on a cacao plantation in Northeast Brazil, owned and farmed by Ernst Gøtsch using a form of innovative agroforestry. When Gøtsch purchased the 1,200 acres of unproductive land in 1985, the land, like much in the region, was degraded and dry, considered unsuitable for cacao production.

Once covered in Atlantic rainforest, decades of timber exploitation and cattle grazing had left the land barren and the wells had run dry. Gøtsch used a unique blend of soil recovery techniques that mimic the natural regeneration of forests and reawaken the biology of the soil; within five years, water was again flowing in the wells. Within 10 years, he was obtaining 4,500 pounds of cacao per acre--1,250 pounds more than average for his region.

When disease destroyed much of the neighboring cacao plantations, his trees were untouched. He was generating 2-3 cm annually of new topsoil. The Atlantic Rainforest resurfaced on his 1200 acres, bringing with it its flora and fauna. Nearly 900 acres of his farm are now a natural heritage reserve. After 20 years, 14 springs have reappeared on the farm.²¹

Diversified Operations

Finca Luena Nueva, Costa Rica

This research project is unique in that it combines all three types of land use management on the same tropical farm - reforestation, cattle/pasture management, and commercial crop production. Crops that are grown include: cassava, turmeric, ginger, taro, sweet potato, banana, plantain, corn, various beans, various leafy greens, and vegetable crops. Cattle are dual-purpose, bred for meat and dairy production. The reforested land was once in open cattle grazing, and now has been replanted to native humid tropical rainforest.²³

Laguna Blanca, Argentina: Tompkins Conservation

7,418 acres; acquired in 2007



Laguna Blanca Farm, Argentina, diversified agriculture with terracing on contour
http://www.tompkinsconservation.org/farm_laguna_blanca.htm

Project of Kris and Doug Tompkins, Dolores Peréa-Muñoz and Eduardo Chorén, Entre Ríos Province, Argentina

Laguna Blanca is in the midst of a dramatic transformation from industrial monoculture to organic polyculture. Comprising more than 7,000 acres at the confluence of the Feliciano and Parana rivers in northeastern Argentina's Entre Ríos Province, Laguna Blanca offers an opportunity to develop a model of diversified organic agriculture for the region.

When it was purchased in 2007, Laguna Blanca was in serious need of restoration: its infrastructure needed attention, and its soils were eroding away. To counter erosion, terraces were built to create level fields in which a variety of grains—including oats, flax, sorghum, barley, and wheat—are now grown. New orchards produce eleven fruit and nut species, including peaches, pears, olives, dates, hazelnuts, pecans, and almonds. Many aromatic and herbal species are being cultivated alongside a wide assortment of horticulture crops, many of which are perennial varieties requiring zero tilling. Sheep graze in restored native pastures, and hay is made to feed them through the winter from the grasses surrounding the orchards. The practicality

of this polyculture style of farming is becoming increasingly evident as interactions between diverse crops, healthy soils, and native wildlife are improving the farm's yields.²²

**Aquaculture/Aquaponics
Hawai'i, USA**

There are several examples of successful commercial aquaponics enterprises in Hawai'i, including Kunia Country Farms²⁴ on Oahu, and Living Aquaponics²⁵ on Hawai'i Island. Both of these enterprises are on the order of one-quarter acre of total land area. Living Aquaponics is generating between \$1,500-\$3,000 gross income per week (\$75,000-150,000/yr) with three people working 20 hrs/wk. They have been in business for five years and have managed to work through the challenges of disease and pest management in organic aquaponics with leafy greens and root crop production even during Hawai'i's warm and wet seasons.

Biofuels


Pongamia silvopasture, Australia

Pongamia pinnata is an exciting legume tree native to Australia which tolerates a variety of climate conditions including salty soil, and can be integrated into silvopasture systems for biofuels production. Trees in trial plots of ~300 acres are producing 16kg of seeds per tree and 'elite' cultivars have produced up to 100kg of seeds per tree annually. In addition to producing biodiesel, the seed cake left after oil extraction can be fed to livestock as a supplement to silvopasture forage and fodder, allowing fertility to be returned to the system.²⁶

Enterprise Systems

Gabe Brown's Nested Enterprise Model^{27,28}

Although from a different climate, this case study showcases a number of successful diversified farming income streams on one farm using sophisticated regenerative agriculture techniques. The cascade of products shown in his Nested Enterprise chart are directly applicable to Maui.



Gabe Brown's Nested Enterprise model

Gabe Brown is rapidly building soil on more than 5,400 acres by merging back-to-basics agrarian practices with innovative, science-based sustainable farming techniques on his diversified family ranch in North Dakota. Beyond converting all cropland to no-till, he constantly seeds with a cocktail of dozens of cover crops. Through Holistic Management, a diverse cropping strategy, rotational grazing and no-till practices the farm has benefited in terms of soil health, mineral and water cycles, greatly reduced inputs, excellent production and profit, and an improvement in quality of life for the farmers.

Gabe Brown has been practicing his form of integrated regenerative agriculture for over 15 years. For decades the cropland had been conventionally farmed with tillage and the use of synthetic fertilizers and herbicides. Tillage had lowered organic matter levels to less than two percent. In 1993 Brown

purchased a no-till drill and converted 100 percent of his cropland to no-till. Brown employs a diverse cropping strategy on his grain and cattle operation which includes over 25 different cash and cover crops, resulting in high yields and strong net profits. The Natural Resources Defense Council awarded one of its 2012 Growing Green Awards to Brown and says, "Gabe's trailblazing work has made him a leader in regenerative ranch management."

**Educational Agritourism
Rancho San Ricardo, México**

Using the magic of Keyline® planning and ingenuity, the Mashumus team has converted a mono-cropped sugarcane plantation in Mexico to one of the best examples of permaculture and organic agriculture in the world. With over 140 acres of sugarcane, the project was conceived by Pablo Ruiz Lavalle and Eugenio Gras of Mashumus to be one of the most

innovative agricultural education sites of our time. Roads, ditches, fruit orchards, and the entire infrastructure were redesigned to harvest water, capture carbon, improve fertility, and beautify the landscape at the same time. Here, local cane growers who manage the surrounding 32,000 acres are able to learn the techniques of Keyline®, the secrets of biofertilizers and microorganisms, natural building techniques, and a myriad of other eco-technologies.²⁹

Keyline planning is based on the natural topography of the land. It uses the form and shape of the land to determine the layout and position of farm dams, irrigation areas, roads, fences, farm buildings and tree lines. Keyline is an agricultural system in which great emphasis is placed on processes designed to increase substantially the fertility of soils. Emphasis is placed on the creation of a soil environment that rapidly accelerates soil biological activity, thus vastly increasing the total organic matter content within the soil. Keyline lay-outs of farm and grazing lands also incorporate designs permitting the storage of run-off water on the farm itself.

The central valley of Maui is a large enough area to influence the local climate. Powerful winds bring energy and nutrients that can be harvested or deflected, including moisture, micro-organisms, minerals, and pollutants. Trees can help to mitigate the effects of the wind on this vast tract of land, and benefit the operation and local environment in many ways.

Regenerating our agriculture offers solutions to many of the “problems” facing the world today- water and food shortages, soil loss, rising energy prices and climate change. It may seem that agriculture has nothing to do with these issues, but in fact it has everything to do with them, and can address them all as a win for everyone.

(Endnotes)

- 1) <http://www.hawaiihistory.org/index.cfm?fuseaction=ig.page&CategoryID=299>
- 2) <http://hawaiitribune-herald.com/sections/news/local-news/bills-promote-food-self-sufficiency.html>
- 4) <http://slmpartners.com/wp-content/uploads/2016/01/SLM-Partners-Investment-case-for-ecological-farming.pdf>
- 5) http://epubs.scu.edu.au/esm_pubs/782/
- 6) <http://www.welmor.org/antioxidant-activity-in-sugarcane-juice.pdf>
- 7) <http://www.fao.org/docrep/003/s8850e/S8850E05.htm>
- 8) http://www.esugartech.com/technology_pubs/Refined_sugar_and_Value_added_Products_from_cane_mills.htm
- 9) http://www.regional.org.au/au/asa/2012/pastures/8052_dollingp.htm
- 10) <http://www.zeri.org/ZERI/Pigs.html>
- 11) <http://www.zeri.org/index.html>
- 12) <http://www.ctahr.hawaii.edu/oc/freepubs/pdf/HG-46.pdf>
- 13) https://www.researchgate.net/publication/267353433_Vermicompost_Production_with_Waste_from_Sugarcane_Industry_Sacharum_officinarum_L
- 14) http://www.soybeansandcorn.com/news/Jun7_13-No-Till-Sugarcane-Planting-Reduces-Costs-Improves-Yields-in-Brazil
- 15) <http://sugarcane.org/resource-library/studies/Wageningen%20-%20Chapter%205.pdf>
- 16) <http://sugarcane.org/sustainability/best-practices>
- 17) <http://mashumus.com/index.php/proyectos>
- 18) <http://www.progressiveforage.com/forage-production/producer-features/florida-ranch-optimizes-use-of-stock-piled-forages>
- 19) <https://www.youtube.com/watch?v=2-M8J3fslmg> <http://www.livingwebfarms.org/regenerative-ranching-hdrg/4591004075>
- 20) <http://www.soilsforlife.org.au/cs-winona>
- 21) <http://agendagotsch.com/project/>
- 22) http://www.tompkinsconservation.org/farm_laguna_blanca.htm
- 23) <http://www.carbontosoil.com/en/farms.html>
- 24) <http://www.kuniacountryfarms.com/>
- 25) <http://www.livingaquaponics.com/>
- 26) <http://www.bioenergyplantations.com.au/update-on-roma-trial-plantation-progress/>
- 27) <https://greencoverseed.com/sites/default/files/attachments/Gabe%20Brown-Livestock%20Integration.pdf>
- 28) <http://www.acresusa.com/gabe-brown-large-scale-rapid-building-of-soil-with-cover-crops>
- 29) <http://mashumus.com/index.php/proyectos>
- 30) while yielding approximately 34 tons of sugarcane per acre, as compared to an average of 11.5 tons per acre at HC&S (http://hcsugar.com/wp-content/uploads/2013/02/hcs_factsheet_2013_130201PDF.pdf).

Water and Soil

Wai: water

Waiwai: true wealth

Mālama `āina: to care for and nurture the land so it can give back all we need to sustain life for ourselves and our future generations

The Constitution of Hawai`i states:

For the benefit of present and future generations, the State and its political subdivisions shall conserve and protect Hawai`i's natural beauty and all natural resources, including land, water, air, minerals and energy sources, and shall promote the development and utilization of these resources in a manner consistent with their conservation and in furtherance of the self-sufficiency of the State.

All public natural resources are held in trust by the State for the benefit of the people.

The State shall conserve and protect agricultural lands, promote diversified agriculture, increase agricultural self-sufficiency and assure the availability of agriculturally suitable lands.

Water

The majority of Hawai`i has a tropical climate and receives copious rainfall. However Maui's central valley is in a rain shadow, and receives between 15 and 60 inches of rain annually, so most of the 35,000 acres being farmed by HC&S are technically in a drylands/sub-humid microclimate. Sugarcane, one of the world's thirstiest crops, could not be grown here without abundant supplemental irrigation.

This irrigation water arrives at the property in a series of long canals that divert streamflow from





Illustration by Silvia Yordanova

Swales are long, level excavations which are constructed on contour across the landscape to slow the flow of water, store water in subsoils, and create fertile and diverse planting microclimates. They are not intended to encourage or allow water to flow but to simply hold the water by promoting infiltration into the soil. Swales will vary greatly in width and length depending on the dictates of the land and design parameters.

The soil is excavated on the contour and normally mounded on the downhill side on the swale. The swale system creates prolonged subsoil moisture, and provides excellent drainage for trees.

more than 100 streams and tributaries in East Maui, and four main streams in West Maui. On average 165 million gallons per day have been diverted from these sources. As a result, these streams no longer support the native habitat they once did, and the communities that live within



East Maui Irrigation ditch
Photo by Will Scullin

their watersheds no longer have access to this water for farming and other uses.

In 2004, over 400 MGD of water, fresh and brackish, is used for domestic, industrial, commercial or agricultural purposes. Only around one-eighth, or 45 MGD of that amount was used for domestic and commercial use. Less than one-tenth of Maui's water resources are actually under public control, although billions of gallons of water originate on public lands. **The vast majority of present use is for agricultural irrigation.**¹

Control of the water is serious business on Maui. It is incumbent upon the next generation of farmers in the central valley to strive for smart water use

WATER AND SOIL

and employ best practices that systematically reduce demands for agricultural irrigation, and that also bank water in the soil across the region. Restoring Maui watersheds must be a priority in any central valley farm design. Regenerative agriculture addresses this issue convincingly.

Strategies of Regenerative Agriculture to Improve Hydrological Cycling:²

- Capturing and storing rainwater through
 - Terracing
 - Building swales on contour
 - Keyline plowing
 - Ponds
- Building healthy soil to improve water holding capacity using
 - compost
 - mulch
 - cover crops
- Planting windbreaks to decrease evapotranspiration and harvest atmospheric moisture
- Using rotational grazing to improve soils
- Planting climate adapted crops that require minimal irrigation
- Using conservation tillage methods
- Using efficient irrigation methods
- Planting perennial crops requiring no tillage

In a 30-year farm systems trial, the Rodale Institute found that corn grown in organic fields had 30 percent greater yields than conventional fields in years of drought. Healthy soil that is rich in organic matter and microbial life serves as a sponge that delivers moisture to plants. **The trial also found that organic fields can recharge groundwater supplies up to 20 percent.**³

A study released by Cornell University Professor

Copyright 2016 by Maui Tomorrow Foundation, Inc.



Coffee under nitrogen fixing leucaena overstory:
<http://afrique-orientale-australe.cirad.fr/en/research-in-partnership/ongoing-projects/ecological-intensification/safse>



Organic citrus orchard with herbaceous understory
<http://www.goldengroveorchard.com.au>

David Pimentel in 2005 reported that organic farming produces the same corn and soybean yields as conventional farming and uses 30 percent less energy and less water. Moreover, because organic farming systems do not use pesticides, they also yield healthier produce and do not contribute to groundwater pollution.⁴

Water Demand of Various Commodity Crops

Water demand varies greatly depending on climate, soils, cultivation practices, and species or cultivar selection. A brief look at alternative crops and livestock management shows significant potential reductions in total water use for the HC&S property.

The data below is derived from conventional systems and does not represent efficiencies possible in regenerative agricultural systems. However, according to the NRCS, a 1% increase in organic matter (carbon) in the top 6" of soil increases its water holding capacity by approximately 27, 000 gallons per acre.⁵ **It is likely that water use would be 10-50% less than these numbers, if the recommended regenerative methods were to be embraced and implemented.**

Sugarcane: HC&S current water use has averaged 165 million gallons per day (mgd) on roughly 35,000 acres. Irrigation requirements vary across the central valley⁶:

- East Maui: 4,844 gallons/acre/day
- Waihe`e-Hopoi: 5,958 gallons/acre/day
- Iao-Waikapu: 5,408 gallons/acre/day

Carob: 0.5-1 acre feet/acre/year

- Low end: ~500 gallons/acre/day
- High end: ~900 gallons/acre/day

Avocado and citrus average: 2-4 acre feet/acre/year

- Low end: ~1,700 gallons/acre/day
- High end: ~3,500 gallons/acre/day

Macadamia = 3-4 acre feet/acre/year

- Low end: ~2,700 gallons/acre/day
- High end: ~3,500 gallons/acre/day

Mango: 5-9 acre feet/acre/year

- Low end: ~4,400 gallons/acre/day
- High end: ~8,000 gallons/acre/day

Sunflower:

- Could be rain fed = no irrigation = 0 gallons/acre/day

Dryland Kalo:

- Planted at the beginning of the rainy season, dryland kalo is rain fed in areas with 6-9 months of rain (the time required for the crop to mature). Supplemental drip irrigation would be required in the central valley

Cattle:

- One cow can drink up to 25 gallons/day
- If stocking rates are 2 head per acre, that is 50 gallons/acre/day for drinking
- Pasture is normally rainfed and requires no supplemental irrigation
- No hay will need to be fed in lean times if trees and shrubs are used as supplement to grass
- The only water needs of holistically managed livestock systems are for the stock themselves
- Compaction of soils through conventional farming activities and the removal of trees and organic matter reduces effective precipitation.

Soils that are bare and hard will not absorb rainfall, leading to erosion and ultimately desertification.

Regenerative agriculture offers many solutions to this crisis, addressing the issue at every level, mauka to makai, from rainfall to aquifers. The thorough integration of elements in the Mainframe Design of the system ensures that less water will be used in the production of crops, and more water will be stored and available, banked in ponds, soils, and plants. Streams and springs may return, and local rainfall may increase through orographic effects if the central valley is designed with the intention to improve the hydrological cycles of the land.

Soil

Bioremediation of Soils

Healthy soil is a dynamic living ecosystem, teeming with billions of microorganisms that continuously create humus, nourish plant growth, hold water, and sequester stunning amounts of carbon. Soils that are exhausted and contaminated from years of conventional agriculture tilling and chemicals have very little microbial action, but can be restored over time using specific targeted strategies depending on the level and types of pollution. The legal requirement to make the transition from conventional to certified organic agriculture is three years. The obvious first step is to stop using chemicals altogether; then begins the journey to robust and healthy soil.

Soil is a complicated mixture, and mechanisms for the metabolism of chemical pollutants are not completely understood. The research shows that healthy populations of microorganisms in the soil have the capability of bioremediating certain pollutants.⁷ Even metals can be bound in the soil

by humic acids.⁸

Definitions:

Bioremediation: The use of soil microbes to remove or neutralize contaminants in polluted soil or water. Bacteria and fungi generally work by breaking down contaminants such as petroleum into less harmful substances. Plants can be used to aerate polluted soil and stimulate microbial action. They can also absorb contaminants such as salts and metals into their tissues, which are then harvested and disposed of.

Biodegradation

Petroleum hydrocarbons will degrade with relative ease as a result of biological metabolism. Although virtually all petroleum hydrocarbons are biodegradable, biodegradability is highly variable and dependent somewhat on the type of hydrocarbon.⁹

Phytoremediation is the direct use of living green plants for in situ removal, degradation, or containment of contaminants in soils and groundwater. Advantages of phytoremediation include that it is generally low cost and has low energy requirements; has a low environmental impact; and contributes to landscape improvement. It provides habitat for animal life, reduces surface runoff and reduces the dispersal of dust and contaminants by wind. It is suitable for large areas of land.¹⁰

Rhizofiltration is a form of phytoremediation that involves filtering water through a mass of roots to remove toxic substances or excess nutrients.

Mycoremediation

Of particular interest are fungi and mycorrhizae, which have the ability to tie up inorganic salts in waxy excretions, and degrade pollutants (many agricultural chemicals are inorganic salts). Mycorrhizae are destroyed by tillage, underscoring

WATER AND SOIL

the importance of no-till systems. Stimulating microbial and enzyme activity, mycelium reduces toxins in situ. Some fungi are hyperaccumulators, capable of absorbing and concentrating heavy metals in the mushroom fruit bodies.¹¹

To understand the extent and type of chemicals present, extensive soil testing is required across a site to measure baseline levels of pollutants. Consistent monitoring is important to document the efficacy of treatments. The most appropriate remediation protocols will depend on the types and quantities of chemicals present.

Soil Building Strategies and Bioremediation

Korean Natural Farming

Korean Natural Farming (KNF) involves the collection and cultivation of indigenous microorganisms (bacteria, fungi, nematodes, and protozoa), and the reintroduction of these microorganisms directly into agricultural systems to build rich and fertile soil. Using on-farm resources and recycling farm wastes, KNF minimizes dependency upon costly external inputs¹² and consistently produces higher yields without the use of chemical fertilizers.

The strategies and techniques of KNF were developed by Master Han Kyo Cho at the Janong Natural Farming Institute in South Korea.¹³ KNF has demonstrated its success to such a degree that it was adopted by the South Korean government. Rice farmers have since experienced larger yields, saved money on inputs, and are able to sell their rice for a premium. It has had the added benefit of cleaning the waterways, rivers, and even coastal waters.¹⁴



Farmer Samson Delos Reyes walks along kalo patches at S&J farms of Wai'anae; Photo by Jamm Aquino <http://www.activistpost.com/wp-content/uploads/2011/01/natural-farm.jpg>



Master Cho teaching on the Big Island of Hawaii <http://natural-farming.weebly.com/about-mr-cho.html>

Since its introduction into Hawai'i in 1999 by Dr. Hoon Park, KNF has been gaining in popularity with local farmers. Numerous trainings and workshops have been conducted on the islands. The University of Hawai'i's College of Tropical Agriculture and Human Resources (CTAHR) has published many articles on the techniques, established a Natural Farming Agent position and conducted field trials that have demonstrated improved plant health, increased yields and improved soil tilth using KNF techniques.¹⁵

Farmers in Hawaii are reporting success with KNF. Samson Delos Reyes of S&J Farms of Waianae stated in an article in the Honolulu Star Advertiser that **since trying Korean Natural Farming, production on his 10-acre plot has doubled.** "This is the first time having earthworms on my farm," he said, scooping up a handful of earth and nutrient-rich worm castings in his fingers. "They're cultivating the soil for me." His land was once classified as 'unsuitable for farming.'¹⁶


Chris Trump and his family have been farming 750 acres of macadamia nuts in North Kohala for 25 years. They began experimenting with Korean Natural Farming 5 years ago. Currently, they have 120 acres in its second year of utilizing KNF techniques. By August of 2016, they intend to utilize KNF on all 750 acres. He states, "This works. It is also organically certifiable and safe."

While field trials in Hawaii have thus far been conducted on a small scale, large scale experiments have been conducted in other parts of the world. Mr. Cho conducted an experiment in the Gobi Desert where previous tree planting efforts had failed three times due to harsh winds and very limited rainfall of the area. The trees he has planted using KNF techniques have had a 97% survival rate and are currently 20 feet tall. Corn and grasses have been

Contaminants are modified along the way and evaporate

PHYTO VOLATILIZATION:



Some plants take up volatile contaminants and release them into the atmosphere through transpiration. The contaminant is transformed or degraded within the plant to create a less toxic substance before and then released into the air.



Enzymes fragment contaminants and produces new plant fiber


PHYTO DEGRADATION:

Plants take up and break down contaminants through the release of enzymes and metabolic processes such as photosynthetic oxidation/reduction. In this process organic pollutants are degraded and incorporated into the plant or broken down in the soil.



PHYTO EXTRACTION:


Plants take up contaminants - mostly metals, metalloids and radionuclides- with their roots and accumulate them in large quantities within their stems and leaves. These plants have to be harvested and disposed as special waste.



Contaminants taken up into plant tissue

PHYTO STABILIZATION:

Some plants can sequester or immobilize contaminants by absorbing them into their roots and releasing a chemical that converts the contaminant to a less toxic state. This mechanism limits the migration of contaminants through water erosion, leaching, wind, and soil dispersion.



Enzymes in the roots immobilize contaminants

WATER AND SOIL

planted as well for livestock feed, and wells have been dug. Watermelon farming now provides a stable income to farmers.

At his January 2016 workshop, Master Cho expressed a keen interest in working with Maui County to teach and implement KNF in the central valley.

Cover Crops

Cover cropping is the strategy of seeding a mixture of plants in a fallow field or within a perennial or annual cropping mix for the purposes of increasing soil fertility and organic matter content. It improves soil structure, controls erosion, holds water in the soil, manages weeds and diseases, and increases biodiversity. Cover cropping is also proven to increase carbon sequestration in the soil. The basic process of cover cropping is to sow a field after harvest with a variety of plant species which are then lightly tilled into the soil when they first start to show flower buds.

Soil fertility is enriched by the variety of plant species in the cover crop mix, which usually includes nitrogen fixing species such as legumes, and dynamic accumulators, which concentrate macro and micronutrients in their leaves. Examples of dynamic accumulators include sunflower, rye, buckwheat, sesbania, and mustard. Soil organics and structure are improved via the plant's rooting and with the tilling in of all plant material.

Cover crops further protect and bind the soil structure from compaction and erosion by rain and wind. The management of water is greatly improved as the vegetative cover vastly reduces any run-off, while significantly increasing infiltration rates due to its roots and improved soil structure, i.e. high percentage of soil air gaps. Covered soil dramatically reduces water loss by reducing exposure to the



Crimson clover cover crop fixes nitrogen, suppresses weeds, prevents erosion, and provides excellent forage for honey bees.

http://media.oregonlive.com/washingtoncounty_impact/photo/clover-001jpg-a027abb64673b3af.jpg

drying effects of the sun and wind.

These effects increase and protect soil biology, and promote a living and dynamic soil ecology. Cover cropping reduces the presence of weeds by reducing their ability to germinate, occupying space that they would usually need, and by making it difficult for the weeds to complete their full life-cycle, thus not being able to produce seed. Certain cover crop species, rye and mustard as examples, have been shown to have allelopathic effects that suppress weeds and disrupt disease cycles.

Water and soil are the foundation for agriculture, and for society as a whole. Nations rise and fall following the health of their soils. It is our imperative to leave a legacy of clean, abundant water and healthy organic soils for our children, our keiki. We must mālama `āina.

Luckily, we have options for improving and restoring these vital resources that sustain life, for us and for all the earth's inhabitants.

(Endnotes)

- 1) Lucienne de Naie and Marty McMahon, M.A., Ka Waiola Project 2002-2004
- 2) <http://www.cuesa.org/article/10-ways-farmers-are-saving-water>
- 3) <http://www.rodalesorganiclife.com/home/organic-methods-hold-water>
- 4) <https://www.organicconsumers.org/news/reports-show-less-water-used-organic-farming>
- 5) http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/stelprdb1082147.pdf
- 6) <http://hcsugar.com/keeping-maui-green/water-conservation/>
- 7) http://dgc.stanford.edu/SCOPE/SCOPE_38/SCOPE_38_3.2_Matsumura_79-90.pdf
- 8) <http://www.ext.colostate.edu/mg/gardennotes/212.html>
- 9) U.S. Environmental Protection Agency, 2009
- 10) <http://www.hawaii.edu/abrp/biotech.html#anchor155092>
- 11) <http://www.hawaii.edu/abrp/Technologies/fungus.html>
- 12) Wang, K.H., DuPont, M., & Chang, K., (2012, Dec—2013, Jan/Feb) Use of Korean Natural Farming for Vegetable Crop Production in Hawai'i. Retrieved from <http://www.ctahr.hawaii.edu/sustainag/news/articles/V14-Wang-KNF.pdf>
- 13) Reddy, Rohini (2011). Cho's Global Natural Farming. Retrieved from <https://ilcasia.files.wordpress.com/2012/02/chos-global-natural-farming-sarra.pdf>
- 14) A Natural Farming Primer. Retrieved from <http://kalapanaorganics.com/natural-farming-with-indigenous-microorganisms/natural-farming/>
- 15) Wang, K.H., DuPont, M., & Chang, K., (2012, Dec—2013, Jan/Feb) Use of Korean Natural Farming for Vegetable Crop Production in Hawai'i. Retrieved from <http://www.ctahr.hawaii.edu/sustainag/news/articles/V14-Wang-KNF.pdf>
- 16) Essoyan, Susan., (2011, Jan. 18th) "Natural Selection", Honolulu Star Advertiser. Retrieved from <https://www.staradvertiser.com/hawaii-news/natural-selection/>

Multi-level Diversified Business Opportunities

Challenges facing local food production include access to affordable land for farmers, competition from foreign markets, a lack of skilled farmers, insufficient local processing and distribution facilities, and the lack of marketing and business skills of farmers. As a result of the above factors, food industry customers are unable to secure consistent supply of quality local produce.

There are, however, significant opportunities for local food production in Hawai'i, specifically on Maui. There is considerable demand for high quality local produce, with many people willing and able to pay a premium price for locally grown organic products. Maui schools, retailers, hotels and restaurants are all seeking local produce, with only a limited supply available. Exact numbers are outside of the scope of this report, but suffice to say that there is opportunity for multi-million dollar yearly contracts to supply local, island-wide, and export markets.

Maui County has extensive resources already in place to facilitate local agricultural enterprise and value-added innovation (manufacturing processes that increase the value of raw agricultural products). These include a large offering of classes, commercial kitchen facilities, small business mentoring, and a robust consortium of partners working to build capacity for agriculture enterprises.

Maui Food Innovation Center

The Maui Food Innovation Center (MFIC) provides business and technological expertise to food and agricultural entrepreneurs throughout the State of Hawai'i. A program of University of Hawai'i Maui College, MFIC helps farmers and food manufacturers increase profitability through



Workshop with interns at Hale Akua Garden Farm

the development of new value-added food products, reduces our dependence on imports, and contributes to the sustainability of island-based agriculture.

MFIC has secured funding through the Hawai'i State Legislature to renovate the former campus cafeteria in the Pilina Building at UH Maui College in Kahului into a state-of-the-art, shared-use food processing facility. This facility will have the capacity to design, test, and produce foods such as sauces, soups, jams, jellies, entrees, bakery products, dehydrated snacks, refrigerated fresh-cut produce, and raw or cooked meat, poultry and seafood products.

Sustainable Living Institute of Maui

The Sustainable Living Institute of Maui (SLIM) is a center with a primary focus on non-credit based community outreach and development activities, as well as complementing UH-Maui College credit-based activities. These activities include the development and dissemination of knowledge and the provision of services to the County of Maui community in various areas of sustainability, particularly renewable energy and sustainable agriculture.

College of Tropical Agriculture and Human Resources (CTAHR)

CTAHR is a land-grant university that provides exceptional education, research, and extension programs in tropical agriculture and food systems, family and consumer science, and natural resource management for Hawai'i and the international community. Topics of recent articles and workshops included moringa, pineapple, beekeeping, soils, legal issues for growers, breadfruit, and aquaculture. Through its CTAHR Extension, the College provides numerous publications, trainings, support staff, and project assistance for farmers.

FARM ENTERPRISE OPPORTUNITIES

The Kohala Center¹ on Hawai'i Island has generated numerous excellent and in-depth reports on agriculture, aquaculture, biofuels, livestock, and other important farming and watershed issues.

There are many professional, non-profit, and government agencies/institutions working together to help create a vital farming future for Maui.


Skilled Farmers

It is a fact that Maui does not currently have the skilled farmers nor the business infrastructure to support a full-fledged transition to scores of diversified agriculture enterprises. On 35,000 acres there is ample room for large commodity crop and livestock operations exceeding 5,000 acres (biofuels, timber, and cereals), other operations from 1,000–5,000 acres (hemp, kenaf, fruit and nut orchards,), and numerous businesses that can span the 5–500 acre range (vegetable, nurseries, seed crops, aquaculture).

Maui County, A&B, HC&S, and non-profit organizations will have to invest in several strategies to jump-start farming businesses. These include providing incentives and assistance to local farmers, recruiting successful farmers from off-island to start businesses on Maui, recruiting and training new farmers, providing mentorship, and financial assistance.

Farm Incubators

Farm incubators provide land-leasing arrangements for beginning farmers who have farming experience and a business idea, but do not have access to land. It is a low-risk environment to launch a farming business and test ideas.



Will Allen from Growing Power teaches aquaponics to students from around the world
http://www.sustainablebrands.com/sites/default/files/imagecache/635x300/article_images/growing-power-fish.jpg

Farmers usually have access to multiple acres of 'shovel-ready' irrigated land, business planning and marketing support, shared equipment and processing facilities, dry storage, greenhouses, and other core infrastructure.

Incubator farms usually have a 3–5 year tenure, expecting the farmers to graduate to their own acreage, thereby making room for new business ventures. Often incubators have a number of permanent anchor businesses who provide stability, mentorship, and profitability.

Economic Multipliers and Job Opportunities


It has been variously estimated that, for every job in the sugar industry, between 1.24 and 2.82 jobs are created in other sectors.² Considering this, the loss of 675 jobs on the sugarcane plantation will result in the loss of around 1,370 jobs total. Considering value-added processing, diversified

production, and increased agritourism, we anticipate the economic multiplier for regenerative agricultural systems to be higher than this.

Businesses must be profitable to survive. What kind of profit is possible for diversified agricultural systems? Can local food production be competitive with imports? It has been shown that sugarcane, when grown just for sugar, is not profitable and cannot compete on the global commodities market.


Economic metrics for regenerative agricultural systems are variable, but conservative estimates for diversified agricultural production put net profits at around \$5000 to \$7500 per acre per year, compared to a potential of \$50 to \$75 per acre per year for sugarcane³ in a monocrop for export, a 100 fold increase. Some farmers claim to gross, and even net, up to \$150,000 per acre per year for direct marketed organic vegetable production.⁴

SAME CANOE • Local Food Challenge



sweet potato 'uala • banana mai'a • taro kalo • breadfruit 'ulu

<http://www.oneisland.org/hawaii/>



Cover of Specialty Crops for Pacific Islands by Craig Elevitch

Farm Enterprise Opportunities

The sampling of farm enterprises listed below is by no means complete, and is meant to demonstrate the potential mosaic of various land uses for the central valley agricultural lands. Each business has its own set of required skills, markets, customers, and distribution. When imagining scores of diverse farm enterprises integrated into a regenerative agricultural mainframe design, it is evident that Maui will create many more than the 675 jobs lost to HC&S closure.

Canoe Crops

Native Hawaiian canoe crops were not traditionally grown in the central valley, but with the reduced water needs of regenerative farming, and some supplemental water, many of these crops will thrive. These crops fed, clothed, housed, and provided medicines in abundance for generations, and are some of the most important crops to consider when analyzing Maui's food systems.

- Kō - Sugar Cane
- `Ohe - Bamboo
- Niu - Coconut Palm
- Kalo - Taro
- Kī - Ti Plant
- Pia - Polynesian Arrowroot
- Uhi - Yam
- Mai`a - Banana
- `Ōlena - Turmeric
- `Awapuhi - Wild Ginger
- `Awa - Kava
- `Ulu - Breadfruit
- Wauke - Paper Mulberry
- Kukui - Candlenut Tree
- Hau - Hibiscus
- `Ōhi`a `Ai - Mountain Apple
- `Uala - Sweet Potato
- Noni - Indian Mulberry
- Ipu - Bottle Gourd

Vegetable Crops and Diversified Fruit and Nut Orchards

Demand for local consumption of vegetables, fruits and nuts far outstrips current production. From direct sales through local farmer's markets up to multi-million dollar annual contracts for Hawai`i's school lunch program, Maui is well-situated to ramp up production. There are cascading job opportunities for value-added products, including:

Dried fruits

- Jams and preserves
- Juices
- Fermented products
- Salsas
- Processed nuts and seeds (salted, dipped in chocolate, in trail mixes)
- Superfood blends
- Alcohol distilleries and brewhouses (rum, vodka, beer, wine)
- Essential oils
- Root-crop chips
- Agritourism

Superfood Crops


There is an enormous demand for fresh and prepared nutrient-rich superfoods, as exemplified by the spending habits of customers seeking healthier lifestyles. A few examples include:

- Moringa
- Turmeric
- Poha berry
- Acai berry
- Cacao

Aquaculture

HC&S has dozens of reservoirs across the central valley, ranging in size from 1 million gallons to 80 million gallons. Converting several reservoirs to fresh-water fish aquaculture in conjunction

FARM ENTERPRISE OPPORTUNITIES



Sorghum is being considered as a biofuel crop for Maui
<https://www.bungenorthamerica.com/products/categories/14-sorghum>



Growing lettuces with aquaponics at Living Aquaponics Inc.
<http://www.livingaquaponics.com>

with neighboring farm operation or aquaponics (growing vegetables in conjunction with fish) would further diversify food supplies and provide skilled job opportunities.

Livestock

To raise meats for the local market Maui needs a full-service slaughterhouse, packing house, butcher services, and a mobile slaughter unit. There is demand for fresh meat, cured specialty meats, dairy, and eggs. Please refer to the section on Holistic Planned Grazing for details.

Biofuels

There is a great opportunity to grow biofuel crops to help make Maui more self-sufficient in energy, reduce air pollution, and cut our emissions of greenhouse gases. Biofuels - fuels made from plants and organic matter - are one way to decrease our consumption of fossil fuels, especially oil. Unlike oil, coal, or natural gas, biofuels are renewable and won't run out.⁸

Biofuels include:

- Ethanol
- Biodiesel
- Biogas/methane from anaerobic digestion
- Syngas from biomass gasification

Ethanol made from bagasse, a byproduct of sugarcane processing, has potential as a transition fuel for the HC&S property. Brazil is considered as a biofuel industry leader, with the world's first sustainable biofuels economy. Touted as a policy model for other countries, its sugarcane ethanol is called "the most successful alternative fuel to date".^{9, 10} Hawai'i had hoped to spur creation of a local ethanol industry, using locally grown feedstocks, with a 2006 requirement that all motor



Polyface Farms mobile chicken coops with noticeably improved grass growing just two weeks after grazing rotation.
<https://www.pinterest.com/pin/34410384625461897/>

gasoline be blended with 10% ethanol, but no ethanol refineries have been built in the state. In 2015, the requirement was repealed (Act 161, Session Laws of Hawai'i 2015).¹⁷


Biodiesel crops include sunflower, safflower, hemp, kenaf, and soybeans, amongst others. Biodiesel burns cleaner than fossil fuels, and releases fewer pollutants and greenhouse gases into the atmosphere. Pacific Biodiesel, headquartered in Kahului, Maui, has 20 years experience internationally in biodiesel, manages a successful operation of biofuel crops on 10,000 acres on the Big Island of Hawai'i, and is considering expanding to Maui.

"Since its inception over 15 years ago, Pacific Biodiesel has built 12 facilities on the mainland U.S. and Japan, and completed expansions on several of those plants. It's newest venture, Big Island Biodiesel, located on Hawai'i Island, began production in the 4th quarter of 2012. Featuring zero-waste processing, this facility produces the highest quality biodiesel available in the country".¹¹

Biodiesel production on Maui could offer lateral job opportunities for former HC&S sugarcane workers. It is a specialized farm business with many jobs in the biodiesel plant itself, and is highly mechanized.

Textiles and Fiber Crops - Kenaf and Hemp

Kenaf (*Hibiscus cannabinis*) and hemp (*Cannabis sativa*) are promising commodity crops that merit further research. Both grow well in the tropics, have multiple high-value yields, help with soil remediation, and have the potential to provide many employment opportunities.



Kenaf bales in warehouse
<http://www.britsauto.co.za/index.php/natural-fibre-mat/>



Harvesting Hemp in Romania:
http://www.hempworld.com/hemp-cyberfarm_com/images/Harvest%20w%20tractor04.jpg

Kenaf is a warm season annual fiber crop closely related to cotton and okra that can be successfully grown on Maui. Kenaf has been used as a cordage crop to produce twine, rope, and sackcloth for over six millennia, and today there is a robust market in paper products, building materials, flotation devices, absorbents, high-tech fine oil for industry, biofuels, viable seed, and livestock feed. Kenaf grows quickly to 9'-12'; the flowers produce a delicious honey, and can likely produce 2 – 3 crops annually on Maui.¹²

Hemp yields many diverse products from foods to medicine, paper and textiles, building materials and more. Hemp is an excellent soil remediation crop and like kenaf has a long history in twine, cloth, burlap, and other textiles.

Hemp's environmental footprint is relatively small; it requires few pesticides and no herbicides. It's an excellent rotation crop, often used to suppress weeds and loosen soil before the planting of cereals. However, it requires a relatively large amount of water (albeit less than sugarcane), and its need for deep, humus-rich, nutrient-dense soil limits growing locales.¹³

Further research is needed to know if hemp is a viable and profitable crop for Maui, and if it would be invasive, as its seeds are easily dispersed by birds. "Feral cannabis is an exceptionally hardy weed, widely dispersing its seeds, which can lie dormant for 7–10 years before sprouting again."¹⁶

Kenaf and hemp also offer lateral job opportunities for retrained HC&S workers.

FARM ENTERPRISE OPPORTUNITIES

Agritourism¹⁴


As part of a whole-farm strategy that complements the farm enterprise itself, farmers can generate significant income by diversifying into agritourism, especially on Maui. Our 2.6 million annual visitors seek out fun, delicious, and educational day-trips to local farms.

It can be difficult to make a living as a small farmer, and the supplemental income from agritourism, in conjunction with real farming, can be an important part of the solution to profitability. However, this will tend to raise the value, and thus the price, of agricultural land.

There are many ways to drive business to local farms including:

- Farm tours
- Farm-to-table lunches and dinners
- Tastings
- Workshops and trainings
- Ziplines
- Dude ranches
- Fishing/hunting
- Wineries/brewhouses/distilleries
- Gift shops
- Concerts
- Fairs
- Festivals
- Weddings

The Hawai'i AgriTourism Association (HATA) connects the agriculture sector with the visitor industry and residents across Hawai'i. They provide educational and economic opportunities to members that include farm diversification, agritourism marketing, developing farm tours, and producing and selling value-added products.



Tourists enjoying the zipline at Maui Tropical Plantation
<http://mauitropicalplantation.com>

(Endnotes)

- 1) <http://kohalacenter.org/research>
- 2) <http://files.hawaii.gov/dlnr/cwrp/cch/cchma0601/DO-HCS.PDF>, page 9
- 3) <https://www.extension.iastate.edu/agdm/wholefarm/html/c3-65.html>
- 4) http://www.nola.com/business/index.ssf/2012/12/high_yields_low_prices_robust.html
- 5) <http://naturalsociety.com/sustainability-on-steroids-organic-farmer-grosses-100k-an-acre/>
- 6) <https://radicalpersonalfinance.com/making-80k-on-13-acre-with-an-urban-farm-without-owning-land-yes-please-interview-with-curtis-stone-rpf0040/>
- 7) http://pvs.kcc.hawaii.edu/ike/moolelo/polynesian_plants.html
- 8) <http://www.climatechwiki.org/technology/agriculture-biofuel-production>
- 9) https://en.wikipedia.org/wiki/Ethanol_fuel_in_Brazil#cite_note-TwoBillion-7
- 10) Sperling, Daniel and Deborah Gordon (2009), Two billion cars: driving toward sustainability
- 11) <http://www.biodiesel.com>
- 12) http://www.cres.gr/biokenaf/files/fs_inferior01_h_files/BOOKLET.pdf
- 1) <http://modernfarmer.com/2013/10/legal-industrial-hemp-wont-matter/>
- 14) <http://www.hiagtourism.org>
- 15) https://en.wikipedia.org/wiki/Ethanol_fuel_in_Hawaii
- 16) https://en.wikipedia.org/wiki/Feral_cannabis
- 17) https://en.wikipedia.org/wiki/Ethanol_fuel_in_Hawaii



Community Engagement and Story of Place

This report has examined many possibilities for how to begin a transition to regenerative agriculture. These options will require a large investment in research, design, implementation, infrastructure, training and much more. The research and potential outcomes outlined in this report are beginnings of a community discussion and vision. The key is to come together as a community, with our largest landowner, and chart this course together.

Moving forward hinges on addressing these important questions:

- How can private farm businesses have long-term access to land owned by A&B?
- Can some of the land be re-zoned to allow farmers to live on their farms? Could that include small farming communities like the plantation villages?
- What water rights will farmers have on these lands?
- Regenerative agriculture will use much less water than sugar cane. How can we guarantee watershed restoration post-sugarcane?
- Will A&B and HC&S continue to own the land and become a diversified multi-farm corporation?
- How will A&B work with the community and provide transparency regarding their farming and development agenda?
- Would A&B sell the land to a consortium of private buyers who are committed to regenerative agriculture? At what price?
- Can the land be held in perpetuity for regenerative agriculture, as a safeguard against development?
- How would the land be managed and distributed to farmers?

If A&B would sell the land at market value, a compelling alternative emerges: forming an island-wide Maui Farm Cooperative. Every citizen of Maui could be either a worker-member or consumer-member with voting rights, profit shares, access to healthy island-grown food, even health care. Under the umbrella of the Maui Farm Cooperative, independently managed divisions would oversee each main business branch: livestock, tree crops, vegetable crops, agritourism, composts, marketing, distribution, irrigation, education, and so on.

Sugarcane production ends this year. Maui needs agriculture jobs. The community has an opportunity to come together and help usher in a new era of farming on Maui. Cultivating beneficial relationships between stakeholders is the foundation for the success of any project. We look forward to hearing your stories, addressing your questions and concerns, and incorporating new ideas.



Legend

Agricultural Soils Productivity Rating

- A (Excellent)
- B (Good)
- C (Fair)
- D (Poor)
- E (Very Poor)
- Not Classified

Important Agricultural Lands

HC&S Reservoirs

Haiku Ditch

Lowrie Ditch

Kaunikoia Ditch

Hamakua Ditch

Rainfall Zones

Prevailing Winds

Land Use Zones

Excellent to Good soils, slopes 1-3%, irrigated
Sited to:

① Annual and Perennial Crops
Orchard Crops
Commodity and Energy Crops
Seed Crops
Pasture

Excellent to Good soils, slopes 3-5%, minimal irrigation
Sited to:

② Annual and Perennial Crops
Orchard Crops
Seed Crops
Pasture

Fair, Poor to Very Poor soils, slopes 5%+, minimal to no irrigation
Sited to:

③ Timber
Pasture
Habitat

④ Farm Hub
Processing Centers
Agricultural Sales Facilities
Plant Nurseries

⑤ Agricultural Park
Educational Facilities


⑥ Aquaculture
Aquaponics


⑦ Small Farm Allotments
Community Gardens

**Conceptual Plan Draft
A&B IAL**

Source:
Land Study Bureau (1987)
U.S. Geological Survey
City/County
This map has been prepared for general planning purposes only.

Data Overlays Prepared by:
**Permaculture Design International
for Maui Tomorrow**
February 2016





Comparison Map Current (2010) HC&S Ag Lands and Lands Committed to Future Ag as IAL.

Current HC&S Lands in Ag Production:
35,000 Acres

HC&S Lands Designated as Important Ag Lands
2009 by LUC:
27,133 Acres

Current Water Use for 35,000 Acres from East
Maui Sources 160mgd

Legend

Crop Type

- Sugar
- Pineapple
- Seed Corn
- Pasture/ Other


Proposed Important Agricultural Lands

Existing HC&S Crop Lands not in IAL Boundary

Development Areas Proposed During Maui Island
Plan Process by A&B Inc. (Approx. 3,000 Acres)

Proposed MLP Development Area

Map Source: HC&S IAL Application to LUC
Composite Map Provided by Maui Tomorrow May 2010



Legume Trees with pods Edible to Livestock				
Common Name	Species Name	Climate	Native Range	Nitrogen Fixation
Carob	* <i>Ceratonia siliqua</i>	Mediterranean	Mediterranean	No
Winter Thorn	* <i>Faidherbia albida</i>	Arid to humid tropical lowlands and highlands	Africa	Yes
African Locust Bean	* <i>Parkia biglobosa</i>	Semiarid to humid tropical lowlands	Africa	Yes
Palo Verde	* <i>Parkinsonia aculeata</i>	Arid to semiarid tropics and subtropics	Americas	No
Monkey Bread	* <i>Piliostigma thonjii</i>	Semiarid tropics	Africa	Yes
Manilla Tamarind	<i>Pithecellobium dulce</i>	Semi-arid to humid tropical lowlands	Americas	Yes
Honey Mesquite	* <i>Prosopis glandulosa</i>	Arid to semi-arid, subtropics to colA	North America	Yes
Kiawe	* <i>Prosopis pallida</i>	Semiarid tropics	South America	Yes
Monkeypod Tree	<i>Samanea (= Albizia) saman</i>	Semi-arid to humid tropical lowlands	tropical Americas	Yes
Wild Cassia	<i>Senna singueana</i>	Semiarid tropics	Africa	No

*pods also edible by humans

Adapted from Martin, Frank. Selecting the best Plants

Animal Feed (Grasses)								
Common Name	Species Name	Food	Feed	Fiber	Const.	Fuel	Soil Amend.	Erosion Control
Bermuda	<i>Cynodon dactylon</i>	0	5	0	0	0	0	4
Guinea	<i>Panicum maximum</i>	0	4	0	1	0	0	2
Kikuyu	<i>Pennisetum clandestin</i>	0	5	0	0	0	0	4
Napier	<i>Pennisetum purpureur</i>	0	5	0	2	1	0	4
Pangola	<i>Digitaria decumbens</i>	0	5	0	0	0	0	4
Star	<i>Cynodon nlemfluensis</i>	0	5	0	0	0	0	4
Sudan	<i>Sorghum sudanense</i>	0	5	0	2	1	0	1
Animal Feed (Legumes)								
Common Name	Species Name	Food	Feed	Fiber	Const.	Fuel	Soil Amend.	Erosion Control
Calliandra	<i>Calliandra calothyrsus</i>		3					
Jack bean	<i>Canavalia ensiformis</i>	1	3	0	0	0	2	2
Leucaena	<i>Leucaena leucacephal</i>	4	4	0	2	4	4	3
Mesquite	<i>Prosopis sp.</i>	2	5	0	3	4	3	4
Mother-of-cacao	<i>Gliricidia sepium</i>	2	3	0	3	3	3	3
Tibet Tree	<i>Albizia lebbeck</i>		4					

Adapted from Martin, Frank. Selecting the best Plants

Fiber					
Common Name	Species Name	Annual or Perennial	Growth Habit	Drought	Other Uses
Abaca	<i>Musa textilis</i>	perennial	large herb	no	cord
Cotton	<i>Gossypium spp.</i>	annual	large herb	no	stuffing
Hemp	<i>Cannabis sativa</i>	annual	large herb	no	yes
Jute	<i>Corchorus capsularis</i>	annual	herb	no	cord
Kapok	<i>Ceiba pendandra</i>	perennial	tree	no	stuffing
Kenaf	<i>Hibiscus spp.</i>	annual	herb	no	cord, leaves
Mahoc	<i>Hibiscus tiliaceus</i>	perennial	tree	yes	no
Sisal	<i>Agava spp.</i>	perennial	herb	yes	cord
Baobab	<i>Adansonia digitate</i>	perennial	large tree	yes	clothing
Paper mulberry	<i>Boussingaltia</i>	perennial	large shrub	yes	clothing

From Toesmeier, E. Carbon Farming Ch 22 industrial oils

Annual Biofuels Plants	
Latin Name	Common Name
<i>Helianthus annus</i>	Sunflower
<i>Carthamus tinctorius L.</i>	Safflower
<i>Glycine max</i>	Soybean
<i>Cannabis sativa l</i>	Hemp
<i>Hibiscus cannabinus</i>	Kenaf

Perennial Oil Biofuels plants				
Inedible oil yields compared				
Latin Name	Common Name	Climate	Seed or fruit yield t/ha	Oil yield t/ha
<i>Jatropha curcas</i>	Jatropha	Tropical lowlands, semi-arid to humid	1-15.0	0.3-5
<i>Aleurites mollucanus</i>	Candlenut	Tropical humid to semiarid	16	3
<i>Ricinus communis</i>	Castor bean	Tropics, subtropics, high or lowlands, semi-arid to humid	0.5-5	0.3-2.7
<i>Pongamia pinnata</i>	Pongamia	Subtropics, tropics, lowlands, highlands, humid	5-8.0	1.8
<i>Simmondsia chinensis</i>	Jojoba	Subtropics, arid to semi-arid	2.2-4.5	0.5-1.1
<i>Azadirachta indica</i>	Neem	Tropics, humid to semi-arid		0.5

Cover Crops							
Common Name	Species Name	Adapt*	Propagation	Nitrogen Fixation	Erosion Control	Mulch	Ground Cover
Cowpea	<i>Vigna unguiculata</i>	I	seeds	4	3	4	4
Hosei	<i>Vigna hosei</i>	I,W	cuttings	4	4	5	5
Indigo	<i>Indigofera spp.</i>	I,W	seeds	4	5	3	5
Jack bean	<i>Canavalia ensiformis</i>	I,W	seeds	4	3	3	4
Lablab bean	<i>Dolichos lablab</i>	I,W	seeds	4	1-5	2	1-5
Perennial peanut	<i>Arachis spp.</i>	I	seeds, cutting	4	5	2	4-5
Sun hemp	<i>Crotalaria juncea</i>	I	seeds	4	3	4	3
Tinaroo	<i>Glycine wightii</i>	I,W	seeds	4	5	3	5
Velvet bean	<i>Stizolobium deeringianum</i>	I,W	seeds	4	5	3	5

Alley Cropping Fodder Trees								
Common Name	Species Name	Adapt*	Alley Crop	Nitrogen Fixing	Erosion Control	Mulch	WindBreak	Shade
Agati	<i>Sesbania grandiflora</i>	I	5	3	4	2	1	1
Calliandra	<i>Calliandra calothyrsus</i>	I,W	5	4	4	3	3	3
Horseradish tree	<i>Moringa oleifera</i>	I	4	0	1	2	0	1
Ice Cream Bean	<i>Inga edulis</i>	I,W	5	3	3	4	2	3
Leucaena	<i>Leucaena leucacephala</i>	I	5	5	2	3	1	1
Mother of cacao	<i>Gliricidia sepium</i>	I	4	4	2	3	1	3
Pigeon pea	<i>Cajanus cajan</i>	I	5	4	3	3	0	0
Tibet Tree	<i>Albizia lebbek</i>	I,W						

*Key to adaptation: W= hot wet tropics; U= upland tropics, D= dry tropics, I = intermediate, neither too wet nor too dry.

Adapted from Martin, Frank. Selecting the best Plants

Specialty crops (beverages, oil, spices, sugar)			
Common Name	Species Name	Annual/Perennial	Growth Habit
BEVERAGES			
Cacao	<i>Theobroma cacao</i>	perennial	small tree
Coffee	<i>Coffea arabica, C. robusta</i>	perennial	small tree
Tea	<i>Camellia sinensis</i>	perennial	shrub
SPECIALTY OILS			
Tung	<i>Aleurites spp.</i>	perennial	tree
Ylang-Ylang	<i>Cananga odorata</i>	perennial	tree
SPICES			
Indonesian Cardamom	<i>Amomum cardamomum</i>	perennial	herb
Cardamom	<i>Elettaria cardamomum</i>	perennial	herb
Turmeric	<i>Curcuma domestica</i>	perennial	herb
Cloves	<i>Syzygium aromaticum</i>	perennial	small tree
Nutmeg & Mace	<i>Myristica fragans</i>	perennial	tree
Pepper	<i>Piper nigrum</i>	perennial	vine
Galangal	<i>Alpinia galanga</i>	perennial	herb
Ginger	<i>Zingiber officinale</i>	perennial	herb
Vanilla	<i>Vanilla fragrans</i>	perennial	vine

Arid/Semi-Arid adapted Multi-purpose windbreak species, Adapted from Elevitch, C. Multipurpose windbreak trees.										
Botanical Name	Common Name	Mature Size (ft)	N-Fixer	Fruit/ Nut/	Fodder	Bee Forage	Wood/ Timber	Wind break	Growth Rate	Potentially Invasive
<i>Acacia koa</i>	Koa	50-80'	Y		Y	Y	Y	Y	F	
<i>Acacia confusa</i>	Formosa Koa	50-80'	Y		Y	Y	Y	Y	F	
<i>Acrocarpus fraxinifolius</i>	Pink Cedar	80-160'					Y	Y	F	
<i>Albizia lebeck</i>	Tibet Tree	25-35'	Y		Y	Y	Y	Y	M	Y
<i>Aleurites moluccana</i>	Kukui, Candlenut	50-80'		P	P			Y	M	
<i>Anacardium occidentale</i>	Cashew	35-40'		Y	Y		Y	Y	S	
<i>Annona muricata</i>	Soursop	<20'		Y				Y	S	
<i>Araucaria bidwillii</i>	bunya-bunya pine	90-120'		Y	Y		P		S	
<i>Artocarpus heterophyllus</i>	Jackfruit	30-70'		Y	Y		Y	Y	S	less drought hardy
<i>Azadirachta indica</i>	Neem	40-60			Y	Y	Y	Y	M	
<i>Bambusa oldhamii</i>	Oldhamii	40-60'		Y			Y	Y	M	
<i>Casimiroa edulis</i>	White Sapote	20-45'		Y			P	Y	S	
<i>Ceratonia siliqua</i>	Carob	45-55'		Y	Y		Y	Y	S	
<i>Chrysophyllum cainito</i>	Star Apple/Caimito	25-50'		Y				Y	S	
<i>Cocos nucifera</i>	Coconut	30-90'		Y		Y	Y	Y	S	less drought hardy
<i>Eucalyptus cameldulensis</i>	Red River Gum	80-120'				Y	Y	Y	F	Y
<i>Eucalyptus sideroxylon</i>	Red Iron Bark	50-60'				Y	Y	Y	F	Y
<i>Eucalyptus robusta</i>	Swamp Mahogany	80-120'				Y	Y	Y	F	
<i>Gliricidia sepium</i>	Madre de cacao	30-35'	Y		Y	Y	Y	Y	F	
<i>Mangifera indica</i>	Mango	80-120'		Y		P	Y	Y	S	
<i>Manilkara zapota</i>	Sapodilla	50-60		Y	Y		Y	Y	S	
<i>Moringa oleifera</i>	horseradish tree	30-45'		Y	Y	Y		Y	F	
<i>Morus nigra</i>	Mulberry	20-25'		Y	Y	Y		Y	M	
<i>Pithecellobium dulce</i>	Manila tamarind	35-50'	Y		P	Y	Y	Y	M	Y
<i>Prosopis glandulosa</i>	Honey Mesquite	20-30'	Y	Y	Y	Y		Y	M	
<i>Pterocarpus indicus</i>	Narra	90-120'	Y		P		Y	Y	M	
<i>Senna siamea</i>	Pheasantwood	50-60'			P		Y	Y	F	
<i>Swietenia macrophylla</i>	Mahogany	90-120'					Y	Y	M	less drought hardy
<i>Tamarindus indica</i>	Tamarind	80-100		Y	Y		Y	Y	S	
<i>Thyostacys siamensis</i>	Monastery Bamboo	20-45'		Y	P		Y	Y	M	

P=potential

Limitations of Livestock in Agroforestry				
Livestock Species	Damage Young Trees	Scratch or Dig	Silvopasture Only	Diverse Perennial Understory
Cattle	Yes		Yes	
Chickens		Yes	Yes	Yes^
Ducks & Muscovies				Yes^
Geese				Yes
Goats	Yes		Yes	
Hogs	Yes	Yes	Yes	Yes**
Sheep	Yes		Yes	
Turkeys		Yes	Yes	Yes*

From Toensmeier, E.

Agroforestry Functions of Livestock						
Livestock Species	Mow & Graze	Clear Brush	Eat Bugs	Till	Weed Grass Only	Clean Drops
Cattle	Yes	Yes*				Yes*
Chickens	Yes		Yes	Yes		Yes
Ducks	Yes		Yes			
Geese	Yes				Yes	
Goats	Yes	Yes				
Hogs	Yes		Yes*	Yes	Yes*	Yes
Sheep	Yes	Yes*				Yes
Turkeys	Yes		Yes			

*Breed dependent

^Sequence and crop dependent

POTENTIAL FOOD CROPS FOR MAUI						
Grains						
CommonName	SpeciesName	Annual/Perennial	Principal Nutrients	Yield (lbs/ac)	Water Use (gal/ac/day)	Reference
Amaranth	<i>A. cruentis</i> <i>A. hypochondr</i>	protein, starch	protein, starch	800	1785	http://tinyurl.com/zvntkw2
Corn, Maize	<i>Zea mays</i>	protein, oil, starch	protein, oil, starch	6,000-12,000	893 - 1785	http://tinyurl.com/huo96od
Pearl Millet	<i>Pennisetum americanum</i>	protein, starch	protein, starch	3,000-4,000	893	http://tinyurl.com/jqk2tn4
Quinoa	<i>Chenopodium quinoa</i>	protein, starch	protein, starch	900-1,200	714 - 1071	http://tinyurl.com/nv4d34o
Sorghum	<i>S. bicolor</i>	protein, starch	protein, starch	4,000-5,000	893 - 1785	http://tinyurl.com/hla42od
Legumes						
Common Name	Species Name	Annual/Perennial	Principal Nutrients	Yield	Water Use	Reference
Bean, common	<i>Phaseolus vulgaris</i>	annual	protein, starch	1,200-1,800	1517 – 1875	http://tinyurl.com/ha4scb9
Chick pea, garbanzo	<i>Cicer arietum</i>	annual	protein, starch	800-2,000	446 - 893	http://tinyurl.com/h9r5vzg
Cowpea	<i>Vigna sinensis</i>	annual	protein, vit. B	1,000-3,000	893 – 1785	http://tinyurl.com/gr4yzyo
Lablab	<i>Dolichos lablab</i>	annual	protein, starch	1,000-2,000	1785 - 3571	http://tinyurl.com/zc8vmdj
Lima bean	<i>Phaseolus vulgaris</i>	annual	protein, vit. B, starch	2,000-3,000	1161 - 1785	http://tinyurl.com/h47cnn5
Mung bean	<i>Vigna radiate</i>	annual	protein, starch	300-2,000	1071 - 1518	http://tinyurl.com/zgcx19f
Pigeon pea	<i>Cajanus cajan</i>	annual or weak perennial	protein	700	1785 - 3303	http://tinyurl.com/hnje8xa
Roots and Tubers						
Common Name	Species Name	Annual, Bi/ Perennial	Principal Nutrients	Yield	Water Use	Reference
Cassava	<i>Manihot esculenta</i>	per. Grown as annual	starch	15,000	3571 - 4465	http://tinyurl.com/h6z2uhy
Jicama	<i>Pachyrhizus erosus</i>	weak per. used as annual	starch, protein	10,000-14,000	N/A	http://tinyurl.com/gt44ka8
Sweet Potato	<i>Ipomea batatus</i>	per. Grown as annual	starch, vit. C, maybe A	28,000-32,000	2233 - 3126	http://tinyurl.com/gmeexfa

POTENTIAL FOOD CROPS FOR MAUI -Continued						
Fruit Vegetables						
CommonName	SpeciesName	Annual/Perennial	Principal Nutrients	Yield (lbs/ac)	Water Use (gal/ac/day)	Reference
Chayote	<i>Sechium edulis</i>	perennial	tips high in vitamins, minerals	40,000-80,000	3571 - 4465	http://tinyurl.com/jeepw3g
Eggplant	<i>Solanum melongena</i>	weak perennial	low nut. Value	15,000-30,000	1339	http://tinyurl.com/jctgl2o
Okra	<i>Abelmoschus esculentus</i>	annual	fair source of most nutrients	7,000-10,000	893 - 1785	http://tinyurl.com/h3kr4wk
Pepper	<i>Capsicum annum</i>	weak perennial	vit. A & C	10,000-20,000	1785 - 2232	http://tinyurl.com/h9odpzh
Pumpkin tropical	<i>Cucurbita moschata</i>	weak perennial	vit. A & C, seed high in oil & protein	9,000-11,000	1518 - 2143	http://tinyurl.com/huqcrxb
Misc. Vegetables						
Common Name	Species Name	Annual/Perennial	Principal Nutrients	Yield	Water Use	Reference
Artichoke	<i>Cynara scolymus</i>	perennial		9,000-11,000	1339 - 1785	http://tinyurl.com/z7jzehu
Asparagus	<i>Asparagus officinale</i>	perennial	vit. C	2,500-3,000	1339 - 2232	http://tinyurl.com/hbcq7k
Tropical Fruit Crops						
Common Name	Species Name	Annual/Perennial	Principal Nutrients	Yield	Water Use	Reference
Avocado	<i>Persia americana</i>	perennial	oil	4,000-7,500	1785 - 3572	http://tinyurl.com/zfjyhmv
Breadfruit	<i>Artocarpus elastica</i>	perennial	starch	12,000-25,000	1339 - 3572	http://tinyurl.com/k8csw63
Carob	<i>Ceratonia siliqua</i>	perennial	starch	6,000-8,000	446 - 893	http://tinyurl.com/j74wfrj
Citrus	<i>Citrus spp.</i>	perennial	vit. A & C	15,000-30,000	1785 - 3572	http://tinyurl.com/z3elcl2
Mango	<i>Mangifera indica</i>	perennial	vit. A & C	20,000-40,000	4465 - 8035	http://tinyurl.com/z7rljuy
Papaya	<i>Carica papaya</i>	perennial	vit. A & C	20,000-30,000	1785 - 2678	http://tinyurl.com/zraf4f2
Tropical Nuts						
Common Name	Species Name	Annual/Perennial	Principal Nutrients	Yield	Water Use	Reference
Cashew	<i>Anacardium occidentale</i>	perennial	protein	800-1,500	893 - 2232	http://tinyurl.com/j9ftwc7
Macadamia	<i>Macadamia spp.</i>	perennial	protein	2,500-3,000	2678 - 3572	http://tinyurl.com/jdxkkg7

General Report References**Maui County:**

Maui General Plan 2030:
<http://www.co.maui.hi.us/DocumentCenter/View/84679>

County of Maui Website with links to General Plan 2030 as well as State and 9 County individual Community Plans:
<http://www.co.maui.hi.us/index.aspx?NID=421>

Maui Ag Design Conference Preliminary Report:
http://h-cx.com/Hawaii_Local_Exchange/Maui_Ag_Design_Conference_files/Maui%20Ag%20Design%20Conf%20-%20-%20Preliminary%20Report.pdf

Maui water:
<https://repositories.lib.utexas.edu/bitstream/handle/2152/ETD-UT-2010-08-1835/GRUBERT-THESIS.pdf?sequence=1>

Maui Agriculture Design Conference 2010 Chart:
http://h-cx.com/Hawaii_Local_Exchange/Maui_Ag_Design_Conference.html

Maui Time article The Sun Sets on Maui Sugar:
<http://mauitime.com/news/business/as-the-sun-sets-on-maui-sugar-we-ask-what-the-closing-of-abs-plantation-means-for-us/>

Maui Watch article about A&B financials:
<http://mauiwatch.com/2016/01/top-stories-of-2015-part-4-the-future-of-sugar-whats-next-for-hcs/>

Hawaii Department of Agriculture - Important Agriculture Lands:
<http://hdoa.hawaii.gov/chair/new-agriculture-initiatives/important-ag-lands-ial/>

Summary of Important Agriculture Lands:
<http://files.hawaii.gov/dlnr/cwrm/cch/cchma1301/CCHMA1301-20141230-HC&S-OB.pdf>

Kohala Center:

The Kohala Center Sustainability and Community Building Center
<http://kohalacenter.org/>

Kohala Center Reports
<http://kohalacenter.org/research>

Local Food Report: p.6 factors affecting local access
http://kohalacenter.org/docs/reports/2015_LocalFoodHawaiiSchools.pdf

Regenerative Agriculture:

Laguna Blanca: http://www.tompkinsconservation.org/who_we_are.htm

Room for Industrial Agriculture on Small Farms! <http://www.civilbeat.com/2016/02/hawaii-needs-all-types-of-farms-including-industrial-operations/>

Silvopasture: <http://www.grazinglandshawaii.org/2015/06/06/some-truths-about-silvopasture/>

Quivira Ranch: http://quiviracoalition.org/Carbon_Ranch/index.html

Cover Crops: <http://www.nytimes.com/2016/02/07/business/cover-crops-a-farming-revolution-with-deep-roots-in-the-past.html?smid=fb-share&r=1>

Regenerative Agriculture - 9 techniques: <http://sheldonfrith.com/2015/12/15/a-brief-introduction-to-most-important-techniques-in-regenerative-agriculture/>

Vetiver: http://www.huffingtonpost.com/jacob-devaney/water-purification-beauty_b_9122230.html

Fao report on biodiversity and the ecosystem:
<http://www.fao.org/docrep/005/y4586e/y4586e05.htm>

Video on Holistic grazing in the Chihuahan Desert:
<https://www.facebook.com/regenerationinternational/videos/1718077591754103/>

Agroforestry in France - 3,000 hectares:
http://www.theguardian.com/environment/2012/aug/21/agroforestry-france-farming-revival?CMP=share_btn_fb

USDA NRCS Soils: <http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>

Regenerative Agriculture and Climate Change:

Rodale Regenerative Agriculture and Climate Change:
<http://rodaleinstitute.org/assets/WhitePaper.pdf>

ZERI – Zero Emissions Research Initiatives:
<http://www.zeri.org/ZERI/Home.html>
http://www.zeri.org/ZERI/Case_Studies.html

Conservation agriculture in the semi arid tropics:
https://www.researchgate.net/publication/279971311_Conservation_Agriculture_in_the_Semi-Arid_Tropics

FAO - The importance of Soil key to drought resistance soil and sustained food production:
<http://www.fao.org/docrep/009/a0100e/a0100e.pdf>

Carbon Farming Solutions lists:
<http://carbonfarmingsolution.com/recommended-readings#http://www.fao.org/docrep/018/i3325e/i3325e00.htm>

Climate Change and the Future of Farming:
<https://cgspace.cgiar.org/bitstream/handle/10568/69225/CCAFS%20info%20note%20AgCop21.pdf?sequence=1&isAllowed=y>

Soil and carbon:
http://www.huffingtonpost.com/larry-kopald/nature-wants-her-carbon-b_b_6173358.html

Permaculture and Climate Change:
<http://www.permacultureclimatechange.org>

Eric Tonsemeier book:
<http://carbonfarmingsolution.com>

Carbon Cycle Institute - data on carbon farming:
<http://www.carboncycle.org/draft-carbon-farm-plan/>

Soil and carbon sequestration:
<http://www.commondreams.org/news/2015/04/24/solution-climate-change-right-under-our-feet>

Farm Economics References

American Farmland Trust:
<https://www.farmland.org>

Earning \$100,000/acre:
<http://craftsmanship.net/drought-fighters/>

Crossroads Resource Center:
<http://www.crcworks.org/?submit=fff>

Socioeconomic analysis of bioenergy:
http://www.globalbioenergy.org/uploads/media/0312_FAO_-_Socio-economic_analysis_of_bioenergy_systems_a_focus_on_employment.pdf

Economic rationale for agricultural regeneration and rural infrastructure investment in South Africa:
<http://www.up.ac.za/media/shared/61/WP/wp117.zp39505.pdf>

FAO green jobs for a revitalized food and agriculture sector:
http://www.fao.org/fileadmin/user_upload/sustainability/pdf/FAO_green_jobs_paper_March_31.pdf

Understanding economic multipliers:
<https://www.agecon.purdue.edu/crd/pdf/ec686.pdf>

APPENDICES

Sequestering carbon in the soil

http://e360.yale.edu/feature/soil_as_carbon_storehouse_new_weapon_in_climate_fight/2744/

<http://www.fastcoexist.com/3052240/restoring-global-soil-quality-is-one-of-the-best-things-we-can-do-for-climate-change>

<http://www.perennialsolutions.org/legume-trees-with-pods-edible-to-livestock>

<http://www.perennialsolutions.org/livestock-integration-reducing-labor-and-fossil-fuel-inputs>

Korean Natural Farming

KNF of Hawaii: <http://naturalfarminghawaii.net>

Cho's Global Korean Natural Farming Handbook:

<https://ilcasia.files.wordpress.com/2012/02/chos-global-natural-farming-sarra.pdf>

Semi-Arid Tropics

CGIAR : consultative group on international agricultural research
<http://www.cgiar.org/>

ICRISAT: international crops research institute for semi arid tropics
<http://www.icrisat.org/>

ICRISAT Dr W. Dar interview about 6 semi-arid staple crops (pasture cropping options)
<http://foodtank.com/news/2013/12/agricultural-improvements-in-the-semi-arid-tropics-an-interview-with-dr-wi>

SOM pools in tropical savanna under agroforestry system in northeastern Brazil:
<http://www.scielo.br/pdf/rarv/v38n4/14.pdf>

SOC accumulation in arid and semi arid areas after afforestation:
<http://www.pjoes.com/pdf/22.2/Pol.J.Environ.Stud.Vol.22.No.2.611-620.pdf>

Marsha Hanzi:
<http://www.marsha.com.br/category/articles/>

Water related technologies for semiarid lands:
<https://www.princeton.edu/~ota/disk3/1983/8331/8331.PDF>

Carbon farming with woody bamboo by R. Lal:
<http://www.sciencedirect.com/science/article/pii/S2351989415000281>

Graphics

xerxes society 10 ac integrated farm graphic

http://www.xerxes.org/wp-content/uploads/2008/09/farming_for_pollinators_brochure.pdf

Uses of Kenaf: <http://www.kenaf-fiber.com/en/infotec-tabella4.asp>

Cover Crops and Green Manures for Hawai'i

<http://www.ctahr.hawaii.edu/sustainag/Database.asp>

Additional information about green manures and cover crops

[Advice on obtaining seeds of green manure and cover crops in Hawaii](#)

[Accelerating the Adoption and Implementation of Proven Cover Crop Technologies in Hawaii](#) by Dr. John McHugh, CropCare Hawaii (posted 7/07)

[Green Manure Crops that can help to control Nematodes in Dryland Taro \(.wmv video\)](#) by Dr. Susan Miyasaka, Alton Arakaki, Dr. Brent Sipes, and Dr. Ray Ming
[UC Davis SARE Program](#) (with searchable cover crop database)
[FAO Grasses & Legumes Index](#)
[Better Pastures for the Tropics and Subtropics](#)

Regenerative Agriculture OCA Bibliography

<https://www.organicconsumers.org/news/regenerative-agriculture-annotated->

World Agroforestry Center

World Agroforestry Center Flickr album:
<https://www.flickr.com/photos/icraf/sets/>

World Agroforestry Center homepage:
<http://www.worldagroforestry.org/bibliography>

Training

IFOAM Training manual for Org. ag in arid and semi-arid tropics
http://www.s-ge.com/sites/default/files/private_files/IFOAM%20Training%20Manual%20for%20Organic%20Agriculture%20in%20the%20Arid%20and%20Semiarid%20Tropics_0.pdf

The case for going 100% organic

Denmark goes all organic:
<http://foodrevolution.org/blog/denmark-organic-country/>

UN calls for end to industrial agriculture:
<http://www.truth-out.org/opinion/item/27636-united-nations-calls-for-an-end-to-industrialized-farming>

Energy independent island:
<http://www.independent.co.uk/news/world/europe/sams-island-to-be-fossil-fuel-free-after-winning-back-danish-government-funds-a6787446.html>

Sikkim 100% organic:
<http://siliguritimes.com/in/sikkim-becomes-the-first-fully-organic-state-of-india/>

Chemical exposure and GMO

Agricultural chemical exposure in children:
<http://thefern.org/2014/03/warning-signs-how-pesticides-harm-the-young-brain/>

Glyphosate Portugal:
<http://sustainablepulse.com/2016/02/09/portuguese-medical-association-president-calls-for-global-ban-on-glyphosate/#VroO7M6Jnwx>

GE Alfalfa - case study of failure:
<http://www.digitaljournal.com/news/environment/new-study-by-usda-proves-they-were-wrong-about-ge-alfalfa/article/455607#tab=comments&sc=0>

Miscellaneous

Generating power from water pipes:
<http://magazine.good.is/articles/portland-pipeline-water-turbine-power>

Maps

Plantation village maps from 1950's: <http://magis.manoa.hawaii.edu/maps/digital/maui.1954-1957.html>

Per Acre Costs for Organic Farming Installations

Detailed chart of per-acre costs for site preparation and installation that includes windbreaks, contour plantings, fodder crops, mulch, buffer zones, and more prepared by USDA Natural Resources Conservation Services., http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs141p2_037112.pdf

CTAHR Seed Initiative

<http://www.ctahr.hawaii.edu/sustainag/news/articles/V11-Valenzuela-seedinitiative.pdf>

Curriculum Vitae

ALBERT PEREZ

1. Academic Degrees and Certificates

Bachelor of Arts in Economics	University of Hawaii at Manoa Honolulu, HI	1981
Master of Urban & Regional Planning	University of Hawaii at Manoa Honolulu, HI	1987
Certificate in Geographic Information Systems	Green River Community College Auburn, WA	1997
Geographic Information Systems Professional Certification (GISP)	GIS Certification Institute	2008

2. Professional Experience

<u>Executive Director</u> June 2015 – present	Maui Tomorrow Foundation, Inc. Wailuku, HI
<u>GIS Analyst</u> Dec 1996 – Aug 2011	Washington State Department of Transportation Olympia, WA
<u>GIS Analyst</u> Jun 1996 – Apr 1998	The Wilderness Society Seattle, WA
<u>Cartographer</u> Apr 1996-Jun 1996	Snohomish County Planning and Development Services Everett, WA
<u>Principal</u> Apr 1995-Dec 1997	Freeland Planning and Development Services Freeland, WA
<u>President</u> Dec 1991-May 1993	Pacific GeoPlanning Kihei, HI and Freeland, WA
<u>Co-Director</u> Nov 1989-Dec 1991	Maui Tomorrow Kahului, HI
<u>Planner</u> Nov 1988-Oct 1989	County of Maui Planning Department Wailuku, HI
<u>Planner</u> Nov 1986-Oct 1988	Hawaii Office of State Planning Honolulu, HI

3. Information Technology Experience

- ▲ Geographic Information Systems: ArcGIS, ArcGIS Server, ARC/INFO, Spatial Analyst, 3D Analyst, Network
- ▲ Programming: Python, Visual Basic, ARC Macro Language (AML)
- ▲ Databases: MS Access, INFO
- ▲ Operating Systems: Windows, UNIX (HP, Solaris and AIX), Macintosh, DOS

4. Information Technology Training

ESRI	Apr 1997	Programming with Avenue
ESRI	Jul 1998	Introduction to GIS Hydrology
ESRI	May 1999	Programming MapObjects with Visual Basic
ESRI	Jul 1999	Environmental Modeling Using Arcview Spatial Analyst
WSDOT	Apr 2001	Managing Project Delivery
Capitol Business Machines	Dec 2001	Database Planning and Design
Washington Geographic Information Council	May 2002	Remote Sensing and LandSat7 Imagery Workshop
Northwest Environmental Training Center	Sep 2002	Introduction to Groundwater Hydrology
ESRI Online	Jan 2003	Exploring the VBA Environment
Capitol Business Machines	Sep 2003	Structured Query Language
Northwest Environmental Training Center	Sep 2004	Introduction to ArcHydro
ESRI Online	Jan 2005	Creating, Editing, and Managing Geodatabases for ArcGIS 8.3
ESRI Online	Jun 2005	Creating and Editing Geodatabase Topology with ArcGIS 9.0-9.1 (for ArcEditor and ArcInfo)
U.S. Geological Survey	Jan 2006	<u>National Hydrography Dataset Applications</u>
ESRI	Mar 2006	<u>Customizing ArcGIS Desktop (for ArcGIS 9.0-9.1)</u>
Urban and Regional Information Systems Association	Sep 2006	Best Practices for Developing Geographic Information Models
ESRI	Jun 2008	Introduction to Geoprocessing Scripts Using Python
Opposite Strengths, Inc.	Feb 2011	The Power of Opposite Strengths (Interpersonal Relationships)
ESRI	Apr 2011	Introduction to ArcGIS Server
ESRI	Apr 2011	Creating Effective Web Applications with ArcGIS Server

5. Publications

Washington Wildlife Habitat Connectivity Working Group. 2010. "Washington Connected Landscapes Project: Statewide Analysis". Washington Departments of Fish and Wildlife and Transportation, Olympia, WA

Xiong, D., Lee, R., Saulsbury, J., Lanzer, E., and Perez, A. "Remote Sensing Applications for Environmental Analysis in Transportation Planning: Application to the Washington State I-405 Corridor". *Oak Ridge National Laboratory and Washington State Department of Transportation*, August 30, 2004, WA-RD-593-1.

Perez, A. Chapter 3 in "Source Water Protection Project: A Comparison of Watershed Delineation Methods in ARC/INFO and ArcView GIS." [Hydrologic and Hydraulic Modeling Support with Geographic Information Systems](#). Compiled and edited by Dr. David Maidment and Dr. Dean Djokic. Redlands, CA. Environmental Systems Research Institute. 2000. Pages 53-64.

Gersib, R., Hilliard, T., Molash E., Neugebauer-Rex, J., Park J., Schanz R., Van Natta, L., Lautz, K., Perez, A., Prosser, K., and Johnson, T. 2005. "Enhancing Transportation Project Delivery Through Watershed Characterization: SR-167 Study - Report to the WSDOT Urban Corridors Office". Washington State Department of Transportation, Olympia, WA.

Gersib, R., Hilliard, T., Molash E., Neugebauer-Rex, J., Park J., Schanz R., Van Natta, L., Lautz, K., Perez, A., Prosser, K., Wood, B., and Johnson, T. 2004. "Enhancing Transportation Project Delivery Through Watershed Characterization: I-405 / SR-520 Study- Report to the WSDOT Urban Corridors Office". Washington State Department of Transportation, Olympia, WA.

Gersib, R., Haddaway, B., Hilliard, T., Molash E., Park J., Perez, A., Schanz R., and Stone, V. 2004. "Enhancing Transportation Project Delivery Through Watershed Characterization: I-405 Case Study - Report to the WSDOT Urban Corridors Office". Washington State Department of Transportation, Olympia, WA.

Perez, A., 1992. "West Maui Regional Carrying Capacity Study". Sponsored by the State of Hawaii, Office of State Planning. Prepared for the County of Maui, Planning Department. Albert Perez, Planning Consultant, Haiku, HI.

Contributor: Bebe, B., Hontz, G., and Swanander, A., 1990. "Maui Coastal Scenic Resources Study". Prepared for the County of Maui, Planning Department. Environmental Planning Associates, Inc., Kihei, HI.

Perez, A., 1990. "South Maui Community Development Forum"

Perez, A., 1989. "Maui: The Built-Out Scenario". Prepared on behalf of Maui Tomorrow, Makawao, HI.

Perez, A. and Quinn, M., 1986. "Helicopter Noise in Haleakala National Park". Prepared in partial fulfillment of Masters in Urban and Regional Planning, Department of Urban and Regional Planning, University of Hawaii at Manoa, Honolulu, HI.

6. Presentations

Perez, A. 2004. "Enhancing Transportation Project Delivery Through Watershed Characterization". 19th Annual Northwest GIS User Conference, Sun Valley, ID.

Perez, A. 1999. "[Source Water Protection Project: A Comparison of Watershed Delineation Methods in ARC/INFO and ArcView](#)". 19th Annual International ESRI User Group Conference, San Diego, CA.

Perez, A. 1998. "[Surface Drinking Water Source Watershed Delineation](#)". 18th Annual International ESRI User Group Conference, San Diego, CA.

Perez, A. 1998. "Surface Drinking Water Source Watershed Delineation". 13th Northwest ESRI Users Group Conference, Seattle, WA

Mills, M., and Perez, A. 1997. "Sensitive Groundwater Mapping Project". 2nd Annual Washington Hydrology Symposium, Olympia, WA

Mills, M., and Perez, A. 1997. "Sensitive Groundwater Mapping Project". American Water Works Association Conference, Seattle, WA.

Perez, A., 1990. "South Maui Development Forum". Kihei, HI

Perez, A. 1989. "Maui: The Built-Out Scenario", Maui Tomorrow, Makawao, HI.

Perez, A., and Sands, F. 1989. "Citizens Development Workshop", Maui Tomorrow, Makawao, HI.