

# Instrument endogeneity and identification-robust tests: some analytical results \*

Firmin Doko <sup>†</sup>  
Université de Montréal

Jean-Marie Dufour <sup>‡</sup>  
McGill University

First version: November 2007

Revised: December 2007, February 2008

This version: February 2008

Compiled: May 31, 2008, 1:36pm

This paper has been published in the *Journal of Statistical Planning and Inference*, 138 (2008), 2649-2661.

---

\* The authors thank Tze Leung Lai and Ingram Olkin for several useful comments. This work was supported by the William Dow Chair in Political Economy (McGill University), the Canada Research Chair Program (Chair in Econometrics, Université de Montréal), the Bank of Canada (Research Fellowship), a Guggenheim Fellowship, a Konrad-Adenauer Fellowship (Alexander-von-Humboldt Foundation, Germany), the Institut de finance mathématique de Montréal (IFM2), the Canadian Network of Centres of Excellence [program on *Mathematics of Information Technology and Complex Systems* (MITACS)], the Natural Sciences and Engineering Research Council of Canada, the Social Sciences and Humanities Research Council of Canada, the Fonds de recherche sur la société et la culture (Québec), and the Fonds de recherche sur la nature et les technologies (Québec), and a Killam Fellowship (Canada Council for the Arts).

<sup>†</sup> Université de Montréal, Département de Sciences Économiques, C.P. 6128, succ. Centre-ville, Montréal, QC H3C 3J7. TEL: (1) 514 343 6111 ext. 5283; Fax: (1) 514 343 7221; e-mail : f.doko.tchatoka@umontreal.ca.

<sup>‡</sup> William Dow Professor of Economics, McGill University, Centre interuniversitaire de recherche en analyse des organisations (CIRANO), and Centre interuniversitaire de recherche en économie quantitative (CIREQ). Mailing address: Department of Economics, McGill University, Leacock Building, Room 519, 855 Sherbrooke Street West, Montréal, Québec H3A 2T7, Canada. TEL: (1) 514 398 8879; FAX: (1) 514 398 4938; e-mail: jean-marie.dufour@mcgill.ca . Web page: <http://www.jeanmariedulfour.com>

## ABSTRACT

When some explanatory variables in a regression are correlated with the disturbance term, instrumental variable methods are typically employed to make reliable inferences. Furthermore, to avoid difficulties associated with weak instruments, identification robust methods are often proposed. However, it is hard to assess whether an instrumental variable is valid in practice because instrument validity is based on the questionable assumption that some of them are exogenous. In this paper, we focus on structural models and analyze the effects of instrument endogeneity on two identification-robust procedures, the Anderson-Rubin (1949, AR) and the Kleibergen (2002, K) tests, with or without weak instruments. Two main setups are considered: (1) the level of “instrument” endogeneity is fixed (does not depend on the sample size), and (2) the instruments are *locally exogenous*, *i.e.* the parameter which controls instrument endogeneity approaches zero as the sample size increases. In the first setup, we show that both test procedures are in general consistent against the presence of invalid instruments (hence asymptotically invalid for the hypothesis of interest), whether the instruments are “strong” or “weak”. We also describe cases where test consistency may not hold, but the asymptotic distribution is modified in a way that would lead to size distortions in large samples. These include, in particular, cases where the 2SLS estimator remains consistent, but the AR and K tests are asymptotically invalid. In the second setup, we find (non-degenerate) asymptotic non-central chi-square distributions in all cases, and describe cases where the non-centrality parameter is zero and the asymptotic distribution remains the same as in the case of valid instruments (despite the presence of invalid instruments). Overall, our results underscore the importance of checking for the presence of possibly invalid instruments when applying “identification-robust” tests.

**Key words:** simultaneous equations; instrumental variables; locally weak instruments; invalid instruments; locally exogenous instruments.

## Contents

<b>1. Introduction</b>	<b>1</b>
<b>2. Framework</b>	<b>2</b>
<b>3. Test statistics</b>	<b>5</b>
<b>4. Asymptotic theory with invalid and weak instruments</b>	<b>7</b>
4.1. Possibly invalid instruments . . . . .	7
4.2. Locally exogenous instruments . . . . .	8
<b>5. Conclusion</b>	<b>10</b>
<b>A. Appendix: Proofs</b>	<b>11</b>

## List of Assumptions, Propositions and Theorems

<b>4.1 Theorem</b> : Asymptotic distribution of the AR statistic . . . . .	7
<b>4.2 Theorem</b> : Asymptotic distribution of the K statistic . . . . .	7
<b>4.3 Theorem</b> : Asymptotic distributions with locally exogenous instruments . . . . .	9
<b>4.4 Theorem</b> : Asymptotic distributions with weak locally exogenous instruments . . . . .	9
Proof of Theorem <b>4.1</b> . . . . .	11
Proof of Theorem <b>4.2</b> . . . . .	12
Proof of Theorem <b>4.3</b> . . . . .	14
Proof of Theorem <b>4.4</b> . . . . .	14

## 1. Introduction

The last decade shows growing interest for so-called *weak instruments* problems in the econometric literature, *i.e.* situations where “instruments” are poorly correlated with endogenous explanatory variables; see the reviews of Dufour (2003) and Stock, Wright and Yogo (2002). More generally, these can be viewed as situations where model parameters are not identified or close not to being *identifiable*, as meant in the econometric literature [see Dufour and Hsiao (2008)]. When instruments are weak, the limiting distributions of standard test statistics – like Student, Wald, likelihood ratio and Lagrange multiplier criteria in structural models – often depend heavily on nuisance parameters; see *e.g.* Phillips (1989), Bekker (1994), Dufour (1997), Staiger and Stock (1997) and Wang and Zivot (1998). In particular, standard Wald-type procedures based on the use of asymptotic standard errors are very unreliable in the presence of weak identification. As a result, several authors have worked on proposing more reliable statistical procedures that would be applicable in such contexts.

Interestingly, in the early days of simultaneous-equations econometrics, Anderson and Rubin (1949, AR) proposed a procedure which is completely robust to weak instruments as well as to other difficulties such as missing instruments [see Dufour (2003), Dufour and Taamouti (2005, 2007)]. But the AR procedure may suffer from power losses when too many instruments are used. So alternative methods largely try to palliate this difficulty, for example: pseudo-pivotal LM-type and LR-type statistics [Wang and Zivot (1998), Kleibergen (2002), Moreira (2003)], sample-splitting methods [Dufour and Jasiak (2001)], approximately optimal instruments [Dufour and Taamouti (2003)], systematic search methods for identifying relevant instruments and excluding unimportant instruments [Hall, Rudebusch and Wilcox (1996), Hall and Peixe (2003), Dufour and Taamouti (2003), Donald and Newey (2001)].

However, all these procedures – including the AR method – rely on the availability on valid (exogenous) instruments. This raises the question: what happens to these procedures when some of the instruments are endogenous? In particular, what happens if an invalid instrument is added to a set of valid instruments? How robust are these inference procedures to instrument endogeneity? Do alternative inference procedures behave differently? If yes, what is their relative performance in the presence of instrument endogeneity?

We view the problem of instrument endogeneity as important because it is hard in practice to assess whether an instrumental variable is valid, *i.e.* whether it is uncorrelated with the disturbance term. Instrument validity or orthogonality tests are built on the availability of a number of undisputed valid instruments, at least as great as the number of coefficients to be estimated, whereas the validity of those initial instruments is not testable.

In the econometric literature, little is known about test procedures when some instruments are both invalid and weak. Hahn and Hausman (2003) deal with both instrument endogeneity and weakness, but they focus on estimation. Ashley (2006) proposed a sensitivity analysis of IV estimators when instruments are imperfect, his results however are only applicable if the covariance between the structural error term and some instruments is known, which is not necessary the case as it is showed in this paper. Analyzing the effect of instrument invalidity on the limiting and empirical distribution of IV estimators, Kivet and Niemczyk (2006) conclude that for the accuracy of asymp-

otic approximations, instrument weakness is much more detrimental than instrument invalidity and that the realizations of IV estimators based on strong but possibly invalid instruments seem usually much closer to the true parameter values than those obtained from valid but weak instruments. However, this finding of Kiviet and Niemczyk leaves open crucial questions: is it really possible to make reliable inference with endogenous instruments? Is instrument endogeneity really more detrimental than its weakness on inference procedures like a general family of Anderson-Rubin-type procedures? Swanson and Chao (2005) proposed a weak-instrument unified framework, but they do not take into account possible invalidity of some instruments. Finally, Small (2007) has recently studied the properties of tests for identifying restrictions [Sargan (1958), Kadane and Anderson (1977)], which can be sensitive to the use of “endogenous instruments”, and he proposed a sensitivity analysis to assess the importance of the issue. These results, however, do not allow for weak identification.

In this paper, we focus on structural models and analyze the effects of instrument endogeneity on the Anderson and Rubin (1949) and Kleibergen (2002) tests, in the presence of possibly weak instruments. After formulating a general asymptotic framework which allows one to study these issues in a convenient way, we consider two main setups: (1) the one where the level of “instrument” endogeneity is fixed (*i.e.*, it does not depend on the sample size), and (2) the one where the instruments are *locally exogenous*, *i.e.* the parameter which controls instrument endogeneity approaches zero (at rate  $T^{-1/2}$ ) as the sample size increases. In the first setup, we show that both test procedures studied are in general consistent against the presence of invalid instruments (hence asymptotically invalid for the hypothesis of interest), whether the instruments are “strong” or “weak”. We also observe there are cases where consistency may not hold, but the asymptotic distribution is modified in a way that would lead to size distortions in large samples. In the second setup, asymptotic non-central chi-square distributions are derived, and we give conditions under which the non-centrality parameter is zero and the asymptotic distribution remains the same as in the case of valid instruments (despite the presence of invalid instruments). Overall, our results underscore the importance of checking for the presence of possibly invalid instruments when applying “identification-robust” tests.

The paper is organized as follows. Section 2 formulates the model considered. Section 3 describes briefly the statistics. Section 4 studies the asymptotic distribution of the statistics (under the null hypothesis) when some instruments are invalid. We conclude in section 5. Proofs are presented in the Appendix.

## 2. Framework

We consider the following standard simultaneous equation framework, which has been the basis of much work on inference in model with possibly weak instruments [see the reviews of Dufour (2003) and Stock et al. (2002)]:

$$y = Y\beta + Z\gamma + u, \tag{2.1}$$

$$Y = X\Pi + Z\Gamma + V, \tag{2.2}$$

where  $y$  is a  $T \times 1$  vector of observations on the dependent variable,  $Y = [Y_1, \dots, Y_T]'$  is a  $T \times G$  matrix of observations on explanatory (possibly) endogenous variables ( $G \geq 1$ ),  $Z$  is a  $T \times r$  matrix of observations on the included exogenous variables,  $X = [X_1, \dots, X_T]'$  is a  $T \times k$  ( $k \geq G$ ) full-column-rank matrix of observations on (supposedly) “exogenous variables” (instruments) excluded from the structural equation (2.1),  $u = [u_1, \dots, u_T]'$  and  $V = [V_1, \dots, V_T]'$  are respectively  $T \times 1$  vector and  $T \times G$  disturbance matrices,  $\beta$  and  $\gamma$  are  $G \times 1$  and  $r \times 1$  vectors of unknown coefficients,  $\Pi$  and  $\Gamma$  are  $k \times G$  and  $r \times G$  matrices of unknown coefficients. The usual necessary and sufficient condition for identification of this model is  $\text{rank}(\Pi) = G$ .

Since we focus on the parameter  $\beta$  in our analysis, we can simplify the presentation of the results without notable loss of generality by setting  $\gamma = 0$  and  $\Gamma = 0$ , so that  $Z$  drops from the model. With this simplification, model (2.1)-(2.2) reduces to

$$y = Y\beta + u, \quad (2.3)$$

$$Y = X\Pi + V. \quad (2.4)$$

We also assume that

$$u_t = V_t' a + \varepsilon_t, \quad t = 1, \dots, T, \quad (2.5)$$

$$X_t = X_{0t} + W_t, \quad t = 1, \dots, T, \quad (2.6)$$

$$u_t = W_t' b + e_t, \quad t = 1, \dots, T, \quad (2.7)$$

where  $X_0 = [X_{01}, \dots, X_{0T}]'$  is a  $T \times k$  matrix of exogenous variables,  $\varepsilon_t$  is uncorrelated with  $V_t$ , and  $e_t$  are uncorrelated with  $W_t$ .  $V_t$  and  $W_t$  have mean zero and covariance matrices  $\Sigma_V$  and  $\Sigma_W$ ,  $\varepsilon_t$  and  $e_t$  have mean zero and variances  $\sigma_\varepsilon^2$  and  $\sigma_e^2$  respectively, while  $a$  and  $b$  are  $G \times 1$  and  $k \times 1$  vectors of unknown coefficients. (2.5)-(2.7) can be rewritten in matrix form as:

$$u = Va + \varepsilon, \quad (2.8)$$

$$X = X_0 + W, \quad (2.9)$$

$$u = Wb + e, \quad (2.10)$$

where  $X_0$  is uncorrelated with  $W$ ,  $V$ ,  $\varepsilon$  and  $e$ , while  $W = [W_1, \dots, W_T]'$  is uncorrelated with  $e$  but may be correlated with  $u$  (when  $b \neq 0$ ). So  $a$  controls the endogeneity of the variable  $Y$ , whereas  $b$  represents the possible endogeneity of the instruments  $X$ . If  $b = 0$ , the instruments  $X$  are valid; otherwise, they are invalid (endogenous). More precisely, if  $b \neq 0$ , *i.e.*, there exists at least one  $i$  such that  $b_i \neq 0$ ,  $i = 1, \dots, k$ , and the corresponding variable  $X_i$  does not constitute a valid instrument.

We also make the following generic assumptions on the asymptotic behaviour of model variables [where  $A > 0$  for a matrix  $A$  means that  $A$  is positive definite (p.d.), and  $\rightarrow$  refers to limits as  $T \rightarrow \infty$ ]:

$$\frac{1}{T} \begin{bmatrix} V & \varepsilon \end{bmatrix}' \begin{bmatrix} V & \varepsilon \end{bmatrix} \xrightarrow{p} \begin{bmatrix} \Sigma_V & 0' \\ 0 & \sigma_\varepsilon^2 \end{bmatrix} > 0, \quad (2.11)$$

$$\frac{1}{T} [ X_0 \ W ]' [ X_0 \ W ] \xrightarrow{p} \begin{bmatrix} \Sigma_0 & 0' \\ 0 & \Sigma_W \end{bmatrix}, \quad \Sigma_0 > 0, \quad (2.12)$$

$$\frac{1}{T} X_0' [ V \ \varepsilon \ e ] \xrightarrow{p} 0, \quad (2.13)$$

$$\frac{1}{T} X' X \xrightarrow{p} \Sigma_X, \quad (2.14)$$

$$\frac{1}{T} [ W \ e ]' [ W \ e ] \xrightarrow{p} \begin{bmatrix} \Sigma_W & 0' \\ 0 & \sigma_e^2 \end{bmatrix}, \quad (2.15)$$

$$\frac{1}{T} W' V \xrightarrow{p} \Sigma_{WV}, \quad (2.16)$$

$$\frac{1}{\sqrt{T}} \begin{bmatrix} X'e \\ (X'W - \Sigma_W)b \end{bmatrix} \xrightarrow{L} \begin{bmatrix} S_e \\ S_b \end{bmatrix} \sim N[0, \Sigma_S], \quad (2.17)$$

$$S_e \sim N[0, \sigma_e^2 \Sigma_X], \quad S_b \sim N[0, \sigma_e^2 \Sigma_b], \quad (2.18)$$

where  $\Sigma_V$  is  $G \times G$  fixed matrix,  $\Sigma_0$  and  $\Sigma_W$  are  $k \times k$  fixed matrices,  $S_e$  and  $S_b$  are  $k \times 1$  random vectors. Note that  $\Sigma_W$  may be singular, and  $S_b$  may not be independent of  $S_e$ .

From the above assumptions, it is easy to see that:

$$\frac{1}{T} X_0' u \xrightarrow{p} 0, \quad \frac{1}{T} X_0' e \xrightarrow{p} 0, \quad (2.19)$$

$$\frac{1}{T} X' u \xrightarrow{p} \varphi = \Sigma_W b, \quad \frac{X' V}{T} \xrightarrow{p} \Sigma_{WV}, \quad (2.20)$$

$$\frac{1}{T} [ u \ V ]' [ u \ V ] \xrightarrow{p} \Sigma = \begin{bmatrix} \sigma_u^2 & \delta' \\ \delta & \Sigma_V \end{bmatrix} > 0, \quad (2.21)$$

$$\frac{1}{T} [ u \ W ]' [ u \ W ] \xrightarrow{p} \begin{bmatrix} \sigma_u^2 & \varphi' \\ \varphi & \Sigma_W \end{bmatrix}, \quad (2.22)$$

$$\frac{1}{T} [ W \ V ]' [ \varepsilon \ e ] \xrightarrow{p} \begin{bmatrix} \delta_{W\varepsilon} & 0 \\ 0 & \delta_{Ve} \end{bmatrix}, \quad (2.23)$$

where

$$\delta = \Sigma_V a, \quad \sigma_u^2 = a' \Sigma_V a + \sigma_\varepsilon^2 = \sigma_\varepsilon^2 + b' \Sigma_W b, \quad (2.24)$$

$$\Sigma_V a = \Sigma'_{WV} b + \delta_{Ve}, \quad \Sigma_W b = \Sigma_{WV} a + \delta_{W\varepsilon}, \quad (2.25)$$

$$\Sigma_X = \Sigma_0 + \Sigma_W > 0, \quad \Sigma_{XY} = \Sigma_X \Pi + \Sigma_{WV}, \quad (2.26)$$

$$\Sigma_Y = \Pi' \Sigma_X \Pi + \Sigma_V + \Sigma'_{WV} \Pi + \Pi' \Sigma_{WV}. \quad (2.27)$$

Finally, we denote by  $\mathcal{N}(\Sigma_W)$  the null set of the linear map on  $\mathbb{R}^k$  characterized by the matrix  $\Sigma_W$  :

$$\mathcal{N}(\Sigma_W) = \{x \in \mathbb{R}^k : \Sigma_W x = 0\}. \quad (2.28)$$

If  $\Sigma_W$  is a full-column-rank matrix, then  $\mathcal{N}(\Sigma_W) = \{0\}$ ; otherwise, there is at least one  $x_0 \neq 0$  such that  $\Sigma_W x_0 = 0$ .

The setup described above is quite wide and does allow one to study several questions associated with the possible presence of “invalid” instruments. In particular, an important practical problem consists in studying the effect on inference of adding an “invalid” instrument to a list of valid (possibly identifying) instruments. Note that this problem is distinct from studying the effect of imposing “incorrect” overidentifying restrictions [as done by Small (2007)]. To better see the issues studied here, it will be useful to consider a simple example.

**Example 2.1** Consider a model with one endogenous explanatory variable ( $G = 1$ ) and two candidate instruments ( $k = 2$ ). Then  $Y$  and  $V$  are  $T \times 1$  vectors,  $X = [X_1, X_2]$  and  $W = [W_1, W_2]$  are  $T \times 2$  matrices,  $\Pi = [\pi_1, \pi_2]'$  and  $b = [b_1, b_2]'$  are vectors of dimension 2, and

$$Y = X\Pi + V = X_1\pi_1 + X_2\pi_2 + V, \quad (2.29)$$

$$u = Wb + e = W_1b_1 + W_2b_2 + e. \quad (2.30)$$

Let us further assume that  $X_1$  is a valid instrument (with  $W_1 = 0$ ),  $E[u | X_1] = 0$ ,  $X_2 = W_2$ ,  $\pi_2 = 0$  and  $b_1 = 0$ , where  $e$  is independent of  $X_1$  and  $X_2$  (with finite mean zero), so that

$$Y = X\Pi + V = X_1\pi_1 + V, \quad (2.31)$$

$$u = Wb + e = W_2b_2 + e. \quad (2.32)$$

Here  $W_2$  is not a “valid” instrument when  $b_2 \neq 0$ . But the structural equation (2.3) may in principle be estimated using only  $X_1$  as an instrument, because  $E[u | X_1] = 0$ ; if  $X_1$  is not a weak instrument ( $\pi_1 \neq 0$ ) and satisfies usual regularity conditions, a consistent estimate of  $\beta$  can be obtained. Among other things, we study below the effect (on some identification-robust tests) of taking  $X_2$  as an instrument when  $b_2 \neq 0$ , *i.e.* when  $X_2$  is correlated with  $u$ . Note that the condition  $E[u | X_1] = 0$  does not entail  $E[e | X_1, X_2] = 0$ , which is a maintained hypothesis used by Small (2007). So the problem considered here is distinct from the problem of testing overidentifying restrictions [studied, for example, by Sargan (1958), Kadane and Anderson (1977) and Small (2007)].

### 3. Test statistics

We consider in this paper the problem of testing

$$H_0 : \beta = \beta_0 \quad (3.1)$$

where some of the “instruments” used are in fact endogenous ( $b \neq 0$ ). We analyze the behavior of the Anderson-Rubin and Kleibergen statistics. The Anderson and Rubin (1949) test for  $H_0$  in equation (2.3) involves considering the transformed equation

$$y - Y\beta_0 = X\Delta + \varepsilon \quad (3.2)$$

where  $\Delta = \Pi(\beta - \beta_0)$  and  $\varepsilon = u + V(\beta - \beta_0)$ .  $H_0$  can then be assessed by testing  $H'_0 : \Delta = 0$ . The AR-statistic for  $H'_0$  is given by

$$AR(\beta_0) = \frac{1}{k} \frac{(y - Y\beta_0)' P_X (y - Y\beta_0)}{(y - Y\beta_0)' M_X (y - Y\beta_0) / (T - k)} \quad (3.3)$$

where  $M_B = I - P_B$  and  $P_B = B(B'B)^{-1}B'$  is the projection matrix on the space spanned by the columns of  $B$ . If  $b = 0$ , the asymptotic distribution of  $AR(\beta_0)$  is a  $\chi^2(k)/k$  under  $H_0$ . If furthermore  $u \sim N[0, \sigma^2 I_T]$  and  $X$  is independent of  $u$ , then  $AR(\beta_0) \sim F(k, T - k)$  under  $H_0$  irrespective of whether the instruments are strong or weak. However, when some instruments are invalid, the distribution of the AR statistic may be affected.

Kleibergen (2002) proposed a modification of the AR statistic to take into account the fact that this statistic may have low power when there are too many instruments in the model. The modified statistic for testing  $H_0$  can be written

$$K(\beta_0) = \frac{(y - Y\beta_0)' P_{\tilde{Y}(\beta_0)} (y - Y\beta_0)}{(y - Y\beta_0)' M_X (y - Y\beta_0) / (T - k)} \quad (3.4)$$

where

$$\tilde{Y}(\beta_0) = X\tilde{\Pi}(\beta_0), \quad \tilde{\Pi}(\beta_0) = (X'X)^{-1}X' \left[ Y - (y - Y\beta_0) \frac{S_{uV}(\beta_0)}{S_{uu}(\beta_0)} \right], \quad (3.5)$$

$$S_{uu}(\beta_0) = \frac{1}{T - k} (y - Y\beta_0)' M_X (y - Y\beta_0), \quad S_{uV}(\beta_0) = \frac{1}{T - k} (y - Y\beta_0)' M_X Y. \quad (3.6)$$

Unlike the AR statistic which projects  $y - Y\beta_0$  on the  $k$  columns of  $X$ , the K statistic projects  $y - Y\beta_0$  on the  $G$  columns of  $X\tilde{\Pi}(\beta_0)$ . If the instruments  $X$  are exogenous,  $\tilde{\Pi}(\beta_0)$  is both a consistent estimator of  $\Pi$  and asymptotically independent of  $X'(y - Y\beta_0)$  under  $H_0$ , and  $K(\beta_0)$  converges to a  $\chi^2(G)$ . However, if some instruments are invalid ( $b \neq 0$ ),  $\tilde{\Pi}(\beta_0)$  may not be asymptotically independent of  $X'(y - Y\beta_0)$  and the asymptotic distribution of the K statistic may not be a  $\chi^2(G)$ .<sup>1</sup>

If the model contains only one instrument and one endogenous variable ( $G = k = 1$ ), the AR and K statistics are equivalent and pivotal even in finite samples whenever  $b = 0$ . When  $k > 1$ , even if  $b = 0$ , the K statistic is not pivotal in finite samples but is asymptotically pivotal, whereas the AR statistic is pivotal even in finite samples (when  $X$  is independent of  $u$ ). Following Staiger and Stock (1997), we refer to the *locally weak-instrument* asymptotic setup by considering a limiting sequence of  $\Pi$  where  $\Pi$  is local-to-zero. We also consider a limiting sequence of  $b$  where  $b$  is local-to-zero. We refer to this later limiting sequence as *locally exogenous* instruments asymptotic.

---

<sup>1</sup>We do not study this paper conditional tests such as those proposed by Moreira (2003), because the distributional theory for such tests is considerably more complex and would go beyond the scope of a short paper like the present one.

## 4. Asymptotic theory with invalid and weak instruments

In this section, we study the large-sample properties of the statistics described above when some of the instruments used are invalid. Two setups are considered. The first is the possibly invalid instrument setup, *i.e.*, the endogeneity parameter  $b$  is a fixed vector. The second is the locally exogenous instrument setup, *i.e.*,  $b$  is local-to-zero.

### 4.1. Possibly invalid instruments

We consider first the case where the endogeneity parameter  $b$  is a constant vector and we analyze the asymptotic distributions of the statistics. Our results cover both strong and weak-instrument asymptotic. Theorem 4.1 below summarizes the asymptotic behavior of the  $AR$  statistic when some instruments may be endogenous. For a random variable  $S$  whose distribution depends on the sample size  $T$ , the notation  $S \xrightarrow{L} +\infty$  means that  $P[S > x] \rightarrow 1$  as  $T \rightarrow \infty$ , for any  $x$ .

**Theorem 4.1** ASYMPTOTIC DISTRIBUTION OF THE  $AR$  STATISTIC. *Suppose that the assumptions (2.3)-(2.18) hold, with  $b = b_0$  and  $\beta = \beta_0$ , where  $b_0$  and  $\beta_0$  are given vectors. If  $b_0 \notin \mathcal{N}(\Sigma_W)$ , then*

$$AR(\beta_0) \xrightarrow{L} +\infty. \quad (4.1)$$

*If  $b_0 \in \mathcal{N}(\Sigma_W)$ , then*

$$AR(\beta_0) \xrightarrow{L} \frac{1}{k\sigma_u^2} (S_e + S_b)' \Sigma_X^{-1} (S_e + S_b) \quad (4.2)$$

*where  $S_e$  and  $S_b$  are defined in (2.17)-(2.18). If  $b_0 = 0$ , then*

$$AR(\beta_0) \xrightarrow{L} \frac{1}{k} \chi^2(k). \quad (4.3)$$

In the above theorem, no restriction is imposed on the rank of  $\Pi$ . In particular, the result holds even if  $\Pi$  is not a full-column rank matrix. When  $b_0 \notin \mathcal{N}(\Sigma_W)$ , the  $AR$  statistic diverges under the null hypothesis  $H_0$ . When  $b_0 \in \mathcal{N}(\Sigma_W)$ , the limiting distribution of the  $AR$  statistic does not diverge, but the  $AR$  test is not valid unless  $S_b = 0$ . Of course, when  $b_0 = 0$  – which is the classical exogenous instrument setup –  $S_b = 0$  and the  $AR$  test is asymptotically valid.

Theorem 4.2 below summarizes the asymptotic behavior of the  $K$  statistic when some instruments are possibly invalid.

**Theorem 4.2** ASYMPTOTIC DISTRIBUTION OF THE  $K$  STATISTIC. *Suppose that the assumptions (2.3)-(2.18) hold, with  $b = b_0$  and  $\beta = \beta_0$ , where  $b_0$  and  $\beta_0$  are given vectors.*

*(A) If  $b_0 \notin \mathcal{N}(\Sigma_W)$  then*

$$K(\beta_0) \xrightarrow{L} +\infty \quad (4.4)$$

*when at least one of the following two conditions holds: (i)  $\Pi = \Pi_0 \neq 0$  with  $\text{rank}(\tilde{\Sigma}_{XY}) = G$ , or (ii)  $\Pi = \Pi_0/\sqrt{T}$  with  $\text{rank}(\Sigma_{XY}^*) = G$ , where*

$$\tilde{\Sigma}_{XY} = \Sigma_{XY} - \Sigma_W b_0 (q_{uV}/\bar{\sigma}_u^2), \quad \Sigma_{XY}^* = \Sigma_{WV} - \Sigma_W b_0 (q_{uV}/\bar{\sigma}_u^2),$$

$$q_{uV} = \delta' - b_0' \Sigma_W \Sigma_X^{-1} \Sigma_{WV}, \quad \bar{\sigma}_u^2 = \sigma_u^2 - b_0' \Sigma_W \Sigma_X^{-1} \Sigma_W b_0.$$

(B) If  $b_0 \in \mathcal{N}(\Sigma_W)$ , then

$$K(\beta_0) \xrightarrow{L} \frac{1}{\sigma_u^2} (S_e + S_b)' \Sigma_X^{-1} \Sigma_{XY} (\Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY})^{-1} \Sigma_{XY}' \Sigma_X^{-1} (S_e + S_b) \quad (4.5)$$

when  $\Pi = \Pi_0 \neq 0$  and  $\text{rank}(\Sigma_{XY}) = G$ , and

$$K(\beta_0) \xrightarrow{L} \frac{1}{\sigma_u^2} (S_e + S_b)' \Sigma_X^{-1} \Sigma_{WV} (\Sigma_{WV}' \Sigma_X^{-1} \Sigma_{WV})^{-1} \Sigma_{WV}' \Sigma_X^{-1} (S_e + S_b) \quad (4.6)$$

when  $\Pi = \Pi_0/\sqrt{T}$  and  $\text{rank}(\Sigma_{WV}) = G$ . (C) If  $b_0 = 0$ , then

$$K(\beta_0) \xrightarrow{L} \chi^2(G) \quad (4.7)$$

when at least one of the following two conditions holds: (i)  $\Pi = \Pi_0 \neq 0$  with  $\text{rank}(\Sigma_{XY}) = G$ , or (ii)  $\Pi = \Pi_0/\sqrt{T}$  with  $\text{rank}(\Sigma_{WV}) = G$ .

Unlike Theorem 4.1 for the AR statistic, Theorem 4.2 requires an additional rank assumption. When  $b_0 \notin \mathcal{N}(\Sigma_W)$ , the null limiting distribution of the K statistic diverges. This means that the K test often rejects  $H_0$  asymptotically when  $b_0 \notin \mathcal{N}(\Sigma_W)$ . Furthermore, when  $b_0 \in \mathcal{N}(\Sigma_W)$ , the K test is not asymptotically valid unless  $S_b = 0$ . As expected, if  $b_0 = 0$  (i.e.,  $S_b = 0$ ), the K statistic converges to a  $\chi^2(G)$ . It is worthwhile to note that the case where the rank assumption fails [e.g., the partial identification of  $\beta$ ] is not covered in this paper.

Finally, it is interesting to observe that the limiting value of the two-stage least-squares (2SLS) estimator of  $\beta$ ,

$$\tilde{\beta} = (\hat{Y}' \hat{Y})^{-1} \hat{Y}' y = [Y' X (X' X)^{-1} X' Y]^{-1} Y' X (X' X)^{-1} X' y, \quad (4.8)$$

is given by

$$\text{plim}_{T \rightarrow \infty} \tilde{\beta} = \beta + [\Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY}]^{-1} \Sigma_{XY}' \Sigma_X^{-1} \Sigma_W b \quad (4.9)$$

provided  $\text{rank}(\Sigma_{XY}) = G$ , so that  $\tilde{\beta}$  is consistent when  $b_0 \in \mathcal{N}(\Sigma_W)$  and  $\Sigma_{XY}$  has full column rank (even if some instruments are invalid). If  $b_0 \in \mathcal{N}(\Sigma_W)$  but  $b_0 \neq 0$ , the asymptotic level of the Anderson-Rubin and Kleibergen tests can be affected.

## 4.2. Locally exogenous instruments

We consider now the case where the endogeneity parameter  $b$  is local-to-zero. As in the previous subsection, we analyze the limiting distributions of the statistics. The results also cover two setups: locally exogenous instruments [ $\Pi = \Pi_0 \neq 0$ ,  $b = b_0/\sqrt{T}$ ], and weak locally exogenous instruments [ $\Pi = \Pi_0/\sqrt{T}$ ,  $b = b_0/\sqrt{T}$ ]. Theorem 4.3 and Theorem 4.4 below derive the distributions of the statistics for both setups.

**Theorem 4.3** ASYMPTOTIC DISTRIBUTIONS WITH LOCALLY EXOGENOUS INSTRUMENTS. Suppose that the assumptions (2.3)-(2.18) hold, with  $b = b_0/\sqrt{T}$ ,  $\Pi = \Pi_0 \neq 0$  and  $\beta = \beta_0$ , where  $b_0$  and  $\beta_0$  are given vectors, and  $\Pi_0$  is a given matrix. If  $b_0 \notin \mathcal{N}(\Sigma_W)$ , then

$$AR(\beta_0) \xrightarrow{L} \frac{1}{k} \chi^2(k, \mu_1), \quad (4.10)$$

$$K(\beta_0) \xrightarrow{L} \chi^2(G, m'm) \quad \text{if } \text{rank}(\Sigma_{XY}) = G, \quad (4.11)$$

where

$$\mu_1 = \frac{1}{\sigma_e^2} b_0' \Sigma_W \Sigma_X^{-1} \Sigma_W b_0, \quad m = \frac{1}{\sigma_e} (\Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY})^{-1/2} \Sigma_{XY}' \Sigma_X^{-1} \Sigma_W b_0, \quad (4.12)$$

and  $\Sigma_X$ ,  $\Sigma_{XY}$ , and  $\Sigma_W$  are given in (2.11)-(2.27). If  $b_0 \in \mathcal{N}(\Sigma_W)$ , then

$$AR(\beta_0) \xrightarrow{L} \frac{1}{k} \chi^2(k), \quad (4.13)$$

$$K(\beta_0) \xrightarrow{L} \chi^2(G) \quad \text{if } \text{rank}(\Sigma_{XY}) = G. \quad (4.14)$$

**Theorem 4.4** ASYMPTOTIC DISTRIBUTIONS WITH WEAK LOCALLY EXOGENOUS INSTRUMENTS. Suppose that the assumptions (2.3)-(2.18) hold, with  $b = b_0/\sqrt{T}$ ,  $\Pi = \Pi_0/\sqrt{T}$  and  $\beta = \beta_0$ , where  $b_0$  and  $\beta_0$  are given vectors, and  $\Pi_0$  is a given matrix ( $\Pi_0 = 0$  is allowed). If  $b_0 \notin \mathcal{N}(\Sigma_W)$ , then

$$AR(\beta_0) \xrightarrow{L} \frac{1}{k} \chi^2(k, \mu_1), \quad (4.15)$$

$$K(\beta_0) \xrightarrow{L} \chi^2(G, \tilde{m}'\tilde{m}) \quad \text{if } \text{rank}(\Sigma_{WV}) = G, \quad (4.16)$$

where

$$\tilde{m} = \frac{1}{\sigma_e} (\Sigma_{WV}' \Sigma_X^{-1} \Sigma_{WV})^{-1/2} \Sigma_{WV}' \Sigma_X^{-1} \Sigma_W b_0, \quad (4.17)$$

and  $\Sigma_X$ ,  $\Sigma_{WV}$ ,  $\Sigma_W$  and  $\mu_1$  are defined in Theorem 4.3. If  $b_0 \in \mathcal{N}(\Sigma_W)$ , then

$$AR(\beta_0) \xrightarrow{L} \frac{1}{k} \chi^2(k), \quad (4.18)$$

$$K(\beta_0) \xrightarrow{L} \chi^2(G) \quad \text{if } \text{rank}(\Sigma_{WV}) = G. \quad (4.19)$$

We make the following remarks concerning Theorem 4.3 and Theorem 4.4. First, the endogeneity parameter  $b$  is local-to-zero, and for  $b_0 \in \mathcal{N}(\Sigma_W)$  the AR and K tests are asymptotically valid. However, unlike the AR test, note that the validity of the K test is established under an additional rank assumption (the case where this additional rank assumption fails is not covered in this paper). So, when  $b_0 \in \mathcal{N}(\Sigma_W)$ , the inference with locally exogenous instruments using the AR and K tests is feasible (at least in large samples). Second, if  $b_0 \notin \mathcal{N}(\Sigma_W)$ , the results in both theorems are different from those of Theorems 4.1 and 4.2 because the limiting distributions of both statistics

do not diverge. Third, even though the AR and K statistics have non-central chi-square limiting distributions when  $b_0 \notin \mathcal{N}(\Sigma_W)$ , they are not pivotal since the non-centrality parameters depend on nuisance parameters. In addition, the limiting distributions of both statistics cannot be bounded by any pivotal distribution.

It will be useful to see how the above theorems apply in a simple example.

**Example 4.1** Consider again model (2.29)-(2.30), which involves one endogenous explanatory variable and two instruments. If the matrix  $\Sigma_W$  is invertible, then  $\mathcal{N}(\Sigma_W) = \{0\}$ , and Theorem 4.1 entails that  $AR(\beta_0) \xrightarrow{L} +\infty$  under the null hypothesis  $\beta = \beta_0$ . Similarly, if  $\tilde{\Sigma}_{XY} \neq 0$ , then  $\text{rank}(\tilde{\Sigma}_{XY}) = G = 1$  and Theorem 4.2 entails that  $K(\beta_0) \xrightarrow{L} +\infty$  when  $\beta = \beta_0$ . If  $X_1$  is a valid instrument (with  $W_1 = 0$ ) and  $X_2 = W_2$  with  $W_2'W_2/T \xrightarrow{p} \sigma_{W_2}^2 > 0$ , we have

$$\Sigma_W = \begin{bmatrix} 0 & 0 \\ 0 & \sigma_{W_2}^2 \end{bmatrix} \quad (4.20)$$

which is a matrix of rank one, and  $\mathcal{N}(\Sigma_W) = \{(x_1, x_2)' : x_2 = 0\}$ . If  $b_2 = 0$ , then  $b_0 \in \mathcal{N}(\Sigma_W)$  and Theorem 4.1 entails that the asymptotic distribution given by (4.2) holds for  $AR(\beta_0)$ , while for  $K(\beta_0)$  part B of Theorem 4.2 is applicable. Of course, when  $b_0 = 0$ ,  $AR(\beta_0)$  follows the usual  $\chi^2(2)/2$  asymptotic distribution, while  $K(\beta_0)$  follows a  $\chi^2(1)$  distribution. For locally exogenous instruments, theorems 4.3 and 4.4 can be applied in a similar way.

## 5. Conclusion

In this paper, we have established conditions under which the AR and K tests are asymptotically valid even if some instruments used are endogenous. We have also showed that when these conditions fail, the limiting distributions of both statistics may diverge. Furthermore, when these conditions fail, under locally exogenous instruments setup, the limiting distributions of the statistics depend on nuisance parameters and cannot be bounded by any pivotal distribution. In consequence, the weak-instrument procedure proposed by Wang and Zivot (1998), the unified weak instruments framework of Swanson and Chao (2005) and the inference with imperfect instruments suggested by Ashley (2006) are not applicable. Overall, our results underscore the importance of checking for the presence of possibly invalid instruments when applying “identification-robust” tests. They also suggest that sensitivity analyses where different sets of instruments are considered [Ashley (2006), Small (2007)] can be quite useful for the interpretation of empirical results based on instrumental variables.

## A. Appendix: Proofs

PROOF OF THEOREM 4.1 Note first that

$$\frac{(y - Y\beta_0)'M_X(y - Y\beta_0)}{T - k} = \frac{u'u}{T - k} - \frac{T}{T - k} \left( \frac{u'X}{T} \right) \left( \frac{X'X}{T} \right)^{-1} \left( \frac{X'u}{T} \right) \quad (\text{A.1})$$

where, by the assumptions (2.3)-(2.18),

$$\frac{u'u}{T - k} \xrightarrow{p} \sigma_u^2 > 0, \quad \frac{X'X}{T} \xrightarrow{p} \Sigma_X > 0, \quad \frac{X'u}{T} = \frac{X'_0u}{T} + \frac{W'W}{T}b_0 + \frac{W'e}{T} \xrightarrow{p} \Sigma_W b_0, \quad (\text{A.2})$$

$$\left( \frac{u'X}{T} \right) \left( \frac{X'X}{T} \right)^{-1} \left( \frac{X'u}{T} \right) \xrightarrow{p} b'_0 \Sigma_W \Sigma_X^{-1} \Sigma_W b_0, \quad (\text{A.3})$$

$$\frac{(y - Y\beta_0)'M_X(y - Y\beta_0)}{T - k} \xrightarrow{p} \bar{\sigma}_u^2 = \sigma_u^2 - b'_0 \Sigma_W \Sigma_X^{-1} \Sigma_W b_0 \geq 0. \quad (\text{A.4})$$

(A) Suppose now that  $b_0 \notin \mathcal{N}(\Sigma_W)$ . Then  $b'_0 \Sigma_W \Sigma_X^{-1} \Sigma_W b_0 > 0$  and the numerator of the AR statistic diverges:

$$(y - Y\beta_0)'P_X(y - Y\beta_0) = T \left( \frac{u'X}{T} \right) \left( \frac{X'X}{T} \right)^{-1} \left( \frac{X'u}{T} \right) \xrightarrow{L} +\infty, \quad (\text{A.5})$$

hence

$$AR(\beta_0) \xrightarrow{L} +\infty. \quad (\text{A.6})$$

(B) If  $b_0 \in \mathcal{N}(\Sigma_W)$ , we have  $\Sigma_W b_0 = 0$  and  $\bar{\sigma}_u^2 = \sigma_u^2$ . Further,

$$X'u = X'(e + Wb_0) = X'e + X'Wb_0, \quad (\text{A.7})$$

$$\frac{1}{\sqrt{T}}X'u = \frac{1}{\sqrt{T}}[X'u - \Sigma_W b_0] = \frac{1}{\sqrt{T}}X'e + \frac{1}{\sqrt{T}}(X'W - \Sigma_W)b_0 \xrightarrow{L} S = S_e + S_b. \quad (\text{A.8})$$

Then,

$$(y - Y\beta_0)'P_X(y - Y\beta_0) = \left( \frac{u'X}{\sqrt{T}} \right) \left( \frac{X'X}{T} \right)^{-1} \left( \frac{X'u}{\sqrt{T}} \right) \xrightarrow{L} S' \Sigma_X^{-1} S, \quad (\text{A.9})$$

$$\frac{(y - Y\beta_0)'M_X(y - Y\beta_0)}{T - k} \xrightarrow{p} \sigma_u^2, \quad (\text{A.10})$$

hence

$$AR(\beta_0) \xrightarrow{L} \frac{1}{k\sigma_u^2} S' \Sigma_X^{-1} S. \quad (\text{A.11})$$

(C) Finally, if  $b_0 = 0$ , we have  $b_0 \in \mathcal{N}(\Sigma_W)$ , with the extra restrictions  $u = e$ ,  $\sigma_u^2 = \sigma_e^2$ ,

$$S = \frac{1}{\sqrt{T}}X'u = \frac{1}{\sqrt{T}}X'e \xrightarrow{L} N[0, \sigma_e^2 \Sigma_X],$$

hence

$$AR(\beta_0) \xrightarrow{L} \frac{1}{k\sigma_e^2} S_e' \Sigma_X^{-1} S_e \sim \frac{1}{k} \chi^2(k). \quad (\text{A.12})$$

□

**PROOF OF THEOREM 4.2** We note first, as in (A.1)-(A.4), that

$$S_{uu}(\beta_0) = \frac{(y - Y\beta_0)' M_X (y - Y\beta_0)}{T - k} \xrightarrow{p} \bar{\sigma}_u^2, \quad \frac{X'X}{T} \xrightarrow{p} \Sigma_X > 0, \quad \frac{X'u}{T} \xrightarrow{p} \Sigma_W b_0. \quad (\text{A.13})$$

(A) Suppose that  $b_0 \notin \mathcal{N}(\Sigma_W)$ . (i) Let  $\Pi = \Pi_0 \neq 0$ . Then, we have

$$S_{uV}(\beta_0) = \frac{1}{T - k} (y - Y\beta_0)' M_X Y \xrightarrow{p} q_{uV} = \delta' - b_0' \Sigma_W \Sigma_X^{-1} \Sigma_{WV}, \quad (\text{A.14})$$

$$\tilde{\Pi}(\beta_0) = \left( \frac{X'X}{T} \right)^{-1} \frac{X'Y}{T} - \left( \frac{X'X}{T} \right)^{-1} \frac{X'u}{T} \frac{S_{uV}(\beta_0)}{S_{uu}(\beta_0)} \xrightarrow{p} \Sigma_X^{-1} \tilde{\Sigma}_{XY}, \quad (\text{A.15})$$

where  $\tilde{\Sigma}_{XY} = \Sigma_{XY} - \Sigma_W b_0 (q_{uV} / \bar{\sigma}_u^2)$ , and

$$\frac{\tilde{Y}(\beta_0)' u}{T} = \tilde{\Pi}(\beta_0)' \frac{X'u}{T} \xrightarrow{p} \tilde{\Sigma}'_{XY} \Sigma_X^{-1} \Sigma_W b_0, \quad (\text{A.16})$$

$$\frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \xrightarrow{p} \tilde{\Sigma}'_{XY} \Sigma_X^{-1} \tilde{\Sigma}_{XY}. \quad (\text{A.17})$$

If  $\text{rank}(\tilde{\Sigma}_{XY}) = G$ , then  $\tilde{\Sigma}'_{XY} \Sigma_X^{-1} \tilde{\Sigma}_{XY} > 0$  and  $\Sigma_X^{-1} \tilde{\Sigma}_{XY} \Sigma_W b_0 \neq 0$  for  $b_0 \notin \mathcal{N}(\Sigma_W)$ , hence

$$\frac{u' \tilde{Y}(\beta_0)}{T} \left[ \frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \right]^{-1} \frac{\tilde{Y}(\beta_0)' u}{T} \xrightarrow{p} b_0' \Sigma_W \Sigma_X^{-1} \tilde{\Sigma}_{XY} (\tilde{\Sigma}'_{XY} \Sigma_X^{-1} \tilde{\Sigma}_{XY})^{-1} \tilde{\Sigma}'_{XY} \Sigma_X^{-1} \Sigma_W b_0 > 0.$$

Consequently, the numerator of the K statistic diverges:

$$(y - Y\beta_0)' P_{\tilde{Y}(\beta_0)} (y - Y\beta_0) = T \frac{u' \tilde{Y}(\beta_0)}{T} \left[ \frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \right]^{-1} \frac{\tilde{Y}(\beta_0)' u}{T} \xrightarrow{p} +\infty \quad (\text{A.18})$$

and

$$K(\beta_0) \xrightarrow{L} +\infty. \quad (\text{A.19})$$

(ii) Let  $\Pi = \Pi_0 / \sqrt{T}$ . Then

$$(y - Y\beta_0)' P_{\tilde{Y}(\beta_0)} (y - Y\beta_0) = T \frac{u' \tilde{Y}(\beta_0)}{T} \left[ \frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \right]^{-1} \frac{\tilde{Y}(\beta_0)' u}{T}, \quad (\text{A.20})$$

where

$$\frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \xrightarrow{L} \Sigma_{XY}^* \Sigma_X^{-1} \Sigma_{XY}^*, \quad \frac{\tilde{Y}(\beta_0)' u}{T} \xrightarrow{p} \Sigma_X^{-1} \Sigma_{XY}^* \Sigma_W b_0,$$

with  $\Sigma_{XY}^* = \Sigma_{WV} - \Sigma_W b_0 (q_{uV} / \bar{\sigma}_u^2)$ . If  $\text{rank}(\Sigma_{XY}^*) = G$ , then the numerator of the K statistic diverges, and  $K(\beta_0) \xrightarrow{L} +\infty$ .

(B) If  $b_0 \in \mathcal{N}(\Sigma_W)$ , we have  $\Sigma_W b_0 = 0$ ,  $\bar{\sigma}_u^2 = \sigma_u^2$  and  $\frac{1}{\sqrt{T}} X' u \xrightarrow{L} S = S_e + S_b$  as in (A.7)-(A.8).

(i) If  $\Pi = \Pi_0 \neq 0$ , we have when  $b_0 \in \mathcal{N}(\Sigma_W)$ , the denominator of the K statistic satisfies

$$\frac{1}{T} (y - Y\beta_0)' M_X (y - Y\beta_0) \xrightarrow{p} \sigma_u^2, \quad (\text{A.21})$$

while the denominator can be written

$$(y - Y\beta_0)' P_{\tilde{Y}(\beta_0)} (y - Y\beta_0) = \frac{u' X}{\sqrt{T}} \tilde{\Pi}(\beta_0) \left[ \frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \right]^{-1} \tilde{\Pi}(\beta_0)' \frac{X' u}{\sqrt{T}} \quad (\text{A.22})$$

where

$$\tilde{\Pi}(\beta_0) \xrightarrow{p} \Sigma_X^{-1} \Sigma_{XY}, \quad \frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \xrightarrow{p} \Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY}, \quad \frac{\tilde{Y}(\beta_0)' u}{\sqrt{T}} \xrightarrow{p} \Sigma_X^{-1} \Sigma_{XY} S. \quad (\text{A.23})$$

If  $\text{rank}(\Sigma_{XY}) = G$ , we have  $\Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY} > 0$ , hence

$$K(\beta_0) \xrightarrow{L} \frac{1}{\sigma_u^2} S' \Sigma_X^{-1} \Sigma_{XY} (\Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY})^{-1} \Sigma_{XY}' \Sigma_X^{-1} S. \quad (\text{A.24})$$

(ii) If  $\Pi = \Pi_0 / \sqrt{T}$ , the numerator of the K statistic is

$$(y - Y\beta_0)' P_{\tilde{Y}(\beta_0)} (y - Y\beta_0) = \frac{u' X}{\sqrt{T}} \tilde{\Pi}(\beta_0) \left[ \frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \right]^{-1} \tilde{\Pi}(\beta_0)' \frac{X' u}{\sqrt{T}}, \quad (\text{A.25})$$

hence

$$\tilde{\Pi}(\beta_0) \xrightarrow{p} \Sigma_X^{-1} \Sigma_{WV}, \quad \frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \xrightarrow{p} \Sigma_{WV}' \Sigma_X^{-1} \Sigma_{WV}, \quad \frac{\tilde{Y}(\beta_0)' u}{\sqrt{T}} \xrightarrow{p} \Sigma_X^{-1} \Sigma_{WV} S. \quad (\text{A.26})$$

If  $\text{rank}(\Sigma_{WV}) = G$ , then

$$K(\beta_0) \xrightarrow{L} \frac{1}{\sigma_u^2} S' \Sigma_X^{-1} \Sigma_{WV} (\Sigma_{WV}' \Sigma_X^{-1} \Sigma_{WV})^{-1} \Sigma_{WV}' \Sigma_X^{-1} S. \quad (\text{A.27})$$

(C) Finally, if  $b_0 = 0$ , we have  $b_0 \in \mathcal{N}(\Sigma_W)$ , with the extra restrictions  $u = e$ ,  $\sigma_u^2 = \sigma_e^2$ ,

$$S = \frac{1}{\sqrt{T}} X' u \xrightarrow{L} N[0, \sigma_e^2 \Sigma_X],$$

hence, if  $\Pi = \Pi_0 \neq 0$ ,

$$K(\beta_0) \xrightarrow{L} \frac{1}{\sigma_e^2} S_e' \Sigma_X^{-1} \Sigma_{XY} (\Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY})^{-1} \Sigma_{XY}' \Sigma_X^{-1} S_e \sim \chi^2(G), \quad (\text{A.28})$$

and if  $\Pi = \Pi_0/\sqrt{T}$  (where  $\Pi_0 = 0$  is allowed),

$$K(\beta_0) \xrightarrow{L} \frac{1}{\sigma_e^2} S_e' \Sigma_X^{-1} \Sigma_{WV} (\Sigma_{WV}' \Sigma_X^{-1} \Sigma_{WV})^{-1} \Sigma_{WV}' \Sigma_X^{-1} S_e \sim \chi^2(G). \quad (\text{A.29})$$

□

**PROOF OF THEOREM 4.3** Since  $b$  is now local-to-zero, we have

$$\frac{X'u}{\sqrt{T}} \xrightarrow{L} S_e + \Sigma_W b_0, \quad \frac{X'X}{T} \xrightarrow{p} \Sigma_X, \quad \frac{X'u}{T} \xrightarrow{p} 0, \quad \frac{(y - Y\beta_0)' M_X (y - Y\beta_0)}{T - k} \xrightarrow{p} \sigma_u^2 > 0. \quad (\text{A.30})$$

Further, we have

$$\begin{aligned} \frac{u'u}{T - k} &= \frac{(e + W \frac{b_0}{\sqrt{T}})' (e + W \frac{b_0}{\sqrt{T}})}{T - k} \\ &= \frac{e'e}{T - k} + \frac{b_0' W' e}{\sqrt{T}(T - k)} + \frac{e' W b_0}{\sqrt{T}(T - k)} + \frac{b_0' e' W b_0}{T(T - k)} \xrightarrow{p} \sigma_e^2 = \sigma_u^2. \end{aligned} \quad (\text{A.31})$$

(A) Let  $b_0 \notin \mathcal{N}(\Sigma_W)$ . Then,

$$AR(\beta_0) \xrightarrow{L} \frac{1}{k\sigma_e^2} (S_e + \Sigma_W b_0)' \Sigma_X^{-1} (S_e + \Sigma_W b_0) \sim \frac{1}{k} \chi^2(k, \mu_1) \quad (\text{A.32})$$

where  $\mu_1 = \frac{1}{\sigma_e^2} b_0' \Sigma_W \Sigma_X^{-1} \Sigma_W b_0 \neq 0$ . Similarly, we have  $\frac{\tilde{Y}(\beta_0)' \tilde{Y}(\beta_0)}{T} \xrightarrow{p} \Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY}$  and  $\frac{\tilde{Y}(\beta_0)' u}{\sqrt{T}} \xrightarrow{L} \Sigma_X^{-1} \Sigma_{XY} (S_e + \Sigma_W b_0)$ . So, if  $\text{rank}(\Sigma_{XY}) = G$ , we have

$$K(\beta_0) \xrightarrow{L} \frac{1}{\sigma_e^2} (S_e + \Sigma_W b_0)' \Sigma_X^{-1} \Sigma_{XY} (\Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY})^{-1} \Sigma_{XY}' \Sigma_X^{-1} (S_e + \Sigma_W b_0) \sim \chi^2(G, m'm) \quad (\text{A.33})$$

where  $m = \frac{1}{\sigma_e} (\Sigma_{XY}' \Sigma_X^{-1} \Sigma_{XY})^{-1/2} \Sigma_{XY}' \Sigma_X^{-1} \Sigma_W b_0 \neq 0$ .

(B) If  $b_0 \in \mathcal{N}(\Sigma_W)$ , we have  $\Sigma_W b_0 = 0$ . Then  $\mu_1 = 0$  and  $m = 0$ , hence  $AR(\beta_0) \xrightarrow{L} \frac{1}{k} \chi^2(k)$  and  $K(\beta_0) \xrightarrow{L} \chi^2(G)$ . □

**PROOF OF THEOREM 4.4** The proof of Theorem 4.3 for the AR statistic covers Theorem 4.4. The proof for the K statistic is similar to the one in Theorem 4.3. □

## References

- Anderson, T. W., Rubin, H., 1949. Estimation of the parameters of a single equation in a complete system of stochastic equations. *Annals of Mathematical Statistics* 20(1), 46–63.
- Ashley, R., 2006. Assessing the credibility of instrumental variables inference with imperfect instruments via sensitivity analysis. Technical report, Economics Department, Virginia Polytechnic Institute Blacksburg, Virginia.
- Bekker, P., 1994. Alternative approximations to the distributions of instrumental variable estimators. *Econometrica* 62, 657–681.
- Donald, S. G., Newey, W. K., 2001. Choosing the number of instruments. *Econometrica* 69, 1161–1191.
- Dufour, J.-M., 1997. Some impossibility theorems in econometrics, with applications to structural and dynamic models. *Econometrica* 65, 1365–1389.
- Dufour, J.-M., 2003. Identification, weak instruments and statistical inference in econometrics. *Canadian Journal of Economics* 36(4), 767–808.
- Dufour, J.-M., Hsiao, C., 2008. Identification. In: L. E. Blume, S. N. Durlauf, eds, *The New Palgrave Dictionary of Economics* 2nd edn. Palgrave Macmillan, Basingstoke, Hampshire, England. forthcoming.
- Dufour, J.-M., Jasiak, J., 2001. Finite sample limited information inference methods for structural equations and models with generated regressors. *International Economic Review* 42, 815–843.
- Dufour, J.-M., Taamouti, M., 2003. Point-optimal instruments and generalized Anderson-Rubin procedures for nonlinear models. Technical report, C.R.D.E., Université de Montréal.
- Dufour, J.-M., Taamouti, M., 2005. Projection-based statistical inference in linear structural models with possibly weak instruments. *Econometrica* 73(4), 1351–1365.
- Dufour, J.-M., Taamouti, M., 2007. Further results on projection-based inference in IV regressions with weak, collinear or missing instruments. *Journal of Econometrics* 139(1), 133–153.
- Hahn, J., Hausman, J. A., 2003. IV estimation with valid and invalid instruments. Technical report, Department of Economics, Massachusetts Institute of Technology Cambridge. Massachusetts.
- Hall, A. R., Peixe, F. P. M., 2003. A consistent method for the selection of relevant instruments. *Econometric Reviews* 2(3), 269–287.
- Hall, A. R., Rudebusch, G. D., Wilcox, D. W., 1996. Judging instrument relevance in instrumental variables estimation. *International Economic Review* 37, 283–298.
- Kadane, J. B., Anderson, T. W., 1977. A comment on the test of overidentifying restrictions. *Econometrica* 45(4), 1027–1031.

- Kivet, J. F., Niemczyk, J., 2006. On the limiting and empirical distribution of IV estimators when some of the instruments are invalid. Technical report, Department of Quantitative Economics, University of Amsterdam Amsterdam, The Netherlands.
- Kleibergen, F., 2002. Pivotal statistics for testing structural parameters in instrumental variables regression. *Econometrica* 70(5), 1781–1803.
- Moreira, M. J., 2003. A conditional likelihood ratio test for structural models. *Econometrica* 71(4), 1027–1048.
- Phillips, P. C. B., 1989. Partially identified econometric models. *Econometric Theory* 5, 181–240.
- Sargan, J. D., 1958. The estimation of economic relationships using instrumental variables. *Econometrica* 26(3), 393–415.
- Small, D. S., 2007. Sensitivity analysis for instrumental variables regression with overidentifying restrictions. *Journal of the American Statistical Association* 102(479), 1049–1058.
- Staiger, D., Stock, J. H., 1997. Instrumental variables regression with weak instruments. *Econometrica* 65(3), 557–586.
- Stock, J. H., Wright, J. H., Yogo, M., 2002. A survey of weak instruments and weak identification in generalized method of moments. *Journal of Business and Economic Statistics* 20(4), 518–529.
- Swanson, N. R., Chao, J. C., 2005. Notes and comments: Consistent estimation with a large number of weak instruments. *Econometrica* 73, 1673–1692.
- Wang, J., Zivot, E., 1998. Inference on structural parameters in instrumental variables regression with weak instruments. *Econometrica* 66(6), 1389–1404.