



Midwest

Federal Coordinating Lead Author

Chris Swanston

USDA Forest Service

Chapter Lead

Jim Angel

Prairie Research Institute, University of Illinois

Chapter Authors

Barbara Mayes Boustead

National Oceanic and Atmospheric Administration

Kathryn C. Conlon

Centers for Disease Control and Prevention

Kimberly R. Hall

The Nature Conservancy

Jenna L. Jorns

University of Michigan, Great Lakes Integrated Sciences and Assessments

Kenneth E. Kunkel

North Carolina State University

Maria Carmen Lemos

University of Michigan, Great Lakes Integrated Sciences and Assessments

Brent Lofgren

National Oceanic and Atmospheric Administration

Todd A. Ontl

USDA Northern Forests Climate Hub

John Posey

East West Gateway Council of Governments

Kim Stone

Great Lakes Indian Fish and Wildlife Commission (through January 2018)

Eugene Takle

Iowa State University

Dennis Todey

USDA Midwest Climate Hub

Review Editor

Thomas Bonnot

University of Missouri

Technical Contributors are listed at the end of the chapter.

Recommended Citation for Chapter

Angel, J., C. Swanston, B.M. Boustead, K.C. Conlon, K.R. Hall, J.L. Jorns, K.E. Kunkel, M.C. Lemos, B. Lofgren, T.A. Ontl, J. Posey, K. Stone, G. Takle, and D. Todey, 2018: Midwest. In *Impacts, Risks, and Adaptation in the United States: Fourth National Climate Assessment, Volume II* [Reidmiller, D.R., C.W. Avery, D.R. Easterling, K.E. Kunkel, K.L.M. Lewis, T.K. Maycock, and B.C. Stewart (eds.)]. U.S. Global Change Research Program, Washington, DC, USA, pp. 872–940. doi: [10.7930/NCA4.2018.CH21](https://doi.org/10.7930/NCA4.2018.CH21)

On the Web: <https://nca2018.globalchange.gov/chapter/midwest>



Key Message 1

Carson, Wisconsin

Agriculture

The Midwest is a major producer of a wide range of food and animal feed for national consumption and international trade. Increases in warm-season absolute humidity and precipitation have eroded soils, created favorable conditions for pests and pathogens, and degraded the quality of stored grain. Projected changes in precipitation, coupled with rising extreme temperatures before mid-century, will reduce Midwest agricultural productivity to levels of the 1980s without major technological advances.

Key Message 2

Forestry

Midwest forests provide numerous economic and ecological benefits, yet threats from a changing climate are interacting with existing stressors such as invasive species and pests to increase tree mortality and reduce forest productivity. Without adaptive actions, these interactions will result in the loss of economically and culturally important tree species such as paper birch and black ash and are expected to lead to the conversion of some forests to other forest types or even to non-forested ecosystems by the end of the century. Land managers are beginning to manage risk in forests by increasing diversity and selecting for tree species adapted to a range of projected conditions.

Key Message 3

Biodiversity and Ecosystems

The ecosystems of the Midwest support a diverse array of native species and provide people with essential services such as water purification, flood control, resource provision, crop pollination, and recreational opportunities. Species and ecosystems, including the important freshwater resources of the Great Lakes, are typically most at risk when climate stressors, like temperature increases, interact with land-use change, habitat loss, pollution, nutrient inputs, and nonnative invasive species. Restoration of natural systems, increases in the use of green infrastructure, and targeted conservation efforts, especially of wetland systems, can help protect people and nature from climate change impacts.

Key Message 4

Human Health

Climate change is expected to worsen existing health conditions and introduce new health threats by increasing the frequency and intensity of poor air quality days, extreme high temperature events, and heavy rainfalls; extending pollen seasons; and modifying the distribution of disease-carrying pests and insects. By mid-century, the region is projected to experience substantial, yet avoidable, loss of life, worsened health conditions, and economic impacts estimated in the billions of dollars as a result of these changes. Improved basic health services and increased public health measures—including surveillance and monitoring—can prevent or reduce these impacts.

Key Message 5

Transportation and Infrastructure

Storm water management systems, transportation networks, and other critical infrastructure are already experiencing impacts from changing precipitation patterns and elevated flood risks. Green infrastructure is reducing some of the negative impacts by using plants and open space to absorb storm water. The annual cost of adapting urban storm water systems to more frequent and severe storms is projected to exceed \$500 million for the Midwest by the end of the century.

Key Message 6

Community Vulnerability and Adaptation

At-risk communities in the Midwest are becoming more vulnerable to climate change impacts such as flooding, drought, and increases in urban heat islands. Tribal nations are especially vulnerable because of their reliance on threatened natural resources for their cultural, subsistence, and economic needs. Integrating climate adaptation into planning processes offers an opportunity to better manage climate risks now. Developing knowledge for decision-making in cooperation with vulnerable communities and tribal nations will help to build adaptive capacity and increase resilience.

Executive Summary



The Midwest is home to over 60 million people, and its active economy represents 18% of the U.S. gross domestic product.¹ The region is probably best known for agricultural production.

Increases in growing-season temperature in the Midwest are projected to be the largest contributing factor to declines in the productivity of U.S. agriculture.² Increases in humidity in spring through mid-century^{3,4} are expected to increase rainfall, which will increase the potential for soil erosion^{5,6} and further reduce planting-season workdays due to waterlogged soil.⁷

Forests are a defining characteristic of many landscapes within the Midwest, covering more than 91 million acres. However, a changing climate, including an increased frequency of late-growing-season drought conditions, is worsening the effects of invasive species, insect pests, and plant disease as trees experience periodic moisture stress. Impacts from human activities, such as logging, fire suppression, and agricultural expansion, have lowered the diversity of the Midwest's forests from the pre-Euro-American settlement period.

Natural resource managers are taking steps to address these issues by increasing the diversity of trees and introducing species suitable for a changing climate.⁸

The Great Lakes play a central role in the Midwest and provide an abundant freshwater resource for water supplies, industry, shipping, fishing, and recreation, as well as a rich and diverse ecosystem. These important ecosystems are under stress from pollution, nutrient and sediment inputs from agricultural systems, and invasive species.^{9,10} Lake surface temperatures are increasing,^{11,12} lake ice cover is declining,^{12,13,14} the seasonal stratification of temperatures in the lakes is occurring earlier in the year,¹⁵ and summer evaporation rates are increasing.^{13,16} Increasing storm impacts and declines in coastal water quality can put coastal communities at risk. While several coastal communities have expressed willingness to integrate climate action into planning efforts, access to useful climate information and limited human and financial resources constrain municipal action.

Land conversion, and a wide range of other stressors, has already greatly reduced biodiversity in many of the region's prairies, wetlands, forests, and freshwater systems. Species are already responding to changes that have

occurred over the last several decades,^{17,18,19} and rapid climate change over the next century is expected to cause or further amplify stress in many species and ecological systems in the Midwest.^{20,21,22} The loss of species and the degradation of ecosystems have the potential to reduce or eliminate essential ecological services such as flood control, water purification, and crop pollination, thus reducing the potential for society to successfully adapt to ongoing changes. However, understanding these relationships also highlights important climate adaptation strategies. For example, restoring systems like wetlands and forested floodplains and implementing agricultural best management strategies that increase vegetative cover (cover crops and riparian buffers) can help reduce flooding risks and protect water quality.^{23,24,25}

Midwestern populations are already experiencing adverse health impacts from climate change, and these impacts are expected to worsen in the future.^{26,27} In the absence of

mitigation, ground-level ozone concentrations are projected to increase across most of the Midwest, resulting in an additional 200–550 premature deaths in the region per year by 2050.²⁸ Exposure to high temperatures impacts workers' health, safety, and productivity.²⁹ Currently, days over 100°F in Chicago are rare. However, they could become increasingly more common by late century in both the lower and higher scenarios (RCP4.5 and RCP8.5).

The Midwest also has vibrant manufacturing, retail, recreation/tourism, and service sectors. The region's highways, railroads, airports, and navigable rivers are major modes for commerce activity. Increasing precipitation, especially heavy rain events, has increased the overall flood risk, causing disruption to transportation and damage to property and infrastructure. Increasing use of green infrastructure (including nature-based approaches, such as wetland restoration, and innovations like permeable pavements) and better engineering practices are beginning to address these issues.



Conservation Practices Reduce Impact of Heavy Rains

Integrating strips of native prairie vegetation into row crops has been shown to reduce sediment and nutrient loss from fields, as well as improve biodiversity and the delivery of ecosystem services.³³ Iowa State University's STRIPS program is actively conducting research into this agricultural conservation practice.³⁴ The inset shows a close-up example of a prairie vegetation strip. *From Figure 21.2 (Photo credits: [main photo] Lynn Betts, [inset] Farnaz Kordbacheh).*

Citizens and stakeholders value their health and the well-being of their communities—all of which are at risk from increased flooding, increased heat, and lower air and water quality under a changing climate.^{30,31} To better prevent and respond to these impacts, scholars and

practitioners highlight the need to engage in risk-driven approaches that not only focus on assessing vulnerabilities but also include effective planning and implementation of adaptation options.³²



The photo shows Menominee Tribal Enterprises staff creating opportunity from adversity by replanting a forest opening caused by oak wilt disease with a diverse array of tree and understory plant species that are expected to fare better under future climate conditions. *From Figure 21.4 (Photo credit: Kristen Schmitt).*

Background

The Midwest is home to more than 60 million people, and its active economy represents 18% of the U.S. gross domestic product.¹ In this report, the Midwest covers Illinois, Indiana, Iowa, Michigan, Minnesota, Missouri, Ohio, and Wisconsin. The region is probably best known for agricultural production. Trends toward warmer, wetter, and more humid conditions provide challenges for field work, increase disease and pest pressure, and reduce yields to an extent that these challenges can be only partially overcome by technology.³⁵ The Midwest contains large tracts of federal, state, and private forests and preserves that provide significant economic and ecological benefits to the region. However, as a changing climate results in shifting precipitation patterns, altered disturbance regimes, and increased frequency of late-growing-season moisture stress, the effects of existing stressors such as invasive species, insect pests, and plant disease are amplified.³⁶ Natural resource managers are taking steps to address these issues by increasing the diversity of trees and introducing species suitable for a changing climate.⁸

The Midwest also has vibrant manufacturing, retail, recreation/tourism, and service sectors. The region's highways, railroads, airports, and navigable rivers are major modes for commercial activity. Increasing precipitation, especially heavy rain events, has increased the overall flood risk, causing disruption to transportation and damage to property and infrastructure (e.g., Winters et al. 2015³⁷). Increasing use of green infrastructure (including nature-based approaches, such as wetland restoration, and innovations like permeable pavements) and better engineering practices are beginning to address these issues (e.g., City of Chicago 2015³⁸).

Tourism and outdoor recreation are major economic activities that may be affected by climate change, particularly in coastal towns that are at risk from algal bloom impacts and in areas that host winter sports that are especially vulnerable to warming winters. For example, ice fishing was limited due to mild temperatures in the winters of 2015–2016 and 2016–2017, and the American Birkebeiner cross-country ski race in Wisconsin was cancelled due to a lack of snow in February 2017. Portions of Michigan, Wisconsin, and Minnesota contain ceded territory of many tribes, and these are used for hunting, fishing, and gathering native plants, all of which play vital roles in maintaining cultural heritage. Projected changes in climate and ecosystems will have strong impacts on these activities.³⁹

The Great Lakes play a central role in the Midwest and provide an abundant freshwater resource for water supplies, industry, shipping, fishing, and recreation, as well as a rich and diverse ecosystem. The same can be said for the upper Mississippi, lower Missouri, Illinois, and Ohio River systems. Episodes of widespread heavy rains in recent years have led to flooding, soil erosion, and water quality issues from nutrient runoff into those systems.¹⁰ Land managers are beginning to change some of their practices (such as increasing the use of cover crops) to better manage excess surface water.⁴⁰

Citizens and stakeholders in the Midwest value their health and the well-being of their communities—all of which are at risk from increased flooding, increased heat, and lower air and water quality under a changing climate.^{30,31}

Energy in the Midwest

The Midwest is a major consumer of coal. In 2015, coal provided 56% of the electricity consumed in the region, and the eight states in the region accounted for 32% of the Nation's coal consumption (in BTUs). Coal's share of electricity production is declining in the Midwest, following the national trend (Ch. 4: Energy, Figure 4.3). In 2008, coal accounted for more than 70% of electricity consumption in the Midwest. Wind power is a small but growing source of electricity for the region. Iowa leads the Nation in per capita consumption of wind power, with wind providing over 30% of the state's electrical needs in 2015.⁴¹

Renewable energy is expanding in the Midwest. As part of a campus-wide initiative to transition to renewable energy sources, in 2017, Michigan State University established five solar carports that have an estimated annual production of 15,000 megawatt hours, representing about 5% of electricity use on campus (Figure 21.1). In addition to reducing carbon emissions, this investment is expected to save the university \$10 million over 25 years.⁴²



Solar Charging Stations

Figure 21.1: Solar carports were recently installed on the Michigan State University campus. Photo credit: David Rothstein.

What Is New in NCA4

Two new Key Messages are introduced (Key Messages 3 and 6). Key Message 3 recognizes the important role that ecosystems of the Midwest play in supporting a diverse array of species and providing important benefits such as flood control, crop pollination, and outdoor recreation. Key Message 6 addresses how at-risk communities in the Midwest are becoming more vulnerable to climate change impacts and how they are working to build adaptive capacity. Tribal nations are especially vulnerable because of their reliance on threatened natural resources for their cultural, subsistence, and economic needs. The four remaining Key Messages address improvements in the understanding of risks and responses to climate change since NCA3. Key Message 1 on agriculture provides more specificity about the risk to agriculture by stating that agricultural productivity (the ratio of outputs to inputs) is projected to decline by 2050 to levels of the 1980s (that is, yields may increase but at the cost of substantial increases in inputs). Key Message 2 on forestry illustrates the progress foresters and land managers have made in climate adaptation through their efforts to incorporate climate change risks into management decision-making. Key Message 5 on transportation and infrastructure highlights a growing interest in green infrastructure—the use of plants and open space in storm water management—as an option for adapting to more frequent episodes of extreme precipitation. Finally, Key Message 4 on human health identifies specific health impacts by naming expected changes in magnitude and occurrence of extreme events, exposures, and economic impacts. The message explicitly states public health actions that can be implemented to avoid or reduce the health impacts.

Key Message 1

Agriculture

The Midwest is a major producer of a wide range of food and animal feed for national consumption and international trade. Increases in warm-season absolute humidity and precipitation have eroded soils, created favorable conditions for pests and pathogens, and degraded the quality of stored grain. Projected changes in precipitation, coupled with rising extreme temperatures before mid-century, will reduce Midwest agricultural productivity to levels of the 1980s without major technological advances.

Recent Agriculturally Important Trends

The two main commodity crops in the Midwest are corn and soybeans, which are grown on 75% of the arable land. Wheat and oats are important crops grown on fewer acres. An increasing number of niche but higher-value crops (such as apples, grapes, cherries, cranberries, blueberries, and pumpkins) also are grown in the region.⁴³

Over the past 30 years, increased rainfall from April to June has been the most impactful climate trend for agriculture in the Midwest,³ providing a favorable supply of soil moisture while also reducing flexibility for timing of spring planting and increasing soil erosion.⁴⁴ In addition, wet conditions at the end of the growing season can create elevated levels of mold, fungus, and toxins.⁴⁵ The last spring frost has occurred earlier, causing the frost-free season to increase by an average of nine days since 1901.⁴⁶ However, daily maximum temperatures in summer in the Midwest have not followed the upward global trend, in part due to higher early summer rainfall on deep, water-holding soils,⁴⁷ thereby avoiding plant stress detrimental to crops. The avoidance of

heat stress and longer growing seasons have favored production in some parts of and some years in the Midwest.

Daily minimum temperatures have increased in all seasons due to increasing humidity.^{48,49} Elevated growing-season minimum daily temperatures are considered a factor in reducing grain weight in corn due to increased nighttime plant respiration.⁵⁰ Warming winters have increased the survival and reproduction of existing insect pests⁵¹ and already are enabling a northward range expansion of new insect pests and crop pathogens into the Midwest.⁵²

A contributing factor underpinning Midwest growing-season trends in both temperature and precipitation is the increase in water vapor (absolute humidity):^{49,53} higher humidity decreases the day–night temperature range and increases warm-season precipitation. Rising humidity also leads to longer dew periods and high moisture conditions that favor many agricultural pests and pathogens for both growing plants and stored grain.

Projected Trends and Agricultural Impacts

Warm-season temperatures are projected to increase more in the Midwest than any other region of the United States.⁵⁴ The frost-free season is projected to increase 10 days by early this century (2016–2045), 20 days by mid-century (2036–2065), and possibly a month by late century (2070–2099) compared to the period 1976–2005 according to the higher scenario (RCP8.5).⁴⁶

By the middle of this century (2036–2065), 1 year out of 10 is projected to have a 5-day period that is an average of 13°F warmer than a comparable period at the end of last century (1976–2005).⁵⁴ Current average annual 5-day maximum temperature values range from about 88°F in Northern Minnesota to 97°F in Southern Missouri. Tables 21.1 and 21.2 show

that by mid-century under the higher scenario (RCP8.5), 5-day maximum temperatures are projected to have moved further above optimum conditions for many crops and closer to the reproductive failure temperature, especially for corn in the southern half of the Midwest. Higher growing-season temperatures also shorten phenological stages in crops (for example, the grain fill period for corn).^{35,50} Under these temperatures, overall yield trends will be reduced because of periodic pollination failures and reduced grain fill during other years.

Increases in humidity in spring through mid-century^{3,4} are expected to increase rainfall, which will increase the potential for soil erosion^{5,6} and further reduce planting-season workdays due to waterlogged soil.⁷ As an example, for the Cedar River Basin in Iowa, the 100-year flood (1% chance of occurring in a given year) of the 20th century is projected to be a 25-year flood (4% chance per year) in the 21st century,⁵⁵ with associated increased frequency of flooding of agricultural land.

Increased spring precipitation and higher temperatures and humidity are expected to increase the number and intensity of fungus and disease outbreaks^{56,57} and the prevalence of bacterial plant diseases,⁵⁸ such as bacterial spot in pumpkin and squash.⁵⁹ Increased precipitation and soil moisture in a warmer climate also lead to increased loss of soil carbon⁶⁰ and degraded surface water quality due to loss of soil particles and nutrients.^{61,62} Transitions from extremes of drought to floods, in particular, increase nitrogen levels in rivers⁶³ and lead to harmful algal blooms.

Current understanding of drought in the Midwest is that human activity has not been a major component in historical droughts, and it remains uncertain how droughts will behave in the future. However, future projections show that Midwest surface soil moisture likely will transition from excessive levels in spring due to increased precipitation to insufficient levels in summer driven by higher temperatures, causing more moisture to be lost through evaporation.⁶⁴

Average Annual 5-Day Maximum Temperature

Geographic Area	Modeled Historical (1976–2005)	Mid-21st Century (2036–2065) for Lower Scenario (RCP4.5)	Mid-21st Century (2036–2065) for Higher Scenario (RCP8.5)
Northern Minnesota	88°F	93°F	95°F
Southern Missouri	97°F	102°F	103°F

Table 21.1: These modeled historical and projected average annual 5-day maximum temperatures illustrate the temperature increases projected for the middle of this century across the Midwest. Sources: NOAA NCEI and CICS-NC.

Optimum and Failure Temperatures for Vegetative Growth and Reproduction

Crop	Optimum Growth	Failure for Growth	Optimum Reproduction	Failure for Reproduction
Corn	80°F	105°F	67°F	95°F
Soybean	86°F	101°F	72°F	102°F

Table 21.2: This table shows the temperatures at which corn and soybeans reach optimum growth and reproduction as well as the temperatures at which growth and reproduction fail.⁵⁰

Projections of mid-century yields of commodity crops^{65,66} show declines of 5% to over 25% below extrapolated trends broadly across the region for corn (also known as maize) and more than 25% for soybeans in the southern half of the region, with possible increases in yield in the northern half of the region. Increases in growing-season temperature in the Midwest are projected to be the largest contributing factor to declines in the productivity of U.S. agriculture.² In particular, heat stress in maize during the reproductive period is projected by crop models to reduce yields in the second half of the 21st century.⁶⁷ These losses may be mitigated by enhanced photosynthesis and reduced crop water use, although the magnitude is uncertain.^{68,69} Elevated atmospheric CO₂ is expected to partially, but not completely, offset yield declines caused by climate extremes, with effects on soybeans less than on maize.⁷⁰

Non-commodity crops produced in the Midwest include tree fruits, sweet corn, and vegetables for farmers markets and canning. While the general impacts of climate change on specialty crops are similar to commodity crops, the more intense heat waves, excessive rain interspersed with drought, and higher humidity of a future climate likely will degrade market quality as well as yield by mid-century.⁷¹ Although data on climate-related losses are sparse, excess moisture is emerging as a major cause of crop loss.⁷² Wild rice is an annual plant harvested by tribes and others in shallow wetlands of northern Minnesota, Wisconsin, and Michigan. Stable production depends on a stable climate that maintains ecosystem diversity. Declines in production are expected, related to increases in climate extremes and climate-related disease and pest outbreaks as well as northward shifts of favorable growing regions.⁷³

Longer growing seasons and the introduction of hoop buildings (low, translucent, fabric-covered structures that protect plants from extreme weather) have allowed local growers of annual vegetable crops to extend the fresh produce season. However, unsheltered perennial crops such as tree fruits may be subjected increasingly to untimely budbreak followed by cold pulses due to earlier and longer occurrences of warm conditions in late winter.

Most animal agriculture in the region is in confinement, rather than range-based without shelter, and therefore offers an opportunity for mitigating some of the effects of climate change. Without adaptive actions, breeding success and production of milk and eggs will be reduced due to projected temperature extremes by mid-century.^{74,75,76}

Adaptation

Soil-erosion suppression methods in row-crop agriculture subjected to more intense rains include use of cover crops, grassed waterways, water management systems, contour farming, and prairie strips.^{6,40} More diversity in planting dates, pollination periods, chemical use, and crop and cultivar selection reduces vulnerability of overall production to specific climate extremes or the changes in pests and pathogens that they cause.

An example of a highly successful program is the Iowa State Science-based Trials of Row-crops Integrated with Prairie Strips (STRIPS) program that demonstrates that replacing 10 percent of cropland with prairie grasses reduced sediment loss 20-fold while total nitrogen concentrations were 3.3 times lower (Figure 21.2).³³ An example of a private-public response is the National Corn Growers Association's Soil Health Partnership (SHP),⁷⁷ a network of working farms across the Midwest



Conservation Practices Reduce Impact of Heavy Rains

Figure 21.2: Integrating strips of native prairie vegetation into row crops has been shown to reduce sediment and nutrient loss from fields, as well as improve biodiversity and the delivery of ecosystem services.³³ Iowa State University's STRIPS program is actively conducting research into this agricultural conservation practice.³⁴ The inset shows a close-up example of a prairie vegetation strip. Photo credits: (main photo) Lynn Betts, (inset) Farnaz Kordbacheh.

engaged in refining techniques for growing cover crops, implementing conservation tillage, and using science-based nutrient management to reduce erosion and nutrient loss while increasing organic matter.

Acreage under irrigation has expanded modestly since 2002,⁷⁸ mostly in the northern part of the Midwest where coarse soils of lower water-holding capacity are more vulnerable to drying under increased temperature. No strategies currently are available for maintaining historical trends in commodity agriculture production to cope with increases in spring rainfall and summer heat waves projected for mid-century.^{2,65}

Key Message 2

Forestry

Midwest forests provide numerous economic and ecological benefits, yet threats from a changing climate are interacting with existing stressors such as invasive species and pests to increase tree mortality and reduce forest productivity. Without adaptive actions, these interactions will result in the loss of economically and culturally important tree species such as paper birch and black ash and are expected to lead to the conversion of some forests to other forest types or even to non-forested ecosystems by the end of the century. Land managers are beginning to manage risk in forests by increasing diversity and selecting for tree species adapted to a range of projected conditions.


Forests are a defining characteristic of many landscapes within the Midwest, covering more than 91 million acres. From the oak–hickory forests of the Missouri Ozarks to the northern hardwood forests of the Upper Midwest, forest ecosystems sustain the people and communities within the region by providing numerous ecological, economic, and cultural benefits. The economic output of the Midwest forestry sector totals around \$122 billion per year.^{79,80,81,82,83,84,85,86} Forest-related recreation such as hunting, fishing, hiking, skiing, camping, wildlife watching, off-highway vehicles, and many other pursuits add to the region's economy. For example, forest-based recreationists spend approximately \$2.5 billion (in 1996 dollars) within Wisconsin communities.⁸⁷ Forests are fundamental to cultural and spiritual practices within tribal communities, supporting plants and animals of central cultural importance and providing food and resources for making items such as baskets, canoes, and shelters.⁸⁸

Climate change is anticipated to have a pervasive influence on forests within this region over the coming decades.^{36,89,90,91,92,93,94} Tree growth rates and forest productivity have benefited from longer growing seasons and higher atmospheric carbon dioxide concentrations, but continued benefits are expected only if adequate moisture and nutrients are available to support enhanced growth rates.⁹⁵ As growing-season temperatures rise, reduced tree growth^{96,97} or widespread tree mortality⁹⁸ is expected as the frequency of drought stress increases from drier air (as a result of increases in vapor pressure deficit [VPD]; Figure 21.3) and changing patterns of precipitation. Greater tree mortality from increased VPD likely will be particularly evident where competition for water is high in dense stands of trees^{99,100} or where forests naturally transition to grasslands due to limited soil moisture.¹⁰¹ Late-growing-season heat- and drought-related vegetation

stress is projected to shift the composition and structure of forests in the region¹⁰² by increasing mortality of younger trees, which are sensitive to drought.¹⁹ Warming winters will reduce snowpack that acts to insulate soil from freezing temperatures, increasing frost damage to shallow tree roots¹⁰³ and reducing tree regeneration.¹⁰⁴ Additionally, increases in existing biological stressors of forests are expected as temperatures rise. Effects of insect pests and tree pathogens are anticipated to intensify as winters warm, increasing winter survival of pests and allowing expansion into new regions.^{105,106} Changing climate conditions and atmospheric carbon dioxide concentrations will likely favor invasive plant species over native species, potentially decreasing tree regeneration.^{107,108} Overall, the increasing stress on trees from rising temperatures, drought, and frost damage raises the susceptibility of individual trees to the negative impacts from invasive plants, insect pests, and disease agents (Ch. 6: Forests, Figure 6.1).^{109,110,111}

Impacts from human activities such as logging, fire suppression, and agricultural expansion have lowered the diversity of the Midwest's forests from the pre-Euro-American settlement period. The forest types that occur within the region have been altered significantly relative to presettlement forests, with greater homogeneity in tree species composition across existing forest types.¹¹² Changes in modern forest types also include reduced structural complexity and less diverse mixes of tree species and tree ages.¹¹³ Forests with reduced diversity are at an increased risk of negative effects from climate change, because the potential for tree species or age classes that are resistant to impacts from biological stressors and climate change is reduced.⁹³ Forests composed of trees of similar size and age or with lower tree diversity are at increased risk of widespread mortality^{114,115} or declines in productivity.¹¹⁶ In many midwestern forests, fire suppression has decreased the prevalence of

Drying Effect of Warmer Air on Plants and Soils



Projected Increases in Vapor Pressure Deficit




Figure 21.3: As air temperature increases in a warming climate, vapor pressure deficit (VPD) is projected to increase. VPD is the difference between how much moisture is in the air and the amount of moisture in the air at saturation (at 100% relative humidity). Increased VPD has a drying effect on plants and soils, as moisture transpires (from plants) and evaporates (from soil) into the air. (a) Cooler air can maintain less water as vapor, putting less demand for moisture on plants, while warmer air can maintain more water as vapor, putting more demand for moisture on plants. (b, c) The maps show the percent change in the moisture deficit of the air based on the projected maximum 5-day VPD by the late 21st century (2070–2099) compared to 1976–2005 for (b) lower and (c) higher scenarios (RCP4.5 and RCP8.5). Sources: U.S. Forest Service, NOAA NCEI, and CICS-NC.

the drought-tolerant tree species, such as oak, hickory, and pine, while increasing the abundance of species with higher moisture requirements, such as maples.^{89,117} This results in greater risk of declines in forest health and productivity as the frequency of drought conditions increases.^{118,119}

Changes in climate and other stressors are projected to result in changes in major forest types and changes in forest composition as tree species at the northern limits of their ranges decline and southern species experience increasingly suitable habitat.¹²⁰ However, the fragmentation of midwestern forests and

the flatness of the terrain raise the possibility that the ranges of particular tree species will not be able to shift to future suitable habitats within the Midwest.¹²¹ For example, to reach areas 1.8°F (1°C) cooler, species in flat terrain must move up to 90 miles (150 km) north to reach cooler habitat, whereas species in mountainous terrain can shift higher in altitude over less latitudinal (north–south) distance.¹²² These changes raise the possibility of future losses of economic and cultural benefits of forests due to conversion to different forest types or the change to non-forest ecosystems.^{119,123,124}

Projected shifts in forest composition in the central hardwood region (southern Missouri, Illinois, Indiana, and Ohio) by the end of the century under a higher scenario (RCP8.5) would result in substantial declines in wildlife habitat and reduce economic value of timber in the region by up to \$788 billion (in 2015 dollars).¹²⁵

Changing climate conditions increasingly cause both cultural and economic impacts within the Midwest, and it is very likely these impacts will worsen in the future. For example, many tree species on which tribes depend for their culture and livelihoods—such as paper birch, northern white cedar, and quaking aspen—are highly vulnerable due to temperature increases.^{90,91,92,126} Populations of the emerald ash borer, a destructive invasive insect pest that attacks native ash trees, will increase due to warming winters in the region. Mortality of black ash trees, which are important for traditional basket-making for many tribes, is highly likely as winter temperatures continue to rise.¹²⁷

Warming winters already have economic impacts on the forest industry, as well. Forest operations (for example, site access, tree harvesting, and product transport) in many northern regions are conducted on snowpack or frozen ground to protect the site from negative impacts such as soil disturbance

and compaction,¹²⁸ but the timing of suitable conditions has become shorter and more variable. In the Upper Midwest, the duration of frozen ground conditions suitable for winter harvest has been shortened by 2 to 3 weeks in the past 70 years.¹²⁹ The contraction of winter snow cover and frozen ground conditions has increased seasonal restrictions on forest operations in these areas,¹³⁰ with resulting economic impacts to both forestry industry and woodland landowners through reduced timber values.¹³¹

Forestry professionals in the Midwest increasingly are considering the risks to forests from climate change¹³² and are responding by incorporating climate adaptation into land management.⁸ There are a growing number of examples of climate adaptation in forest management developed by more than 150 organizations that have participated in the Climate Change Response Framework, an approach to climate change adaptation led by the U.S. Forest Service.^{133,134,135} Management actions intended to maintain healthy and productive forests in a changing climate include a diverse suite of actions¹³⁵ but largely focus on activities that enhance species and structural diversity of existing forest communities and on management approaches that aim to increase the prevalence of species that are better suited to future climatic conditions.⁸ Forest management on tribal lands and ceded territory within the region increasingly integrates Scientific Ecological Knowledge of natural resource management with Traditional Ecological Knowledge, a highly localized, place-based system of knowledge learned and observed over many generations.¹³⁶ This integration can inform the co-creation of approaches to climate adaptation important for maintaining healthy, functioning forests that continue to provide cultural and spiritual benefits (see Case Study “Adaptation in Forestry”).

Case Study: Adaptation in Forestry

The Menominee Forest is well known as an exemplary forest; for generations, the Menominee Tribe has pioneered practices that have preserved nearly 220,000 acres with numerous species and varied habitats while maximizing the sustainable production of forest products. However, climate change—along with invasive species and insect pests and diseases—is creating new challenges for maintaining these diverse habitats and the sustainable supply of timber.

In response to tree mortality caused by oak wilt disease, an introduced exotic disease first identified in 1944 in Wisconsin, foresters at Menominee Tribal Enterprises (MTE) have integrated climate change adaptation into reforestation activities on severely disturbed areas created by the disease.¹³⁴ Using science guided by Traditional Ecological Knowledge of forest communities, forest openings created by oak wilt disease were replanted with a diverse array of tree and understory plant species that are expected to fare better under future climate conditions. Many of these species tolerate late-growing-season heat- and drought-related stress, while also providing important cultural benefits to the tribe such as food and medicine. The selection of locally collected plants and seeds used for restoring the oak wilt-affected openings combined scientific information on the future habitat of tree species with Indigenous knowledge of the forest communities necessary for guiding the development of diverse and healthy forests.



Figure 21.4: The photo shows Menominee Tribal Enterprises staff creating opportunity from adversity by replanting a forest opening caused by oak wilt disease with a diverse array of tree and understory plant species that are expected to fare better under future climate conditions. Photo credit: Kristen Schmitt.

The grass, plant, and shrub species are put together to strengthen the immune system of the deep-rooted trees. We tried to emphasize the underground biotic community within these openings. A healthy underground community ensures a healthy aboveground community. The shrubs hold the key to a healthy change of species within the local plant communities.

—MTE forester and tribal member

Key Message 3

Biodiversity and Ecosystems

The ecosystems of the Midwest support a diverse array of native species and provide people with essential services such as water purification, flood control, resource provision, crop pollination, and recreational opportunities. Species and ecosystems, including the important freshwater resources of the Great Lakes, are typically most at risk when climate stressors, like temperature increases, interact with land-use change, habitat loss, pollution, nutrient inputs, and nonnative invasive species. Restoration of natural systems, increases in the use of green infrastructure, and targeted conservation efforts, especially of wetland systems, can help protect people and nature from climate change impacts.

Species already are responding to environmental changes that have occurred over the last several decades,^{17,18,19} and rapid climate change over the next century is expected to cause or further amplify stress in many species and ecological systems in the Midwest.^{20,21,22} Land conversion and a wide range of other stressors have already greatly reduced biodiversity in many of the region's prairies, wetlands, forests, and freshwater systems. High rates of change in climate factors like air and water temperature and increasing drought risk likely will accelerate the rate of species declines and extinctions.^{18,137} The Midwest region supports the world's largest freshwater ecosystem, the Great Lakes, which are at risk from rising temperatures, changes in seasonal stratification of lake temperatures, and increased summer evaporation rates, combined with stresses from pollution, nutrient inputs that promote harmful algal blooms, and invasive species (Box 21.1).

The loss of species and degradation of ecosystems have the potential to reduce or eliminate essential ecological services such as flood control, water purification, and crop pollination, thus reducing the potential for society to successfully adapt to ongoing changes.

Observations, ecological theory, experimental studies, and predictive models provide insights into how shifts in several climate factors (temperature, precipitation patterns, humidity, and moisture stress) may interact over the next several decades.^{120,138,139} Vulnerability assessments for species and ecosystems quickly become complex, as species in the same ecosystem may have different climate sensitivities, and interactions with land-use change and other factors can strongly influence the level of impact (Ch. 5: Land Changes, KM 2; Ch. 17: Complex Systems, KM 1). Local expertise, input from multiple stakeholders, and tools like scenario planning can help improve assessment of vulnerability so that risks can be connected to management actions.^{132,140} Changes observed in the Midwest include species range shifts (avoiding exposure to new climatic conditions by shifting location), changes in population size (indicating a change in viability in a given place), shifts in body size and growth rates, and changes in the timing of seasonal events (phenology). Since the Third National Climate Assessment,²⁷ the number of studies documenting these types of changes has continued to grow. For example, climate change appears to have contributed to the apparent local extinction of populations of the Federally Endangered Karner blue butterfly at sites in the southern end of its range in northern Indiana, despite active management and extensive habitat restoration efforts. While climate change cannot be singled out as the only cause, the populations disappeared following multiple years of warming conditions and a very early onset of spring in 2012.¹³⁹ New evidence of shifting ranges comes from

Wisconsin forests, where a set of 78 understory plant species sampled in the 1950s and again in the 2000s have demonstrated shifts in their abundance centroids (a measure of the distribution and local abundance of populations) of about 30 miles ($49 \text{ km} \pm 29 \text{ km}$) over this 50-year period (Figure 21.5).¹⁴¹ The dominant direction of this shift was to the northwest, which matches the direction of change in important climatic conditions associated with the distributions of these species. While this shift suggests the potential for successful adaptation to changing conditions, the rate of change for most species was much less than the amount of change in the climate metrics over the same time period, raising the concern that the climate is changing too fast for these species to keep up.¹⁴¹ Similarly, a study of shifts in the timing of spring green-up, an indicator of when plant-feeding insects emerge, and the timing of migratory bird arrivals found that while both are shifting earlier in the Midwest, the arrival of birds is not advancing as quickly as the plants.¹⁴² Risks to birds from this mismatch in phenology include the potential for birds to arrive after food availability has peaked or for later arrivals to be less able to compete for territories or mates. Land protection and management strategies that help maintain or increase phenological variation of plants within key migratory and breeding habitats like

the Great Lakes coastlines may help increase the odds that birds can find the resources they need.¹⁴³

The drivers of changes in species ranges or abundance can be complex and difficult to detect until key thresholds are crossed. For example, in the Midwest region, cool- and coldwater fishes in inland lakes are particularly susceptible to changes in climate because habitat with appropriate temperatures and oxygen concentrations is often limited during summer months. In lakes at the southern (warmer) end of their ranges, these fish experience a squeezing of available habitat during summer months as the water near the lake surface becomes too warm and the dissolved oxygen levels in deeper waters drop (Figure 21.6).^{144,145,146} This “invisible” loss of habitat is driven by increases in water temperatures, longer duration of the stratified period (which delays the mixing of oxygen-rich water into the deeper waters), and declines in ice cover.^{147,148,149,150} Recent research has identified fish kill events tied to temperature and oxygen stress from increased air temperatures, and modeling results forecast increased numbers of these events, likely leading to local extinction of cool- and coldwater fish species in some lakes and reduced geographic distribution across the Midwest.^{151,152,153,154}

Climate Change Outpaces Plants' Ability to Shift Habitat Range




Figure 21.5: While midwestern species, such as understory plants in Wisconsin, are showing changes in range, they may not be shifting quickly enough to keep up with changes in climate. The panels here represent 78 plant species, showing (a) observed changes in the center of plant species abundances (centroids) from the 1950s to 2000s, (b) the direction and magnitude of changes in climate factors associated with those species, and (c) the lag, or difference, between where the species centroid is now located and where the change in climate factors suggests it should be located in order to keep pace with a changing climate. Source: adapted from Ash et al. 2017.¹⁴¹ ©John Wiley & Sons, Ltd.

Coldwater Fish at Risk




Figure 21.6: The graphic shows the oxythermal (oxygen and temperature) habitat of coldwater fish in midwestern inland lakes, illustrated by water depth under (left) a typical ice-free period and (right) a warm ice-free period (right). The top plots show water temperatures during the ice-free period, and the bottom plots show the dissolved oxygen concentrations. The schematics at the bottom illustrate the area of the lake that is ideal habitat for coldwater fish (in blue) and areas that represent water outside of the temperature or dissolved oxygen limit (in yellow and red, respectively). The left plots show how available habitat “squeezes” during a typical year, while the right plots illustrate a complete loss of suitable habitat during very warm years. Source: Madeline Magee, University of Wisconsin.

Taken individually, responses like range shifts, changes in local abundance, or changes in phenology may indicate that a species is successfully adapting to new conditions, or conversely may indicate a species is under stress. The extent to which responses indicate risk and the challenge of attributing changes to climate drivers when systems are exposed to many additional stressors are important sources of uncertainty that likely slow progress on climate change adaptation within the resource management sector.^{155,156} Further, while evidence of species- and ecosystem-level responses to direct climate change impacts is increasing, many of the most immediate risks are even more challenging to track, because they relate to climate-driven enhancement of existing stressors, such as habitat loss and degradation, pollution, the spread of invasive

species, and drainage and irrigation practices in agricultural landscapes.^{138,157} As species are lost from midwestern ecosystems, there likely will be a net loss of biodiversity, as numerous additional stressors, especially widespread land conversion across the southern Midwest, limit opportunities for these gaps to be filled by species moving in from other regions (Ch. 7: Ecosystems, KM 1 and 2).^{158,159}

While movement of species from the south-central United States could help sustain species-diverse ecosystems as some of the Midwest’s current species move north, these range expansions can further stress current species. Many species and ecosystems in the Midwest, especially the Upper Midwest, are best suited to survive and compete for resources when winter conditions are harsh

and growing seasons are short. As winter warms and the growing season extends, species from the south-central United States, as well as species from outside the country that are more traditionally viewed as invasive species, are expected to be able to grow faster and take advantage of these changes, increasing the rate of loss of the region's native species.^{160,161} For invasive insect pests, these impacts may be compounded as extended growing seasons allow time for additional generations to be produced in a single season;¹⁶² the same mechanism can promote higher impacts from native insect pests, as well. Given that some native species will decline in the region, to maintain or increase species diversity, some managers are beginning to plan for and even promote some native plant species that are present in a region, but more common to the south, as conditions change. While these can be important strategies for maintaining diversity and ecosystem functions, especially in isolated habitats where inward migration is not likely, careful consideration of the source of plant stocks is important when seeking to avoid introducing new or more competitive genotypes.¹⁶³ Further, as some native species decline, managers will benefit from increased vigilance in keeping potential invasive species from outside of North America from gaining a foothold.

Declines in native pollinator species are another important concern in the Midwest, as both native and managed pollinator species (typically nonnative bee species) play vital roles in supporting food production and farmer livelihoods and are critical for supporting wild plant reproduction and the diversity of ecosystems.^{164,165} Key threats to this diverse group of insects, mammals, and birds include habitat loss and degradation, pathogens, pesticide use, and invasive species.^{164,165,166} Most native and agricultural crops that require a pollinator are pollinated by insects, and where information is available, declines in populations of pollinator

insects in the Midwest have primarily been linked to the expansion of intensive agriculture.^{167,168,169,170} In addition to habitat loss, climate change is likely to act as an added stressor for many species, through many different mechanisms.¹⁶⁴ Many insects may be limited by their ability to shift to new habitats as conditions change; for example, many bumble bee species are showing population declines at southern range edges but not expanding as quickly at northern range edges.¹⁷¹ It is likely that pollinators that specialize on one or a few species for some aspect of their life history will be particularly vulnerable.¹⁷² Within the Midwest, observed high rates of decline in the monarch butterfly,¹⁶⁷ which relies on milkweed species as a host plant, are the focus of a network of outreach and ambitious multi-partner conservation efforts that are helping raise awareness of pollinator declines and links between pollinators and habitat availability.¹⁷³ These efforts, boosted by research demonstrating that habitat restoration can help sustain pollinator populations,^{174,175} provide examples of how to help support the adaptation of this critical group of species.

Perhaps more than in any other region of the United States, human land use has influenced the structure and function of natural systems of the Midwest. Widespread conversion of natural systems to agriculture has changed much of the region's water and energy balance (Ch. 5: Land Changes, KM 1). When vegetation has been removed or undergoes a major change, runoff and flooding both tend to increase.^{24,176,177} As land has been cleared for agriculture and cities, it simultaneously has lost the capacity to store water due to the resulting conversion to pavement, compaction of soils, and widespread loss of wetlands. More than half of the region's wetlands have been drained (Ch. 22: N. Great Plains, Case Study "Wetlands and the Birds of the Prairie Pothole Region"); in states at the southern end of the region, fewer than

10%–15% of presettlement wetlands remained in the 1980s.¹⁷⁸ The growth of agriculture and loss of wetlands in the Midwest mean that changes to the timing, type (snow or rain), and amount of precipitation are acting on a system that is already highly altered in ways that tend to promote flooding.²⁴ Climate change modeling suggests that the southern half of the Midwest likely will see increases in saturated soils, which also indicates risks to agriculture and property from inundation and flooding;¹⁷⁹ recent work incorporating land-use

change and population changes also suggests the number of people at risk from flooding will increase across much of the Midwest.¹⁸⁰ However, understanding these relationships also highlights important climate adaptation strategies. For example, restoring systems like wetlands and forested floodplains and implementing agricultural best management strategies that increase vegetative cover (such as cover crops and riparian buffers) can help reduce flooding risks and protect water quality (Figure 21.7).^{23,24,25}



Wetland Restoration Projects Can Help Reduce Impacts

Figure 21.7: The Blausey Tract restoration project on the U.S. Fish and Wildlife Service’s Ottawa National Wildlife Refuge (Ohio) restored 100 acres of former Lake Erie coastal wetlands that were previously in row crop production. In addition to providing habitat for wildlife and fish, these wetlands help reduce climate change impacts by storing water from high-water events and by filtering nutrients and sediments out of water pumped from an adjacent farm ditch. This work was carried out by two conservation groups, The Nature Conservancy and Ducks Unlimited, in partnership with the U.S. Fish and Wildlife Service, and was funded by The Great Lakes Restoration Initiative.^{186,187} (top) Shown here is the Blausey Tract restoration site in early spring of 2011, prior to the restoration activities. (bottom) In the spring of 2013, just two years after the start of restoration, the site already was providing important habitat for wildlife and fish. Photo credits: (top) ©The Nature Conservancy, (bottom) Bill Stanley, ©The Nature Conservancy.

As the flooding risk example above illustrates, understanding both the history of change and how future climate patterns can drive additional changes is useful for identifying meaningful strategies for reducing risks to both people and biodiversity through strategically protecting and restoring ecosystems. Since the Third National Climate Assessment,²⁷ the recognition, promotion, and implementation of green or ecosystem-based climate change adaptation solutions have expanded. While the idea of using natural systems to reduce risks and provide benefits to society is not new, efforts to document and quantify benefits, costs, and costs savings (relative to hard, or “gray,” infrastructure) of these types of approaches are increasing.¹⁸¹ These approaches often help replace systems that

have been lost, such as Great Lakes coastal wetlands, prairies, and vegetated floodplains along rivers and streams that slow water flows and act as sponges that keep floodwaters from people, property, and infrastructure (Figure 21.7),^{182,183} or tree cover that increases shade and improves urban air quality.^{181,184} The important role of nature-based solutions like reforestation for mitigating climate change is also increasingly being recognized and quantified.¹⁸⁵ From the perspective of protecting the biodiversity of the Midwest, adaptation and mitigation strategies that incorporate protection or restoration of natural systems can be a great win-win approach, because they often add habitat and restore ecological and hydrological functions that were reduced as a result of land conversion.

Box 21.1: Focus on the Great Lakes

The Great Lakes contain 20% of the world’s surface freshwater, provide drinking water and livelihood to more than 35 million people,¹⁸⁸ and allow for important economic and cultural services such as shipping and recreation. The Great Lakes influence regional weather and climate conditions and impact climate variability and change across the region. The lakes influence daily weather by 1) moderating maximum and minimum temperatures of the region in all seasons, 2) increasing cloud cover and precipitation over and just downwind of the lakes during winter, and 3) decreasing summertime convective clouds and rainfall over the lakes.^{189,190} In recent decades, the Great Lakes have exhibited notable changes that are impacting and will continue to impact people and the environment within the region.¹⁹¹ In particular, lake surface temperatures are increasing,^{11,12} lake ice cover is declining,^{12,13,14} the seasonal stratification of temperatures in the lakes is occurring earlier in the year,¹⁵ and summer evaporation rates are increasing.^{13,16}

Along the Great Lakes, lake-effect snowfall has increased overall since the early 20th century. However, studies have shown that the increase has not been steady, and it generally peaked in the 1970s and early 1980s before decreasing.¹⁹³ As the warming in the Midwest continues, reductions in lake ice may increase the frequency of lake-effect snows until winters become so warm that snowfall events shift to rain.^{194,195}

Lake-surface temperatures increased during the period 1985–2009 in most lakes worldwide, including the Great Lakes.¹⁹⁶ The most rapid increases in lake-surface temperature occur during the summer and can greatly exceed temperature trends of air at locations surrounding the lakes.¹⁹⁷ From 1973 to 2010, ice cover on the Great Lakes declined an average of 71%;¹⁴ although ice cover was again high in the winters of 2014 and 2015,¹⁹² a continued decrease in ice cover is expected in the future.^{198,199}

Water levels in the Great Lakes fluctuate naturally, though levels more likely than not will decline with the changing climate.²⁰⁰ A period of low water levels persisted from 1998 to early 2013. A single warm winter in

Box 21.1: Focus on the Great Lakes, *continued*

The Changing Great Lakes




Figure 21.8: The duration of seasonal ice cover decreased in most areas of the Great Lakes between 1973 and 2013, while summer surface water temperature (SWT) increased in most areas between 1994 and 2013. (a) The map shows the rate of change in ice cover duration. The greatest rate of decrease in seasonal ice cover duration is seen near shorelines, with smaller rates occurring in the deeper central parts of Lakes Michigan and Ontario, which rarely have ice cover. (b) The map shows the rate of change in summer SWT. The greatest rates of increase in summer SWT occurred in deeper water, with smaller increases occurring near shorelines. Source: adapted from Mason et al. 2016.¹⁹² Used with permission from Springer.

1997–1998 (corresponding to a major El Niño event) and ongoing increases in sunlight reaching the lake surface (due to reduced cloud cover) were likely strong contributors to these low water levels.¹¹ Following this period, water levels rose rapidly. Between January 2013 and December 2014, Lake Superior’s water rose by about 2 feet (0.6 meters) and Lakes Michigan and Huron’s by about 3.3 feet (1.0 meter).²⁰¹ Recent projections with updated methods of lake levels for the next several decades under 64 global model-based climate change simulations (from the Coupled Model Intercomparison Project Phase 5, or CMIP5 database, using the RCP4.5, RCP6.0, and RCP8.5 scenarios) on average show small drops in water levels over the 21st century (approximately 6 inches for Lakes Michigan and Huron and less for the other lakes), with a wide range of uncertainty.²⁰⁰

An important seasonal event for biological activity in the Great Lakes is the turnover of water, or destratification, which historically has occurred twice per year. Destratification occurs during the fall as the water temperature drops below a threshold of 39°F, the point at which freshwater attains its maximum density, and again during the spring when the water temperature rises above that threshold. The resultant mixing carries oxygen down from the lake surface and nutrients up from the lake bottom and into the water column. In a pattern that is similar to changes in duration of the growing season on land, the climate projections suggest that the overturn in spring that triggers the start of the aquatic “growing season” will happen earlier, and the fall overturn will happen later.^{198,202} This trend toward a longer stratified season has been documented at locations in Lake Superior.^{197,203} As the duration of the stratified period increases, the risk of impacts from low oxygen levels at depth and a lack of nutrient inputs at the surface increases, potentially leading to population declines of species in both zones. As warming trends continue, it is possible that a full overturning may not occur each year.²⁰⁴ For example, lake surface temperatures failed to drop below the 39°F threshold during the winters of 2012 and 2017 in parts of southern Lake Michigan and Lake Ontario (see <https://coastwatch.glerl.noaa.gov/glsea/glsea.html>). When this lack of water mixing contributes to persistently low oxygen levels, the result may be reductions in the growth of phytoplankton (algae) and zooplankton (microscopic animals) that form the basis of aquatic food webs, potentially leading to cascading effects on the health and abundance of species across all levels of Great Lakes food webs.^{202,205,206}

Box 21.1: Focus on the Great Lakes, *continued*

Ecological impacts of climate change in the Great Lakes occur in the context of multiple stressors, as these important ecosystems are under stress from pollution, nutrient and sediment inputs from agricultural systems, and invasive species (Ch. 17: Complex Systems, KM 1).^{9,10} Human influence on habitats is another stressor. Examples include coastal wetland damage²⁰⁷ and disturbance by human structures that change habitat conditions and water flow patterns.²⁰⁸ Fish harvest and other management activities also have influences on populations.²⁰⁹ Especially in Lake Erie, runoff from agricultural watersheds can carry large volumes of nutrients and sediments that can reduce water quality, potentially leading to hypoxia (inadequate oxygen supply),^{210,211} an occurrence that is predicted to be more likely as the climate continues to change.¹⁰ Increased water temperatures and nutrient inputs also contribute to algal blooms, including harmful cyanobacterial algae that are toxic to people, pets, and many native species.^{212,213}

As with the inland lake fish described above (see Figure 21.6), climate change is expected to impact the species and fisheries of the Great Lakes.²¹⁴ However, the vast size and low temperatures in these lakes suggest that mortality events from temperature are a much lower risk. One key aspect of the influence of warming lakes on fish growth is the availability of suitable thermal habitat, as ectotherms, or cold-blooded species, can grow faster in warmer water due to temperature impacts on metabolic rates. Fish can behaviorally thermoregulate, meaning they can migrate to the portion of the water column that contains water of the particular species' preferred temperature.²¹⁵ Bottom-water temperatures in the deep parts of the lakes are expected to remain close to 39°F, while temperatures above the seasonal thermocline (the distinct temperature transition zone separating warmer surface waters from colder waters below) are expected to warm considerably.²⁰² This means that fish will be able to find habitats that favor higher growth rates for a longer period of time during the year. This same growth rate increase may occur for some species in smaller lakes, but the potential for exceeding critical thresholds is likely higher (Figure 21.6). If sufficient food is available, this will enhance the growth rates for economically important species like yellow perch and lake whitefish even though they are classed as cool-water and coldwater fishes, respectively.²¹⁶ It remains unclear, however, if a sufficient food supply will be available to sustain this increase in growth rates.

While some native fish may show enhanced growth, these same changes can influence the survival and growth of invasive species. Nonnative species such as alewife²¹⁷ and zebra and quagga mussels²¹⁸ have had dramatic impacts on the Great Lakes. Warmer conditions may lead to increases in invasion success and may increase the impact of invasive species that are already present. For example, sea lamprey are parasitic fish that are native to the Atlantic Ocean, and in the Great Lakes, they are the focus of several forms of control efforts.²¹⁹ Climate change has potential to reduce the effectiveness of these efforts. In the Lake Superior watershed, in years with longer growing seasons (defined as the number of days with water temperatures above 50°F), lamprey reach larger weights before spawning.¹⁶¹ Larger body sizes suggest a greater impact on other fish species, because larger lamprey produce more eggs and require more food to survive.¹⁶¹

Coastal communities and several economic sectors, including shipping, transportation, and tourism, are vulnerable to the aforementioned climate impacts (Ch. 8: Coastal, KM 1). While the most recent research²⁰⁰ underscores the great uncertainty in future lake levels, earlier research showed that scenarios of decreasing lake levels will increase shipping costs even if the shipping season is longer,²²⁰ or that lower ice cover could increase the damage to coastal infrastructure caused by winter storms.^{221,222} While several coastal communities have expressed willingness to integrate climate action into planning efforts, access to useful climate information and limited human and financial resources constrain municipal action. Producers and users of climate

Box 21.1: Focus on the Great Lakes, *continued*

information are working together to create customized climate information and resources, which increases trust and legitimacy, addressing this challenge (see Case Study “Great Lakes Climate Adaptation Network”). This has been demonstrated in projects, for instance, with marinas and harbors in Michigan, with ravine management in Illinois and Wisconsin, and with the Chicago Climate Action Plan in Illinois.^{223,224,225,226} Although many communities in the region are taking steps to incorporate climate change and related impacts into policy and planning decisions, many more may benefit from using their existing stakeholder networks to engage with producers of climate information and build upon lessons learned from leaders in the region.²²⁷

Key Message 4**Human Health**

Climate change is expected to worsen existing health conditions and introduce new health threats by increasing the frequency and intensity of poor air quality days, extreme high temperature events, and heavy rainfalls; extending pollen seasons; and modifying the distribution of disease-carrying pests and insects. By mid-century, the region is projected to experience substantial, yet avoidable, loss of life, worsened health conditions, and economic impacts estimated in the billions of dollars as a result of these changes. Improved basic health services and increased public health measures—including surveillance and monitoring—can prevent or reduce these impacts.

Climate change directly and indirectly impacts human health (Ch. 14: Human Health, KM 1). Midwestern populations are already experiencing adverse health impacts from climate change, and these impacts are expected to worsen in the future.^{26,27} The risks are especially high for people who are less able to cope because characteristics like age, income, or social connectivity make them more vulnerable.²²⁸

Air Quality

Degraded air quality impacts people living in the Midwest. Increases in ground-level ozone and particulate matter are associated with the prevalence of various lung and cardiovascular diseases, which can lead to missed school days, hospitalization, and premature death (Ch. 13: Air Quality, KM 1).^{26,28} Despite successful efforts to reduce particulate matter and ozone pollution, climate change could increase the frequency of meteorological conditions that lead to poor air quality.^{26,229} In the absence of mitigation, ground-level ozone concentrations are projected to increase across most of the Midwest, resulting in an additional 200 to 550 premature deaths in the region per year by 2050.²⁸ These account for almost half of the total projected deaths due to the climate-related increase in ground-level ozone nationwide and may cost an estimated \$4.7 billion (in 2015 dollars).²⁸

Pollen production has been on the rise in the Midwest in recent years, with pollen seasons starting earlier and lasting longer (Ch. 13: Air Quality, KM 3).^{28,230} People, particularly children, with asthma and other respiratory diseases are especially vulnerable to aeroallergens.²³¹ Aeroallergens can cause allergic rhinitis and exacerbate asthma and sinusitis.²³¹ Oak pollen may be responsible for an increase of 88 to 350 asthma-related emergency room visits by 2050 under the higher scenario (RCP8.5), with an estimated average annual cost ranging between \$43,000 and \$170,000 (in 2015 dollars).²⁸

Projected Changes in Ozone-Related Premature Deaths




Figure 21.9: Maps show county-level estimates for the change in average annual ozone-related premature deaths over the summer months in 2050 (2045–2055) and 2090 (2085–2095) compared to 2000 (1995–2005) under the lower and higher scenarios (RCP4.5 and RCP8.5) in the Midwest. The results represent the average of five global climate models. Source: adapted from EPA 2017.²⁸

Temperature

Increased daytime and nighttime temperatures are associated with heat-related diseases (for example, dehydration and heatstroke) and death in the Midwest.^{26,232} Extreme heat in urban centers like Chicago, St. Louis, Cincinnati, Minneapolis/St. Paul, Milwaukee, and Detroit can cause dangerous living conditions.^{26,232,233,234,235,236} High rates of heat-related illness also have been observed in rural populations,²³⁵ where occupational exposure to heat and access to care is a concern. Exposure to high temperatures impacts workers' health, safety, and productivity.²⁹

Future risk of heat-related disease could be significantly higher. As an example, Figure 21.10 shows the projected number of days over 100°F in Chicago over the 21st century using 32 models and two scenarios. Currently, days over 100°F in Chicago are rare. However, they could become increasingly more common in both the lower and higher scenarios (RCP4.5 and RCP8.5). The higher scenario (RCP8.5) yields a wider range and a higher number of days over 100°F than the lower scenario (RCP4.5), especially by 2070–2090. Near the upper end of the model results (95th percentile) at late-century, with the potential for almost 60 days per year

over 100°F, conditions could be more typical of present-day Las Vegas than Chicago. While the degree of uncertainty becomes larger further into the future, all model results show an increase in heat in the last two periods of the 21st century—changes that would pose a significant challenge to Chicago and other midwestern cities.

Compared to other regions where worsening heat is also expected to occur, the Midwest is projected to have the largest increase in extreme temperature-related premature deaths under the higher scenario (RCP8.5): by 2090, 2,000 additional premature deaths per year, compared to the base period of 1989–2000, are projected due to heat alone without adaptation efforts.²⁸ Northern midwestern communities and vulnerable populations (see Key Message 6) that historically have

not experienced high temperatures may be at risk for heat-related disease and death. Risk of death from extremely cold temperatures will decrease under most climate projection scenarios.²⁸

Unabated climate change will translate into costs among the workforce and in utility bills, potentially exacerbating existing health disparities among those most at risk. By 2050, increased temperatures under the higher scenario (RCP8.5) are estimated to cost around \$10 billion (in 2015 dollars) due to premature deaths and lost work hours.²⁸ Increased electricity demand is estimated to amount to \$1.2 billion by 2090 (in 2015 dollars).²⁸ For those who are chronically ill or reliant on electronic medical devices, the increased cost of electricity, which contributes to energy insecurity,²⁸ may introduce financial and health burdens.




Figure 21.10: This graph shows the annual number of days above 100°F in Chicago for the historical period of 1976–2005 (black dot) and projected throughout the 21st century under lower (RCP4.5, teal) and higher (RCP8.5, red) scenarios. Increases at the higher end of these ranges would pose major heat-related health problems for people in Chicago. As shown by the black dot, the average number of days per year above 100°F for 1976–2005 was essentially zero. By the end of the century (2070–2099), the projected number of these very hot days ranges from 1 to 23 per year under the lower scenario and 3 to 63 per year under the higher scenario. For the three future periods, the teal and red dots represent the model-weighted average for each scenario, while the vertical lines represent the range of values (5th to 95th percentile). Both scenarios show an increasing number of days over 100°F with time but increasing at a faster rate under the higher scenario. Sources: NOAA NCEI and CICS-NC.

Precipitation

An increase in localized extreme precipitation and storm events can lead to an increase in flooding.²⁷ River flooding in large rivers like the Mississippi, Ohio, and Missouri Rivers and their tributaries can flood surface streets and low-lying areas, resulting in drinking water contamination, evacuations, damage to buildings, injury, and death.²⁶ Flooded buildings can experience mold growth that can trigger asthma attacks and allergies during cleanup efforts.²³⁷ Mental stress following flooding events can cause substantial health impacts, including sleeplessness, anxiety, depression, and post-traumatic stress disorder.²³⁸ Similarly, drought has been identified as a slow-moving stressor that contributes to acute and chronic mental health impacts such as anxiety and depression.²³⁹

Precipitation events can transport pathogens that cause gastrointestinal illnesses, putting populations who rely on untreated groundwater (such as wells) at an increased risk of disease,²⁴⁰ particularly following large rainfall events.²⁴¹ Many midwestern communities use wells as their drinking water sources. Adaptive measures, such as water treatment installations, may substantially reduce the risk of gastrointestinal illness, in spite of climate change.²⁴⁰

Habitat Conditions

Climate-related changes in habitats (see Key Message 3) for disease-carrying insects like the mosquito found in the Midwest (*Culex pipiens* and *Culex tarsalis*) that transmits West Nile virus (WNV) and the blacklegged, or deer, tick (*Ixodes scapularis*) that transmits Lyme disease have been associated with higher rates of infection.^{242,243} Northern expansion of the *Culex* species in the Midwest is expected to result in upwards of 450 additional WNV cases above the 1995 baseline by 2090 absent greenhouse gas mitigation.²⁸

Harmful algal blooms (Box 21.1), such as one that occurred in August 2014 in Lake Erie, can introduce cyanobacteria into drinking and recreational water sources, resulting in restrictions on access and use.²⁸ Contact with and consumption of water contaminated with cyanobacteria have been associated with skin and eye irritation, respiratory illness, gastrointestinal illness, and liver and kidney damage.²⁶ The occurrence of conditions that encourage cyanobacteria growth, such as higher water temperatures, increased runoff, and nutrient-rich habitats, are projected to increase in the Midwest.²⁸

Challenges and Opportunities

Climate-sensitive health impacts are complex and dynamic. Coordination across public health, emergency preparedness, planning, and communication agencies can maximize outreach to the most at-risk populations while directing activities to reduce health disparities and impacts.²⁴⁴ Public health agencies in the Midwest have developed interdisciplinary communities of practice around climate and health adaptation efforts, effectively enhancing the resilience of the region's public health systems.^{244,245,246,247,248} Activities around increased surveillance of climate-sensitive exposures and disease are gaining momentum and interest among practitioners and researchers.^{249,250}

Actions tied to reducing contributions to global climate change can result in direct co-benefits related to health and other outcomes (such as economic development).²⁵¹ Reducing emissions related to energy production and transportation may involve changes to fuel sources, vehicle technology, land use, and infrastructure.²⁵¹ Active transportation, such as biking and walking, has been found to significantly decrease disease burden.^{252,253,254} A study of the 11 largest midwestern metropolitan areas estimated a health benefit of nearly 700 fewer deaths per year by swapping half of short trips

from car to bike.²⁵⁵ As Midwest Rust Belt metropolitan areas revitalize and reinvest, there are opportunities to prioritize active living to maximally reduce climate change drivers and improve health.

Key Message 5

Transportation and Infrastructure

Storm water management systems, transportation networks, and other critical infrastructure are already experiencing impacts from changing precipitation patterns and elevated flood risks. Green infrastructure is reducing some of the negative impacts by using plants and open space to absorb storm water. The annual cost of adapting urban storm water systems to more frequent and severe storms is projected to exceed \$500 million for the Midwest by the end of the century.

Climate change poses several challenges to transportation and storm water systems in the Midwest. Annual precipitation in the Midwest has increased by 5% to 15% from the first half of the last century (1901–1960) compared to present day (1986–2015).¹⁹³ Winter and spring precipitation are important to flood risk in the Midwest and are projected to increase by up to 30% by the end of this century. Heavy precipitation events in the Midwest have increased in frequency and intensity since 1901 and are projected to increase through this century.¹⁹³

There has been an increase in extreme precipitation events that overwhelm storm water sewage systems, disrupt transportation networks, and cause damage to infrastructure and property. Runoff from extreme precipitation events can exceed the capacity of storm water systems, resulting in property damage, including basement backups (Ch. 11: Urban,

KM 2).^{37,256} In addition, in metropolitan areas with older sewer systems that combine sanitary sewage with storm water, extreme rain can result in the release of raw sewage into rivers and streams, posing both health and ecological risks.²⁵⁷ These releases, known as combined sewer overflows (CSO), pose challenges to major sources of drinking water including the Mississippi River²⁵⁸ and the Great Lakes.^{259,260} On the Great Lakes, increases in CSO frequency and volume are projected under mid-high and higher scenarios (RCP6.0 and RCP8.5).²⁶¹ The U.S. Environmental Protection Agency (EPA) estimates that the cost of adapting urban storm water systems to handle more intense and frequent storms in the Midwest could exceed \$480 million per year (in 2015 dollars) by the end of the century under either the lower or higher scenario (RCP4.5 or RCP8.5).²⁸ Extreme precipitation events also affect transportation systems (Ch. 12: Transportation, KM 1). Heavy rainstorms can result in the temporary closure of roadways. In addition, faster streamflow caused by extreme precipitation can erode the bases of bridges, a condition known as scour. A study of six Iowa bridges deemed to be critical infrastructure found that under all emissions scenarios (in the Coupled Model Intercomparison Project Phase 3), each location was projected to have increased vulnerability from more frequent episodes of overtopping and potential scour.⁵⁵ The EPA estimates that the annual cost of maintaining current levels of service on midwestern bridges in the face of increased scour damage from climate change could reach approximately \$400 million in the year 2050 under either the lower or higher scenario (RCP4.5 or RCP8.5).²⁸

In addition to its impacts on infrastructure, heavy precipitation also affects the operation of roadways by reducing safety and capacity while increasing travel times (Ch. 12: Transportation, KM 1). Projected increases in the number of extreme precipitation events have

been linked to an increased risk of traffic crashes.²⁶² Intelligent Transportation Systems (ITS) use sensors and cameras to monitor road conditions. This allows for rapid deployment of emergency response vehicles and use of electronic signage to reroute traffic. Such systems allow transportation agencies to minimize the adverse impacts associated with extreme weather.²⁶³

Flooding on major rivers also poses a challenge to Midwest communities. Major river floods differ from flash floods on smaller streams in that they affect a larger area and require longer periods of heavy precipitation to create flood conditions. The Nation's two largest rivers, the Mississippi and the Missouri, flow through the Midwest. River floods can cause loss of life, as well as significant property damage. River floods have caused the closure of interstate highways in the Midwest and temporary inundation of secondary roads. During floods in May 2017, more than 400 state roads in Missouri were closed due to flooding, including several stretches of Interstate 44 (Figure 21.11).²⁶⁴ High water also disrupts barge traffic on the Mississippi River.^{265,266,267,268,269,270} Billion-dollar floods in the Midwest have occurred three times in the last quarter-century.²⁷¹ Climate projections suggest an increased risk of inland flooding under either the lower or higher scenario (RCP4.5 or RCP8.5). Average annual damages from heightened flooding risk in the Midwest are projected to be in excess of \$500 million (in 2015 dollars) by 2050.²⁸

Changes in temperature also can pose challenges to infrastructure. Extreme heat creates material stress on road pavements, bridge expansion joints, and railroad tracks. Milder winter temperatures, however, may be expected to partially offset these damages by reducing the amount of rutting caused by the freeze-thaw cycle. Even taking into account



River Flooding in the Midwest

Figure 21.11: This composite image shows portions of Interstate 44 near St. Louis that were closed by Meramec River flooding in both 2015 and 2017. The flooding shown here occurred in May 2017. Image credit: Surdex Corporation.

the benefits of milder winters for paved surfaces, the EPA estimates that higher temperatures associated with unmitigated climate change would result in approximately \$6 billion annually in added road maintenance costs and over \$1 billion in impacts to rail transportation by 2090 (in 2015 dollars).²⁸

Green infrastructure—the use of plants and open space to manage storm water—is helping communities in the Midwest become more resilient to challenges associated with heavy precipitation. At the site or neighborhood level, rain gardens and other planted landscape elements collect and filter rainwater in the soil, slowing runoff into sewer systems. Permeable pavements on parking lots allow water to be stored in the soil. Trees planted next to streets also provide important storm water management benefits. Larger-scale projects include preservation of wetlands. In addition to their storm water management benefits, some types of green infrastructure, such as urban trees and green roofs, contribute to climate change mitigation by acting as carbon sinks.^{272,273,274}

There are many examples of green infrastructure projects in the Midwest, though not all explicitly identify climate change as a rationale. The examples below enhance resilience to the heavy rains that are projected to become more frequent.

- The Cermak/Blue Island Sustainable Streetscape Project in the Pilsen neighborhood of Chicago uses bioswales, rain gardens, and permeable pavements to reduce up to 80% of storm water runoff. It also uses street trees and other vegetation to reduce the urban heat island effect while also providing an attractive public space.²⁷⁵
- The Metropolitan Sewer District in St. Louis has embarked upon a \$100 million rain-scaping project designed to divert storm water runoff in the northern portion of the City of St. Louis and adjacent north St. Louis County.²⁷⁶
- The City of Minneapolis uses street trees to reduce storm water runoff through enhanced evaporation and infiltration of water into the soil.²⁷⁷ The City of Cleveland also prioritizes tree planting as an adaptation strategy, with an emphasis on increasing the tree canopy in low-income neighborhoods. In addition to its storm water management benefits, urban forestry also reduces the urban heat island effect and acts as a carbon sink.²⁷⁸

At the scale of a metropolitan region, preservation and restoration of streams, floodplains, and watersheds are enhancing biodiversity while also reducing storm water runoff.

- *Open Space Preservation:* Many communities in the Midwest are recognizing that preservation of open space, particularly in floodplains, is a cost-effective method for

managing storm water. Ducks Unlimited, a non-profit organization, has purchased conservation easements that restrict future development on nearly 10,000 acres of floodplain around the confluence of the Mississippi and Missouri Rivers. In the Milwaukee area, the Ozaukee Washington Land Trust has preserved more than 6,000 acres of forests, wetlands, and open space through acquisitions and the purchase of conservation easements, preserving lands important for absorbing rainwater and filtering toxins from sediment.^{279,280}

- *Stream Restoration:* Several midwestern communities are turning to dechannelization (the removal of concrete linings placed in waterways) and daylighting (bringing back to the surface streams that had been previously buried in pipes) as methods of storm water management. The Milwaukee Metropolitan Sewerage District is currently undertaking a dechannelization of the Kinnickinnic River. According to the District, the concrete lining of the waterway actually makes the waterway more dangerous during heavy rain. Flooding motivated the City of Kalamazoo to daylight a 1,500-foot section of Arcadia Creek in the downtown district.^{281,282}
- *Ravine Restoration:* Lake Michigan's western shore in Wisconsin and northern Illinois holds more than 50 small watersheds, known locally as ravines. Storm water runoff subjects these ravines to serious erosion, which threatens property and infrastructure. The Great Lakes Alliance has produced guides to reduce erosion through best management practices, including stream buffers, use of native plants for stabilization, and reducing the steepness or gradient of the stream bank.²²³

Key Message 6

Community Vulnerability and Adaptation

At-risk communities in the Midwest are becoming more vulnerable to climate change impacts such as flooding, drought, and increases in urban heat islands. Tribal nations are especially vulnerable because of their reliance on threatened natural resources for their cultural, subsistence, and economic needs. Integrating climate adaptation into planning processes offers an opportunity to better manage climate risks now. Developing knowledge for decision-making in cooperation with vulnerable communities and tribal nations will help to build adaptive capacity and increase resilience.

Vulnerability and Adaptation

In the Midwest, negative impacts related to climate change are projected to affect human systems, including cities, rural and coastal communities, and tribes.^{28,283,284} Higher temperatures, increasing variation in precipitation patterns, and changes in lake levels are likely to increase the vulnerability of these systems to extreme events (including flooding, drought, heat waves, and more intense urban heat island effects), compounding already existing stressors such as economic downturns, shrinking cities, and deteriorating infrastructure.²⁸⁵ Extreme heat such as that experienced in July 2011 (with temperatures reaching over 100°F in the majority of the Midwest) is expected to intensify,²⁸⁶ and urban heat islands may cause hardships to those most vulnerable, such as the old and infirm and those without resources to control their microclimate (for example, through the use of air conditioning).²⁸⁷ Under the higher scenario (RCP8.5), extreme heat is

projected to result in losses in labor and associated losses in economic revenue up to \$9.8 billion per year in 2050 and rising to \$33 billion per year in 2090 (in 2015 dollars).²⁸ Expanding the use of green infrastructure and locating it properly may mitigate the negative impact of heat islands in urban settings (see Key Messages 4 and 5) (see also Ch. 11: Urban, KM 4).

To mitigate or better respond to these impacts, scholars and practitioners highlight the need to engage in risk-based approaches that not only focus on assessing vulnerabilities but also include effective planning and implementation of adaptation options (Ch. 28: Adaptation, KM 3).³² These place-based approaches actively rely on participatory methodologies to evaluate and manage risk and to monitor and evaluate adaptation actions.³² However, documented implementation of climate change planning and action in Midwest cities and rural communities remains low. For example, in 2015, only four counties and cities in the region—Marquette and Grand Rapids in Michigan and Dane County and Milwaukee in Wisconsin—had created formal climate adaptation plans, none of which have been implemented.²⁸⁸ Moreover, a recent study of 371 cities in the Great Lakes region found that only 36 of them could identify a climate entrepreneur, that is, a public official clearly associated with pushing for climate action.²⁸⁵ Attempts to assess vulnerabilities, especially for poor urban communities, face persisting environmental and social justice barriers, such as lack of participation and historical disenfranchisement,²⁸⁹ despite evidence that these communities are going to be disproportionately affected by climate impacts.²⁹⁰ Additionally, in-depth interviews with local decision-makers on water management across scales have suggested that a lack of political and financial support at the state and federal levels is a barrier to adaptation action in cities and counties.²⁹¹ While initiatives are underway in the Midwest to mainstream

adaptation action—that is, embed and integrate climate adaptation action in what cities already do (see Case Study “Great Lakes Climate Adaptation Network”) (see also Ch. 28: Adaptation,

KM 5)—there are few examples in the published literature that document failure or success (but see Kalafatis et al. 2015, Vogel et al. 2016^{292,293}).

Case Study: Great Lakes Climate Adaptation Network

The Great Lakes Climate Adaptation Network (GLCAN) is a regional, member-driven peer network of local government staff who work together to identify and act on the unique climate adaptation challenges of the Great Lakes region. GLCAN formed in 2015 as a regional network of the Urban Sustainability Directors’ Network (USDN) to unite Great Lakes cities with universities in the region. It has been cooperating actively with a regional climate organization, the Great Lakes Integrated Sciences and Assessments (GLISA), a NOAA-supported program housed at the University of Michigan and Michigan State University, to create climate information in support of decision-making in member cities. In this example of sustained engagement, GLCAN and GLISA work as a boundary chain that moves climate information from producers at the Universities to users in the cities, as well as across cities. This minimizes transaction costs, in terms of human and financial resources, while building trust and legitimacy.^{292,294} In one example of this partnership, with funding from USDN, GLCAN and GLISA worked with the Huron River Watershed Council and five Great Lakes cities (Ann Arbor, Dearborn, Evanston, Indianapolis, and Cleveland) to develop a universal vulnerability assessment template that mainstreams the adaptation planning process and results in the integration of climate-smart and equity-focused information into all types of city planning.²⁹⁵ The template is publicly available;²⁹⁶ its purpose is to reduce municipal workloads and save limited resources by mainstreaming existing, disparate planning domains (such as natural hazards, infrastructure, and climate action), regardless of city size or location. Based on this work, USDN funded a follow-up project for GLISA to work with additional Great Lakes and Mid-Atlantic cities and a nonprofit research group (Headwaters Economics) to develop a socioeconomic mapping tool for climate risk planning.

Linked Boundary Chain Model




Figure 21.12: Shown here is a configuration of the boundary chain employed in the Great Lakes Climate Adaptation Network (GLCAN) Case Study. The information is tailored and moves through different boundary organizations (links in the chain) to connect science to users. By co-creating information and pooling resources throughout the chain, trust and legitimacy are built and cost is decreased. Source: adapted from Lemos et al. 2014.²⁹⁴ ©American Meteorological Society.

In addition, work on estimating the cost of adaptation nationally and in the Midwest remains limited, though the EPA has estimated that the Midwest is among the regions with the largest expected damages to infrastructure, including the highest estimated damages to roads, rising from \$3.3 billion per year in 2050 to \$6 billion per year in 2090 (in 2015 dollars) under a higher scenario (RCP8.5), and highest number of vulnerable bridges (Key Message 5).²⁸ Additionally, economic models that value climate amenities—for example, offering residents the benefits of warmer winters or cooler summers—indicate that while the Midwest is among the regions with the largest predicted amenity loss, certain cities (such as Minneapolis and Minnesota) and subregions (such as upper Michigan) will be among the few places where the value of warmer winters outweighs the cost of hotter summers.^{297,298} Limited evidence indicates that household consideration of climate amenities may contribute to reversing long-standing trends in out-migration from the Midwest²⁹⁸ and that changes in national migration patterns will contribute to population growth in the region.²⁸ More research is needed to understand how cities in the Midwest might be affected by long-term migration to the region.³¹

Collaboratively Developing Knowledge and Building Adaptive Capacity

Interactions among producers of climate information (for example, universities and research institutes), end users (such as city planners, watershed managers, and natural resource managers), and intermediaries (for example, information brokers and organizations) play a critical role in increasing the integration and use of climate knowledge for adaptation.²⁹⁹ In the Midwest, organizations such as the Great Lakes Integrated Sciences and Assessments (GLISA; glisa.umich.edu) and the Wisconsin Initiative on Climate Impacts (wicci.wisc.edu), and research projects such as Useful to Usable (U2U), have created mechanisms and tools, such as climate scenarios, decision support tools, and climate

data, that promote the joint development of usable climate information across different types of stakeholders, including city officials, water managers, farmers, and tribal officials.^{224,294,300} For example, working closely with corn farmers and climate information intermediaries, including extension agents and crop consultants, in Iowa, Nebraska, Michigan, and Indiana, an interdisciplinary team of climate scientists, agronomists, computer scientists, and social scientists have not only created a suite of decision support tools (see Key Message 1) but also significantly advanced understanding of corn farmers' perceptions of climate change,³⁰¹ willingness to adapt,³⁰² and opportunities for and limitations of the use of climate information in the agricultural sector.^{294,303} Strategies being implemented as a result of these collaborations, including the use of green infrastructure and water conservation efforts, are proving effective at reducing sensitivity to the impacts of climate change in the Midwest.^{304,305,306} In addition, binational partnerships between the United States and Canada, in support of the Great Lakes Water Quality Agreement, synthesized annual climate trends and impacts for a general audience in a pilot product for 2017 to provide a timely and succinct summary in an easy-to-understand format (Ch. 16: International, KM 4).³⁰⁷ However, these organizations face challenges including the high costs in interacting with users, contextualizing and customizing climate information, and building trust.³⁰⁸ The development of new forms of sustained engagement likely would increase the use of climate information in the region.

Tribal Adaptation

Tribes and Indigenous communities in the Midwest have been among the first to feel the effects of climate change as it impacts their culture, sovereignty, health, economies, and ways of life.³⁹ The Midwest contains ceded territory—large swaths of land in Minnesota, Wisconsin, and Michigan in which Ojibwe tribes reserved hunting, fishing, and gathering

rights in treaties with the United States government.⁸⁸ Climate change presents challenges to the Ojibwe tribes in co-managing these resources with other land managers; as the climate changes, various species utilized by tribes are declining and may shift entirely outside of treaty boundaries and reserved lands.^{127,309,310} In certain tribal cultures, all beings (species) are important; climate adaptation efforts that favor certain beings at the detriment of others can be problematic. Adaptation to climate change might also mean giving up on something deeply embedded in tribal culture for which no substitute exists.³¹ A family sugarbush (a forest stand used for maple syrup), for example, cannot be replaced culturally, spiritually, or economically if the sugar maple range were to shift outside of treaty or reservation boundaries. As the effects of climate change become more pronounced, further research can shed light on how tribal nations are being affected.

Projected changes in climate, particularly increases in extreme precipitation events, will have pronounced impacts on tribal culture and tribal people in the Midwest.²⁸³ Reservations often are located in isolated rural communities, meaning emergency response to flooding presents challenges in getting help to tribal citizens. Additionally, in areas of the Midwest, infestations of the invasive emerald ash borer already are devastating ash tree populations and corresponding Indigenous cultural and economic traditions.¹²⁷

Across the United States, a number of tribal nations are developing adaptation plans, including in the Midwest (Ch. 15: Tribes, KM 3).²⁸³ These plans bring together climate data and projections with Traditional Ecological Knowledge^{311,312} of tribal members. Within Indigenous oral history lies a complex and rich documentation of local ecosystems—not found in books—that can be used to understand and document the changes that are occurring.³¹³ Climate change effects are not typically immediate or dramatic because they

occur over a relatively long period of time, but tribal elders and harvesters have been noticing changes, such as declining numbers of waabooz (snowshoe hare), many of which Scientific Ecological Knowledge has been slower to document. The Traditional Ecological Knowledge of elders and harvesters who have lived and subsisted in a particular ecosystem can provide a valuable and nuanced understanding of ecological conditions on a smaller, more localized scale. Integrating this Traditional Ecological Knowledge with Scientific Ecological Knowledge in climate change initiatives provides a more complete understanding of climate change impacts.¹³⁶ Community input to tribal adaptation plans ensures that Traditional Ecological Knowledge can be used to produce adaptation strategies trusted by community members.³¹⁴

Acknowledgments

Technical Contributors

Katherine Browne

University of Michigan

Melonee Montano

Great Lakes Indian Fish and Wildlife Commission

Hannah Panci

Great Lakes Indian Fish and Wildlife Commission

Jason Vargo

University of Wisconsin

Madeline R. Magee

University of Wisconsin–Madison

USGCRP Coordinators

Kristin Lewis

Senior Scientist

Allyza Lustig

Program Coordinator

Katie Reeves

Engagement and Communications Lead

Opening Image Credit

Carson, Wisconsin: © William Garrett/Flickr (CC BY 2.0). Adaptation: cropped top and bottom to conform to the size needed for publication.

Traceable Accounts

Process Description

The chapter lead authors were identified in October 2016, and the author team was recruited in October and November 2016. Authors were selected for their interest and expertise in areas critical to the Midwest with an eye on diversity in expertise, level of experience, and gender. The writing team engaged in conference calls starting in December 2016, and calls continued on a regular basis to discuss technical and logistical issues related to the chapter. The Midwest chapter hosted an engagement workshop on March 1, 2017, with the hub in Chicago and satellite meetings in Iowa, Indiana, Michigan, and Wisconsin. The authors also considered other outreach with stakeholders, inputs provided in the public call for technical material, and incorporated the available recent scientific literature to write the chapter. Additional technical authors were added as needed to fill in the gaps in knowledge.

Discussion amongst the team members, along with reference to the Third National Climate Assessment and conversations with stakeholders, led to the development of six Key Messages based on key economic activities, ecology, human health, and the vulnerability of communities. In addition, care was taken to consider the concerns of tribal nations in the northern states of the Midwest. The Great Lakes were singled out as a special case study based on the feedback of the engagement workshop and the interests of other regional and sector chapters.

Note on regional modeling uncertainties

Interaction between the lakes and the atmosphere in the Great Lakes region (e.g., through ice cover, evaporation rates, moisture transport, and modified pressure gradients) is crucial to simulating the region's future climate (i.e., changes in lake levels or regional precipitation patterns).^{315,316} Globally recognized modeling efforts (i.e., the Coupled Model Intercomparison Project, or CMIP) do not include a realistic representation of the Great Lakes, simulating the influence of the lakes poorly or not at all.^{192,198,317,318,319} Ongoing work to provide evaluation, analysis, and guidance for the Great Lakes region includes comparing this regional model data to commonly used global climate model data (CMIP) that are the basis of many products practitioners currently use (i.e., NCA, IPCC, NOAA State Climate Summaries). To address these challenges, a community of regional modeling experts are working to configure and utilize more sophisticated climate models that more accurately represent the Great Lakes' lake-land-atmosphere system to enhance the understanding of uncertainty to inform better regional decision-making capacity (see <http://glisa.umich.edu/projects/great-lakes-ensemble> for more information).

Key Message 1

Agriculture

The Midwest is a major producer of a wide range of food and animal feed for national consumption and international trade. Increases in warm-season absolute humidity and precipitation have eroded soils, created favorable conditions for pests and pathogens, and degraded the quality of stored grain (*very likely, very high confidence*). Projected changes in precipitation, coupled with rising extreme temperatures before mid-century, will reduce Midwest agricultural productivity to levels of the 1980s without major technological advances (*likely, medium confidence*).

Description of evidence base

Humidity is increasing. Feng et al. (2016)³ show plots of trends in surface and 850 hPa specific humidity of 0.4 and 0.2 g/kg/decade, respectively, from 1979–2014 for the April–May–June period across the Midwest. These represent increases of approximately 5% and 3% per decade, respectively. Automated Surface Observing Stations in Iowa³²⁰ having dew point records of this length and season show dew point temperature increases of about 1°F per decade. Brown and DeGaetano (2013)⁴⁹ show increasing dew points in all seasons throughout the Midwest. Observed changes in annual average maximum temperature for the Midwest over the 20th century (Vose et al. 2017,⁵⁴ Table 6.1) have been less than 1°F. However, future projected changes in annual average temperature (Vose et al. 2017,⁵⁴ Table 6.4), as well as in both warmest day of the year and warmest 5-day 1-in-10 year events (Vose et al. 2017,⁵⁴ Table 6.5), are higher for the Midwest than in any other region of the United States.

Garbrecht et al. (2007)³²¹ state that precipitation changes are sufficient to require U.S. policy changes for agricultural lands. The Soil Erosion Site (http://soilerosion.net/water_erosion.html) describes the soil erosion process and provides links to soil erosion models.³²² Nearing et al. (2004)⁴⁴ report that global climate models project increases in erosivity (the ability or power of rain to cause soil loss) across the northern states of the United States over the 21st century.

Spoilage in stored grain is caused by mold growth and insect activity, which are related to the moisture content and temperature of the stored grain.³²³ The ability of fungi to produce mycotoxins, including aflatoxin and fumonisins, is largely influenced by temperature, relative humidity, insect attack, and stress conditions of the plants.^{57,324} Humidity has a determining influence on the growth rate of these degradation agents.³²⁵

Germination of wheat declined in storage facilities where moisture level increased with time.³²⁶ Freshly harvested, high-moisture content grain must be dried to minimize (or prevent) excessive respiration and mold growth on grains.³²⁷ The storage life of grain is shortened significantly when stored at warm temperatures. One day of holding warm, wet corn before drying can decrease storage life by 50%.⁴⁵

Feng et al. (2016)³ show humidity is rising in the Midwest in the warm season. Cook et al. (2008)⁴ show that the factors leading to these humidity increases (warming Gulf of Mexico and strengthening of the Great Plains Low-Level Jet) will increase in a warming climate.

The ability of fungi to produce mycotoxins is largely influenced by temperature, relative humidity, insect attack, and stress conditions of the plants.³²⁴ More extreme rainfall events would favor formation of Deoxynivalenol, also known as vomitoxin.⁵⁷

Hatfield et al. (2011,⁵⁰ Table 1) give the relationships between temperature and vegetative function as well as reproductive capacity. This work was expanded and updated in Walthall et al. (2012).³²⁸

Mader et al. (2010)⁷⁴ report a comprehensive climate index for describing the effect of ambient temperature, relative humidity, radiation, and wind speed on environmental stress in animals. St-Pierre et al. (2003)³²⁹ provide tables estimating economic losses in dairy due to reduced reproduction. The data show a strong gradient across the Midwest (with losses in Iowa, Illinois, and Indiana being three times the losses in Minnesota, Wisconsin, and Michigan under the current

climate). Temperature and humidity increases projected for the Midwest will increase economic losses across the entire region. Lewis and Bunter (2010)³³⁰ document heat stress effects of temperature on pig production and reproduction.

St-Pierre et al. (2003)³²⁹ provide tables estimating economic losses in dairy, beef, swine, and poultry, resulting in declines from both meat/milk/egg production. The data show a strong gradient across the Midwest (with losses in Iowa, Illinois, and Indiana being twice the losses in Minnesota, Wisconsin, and Michigan under the current climate). Temperature and humidity increases projected for the Midwest will increase losses across the entire region. Babinszky et al. (2011)⁷⁵ identified temperature thresholds for meat/egg/milk production, beyond which performance declines. The adverse effects of heat stress include high mortality, decreased feed consumption, poor body weight gain and meat quality in broiler chickens, and poor laying rate, egg weight, and shell quality in laying hens.⁷⁶

Takle et al. (2013)⁶⁵ found that by mid-century, yields of corn and soybean are projected to fall well below projections based on extrapolation of trends since 1970 even under an optimistic economic scenario, with larger interannual variability in yield and total production. Liang et al. (2017)² report that the ratio of measured agricultural output to measured inputs would drop by an average 3% to 4% per year under medium to high emissions scenarios and could fall to pre-1980 levels by 2050 even when accounting for present rates of innovation. Schauburger et al. (2017)⁶⁶ found that the impact of exposure to temperatures from 30°C to 36°C projected for the end of the century under RCP8.5 creates yield losses of 49% for maize and 40% for soybean.

According to Easterling et al. (2017),¹⁹³ evidence suggests that droughts have become less frequent in the Midwest as the region has become wetter. However, they note that “future higher temperatures will likely lead to greater frequencies and magnitudes of agricultural droughts throughout the continental United States as the resulting increases in evapotranspiration outpace projected precipitation increases.”

Major uncertainties

Global and regional climate models do not simulate well the dynamical structure of mesoscale convective systems in the Midwest, which are the critical “end processes” that create intense precipitation from increasing amounts of moisture evaporated over the Gulf of Mexico and transported by low-level jets (LLJs) into the Midwest. Secondly, the strengthening of future LLJs depends on strengthening of both the Bermuda surface high pressure and the lee surface low over the eastern Rocky Mountains. Confirming simulations of this in future climates are needed. Global and regional climate models do simulate future scenarios having increasing temperatures for the region with high confidence (a necessary ingredient for increased humidity). There is uncertainty of the temperature thresholds for crops because, as pointed out by Schauburger et al. (2017),⁶⁶ some negative impacts of higher temperatures can be overcome through increased water availability. Agricultural yield models, productivity models, and integrated assessment models each provide different ways of looking at agricultural futures, and each of these three types of models has high levels of uncertainty. However, all point to agriculture futures that fail to maintain upward historical trends.

Description of confidence and likelihood

There is *very high confidence* that increases in warm-season absolute humidity and precipitation *very likely* have eroded soils, created favorable conditions for pests and pathogens, and degraded quality of stored grain. There is *medium confidence* that projected increases in moisture, coupled with rising mid-summer temperatures, *likely* will be detrimental to crop and livestock production and put future gains in commodity grain production at risk by mid-century. Projected changes in precipitation, coupled with rising extreme temperatures, provide *medium confidence* that by mid-century Midwest agricultural productivity *likely* will decline to levels of the 1980s without major technological advances.

Key Message 2

Forestry

Midwest forests provide numerous economic and ecological benefits, yet threats from a changing climate are interacting with existing stressors such as invasive species and pests to increase tree mortality and reduce forest productivity (*likely, high confidence*). Without adaptive actions, these interactions will result in the loss of economically and culturally important tree species such as paper birch and black ash (*very likely, very high confidence*) and are expected to lead to the conversion of some forests to other forest types (*likely, high confidence*) or even to non-forested ecosystems by the end of the century (*as likely as not, medium confidence*). Land managers are beginning to manage risk in forests by increasing diversity and selecting for tree species adapted to a range of projected conditions.

Description of evidence base

Multiple ecosystem vulnerability assessments that have been conducted for major forested ecoregions within the Midwest^{89,90,91,92,93} suggest that climate change is expected to have significant direct impacts to forests through effects of warming and changes in the timing and amounts of precipitation.^{96,98,103,104}

Significant indirect impacts to forests are expected as warming increases the negative effects of invasive plants, insect pests, and tree pathogens of forests.^{105,106} Increasing stress on individual trees from climate changes (warming temperatures, drought, and frost damage) increases the susceptibility of trees to the impacts from invasive plants, insect pests, and disease agents.^{109,111}

Direct and indirect impacts of climate change may lead to the decline of culturally^{88,127} and economically important tree species,¹²⁵ as well as leading to shifts in major forest types and altered forest composition as tree species at the northern limits of their ranges decline and southern species experience increasing suitable habitat.¹²⁰ These shifts raise the possibility of future losses of economic and cultural benefits of forests due to conversion to different forest types or the change to non-forest ecosystems.^{119,123,124}

Many examples of land managers implementing climate adaptation in forest management exist, suggesting significant willingness to address the impacts of a changing climate across diverse land ownerships in managed forests¹³⁴ and urban forests.¹³³ Forest management strategies to adapt to a changing climate highlight the importance of increasing forest diversity and managing for

tree species adapted to a range of climate conditions.⁸ The importance of Traditional Ecological Knowledge for informing approaches for climate adaptation on tribal lands and within ceded territory is recognized.³³¹

Major uncertainties

There is significant uncertainty surrounding the ability of tree species migration rates to keep pace with changes in climate (based on temperature and precipitation) due to existing forest fragmentation and loss of habitat. Uncertainty in forest management responses, including active and widespread adaptation efforts that alter forest composition, add to the uncertainty of tree species movements. This leads to considerable uncertainty in the extent to which shifts in tree species ranges may lead to altered forest composition or loss of forest ecosystems in the future.

Due to the complex interactions among species, there is uncertainty in the extent that longer growing seasons, warming temperatures, and increased CO₂ concentrations will benefit tree species, due to both limitations in available water and nutrients, as well as limited benefits for trees relative to the positive influences of these changes on stressors (invasives, insect pests, pathogens).

Description of confidence and likelihood

There is *high confidence* that the interactions of warming temperatures, precipitation changes, and drought with insect pests, invasive plants, and tree pathogens will *likely* lead to increased tree mortality of some species, reducing productivity of some forests. There is *very high confidence* that these interactions will *very likely* result in the decline of some economically or culturally important tree species. Additionally, there is *high confidence* that suitable habitat conditions for tree species will change as temperatures increase and precipitation patterns change, making it *likely* that forest composition will be altered and forest ecosystems may shift to new forest types. Due to uncertainties on species migration rates and forest management responses to climate changes, there is *medium confidence* that by the end of the century, some forest ecosystems are *as likely as not* to convert to non-forest ecosystems.

Key Message 3

Biodiversity and Ecosystems

The ecosystems of the Midwest support a diverse array of native species and provide people with essential services such as water purification, flood control, resource provision, crop pollination, and recreational opportunities. Species and ecosystems, including the important freshwater resources of the Great Lakes, are typically most at risk when climate stressors, like temperature increases, interact with land-use change, habitat loss, pollution, nutrient inputs, and nonnative invasive species (*very likely, very high confidence*). Restoration of natural systems, increases in the use of green infrastructure, and targeted conservation efforts, especially of wetland systems, can help protect people and nature from climate change impacts (*likely, high confidence*).

Description of evidence base

Changes in climate will very likely stress many species and ecological systems in the Midwest. As a result of increases in climate stressors, which typically interact with multiple other stressors, especially in the southern half of the Midwest region, both the ecological systems and the ecological services (water purification, pollination of crops and wild species, recreational opportunities, etc.) they provide to people are at risk. We draw from a wide range of national and global scale assessments of risks to biodiversity (e.g., Maclean and Wilson 2011, Pearson et al. 2014, and the review by Staudinger et al. 2013 that covered literature included in the Third National Climate Assessment^{20,18,22}), which all agree that on the whole, we are highly likely to see increases in species declines and extinctions as a result of climate change. It is very challenging to say specifically what combination of factors will drive these responses, but the weight of evidence suggests very high confidence in the overall trends. The link to interactions with other stressors is also very strong and is described in Brook et al. (2008)¹⁵⁷ and Cahill et al. (2013),¹⁷ among others. Terrestrial ecosystem connectivity, thought to be important for the adaptive capacity of many species, is very low in the southern half of the Midwest region.^{158,159} This may limit the movement of species to more suitable habitats or for species from the southern United States to migrate into the Midwest. These connectivity/movement potential studies also support the idea that land-use change will constrain the potential for retaining function and overall diversity levels. The last section refers to the benefits of restoration as a mechanism for protecting people and nature from climate change impacts. While it is not possible to fully demonstrate that protection of people and nature is indeed occurring now from climate change impacts (we would need attribution of current floods, etc.), there is strong evidence that actions like restoring wetlands can reduce flooding impacts¹⁸² and that protecting forests protects water quality and supply.

Major uncertainties

There is significant uncertainty surrounding the ability of species and ecosystems to persist and thrive under climate change, and we expect to see many different types of responses (population increases, declines, local and regional extinctions).¹⁷ In some cases, climate change does have the potential to benefit species; for example, fish in the coldest regions of the Great Lakes (i.e., Lake Superior) are likely to show increases in productivity, at least in the short run.³³² However, as a whole, given the environmental context upon which climate change is operating, and the presence of many cold-adapted species that are close to the southern edge of their distributional range, we expect more declines than increases.

The last section of the Key Message focuses on land protection and restoration—conservation strategies intended to reduce the impacts of land-use change. Many modeling studies have called out loss of habitat in the Midwest as a key barrier to both local survival and species movement in response to climate change (Schloss et al. 2012 and Carroll et al. 2015 are two of the most recent^{158,159}). Restoring habitat can restore connectivity and protect key ecological functions like pollination services and water purification. Restoring wetlands also can help protect ecosystems and people from flooding, which is the rationale for the last line in the Key Message.

Description of confidence and likelihood

In the Midwest, we already have seen very high levels of habitat loss and conversion, especially in grasslands, wetlands, and freshwater systems. This habitat degradation, in addition to the

pervasive impacts of invasive species, pollution, water extraction, and lack of connectivity, all suggest that the adaptive capacity of species and systems is compromised relative to systems that are more intact and under less stress. Over time, this pervasive habitat loss and degradation has contributed to population declines, especially for wetland, prairie, and stream species. A reliance on cold surface-water systems, which often have compromised connectivity (due to dams, road-stream crossings with structures that impede stream flow, and other barriers) suggests that freshwater species, especially less mobile species like mussels, which are already rare, are at particular risk of declines and extinction. Due to the variety of life histories and climate sensitivities of species within the region, it is very challenging to specify what mechanisms will be most important in terms of driving change. However, knowing that drivers like invasive species, habitat loss, pollution, and hydrologic modifications promote species declines, it is *very likely* that the effects of climate change will interact, and we have *very high confidence* that these interactions will tend to increase, rather than decrease, stresses on species that are associated with these threats. While there is strong evidence that investments in restoring habitat can benefit species, we currently do not have strong observational evidence of the use of these new habitats, or benefits of restored wetlands, in response to isolated climate drivers. Thus, the confidence level for this statement is lower than for the first half of the message.

Key Message 4

Human Health

Climate change is expected to worsen existing conditions and introduce new health threats by increasing the frequency and intensity of poor air quality days, extreme high temperature events, and heavy rainfalls; extending pollen seasons; and modifying the distribution of disease-carrying pests and insects (*very likely, very high confidence*). By mid-century, the region is projected to experience substantial, yet avoidable, loss of life, worsened health conditions, and economic impacts estimated in the billions of dollars as a result of these changes (*likely, high confidence*). Improved basic health services and increased public health measures—including surveillance and monitoring—can prevent or reduce these impacts (*likely, high confidence*).

Description of evidence base

There is strong evidence that increasing temperatures and precipitation in the Midwest will occur by the middle and end of the 21st century.²⁷ The impacts of these changes on human health are broadly captured in the 2016 U.S. Global Change Research Program's Climate and Health Assessment.²⁶ Air quality, including particulate matter and ground-level ozone, is positively associated with increased temperatures and has been well-documented to show deleterious impacts on morbidity and mortality.²³¹ Likewise, increased temperatures have been shown in communities in the Midwest, as well as across the United States, to have substantial impacts on health and well-being.^{232,233,235,236,333,334} The frequency of extreme rainfall events in the Midwest has increased in recent decades, and this trend is projected to continue.¹⁹³ Studies have shown that extreme rainfall events lead to disease, injury, and death.²³⁷ Increases in seasonal temperatures and shifting precipitation patterns have been well documented to be correlated with increased pollen production, allergenicity, and pollen season length.^{230,231} Similarly, there is agreement that shifting temperature and precipitation patterns are making habitats more suitable for disease-carrying vectors to move

northward toward the Midwest region.^{242,243,250,335,336,337} The disease burden and economic projections primarily are based on EPA estimates.²⁸

Access to basic preventive care measures quantifiably reduces disease burden for climate-sensitive exposures.^{238,240} Gray literature indicates that public health practitioners are dedicated to increasing capacity for adapting to climate change through classic public health activities such as conducting vulnerability assessments, employing communication and outreach campaigns, and investing in surveillance efforts.^{26,244,245,246,247,248}

Major uncertainties

While the modeling performed by the EPA was completed using the best available information, there is uncertainty around the extent to which biophysical adaptations will protect midwestern populations from heat-, air pollution-, aeroallergen-, and vector-related illness and death. Likewise, while there is a general consensus regarding habitat suitability for disease-carrying vectors in the eastern and western United States, the degree to which the disease burden may increase or decrease is largely uncertain.

Description of confidence and likelihood

Based on the evidence, there is *very high confidence* that climate change is *very likely* to impact midwesterners' health.

Key Message 5

Transportation and Infrastructure

Storm water management systems, transportation networks, and other critical infrastructure are already experiencing impacts from changing precipitation patterns and elevated flood risks (*medium confidence*). Green infrastructure is reducing some of the negative impacts by using plants and open space to absorb storm water (*medium confidence*). The annual cost of adapting urban storm water systems to more frequent and severe storms is projected to exceed \$500 million for the Midwest by the end of the century (*medium confidence*).

Description of evidence base

The patterns of increased annual precipitation, and the size and frequency of heavy precipitation events in the Midwest, are shown in numerous studies and highlighted in Melillo et al. (2014)²⁷ and Easterling et al. (2017).¹⁹³ Increases in annual precipitation of 5% to 15% are reported across the Midwest region.¹⁹³ In addition, both the frequency and the intensity of heavy precipitation events in the Midwest have increased since 1901.¹⁹³

For the early 21st century (2016–2045), both lower and higher scenarios (RCP4.5 and RCP8.5) indicate that average annual precipitation could increase by 1% to 5% across the Midwest, suggesting that the observed increases are likely to continue. By mid-century (2036–2065), both scenarios (RCP4.5 and RCP8.5) indicate precipitation increases of 1% to 5% in Missouri and Iowa and 5% to 10% increases in states to the north and east. By late century (2070–2089), precipitation is expected to increase by 5% to 15% over present day, with slightly larger increases in the higher scenario (RCP8.5). Model simulations suggest that most of these increases will occur in winter and spring

over the 21st century. Similar to annual precipitation, the amounts from the annual maximum one-day precipitation events (a measure of heavy precipitation events) are projected to increase over time in the Midwest. The size of the events could increase by 5% to 15% by late century.¹⁹³

Gray literature documents that heavy rains in the Midwest are overwhelming storm water management systems, leading to property damage. Kenward et al. (2016)²⁵⁶ provide examples of rain-related sewage overflows in the Midwest. These include an overflow of 681 million gallons during heavy rains in April 2015 in Milwaukee and an overflow of over 100 million gallons from December 26–28, 2015, in St. Louis. Winters et al. (2015)³⁷ document that failure of storm water management systems in heavy rain leads to property damage, including basement backups.

The disruption of transportation networks by heavy precipitation in the Midwest has been documented by collecting contemporary news reports and by compiling state government reports. Posey (2016)³³⁸ relates that four storms between April 2013 and April 2014 forced evacuations or damaged cars in St. Louis, Missouri. In the same period, there were 18 flood-related closures on Missouri roads, a figure that excludes closures on small local roads. Flooding in May 2017 led to the closure of more than 400 roads across Missouri, a figure that again excludes local roads. Closed roadways included multiple stretches of Interstate 44, as well as sections of I-55, affecting interstate traffic between St. Louis and Memphis.³³⁹ News reports document that the same stretch of I-44 was shut down during the floods of December 2015–January 2016.³⁴⁰

Flood-related disruptions to Midwest barge and rail traffic in 2013 were documented by several articles in *Journal of Commerce*, a shipping trade magazine.^{265,266} *WorkBoat*, a trade journal of the inland shipping industry, documents that Mississippi River navigation has been halted by flooding in 2013, 2015, 2016, and 2017. It also documents low river conditions affecting navigation in 2012 and 2015.^{267,268,269,270,341} Disruptions to rail service caused by the floods of 2017 were documented in news media accounts.³⁴² Changon (2009)³⁴³ documents that flooding in 2008 resulted in extensive damage to railroads in Illinois and adjacent states, with costs exceeding \$150 million due to direct damage and lost revenue.

Although there is ample documentation of transportation systems in the Midwest being disrupted by floods in recent years, there is a lack of long-term time series data on disruptions with which to determine whether these incidents are becoming more frequent. Development of long-term data on transportation disruptions in the Midwest is a research need. It is clear that flood frequency and severity on major rivers in the Midwest have increased in recent decades, although additional research is needed on the relative contributions of climate change and land-use change to increases in flood risk.^{344,345,346}

The EPA estimated economic costs related to infrastructure and transportation in the Midwest, including costs associated with bridge scour and pavement degradation.²⁸ The use of green infrastructure to reduce impacts associated with heavy precipitation is also documented in gray literature, including municipal planning documents. Using planted areas to absorb rainfall and reduce runoff has become a common approach to storm water management.^{223,275,276,347,348,349,350} Dechannelization and restoration of streams as a technique for improving storm water management is described in Trice (2013)²⁸² and Milwaukee Metropolitan Sewer District (2017).²⁸¹ Preservation of open space is described in Ducks Unlimited (2017)²⁷⁹ and the Ozaukee Washington

Land Trust (2016).²⁸⁰ The use of urban forestry as an adaptation method is documented in the Minneapolis Marq2 Project (2017)²⁷⁷ and the Cleveland Tree Plan (2015).²⁷⁸ Projected costs to storm water systems are based on EPA projections.²⁸

Major uncertainties

Although there is *very high confidence* that flood risk is increasing in the Midwest, there remains uncertainty about the relative contributions of climate change and land-use change. There is, however, sufficient evidence that changing precipitation patterns are leading to changes in hydrology in the Midwest,^{351,352,353,354,355} and that heavier precipitation patterns are consistent with projections from climate models, to justify a rating of *medium confidence* to the assertion that climate change is contributing to changes in flooding risk. There is *high confidence* that local governments and nongovernmental organizations are turning to green infrastructure solutions as a response to increased flooding risk. Additional research is needed to quantify the aggregate benefits of these approaches.

While it is clear that flood frequency and severity on major rivers in the Midwest have increased in recent decades, it must be emphasized that the change in precipitation levels is not the only factor contributing to the increase in flood risk. Land-use change, particularly the destruction of floodplains by levee systems, has also been documented as a key contributor to increasing flood risk in the Midwest.^{344,345,346} On smaller streams, tile drainage systems have been shown to exacerbate flood risk.²⁴ Determining the relative contribution of land-use change and climate change to increases in riverine flood risk is an important research need.

Description of confidence and likelihood

There is *medium confidence* that climate change is contributing to increased flood risk in the Midwest; there is *medium confidence* that green infrastructure is reducing flood risk. There is much uncertainty associated with specific numerical projections. This leads to *medium confidence* that costs will exceed \$500 million. However, the EPA projections are sufficient to provide *high confidence* that increasing the capacity of existing storm water systems in order to maintain current levels of service would require significant expenditures on the part of urban sewer districts.

Key Message 6

Community Vulnerability and Adaptation

At-risk communities in the Midwest are becoming more vulnerable to climate change impacts such as flooding, drought, and increases in urban heat islands (*as likely as not, high confidence*). Tribal nations are especially vulnerable because of their reliance on threatened natural resources for their cultural, subsistence, and economic needs (*likely, medium confidence*). Integrating climate adaptation into planning processes offers an opportunity to better manage climate risks now (*medium confidence*). Developing knowledge for decision-making in cooperation with vulnerable communities and tribal nations will help to build adaptive capacity and increase resilience (*high confidence*).

Description of evidence base

Limited evidence in the scientific literature indicates that at-risk communities in the Midwest will be increasingly vulnerable to the impacts of climate change, including increased flooding resulting from increased variation in precipitation patterns and changing lake levels,²⁸⁵ urban heat islands,²⁸⁷ and an intensification of heat and drought (see also the impacts and associated references in the previous sections).²⁸⁶

Several recent survey reports^{28,283,284} project negative climate impacts for tribal nations and Indigenous communities, especially as a result of an increased frequency of extreme precipitation events.²⁸³ Tribal nations are especially vulnerable to climate impacts because of their reliance on natural resources,¹²⁷ the isolation of rural communities, and potential shifts of species out of sovereign land.^{309,310} Climate change thus poses a threat to tribal culture, sovereignty, health, and way of life.³⁹

Gray literature,²⁹³ survey reports,³² and scientific literature²⁹² point to a few initiatives to integrate adaptation into municipal planning processes and utilize participatory methodologies to evaluate and manage climate risk.

A growing body of research indicates that interaction between producers of climate information, intermediaries, and end users plays a critical role in increasing climate knowledge integration and use for adaptation in the Midwest.^{224,294,300,308} Limited evidence links the implementation of adaptation actions identified as a result of these collaborations to reduced sensitivity.^{304,305,306}

Major uncertainties

Limited research specific to the Midwest region contributes to uncertainty around the specific vulnerabilities of at-risk communities, including urban and rural communities and tribal nations. Though climate change planning and action in both Midwest cities and rural areas are underway, documentation remains low, few examples exist in the public literature of the failure or success of efforts to mainstream climate action into municipal governance, and attempts to assess vulnerabilities, especially in poor urban communities, frequently encounter climate justice barriers. Likewise, the number, scope, and nature of tribal adaptation plans remain undocumented, as does the degree of implementation of these plans and the manner in which Traditional Ecological Knowledge is incorporated.

Description of confidence and likelihood

There is *high confidence* that communities in the Midwest will *as likely as not* be increasingly vulnerable to climate change impacts such as flooding, urban heat islands, and drought. Similarly, there is *medium confidence* that tribal nations in the Midwest are *likely* to be especially vulnerable because of their reliance on threatened natural resources for their cultural, subsistence, and economic needs. Due to limited documentation in the literature, there is *medium confidence* that integrating adaptation into planning processes will offer an opportunity to manage climate risk better. Finally, there is *high confidence* that developing knowledge for decision-making in cooperation with vulnerable communities and tribal nations will help to decrease sensitivity and build adaptive capacity.

References

1. U.S. Bureau of Economic Analysis, 2017: Gross Domestic Product by State: Fourth Quarter and Annual 2016. U.S. Department of Commerce, Washington, DC, 11 May 2017. https://apps.bea.gov/newsreleases/regional/gdp_state/2017/pdf/qgsp0517.pdf
2. Liang, X.-Z., Y. Wu, R.G. Chambers, D.L. Schmoldt, W. Gao, C. Liu, Y.-A. Liu, C. Sun, and J.A. Kennedy, 2017: Determining climate effects on US total agricultural productivity. *Proceedings of the National Academy of Sciences of the United States of America*, **114** (12), E2285-E2292. <http://dx.doi.org/10.1073/pnas.1615922114>
3. Feng, Z., L.R. Leung, S. Hagos, R.A. Houze, C.D. Burleyson, and K. Balaguru, 2016: More frequent intense and long-lived storms dominate the springtime trend in central US rainfall. *Nature Communications*, **7**, 13429. <http://dx.doi.org/10.1038/ncomms13429>
4. Cook, K.H., E.K. Vizy, Z.S. Launer, and C.M. Patricola, 2008: Springtime intensification of the Great Plains low-level jet and midwest precipitation in GCM simulations of the twenty-first century. *Journal of Climate*, **21** (23), 6321-6340. <http://dx.doi.org/10.1175/2008jcli2355.1>
5. Pruski, F.F. and M.A. Nearing, 2002: Climate-induced changes in erosion during the 21st century for eight U.S. locations. *Water Resources Research*, **38** (12), 34-1 - 34-11. <http://dx.doi.org/10.1029/2001WR000493>
6. Delgado, J.A., M.A. Nearing, and C.W. Rice, 2013: Conservation practices for climate change adaptation. *Advances in Agronomy*. Sparks, D.L., Ed. Academic Press, 47-115. <http://dx.doi.org/10.1016/B978-0-12-407685-3.00002-5>
7. Rosenzweig, C., F.N. Tubiello, R. Goldberg, E. Mills, and J. Bloomfield, 2002: Increased crop damage in the US from excess precipitation under climate change. *Global Environmental Change*, **12**, 197-202. [http://dx.doi.org/10.1016/S0959-3780\(02\)00008-0](http://dx.doi.org/10.1016/S0959-3780(02)00008-0)
8. Ontl, T.A., C. Swanston, L.A. Brandt, P.R. Butler, A.W. D'Amato, S.D. Handler, M.K. Janowiak, and P.D. Shannon, 2018: Adaptation pathways: Ecoregion and land ownership influences on climate adaptation decision-making in forest management. *Climatic Change*, **146** (1), 75-88. <http://dx.doi.org/10.1007/s10584-017-1983-3>
9. Allan, J.D., P.B. McIntyre, S.D.P. Smith, B.S. Halpern, G.L. Boyer, A. Buchsbaum, G.A. Burton, L.M. Campbell, W.L. Chadderton, J.J.H. Ciborowski, P.J. Doran, T. Eder, D.M. Infante, L.B. Johnson, C.A. Joseph, A.L. Marino, A. Prusevich, J.G. Read, J.B. Rose, E.S. Rutherford, S.P. Sowa, and A.D. Steinman, 2013: Joint analysis of stressors and ecosystem services to enhance restoration effectiveness. *Proceedings of the National Academy of Sciences of the United States of America*, **110** (1), 372-377. <http://dx.doi.org/10.1073/pnas.1213841110>
10. Michalak, A.M., E.J. Anderson, D. Beletsky, S. Boland, N.S. Bosch, T.B. Bridgeman, J.D. Chaffin, K. Cho, R. Confesor, I. Daloğlu, J.V. DePinto, M.A. Evans, G.L. Fahnenstiel, L. He, J.C. Ho, L. Jenkins, T.H. Johengen, K.C. Kuo, E. LaPorte, X. Liu, M.R. McWilliams, M.R. Moore, D.J. Posselt, R.P. Richards, D. Scavia, A.L. Steiner, E. Verhamme, D.M. Wright, and M.A. Zagorski, 2013: Record-setting algal bloom in Lake Erie caused by agricultural and meteorological trends consistent with expected future conditions. *Proceedings of the National Academy of Sciences of the United States of America*, **110** (16), 6448-6452. <http://dx.doi.org/10.1073/pnas.1216006110>
11. Zhong, Y., M. Notaro, S.J. Vavrus, and M.J. Foster, 2016: Recent accelerated warming of the Laurentian Great Lakes: Physical drivers. *Limnology and Oceanography*, **61** (5), 1762-1786. <http://dx.doi.org/10.1002/lno.10331>
12. EPA, 2016: Report to Congress: Combined Sewer Overflows into the Great Lakes Basin. EPA 833-R-16-006. U.S. EPA, Office of Wastewater Management, Washington, DC, 92 pp. https://www.epa.gov/sites/production/files/2016-05/documents/gls_cso_report_to_congress_-_4-12-2016.pdf
13. Van Cleave, K., J.D. Lenters, J. Wang, and E.M. Verhamme, 2014: A regime shift in Lake Superior ice cover, evaporation, and water temperature following the warm El Niño winter of 1997-1998. *Limnology and Oceanography*, **59** (6), 1889-1898. <http://dx.doi.org/10.4319/lno.2014.59.6.1889>
14. Wang, J., X. Bai, H. Hu, A. Clites, M. Colton, and B. Lofgren, 2012: Temporal and spatial variability of Great Lakes ice cover, 1973-2010. *Journal of Climate*, **25**, 1318-1329. <http://dx.doi.org/10.1175/2011JCLI4066.1>

15. Mishra, V., K.A. Cherkauer, and L.C. Bowling, 2011: Changing thermal dynamics of lakes in the Great Lakes region: Role of ice cover feedbacks. *Global and Planetary Change*, **75** (3), 155-172. <http://dx.doi.org/10.1016/j.gloplacha.2010.11.003>
16. Hanrahan, J.L., S.V. Kravtsov, and P.J. Roebber, 2010: Connecting past and present climate variability to the water levels of Lakes Michigan and Huron. *Geophysical Research Letters*, **37** (1), L01701. <http://dx.doi.org/10.1029/2009GL041707>
17. Cahill, A.E., M.E. Aiello-Lammens, M.C. Fisher-Reid, X. Hua, C.J. Karanewsky, H. Yeong Ryu, G.C. Sbeglia, F. Spagnolo, J.B. Waldron, O. Warsi, and J.J. Wiens, 2013: How does climate change cause extinction? *Proceedings of the Royal Society B: Biological Sciences*, **280** (1750). <http://dx.doi.org/10.1098/rspb.2012.1890>
18. Pearson, R.G., J.C. Stanton, K.T. Shoemaker, M.E. Aiello-Lammens, P.J. Ersts, N. Horning, D.A. Fordham, C.J. Raxworthy, H.Y. Ryu, J. McNees, and H.R. Akcakaya, 2014: Life history and spatial traits predict extinction risk due to climate change. *Nature Climate Change*, **4** (3), 217-221. <http://dx.doi.org/10.1038/nclimate2113>
19. Fei, S., J.M. Desprez, K.M. Potter, I. Jo, J.A. Knott, and C.M. Oswalt, 2017: Divergence of species responses to climate change. *Science Advances*, **3** (5), e1603055. <http://dx.doi.org/10.1126/sciadv.1603055>
20. Maclean, I.M.D. and R.J. Wilson, 2011: Recent ecological responses to climate change support predictions of high extinction risk. *Proceedings of the National Academy of Sciences of the United States of America*, **108** (30), 12337-12342. <http://dx.doi.org/10.1073/pnas.1017352108>
21. Duffinbaugh, N.S. and C.B. Field, 2013: Changes in ecologically critical terrestrial climate conditions. *Science*, **341** (6145), 486-92. <http://dx.doi.org/10.1126/science.1237123>
22. Staudinger, M.D., S.L. Carter, M.S. Cross, N.S. Dubois, J.E. Duffy, C. Enquist, R. Griffis, J.J. Hellmann, J.J. Lawler, J. O'Leary, S.A. Morrison, L. Sneddon, B.A. Stein, L.M. Thompson, and W. Turner, 2013: Biodiversity in a changing climate: A synthesis of current and projected trends in the US. *Frontiers in Ecology and the Environment*, **11** (9), 465-473. <http://dx.doi.org/10.1890/120272>
23. Villarini, G., E. Scoccimarro, K.D. White, J.R. Arnold, K.E. Schilling, and J. Ghosh, 2015: Projected changes in discharge in an agricultural watershed in Iowa. *JAWRA Journal of the American Water Resources Association*, **51** (5), 1361-1371. <http://dx.doi.org/10.1111/1752-1688.12318>
24. Kelly, S.A., Z. Takbiri, P. Belmont, and E. Foufoula-Georgiou, 2017: Human amplified changes in precipitation-runoff patterns in large river basins of the Midwestern United States. *Hydrology and Earth System Sciences*, **21** (10), 5065-5088. <http://dx.doi.org/10.5194/hess-21-5065-2017>
25. Hall, K.R., M.E. Herbert, S.P. Sowa, S. Mysorekar, S.A. Woznicki, P.A. Nejadhashemi, and L. Wang, 2017: Reducing current and future risks: Using climate change scenarios to test an agricultural conservation framework. *Journal of Great Lakes Research*, **43** (1), 59-68. <http://dx.doi.org/10.1016/j.jglr.2016.11.005>
26. USGCRP, 2016: *The Impacts of Climate Change on Human Health in the United States: A Scientific Assessment*. U.S. Global Change Research Program, Washington, DC, 312 pp. <http://dx.doi.org/10.7930/J0R49NQX>
27. Melillo, J.M., T.C. Richmond, and G.W. Yohe, Eds., 2014: *Climate Change Impacts in the United States: The Third National Climate Assessment*. U.S. Global Change Research Program, Washington, DC, 841 pp. <http://dx.doi.org/10.7930/J0Z31WJ2>
28. EPA, 2017: Multi-model Framework for Quantitative Sectoral Impacts Analysis: A Technical Report for the Fourth National Climate Assessment. EPA 430-R-17-001. U.S. Environmental Protection Agency (EPA), Washington, DC, 271 pp. https://cfpub.epa.gov/si/si_public_record_Report.cfm?dirEntryId=335095
29. Sarofim, M.C., S. Saha, M.D. Hawkins, D.M. Mills, J. Hess, R. Horton, P. Kinney, J. Schwartz, and A. St. Juliana, 2016: Ch. 2: Temperature-related death and illness. *The Impacts of Climate Change on Human Health in the United States: A Scientific Assessment*. U.S. Global Change Research Program, Washington, DC, 43-68. <http://dx.doi.org/10.7930/J0MG7MDX>
30. USGCRP, 2017: Climate Science Special Report: Fourth National Climate Assessment, Volume I. Wuebbles, D.J., D.W. Fahey, K.A. Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock, Eds. U.S. Global Change Research Program, Washington, DC, 470 pp. <http://dx.doi.org/10.7930/J0J964J6>

31. USGCRP, 2017: Regional Engagement Workshop Summary Report: Midwest Region U.S. Global Change Research Program, Washington, DC, 9 pp. http://www.globalchange.gov/sites/globalchange/files/REW_Midwest.pdf
32. EPA, 2014: Being Prepared for Climate Change: A Workbook for Developing Risk-Based Adaptation Plans. U.S. EPA, Office of Water, Washington, DC, 120 pp. https://www.epa.gov/sites/production/files/2014-09/documents/being_prepared_workbook_508.pdf
33. Schulte, L.A., J. Niemi, M.J. Helmers, M. Liebman, J.G. Arbuckle, D.E. James, R.K. Kolka, M.E. O'Neal, M.D. Tomer, J.C. Tyndall, H. Asbjornsen, P. Drobney, J. Neal, G. Van Ryswyk, and C. Witte, 2017: Prairie strips improve biodiversity and the delivery of multiple ecosystem services from corn-soybean croplands. *Proceedings of the National Academy of Sciences of the United States of America*, **114** (42), 11247-11252. <http://dx.doi.org/10.1073/pnas.1620229114>
34. ISU, 2018: STRIPS (Science-based Trials of Rowcrops Integrated with Prairie Strips) Project [web site]. Iowa State University (ISU), Ames, IA. <https://www.nrem.iastate.edu/research/STRIPS/>
35. Hatfield, J., C. Swanston, M. Janowiak, R.F. Steele, J. Hempel, J. Bochicchio, W. Hall, M. Cole, S. Hestvik, and J. Whitaker, 2015: USDA Midwest and Northern Forests Regional Climate Hub: Assessment of Climate Change Vulnerability and Adaptation and Mitigation Strategies. Anderson, T., Ed. U.S. Department of Agriculture, 55 pp. <https://www.climatehubs.occ.usda.gov/content/usda-midwest-and-northern-forests-regional-climate-hub-assessment-climate-change>
36. Swanston, C., L.A. Brandt, M.K. Janowiak, S.D. Handler, P. Butler-Leopold, L. Iverson, F.R. Thompson III, T.A. Ontl, and P.D. Shannon, 2018: Vulnerability of forests of the Midwest and Northeast United States to climate change. *Climatic Change*, **146** (1), 103-116. <http://dx.doi.org/10.1007/s10584-017-2065-2>
37. Winters, B.A., J. Angel, C. Ballerine, J. Byard, A. Flegel, D. Gambill, E. Jenkins, S. McConkey, M. Markus, B.A. Bender, and M.J. O'Toole, 2015: Report for the Urban Flooding Awareness Act. Illinois Department of Natural Resources, Springfield, IL, 89 pp. https://www.dnr.illinois.gov/WaterResources/Documents/Final_UFAA_Report.pdf
38. City of Chicago, 2014: Green Stormwater Infrastructure Strategy. 44 pp. <https://www.cityofchicago.org/content/dam/city/progs/env/ChicagoGreenStormwaterInfrastructureStrategy.pdf>
39. Bennett, T.M.B., N.G. Maynard, P. Cochran, R. Gough, K. Lynn, J. Maldonado, G. Voggesser, S. Wotkyns, and K. Cozzetto, 2014: Ch. 12: Indigenous peoples, lands, and resources. *Climate Change Impacts in the United States: The Third National Climate Assessment*. Melillo, J.M., T.C. Richmond, and G.W. Yohe, Eds. U.S. Global Change Research Program, Washington, DC, 297-317. <http://dx.doi.org/10.7930/J09G5JR1>
40. USDA, 2017: Climate Change: Cover Crops and Soil Health. USDA National Resources Conservation Service. <https://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/climatechange/?cid=stelprdb1077238>
41. EIA, 2016: U.S. States: Table C9. Electric Power Sector Consumption Estimates, 2016 [web site]. U.S. Energy Information Administration, Washington, DC. https://www.eia.gov/state/seds/data.php?incfile=/state/seds/sep_sum/html/sum_btu_eu.html&sid=US
42. MSU, 2018: Solar Carport Initiative [web site]. Michigan State University (MSU), East Lansing, MI, accessed March 28. <http://ipf.msu.edu/green/practices/solar-carport-initiative.html>
43. Hatfield, J.L., L. Wright-Morton, and B. Hall, 2018: Vulnerability of grain crops and croplands in the Midwest to climatic variability and adaptation strategies. *Climatic Change*, **146** (1-2), 263-275. <http://dx.doi.org/10.1007/s10584-017-1997-x>
44. Nearing, M., F.F. Pruski, and M.R. O'Neal, 2004: Expected climate change impacts on soil erosion rates: A review. *Journal of Soil and Water Conservation*, **59** (1), 43-50. <http://www.jswconline.org/content/59/1/43.abstract>
45. Hurburgh, C., 2016: Wet Weather Creates Challenges for Harvest. Iowa State University, Extension and Outreach, Ames, IA. <https://crops.extension.iastate.edu/cropnews/2016/09/wet-weather-creates-challenges-harvest>

46. Hibbard, K.A., F.M. Hoffman, D. Huntzinger, and T.O. West, 2017: Changes in land cover and terrestrial biogeochemistry. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*. Wuebbles, D.J., D.W. Fahey, K.A. Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock, Eds. U.S. Global Change Research Program, Washington, DC, USA, 277-302. <http://dx.doi.org/10.7930/J0416V6X>
47. Pan, Z., R.W. Arritt, E.S. Takle, W.J. Gutowski, Jr., C.J. Anderson, and M. Segal, 2004: Altered hydrologic feedback in a warming climate introduces a "warming hole." *Geophysical Research Letters*, **31** (17), L17109. <http://dx.doi.org/10.1029/2004GL020528>
48. Kunkel, K.E., L.E. Stevens, S.E. Stevens, L. Sun, E. Janssen, D. Wuebbles, S.D. Hilberg, M.S. Timlin, L. Stoecker, N.E. Westcott, and J.G. Dobson, 2013: Regional Climate Trends and Scenarios for the U.S. National Climate Assessment: Part 3. Climate of the Midwest U.S. NOAA Technical Report NESDIS 142-3. National Oceanic and Atmospheric Administration, National Environmental Satellite, Data, and Information Service, Washington, DC, 103 pp. http://www.nesdis.noaa.gov/technical_reports/NOAA_NESDIS_Tech_Report_142-3-Climate_of_the_Midwest_U.S.pdf
49. Brown, P.J. and A.T. DeGaetano, 2013: Trends in U.S. surface humidity, 1930–2010. *Journal of Applied Meteorology and Climatology*, **52** (1), 147-163. <http://dx.doi.org/10.1175/jamc-d-12-035.1>
50. Hatfield, J.L., K.J. Boote, B.A. Kimball, L.H. Ziska, R.C. Izaurralde, D. Ort, A.M. Thomson, and D. Wolfe, 2011: Climate impacts on agriculture: Implications for crop production. *Agronomy Journal*, **103** (2), 351-370. <http://dx.doi.org/10.2134/agronj2010.0303>
51. Tobin, P.C., S. Nagarkatti, G. Loeb, and M.C. Saunders, 2008: Historical and projected interactions between climate change and insect voltinism in a multivoltine species. *Global Change Biology*, **14** (5), 951-957. <http://dx.doi.org/10.1111/j.1365-2486.2008.01561.x>
52. Bebbler, D.P., M.A.T. Ramotowski, and S.J. Gurr, 2013: Crop pests and pathogens move polewards in a warming world. *Nature Climate Change*, **3** (11), 985-988. <http://dx.doi.org/10.1038/nclimate1990>
53. Andresen, J., S. Hilberg, and K. Kunkel, 2012: Historical Climate and Climate Trends in the Midwestern USA. U.S. National Climate Assessment Midwest Technical Input Report. Great Lakes Integrated Sciences and Assessments (GLISA) Center, Ann Arbor, MI, 18 pp. http://glisa.umich.edu/media/files/NCA/MTIT_Historical.pdf
54. Vose, R.S., D.R. Easterling, K.E. Kunkel, A.N. LeGrande, and M.F. Wehner, 2017: Temperature changes in the United States. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*. Wuebbles, D.J., D.W. Fahey, K.A. Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock, Eds. U.S. Global Change Research Program, Washington, DC, USA, 185-206. <http://dx.doi.org/10.7930/J0N29V45>
55. Anderson, C., D. Claman, and R. Mantilla, 2015: Iowa's Bridge and Highway Climate Change and Extreme Weather Vulnerability Assessment Pilot. HEPN-707. Iowa State University, Institute for Transportation, Ames, IA, 45 pp. http://www.intrans.iastate.edu/research/documents/research-reports/IA_climate_change_vulnerability_assess_w_cvr1.pdf
56. Munkvold, G.P. and X.B. Yang, 1995: Crop damage and epidemics associated with 1993 floods in Iowa. *Plant Disease*, **79** (1), 95-101. <http://dx.doi.org/10.1094/PD-79-0095>
57. Wu, F., D. Bhatnagar, T. Bui-Klimke, I. Carbone, R. Hellmich, G. Munkvold, P. Paul, G. Payne, and E. Takle, 2011: Climate change impacts on mycotoxin risks in US maize. *World Mycotoxin Journal*, **4** (1), 79-93. <http://dx.doi.org/10.3920/WMJ2010.1246>
58. Anderson, P.K., A.A. Cunningham, N.G. Patel, F.J. Morales, P.R. Epstein, and P. Daszak, 2004: Emerging infectious diseases of plants: Pathogen pollution, climate change and agrotechnology drivers. *Trends in Ecology & Evolution*, **19** (10), 535-544. <http://dx.doi.org/10.1016/j.tree.2004.07.021>
59. Liu, Q., A. Ravanlou, and M. Babadoost, 2016: Occurrence of bacterial spot on pumpkin and squash fruit in the north central region of the United States and bacteria associated with the spots. *Plant Disease*, **100** (12), 2377-2382. <http://dx.doi.org/10.1094/PDIS-01-16-0107-RE>
60. Pan, Z., D. Andrade, M. Segal, J. Wimberley, N. McKinney, and E. Takle, 2010: Uncertainty in future soil carbon trends at a central US site under an ensemble of GCM scenario climates. *Ecological Modelling*, **221** (5), 876-881. <http://dx.doi.org/10.1016/j.ecolmodel.2009.11.013>

61. Cai, X., X. Zhang, P.H. Noël, and M. Shafiee-Jood, 2015: Impacts of climate change on agricultural water management: A review. *Wiley Interdisciplinary Reviews: Water*, **2** (5), 439-455. <http://dx.doi.org/10.1002/wat2.1089>
62. Takle, E.S., C. Anderson, M. Jha, and P.W. Gassman, 2006: Upper Mississippi River Basin Modeling Systems Part 4: Climate change impacts on flow and water quality. *Coastal Hydrology and Processes*. Singh, V.P. and Y.J. Xu, Eds. Water Resources Publications LLC, Highlands Ranch, CO, 135-142.
63. Loecke, T.D., A.J. Burgin, D.A. Riveros-Iregui, A.S. Ward, S.A. Thomas, C.A. Davis, and M.A.S. Clair, 2017: Weather whiplash in agricultural regions drives deterioration of water quality. *Biogeochemistry*, **133** (1), 7-15. <http://dx.doi.org/10.1007/s10533-017-0315-z>
64. Wehner, M.F., J.R. Arnold, T. Knutson, K.E. Kunkel, and A.N. LeGrande, 2017: Droughts, floods, and wildfires. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*. Wuebbles, D.J., D.W. Fahey, K.A. Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock, Eds. U.S. Global Change Research Program, Washington, DC, USA, 231-256. <http://dx.doi.org/10.7930/J0CJ8BNN>
65. Takle, E.S.T., D. Gustafson, R. Beachy, G.C. Nelson, D. Mason-D'Croz, and A. Palazzo, 2013: US food security and climate change: Agricultural futures. *Economics: The Open-Access, Open-Assessment E-Journal*, **7** (2013-34), 1-41. <http://dx.doi.org/10.5018/economics-ejournal.ja.2013-34>
66. Schauburger, B., S. Archontoulis, A. Arneth, J. Balkovic, P. Ciais, D. Deryng, J. Elliott, C. Folberth, N. Khabarov, C. Müller, T.A.M. Pugh, S. Rolinski, S. Schaphoff, E. Schmid, X. Wang, W. Schlenker, and K. Frieler, 2017: Consistent negative response of US crops to high temperatures in observations and crop models. *Nature Communications*, **8**, 13931. <http://dx.doi.org/10.1038/ncomms13931>
67. Jin, Z., Q. Zhuang, J. Wang, S.V. Archontoulis, Z. Zobel, and V.R. Kotamarthi, 2017: The combined and separate impacts of climate extremes on the current and future U.S. rainfed maize and soybean production under elevated CO₂. *Global Change Biology*, **23** (7), 2687-2704. <http://dx.doi.org/10.1111/gcb.13617>
68. Deryng, D., J. Elliott, C. Folberth, C. Muller, T.A.M. Pugh, K.J. Boote, D. Conway, A.C. Ruane, D. Gerten, J.W. Jones, N. Khabarov, S. Olin, S. Schaphoff, E. Schmid, H. Yang, and C. Rosenzweig, 2016: Regional disparities in the beneficial effects of rising CO₂ concentrations on crop water productivity. *Nature Climate Change*, **6** (8), 786-790. <http://dx.doi.org/10.1038/nclimate2995>
69. Jaggard, K.W., A. Qi, and E.S. Ober, 2010: Possible changes to arable crop yields by 2050. *Philosophical Transactions of the Royal Society B Biological Sciences*, **365** (1554), 2835-2851. <http://dx.doi.org/10.1098/rstb.2010.0153>
70. Deryng, D., D. Conway, N. Ramankutty, J. Price, and R. Warren, 2014: Global crop yield response to extreme heat stress under multiple climate change futures. *Environmental Research Letters*, **9** (3), 034011. <http://dx.doi.org/10.1088/1748-9326/9/3/034011>
71. Kistner, E., O. Kellner, J. Andresen, D. Todey, and L.W. Morton, 2018: Vulnerability of specialty crops to short-term climatic variability and adaptation strategies in the Midwestern USA. *Climatic Change*, **146** (1), 145-158. <http://dx.doi.org/10.1007/s10584-017-2066-1>
72. Babadoost, M., 2012: The Fruit Rots of Pumpkin. Report on Plant Disease RPD No. 950. University of Illinois Extension, Urbana-Champaign, IL, 7 pp. http://extension.cropsciences.illinois.edu/fruitveg/pdfs/950_fruits_rot_pumpkin.pdf
73. MDNR, 2008: Natural Wild Rice in Minnesota. Minnesota Department of Natural Resources, St. Paul, MN, 114 pp. http://files.dnr.state.mn.us/fish_wildlife/wildlife/shallowlakes/natural-wild-rice-in-minnesota.pdf
74. Mader, T.L., L.J. Johnson, and J.B. Gaughan, 2010: A comprehensive index for assessing environmental stress in animals. *Journal of Animal Science*, **88** (6), 2153-2165. <http://dx.doi.org/10.2527/jas.2009-2586>
75. Babinszky, L., V. Halas, and M.W.A. Verstegen, 2011: Impacts of climate change on animal production and quality of animal food products. *Climate Change—Socioeconomic Effects*. Blanco, J. and H. Kheradmand, Eds. InTech, Rijeka, Croatia, Ch. 10. <http://dx.doi.org/10.5772/23840>
76. Lin, H., H.C. Jiao, J. Buyse, and E. Decuyper, 2007: Strategies for preventing heat stress in poultry. *World's Poultry Science Journal*, **62** (1), 71-86. <http://dx.doi.org/10.1079/WPS200585>

77. NCGA, 2018: Soil Health Partnership [web page]. National Corn Growers Association (NCGA), Chesterfield, MO. <https://www.soilhealthpartnership.org/>
78. National Agricultural Statistics Service, 2014: 2012 Census of Agriculture: 2013 Farm and Ranch Irrigation Survey. AC-12-SS-1. U.S. Department of Agriculture, 249 pp. https://www.agcensus.usda.gov/Publications/2012/Online_Resources/Farm_and_Ranch_Irrigation_Survey/
79. Ballweg, J., 2016: Forest Economy Wisconsin. Wisconsin Department of Natural Resources, Madison, WI, 1 p. <http://dnr.wi.gov/topic/ForestBusinesses/documents/factSheets/FactSheetWisconsin.pdf>
80. Decision Innovation Solutions, 2016: Economic Contributions of Missouri Agriculture and Forestry. Missouri Department of Agriculture, 30 pp. <http://agriculture.mo.gov/economicimpact/county-pdf/MissouriAgForestryEconomicContributionStudy.pdf>
81. Deckard, D.L. and J.A. Skurla, 2011: Economic Contributions of Minnesota's Forest Products Industry—2011 Edition. Minnesota Department of Natural Resources, St. Paul, MN, 18 pp. <https://bit.ly/1CVw9cx>
82. Leefers, L.A., 2015: Forest Products Industries' Economic Contributions to Michigan's Economy in 2013. Michigan Department of Natural Resources. Forest Resource Division, Lansing, MI, 32 pp. https://www.michigan.gov/documents/dnr/FPIECME2013-Leefers_513869_7.pdf
83. Henderson, J.E. and I.A. Munn, 2012: Forestry in Illinois—The Impact of the Forest Products Industry on the Illinois Economy: An Input-Output Analysis. Illinois Forestry Development Council, Springfield, IL, 22 pp. http://ifdc.nres.illinois.edu/wp-content/uploads/2013/10/illinois-forest-products-impact_2012.pdf
84. Leatherberry, E.C., W.K. Moser, C. Perry, C. Woodall, E. Jepsen, S. Pennington, and A. Flickinger, 2006: Iowa's Forests 1999-2003 (Part A). Resource Bulletin NC-266A. U.S. Department of Agriculture, Forest Service, North Central Research Station, St. Paul, MN, 84 pp. https://www.nrs.fs.fed.us/pubs/rb/rb_nc266a.pdf
85. McConnell, E., 2012: Ohio's Forest Economy. Fact Sheet F-80. Ohio State University Extension, Columbus, OH, 8 pp. <https://ohioline.osu.edu/factsheet/F-80>
86. Settle, J., C. Gonso, and M. Seidl, 2016: Indiana's Hardwood Industry: Its Economic Impact (Update of the 2010 Hoover/Settle Report). Indiana State Department of Agriculture, Indianapolis, IN, 25 pp. https://in.gov/isda/files/Indiana_Hardwoods_and_Their_Economic_Impact.pdf
87. Marcouiller, D. and T. Mace, 2005: Forests and Regional Development: Economic Impacts of Woodland Use For Recreation and Timber in Wisconsin. G3694 RP-10/05. University of Wisconsin Cooperative Extension, Madison, WI, 43 pp. <http://learningstore.uwex.edu/Assets/pdfs/G3694.pdf>
88. Stults, M., S. Petersen, J. Bell, W. Baule, E. Nasser, E. Gibbons, and M. Fougerat, 2016: Climate Change Vulnerability Assessment and Adaptation Plan: 1854 Ceded Territory Including the Bois Forte, Fond du Lac, and Grand Portage Reservations. 1854 Treaty Authority, Duluth, MN, 146 pp. [http://www.1854treatyauthority.org/images/ClimateAdaptationPlan_Final-July_2016-optimized\(1\).pdf](http://www.1854treatyauthority.org/images/ClimateAdaptationPlan_Final-July_2016-optimized(1).pdf)
89. Brandt, L., H. He, L. Iverson, F.R. Thompson, P. Butler, S. Handler, M. Janowiak, P.D. Shannon, C. Swanston, M. Albrecht, R. Blume-Weaver, P. Deizman, J. DePuy, W.D. Dijk, G. Dinkel, S. Fei, D.T. Jones-Farrand, M. Leahy, S. Matthews, P. Nelson, B. Oberle, J. Perez, M. Peters, A. Prasad, J.E. Schneiderman, J. Shuey, A.B. Smith, C. Studyvin, J.M. Tirpak, J.W. Walk, W.J. Wang, L. Watts, D. Weigel, and S. Westin, 2014: Central Hardwoods Ecosystem Vulnerability Assessment and Synthesis: A Report from the Central Hardwoods Climate Change Response Framework Project. Gen. Tech. Rep. NRS-124. USDA Forest Service, Newtown Square, PA, 254 pp. <https://www.nrs.fs.fed.us/pubs/45430>
90. Handler, S., M.J. Duveneck, L. Iverson, E. Peters, R.M. Scheller, K.R. Wythers, L. Brandt, P. Butler, M. Janowiak, P.D. Shannon, C. Swanston, K. Barrett, R. Kolka, C. McQuiston, B. Palik, P.B. Reich, C. Turner, M. White, C. Adams, A. D'Amato, S. Hagell, P. Johnson, R. Johnson, M. Larson, S. Matthews, R. Montgomery, S. Olson, M. Peters, A. Prasad, J. Rajala, J. Daley, M. Davenport, M.R. Emery, D. Fehringer, C.L. Hoving, G. Johnson, L. Johnson, D. Neitzel, A. Rissman, C. Rittenhouse, and R. Ziel, 2014: Minnesota Forest Ecosystem Vulnerability Assessment and Synthesis: A Report from the Northwoods Climate Change Response Framework Project. General Technical Report NRS-133. U.S. Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA, 228 pp. https://www.fs.fed.us/nrs/pubs/gtr/gtr_nrs133.pdf

91. Handler, S., M.J. Duveneck, L. Iverson, E. Peters, R.M. Scheller, K.R. Wythers, L. Brandt, P. Butler, M. Janowiak, P.D. Shannon, C. Swanston, A.C. Eagle, J.G. Cohen, R. Corner, P.B. Reich, T. Baker, S. Chhin, E. Clark, D. Fehring, J. Fosgitt, J. Gries, C. Hall, K.R. Hall, R. Heyd, C.L. Hoving, I. Ibáñez, D. Kuhr, S. Matthews, J. Muladore, K. Nadelhoffer, D. Neumann, M. Peters, A. Prasad, M. Sands, R. Swaty, L. Wonch, J. Daley, M. Davenport, M.R. Emery, G. Johnson, L. Johnson, D. Neitzel, A. Rissman, C. Rittenhouse, and R. Ziel, 2014: Michigan Forest Ecosystem Vulnerability Assessment and Synthesis: A Report from the Northwoods Climate Change Response Framework Project. General Technical Report NRS-129. U.S. Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA, 229 pp. <https://www.nrs.fs.fed.us/pubs/45688>
92. Janowiak, M.K., L.R. Iverson, D.J. Mladenoff, E. Peters, K.R. Wythers, W. Xi, L.A. Brandt, P.R. Butler, S.D. Handler, P.D. Shannon, C. Swanston, L.R. Parker, A.J. Amman, B. Bogaczyk, C. Handler, E. Lesch, P.B. Reich, S. Matthews, M. Peters, A. Prasad, S. Khanal, F. Liu, T. Bal, D. Bronson, A. Burton, J. Ferris, J. Fosgitt, S. Hagan, E. Johnston, E. Kane, C. Matula, R. O'Connor, D. Higgins, M. St. Pierre, J. Daley, M. Davenport, M.R. Emery, D. Fehring, C.L. Hoving, G. Johnson, D. Neitzel, M. Notaro, A. Rissman, C. Rittenhouse, and R. Ziel, 2014: Forest Ecosystem Vulnerability Assessment and Synthesis for Northern Wisconsin and Western Upper Michigan: A Report from the Northwoods Climate Change Response Framework Project. Gen. Tech. Rep. NRS-136. U.S. Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA, 247 pp. https://www.fs.fed.us/nrs/pubs/gtr/gtr_nrs136.pdf
93. Swanston, C., M. Janowiak, L. Iverson, L. Parker, D. Mladenoff, L. Brandt, P. Butler, M. St. Pierre, A. Prasad, S. Matthews, M. Peters, D. Higgins, and A. Dorland, 2011: Ecosystem Vulnerability Assessment and Synthesis: A Report from the Climate Change Response Framework Project in Northern Wisconsin. Gen. Tech. Rep. NRS-82. U.S. Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA, 142 pp. https://www.fs.fed.us/nrs/pubs/gtr/gtr_nrs82.pdf
94. Vose, J.M., D.L. Peterson, and T. Patel-Weyand, Eds., 2012: *Effects of Climatic Variability and Change on Forest Ecosystems: A Comprehensive Science Synthesis for the U.S. Forest Sector*. General Technical Report PNW-GTR-870. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, OR, 265 pp. http://www.usda.gov/oce/climate_change/effects_2012/FS_Climate1114%20opt.pdf
95. Boisvenue, C. and S.W. Running, 2006: Impacts of climate change on natural forest productivity—Evidence since the middle of the 20th century. *Global Change Biology*, **12** (5), 862-882. <http://dx.doi.org/10.1111/j.1365-2486.2006.01134.x>
96. Chhin, S., 2010: Influence of climate on the growth of hybrid poplar in Michigan. *Forests*, **1** (4), 209-229. <http://dx.doi.org/10.3390/fl040209>
97. Restaino, C.M., D.L. Peterson, and J. Littell, 2016: Increased water deficit decreases Douglas fir growth throughout western US forests. *Proceedings of the National Academy of Sciences of the United States of America*, **113** (34), 9557-9562. <http://dx.doi.org/10.1073/pnas.1602384113>
98. Worrall, J.J., G.E. Rehfeldt, A. Hamann, E.H. Hogg, S.B. Marchetti, M. Michaelian, and L.K. Gray, 2013: Recent declines of *Populus tremuloides* in North America linked to climate. *Forest Ecology and Management*, **299**, 35-51. <http://dx.doi.org/10.1016/j.foreco.2012.12.033>
99. Bottero, A., A.W. D'Amato, B.J. Palik, J.B. Bradford, S. Fraver, M.A. Battaglia, and L.A. Asherin, 2017: Density-dependent vulnerability of forest ecosystems to drought. *Journal of Applied Ecology*, **54** (6), 1605-1614. <http://dx.doi.org/10.1111/1365-2664.12847>
100. D'Amato, A.W., J.B. Bradford, S. Fraver, and B.J. Palik, 2013: Effects of thinning on drought vulnerability and climate response in north temperate forest ecosystems. *Ecological Applications*, **23** (8), 1735-1742. <http://dx.doi.org/10.1890/13-0677.1>
101. Will, R.E., S.M. Wilson, C.B. Zou, and T.C. Hennessey, 2013: Increased vapor pressure deficit due to higher temperature leads to greater transpiration and faster mortality during drought for tree seedlings common to the forest-grassland ecotone. *New Phytologist*, **200** (2), 366-374. <http://dx.doi.org/10.1111/nph.12321>

102. Vose, J.M., C.F. Miniati, C.H. Luce, H. Asbjornsen, P.V. Caldwell, J.L. Campbell, G.E. Grant, D.J. Isaak, S.P. Loheide II, and G. Sun, 2016: Ecohydrological implications of drought for forests in the United States. *Forest Ecology and Management*, **380**, 335-345. <http://dx.doi.org/10.1016/j.foreco.2016.03.025>
103. Auclair, A.N.D., W.E. Heilman, and B. Brinkman, 2010: Predicting forest dieback in Maine, USA: A simple model based on soil frost and drought. *Canadian Journal of Forest Research*, **40** (4), 687-702. <http://dx.doi.org/10.1139/X10-023>
104. Groffman, P.M., L.E. Rustad, P.H. Templer, J.L. Campbell, L.M. Christenson, N.K. Lany, A.M. Soggi, M.A. Vadeboncoeur, P.G. Schaberg, G.F. Wilson, C.T. Driscoll, T.J. Fahey, M.C. Fisk, C.L. Goodale, M.B. Green, S.P. Hamburg, C.E. Johnson, M.J. Mitchell, J.L. Morse, L.H. Pardo, and N.L. Rodenhouse, 2012: Long-term integrated studies show complex and surprising effects of climate change in the northern hardwood forest. *BioScience*, **62** (12), 1056-1066. <http://dx.doi.org/10.1525/bio.2012.62.12.7>
105. Ramsfield, T.D., B.J. Bentz, M. Faccoli, H. Jactel, and E.G. Brockerhoff, 2016: Forest health in a changing world: Effects of globalization and climate change on forest insect and pathogen impacts. *Forestry: An International Journal of Forest Research*, **89** (3), 245-252. <http://dx.doi.org/10.1093/forestry/cpw018>
106. Weed, A.S., M.P. Ayres, and J.A. Hicke, 2013: Consequences of climate change for biotic disturbances in North American forests. *Ecological Monographs*, **83** (4), 441-470. <http://dx.doi.org/10.1890/13-0160.1>
107. Aronson, M.F.J. and S.N. Handel, 2011: Deer and invasive plant species suppress forest herbaceous communities and canopy tree regeneration. *Natural Areas Journal*, **31** (4), 400-407. <http://dx.doi.org/10.3375/043.031.0410>
108. Liu, Y., A.M.O. Oduor, Z. Zhang, A. Manea, I.M. Tooth, M.R. Leishman, X. Xu, and M. Kleunen, 2017: Do invasive alien plants benefit more from global environmental change than native plants? *Global Change Biology*, **23** (8), 3363-3370. <http://dx.doi.org/10.1111/gcb.13579>
109. Sturrock, R.N., S.J. Frankel, A.V. Brown, P.E. Hennon, J.T. Kliejunas, K.J. Lewis, J.J. Worrall, and A.J. Woods, 2011: Climate change and forest diseases. *Plant Pathology*, **60** (1), 133-149. <http://dx.doi.org/10.1111/j.1365-3059.2010.02406.x>
110. Dale, A.G. and S.D. Frank, 2017: Warming and drought combine to increase pest insect fitness on urban trees. *PLOS ONE*, **12** (3), e0173844. <http://dx.doi.org/10.1371/journal.pone.0173844>
111. Seidl, R., D. Thom, M. Kautz, D. Martin-Benito, M. Peltoniemi, G. Vacchiano, J. Wild, D. Ascoli, M. Petr, J. Honkaniemi, M.J. Lexer, V. Trotsiuk, P. Mairota, M. Svoboda, M. Fabrika, T.A. Nagel, and C.P.O. Reyher, 2017: Forest disturbances under climate change. *Nature Climate Change*, **7** (6), 395-402. <http://dx.doi.org/10.1038/nclimate3303>
112. Goring, S.J., D.J. Mladenoff, C.V. Cogbill, S. Record, C.J. Paciorek, S.T. Jackson, M.C. Dietze, A. Dawson, J.H. Matthes, J.S. McLachlan, and J.W. Williams, 2016: Novel and lost forests in the upper midwestern United States, from new estimates of settlement-era composition, stem density, and biomass. *PLOS ONE*, **11** (12), e0151935. <http://dx.doi.org/10.1371/journal.pone.0151935>
113. Shifley, S.R. and W.K. Moser, Eds., 2016: *Future Forests of the Northern United States*. U.S. Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA, 388 pp. <https://www.nrs.fs.fed.us/pubs/50448>
114. O'Hara, K.L. and B.S. Ramage, 2013: Silviculture in an uncertain world: Utilizing multi-aged management systems to integrate disturbance. *Forestry: An International Journal of Forest Research*, **86** (4), 401-410. <http://dx.doi.org/10.1093/forestry/cpt012>
115. Cadotte, M.W., R. Dinnage, and D. Tilman, 2012: Phylogenetic diversity promotes ecosystem stability. *Ecology*, **93** (sp8), S223-S233. <http://dx.doi.org/10.1890/11-0426.1>
116. Duveneck, M.J., R.M. Scheller, M.A. White, S.D. Handler, and C. Ravenscroft, 2014: Climate change effects on northern Great Lake (USA) forests: A case for preserving diversity. *Ecosphere*, **5** (2), 1-26. <http://dx.doi.org/10.1890/ES13-00370.1>
117. McEwan, R.W., J.M. Dyer, and N. Pederson, 2011: Multiple interacting ecosystem drivers: Toward an encompassing hypothesis of oak forest dynamics across eastern North America. *Ecography*, **34** (2), 244-256. <http://dx.doi.org/10.1111/j.1600-0587.2010.06390.x>

118. Clark, J.S., L. Iverson, C.W. Woodall, C.D. Allen, D.M. Bell, D.C. Bragg, A.W. D'Amato, F.W. Davis, M.H. Hersh, I. Ibanez, S.T. Jackson, S. Matthews, N. Pederson, M. Peters, M.W. Schwartz, K.M. Waring, and N.E. Zimmermann, 2016: The impacts of increasing drought on forest dynamics, structure, and biodiversity in the United States. *Global Change Biology*, **22** (7), 2329-2352. <http://dx.doi.org/10.1111/gcb.13160>
119. Millar, C.I. and N.L. Stephenson, 2015: Temperate forest health in an era of emerging megadisturbance. *Science*, **349** (6250), 823-826. <http://dx.doi.org/10.1126/science.aaa9933>
120. Iverson, L.R., F.R. Thompson, S. Matthews, M. Peters, A. Prasad, W.D. Dijk, J. Fraser, W.J. Wang, B. Hanberry, H. He, M. Janowiak, P. Butler, L. Brandt, and C. Swanston, 2017: Multi-model comparison on the effects of climate change on tree species in the eastern U.S.: Results from an enhanced niche model and process-based ecosystem and landscape models. *Landscape Ecology*, **32** (7), 1327-1346. <http://dx.doi.org/10.1007/s10980-016-0404-8>
121. Jump, A.S., L. Cavin, and P.D. Hunter, 2010: Monitoring and managing responses to climate change at the retreating range edge of forest trees. *Journal of Environmental Monitoring*, **12** (10), 1791-1798. <http://dx.doi.org/10.1039/B923773A>
122. Jump, A.S., C. Mátyás, and J. Peñuelas, 2009: The altitude-for-latitude disparity in the range retractions of woody species. *Trends in Ecology & Evolution*, **24** (12), 694-701. <http://dx.doi.org/10.1016/j.tree.2009.06.007>
123. Frelich, L.E. and P.B. Reich, 2010: Will environmental changes reinforce the impact of global warming on the prairie-forest border of central North America? *Frontiers in Ecology and the Environment*, **8** (7), 371-378. <http://dx.doi.org/10.1890/080191>
124. Lenihan, J.M., D. Bachelet, R.P. Neilson, and R. Drapek, 2008: Simulated response of conterminous United States ecosystems to climate change at different levels of fire suppression, CO₂ emission rate, and growth response to CO₂. *Global and Planetary Change*, **64** (1-2), 16-25. <http://dx.doi.org/10.1016/j.gloplacha.2008.01.006>
125. Ma, W., J. Liang, J.R. Cumming, E. Lee, A.B. Welsh, J.V. Watson, and M. Zhou, 2016: Fundamental shifts of central hardwood forests under climate change. *Ecological Modelling*, **332**, 28-41. <http://dx.doi.org/10.1016/j.ecolmodel.2016.03.021>
126. Fisichelli, N.A., S.R. Abella, M. Peters, and F.J. Krist, 2014: Climate, trees, pests, and weeds: Change, uncertainty, and biotic stressors in eastern U.S. national park forests. *Forest Ecology and Management*, **327**, 31-39. <http://dx.doi.org/10.1016/j.foreco.2014.04.033>
127. Voggesser, G., K. Lynn, J. Daigle, F.K. Lake, and D. Ranco, 2013: Cultural impacts to tribes from climate change influences on forests. *Climatic Change*, **120** (3), 615-626. <http://dx.doi.org/10.1007/s10584-013-0733-4>
128. Grigal, D.F., 2000: Effects of extensive forest management on soil productivity. *Forest Ecology and Management*, **138** (1), 167-185. [http://dx.doi.org/10.1016/S0378-1127\(00\)00395-9](http://dx.doi.org/10.1016/S0378-1127(00)00395-9)
129. Rittenhouse, C.D. and A.R. Rissman, 2015: Changes in winter conditions impact forest management in north temperate forests. *Journal of Environmental Management*, **149**, 157-167. <http://dx.doi.org/10.1016/j.jenvman.2014.10.010>
130. Evans, A.M., M. Lynch, F. Clark, G.M. Mickel, K. Chapman, E.R. Tiller, and M. Haynes, 2016: Economic and Ecological Effects of Forest Practices and Harvesting Constraints on Wisconsin's Forest Resources and Economy. Forest Stewards Guild, Madison, WI, various pp. <https://councilonforestry.wi.gov/Documents/PracticesStudy/WFPSForestStewardsGuild2016.pdf>
131. Conrad IV, J.L., M.C. Demchik, M.M. Vokoun, A.M. Evans, and M.P. Lynch, 2017: Foresters' perceptions of the frequency, cost, and rationale for seasonal timber harvesting restrictions in Wisconsin. *Forest Science*. <http://dx.doi.org/10.5849/FS-2016-051>
132. Brandt, L.A., P.R. Butler, S.D. Handler, M.K. Janowiak, P.D. Shannon, and C.W. Swanston, 2017: Integrating science and management to assess forest ecosystem vulnerability to climate change. *Journal of Forestry*, **115** (3), 212-221. <http://dx.doi.org/10.5849/jof.15-147>
133. Brandt, L., A. Derby Lewis, R. Fahey, L. Scott, L. Darling, and C. Swanston, 2016: A framework for adapting urban forests to climate change. *Environmental Science & Policy*, **66**, 393-402. <http://dx.doi.org/10.1016/j.envsci.2016.06.005>

134. Janowiak, M.K., C.W. Swanston, L.M. Nagel, L.A. Brandt, P.R. Butler, S.D. Handler, P.D. Shannon, L.R. Iverson, S.N. Matthews, A. Prasad, and M.P. Peters, 2014: A practical approach for translating climate change adaptation principles into forest management actions. *Journal of Forestry*, **112** (5), 424-433. <http://dx.doi.org/10.5849/jof.13-094>
135. Swanston, C., M. Janowiak, L. Brandt, P. Butler, S.D. Handler, P.D. Shannon, A. Derby Lewis, K. Hall, R.T. Fahey, L. Scott, A. Kerber, J.W. Miesbauer, and L. Darling, 2016: Forest Adaptation Resources: Climate Change Tools and Approaches for Land Managers, 2nd ed. Gen. Tech. Rep. NRS-87-2. U.S. Department of Agriculture, Forest Service, Northern Research Station, Newtown Square, PA, 161 pp. https://www.fs.fed.us/nrs/pubs/gtr/gtr_nrs87-2.pdf
136. Vinyeta, K. and K. Lynn, 2013: Exploring the Role of Traditional Ecological Knowledge in Climate Change Initiatives. General Technical Report PNW-GTR-879. U.S. Department of Agriculture Pacific Northwest Research Station, Portland, OR, 37 pp. https://www.fs.fed.us/pnw/pubs/pnw_gtr879.pdf
137. Crausbay, S.D., A.R. Ramirez, S.L. Carter, M.S. Cross, K.R. Hall, D.J. Bathke, J.L. Betancourt, S. Colt, A.E. Cravens, M.S. Dalton, J.B. Dunham, L.E. Hay, M.J. Hayes, J. McEvoy, C.A. McNutt, M.A. Moritz, K.H. Nislow, N. Raheem, and T. Sanford, 2017: Defining ecological drought for the twenty-first century. *Bulletin of the American Meteorological Society*, **98** (12), 2543-2550. <http://dx.doi.org/10.1175/bams-d-16-0292.1>
138. Hall, K.R. and T.L. Root, 2012: Climate change and biodiversity in the Great Lakes Region from “fingerprints” of change to helping safeguard species. *Climate Change in the Great Lakes Region: Navigating an Uncertain Future*. Dietz, T. and D. Bidwell, Eds. Michigan State University Press, 63-96.
139. Hellmann, J.J., R. Grundel, C. Hoving, and G.W. Schuurman, 2016: A call to insect scientists: Challenges and opportunities of managing insect communities under climate change. *Current Opinion in Insect Science*, **17**, 92-97. <http://dx.doi.org/10.1016/j.cois.2016.08.005>
140. Dawson, T.P., S.T. Jackson, J.I. House, I.C. Prentice, and G.M. Mace, 2011: Beyond predictions: Biodiversity conservation in a changing climate. *Science*, **332** (6025), 53-58. <http://dx.doi.org/10.1126/science.1200303>
141. Ash, J.D., T.J. Givnish, and D.M. Waller, 2017: Tracking lags in historical plant species' shifts in relation to regional climate change. *Global Change Biology*, **23** (3), 1305-1315. <http://dx.doi.org/10.1111/gcb.13429>
142. Mayor, S.J., R.P. Guralnick, M.W. Tingley, J. Otegui, J.C. Withey, S.C. Elmendorf, M.E. Andrew, S. Leyk, I.S. Pearse, and D.C. Schneider, 2017: Increasing phenological asynchrony between spring green-up and arrival of migratory birds. *Scientific Reports*, **7** (1), 1902. <http://dx.doi.org/10.1038/s41598-017-02045-z>
143. Ewert, D., K. Hall, R. Smith, and P. Rodewald, 2015: Landbird stopover in the Great Lakes region: Integrating habitat use and climate change in conservation. *Phenological Synchrony and Bird Migration*. Wood, E.M. and J.L. Kellermann, Eds. CRC Press, 17-46.
144. Lyons, J., T.P. Parks, K.L. Minahan, and A.S. Ruesch, 2017: Evaluation of oxythermal metrics and benchmarks for the protection of cisco (*Coregonus artedii*) habitat quality and quantity in Wisconsin lakes. *Canadian Journal of Fisheries and Aquatic Sciences*, **75** (4), 600-608. <http://dx.doi.org/10.1139/cjfas-2017-0043>
145. Jacobson, P.C., T.S. Jones, P. Rivers, and D.L. Pereira, 2008: Field estimation of a lethal oxythermal niche boundary for adult ciscoes in Minnesota lakes. *Transactions of the American Fisheries Society*, **137** (5), 1464-1474. <http://dx.doi.org/10.1577/T07-148.1>
146. Jacobson, P.C., H.G. Stefan, and D.L. Pereira, 2010: Coldwater fish oxythermal habitat in Minnesota lakes: Influence of total phosphorus, July air temperature, and relative depth. *Canadian Journal of Fisheries and Aquatic Sciences*, **67** (12), 2002-2013. <http://dx.doi.org/10.1139/F10-115>
147. Magee, M.R. and C.H. Wu, 2017: Response of water temperatures and stratification to changing climate in three lakes with different morphometry. *Hydrology and Earth System Sciences*, **21** (12), 6253-6274. <http://dx.doi.org/10.5194/hess-21-6253-2017>
148. Magee, M.R. and C.H. Wu, 2017: Effects of changing climate on ice cover in three morphometrically different lakes. *Hydrological Processes*, **31** (2), 308-323. <http://dx.doi.org/10.1002/hyp.10996>

149. Honsey, A.E., S.B. Donabauer, and T.O. Höök, 2016: An analysis of lake morphometric and land-use characteristics that promote persistence of Cisco in Indiana. *Transactions of the American Fisheries Society*, **145** (2), 363-373. <http://dx.doi.org/10.1080/00028487.2015.1125949>
150. Hewitt, B., L. Lopez, K. Gaibisels, A. Murdoch, S. Higgins, J. Magnuson, A. Paterson, J. Rusak, H. Yao, and S. Sharma, 2018: Historical trends, drivers, and future projections of ice phenology in small north temperate lakes in the Laurentian Great Lakes region. *Water*, **10** (1), [16]. <http://dx.doi.org/10.3390/w10010070>
151. Herb, W.R., L.B. Johnson, P.C. Jacobson, and H.G. Stefan, 2014: Projecting cold-water fish habitat in lakes of the glacial lakes region under changing land use and climate regimes. *Canadian Journal of Fisheries and Aquatic Sciences*, **71** (9), 1334-1348. <http://dx.doi.org/10.1139/cjfas-2013-0535>
152. Jiang, L., X. Fang, H.G. Stefan, P.C. Jacobson, and D.L. Pereira, 2012: Oxythermal habitat parameters and identifying cisco refuge lakes in Minnesota under future climate scenarios using variable benchmark periods. *Ecological Modelling*, **232**, 14-27. <http://dx.doi.org/10.1016/j.ecolmodel.2012.02.014>
153. Magee, M.R., P.B. McIntyre, and C.H. Wu, 2017: Modeling oxythermal stress for cool-water fishes in lakes using a cumulative dosage approach. *Canadian Journal of Fisheries and Aquatic Sciences*, **75** (8), 1303-1312. <http://dx.doi.org/10.1139/cjfas-2017-0260>
154. Jiang, L. and X. Fang, 2016: Simulation and validation of cisco lethal conditions in Minnesota lakes under past and future climate scenarios using constant survival limits. *Water*, **8** (7), 279. <http://dx.doi.org/10.3390/w8070279>
155. Petersen, B., K.R. Hall, K. Kahl, and P.J. Doran, 2013: Research articles: In their own words: Perceptions of climate change adaptation from the Great Lakes region's resource management community. *Environmental Practice*, **15** (4), 377-392. <http://dx.doi.org/10.1017/S1466046613000446>
156. Anhalt-Depies, C.M., T.G. Knoot, A.R. Rissman, A.K. Sharp, and K.J. Martin, 2016: Understanding climate adaptation on public lands in the Upper Midwest: Implications for monitoring and tracking progress. *Environmental Management*, **57** (5), 987-997. <http://dx.doi.org/10.1007/s00267-016-0673-7>
157. Brook, B.W., N.S. Sodhi, and C.J.A. Bradshaw, 2008: Synergies among extinction drivers under global change. *Trends in Ecology & Evolution*, **23** (8), 453-460. <http://dx.doi.org/10.1016/j.tree.2008.03.011>
158. Schloss, C.A., T.A. Nunez, and J.J. Lawler, 2012: Dispersal will limit ability of mammals to track climate change in the Western Hemisphere. *Proceedings of the National Academy of Sciences of the United States of America*, **109** (22), 8606-11. <http://dx.doi.org/10.1073/pnas.1116791109>
159. Carroll, C., J.J. Lawler, D.R. Roberts, and A. Hamann, 2015: Biotic and climatic velocity identify contrasting areas of vulnerability to climate change. *PLOS ONE*, **10** (10), e0140486. <http://dx.doi.org/10.1371/journal.pone.0140486>
160. Hellmann, J.J., J.E. Byers, B.G. Bierwagen, and J.S. Dukes, 2008: Five potential consequences of climate change for invasive species. *Conservation Biology*, **22** (3), 534-543. <http://dx.doi.org/10.1111/j.1523-1739.2008.00951.x>
161. Cline, T.J., J.F. Kitchell, V. Bennington, G.A. McKinley, E.K. Moody, and B.C. Weidel, 2014: Climate impacts on landlocked sea lamprey: Implications for host-parasite interactions and invasive species management. *Ecosphere*, **5** (6), 1-13. <http://dx.doi.org/10.1890/ES14-00059.1>
162. Forrest, J.R.K., 2016: Complex responses of insect phenology to climate change. *Current Opinion in Insect Science*, **17**, 49-54. <http://dx.doi.org/10.1016/j.cois.2016.07.002>
163. Holmstrom, R.M., J.R. Etterson, and D.J. Schimpf, 2010: Dune restoration introduces genetically distinct American beachgrass, *Ammophila breviligulata*, into a threatened local population. *Restoration Ecology*, **18** (s2), 426-437. <http://dx.doi.org/10.1111/j.1526-100X.2009.00593.x>
164. IPBES, 2017: The Assessment Report on Pollinators, Pollination and Food Production. Potts, S.G., V. Imperatriz-Fonseca, and H.T. Ngo, Eds. Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES), Bonn, Germany, 502 pp. https://www.ipbes.net/sites/default/files/downloads/pdf/individual_chapters_pollination_20170305.pdf

165. Potts, S.G., V. Imperatriz-Fonseca, H.T. Ngo, M.A. Aizen, J.C. Biesmeijer, T.D. Breeze, L.V. Dicks, L.A. Garibaldi, R. Hill, J. Settele, and A.J. Vanbergen, 2016: Safeguarding pollinators and their values to human well-being. *Nature*, **540**, 220-229. <http://dx.doi.org/10.1038/nature20588>
166. Vanbergen, A.J., A. Espíndola, and M.A. Aizen, 2018: Risks to pollinators and pollination from invasive alien species. *Nature Ecology & Evolution*, **2** (1), 16-25. <http://dx.doi.org/10.1038/s41559-017-0412-3>
167. Flockhart, D.T.T., J.-B. Pichancourt, D.R. Norris, and T.G. Martin, 2015: Unravelling the annual cycle in a migratory animal: Breeding-season habitat loss drives population declines of monarch butterflies. *Journal of Animal Ecology*, **84** (1), 155-165. <http://dx.doi.org/10.1111/1365-2656.12253>
168. Koh, I., E.V. Lonsdorf, N.M. Williams, C. Brittain, R. Isaacs, J. Gibbs, and T.H. Ricketts, 2016: Modeling the status, trends, and impacts of wild bee abundance in the United States. *Proceedings of the National Academy of Sciences of the United States of America*, **113** (1), 140-145. <http://dx.doi.org/10.1073/pnas.1517685113>
169. Landis, D.A., 2017: Productive engagement with agriculture essential to monarch butterfly conservation. *Environmental Research Letters*, **12** (10), 101003. <http://dx.doi.org/10.1088/1748-9326/aa825c>
170. Kovács-Hostyánszki, A., A. Espíndola, A.J. Vanbergen, J. Settele, C. Kremen, and L.V. Dicks, 2017: Ecological intensification to mitigate impacts of conventional intensive land use on pollinators and pollination. *Ecology Letters*, **20** (5), 673-689. <http://dx.doi.org/10.1111/ele.12762>
171. Kerr, J.T., A. Pindar, P. Galpern, L. Packer, S.G. Potts, S.M. Roberts, P. Rasmont, O. Schweiger, S.R. Colla, L.L. Richardson, D.L. Wagner, L.F. Gall, D.S. Sikes, and A. Pantoja, 2015: Climate change impacts on bumblebees converge across continents. *Science*, **349** (6244), 177-180. <http://dx.doi.org/10.1126/science.aaa7031>
172. Bartomeus, I., J.S. Ascher, J. Gibbs, B.N. Danforth, D.L. Wagner, S.M. Hedtke, and R. Winfree, 2013: Historical changes in northeastern US bee pollinators related to shared ecological traits. *Proceedings of the National Academy of Sciences of the United States of America*, **110** (12), 4656-4660. <http://dx.doi.org/10.1073/pnas.1218503110>
173. Thogmartin, W.E., L. López-Hoffman, J. Rohweder, J. Diffendorfer, R. Drum, D. Semmens, S. Black, I. Caldwell, D. Cotter, P. Drobney, L.L. Jackson, M. Gale, D. Helmers, S. Hilburger, E. Howard, K. Oberhauser, J. Pleasants, B. Semmens, O. Taylor, P. Ward, J.F. Weltzin, and R. Wiederholt, 2017: Restoring monarch butterfly habitat in the Midwestern US: "All hands on deck." *Environmental Research Letters*, **12** (7), 074005. <http://dx.doi.org/10.1088/1748-9326/aa7637>
174. Kaiser-Bunbury, C.N., J. Mougil, A.E. Whittington, T. Valentin, R. Gabriel, J.M. Olesen, and N. Blüthgen, 2017: Ecosystem restoration strengthens pollination network resilience and function. *Nature*, **542**, 223-227. <http://dx.doi.org/10.1038/nature21071>
175. Toniello, R.K. and D.J. Larkin, 2018: Habitat restoration benefits wild bees: A meta-analysis. *Journal of Applied Ecology*, **55** (2), 582-590. <http://dx.doi.org/10.1111/1365-2664.13012>
176. Mao, D. and K.A. Cherkauer, 2009: Impacts of land-use change on hydrologic responses in the Great Lakes region. *Journal of Hydrology*, **374** (1), 71-82. <http://dx.doi.org/10.1016/j.jhydrol.2009.06.016>
177. Mishra, V., K.A. Cherkauer, D. Niyogi, M. Lei, B.C. Pijanowski, D.K. Ray, L.C. Bowling, and G. Yang, 2010: A regional scale assessment of land use/land cover and climatic changes on water and energy cycle in the upper Midwest United States. *International Journal of Climatology*, **30** (13), 2025-2044. <http://dx.doi.org/10.1002/joc.2095>
178. Mitsch, W.J. and J.G. Gosselink, 2015: Appendix A. Wetland losses by state in the United States, 1780s-1980s. *Wetlands*, 5th ed. Wiley, Hoboken, NJ, 701-702.
179. Garris, H.W., R.J. Mitchell, L.H. Fraser, and L.R. Barrett, 2015: Forecasting climate change impacts on the distribution of wetland habitat in the Midwestern United States. *Global Change Biology*, **21** (2), 766-776. <http://dx.doi.org/10.1111/gcb.12748>
180. Wing, O.E.J., P.D. Bates, A.M. Smith, C.C. Sampson, K.A. Johnson, J. Fargione, and P. Morefield, 2018: Estimates of present and future flood risk in the conterminous United States. *Environmental Research Letters*, **13** (3), 034023. <http://dx.doi.org/10.1088/1748-9326/aaac65>

181. Brink, E., T. Aalders, D. Ádám, R. Feller, Y. Henselek, A. Hoffmann, K. Ibe, A. Matthey-Doret, M. Meyer, N.L. Negrut, A.-L. Rau, B. Riewerts, L. von Schuckmann, S. Törnros, H. von Wehrden, D.J. Abson, and C. Wamsler, 2016: Cascades of green: A review of ecosystem-based adaptation in urban areas. *Global Environmental Change*, **36**, 111-123. <http://dx.doi.org/10.1016/j.gloenvcha.2015.11.003>
182. Kousky, C., 2010: Using natural capital to reduce disaster risk. *Journal of Natural Resources Policy Research*, **2** (4), 343-356. <http://dx.doi.org/10.1080/19390459.2010.511451>
183. Kousky, C., S. Olmstead, M. Walls, A. Stern, and M. Macauley, 2011: The Role of Land Use in Adaptation to Increased Precipitation and Flooding: A Case Study in Wisconsin's Lower Fox River Basin. RFF Report. Resources for the Future, Washington, DC, 72 pp. [http://www.rff.org/files/sharepoint/WorkImages/Download/RFF-Rpt-Kousky%20etal%20GreatLakes%20\(2\).pdf](http://www.rff.org/files/sharepoint/WorkImages/Download/RFF-Rpt-Kousky%20etal%20GreatLakes%20(2).pdf)
184. Meerow, S. and J.P. Newell, 2017: Spatial planning for multifunctional green infrastructure: Growing resilience in Detroit. *Landscape and Urban Planning*, **159**, 62-75. <http://dx.doi.org/10.1016/j.landurbplan.2016.10.005>
185. Griscom, B.W., J. Adams, P.W. Ellis, R.A. Houghton, G. Lomax, D.A. Miteva, W.H. Schlesinger, D. Shoch, J.V. Siikamäki, P. Smith, P. Woodbury, C. Zganjar, A. Blackman, J. Campari, R.T. Conant, C. Delgado, P. Elias, T. Gopalakrishna, M.R. Hamsik, M. Herrero, J. Kiesecker, E. Landis, L. Laestadius, S.M. Leavitt, S. Minnemeyer, S. Polasky, P. Potapov, F.E. Putz, J. Sanderman, M. Silvius, E. Wollenberg, and J. Fargione, 2017: Natural climate solutions. *Proceedings of the National Academy of Sciences of the United States of America*, **114** (44), 11645-11650. <http://dx.doi.org/10.1073/pnas.1710465114>
186. Dierkes, C., 2012: From farm fields to wetlands. *Twine Line*, **34** (4), 3-5. <https://ohioseagrant.osu.edu/p/3un80>
187. Staff Writer, 2015: Country life: Wetland rehabilitation effort paying off. Ohio Ag Net, September 30. Agri Communicators Inc., Columbus, OH. <http://ocj.com/2015/09/wetland-rehabilitation-effort-paying-off/>
188. Wisconsin Sea Grant Institute, 2013: Great Lakes and Wisconsin Water Facts: Great Lakes and Fresh Water. University of Wisconsin Sea Grant Institute, Madison, WI. <http://www.seagrant.wisc.edu/Home/AboutUsSection/PressRoom/Details.aspx?PostID=796>
189. Scott, R.W. and F.A. Huff, 1996: Impacts of the Great Lakes on regional climate conditions. *Journal of Great Lakes Research*, **22** (4), 845-863. [http://dx.doi.org/10.1016/S0380-1330\(96\)71006-7](http://dx.doi.org/10.1016/S0380-1330(96)71006-7)
190. Notaro, M., K. Holman, A. Zarrin, E. Fluck, S. Vavrus, and V. Bennington, 2013: Influence of the Laurentian Great Lakes on regional climate. *Journal of Climate*, **26** (3), 789-804. <http://dx.doi.org/10.1175/jcli-d-12-00140.1>
191. McDermid, J.L., S.K. Dickin, C.L. Winsborough, H. Switzman, S. Barr, J.A. Gleeson, G. Krantzberg, and P.A. Gray, 2015: State of Climate Change Science in the Great Lakes Basin: A Focus on Climatological, Hydrological, and Ecological Effects. Prepared Jointly by the Ontario Climate Consortium and Ontario Ministry of Natural Resources and Forestry to Advise Annex 9—Climate Change Impacts Under the Great Lakes Water Quality Agreement, October 2015. https://binational.net/wp-content/uploads/2016/09/OCC_GreatLakes_Report_ExecSummary%20ENGLISH.pdf
192. Mason, L.A., C.M. Riseng, A.D. Gronewold, E.S. Rutherford, J. Wang, A. Clites, S.D.P. Smith, and P.B. McIntyre, 2016: Fine-scale spatial variation in ice cover and surface temperature trends across the surface of the Laurentian Great Lakes. *Climatic Change*, **138** (1), 71-83. <http://dx.doi.org/10.1007/s10584-016-1721-2>
193. Easterling, D.R., K.E. Kunkel, J.R. Arnold, T. Knutson, A.N. LeGrande, L.R. Leung, R.S. Vose, D.E. Waliser, and M.F. Wehner, 2017: Precipitation change in the United States. *Climate Science Special Report: Fourth National Climate Assessment, Volume I*. Wuebbles, D.J., D.W. Fahey, K.A. Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock, Eds. U.S. Global Change Research Program, Washington, DC, USA, 207-230. <http://dx.doi.org/10.7930/J0H993CC>
194. Vavrus, S., M. Notaro, and A. Zarrin, 2013: The role of ice cover in heavy lake-effect snowstorms over the Great Lakes Basin as simulated by RegCM4. *Monthly Weather Review*, **141** (1), 148-165. <http://dx.doi.org/10.1175/mwr-d-12-00107.1>

195. Wright, D.M., D.J. Posselt, and A.L. Steiner, 2013: Sensitivity of lake-effect snowfall to lake ice cover and temperature in the Great Lakes region. *Monthly Weather Review*, **141** (2), 670-689. <http://dx.doi.org/10.1175/mwr-d-12-00038.1>
196. O'Reilly, C.M., S. Sharma, D.K. Gray, S.E. Hampton, J.S. Read, R.J. Rowley, P. Schneider, J.D. Lenters, P.B. McIntyre, B.M. Kraemer, G.A. Weyhenmeyer, D. Straile, B. Dong, R. Adrian, M.G. Allan, O. Anneville, L. Arvola, J. Austin, J.L. Bailey, J.S. Baron, J.D. Brookes, E. de Eyto, M.T. Dokulil, D.P. Hamilton, K. Havens, A.L. Hetherington, S.N. Higgins, S. Hook, L.R. Izmet'seva, K.D. Joehnk, K. Kangur, P. Kasprzak, M. Kumagai, E. Kuusisto, G. Leshkevich, D.M. Livingstone, S. MacIntyre, L. May, J.M. Melack, D.C. Mueller-Navarra, M. Naumenko, P. Noges, T. Noges, R.P. North, P.-D. Plisnier, A. Rigosi, A. Rimmer, M. Rogora, L.G. Rudstam, J.A. Rusak, N. Salmaso, N.R. Samal, D.E. Schindler, S.G. Schladow, M. Schmid, S.R. Schmidt, E. Silow, M.E. Soylu, K. Teubner, P. Verburg, A. Voutilainen, A. Watkinson, C.E. Williamson, and G. Zhang, 2015: Rapid and highly variable warming of lake surface waters around the globe. *Geophysical Research Letters*, **42** (24), 10,773-10,781. <http://dx.doi.org/10.1002/2015GL066235>
197. Austin, J. and S. Colman, 2008: A century of temperature variability in Lake Superior. *Limnology and Oceanography*, **53** (6), 2724-2730. <http://dx.doi.org/10.4319/lo.2008.53.6.2724>
198. Notaro, M., V. Bennington, and B. Lofgren, 2015: Dynamical downscaling-based projections of Great Lakes water levels. *Journal of Climate*, **28** (24), 9721-9745. <http://dx.doi.org/10.1175/jcli-d-14-00847.1>
199. MacKay, M. and F. Seglenieks, 2012: On the simulation of Laurentian Great Lakes water levels under projections of global climate change. *Climatic Change*, **117** (1-2), 55-67. <http://dx.doi.org/10.1007/s10584-012-0560-z>
200. Lofgren, B.M. and J. Rouhana, 2016: Physically plausible methods for projecting changes in Great Lakes water levels under climate change scenarios. *Journal of Hydrometeorology*, **17** (8), 2209-2223. <http://dx.doi.org/10.1175/jhm-d-15-0220.1>
201. Gronewold, A.D., A.H. Clites, J. Bruxer, K.W. Kompoltowicz, J.P. Smith, T.S. Hunter, and C. Wong, 2015: Water levels surge on Great Lakes. *Eos, Earth & Space Science News*, **61**, 14-17. <http://dx.doi.org/10.1029/2015EO026023>
202. Trumpickas, J., B.J. Shuter, and C.K. Minns, 2009: Forecasting impacts of climate change on Great Lakes surface water temperatures. *Journal of Great Lakes Research*, **35** (3), 454-463. <http://dx.doi.org/10.1016/j.jglr.2009.04.005>
203. Austin, J.A. and S.M. Colman, 2007: Lake Superior summer water temperatures are increasing more rapidly than regional air temperatures: A positive ice-albedo feedback. *Geophysical Research Letters*, **34** (6), L06604. <http://dx.doi.org/10.1029/2006GL029021>
204. Croley II, T.E., 2003: Great Lakes Climate Change Hydrologic Impact Assessment I.J.C. Lake Ontario-St. Lawrence River Regulation Study. NOAA Technical Memorandum GLERL-126. Great Lakes Environmental Research Laboratory, Ann Arbor, MI, 77 pp. https://www.glerl.noaa.gov/pubs/tech_reports/glerl-126/tm-126.pdf
205. Magnuson, J.J., K.E. Webster, R.A. Assel, C.J. Bowser, P.J. Dillon, J.G. Eaton, H.E. Evans, E.J. Fee, R.I. Hall, L.R. Mortsch, D.W. Schindler, and F.H. Quinn, 1997: Potential effects of climate changes on aquatic systems: Laurentian Great Lakes and Precambrian shield region. *Hydrological Processes*, **11** (8), 825-871. [http://dx.doi.org/10.1002/\(SICI\)1099-1085\(19970630\)11:8<825::AID-HYP509>3.0.CO;2-G](http://dx.doi.org/10.1002/(SICI)1099-1085(19970630)11:8<825::AID-HYP509>3.0.CO;2-G)
206. Jones, M.L., B.J. Shuter, Y. Zhao, and J.D. Stockwell, 2006: Forecasting effects of climate change on Great Lakes fisheries: Models that link habitat supply to population dynamics can help. *Canadian Journal of Fisheries and Aquatic Sciences*, **63** (2), 457-468. <http://dx.doi.org/10.1139/f05-239>
207. Trebitz, A.S. and J.C. Hoffman, 2015: Coastal wetland support of Great Lakes fisheries: Progress from concept to quantification. *Transactions of the American Fisheries Society*, **144** (2), 352-372. <http://dx.doi.org/10.1080/00028487.2014.982257>
208. Trebitz, A.S., J.C. Brazner, N.P. Danz, M.S. Pearson, G.S. Peterson, D.K. Tanner, D.L. Taylor, C.W. West, and T.P. Hollenhorst, 2009: Geographic, anthropogenic, and habitat influences on Great Lakes coastal wetland fish assemblages. *Canadian Journal of Fisheries and Aquatic Sciences*, **66** (8), 1328-1342. <http://dx.doi.org/10.1139/F09-089>

209. Brenden, T.O., R.W. Brown, M.P. Ebener, K. Reid, and T.J. Newcomb, 2013: Great Lakes commercial fisheries: Historical overview and prognoses for the future. *Great Lakes Fisheries Policy and Management: A Binational Perspective*, 2nd ed. Taylor, W.W., A.J. Lynch, and N.J. Leonard, Eds. Michigan State University Press, Lansing, MI, 339-397.
210. Ludsin, S.A., M.W. Kershner, K.A. Blocksom, R.L. Knight, and R.A. Stein, 2001: Life after death in Lake Erie: Nutrient controls drive fish species richness, rehabilitation. *Ecological Applications*, **11** (3), 731-746. [http://dx.doi.org/10.1890/1051-0761\(2001\)011\[0731:LADILE\]2.0.CO;2](http://dx.doi.org/10.1890/1051-0761(2001)011[0731:LADILE]2.0.CO;2)
211. Dolan, D.M. and S.C. Chapra, 2012: Great Lakes total phosphorus revisited: 1. Loading analysis and update (1994–2008). *Journal of Great Lakes Research*, **38** (4), 730-740. <http://dx.doi.org/10.1016/j.jglr.2012.10.001>
212. Carmichael, W.W. and G.L. Boyer, 2016: Health impacts from cyanobacteria harmful algae blooms: Implications for the North American Great Lakes. *Harmful Algae*, **54**, 194-212. <http://dx.doi.org/10.1016/j.hal.2016.02.002>
213. Chapra, S.C., B. Boehlert, C. Fant, V.J. Bierman, J. Henderson, D. Mills, D.M.L. Mas, L. Rennels, L. Jantarasami, J. Martinich, K.M. Strzepak, and H.W. Paerl, 2017: Climate change impacts on harmful algal blooms in U.S. freshwaters: A screening-level assessment. *Environmental Science & Technology*, **51** (16), 8933-8943. <http://dx.doi.org/10.1021/acs.est.7b01498>
214. Collingsworth, P.D., D.B. Bunnell, M.W. Murray, Y.-C. Kao, Z.S. Feiner, R.M. Claramunt, B.M. Lofgren, T.O. Höök, and S.A. Ludsin, 2017: Climate change as a long-term stressor for the fisheries of the Laurentian Great Lakes of North America. *Reviews in Fish Biology and Fisheries*, **27** (2), 363-391. <http://dx.doi.org/10.1007/s11160-017-9480-3>
215. Allan, J.D., M. Palmer, and N.L. Poff, 2005: Climate change and freshwater ecosystems. *Climate Change and Biodiversity*. Lovejoy, T.E. and L. Hannah, Eds. Yale University Press, Ann Arbor, MI, 274-290.
216. Kao, Y.-C., C.P. Madenjian, D.B. Bunnell, B.M. Lofgren, and M. Perroud, 2015: Potential effects of climate change on the growth of fishes from different thermal guilds in Lakes Michigan and Huron. *Journal of Great Lakes Research*, **41** (2), 423-435. <http://dx.doi.org/10.1016/j.jglr.2015.03.012>
217. Madenjian, C.P., R. O’Gorman, D.B. Bunnell, R.L. Argyle, E.F. Roseman, D.M. Warner, J.D. Stockwell, and M.A. Stapanian, 2008: Adverse effects of alewives on Laurentian Great Lakes fish communities. *North American Journal of Fisheries Management*, **28** (1), 263-282. <http://dx.doi.org/10.1577/M07-012.1>
218. Higgins, S.N. and M.J.V. Zanden, 2010: What a difference a species makes: A meta-analysis of dreissenid mussel impacts on freshwater ecosystems. *Ecological Monographs*, **80** (2), 179-196. <http://dx.doi.org/10.1890/09-1249.1>
219. Hansen, M.J., C.P. Madenjian, J.W. Slade, T.B. Steeves, P.R. Almeida, and B.R. Quintella, 2016: Population ecology of the sea lamprey (*Petromyzon marinus*) as an invasive species in the Laurentian Great Lakes and an imperiled species in Europe. *Reviews in Fish Biology and Fisheries*, **26** (3), 509-535. <http://dx.doi.org/10.1007/s11160-016-9440-3>
220. Millerd, F., 2011: The potential impact of climate change on Great Lakes international shipping. *Climatic Change*, **104** (3-4), 629-652. <http://dx.doi.org/10.1007/s10584-010-9872-z>
221. Howk, F., 2009: Changes in Lake Superior ice cover at Bayfield, Wisconsin. *Journal of Great Lakes Research*, **35** (1), 159-162. <http://dx.doi.org/10.1016/j.jglr.2008.11.002>
222. Forbes, D.L., G.K. Manson, R. Chagnon, S.M. Solomon, J.J.v.d. Sanden, and T.L. Lynds, 2002: Nearshore ice and climate change in the southern Gulf of St. Lawrence. *Ice in the Environment: Proceedings of the 16th IAHR International Symposium on Ice*, Dunedin, New Zealand, December 2-6, 344-351.
223. Larsen, A., A. Derby Lewis, O. Lyandres, T. Chen, and K. Frank, 2014: Developing a Community of Climate-Informed Conservation Practitioners to Protect a Priority Landscape in Illinois and Wisconsin. *Great Lakes Integrated Sciences + Assessments (GLISA)*, 18 pp. http://glisa.umich.edu/media/files/projectreports/GLISA_ProjRep_ILWI_Ravines.pdf
224. Briley, L., D. Brown, and S.E. Kalafatis, 2015: Overcoming barriers during the co-production of climate information for decision-making. *Climate Risk Management*, **9**, 41-49. <http://dx.doi.org/10.1016/j.crm.2015.04.004>
225. Samples, A., 2015: Engaging marina and harbor operators in climate adaptation. *Michigan Journal of Sustainability*, **3**, 65-72. <http://dx.doi.org/10.3998/mjs.12333712.0003.004>

226. City of Chicago, 2008: Chicago Climate Action Plan: Our City. Our Future. 57 pp. <http://www.chicagoclimataction.org/filebin/pdf/finalreport/CCAPREPORTFINALv2.pdf>
227. Norton, R.K., N.P. David, S. Buckman, and P.D. Koman, 2018: Overlooking the coast: Limited local planning for coastal area management along Michigan's Great Lakes. *Land Use Policy*, **71**, 183-203. <http://dx.doi.org/10.1016/j.landusepol.2017.11.049>
228. Gronlund, C.J., V.J. Berrocal, J.L. White-Newsome, K.C. Conlon, and M.S. O'Neill, 2015: Vulnerability to extreme heat by socio-demographic characteristics and area green space among the elderly in Michigan, 1990-2007. *Environmental Research*, **136**, 449-461. <http://dx.doi.org/10.1016/j.envres.2014.08.042>
229. Fann, N., C.G. Nolte, P. Dolwick, T.L. Spero, A. Curry Brown, S. Phillips, and S. Anenberg, 2015: The geographic distribution and economic value of climate change-related ozone health impacts in the United States in 2030. *Journal of the Air & Waste Management Association*, **65** (5), 570-580. <http://dx.doi.org/10.1080/10962247.2014.996270>
230. Ziska, L., K. Knowlton, C. Rogers, D. Dalan, N. Tierney, M.A. Elder, W. Filley, J. Shropshire, L.B. Ford, C. Hedberg, P. Fleetwood, K.T. Hovanky, T. Kavanaugh, G. Fulford, R.F. Vrtis, J.A. Patz, J. Portnoy, F. Coates, L. Bielory, and D. Frenz, 2011: Recent warming by latitude associated with increased length of ragweed pollen season in central North America. *Proceedings of the National Academy of Sciences of the United States of America*, **108** (10), 4248-4251. <http://dx.doi.org/10.1073/pnas.1014107108>
231. Fann, N., T. Brennan, P. Dolwick, J.L. Gamble, V. Ilacqua, L. Kolb, C.G. Nolte, T.L. Spero, and L. Ziska, 2016: Ch. 3: Air quality impacts. *The Impacts of Climate Change on Human Health in the United States: A Scientific Assessment*. U.S. Global Change Research Program, Washington, DC, 69-98. <http://dx.doi.org/10.7930/J0GQ6VP6>
232. Palecki, M.A., S.A. Changnon, and K.E. Kunkel, 2001: The nature and impacts of the July 1999 heat wave in the midwestern United States: Learning from the lessons of 1995. *Bulletin of the American Meteorological Society*, **82**, 1353-1368. [http://dx.doi.org/10.1175/1520-0477\(2001\)082<1353:TNAIOT>2.3.CO;2](http://dx.doi.org/10.1175/1520-0477(2001)082<1353:TNAIOT>2.3.CO;2)
233. Bobb, J.F., R.D. Peng, M.L. Bell, and F. Dominici, 2014: Heat-related mortality and adaptation to heat in the United States. *Environmental Health Perspectives*, **122** (8), 811-816. <http://dx.doi.org/10.1289/ehp.1307392>
234. Hondula, D.M., R.E. Davis, M.V. Saha, C.R. Wegner, and L.M. Veazey, 2015: Geographic dimensions of heat-related mortality in seven U.S. cities. *Environmental Research*, **138**, 439-452. <http://dx.doi.org/10.1016/j.envres.2015.02.033>
235. Jagai, J.S., E. Grossman, L. Navon, A. Sambanis, and S. Dorevitch, 2017: Hospitalizations for heat-stress illness varies between rural and urban areas: An analysis of Illinois data, 1987-2014. *Environmental Health*, **16** (1), 38. <http://dx.doi.org/10.1186/s12940-017-0245-1>
236. Sheridan, S.C. and P.G. Dixon, 2017: Spatiotemporal trends in human vulnerability and adaptation to heat across the United States. *Anthropocene*, **20**, 61-73. <http://dx.doi.org/10.1016/j.ancene.2016.10.001>
237. Chew, G.L., J. Wilson, F.A. Rabito, F. Grimsley, S. Iqbal, T. Reponen, M.L. Muilenberg, P.S. Thorne, D.G. Dearborn, and R.L. Morley, 2006: Mold and endotoxin levels in the aftermath of Hurricane Katrina: A pilot project of homes in New Orleans undergoing renovation. *Environmental Health Perspectives*, **114** (12), 1883-1889. <http://dx.doi.org/10.1289/ehp.9258>
238. Adeola, F.O., 2009: Mental health & psychosocial distress sequelae of Katrina: An empirical study of survivors. *Human Ecology Review*, **16** (2), 195-210. <http://www.jstor.org/stable/24707543>
239. Vins, H., J. Bell, S. Saha, and J. Hess, 2015: The mental health outcomes of drought: A systematic review and causal process diagram. *International Journal of Environmental Research and Public Health*, **12** (10), 13251. <http://dx.doi.org/10.3390/ijerph121013251>
240. Uejio, C.K., M. Christenson, C. Moran, and M. Gorelick, 2017: Drinking-water treatment, climate change, and childhood gastrointestinal illness projections for northern Wisconsin (USA) communities drinking untreated groundwater. *Hydrogeology Journal*, **25** (4), 969-979. <http://dx.doi.org/10.1007/s10040-016-1521-9>
241. Drayna, P., S.L. McLellan, P. Simpson, S.-H. Li, and M.H. Gorelick, 2010: Association between rainfall and pediatric emergency department visits for acute gastrointestinal illness. *Environmental Health Perspectives*, **118** (10), 1439-1443. <http://dx.doi.org/10.1289/ehp.0901671>

242. Hahn, M.B., A.J. Monaghan, M.H. Hayden, R.J. Eisen, M.J. Delorey, N.P. Lindsey, R.S. Nasci, and M. Fischer, 2015: Meteorological conditions associated with increased incidence of West Nile virus disease in the United States, 2004–2012. *The American Journal of Tropical Medicine and Hygiene*, **92** (5), 1013-1022. <http://dx.doi.org/10.4269/ajtmh.14-0737>
243. Lantos, P.M., J. Tsao, L.E. Nigrovic, P.G. Auwaerter, V.G. Fowler, F. Ruffin, E. Foster, and G. Hickling, 2017: Geographic expansion of Lyme disease in Michigan, 2000–2014. *Open Forum Infectious Diseases*, **4** (1), Art. ofw269. <http://dx.doi.org/10.1093/ofid/ofw269>
244. Rajkovich, N.B., 2016: A system of professions approach to reducing heat exposure in Cuyahoga County, Ohio. *Michigan Journal of Sustainability*, **4**, 81-101. <http://dx.doi.org/10.3998/mjs.12333712.0004.007>
245. Cameron, L., A. Ferguson, R. Walker, L. Briley, and D. Brown, 2015: Michigan Climate and Health Profile Report 2015: Building Resilience Against Climate Effects on Michigan's Health. Michigan Department of Health & Human Services, Lansing, MI, 97 pp. http://www.michigan.gov/documents/mdhhs/MI_Climate_and_Health_Profile_517517_7.pdf
246. BRACE-Illinois, 2016: Climate and Health in Illinois. University of Illinois at Chicago School of Public Health, Chicago, IL, 15 pp. <http://www.dph.illinois.gov/sites/default/files/publications/publicationsopclimatehealthreport.pdf>
247. Minnesota Department of Health, 2015: Minnesota Climate and Health Profile Report 2015: An Assessment of Climate Change Impacts on the Health & Well-Being of Minnesotans. Minnesota Department of Health, St. Paul, MN, 100 pp. <http://www.health.state.mn.us/divs/climatechange/docs/mnprofile2015.pdf>
248. Wisconsin Climate and Health Program, 2015: Understanding the Link Between Climate and Health. P-00709. Wisconsin Department of Health Services, Madison, WI, 2 pp. <https://www.dhs.wisconsin.gov/publications/p0/p00709.pdf>
249. Council for State and Territorial Epidemiologists (CSTE), 2016: Heat-Related Illness Syndrome Query: A Guidance Document for Implementing Heat-Related Illness Syndromic Surveillance in Public Health Practice. CSTE, 12 pp. http://c.ymcdn.com/sites/www.cste.org/resource/resmgr/pdfs/pdfs2/CSTE_Heat_Syndrome_Case_Defi.pdf
250. Hahn, M.B., R.J. Eisen, L. Eisen, K.A. Boegler, C.G. Moore, J. McAllister, H.M. Savage, and J.-P. Mutebi, 2016: Reported distribution of *Aedes (Stegomyia) aegypti* and *Aedes (Stegomyia) albopictus* in the United States, 1995–2016 (Diptera: Culicidae). *Journal of Medical Entomology*, **53** (5), 1169-1175. <http://dx.doi.org/10.1093/jme/tjw072>
251. Abel, D., T. Holloway, M. Harkey, A. Rrushaj, G. Brinkman, P. Duran, M. Janssen, and P. Denholm, 2018: Potential air quality benefits from increased solar photovoltaic electricity generation in the Eastern United States. *Atmospheric Environment*, **175**, 65-74. <http://dx.doi.org/10.1016/j.atmosenv.2017.11.049>
252. Maizlish, N., J. Woodcock, S. Co, B. Ostro, A. Fanai, and D. Fairley, 2013: Health cobenefits and transportation-related reductions in greenhouse gas emissions in the San Francisco Bay area. *American Journal of Public Health*, **103** (4), 703-709. <http://dx.doi.org/10.2105/ajph.2012.300939>
253. Whitfield, G.P., L.A. Meehan, N. Maizlish, and A.M. Wendel, 2017: The integrated transport and health impact modeling tool in Nashville, Tennessee, USA: Implementation steps and lessons learned. *Journal of Transport & Health*, **5**, 172-181. <http://dx.doi.org/10.1016/j.jth.2016.06.009>
254. Woodcock, J., M. Givoni, and A.S. Morgan, 2013: Health impact modelling of active travel visions for England and Wales using an integrated transport and health impact modelling tool (ITHIM). *PLOS ONE*, **8** (1), e51462. <http://dx.doi.org/10.1371/journal.pone.0051462>
255. Grabow, M.L., S.N. Spak, T. Holloway, B. Stone, Jr., A.C. Mednick, and J.A. Patz, 2012: Air quality and exercise-related health benefits from reduced car travel in the midwestern United States. *Environmental Health Perspectives*, **120** (1), 68-76. <http://dx.doi.org/10.1289/ehp.1103440>
256. Kenward, A., N. Zenes, J. Bronzan, J. Brady, and K. Shah, 2016: Overflow: Climate Change, Heavy Rain, and Sewage. Climate Central, Princeton, NJ, 12 pp. http://assets.climatecentral.org/pdfs/Overflow_sewagereport_update.pdf
257. EPA, 2016: Climate Change Indicators in the United States, 2016. 4th edition. EPA 430-R-16-004. U.S. Environmental Protection Agency, Washington, DC, 96 pp. https://www.epa.gov/sites/production/files/2016-08/documents/climate_indicators_2016.pdf

258. Norton, M.D. and G.T. Moore, 2017: St. Louis MSD CSO Volume Reduction Green Infrastructure Program. *Proceedings of the Water Environment Federation*, **2017** (2), 61-81. <http://dx.doi.org/10.2175/193864717821494853>
259. Sekaluvu, L., L. Zhang, and M. Gitau, 2018: Evaluation of constraints to water quality improvements in the Western Lake Erie Basin. *Journal of Environmental Management*, **205**, 85-98. <http://dx.doi.org/10.1016/j.jenvman.2017.09.063>
260. Tavakol-Davani, H., S.J. Burian, J. Devkota, and D. Apul, 2016: Performance and cost-based comparison of green and gray infrastructure to control combined sewer overflows. *Journal of Sustainable Water in the Built Environment*, **2** (2), 04015009. <http://dx.doi.org/10.1061/JSWBAY.0000805>
261. Tavakol-Davani, H., E. Goharian, C.H. Hansen, H. Tavakol-Davani, D. Apul, and S.J. Burian, 2016: How does climate change affect combined sewer overflow in a system benefiting from rainwater harvesting systems? *Sustainable Cities and Society*, **27**, 430-438. <http://dx.doi.org/10.1016/j.scs.2016.07.003>
262. Leard, B. and K. Roth, 2017: Voluntary Exposure Benefits and the Costs of Climate Change. RFF DP 15-19-REV2. Resources for the Future Washington DC, 57 pp. <http://www.rff.org/files/document/file/RFF-WP-15-19-REV2.pdf>
263. Dey, K.C., A. Mishra, and M. Chowdhury, 2015: Potential of intelligent transportation systems in mitigating adverse weather impacts on road mobility: A review. *IEEE Transactions on Intelligent Transportation Systems*, **16** (3), 1107-1119. <http://dx.doi.org/10.1109/TITS.2014.2371455>
264. Missouri Department of Transportation, 2017: Traveler Information Report [web site], accessed 7:53 AM; May 4, 2017. http://traveler.modot.org/report/modottext.aspx?type=all#tag_flood_closed
265. JOC 2013: High Water Forces Upper Mississippi River Closure. *Journal of Commerce*, **04 Jun.**
266. JOC, 2013: North American rail traffic slips. *Journal of Commerce*, **25 Apr.**
267. Workboat Staff, 2017: Portion of Upper Mississippi River Closed Near St. Louis. <https://www.workboat.com/news/coastal-inland-waterways/portion-upper-mississippi-river-closed-near-st-louis/>
268. Moore, K., 2016: High River Water Creates Navigation Turmoil. *WorkBoat.com*. <https://www.workboat.com/archive/high-river-water-creates-navigation-turmoil/>
269. Workboat Staff, 2015: Flooding Delays Barge Traffic. *WorkBoat.com*. <https://www.workboat.com/news/coastal-inland-waterways/flooding-delays-barge-traffic/>
270. DuPont, D.K., 2013: High Water Closes River Near St. Louis. *WorkBoat.com*. <https://www.workboat.com/archive/high-water-closes-river-near-st-louis/>
271. NOAA NCEI, 2018: Billion-Dollar Weather and Climate Disasters [web page]. NOAA National Centers for Environmental Information, Asheville, NC. <https://www.ncdc.noaa.gov/billions/events/US/1980-2017>
272. Nowak, D.J., E.J. Greenfield, R.E. Hoehn, and E. Lapoint, 2013: Carbon storage and sequestration by trees in urban and community areas of the United States. *Environmental Pollution*, **178**, 229-236. <http://dx.doi.org/10.1016/j.envpol.2013.03.019>
273. Nowak, D.J., R.E. Hoehn III, A.R. Bodine, D.E. Crane, J.F. Dwyer, V. Bonnewell, and G. Watson, 2013: Urban Trees and Forests of the Chicago Region. Resource Bulletin NRS-84. USDA, Forest Service, Northern Research Station, Newtown Square, PA, 106 pp. <http://dx.doi.org/10.2737/NRS-RB-84>
274. Getter, K.L., D.B. Rowe, G.P. Robertson, B.M. Cregg, and J.A. Andresen, 2009: Carbon sequestration potential of extensive green roofs. *Environmental Science & Technology*, **43** (19), 7564-7570. <http://dx.doi.org/10.1021/es901539x>
275. City of Chicago, 2012: City Unveils "Greenest Street in America" in Pilsen Neighborhood. City of Chicago Department of Transportation, Chicago, IL. https://www.cityofchicago.org/city/en/depts/cdot/provdrs/conservation_outreachgreenprograms/news/2012/oct/cdot_opens_the_pilsensustainablestreet.html
276. Metropolitan Sewer District, 2017: Rainscaping. MSD Project Clear, St. Louis, MO. <http://www.projectclearstl.org/get-the-rain-out/rainscaping/>
277. City of Minneapolis, 2009: City of Minneapolis Tree Cell Installation—Marq2 Project. Public Works, Minneapolis, MN. http://www.ci.minneapolis.mn.us/publicworks/stormwater/green/stormwater_green-initiatives_marq2-tree-install

278. Cleveland, 2015: The Cleveland Tree Plan. Cleveland Forest Coalition, Cleveland, OH, 57 pp. http://www.city.cleveland.oh.us/sites/default/files/forms_publications/ClevelandTreePlan.pdf
279. Ducks Unlimited, 2016: Missouri State Conservation Report. Ducks Unlimited Great Lakes/Atlantic Region, Ann Arbor, MI, 2 pp. <http://www.ducks.org/missouri/missouri-conservation-projects>
280. Ozaukee Washington Land Trust, 2016: Open Spaces: 2016 Annual Report. Ozaukee Washington Land Trust, West Bend, WI, 8 pp.
281. GRAEF, Hey and Associates Inc., and CDM Smith, 2017: Kinnickinnic River Watershed Flood Management Plan: Final Report. Executive Summary. Milwaukee Metropolitan Sewerage District, Milwaukee, WI, 9 pp. https://www.mmsd.com/application/files/4314/9522/1491/KK_Watershed_Flood_Management_Plan_05_04_17_-_EXECUTIVE_SUMMARY_002.pdf
282. Trice, A., 2016: Daylighting Streams: Breathing Life into Urban Streams and Communities. American Rivers, Washington, DC, 32 pp. http://americanrivers.org/wp-content/uploads/2016/05/AmericanRivers_daylighting-streams-report.pdf
283. Norton-Smith, K., K. Lynn, K. Chief, K. Cozzetto, J. Donatuto, M.H. Redsteer, L.E. Kruger, J. Maldonado, C. Viles, and K.P. Whyte, 2016: Climate Change and Indigenous Peoples: A Synthesis of Current Impacts and Experiences. Gen. Tech. Rep. PNW-GTR-944. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station, Portland, OR, 136 pp. <https://www.fs.usda.gov/treearch/pubs/53156>
284. Whyte, K.P., 2014: A concern about shifting interactions between indigenous and non-indigenous parties in US climate adaptation contexts. *Interdisciplinary Environmental Review*, **15** (2/3), 114-133. <http://dx.doi.org/10.1504/IER.2014.063658>
285. Kalafatis, S.E. and M.C. Lemos, 2017: The emergence of climate change policy entrepreneurs in urban regions. *Regional Environmental Change*, **17** (6), 1791-1799. <http://dx.doi.org/10.1007/s10113-017-1154-0>
286. Pryor, S.C., Ed. 2013: *Climate Change in the Midwest: Impacts, Risks, Vulnerability and Adaptation*. Indiana University Press, Bloomington, IN, 288 pp.
287. Larsen, L., 2015: Urban climate and adaptation strategies. *Frontiers in Ecology and the Environment*, **13** (9), 486-492. <http://dx.doi.org/10.1890/150103>
288. Woodruff, S.C. and M. Stults, 2016: Numerous strategies but limited implementation guidance in US local adaptation plans. *Nature Climate Change*, **6** (8), 796-802. <http://dx.doi.org/10.1038/nclimate3012>
289. Phadke, R., C. Manning, and S. Burlager, 2015: Making it personal: Diversity and deliberation in climate adaptation planning. *Climate Risk Management*, **9**, 62-76. <http://dx.doi.org/10.1016/j.crm.2015.06.005>
290. Barclay, P., C. Bastoni, D. Eisenhauer, M. Hassan, M. Lopez, L. Mekias, S. Ramachandran, and R. Stock. 2013: Climate Change Adaptation in Great Lakes Cities. M.Sc. project, Natural Resources and Environment, University of Michigan, 99 pp. <http://hdl.handle.net/2027.42/97435>
291. Rasmussen, L.V., C.J. Kirchoff, and M.C. Lemos, 2017: Adaptation by stealth: Climate information use in the Great Lakes region across scales. *Climatic Change*, **140** (3), 451-465. <http://dx.doi.org/10.1007/s10584-016-1857-0>
292. Kalafatis, S.E., A. Grace, and E. Gibbons, 2015: Making climate science accessible in Toledo: The linked boundary chain approach. *Climate Risk Management*, **9**, 30-40. <http://dx.doi.org/10.1016/j.crm.2015.04.003>
293. Vogel, J., K.M. Carney, J.B. Smith, C. Herrick, M. Stults, M. O'Grady, A.S. Juliana, H. Hosterman, and L. Giangola, 2016: Climate Adaptation—The State of Practice in U.S. Communities. Kresge Foundation, Detroit. <http://kresge.org/sites/default/files/library/climate-adaptation-the-state-of-practice-in-us-communities-full-report.pdf>
294. Lemos, M.C., C.J. Kirchoff, S.E. Kalafatis, D. Scavia, and R.B. Rood, 2014: Moving climate information off the shelf: Boundary chains and the role of RISAs as adaptive organizations. *Weather, Climate, and Society*, **6** (2), 273-285. <http://dx.doi.org/10.1175/WCAS-D-13-00044.1>
295. HRWC, 2018: Assessing Urban Vulnerability [web site]. Huron River Watershed Council (HRWC), Ann Arbor, MI. <https://www.hrwc.org/what-we-do/programs/climate-change/assessing-urban-vulnerability/>
296. USDN, 2018: Urban Sustainability Directors Network [web site]. <https://www.usdn.org/>

297. Albouy, D., W. Graf, R. Kellogg, and H. Wolff, 2016: Climate amenities, climate change, and American quality of life. *Journal of the Association of Environmental and Resource Economists*, **3** (1), 205-246. <http://dx.doi.org/10.1086/684573>
298. Sinha, P. and M.L. Cropper, 2013: The Value of Climate Amenities: Evidence from US Migration Decisions. NBER Working Paper No. 18756. National Bureau of Economic Research, Cambridge, MA, 49 pp. <http://dx.doi.org/10.3386/w18756>
299. Brugger, J., A. Meadow, and A. Horangic, 2016: Lessons from first-generation climate science integrators. *Bulletin of the American Meteorological Society*, **97** (3), 355-365. <http://dx.doi.org/10.1175/bams-d-14-00289.1>
300. Prokopy, L.S., J.S. Carlton, T. Haigh, M.C. Lemos, A.S. Mase, and M. Widhalm, 2017: Useful to usable: Developing usable climate science for agriculture. *Climate Risk Management*, **15**, 1-7. <http://dx.doi.org/10.1016/j.crm.2016.10.004>
301. Arbuckle, J.G., J. Hobbs, A. Loy, L.W. Morton, L.S. Prokopy, and J. Tyndall, 2014: Understanding Corn Belt farmer perspectives on climate change to inform engagement strategies for adaptation and mitigation. *Journal of Soil and Water Conservation*, **69** (6), 505-516. <http://dx.doi.org/10.2489/jswc.69.6.505>
302. Carlton, J.S., A.S. Mase, C.L. Knutson, M.C. Lemos, T. Haigh, D.P. Todey, and L.S. Prokopy, 2016: The effects of extreme drought on climate change beliefs, risk perceptions, and adaptation attitudes. *Climatic Change*, **135** (2), 211-226. <http://dx.doi.org/10.1007/s10584-015-1561-5>
303. Haigh, T., E. Takle, J. Andresen, M. Widhalm, J.S. Carlton, and J. Angel, 2015: Mapping the decision points and climate information use of agricultural producers across the U.S. corn belt. *Climate Risk Management*, **7**, 20-30. <http://dx.doi.org/10.1016/j.crm.2015.01.004>
304. Heltberg, R., P.B. Siegel, and S.L. Jorgensen, 2009: Addressing human vulnerability to climate change: Toward a “no-regrets” approach. *Global Environmental Change*, **19** (1), 89-99. <http://dx.doi.org/10.1016/j.gloenvcha.2008.11.003>
305. Henstra, D., 2012: Toward the climate-resilient city: Extreme weather and urban climate adaptation policies in two Canadian provinces. *Journal of Comparative Policy Analysis: Research and Practice*, **14** (2), 175-194. <http://dx.doi.org/10.1080/13876988.2012.665215>
306. van Vuuren, D.P., K. Riahi, R. Moss, J. Edmonds, A. Thomson, N. Nakicenovic, T. Kram, F. Berkhout, R. Swart, A. Janetos, S.K. Rose, and N. Arnell, 2012: A proposal for a new scenario framework to support research and assessment in different climate research communities. *Global Environmental Change*, **22** (1), 21-35. <http://dx.doi.org/10.1016/j.gloenvcha.2011.08.002>
307. Environment and Climate Change Canada and the U.S. National Oceanic and Atmospheric Administration, 2018: 2017 Annual Climate Trends and Impacts Summary for the Great Lakes Basin. <https://binational.net/2018/07/10/ctis-ctic-2017/>
308. Kirchhoff, C.J., M.C. Lemos, and S. Dessai, 2013: Actionable knowledge for environmental decision making: Broadening the usability of climate science. *Annual Review of Environment and Resources*, **38** (1), 393-414. <http://dx.doi.org/10.1146/annurev-environ-022112-112828>
309. Lynn, K., J. Daigle, J. Hoffman, F. Lake, N. Michelle, D. Ranco, C. Viles, G. Voggeser, and P. Williams, 2013: The impacts of climate change on tribal traditional foods. *Climatic Change*, **120** (3), 545-556. <http://dx.doi.org/10.1007/s10584-013-0736-1>
310. Montag, J.M., K. Swan, K. Jenni, T. Nieman, J. Hatten, M. Mesa, D. Graves, F. Voss, M. Mastin, J. Hardiman, and A. Maule, 2014: Climate change and Yakama Nation tribal well-being. *Climatic Change*, **124** (1), 385-398. <http://dx.doi.org/10.1007/s10584-013-1001-3>
311. Brubaker, M., J. Bell, J. Berner, M. Black, R. Chaven, J. Smith, and J. Warren, 2011: Climate Change in Noatak, Alaska: Strategies for Community Health. Alaska Native Tribal Health Consortium, Anchorage, AK, 54 pp. https://anthc.org/wp-content/uploads/2016/01/CCH_AR_062011_Climate-Change-in-Noatak.pdf
312. Peterson, K. and J.K. Maldonado, 2016: When adaptation is not enough: “Between now and then” of community-led resettlement. *Anthropology and Climate Change: From Encounters to Actions*, 2nd ed. Crate, S.A. and M. Nuttall, Eds. Taylor & Francis, New York, NY, 336-353.

313. Emery, M.R., A. Wrobel, M.H. Hansen, M. Dockry, W.K. Moser, K.J. Stark, and J.H. Gilbert, 2014: Using traditional ecological knowledge as a basis for targeted forest inventories: Paper birch (*Betula papyrifera*) in the US Great Lakes region. *Journal of Forestry*, **112** (2), 207-214. <http://dx.doi.org/10.5849/jof.13-023>
314. Werkheiser, I., 2016: Community epistemic capacity. *Social Epistemology*, **30** (1), 25-44. <http://dx.doi.org/10.1080/02691728.2014.971911>
315. Vavrus, S.J., M. Notaro, and D.J. Lorenz, 2015: Interpreting climate model projections of extreme weather events. *Weather and Climate Extremes*, **10** (Part B), 10-28. <http://dx.doi.org/10.1016/j.wace.2015.10.005>
316. Harding, K.J., P.K. Snyder, and S. Liess, 2013: Use of dynamical downscaling to improve the simulation of Central U.S. warm season precipitation in CMIP5 models. *Journal of Geophysical Research Atmospheres*, **118** (22), 12,522-12,536. <http://dx.doi.org/10.1002/2013JD019994>
317. Bryan, A.M., A.L. Steiner, and D.J. Posselt, 2015: Regional modeling of surface-atmosphere interactions and their impact on Great Lakes hydroclimate. *Journal of Geophysical Research Atmospheres*, **120** (3), 1044-1064. <http://dx.doi.org/10.1002/2014JD022316>
318. Briley, L.J., W.S. Ashley, R.B. Rood, and A. Krmenc, 2017: The role of meteorological processes in the description of uncertainty for climate change decision-making. *Theoretical and Applied Climatology*, **127** (3), 643-654. <http://dx.doi.org/10.1007/s00704-015-1652-2>
319. Mallard, M.S., C.G. Nolte, T.L. Spero, O.R. Bullock, K. Alapaty, J.A. Herwehe, J. Gula, and J.H. Bowden, 2015: Technical challenges and solutions in representing lakes when using WRF in downscaling applications. *Geoscientific Model Development*, **8** (4), 1085-1096. <http://dx.doi.org/10.5194/gmd-8-1085-2015>
320. ISU, 2017: Iowa Environmental Mesonet (IEM). Iowa State University (ISU), Ames, IA. <https://mesonet.agron.iastate.edu/>
321. Garbrecht, J.D., J.L. Steiner, and C.A. Cox, 2007: Climate change impacts on soil and water conservation. *Eos, Transactions, American Geophysical Union*, **88** (11), 136-136. <http://dx.doi.org/10.1029/2007EO110016>
322. Favis-Mortlock, D., 2017: *The Soil Erosion Site: Soil Erosion by Water*, Oxford, UK. http://soilerosion.net/water_erosion.html
323. Cloud, H.A. and R.V. Morey, 2017: Management of Stored Grain with Aeration [web site]. University of Minnesota Extension, St. Paul, MN. <https://extension.umn.edu/corn-harvest/managing-stored-grain-aeration>
324. Miraglia, M., H.J.P. Marvin, G.A. Kleter, P. Battilani, C. Brera, E. Coni, F. Cubadda, L. Croci, B. De Santis, S. Dekkers, L. Filippi, R.W.A. Hutjes, M.Y. Noordam, M. Pisante, G. Piva, A. Prandini, L. Toti, G.J. van den Born, and A. Vespermann, 2009: Climate change and food safety: An emerging issue with special focus on Europe. *Food and Chemical Toxicology*, **47** (5), 1009-1021. <http://dx.doi.org/10.1016/j.fct.2009.02.005>
325. De Lucia, M. and D. Assennato, 1994: Agricultural engineering in development: Post-harvest operations and management of foodgrains. FAO Agricultural Services Bulletin No. 93. Food and Agriculture Organization of the United Nations, Rome, Italy. <http://www.fao.org/docrep/t0522e/t0522e00.htm>
326. Sawant, A.A., S.C. Patil, S.B. Kalse, and N.J. Thakor, 2012: Effect of temperature, relative humidity and moisture content on germination percentage of wheat stored in different storage structures. *Agricultural Engineering International: CIGR Journal*, **14** (2), 110-118. <http://www.cigrjournal.org/index.php/Ejournal/article/view/2019>
327. Atungulu, G.R., 2017: Management of in-bin grain drying and storage systems for improved grain quality and prevention of mycotoxins. USDA Research, Education & Economics Information System. <https://portal.nifa.usda.gov/web/crisprojectpages/1002599-management-of-in-bin-grain-drying-and-storage-systems-for-improved-grain-quality-and-prevention-of-mycotoxins.html>

328. Walthall, C., P. Backlund, J. Hatfield, L. Lengnick, E. Marshall, M. Walsh, S. Adkins, M. Aillery, E.A. Ainsworth, C. Amman, C.J. Anderson, I. Bartomeus, L.H. Baumgard, F. Booker, B. Bradley, D.M. Blumenthal, J. Bunce, K. Burkey, S.M. Dabney, J.A. Delgado, J. Dukes, A. Funk, K. Garrett, M. Glenn, D.A. Grantz, D. Goodrich, S. Hu, R.C. Izaurralde, R.A.C. Jones, S.-H. Kim, A.D.B. Leaky, K. Lewers, T.L. Mader, A. McClung, J. Morgan, D.J. Muth, M. Nearing, D.M. Oosterhuis, D. Ort, C. Parmesan, W.T. Pettigrew, W. Polley, R. Rader, C. Rice, M. Rivington, E. Rosskopf, W.A. Salas, L.E. Sollenberger, R. Srygley, C. Stöckle, E.S. Takle, D. Timlin, J.W. White, R. Winfree, L. Wright-Morton, and L.H. Ziska, 2012: Climate Change and Agriculture in the United States: Effects and Adaptation. USDA Technical Bulletin 1935. U.S. Department of Agriculture, Washington, DC, 186 pp. [http://www.usda.gov/oce/climate_change/effects_2012/CC%20and%20Agriculture%20Report%20\(02-04-2013\)b.pdf](http://www.usda.gov/oce/climate_change/effects_2012/CC%20and%20Agriculture%20Report%20(02-04-2013)b.pdf)
329. St-Pierre, N.R., B. Cobanov, and G. Schnitkey, 2003: Economic losses from heat stress by US livestock industries. *Journal of Dairy Science*, **86**, E52-E77. [http://dx.doi.org/10.3168/jds.S0022-0302\(03\)74040-5](http://dx.doi.org/10.3168/jds.S0022-0302(03)74040-5)
330. Lewis, C.R.G. and K.L. Bunter, 2010: Heat stress: The effects of temperature on production and reproduction traits. In *AGBU Pig Genetics Workshop*, October 2010, 87-96. http://agbu.une.edu.au/pig_genetics/pdf/2010/P12-Craig-Heat%20stress.pdf
331. Bussey, J., M.A. Davenport, M.R. Emery, and C. Carroll, 2016: "A lot of it comes from the heart": The nature and integration of ecological knowledge in tribal and nontribal forest management. *Journal of Forestry*, **114** (2), 97-107. <http://dx.doi.org/10.5849/jof.14-130>
332. Cline, T.J., V. Bennington, and J.F. Kitchell, 2013: Climate change expands the spatial extent and duration of preferred thermal habitat for Lake Superior fishes. *PLOS ONE*, **8** (4), e62279. <http://dx.doi.org/10.1371/journal.pone.0062279>
333. Basu, R., 2009: High ambient temperature and mortality: A review of epidemiologic studies from 2001 to 2008. *Environmental Health*, **8**, 40. <http://dx.doi.org/10.1186/1476-069X-8-40>
334. Kovats, R.S. and S. Hajat, 2008: Heat stress and public health: A critical review. *Annual Review of Public Health*, **29**, 41-55. <http://dx.doi.org/10.1146/annurev.publhealth.29.020907.090843>
335. Brownstein, J.S., T.R. Holford, and D. Fish, 2003: A climate-based model predicts the spatial distribution of the Lyme disease vector *Ixodes scapularis* in the United States. *Environmental Health Perspectives*, **111** (9), 1152-1157. <https://www.jstor.org/stable/3435502>
336. Hamer, S.A., G.J. Hickling, E.D. Walker, and J.I. Tsao, 2014: Increased diversity of zoonotic pathogens and *Borrelia burgdorferi* strains in established versus incipient *Ixodes scapularis* populations across the Midwestern United States. *Infection, Genetics and Evolution*, **27**, 531-542. <http://dx.doi.org/10.1016/j.meegid.2014.06.003>
337. Ogden, N.H., L.R. Lindsay, and P.A. Leighton, 2013: Predicting the rate of invasion of the agent of Lyme disease *Borrelia burgdorferi*. *Journal of Applied Ecology*, **50** (2), 510-518. <http://dx.doi.org/10.1111/1365-2664.12050>
338. Posey, J., 2016: St. Louis in the Anthropocene: Responding to Global Environmental Change. *St. Louis Currents: The Fifth Edition*. Theising, A. and E.T. Jones, Eds. Reedy Press, St. Louis, MO.
339. Missouri Department of Transportation, 2017: Traveler Information Report [web site], accessed May 24, 2017. http://traveler.modot.org/report/modottext.aspx?type=all#tag_flood_closed
340. Smith, A., E. Chuck, and A. Gostanian, 2015: Swollen Midwest Rivers Bring Transportation to Standstill. NBC News, New York. <https://www.nbcnews.com/news/weather/missouri-illinois-face-slow-motion-disaster-swollen-rivers-rise-n488376>
341. Workboat Staff, various: Aggregation of articles documenting Mississippi River flood-related closures. WorkBoat.com. <https://www.workboat.com/?s=mississippi+river+closed+flood>
342. Associated Press, 2017: "Amtrak suspends rail service across Missouri." May 2. <http://fox2now.com/2017/05/02/amtrak-suspends-rail-service-across-missouri/>
343. Changnon, S., 2009: Impacts of the 2008 floods on railroads in Illinois and adjacent states. *Transactions of the Illinois State Academy of Science*, **102** (3-4), 181-190. <http://ilacadofsci.com/wp-content/uploads/2013/03/102-17MS2819-print.pdf>
344. Criss, R.E. and W.E. Winston, 2008: Public safety and faulty flood statistics. *Environmental Health Perspectives*, **116** (12), A516-A516. <http://dx.doi.org/10.1289/ehp.12042>

345. Criss, R.E. and M. Luo, 2017: Increasing risk and uncertainty of flooding in the Mississippi River basin. *Hydrological Processes*, **31** (6), 1283-1292. <http://dx.doi.org/10.1002/hyp.11097>
346. Criss, R.E., 2016: Statistics of evolving populations and their relevance to flood risk. *Journal of Earth Science*, **27** (1), 2-8. <http://dx.doi.org/10.1007/s12583-015-0641-9>
347. Asam, S., D. Spindler, S. Julius, and B. Beierwagen, 2016: Stormwater Management in Response to Climate Change Impacts: Lessons from the Chesapeake Bay and Great Lakes Regions. EPA/600/R-15/087F. U.S. Environmental Protection Agency, Washington, DC. <https://cfpub.epa.gov/ncea/global/recordisplay.cfm?deid=310045>
348. Chicago Metropolitan Agency for Planning (CMAP), 2013: Climate Adaptation Guidebook for Municipalities in the Chicago Region. CMAP, Chicago, IL. <http://www.cmap.illinois.gov/documents/10180/14136/FY13-0119%20Climate%20Adaptation%20toolkit.pdf/fa5e3867-8278-4867-841a-aad4e090847a>
349. Delta Institute, 2015: Green Infrastructure Designs: Scalable Solutions to Local Challenges. Delta Institute, Chicago, IL, 70 pp. <http://delta-institute.org/delta/wp-content/uploads/Green-Infrastructure-Designs-July-2015.pdf>
350. Lichten, N., J.I. Nassauer, M. Dewar, N.R. Sampson, and N.J. Webster, 2016: Green Infrastructure on Vacant Land: Achieving Social and Environmental Benefits in Legacy Cities. NEW-GI White Paper No. 1. University of Michigan Water Center, Ann Arbor, MI. <https://static1.squarespace.com/static/52a213fce4b0a5794c59856f/t/58d42d0f725e25f7c64240e3/1490300177284/Green+Infrastructure+on+Vacant+Land.pdf>
351. Tomer, M.D. and K.E. Schilling, 2009: A simple approach to distinguish land-use and climate-change effects on watershed hydrology. *Journal of Hydrology*, **376** (1), 24-33. <http://dx.doi.org/10.1016/j.jhydrol.2009.07.029>
352. Frans, C., E. Istanbuluoglu, V. Mishra, F. Munoz-Arriola, and D.P. Lettenmaier, 2013: Are climatic or land cover changes the dominant cause of runoff trends in the Upper Mississippi River Basin? *Geophysical Research Letters*, **40** (6), 1104-1110. <http://dx.doi.org/10.1002/grl.50262>
353. Slater, L.J., M.B. Singer, and J.W. Kirchner, 2015: Hydrologic versus geomorphic drivers of trends in flood hazard. *Geophysical Research Letters*, **42** (2), 370-376. <http://dx.doi.org/10.1002/2014GL062482>
354. Ryberg, K.R., W. Lin, and A.V. Vecchia, 2014: Impact of climate variability on runoff in the north-central United States. *Journal of Hydrologic Engineering*, **19** (1), 148-158. [http://dx.doi.org/10.1061/\(ASCE\)HE.1943-5584.0000775](http://dx.doi.org/10.1061/(ASCE)HE.1943-5584.0000775)
355. Mallakpour, I. and G. Villarini, 2015: The changing nature of flooding across the central United States. *Nature Climate Change*, **5** (3), 250-254. <http://dx.doi.org/10.1038/nclimate2516>