



Modes of Antarctic tidal grounding line migration revealed by Ice, Cloud, and land Elevation Satellite-2 (ICESat-2) laser altimetry

Bryony I. D. Freer^{1,2}, Oliver J. Marsh¹, Anna E. Hogg², Helen Amanda Fricker³, and Laurie Padman⁴

¹British Antarctic Survey, Cambridge, CB3 0ET, UK

²School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK

³Scripps Polar Center, Scripps Institution of Oceanography, UC San Diego, La Jolla, California 92093-0225, USA

⁴Earth and Space Research, Corvallis, Oregon 97333-1536, USA

Correspondence: Bryony I. D. Freer (breer90@bas.ac.uk)

Received: 23 December 2022 – Discussion started: 24 February 2023

Revised: 15 June 2023 – Accepted: 9 July 2023 – Published: 26 September 2023

Abstract. Tide-forced short-term migration of the grounding line (GL) of Antarctic ice shelves can impact ice dynamics at the ice sheet margins and obscures assessments of long-term GL advance or retreat. However, the magnitude of tidally induced GL migration is poorly known, and the spatial patterns and modes of variability are not well characterised. Here we develop and apply a technique that uses Ice, Cloud, and land Elevation Satellite-2 (ICESat-2) repeat-track laser altimetry to locate the inland limit of tidal ice shelf flexure for each sampled tide, enabling the magnitude and temporal variability of tidal GL migration to be resolved. We demonstrate its application at an ice plain north of Bungenstockrücken, in a region of the southern Ronne Ice Shelf subject to large ocean tides. We observe a 1300 km² area of ephemeral grounding over which the GL migrates by up to 15 km between low and high tide and identify four distinct modes of migration: *linear*, *asymmetric*, *threshold* and *hysteresis*. The short-term movement of the GL dominates any long-term migration signal in this location, and the distribution of GL positions and modes contains information about spatial variability in the ice–bed interface. We discuss the impact of extreme tidal GL migration on ice shelf–ocean–subglacial systems in Antarctica and make recommendations for how GLs should be more precisely defined and documented in future by the community.

1 Introduction

Recent mass loss from the Antarctic Ice Sheet has been attributed to changes in the floating ice shelves that fringe 74 % of its margins (Gudmundsson et al., 2019; Joughin et al., 2012) and buttress upstream grounded ice (Dupont and Alley, 2005). Accurate representation of the ice sheet in coupled Earth system models is essential to predict its evolution and the magnitude and rate of its future contribution to sea level rise. This requires knowledge of ice sheet processes and boundary conditions, including the configuration and behaviour of ice shelves. One important parameter for ice sheet monitoring and modelling is the location of the grounding line (GL), which marks the boundary between the grounded ice sheet and floating ice shelf (Thomas, 1979). As the ice detaches from the underlying bed, ice flow across the wider grounding zone (GZ) transitions from a regime dominated by vertical shear and basal drag to that of primarily buoyancy-driven flow dominated by longitudinal stretching and lateral shear with no basal drag (Schoof, 2007). There can be high rates of basal melt at the GL as ocean water comes into contact with the base of the ice shelf (Jenkins et al., 2006; Depoorter et al., 2013). The location of the GL and change in its position over time is, therefore, a sensitive indicator of ice sheet stability and local environmental forcing, with sustained ice thinning causing GL retreat and thickening causing GL advance (Joughin et al., 2010, 2012; Dutrieux et al., 2014), and it has been identified as an essential climate variable (Bojinski et al., 2014)

Several satellite-based techniques have been used to map features in the GZ, with the primary goal of monitoring long-term change in GL position (Sect. 2). The GL is typically located by identifying the inland limit of ice shelf flexure, but this is complicated by the effect of short-term sea level variations, primarily driven by ocean tides, which can cause GL migration on hourly to daily timescales. During a rising tide, increased buoyancy causes more of the ice shelf to lift off the bed and the GL temporarily migrates inland, returning to its most seaward position at low tide (Brancato et al., 2020), with some slight time lag due to the viscoelastic effects of the ice (Reeh et al., 2003). Prior satellite observations have shown that the extent of this short-term tidal GL migration can range from a few hundred metres to several kilometres (Hogg et al., 2016; Brunt et al., 2011; Milillo et al., 2017, 2019, 2022), with the magnitude of this migration controlled by the tide amplitude, local bed topography, and thickness and strength of the ice. The signal of short-term tidal GL migration is superimposed on the long-term signal of GL migration associated with ice dynamic change, which is traditionally calculated by comparing sparsely sampled, individual measurements of GL position through time (e.g. Rignot et al., 2014). Therefore, in order to increase confidence in measurements of long-term GL migration rates, we must improve our knowledge of the spatially variable pattern of short-term tidal GL migration.

Despite the importance of understanding tidal GL migration, there is currently no pan-Antarctic assessment of its extent or variability. This is mostly because there is a lack of acquired or suitable satellite data in the historical archive with sufficient spatial coverage and temporal resolution. We currently also have a limited understanding of the *mechanisms* of tidal ice shelf flexure and GL migration. Various representations of tidal flexure have been modelled using elastic and viscoelastic frameworks (Holdsworth, 1969; Vaughan, 1995; Schmeltz et al., 2002; Walker et al., 2013), some of which allow for tidal GL migration (Sayag and Worster, 2013; Tsai and Gudmundsson, 2015; Stubblefield et al., 2021), but results vary considerably, and the choice of representation of these processes can have a large impact on results of modelling studies (e.g. Mosbeux et al., 2023). Whilst some localised in situ measurements of tidal flexure have been made to examine these processes (Smith, 1991; Vaughan, 1995), there are very few satellite observations with sufficient sampling of different tidal states to test and validate these models. Furthermore, modelled projections of future mass loss from West Antarctica are very sensitive to the amount of basal melt and ice flux at the GL (Arthern and Williams, 2017; Goldberg et al., 2019), yet most numerical ice sheet models typically assume a fixed or slowly moving GL determined by flotation conditions alone with zero basal melt (Milillo et al., 2017; Favier et al., 2014). This is an oversimplification in areas subject to extreme tidal variability, where short-term GL migration is likely to impact both ice dynamics through rapid variations in basal shear stress and basal melt rate through

changes in cavity geometry, enhancing tidal mixing (Chen et al., 2023; Ciraci et al., 2023; Milillo et al., 2019). Quantifying the extent and influence of tidal GL migration around the margin of Antarctica is valuable for these processes to be more accurately parameterised in ice sheet models.

In this study we present a new approach to observe tidal GL migration that takes advantage of the high along-track resolution and repeat-track configuration of NASA's Ice, Cloud, and land Elevation Satellite-2 (ICESat-2) laser altimeter mission to improve temporal sampling of tidal ice flexure and GL migration. We apply the method to the ice plain north of the Bungenstockrücken ice rise, on the southern Ronne Ice Shelf. An ice plain is defined as an area of low surface slope close to the GZ where the ice is close to flotation (Alley et al., 1989) and which can sometimes experience "ephemeral grounding" between high and low tide (Schmeltz et al., 2001; Brunt et al., 2011). We show that with ICESat-2 repeat-track laser altimetry (RTLA) it is possible to monitor the time-varying GZ structure at the Bungenstockrücken ice plain in unprecedented detail, and we identify four distinct modes of tidal GL migration. We discuss the implications both for long-term GL monitoring and how these findings inform our process understanding of tidal flexure and GL migration mechanisms in Antarctica.

2 Grounding lines and their detection

2.1 Features of the grounding zone

The GZ is the region over which the ice transitions from being in constant contact with the bed to floating in hydrostatic equilibrium with the ocean (see Fig. 1). Ice in the GZ is supported by both the hydrostatic pressure from the underlying ocean and internal stresses, and it undergoes tidal flexure (Vaughan, 1995). The true GL location, Point *G*, refers to the point at which the ice base first detaches from the bed as it flows from the ice sheet. As a subglacial feature within the GZ, Point *G* can only be directly observed using ground-based radar (MacGregor et al., 2011; Catania et al., 2010) or using remotely operated vehicles deployed from the shelf edge or through boreholes (Schmidt et al., 2023). However, there are a number of observable surface features in the GZ that relate to the location of Point *G*, which can be more readily detected in situ or in satellite data and used as GL proxies (Fig. 1). Point *F* is the landward limit of tidal flexure and is often located slightly inland of Point *G* (on the order of hundreds of metres) due to the elastic properties of the ice (Padman et al., 2018; Rignot et al., 2011). Point *H* is the seaward limit of tidal flexure and inland limit of hydrostatic equilibrium. The break in slope, Point *I_b*, marks the sharp reduction in surface slope that occurs due to the abrupt change in basal shear stress across the GL, and in typical GZs it is located slightly seaward of Point *G* (Schoof, 2011). However, where a GZ has an ice plain, Point *I_b* has been observed several




Figure 1. (a) Idealised cross section of an ice plain GZ, showing the location of typical satellite-derived GZ proxies: Point I_b , the first break in slope; Point F , the inland limit of tidal flexure (represented here as approximately equal to Point G , the true GL); and Point H , the inshore limit of hydrostatic equilibrium. (b) Idealised cross section of an ice plain GZ experiencing tidal GL migration, showing the difference in position of the ice shelf and migration of Point F between high tide (F_{\max}) and low tide (F_{\min}), across 24 h of a simplified ~ 12 h semidiurnal tide cycle.

kilometres inland of Point G (Corr et al., 2001). We show the locations of the key GZ surface proxies as they relate to Point G for an idealised ice plain in Fig. 1a, and in Fig. 1b we depict how Point F can migrate by several kilometres across the tide cycle at an ice plain experiencing ephemeral grounding.

2.2 Methods for detecting the grounding zone

GZ surface proxies can be measured in situ using instruments such as tiltmeters and GNSS (global navigation satellite system) (Stephenson et al., 1979; Smith, 1991; Vaughan, 1995) or using satellite techniques to obtain more widespread measurements (e.g. Rignot, 1996; Friedl et al., 2020). Static satellite methods identify the break in slope, Point I_b , from either shadows in a single-epoch optical satellite image (Scambos et al., 2007; Bindenschadler et al., 2011) or single-epoch ele-

vation profiles from radar and laser satellite altimetry (Hogg et al., 2018; Brunt et al., 2010a; Li et al., 2022a). Dynamic satellite methods use data acquired at two or more distinct phases of the tidal cycle to locate the limits of tidal flexure, Points F and H . CryoSat-2 radar altimetry has been used to map Points F and H by applying a pseudo-crossover method to detect tidal flexure (Dawson and Bamber, 2017, 2020). This provides good spatial coverage by utilising large data volumes over a multi-year record but cannot be used to detect short-term GL migration between specific tides due to its long repeat cycle (369 d). Differential interferometric synthetic-aperture radar (DInSAR) is one of the most commonly used dynamic satellite methods and can produce quasi-continuous maps of Points F and H from double-differenced interferograms produced from three or four SAR images acquired at different times in the tidal cycle (Rignot, 1996). However, the temporal sampling and spatial coverage

of DInSAR in Antarctica is limited, in part due to its reliance on a short repeat time between acquisitions to retain interferometric coherence. Sentinel-1 DInSAR has been used to detect tidal GL migration (Mohajerani et al., 2021; Chen et al., 2023), but the coverage only extends to $\sim 77^\circ$ S, which excludes the majority of the Ross and Filchner–Ronne GLs. Recent studies using data from the COSMO-SkyMed (COntstellatIon of small SatellItes for the Mediterranean basin Observation, CSK) constellation, with a short repeat period of 1, 3, 4 or 8 d and up to 3 m spatial resolution, have improved DInSAR sampling of tidal GL migration in specific locations (Milillo et al., 2017; Minchew et al., 2017; Brancato et al., 2020) but are still complicated by the fact that interferograms are produced from oblique imagery from at least three different tide times. Moreover, while the relatively short repeat period of CSK enables interferometric coherence to be maintained on faster floating ice streams, the spatial coverage of coherent SAR image pairs is still limited, constraining the period where continent-wide studies can be performed.

An alternative dynamic satellite technique is repeat-track satellite laser altimetry (RTLA), which was first introduced for GZ detection by Fricker and Padman (2006). For this method, comparison of repeat tracks of ice shelf elevation profiles sampled at different tidal states identifies elevation anomalies that relate to Points F and H along discrete ground tracks. RTLA was pioneered using data from the Geoscience Laser Altimeter System instrument on board the Ice, Cloud, and Land Elevation Satellite (ICESat) that was in orbit from 2003 to 2009. However, ICESat only had a single ground track that only repeated to about ± 100 m, which led to unrecoverable topographic biases across GZs due to the typically large across-track slopes (Fricker et al., 2009). In contrast, the Advanced Topographic Laser Altimeter System (ATLAS) that launched on board ICESat-2 in 2018 has a six-beam design with more accurate pointing, which reduces across-track deviation from each reference pair track (RPT) to within 4.4 ± 6.0 m (Luthcke et al., 2021), providing better spatial sampling of the GZ. Only two repeat measurements are required to locate Point F using RTLA, providing an advantage over DInSAR which requires at least three repeats, allowing us to attribute the migration of Point F to individual tidal states. Li et al. (2022b) have used these data to locate a single Point F , H and I_b position (Fig. 1a) along most ICESat-2 ground tracks around Antarctica. This is a highly valuable and comprehensive dataset for the identification of long-term grounding line change, but as only a single Point F has been located per ICESat-2 ground track, it is limited in use for studying tidal GZ processes. Therefore, here we extend the Li et al. (2022b) record at Bungenstockrücken to locate *multiple* positions of Point F along each ICESat-2 ground track as it migrates over the tide cycle (Fig. 1b), providing novel observations of tidal GZ behaviour.

3 Data and methods

3.1 Datasets

The ATLAS instrument on board ICESat-2 is the first spaceborne photon-counting laser altimeter (Markus et al., 2017). It samples 11 m diameter footprints (Magruder et al., 2021), estimating surface heights every 0.7 m along its 1387 reference ground tracks (RGTs). ATLAS transmits a single beam of 532 nm (green) laser light pulsing at 10 kHz along each RGT, which is split into six beams organised into three pairs, each separated by 3.3 km. The beams in each pair, one “weak” and one “strong”, are separated by 90 m across-track and each beam follows its own ground track (GT1L, GT1R, GT2R, ... GT3L). The system operates in a 91 d exact-repeat orbit, providing repeat surface height measurements along each ground track about four times per year (Neumann et al., 2019). We obtained data from version 5 of the Land Ice Height (ATL06) product along all ground tracks crossing the Bungenstockrücken GL (Fig. 2b). This included data from repeat cycles 3 to 15 (April 2019 to May 2022; Smith et al., 2021), with cycles 1 and 2 omitted due to instrument off-pointing at the start of the mission. ATL06 provides surface height measurements every 20 m along-track, calculated by averaging the level-2 geolocated photon data over 40 m segments (Smith et al., 2019).

We obtained coincident tide amplitudes at the most seaward point of each ICESat-2 ground track per cycle from the circum-Antarctic inverse tide model CATS2008 (Howard et al., 2019) using the pyTMD software (Sutterley et al., 2019). CATS2008 is an update of the model described by Padman et al. (2002) that is widely considered the best performing tidal model in the region, in part due to its assimilation of ICESat altimetry data over large ice shelves (King et al., 2011). We note that this tide information was just used for reference and not directly in the calculation of locations of Point F . We also selected three locations 35–40 km seaward of Bungenstockrücken (Fig. 2a) to extract a time series of tide heights over a single year (centred on 1 January 2020). These are used to calculate annual tide distributions across the width of the region, which is important to give context to the range of tides (and therefore locations of Point F) that have been sampled by ICESat-2 along each RGT over the study period. All modelled tides were extracted from locations seaward of the Making Earth System Data Records for Use in Research Environments (MEaSUREs) Point H to ensure that they represented the ice shelf in hydrostatic equilibrium.

3.2 Repeat-track laser altimetry

RTLA is a dynamic method for GL estimation that uses repeat measurements made at different phases of the tidal cycle to estimate the temporal change in the surface elevation of the ice shelf caused by ocean tides to identify the limits of tidal flexure (Points F and H) as proxies for GL location




Figure 2. Method used to define three types of reference profiles for elevation anomaly calculations along ICESat-2 ground tracks, as illustrated for RGT 559 GT3L at the Bungenstockrücken ice plain. (a) Location of Bungenstockrücken on the Ronne Ice Shelf. Red diamonds show the locations used to calculate annual tide distributions. (b) ICESat-2 ground tracks crossing the MEaSUREs and Antarctic Surface Accumulation and Ice Discharge (ASAID) GLs in the study area (Rignot et al., 2016; Bindschadler and Choi, 2011). (c–e) Repeat surface elevation profiles and anomalies of RGT 559 GT3L, each using a different reference profile: (c) *mean* profile calculated using cycles 4, 6, 8, 9, 11, 12 and 13; (d) *neutral* profile of cycle 13; and (e) *lowest-sampled* tide profile of cycle 9. The reference profile at the zero-elevation anomaly is shown as a dashed black line, and the latitude of the derived positions of Point *F* for each cycle along-track are shown as coloured circles along the *x* axis. Dashed coloured lines show the modelled tide height at the time of each cycle at the most seaward point along-track (in e these are differenced from the tide of cycle 9). (f) Histogram of CATS2008-modelled tide heights experienced over a single year, centred on 1 January 2020 (Howard et al., 2019), with vertical dashed coloured lines showing the tide height at each repeat cycle. (g) Tidal time series showing tide height, velocity and phase during the 24 h period before and after the ICESat-2 overpass for each repeat cycle at RGT 559 GT3L.

(Fricker and Padman, 2006). This method typically involves the following three steps:

- i. preparing data,
- ii. defining the reference elevation profile,
- iii. locating the limits of tidal flexure by calculating elevation anomalies relative to the reference profile.

In this study we present updates to steps ii and iii that allow us to improve the temporal sampling of short-term tidal GL migration. We describe each of these steps below.

3.2.1 Preparing data

We obtained ATL06 surface height data along all ICESat-2 ground tracks in the study area (Smith et al., 2021) using the `icepyx` software library (Scheick et al., 2023, version 0.3.3). We removed low-quality measurements caused by cloud cover, blowing snow or background photon clustering using the `ATL06_quality_summary` parameter. We rejected all values without a valid associated across-track slope measurement and/or where the minimum segment difference exceeds 1 m, which is calculated as the minimum absolute difference between each segment's endpoints and those of its two neighbours (Arendt et al., 2020). Following the method described by Li et al. (2020), we applied a cross-track slope correction to minimise the effects of rough terrain (Smith et al., 2019), and for each ground track we omitted repeat cycles with fewer than 50 % valid measurements along-track, as they are deemed unreliable for GZ calculation. We then performed RTLA analysis along each ground track using the remaining high-quality repeat cycles, as described in Sect. 3.2.2 and 3.2.3.

3.2.2 Defining the reference elevation profile

The reference profile is important for identifying the limits of tidal flexure, as it is used as a baseline from which to calculate elevation anomalies for each ICESat-2 repeat sampled at different tidal states. The *mean* elevation profile from all sampled repeats has been used as the reference profile in all previous RTLA studies to locate one or two Point *F* positions per ground track (Fricker and Padman, 2006; Fricker et al., 2009; Brunt et al., 2010b, 2011; Li et al., 2020, 2022a, b). This allows us to derive one Point *F* per ground track, which is useful for measuring long-term change but does not make sense for mapping short-term tidal GL migration (for the reasons discussed below). Here we propose two alternative approaches: using a reference profile sampled at a *neutral* tide or at the *lowest-sampled* tide. These are illustrated for RGT 559 GT3L in Fig. 2. Before selecting one of these approaches for our study, we first evaluated the results from each and considered the situations for which each would be appropriate. To test each approach, we fitted a cubic B-spline approximation to the valid repeat surface elevation profiles

per ground track, using a smoothing parameter of 0.7. The use of an interpolated profile rather than the raw data points is more robust, providing a method for handling small along-track data gaps in individual profiles, which can be consistently applied across all repeat tracks.

- i. *Mean elevation profile* (Fig. 2c). The mean elevation profile in all prior studies has been calculated as the mean of the interpolated profiles of all sampled repeat cycles along each ground track. However, this reference profile does not represent the true mean ice shelf surface corresponding to zero (neutral) tide. Instead, it gives us the along-track mean of the ice shelf surfaces at the tide phases sampled by ICESat-2. This can introduce an observation bias; for example, the mean reference elevation profile would be biased high if most sampled repeats of a single RGT are acquired at positive tides or during one or two extreme high tides. This particularly affects areas experiencing large tidal GL migration, as can be seen clearly in Fig. 2c, where the anomalies for cycles 6 and 13 fall below 0 m between 80.96 and 80.88° S. This might incorrectly suggest that tidal flexure extends this far inland, when in fact it reflects the skew of the mean reference profile caused by GL migration further inland at higher tides (cycles 4 and 12). Given that we locate Point *F* along-track where the anomaly gradient for each cycle first deviates from 0 m, this would lead us to incorrectly locate just two Point *F* positions along this ground track: one for cycle 4 at ~ 80.99° S and one for all other cycles together at ~ 81.96° S (Fig. 2c). As a result, we see a systematic inland bias in the derived GL position when using the mean reference profile, leading to a large underestimation of the total extent of tidal GL migration along this ground track. We note that this ambiguity could be minimised by reducing the number of repeats used to calculate the mean reference profile to two; however this would require the mean tide profile to be recalculated several times per track with different pairs of repeat cycles, which would both increase the processing time and prevent a direct comparison of anomaly profiles between cycles.

- ii. *Neutral tide profile* (Fig. 2d). To more closely represent the true mean ice shelf surface, we can use the elevation profile from a single cycle sampled for a neutral tide profile as the reference profile. Ideally, this would use the profile sampled close to a 0 m tide during the neap phase, such as cycle 13 on Bungenstockrücken for RGT 559 GT3L (Fig. 2g). Using the neutral tide profile removes the effect of the skewed mean in the inner regions of the GZ and improves our interpretation of flexure between tides, as we are now directly comparing the surface profiles between two individual tidal states for each cycle. However, it still raises the same issue that for all cycles with a negative elevation anomaly (i.e. here with

tides below neutral) we would locate Point *F* too far inland, where anomalies first deviate from 0 m (e.g. cycles 9 and 11 in Fig. 2d). We are also unable to locate Point *F* for cycles sampled close to the neutral tide profile as there is no deviation in elevation from the reference profile (cycles 6 and 13 in Fig. 2d). Again, this would mean we would underestimate the total extent of tidal GL migration by overlooking migration that occurs at tides below a neutral level. Moreover, many ICESat-2 RGTs have not been sampled close to a neutral tide, which itself is also difficult to accurately define, because at deep grounding zones the ice surface may not respond adequately in phase with the ocean tide. This means it would not be possible to apply this approach consistently across multiple ground tracks.

- iii. *Lowest-sampled tide profile* (Fig. 2e). Alternatively, we propose that the elevation profile of the repeat cycle sampled at the lowest sampled tide can be used as the reference profile, identified as the cycle with the lowest elevation anomaly at the most seaward sampled point. Again, by using an elevation profile from a single cycle as the reference, we are calculating anomalies against a “real” ice shelf surface profile, giving a more realistic representation of the impact of tidal forcing on ice shelf flexure. This overcomes the issue of the mean reference profile (calculated from multiple cycles of data) being skewed by inland flexure at higher tides, but, unlike the method using the neutral tide profile, it has the additional advantage that it can be applied consistently across RGTs, enabling automation. Most importantly, it allows us to locate Point *F* for cycles sampled at lower tides that are missed when using both the mean and neutral reference profiles. For RGT 559 GT3L we identify five individual Point *F* positions using the lowest-sampled reference profile (Fig. 2e) as opposed to two with the traditional mean approach (Fig. 2c) or four using the neutral tide profile (Fig. 2d). Locating the most seaward Point *F* (from cycle 11 in Fig. 2e) gives us a more realistic baseline for the low-tide GL position against which to quantify the extent of migration at each higher sampled tide. Note that we cannot locate Point *F* for the lowest-sampled tide itself (e.g. cycle 9 in Fig. 2e) as there is no lower profile to compare to, but this is a common problem across all methods.

The type of reference profile (mean, neutral or lowest sampled tide) chosen to calculate elevation anomalies ultimately depends on the scientific question being posed and the tidal sampling by the ICESat-2 RGTs. We must also consider the impact of signals in the repeat elevation profiles that are not associated with tidal displacement, which can be especially prominent where there are large time gaps between repeat cycles. For example, in regions of fast-flowing ice, elevation gradients will advect through the GZ between repeat passes, or alternatively surface mass balance or long-term ice dy-

namic thickness change could alter the repeat-track elevation profiles and obscure the tidal signal. These issues could be addressed by using shorter subsections of the whole time series to define the choice of reference profile. For this study we chose to apply the reference track of the lowest-sampled tide profile for RTLA analysis across the whole 3-year time series as we found it to be the most effective for assessing *tidal* GL migration. Regardless of the chosen reference profile, the results will be limited by the range of the tidal amplitudes that have been sampled by ICESat-2 along each RGT. The CATS2008 tide model can provide context for this; for example Fig. 2f shows that along RGT 559 the lowest 3 % and highest 7 % of tides have not been sampled, meaning we are unable to locate the full extent of the migration of Point *F* at these most extreme tides. The tide phase and velocity at each repeat cycle can also affect the observed flexure profile; therefore we also extract this information from CATS2008 to aid our interpretations (Fig. 2g).

3.2.3 Locating the limits of flexure in regions of ephemeral grounding

Most previous GZ studies using RTLA located a single Point *F* per track (Fricker and Padman, 2006; Fricker et al., 2009; Brunt et al., 2010b; Li et al., 2020, 2022a, b). However, with our method using the lowest-sampled tide profile as the reference profile, it is possible to locate multiple positions of Point *F* along each ICESat-2 ground track, each sampled at a different tidal state. To achieve this, we calculated the elevation anomalies per repeat cycle against the reference profile for each ground track, as is shown for RGT 559 GT3L in Fig. 3. The region where the elevation anomaly is close to 0 m is interpreted as fully grounded ice, and where it is close to the modelled tide prediction with zero gradient it is interpreted as freely floating ice in hydrostatic equilibrium. We compared this with modelled tide predictions from CATS2008 to distinguish the tidal signal from other causes of elevation change, including surface mass balance and ice dynamics. We then located Point *F* for each repeat cycle using an automated technique adapted from Li et al. (2022a). First, we applied a low-pass Butterworth filter to the elevation anomalies per cycle, using a normalised cut-off frequency of 0.016 and an order of 5 to remove the high-frequency noise whilst retaining the shape of each anomaly curve. From this, we located Point *F* where the anomaly gradient first deviates from 0 m and the second derivative of the anomaly peaks (Fig. 3b–g). To ensure the correct estimation of Point *F*, we restricted the choice of peak to where anomaly values are < 0.25 m. In certain cases, the chosen peak of the second derivative was visibly incorrect; i.e. the peak related to Point *F* was not selected but instead a peak related to another cause of elevation change (e.g. crevasse migration). Since this only affected a small minority of tracks in this study area, it was dealt with by manually adjusting the choice of peak to ensure the peak relating to Point *F* was




Figure 3. Method used to locate Point *F* per repeat cycle from ICESat-2 elevation anomalies, as illustrated for RGT 559 GT3L. (a) Elevation anomalies for repeat cycles 4, 6, 8, 11, 12 and 13, each calculated against the lowest-sampled tide reference profile of cycle 9. Dashed coloured lines indicate the CATS2008-modelled tide height per cycle, differenced from the tide of cycle 9 (Howard et al., 2019). (b–g) Individual elevation anomalies for each repeat cycle, with the low-pass-filtered anomalies in black and the second derivative of the filtered anomalies in green. Vertical dashed lines show the along-track latitude where Point *F* has been located from the peak(s) of the second derivative. (h) Map showing the derived position of Point *F* per repeat cycle along RGT 559 GT3L. (i) Along-track migration distance of Point *F* per cycle for RGT 559 GT3L as a function of coincident tide height, overlain on the histogram of tide heights experienced over a single year (centred on 1 January 2020).

selected consistently. We note that more advanced filtering could be applied to address this problem in a more automated way if scaled up to a larger area, but it would likely cause some loss of useful data. For each Point *F* we then extracted the latitude and longitude as well as the coincident tide height and phase at the time of ICESat-2 overpass from CATS2008, which allows us to map the migration of Point *F* across the tide cycle (Fig. 3h). Finally, to investigate modes of tidal GL migration along each ground track we plotted the distance between Point *F* and the location of the furthest seaward Point *F* against the coincident tide height for each repeat cycle, distinguishing between those sampled during rising vs. falling tides (Fig. 3i).

3.3 Study region

We tested our updated RTLA methodology along all ICESat-2 ground tracks crossing a 220 km section of GL at the ice plain north of Bungenstockrücken, located on the southern Ronne Ice Shelf between the Institute and Möller ice streams (Fig. 2). This is an ideal location to study tidal GL migration for the following four reasons.

1. *Large tidal range with ephemeral grounding.* The southern Ronne Ice Shelf experiences predominantly semidiurnal tides with a 7–8 m total tidal range, one of the largest in Antarctica (Padman et al., 2018). Ephemeral grounding has already been identified over this ice plain in earlier ICESat studies (Brunt et al., 2011).
2. *ICESat-2 sampling.* The high-latitude location, GZ orientation and low cloud cover leads to a high density of unobstructed ICESat-2 tracks crossing normally to the GL (67 RGTs, each with six separate ground tracks).
3. *Low ice velocity.* Ice flows across the GL at $\sim 5 \text{ m yr}^{-1}$ (Rignot et al., 2017), resulting in little advection of surface features between repeat profiles over the 3-year study period, which could otherwise affect the success of RTLA.
4. *Evidence for past change.* There is evidence that this region has undergone significant GL retreat in the past, and as the upstream ice sheet is currently grounded over a $> 1.5 \text{ km}$ deep basin with a retrograde slope, it is potentially susceptible to future ice loss (Ross et al., 2012; Siegert et al., 2013). In general, ice plains are a good location to monitor early signs of change as they are highly sensitive to small perturbations in ice dynamics, sea level, surface mass balance, basal melt and sediment deposition (Brunt et al., 2011; Horgan et al., 2013b).

We compared our results at Bungenstockrücken against five existing GL datasets (Table 1).

4 Results

4.1 Extent of tidal grounding line migration at Bungenstockrücken

Our results reveal the spatial pattern of tidal GL migration at Bungenstockrücken across a $\sim 1300 \text{ km}^2$ zone of ephemeral grounding (Fig. 4). Across the central part of the study area (between Areas A and B), we observe the inland limit of tide flexure (Point *F*) migrating by 5 to 15 km across the GZ, although the total migration distance may be greater in regions where the full tide range has not yet been sampled by ICESat-2. In this central region, Point *F* measured at the highest tides consistently reaches several kilometres further inland than the MEaSURES GL and as far inland as the ICESat-2 break in slope. The spatial pattern of short-term tidal GL migration is not consistent across the region, with the most migration observed in Areas A and B (Fig. 4a).

In Area A and across the central section we observe a fairly consistent migration of Point *F* with the tide. This is exemplified by the pattern of RGT 559 GT3L (Fig. 4b, c), where we observe Point *F* migrating progressively further inland as tides increase, totalling 10.4 km of GL migration across the $\sim 4.5 \text{ m}$ sampled tide range (90.3 % of total tide range). At the highest sampled tide of $+2.08 \text{ m}$ (cycle 4), Point *F* reaches as far inland as Point *I_b*, but since 7 % of tides experienced in the region exceed $+2.08 \text{ m}$, it is possible that flexure extends even further inland at these higher tides. In Area B, we observe two main clusters of Point *F*, possibly indicating the presence of two stable GL locations (one landward and one seaward), depending on the tide. This pattern can be seen in the surface and anomaly profiles of RGT 574 GT3L (Fig. 4d, e), which show $\sim 11.5 \text{ km}$ of separation between the seaward GL position near 81.45° S and the landward GL position near 81.55° S . Similarly to RGT 559 GT3L, the landward GL position at the highest sampled tide for RGT 574 GT3L coincides with the ICESat-2 Point *I_b*, just inland of the large surface undulation visible in the elevation profile (Fig. 4d). We note that in parts of Area B the landward GL position extends up to 5 km further inland than the ASAIID Point *I_b*, up to 3 km beyond the ICESat-2 Point *I_b*, and is positioned around the prominent surface undulations visible in the ice surface topography.

In the areas west of Area A and east of Area B, Point *F* is consistently located just inland ($< 2 \text{ km}$) of the ASAIID and ICESat-2 break in slope with little tidal migration. This is consistent with the surface features of a typical GZ (i.e. non-ice-plain) (Schoof, 2011), indicating that these regions are beyond the edges of the ice plain.

4.2 Long-term grounding zone change at Bungenstockrücken

To assess if there has been any long-term GZ change in our study region, we compared our results to previous GL




Figure 4. Tidal GL migration at Bungenstockrücken, measured by ICESat-2 RTLA. **(a)** The location of Point F at every sampled repeat cycle per ICESat-2 ground track is coloured by percent of maximum tide height, overlain on the Reference Elevation Model of Antarctica (REMA) 8 m DEM (Howat et al., 2018), MEaSURES GL (Rignot et al., 2016), ASAID break-in-slope GL (Bindschadler and Choi, 2011) and the most recent ICESat-2-derived break in slope (Point I_b) along each ground track (Li et al., 2022b). The two areas with the largest observed GL migration are highlighted as Areas A and B, and the locations for RGT 559 GT3L, 537 GT3L and 574 GT3L are shown. **(b–c)** The repeat elevation profiles and anomalies of RGT 559 GT3L, with repeat cycles coloured by tide (as percent of maximum tide height) and labelled in **(c)**. The location of ICESat-2 Point I_b is marked with a vertical dashed line (Li et al., 2022b). **(d–e)** The equivalent repeat elevation profiles and anomalies for RGT 574 GT3L.

datasets (Fig. 5). By locating multiple Point F positions along each ground track between high and low tide, we have been able to assess almost the entire width of the zone of ephemeral grounding at Bungenstockrücken for the first time. It is difficult to make a direct comparison to other GL products (Fig. 5a, b), which only locate a single position of Point F (Rignot et al., 2016; Brunt et al., 2010a; Li et al., 2022b). Nonetheless, we see that across the central section of Bungenstockrücken, the ICESat-2-derived Point F positions for the highest $\sim 60\%$ of tides are located consistently inland of the MEaSURES GL, by $\sim 2\text{--}10\text{ km}$ (Fig. 5a). The ICESat-derived Point F positions are mostly located towards the middle of the band of our ICESat-2-derived Point F positions (Fig. 5a), although the maximum inland Point “ F_2 ” positions (Brunt et al., 2011) match more closely our ICESat-2-derived Point F positions at high tide. There has been little long-term change in the position of Point I_b between ICESat, ASAID and ICESat-2, apart from a few locations in Areas A

and B where we observe several kilometres of landward migration (Fig. 5c).

4.3 Modes of tidal grounding line migration

We investigated the relationship between modelled tide height, tide phase and GL migration distance for the ICESat-2 ground tracks at Bungenstockrücken and identified four distinct modes of tidal GL migration (Fig. 6). We manually classified these modes based on visual assessment of the most dominant behaviour. To increase confidence in the classification, we included only tracks where $> 2\text{ m}$ of the tide range was sampled with a total distance of tidal GL migration of $> 1\text{ km}$. This removes tracks where small uncertainties in GL position associated with tide phase, basal properties and long-term surface change are sufficient to obscure the detail in the patterns of GL migration required for this type of classification.

Table 1. Existing grounding line datasets used in this study to compare to our ICESat-2 RTLA results.

Grounding line dataset	Continuous line/discrete points	Satellite(s)	Method	Acquisition date	Reference
MEaSURES Point F	Line	RADARSAT-2	DInSAR	2009	Rignot et al. (2016)
ICESat Point F	Points	ICESat	RTLA	2003–2009	Brunt et al. (2010a)
ICESat Point F_2 (the maximum inland flexure limit at high tide along eight individual ICESat ground tracks at Bungenstockrücken)	Points	ICESat	RTLA	2003–2009	Brunt et al. (2011)
ICESat-2 Point F	Points	ICESat-2	RTLA	March 2019–September 2020	Li et al. (2022b)
ASAID Point I_b (Antarctic Surface Accumulation and Ice Discharge)	Line	Landsat 7, ICESat	Photoclinometry and laser altimetry	Landsat 7: 1999–2003, ICESat: 2003–2009	Bindschadler and Choi (2011)
ICESat-2 Point I_b	Points	ICESat-2	Laser altimetry	March 2019–September 2020	Li et al. (2022b)




Figure 5. (a) Comparison of the ICESat-2-derived Point F positions at Bungenstockrücken to the following: (i) the ICESat-derived Point F (Brunt et al., 2010a); (ii) the ICESat-derived maximum inland Point F_2 (Brunt et al., 2011); and (iii) the MEaSURES GL (Rignot et al., 2016), derived from 2009 RADARSAT-2 data. (b) Comparison of the ICESat-2-derived Point F positions per repeat cycle from this study to the ICESat-2-derived Point F dataset from Li et al. (2022b) that locates a single Point F per ground track. (c) Comparison between Point I_b from ASAID (Bindschadler and Choi (2011); derived from Landsat 7 and ICESat), ICESat (Brunt et al., 2010a) and ICESat-2 (Li et al., 2022b).




Figure 6. Four modes of tidal grounding line migration observed along ICESat-2 ground tracks at Bungenstockrücken. Three examples are shown per mode, which are classified from top to bottom as the following: (a–c) linear, with a linear relationship between tide height and GL migration distance; (d–f) asymmetric, with higher tides causing an increased rate of GL migration; (g–i) threshold, with a sharp tide threshold above which there is significant GL migration; and (j–l) hysteresis, with a difference in the impact of tidal variability on GL migration during the rising- vs. falling-tide phase. In each example we show the along-track migration distance between each Point F and Point F measured at the lowest-sampled tide as a function of coincident tide height from CATS2008 (Howard et al., 2019). Note the variable y-axis scale between ground tracks. The orientation of the triangle indicates whether the repeat cycle was sampled during a rising- or falling-tide phase. Histograms show the distribution of tide heights experienced across a single year, centred on 1 January 2020. (m) Map of Bungenstockrücken showing ICESat-2 ground tracks coloured by migration mode (excluding tracks with < 1 km tidal migration or < 2 m tide range sampled).

1. *Linear* (Fig. 6a–c). In this mode we observe a *linear* relationship between tide height and GL migration distance. Within the sampled tide range, the GL migrates by 2.2 km with every metre of tidal change in Fig. 6a–b and by 1.7 km m^{-1} in Fig. 6c. Considering a simplified semidiurnal spring tide regime where the tide rises from its minimum (–3 m) to maximum (+3 m) within 6 h, this could translate to an average GL retreat rate of about 2 km h^{-1} . We observe little difference between GL positions sampled at rising vs. falling tides. However, we note that, for RGT 781 GT1R (Fig. 6b) there is only one Point F measured during falling tide, and for all ground tracks we cannot determine whether the lin-

ear response extends out to the most extreme tides. The linear mode of tidal GL migration is seen on ground tracks across the study area but is particularly clustered east of Area A (Fig. 6m).

2. *Asymmetric* (Fig. 6d–f). In this mode there is an *asymmetric* relationship between tide height and GL migration, with a large increase in the GL migration rate observed at higher tides. For tides below about +0.5 m in Fig. 6d and below 0 m in Fig. 6e–f, Point F positions are clustered close together, indicating that there is little tidal GL migration within the lower tidal range. However, as the tide increases beyond this point the

rates of GL migration increases sharply to linear rates of $+2.5 \text{ km m}^{-1}$ in Fig. 6d, $+1.8 \text{ km m}^{-1}$ in Fig. 6e and $+4.1 \text{ km m}^{-1}$ in Fig. 6f. Again, outside of the sampled tide range the magnitudes and modes of GL migration are unknown. Ground tracks with an asymmetric mode of tidal GL migration are mainly clustered in Area A and just west of Area B (Fig. 6m).

3. *Threshold* (Fig. 6g–i). In this mode there is little GL migration up to a certain tide *threshold*, beyond which the GL migrates significantly with just a small increase in tide. The tide threshold in Fig. 6g is at $\sim +0.7 \text{ m}$ and at $\sim +0.4 \text{ m}$ in Fig. 6h, and in both examples a $< 0.3 \text{ m}$ rise in the tide across the threshold causes the GL to migrate inland by over 10 km. At peak spring tide (across the full 6 m tide range) this could feasibly occur within 18 min, translating to a possible maximum GL migration rate exceeding 30 km h^{-1} . On either side of the threshold in Fig. 6g and i, further tide change does not cause much more GL migration. This supports the interpretation that in some locations two fairly stable GL positions (landward and seaward) can exist, as observed in Area B (Fig. 4). Considering the histogram of tide distributions, we can infer from this that the GL is at the landward position $\sim 26\%$ of the time in Fig. 6g, $\sim 38\%$ in Fig. 6h and $\sim 22\%$ in Fig. 6i. The majority of ground tracks with the threshold mode of tidal GL migration are located in Area B (Fig. 6m).
4. *Hysteresis* (Fig. 6j–l). In this mode there is a *hysteresis* between tidal forcing and GL migration, related to the phase of the sampled tide at each measurement of Point *F*. This means we observe a different GL migration behaviour between rising and falling tides. For example, along RGT 34 GT1R (Fig. 6j) we note a landward and seaward cluster of Point *F* positions separated by about 15 km, similar to the threshold pattern seen in Fig. 6g–i. However, instead of a single tidal threshold that defines if the GL is in the landward or seaward position, in Fig. 6j we observe two different thresholds for the rising- and falling-tide phases. From this we infer that at the lowest tides the GL is located in the seaward position, but as the tide rises and crosses some threshold between $+0.5$ and $+1.2 \text{ m}$ there is enough buoyant uplift force to unground the ice. This then causes ocean water to intrude into the GZ cavity and, due to the likely very flat bed, allows for the GL to migrate by up to 15 km to its landward position. As the tide rises beyond $+1.2 \text{ m}$, the GL does not migrate any further within the sampled tide range. However, as the tide turns and falls below $+0.5 \text{ m}$ there is not an immediate readvance of the GL to its seaward position. Instead, the GL is observed to remain at its landward position until the tide has fallen beneath some secondary threshold at about -0.8 m . A similar, if less pronounced, hysteresis mode is observed in Fig. 6k–l. Hysteresis is observed

on ground tracks across the study area but particularly in Areas A and B and close to ground tracks with asymmetric and threshold modes (Fig. 6m).

The spatial pattern of tidal GL migration at Area B confirms the presence of a landward and seaward GL, with the location at any given epoch varying between these depending on the tide (Fig. 7a–c). The repeat elevation profiles and anomalies of RGT 537 GT3L reveal more detail about the hysteresis mechanism observed in this region (Fig. 7d–g). Along this ground track, cycles 9 and 12 are sampled at a very similar absolute tide height ($\sim 0.25 \text{ m}$ higher at cycle 9 vs. 12), but their elevation anomaly profiles look very different (Fig. 7f–g). This can be explained by comparing the coincident tide phases, which shows that cycle 9 was sampled during a rising tide, whereas cycle 12 was sampled during a falling tide. The extra bump further inland in the anomalies for cycle 12 (between 81.5 and 81.57°S ; Fig. 7g) indicates that the ice shelf surface between the seaward and landward GLs has not yet fully recovered from its high-tide position. We discuss the potential causes and implications of this in Sect. 5.

We note that the four modes of tidal GL migration described here are not necessarily discrete categories. This is highlighted in Fig. 6h, which shows that along RGT 1016 GT3R we observe both linear and threshold behaviours in different parts of the tidal cycle. Hysteresis may also be superimposed on top of the other modes, as is arguably visible in Fig. 6a, but will become much more clearly identifiable as more repeats are acquired by ICESat-2. We emphasise that there is greater uncertainty in the categorisation of these modes where there are gaps in the sampling of certain tidal ranges, for example above $+0.5 \text{ m}$ in Fig. 6h and k, and there is subjectivity introduced by the manual classification of modes. This could explain the overlap between tracks with different modes in Fig. 6m and particularly those with hysteresis. Our ability to understand and identify these different tidal behaviours will greatly improve as ICESat-2 collects an increasing volume of repeat measurements over the GZ, which should sample more of the tide range. This will improve our confidence in the classification of different modes of tidal GL migration.

5 Discussion

The improved repeat-track sampling capabilities of ICESat-2 together with our updated RTLA methodology offers several advantages over other satellite-based techniques for monitoring tidal GZ behaviour. Using this approach, our observation of widespread short-term tidal GL migration at Bungenstockrücken has implications for how long-term change is assessed and interpreted and for how the GL is represented in models. Furthermore, monitoring the dynamic movement of the GL provides an independent satellite method to illuminate bed characteristics and processes that can influence




Figure 7. (a) The spatial pattern of tidal GL migration observed in Area B, Bungenstockkrücken. Point *F* identified for every repeat cycle of each ICESat-2 ground track is coloured by percent of maximum tide height. (b–c) Schematics depicting the flow of water in between the seaward and landward GL positions inferred from ICESat-2 RTLA results in Area B, during the rising- and falling-tide phase. (d–e) The repeat elevation profiles and anomalies of RGT 537 GT3L, calculated using the lowest-sampled tide reference profile from cycle 4. Repeat cycles are coloured by tide (as percent of maximum tide height), and the cycle numbers are labelled in (e) alongside the CATS2008-modelled tide heights, differenced from the tide at cycle 4. (f–g) The individual elevation anomalies for cycles 9 and 12 against the lowest-sampled tide profile of cycle 4. Inset panels show the time series of tide heights for the 24 h before and after the time of the ICESat-2 overpass at each cycle, showing the coincident tide phase and velocity.

stability in the GZ region. We expand on each of these implications below.

5.1 Advantages of our ICESat-2 RTLA method for monitoring tidal grounding line behaviour

First, the use of ICESat-2 itself offers several advantages that improve the spatial sampling of tidal GL behaviour compared to other satellites. Its track spacing and six-beam pattern greatly increases the density of measurements compared to ICESat, with 402 separate ground tracks crossing the Bungenstockkrücken GZ (using the six beams from 67 ICESat-

2 RGTs), compared to 23 ICESat tracks. There is also an improvement in along-track sampling, with ATL06 surface height measurements calculated every 20 m along-track (averaged from 40 m overlapping segments), compared to a spacing of 172 m with ICESat (Schutz et al., 2005). This allows us to better resolve finer-scale details of the shape and configuration of the GL and observe how these may be affected by the tide. Furthermore, ICESat-2's repeat-track configuration and sensitivity to low-amplitude tidal variations allow us to attribute GL migration to specific tidal states, which is not possible with CryoSat-2 radar altimetry (Dawson and Bamber, 2017). ICESat-2 coverage also extends to 88° S, en-

abling observations at GLs located further south than the limit of most DInSAR coverage.

Second, the use of our updated RTLA technique improves the temporal sampling of tidal GL behaviour. All previous RTLA approaches with both ICESat and ICESat-2 used a mean reference profile calculated from all sampled cycles to derive a single Point F per track (Fricker and Padman, 2006; Fricker et al., 2009; Brunt et al., 2010b; Li et al., 2020, 2022a, b). Our method, by using the lowest-sampled tide as the reference profile, allows us to locate multiple positions of Point F along each ground track as it migrates with the tide. This produces GL measurements at fine temporal resolution, which enables a closer investigation of tidal GZ processes that occur over hourly to daily timescales. Within the time period of this study, we locate up to 11 Point F positions per ground track, each sampled at a different tide, totalling 2402 measurements across the 402 ground tracks. The number of valid measurements per ground track (of a maximum possible 13) largely depends on the number of repeats obscured by cloud cover. As more cycles of ICESat-2 data are acquired in the future, the temporal sampling of tidal GZ behaviour will continue to increase.

5.2 Implications for grounding line monitoring

Our observation of up to 15 km of tidal GL migration across the Bungenstockrücken ice plain is one of the largest migration values reported in Antarctica, comparable to the ~ 15 km found at the Amery GL in DInSAR by Chen et al. (2023). The signal is an order of magnitude higher than the fastest annual average GL retreat of 1.8 km yr^{-1} observed at Pine Island Glacier between 1992 and 2011 (Park et al., 2013) and several orders of magnitude higher than average long-term Antarctic GL migration rates (Konrad et al., 2018). For comparison, the highest estimated deglaciation retreat rates from palaeo-records are > 2.1 and $> 10 \text{ km yr}^{-1}$ for Thwaites and Larsen A, respectively (Dowdeswell et al., 2020; Graham et al., 2022). Therefore, to accurately measure long-term GZ change it is necessary to isolate the spatially variable pattern of tidal GL migration. At Bungenstockrücken, the observed difference between the MEaSURES GL and the ICESat-2-derived Point F positions (Fig. 5a) could suggest that there has been either up to 10 km of GL retreat since 2009 or no change, depending on whether the MEaSURES GL was produced from data sampled at high or low tides. This shows the value of providing a precise timestamp in addition to the acquisition date when producing a GL data product, as the tide amplitude at the time of measurement will inform our interpretation about whether a short-term, temporary tidal fluctuation or a longer-term, dynamic change has occurred. Similarly, in this particular location, it is also difficult to quantify any long-term GL migration between the ICESat and ICESat-2 records (Fig. 5a) because the signal is dominated by tidal variability and different methods have been used to derive Point F . The fact that the small number

of ICESat-derived maximum inland Point F_2 positions from Brunt et al. (2011) are broadly consistent with our highest tide ICESat-2 Point F positions suggests that there has not been significant retreat of the maximum inland flexure limit over the past 10 to 15 years.

Bungenstockrücken provides a reasonable upper bound for rates of tidal GL migration around the ice sheet, as it is located in an ice plain region with low bed slopes and one of the highest tide ranges in Antarctica. Although tidal migration is likely to be much smaller at most Antarctic GLs (depending on tide range and bed properties, including slope and friction coefficient), even an order of magnitude less may still be sufficient to obscure early signs of long-term retreat in response to ice shelf thinning (Milillo et al., 2017; Li et al., 2023) and to impact ice dynamics. Therefore, when assessing long-term GL migration rates from sparse observations of GL location through time, we must consider the tidal state associated with each measurement epoch. Ideally, long-term GL location estimates should be made at the same tidal state or tide difference; however, given the scarcity of GL measurements in most locations, this has historically not been feasible. In lieu of this, our updated ICESat-2 RTLA methodology could be applied around the ice sheet margin to provide a constraint on the expected tidal range and mechanisms of GL movement. This will enable better evaluation of the impact of tides on previously derived estimates of GL position and ultimately improve the confidence with which we can assess long-term GL change.

The identification of different modes of tidal GL migration using ICESat-2 RTLA (Fig. 6) could be used to identify regions that are particularly sensitive to ice thickness changes, similar to Schmeltz et al. (2001) who showed that areas of ephemeral grounding can reveal subtle changes in ice thickness. In regions with a threshold or hysteresis mode of tidal GL migration, an observed decrease in the tide threshold that defines whether the GL is located in the landward or seaward position could indicate a retreat of the GL and signal dynamic ice thinning. Conversely, if the tide threshold rises it could signal GL advance and ice thickening. We suggest that Area B at Bungenstockrücken (Fig. 7) would be an ideal place to monitor this, with its wide zone of ephemeral grounding (up to 15 km) and numerous ICESat-2 tracks exhibiting either threshold or hysteresis GL migration modes. Monitoring ephemeral grounding in this way could enable us to discriminate between inferred thickness changes due to changing firn-air content and true ice thickness change (Moholdt et al., 2014). This is useful for detecting early signs of ice sheet change, particularly as the accuracy with which altimeters can measure long-term ice thickness change at the GL is limited by our lack of knowledge about the flotation state of the ice across the GZ (further complicated by tidal GL migration).

5.3 Implications for modelling tidal ice shelf flexure and grounding line migration

The representation of GZ processes in ice sheet models strongly impacts projections of future ice sheet mass balance (Arthern and Williams, 2017). The 15 km observed tidal GL migration at Bungenstockrücken is much larger than standard model grid spacing at the GL, which is typically less than 2 km and down to 250 m where mesh refinement is applied (Cornford et al., 2020). Similarly, the time step of an ice sheet model is generally much longer than 1 d; therefore, prescribing a subdaily change in GL position is not generally possible. It is beyond the scope of this study to parameterise tidal GL effects in ice sheet models at dissimilar scales, but any “short-term fixed” GL derived from observations must be applied in a consistent manner, whether this is using the highest, lowest or mean tide position. This choice of a consistent GL is complicated by the fact that different satellite-derived GL products use different surface proxies (i.e. Points F vs. I_b), each of which relate differently to the basal-stress boundary, which is the parameter that we generally want to represent in models. Moreover, the extent to which tidal GL migration can blur the basal-stress boundary in areas of ephemeral grounding is largely unknown.

Our results at Bungenstockrücken show that the use of ICESat-2 RTLA to map Point F across the tide cycle can improve our understanding of the relationship between short-term GL migration and the long-term ice sheet surface profile. We observe that Point F is located seaward of Point I_b for most of the tidal cycle but see that at the highest tides it consistently migrates as far inland as Point I_b (Fig. 4). As the break in slope (I_b) is one of the key surface signatures of the change in basal shear stress at the GL (Schoof, 2011), these observations imply that from a stress-balance perspective the ice sheet geometry is controlled by the furthest inland position that Point F reaches at the highest tides. Although the ice shelf regrounds seaward of the break in slope at low tides, the variable stresses over the ice plain area do not appear to substantially affect the surface profile. There are substantial deviations between flexure-derived GLs and those derived from the break in slope, which may therefore be partly explained by sampling of GL positions at either mean or non-maximum tides. We propose that the information derived from ICESat-2 RTLA, by better constraining the extent and pattern of tidal GL migration, could be used to better inform the choice of GL in models where observations are used.

Our results provide observational validations for different numerical models of tidal ice flexure and GL migration. The existence of different modes of tidal GL migration within a region experiencing similar tidal forcing indicates that there is a spatial heterogeneity in local bed properties. Therefore, in regions like Bungenstockrücken that experience large tidal GL migration, models of tide flexure that do not allow for movement of the GL (Holdsworth, 1969; Schmeltz et al., 2002; Walker et al., 2013) cannot produce realistic repre-

sentations of the basal stress or GL velocity change. Sayag and Worster (2013) developed an elastic beam model coupled to an elastically deforming bed that allows for the GL to migrate proportionally with tidal forcing. We observe a similar linear mode of GL migration within the sampled tide range along numerous ICESat-2 ground tracks at Bungenstockrücken (Fig. 6a–c), as has also recently been observed in parts of West Antarctica using CSK DInSAR (Milillo et al., 2022). In contrast, Tsai and Gudmundsson (2015) modelled the GL as an elastic fracture problem forced by the tidally induced increase in ocean water pressure, concluding that tidally induced GL migration in areas with prograde bed slopes is asymmetric, with the GL migrating up to 9 times further inland at high tide than it migrates seaward at low tide. Along several ICESat-2 ground tracks at Bungenstockrücken we observe a similar kind of asymmetry in the mode of tidal GL migration (Fig. 6d–f). It is likely that the shape and stiffness of the bed exert strong controls over the mode of GL migration (Sayag and Worster, 2013), but independent bed information is generally not of high enough resolution to further inform our interpretation of the flexure patterns observed by our ICESat-2 RTLA method.

Our ICESat-2 observations may also inform the representation of elastic and viscous stresses in these models of tidal flexure and GL migration. Both the Sayag and Worster (2013) and Tsai and Gudmundsson (2015) models use purely elastic frameworks, justified due to the short timescales (< 12 h) over which the forcing acts. In reality, ice deforms viscoelastically (Reeh et al., 2003), and so these models overlook any time-dependent viscoelastic deformation that follows the initial elastic response to tides. The hysteresis mode of tidal GL migration observed at Bungenstockrücken (Figs. 5j–l and 6) reveals a time delay of the response of the ice shelf surface (and perhaps also the subglacial till) to tidal forcing. Similar observations in Greenland have shown a difference in the surface deflection profile between rising and falling tides and have been described well by viscoelastic beam models that capture this lag between the change in the tide and the ice shelf response (Reeh et al., 2000, 2003; Wild et al., 2017). Currently no models of tidal flexure exist that use both a viscoelastic framework and allow for migration of the GL, but this may be necessary to better represent these kinds of processes in GZs subject to high tidal variability and could account for the discrepancies between model predictions and observations.

Modes of tidal GL migration and why they differ spatially can also provide insights into the underlying bed topography. As a specific example, we expect that the threshold mode of GL migration (Fig. 6g–i) is likely to only exist in regions with a very flat bed where the ice is very close to flotation, whereby a small increase in tide across this threshold can provide enough buoyant force to unground the ice and cause large GL migration. Area B at Bungenstockrücken is likely grounded over a very flat bed, as we observe a cluster of ground tracks with a threshold mode of tidal GL migration

(Fig. 6m). While Brunt et al. (2011) presented a method to calculate bed slopes from tidal GL migration along ICESat tracks based on a hydrostatic assumption, this assumed a linear relationship between migration distance and tide height, which, as our findings indicate, is not necessarily valid in the GZ. Therefore, our results support the conclusion of Tsai and Gudmundsson (2015) of urging caution when quantitatively inferring bed slopes from measurements of tidal GL migration. We note that changes in tidal GL behaviour on longer timescales can also provide insight into bed slopes, as was observed at Thwaites, where the total width of the zone of ephemeral grounding increased from 0.5 km in 1996 to 2.5 km in 2016–2017 as the GL retreated into a region of flatter bed topography (Milillo et al., 2019). There is, therefore, potential for gaining further information about the bed by identifying any changes in tidal migration modes over time.

The distribution of *modes* of tidal GL migration as well as the *width* and *rate* of tidal GL migration around Antarctic GZs is valuable for informing the choice of parameterisation of tidal flexure mechanisms, basal shear stress and/or basal melt in larger-scale ice sheet models. For example, Mosbeux et al. (2023) considered two different GL migration parameterisations to assess the impact of seasonal sea surface height variability on ice velocities over the Ross Ice Shelf. The first considered an asymmetric treatment of GL migration with elastic fracture mechanics introduced by Tsai and Gudmundsson (2015) and a constant bed slope, and the second defined migration using hydrostatic equilibrium of the GL with Bedmap2 bed slopes. They found that the asymmetric treatment led to larger modelled ice shelf velocity variability that more closely matched GNSS observations. Given that the mechanism of migration varies spatially, even across relatively small distances (as we have demonstrated), obtaining reliable observations of the spatial distribution of these GL migration modes around Antarctica is needed to inform the parameterisation of GL processes in larger ice sheet modelling studies. Our study at Bungenstockrücken has successfully demonstrated how ICESat-2 RTLA can be used as a first step towards this.

5.4 Insights into tidal processes in the ice shelf–ocean–subglacial system

The cyclical, twice-daily ungrounding and tidal flushing of ocean water up to 15 km into the GZ has implications for several processes contributing to the net dynamics of this system (Fig. 8). Many of these processes are poorly understood with few observations, and so the use of ICESat-2 RTLA to study tidal GL behaviour in settings like Bungenstockrücken could provide important new insights and constraints.

Little is known about the oceanography close to most of the Antarctic GZ, but the ability to observe the tidal variability in cavity geometry with ICESat-2 RTLA will allow us to model the potential melt rates in these anomalous regions. At Bungenstockrücken, our results indicate that an ad-

ditional $\sim 1300 \text{ km}^2$ of ice ungrounds twice-daily between low and high tide (Fig. 4), allowing for the relatively warm ocean water to reach deeper into the GZ cavity and exposing this additional area to basal melt. The cold High-Salinity Shelf Water (HSSW) beneath the Ronne Ice Shelf causes relatively low basal melt rates (compared to warmer regions like the Amundsen Sea that are affected by Circumpolar Deep Water), but depression of the seawater freezing point under pressure means it still has the capacity to melt ice at the GL (Nicholls and Østerhus, 2004; Makinson et al., 2011; Naughten et al., 2021). The very shallow water column ($< 100 \text{ m}$) immediately seaward of Bungenstockrücken (Johnson and Smith, 1997) suggests that tidal mixing would be strong here and is likely to be enhanced as water is “pumped” in and out of the cavity as the GL migrates with the tides. Milillo et al. (2019) report significant basal melt over the entire zone of tidal GL migration at Thwaites, and such short-term tidal fluctuations in melt rate and tidal mixing have been shown to impact long-term GL retreat in less stable regions (Graham et al., 2022). Given that projections of future ice sheet mass loss are highly sensitive to basal melt at the GL (Arthern and Williams, 2017; Goldberg et al., 2019; Adusumilli et al., 2020), this once again calls into question the use of fixed GLs with zero melt in ice sheet models. The type of information about tidal GZ behaviour that we can derive from ICESat-2 RTLA will therefore be valuable for improving parameterisations of melt at the GL.

Variation in basal stresses in the GZ caused by tidal flexure and GL migration can also impact the flow of upstream and downstream ice. Grounding and ungrounding of the ice shelf throughout the tidal cycle has several effects: it changes the area affected by basal drag, modifies the driving stress through changes in surface slope (e.g. Makinson et al., 2012; Mosbeux et al., 2023), modulates hydrostatic and flexural stresses (e.g. Rosier and Gudmundsson, 2016), and impacts buttressing (e.g. Robel et al., 2017). Rosier and Gudmundsson (2020) concluded that the presence of tides enhances ice flow across the entire Filchner–Ronne Ice Shelf by 21%. Since ice velocities across the Bungenstockrücken GL do not exceed $\sim 5 \text{ m yr}^{-1}$ (Rignot et al., 2017), the impact of tidal GL migration on absolute flow speed here is likely to be low, but it may be more significant in faster-flowing regions, such as the Rutford Ice Stream (Minchew et al., 2017).

Our results have implications for testing models and hypotheses in subglacial hydrology. It has been suggested that tidal ice shelf flexure acts to pump ocean water upstream of the GL into the subglacial hydrological system, with implications for basal lubrication, basal melt and till compaction (Christianson et al., 2013; Horgan et al., 2013a; Sayag and Worster, 2013; Walker et al., 2013; Drews et al., 2017; Robel et al., 2022). Airborne radar data indicate the possible presence of brackish water several kilometres inland of the Bungenstockrücken GL (Corr, 2021), which may provide evidence for upstream seawater incursion driven by tidal GL migration. Modelling of this process by Warburton et al. (2020)




Figure 8. Schematic summarising impacts of tidal flexure and GL migration on the ice shelf–ocean–subglacial system. These include impacts on the following: ice velocity (Gudmundsson, 2007; Thompson et al., 2014; Robel et al., 2017, 2022; Mosbeux et al., 2023); the GZ surface profile (e.g. Schoof, 2011); inland seawater incursion (Warburton et al., 2020; Robel et al., 2022); variations in subglacial water pressure (Rosier et al., 2015; Begeman et al., 2020); the formation of estuarine features in the GZ (Horgan et al., 2013a); enhanced basal melt (Makinson et al., 2011; Mueller et al., 2012); basal shear stress (Anandakrishnan et al., 2003); till compaction, leading to changes in basal lubrication (Walker et al., 2013; Christianson et al., 2013; Sayag and Worster, 2013); the formation of tidal seabed ridges (Dowdeswell et al., 2020; Graham et al., 2022); and ocean circulation through tidal mixing. The landward and seaward tidal GL positions are labelled as F_{\max} and F_{\min} .

indicated that water is pumped upstream into the subglacial environment much faster during a rising tide than it drains as the tide falls. Similarly, Stubblefield et al. (2021) model the formation of an extended grounding zone during the falling tide phase, comprising a thin layer of water that drains more slowly than the seaward GL migration due to the flexural response of the ice. The hysteresis mode of GL migration observed at Bungenstockrücken (Figs. 6j–l and 7) suggests there is a similar lag in the response of the ice shelf surface to tidal forcing between rising- and falling-tide phases, which could provide observational support for this theory. For example, the pattern observed in Area B (Fig. 7) is likely to indicate that a subglacial channel or cavity exists in between the seaward and landward GL positions that fills with water at high tide (above a certain threshold). As the tide falls, the water drains away more slowly and the ice around the seaward GL position (a possible topographic high) regrounds before the landward cavity has fully emptied, forcing the water to be flushed out from this channel to the east (Fig. 7b, c). This could be a pattern similar to tidal beach drainage and supports the idea of “estuary” type features at the ice sheet margins (Horgan et al., 2013a). Knowledge of the modes of tidal GL migration around the coast derived from ICESat-2 RTLA will be useful for constraining such regional models.

6 Summary and outlook

We have presented an updated method for determining Antarctic ice shelf grounding line (GL) location from ICESat-2 repeat-track laser altimetry (RTLA). We show that, by using the elevation profile at the lowest-sampled tide as the reference profile for calculating anomalies, it is possible to locate the inland limit of tidal flexure at different stages of the tidal cycle. This temporal resolution provides insights into grounding zone (GZ) processes on tidal timescales that are difficult to observe with other methods. We have applied the technique to the GZ of an ice plain north of Bungenstockrücken on the southern Ronne Ice Shelf, where we observe > 15 km of tidal GL migration. This is one of the largest reported distances of ephemeral grounding anywhere in Antarctica, and it demonstrates the necessity of accounting for short-term tidal GL migration to accurately measure long-term migration. It also calls into question the use of a fixed or slowly moving GL boundary in ice sheet models, which does not represent the full range of GL migration behaviour. Our updated ICESat-2 RTLA technique also allows us to observe different modes of tidal GL migration, and we classify four modes at the Bungenstockrücken ice plain: linear, asymmetric, threshold and hysteresis. This can provide observational validation for models of tidal ice shelf flexure, GL migration and subglacial hydrology at the GZ and offers a tool for future explorations of bed characteristics, basal melt and basal

shear stress in these sensitive parts of the ice sheet. We suggest that monitoring the change in extent and modes of tidal GL migration around Antarctica could provide vital early indications of wider ice sheet change.

Based on the findings of this study, we make four recommendations for future work.

1. Any future satellite-derived measurement of GL position should be accompanied with a timestamp (at least to the closest hour), ideally also with coincident tide height and phase. For DInSAR this information should be provided per image, and for RTLA it should be provided for both the reference and repeat cycle tides. This will allow for a more robust assessment of the impact of tidal processes on individual measurements of the GL position.
2. Future studies should scale this analysis up across Antarctica to derive a continent-wide dataset of the extent and mode of tidal GL migration. This could be used to improve confidence in long-term records of GL migration.
3. Targeted ground-based surveys in regions subject to extreme tidal variability would be valuable for better understanding the link between these surface observations and the processes taking place within and beneath the ice in GZs on tidal timescales. We suggest that Bungenstockrücken would be a suitable location, due to its high tidal signal, low ice velocity and crevassing and current long-term stability which isolates the tidal signal from long-term change.
4. The observations presented here have only been possible to obtain using ICESat-2. Therefore, it is crucial that we extend our satellite-based *repeat-track* sampling of ice sheet surface elevation beyond this satellite mission to continue this valuable record as ice sheets continue to respond to the changing climate.

Code availability. The code developed for this study can be found at <https://doi.org/10.5281/zenodo.8037209> (Freer, 2023).

Data availability. The ICESat-2 ATL06 data (<https://doi.org/10.5067/ATLAS/ATL06.005>, Smith et al., 2021), ICESat-2 grounding zone products (<https://doi.org/10.5067/RI167B92708M9>, Li et al., 2022b) and MEaSURES grounding line product (<https://doi.org/10.5067/IKBWW4RYHF1Q>, Rignot et al., 2016) used in this study are available from the National Snow and Ice Data Center (NSIDC). The ASAIID grounding line (<https://doi.org/10.7265/N56T0JK2>, Bindenschadler and Choi, 2011) and ICESat-derived grounding zone products (<https://doi.org/10.7265/N5CF9N19>, Brunt et al., 2010a) are available from the U.S. Antarctic Program Data Center.

Author contributions. BIDF and OJM planned the research; BIDF conducted the data analysis and wrote the manuscript. All authors provided insights regarding the interpretation of data and reviewed and edited the manuscript.

Competing interests. The contact author has declared that none of the authors has any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. This work was led by Bryony I. D. Freer at the British Antarctic Survey (BAS) and the School of Earth and Environment at the University of Leeds. Bryony I. D. Freer was supported by the Natural Environment Research Council (NERC) Satellite Data in Environmental Science (SENSE) Centre for Doctoral Training (grant no. NE/T00939X/1). Anna E. Hogg was supported by the NERC Drivers of Oceanic Change in the Amundsen Sea (DeCAdeS) project (no. NE/T012757/1) and the ESA Polar+ Ice Shelves project (no. ESA-IPL-POE-EF-cb-LE-2019-834). Helen Amanda Fricker and Laurie Padman were supported by NASA (grant no. 80NSSC20K0977). Laurie Padman was also supported by NASA (grant no. 80NSSC21K0911). The authors gratefully acknowledge the National Aeronautics and Space Administration for acquiring the ICESat-2 satellite data.

Financial support. This research has been supported by the Natural Environment Research Council (grant nos. NE/T00939X/1 and NE/T012757/1), the National Aeronautics and Space Administration (grant nos. 80NSSC20K0977 and 80NSSC21K0911), and the European Space Agency (grant no. ESA-IPL-POE-EF-cb-LE-2019-834).

Review statement. This paper was edited by Bert Wouters and reviewed by Tian Li, Pietro Milillo, and Kasia Warburton.

References

- Adusumilli, S., Fricker, H. A., Medley, B., Padman, L., and Siegfried, M. R.: Interannual variations in meltwater input to the Southern Ocean from Antarctic ice shelves, *Nat. Geosci.*, 13, 616–620, <https://doi.org/10.1038/s41561-020-0616-z>, 2020.
- Alley, R. B., Blankenship, D. D., Rooney, S. T., and Bentley, C. R.: Sedimentation beneath ice shelves – the view from ice stream B, *Mar. Geol.*, 85, 101–120, [https://doi.org/10.1016/0025-3227\(89\)90150-3](https://doi.org/10.1016/0025-3227(89)90150-3), 1989.
- Anandakrishnan, S., Voigt, D. E., Alley, R. B., and King, M. A.: Ice stream D flow speed is strongly modulated by the tide beneath the Ross Ice Shelf, *Geophys. Res. Lett.*, 30, 1361, <https://doi.org/10.1029/2002GL016329>, 2003.

- Arendt, A., Scheick, J., Shean, D., Buckley, E., Grigsby, S., Haley, C., Heagy, L., Mohajerani, Y., Neumann, T., Nilsson, J., Markus, T., Paolo, F. S., Perez, F., Petty, A., Schweiger, A., Smith, B., Steiker, A., Alvis, S., Henderson, S., Holschuh, N., Liu, Z., and Sutterly, T.: 2020 ICESat-2 Hackweek Tutorials, Zenodo [code], <https://doi.org/10.5281/zenodo.3966463>, 2020.
- Arthern, R. J. and Williams, C. R.: The sensitivity of West Antarctica to the submarine melting feedback, *Geophys. Res. Lett.*, 44, 2352–2359, <https://doi.org/10.1002/2017GL072514>, 2017.
- Begeman, C. B., Tulaczyk, S., Padman, L., King, M., Siegfried, M. R., Hodson, T. O., and Fricker, H. A.: Tidal Pressurization of the Ocean Cavity Near an Antarctic Ice Shelf Grounding Line, *J. Geophys. Res.-Oceans*, 125, e2019JC015562, <https://doi.org/10.1029/2019JC015562>, 2020.
- Bindschadler, R. and Choi, H.: High-resolution Image-derived Grounding and Hydrostatic Lines for the Antarctic Ice Sheet, U.S. Antarctic Program (USAP) Data Center [data set], <https://doi.org/10.7265/N56T0JK2>, 2011.
- Bindschadler, R., Choi, H., Wichlacz, A., Bingham, R., Bohlander, J., Brunt, K., Corr, H., Drews, R., Fricker, H., Hall, M., Hindmarsh, R., Kohler, J., Padman, L., Rack, W., Rotschky, G., Urbini, S., Vornberger, P., and Young, N.: Getting around Antarctica: new high-resolution mappings of the grounded and freely-floating boundaries of the Antarctic ice sheet created for the International Polar Year, *The Cryosphere*, 5, 569–588, <https://doi.org/10.5194/tc-5-569-2011>, 2011.
- Bojinski, S., Verstraete, M., Peterson, T. C., Richter, C., Simmons, A., and Zemp, M.: The Concept of Essential Climate Variables in Support of Climate Research, Applications, and Policy, *B. Am. Meteorol. Soc.*, 95, 1431–1443, <https://doi.org/10.1175/BAMS-D-13-00047.1>, 2014.
- Brancato, V., Rignot, E., Milillo, P., Morlighem, M., Mouginot, J., An, L., Scheuchl, B., Jeong, S., Rizzoli, P., Bueso Bello, J. L., and Prats-Iraola, P.: Grounding Line Retreat of Denman Glacier, East Antarctica, Measured With COSMO-SkyMed Radar Interferometry Data, *Geophys. Res. Lett.*, 47, e2019GL086291, <https://doi.org/10.1029/2019GL086291>, 2020.
- Brunt, K. M., Fricker, H. A., Padman, L., and O’Neel, S.: ICESat-Derived Grounding Zone for Antarctic Ice Shelves, U.S. Antarctic Program (USAP) Data Center [data set], <https://doi.org/10.7265/N5CF9N19>, 2010a.
- Brunt, K. M., Fricker, H. A., Padman, L., Scambos, T. A., and O’Neel, S.: Mapping the grounding zone of the Ross Ice Shelf, Antarctica, using ICESat laser altimetry, *Ann. Glaciol.*, 51, 71–79, <https://doi.org/10.3189/172756410791392790>, 2010b.
- Brunt, K. M., Fricker, H. A., and Padman, L.: Analysis of ice plains of the Filchner–Ronne Ice Shelf, Antarctica, using ICESat laser altimetry, *J. Glaciol.*, 57, 965–975, <https://doi.org/10.3189/002214311798043753>, 2011.
- Catania, G., Hulbe, C., and Conway, H.: Grounding-line basal melt rates determined using radar-derived internal stratigraphy, *J. Glaciol.*, 56, 545–554, <https://doi.org/10.3189/002214310792447842>, 2010.
- Chen, H., Rignot, E., Scheuchl, B., and Ehrenfeucht, S.: Grounding Zone of Amery Ice Shelf, Antarctica, From Differential Synthetic-Aperture Radar Interferometry, *Geophys. Res. Lett.*, 50, e2022GL102430, <https://doi.org/10.1029/2022GL102430>, 2023.
- Christianson, K., Parizek, B. R., Alley, R. B., Horgan, H. J., Jacobel, R. W., Anandakrishnan, S., Keisling, B. A., Craig, B. D., and Muto, A.: Ice sheet grounding zone stabilization due to till compaction, *Geophys. Res. Lett.*, 40, 5406–5411, <https://doi.org/10.1002/2013GL057447>, 2013.
- Ciraci, E., Rignot, E., Scheuchl, B., Tolpekin, V., Wollersheim, M., An, L., Milillo, P., Bueso-Bello, J.-L., Rizzoli, P., and Dini, L.: Melt rates in the kilometer-size grounding zone of Petermann Glacier, Greenland, before and during a retreat, *P. Natl. Acad. Sci. USA*, 120, e2220924120, <https://doi.org/10.1073/pnas.2220924120>, 2023.
- Cornford, S. L., Seroussi, H., Asay-Davis, X. S., Gudmundsson, G. H., Arthern, R., Borstad, C., Christmann, J., Dias dos Santos, T., Feldmann, J., Goldberg, D., Hoffman, M. J., Humbert, A., Kleiner, T., Leguy, G., Lipscomb, W. H., Merino, N., Durand, G., Morlighem, M., Pollard, D., Rückamp, M., Williams, C. R., and Yu, H.: Results of the third Marine Ice Sheet Model Intercomparison Project (MISMIP+), *The Cryosphere*, 14, 2283–2301, <https://doi.org/10.5194/tc-14-2283-2020>, 2020.
- Corr, H.: Processed airborne radio-echo sounding data from the GRADES-IMAGE survey covering the Evans and Rutford Ice Streams, and ice rises in the Ronne Ice Shelf, West Antarctica (2006/2007) (1.0), NERC EDS UK Polar Data Centre [data set], <https://doi.org/10.5285/C7EA5697-87E3-4529-A0DD-089A2ED638FB>, 2021.
- Corr, H. F. J., Doake, C. S. M., Jenkins, A., and Vaughan, D. G.: Investigations of an “ice plain” in the mouth of Pine Island Glacier, Antarctica, *J. Glaciol.*, 47, 51–57, <https://doi.org/10.3189/172756501781832395>, 2001.
- Dawson, G. J. and Bamber, J. L.: Antarctic Grounding Line Mapping From CryoSat-2 Radar Altimetry, *Geophys. Res. Lett.*, 44, 11886–11893, <https://doi.org/10.1002/2017GL075589>, 2017.
- Dawson, G. J. and Bamber, J. L.: Measuring the location and width of the Antarctic grounding zone using CryoSat-2, *The Cryosphere*, 14, 2071–2086, <https://doi.org/10.5194/tc-14-2071-2020>, 2020.
- Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T. M., Ligtenberg, S. R. M., van den Broeke, M. R., and Moholdt, G.: Calving fluxes and basal melt rates of Antarctic ice shelves, *Nature*, 502, 89–92, <https://doi.org/10.1038/nature12567>, 2013.
- Dowdeswell, J. A., Batchelor, C. L., Montelli, A., Ottesen, D., Christie, F. D. W., Dowdeswell, E. K., and Evans, J.: Delicate seafloor landforms reveal past Antarctic grounding-line retreat of kilometers per year, *Science*, 368, 1020–1024, <https://doi.org/10.1126/science.aaz3059>, 2020.
- Drews, R., Pattyn, F., Hewitt, I. J., Ng, F. S. L., Berger, S., Matsuoka, K., Helm, V., Bergeot, N., Favier, L., and Neckel, N.: Actively evolving subglacial conduits and eskers initiate ice shelf channels at an Antarctic grounding line, *Nat. Commun.*, 8, 15228, <https://doi.org/10.1038/ncomms15228>, 2017.
- Dupont, T. K. and Alley, R. B.: Assessment of the importance of ice-shelf buttressing to ice-sheet flow, *Geophys. Res. Lett.*, 32, L04503, <https://doi.org/10.1029/2004GL022024>, 2005.
- Dutrieux, P., De Rydt, J., Jenkins, A., Holland, P. R., Ha, H. K., Lee, S. H., Steig, E. J., Ding, Q., Abrahamsen, E. P., and Schroder, M.: Strong Sensitivity of Pine Island Ice-Shelf Melting to Climatic Variability, *Science*, 343, 174–178, <https://doi.org/10.1126/science.1244341>, 2014.

- Favier, L., Durand, G., Cornford, S. L., Gudmundsson, G. H., Gagliardini, O., Gillet-Chaulet, F., Zwinger, T., Payne, A. J., and Le Brocq, A. M.: Retreat of Pine Island Glacier controlled by marine ice-sheet instability, *Nat. Clim. Change*, 4, 117–121, <https://doi.org/10.1038/nclimate2094>, 2014.
- Freer, B.: s2002365/Tidal-GL-Migration-ICESat-2: v1.0.0 (v1.0.0), Zenodo [code], <https://doi.org/10.5281/zenodo.8037209>, 2023.
- Fricker, H. A. and Padman, L.: Ice shelf grounding zone structure from ICESat laser altimetry, *Geophys. Res. Lett.*, 33, L15502, <https://doi.org/10.1029/2006GL026907>, 2006.
- Fricker, H. A., Coleman, R., Padman, L., Scambos, T. A., Bohlander, J., and Brunt, K. M.: Mapping the grounding zone of the Amery Ice Shelf, East Antarctica using InSAR, MODIS and ICESat, *Antarct. Sci.*, 21, 515–532, <https://doi.org/10.1017/S095410200999023X>, 2009.
- Friedl, P., Weiser, F., Fluhrer, A., and Braun, M. H.: Remote sensing of glacier and ice sheet grounding lines: A review, *Earth-Sci. Rev.*, 201, 102948, <https://doi.org/10.1016/j.earscirev.2019.102948>, 2020.
- Goldberg, D. N., Gourmelen, N., Kimura, S., Millan, R., and Snow, K.: How Accurately Should We Model Ice Shelf Melt Rates?, *Geophys. Res. Lett.*, 46, 189–199, <https://doi.org/10.1029/2018GL080383>, 2019.
- Graham, A. G. C., Wählin, A., Hogan, K. A., Nitsche, F. O., Heywood, K. J., Totten, R. L., Smith, J. A., Hillenbrand, C.-D., Simkins, L. M., Anderson, J. B., Wellner, J. S., and Larter, R. D.: Rapid retreat of Thwaites Glacier in the pre-satellite era, *Nat. Geosci.*, 706–713, <https://doi.org/10.1038/s41561-022-01019-9>, 2022.
- Gudmundsson, G. H.: Tides and the flow of Rutford Ice Stream, West Antarctica, *J. Geophys. Res.*, 112, F04007, <https://doi.org/10.1029/2006JF000731>, 2007.
- Gudmundsson, G. H., Paolo, F. S., Adusumilli, S., and Fricker, H. A.: Instantaneous Antarctic ice-sheet mass loss driven by thinning ice shelves, *Geophys. Res. Lett.*, 46, 13903–13909, <https://doi.org/10.1029/2019GL085027>, 2019.
- Hogg, A. E., Shepherd, A., Gourmelen, N., and Engdahl, M.: Grounding line migration from 1992 to 2011 on Petermann Glacier, North-West Greenland, *J. Glaciol.*, 62, 1104–1114, <https://doi.org/10.1017/jog.2016.83>, 2016.
- Hogg, A. E., Shepherd, A., Gilbert, L., Muir, A., and Drinkwater, M. R.: Mapping ice sheet grounding lines with CryoSat-2, *Adv. Space Res.*, 62, 1191–1202, <https://doi.org/10.1016/j.asr.2017.03.008>, 2018.
- Holdsworth, G.: Flexure of a Floating Ice Tongue, *J. Glaciol.*, 8, 385–397, <https://doi.org/10.3189/S0022143000026976>, 1969.
- Horgan, H. J., Alley, R. B., Christianson, K., Jacobel, R. W., Anandakrishnan, S., Muto, A., Beem, L. H., and Siegfried, M. R.: Estuaries beneath ice sheets, *Geology*, 41, 1159–1162, <https://doi.org/10.1130/G34654.1>, 2013a.
- Horgan, H. J., Christianson, K., Jacobel, R. W., Anandakrishnan, S., and Alley, R. B.: Sediment deposition at the modern grounding zone of Whillans Ice Stream, West Antarctica, *Geophys. Res. Lett.*, 40, 3934–3939, <https://doi.org/10.1002/grl.50712>, 2013b.
- Howard, S. L., Padman, L., and Erofeeva, S. Y.: CATS2008: Circum-Antarctic Tidal Simulation version 2008 (1), U.S. Antarctic Program (USAP) Data Center [code], <https://doi.org/10.15784/601235>, 2019.
- Howat, I., Morin, P., Porter, C., and Noh, M.-J.: The Reference Elevation Model of Antarctica, Version 1, Harvard Dataverse [data set], <https://doi.org/10.7910/DVN/SAIK8B>, 2018.
- Jenkins, A., Corr, H. F. J., Nicholls, K. W., Stewart, C. L., and Doake, C. S. M.: Interactions between ice and ocean observed with phase-sensitive radar near an Antarctic ice-shelf grounding line, *J. Glaciol.*, 52, 325–346, <https://doi.org/10.3189/172756506781828502>, 2006.
- Johnson, M. R. and Smith, A. M.: Seabed topography under the southern and western Ronne Ice Shelf, derived from seismic surveys, *Antarct. Sci.*, 9, 201–208, <https://doi.org/10.1017/S0954102097000254>, 1997.
- Joughin, I., Smith, B. E., and Holland, D. M.: Sensitivity of 21st century sea level to ocean-induced thinning of Pine Island Glacier, Antarctica, *Geophys. Res. Lett.*, 37, L20502, <https://doi.org/10.1029/2010GL044819>, 2010.
- Joughin, I., Alley, R. B., and Holland, D. M.: Ice-Sheet Response to Oceanic Forcing, *Science*, 338, 1172–1176, <https://doi.org/10.1126/science.1226481>, 2012.
- King, M. A., Padman, L., Nicholls, K., Clarke, P. J., Gudmundsson, G. H., Kulesa, B., and Shepherd, A.: Ocean tides in the Weddell Sea: New observations on the Filchner-Ronne and Larsen C ice shelves and model validation, *J. of Geophys. Res.-Oceans*, 116, C06006, <https://doi.org/10.1029/2011JC006949>, 2011.
- Konrad, H., Shepherd, A., Gilbert, L., Hogg, A. E., McMillan, M., Muir, A., and Slater, T.: Net retreat of Antarctic glacier grounding lines, *Nat. Geosci.*, 11, 258–262, <https://doi.org/10.1038/s41561-018-0082-z>, 2018.
- Li, T., Dawson, G. J., Chuter, S. J., and Bamber, J. L.: Mapping the grounding zone of Larsen C Ice Shelf, Antarctica, from ICESat-2 laser altimetry, *The Cryosphere*, 14, 3629–3643, <https://doi.org/10.5194/tc-14-3629-2020>, 2020.
- Li, T., Dawson, G. J., Chuter, S. J., and Bamber, J. L.: A high-resolution Antarctic grounding zone product from ICESat-2 laser altimetry, *Earth Syst. Sci. Data*, 14, 535–557, <https://doi.org/10.5194/essd-14-535-2022>, 2022a.
- Li, T., Dawson, Geoffrey J., Chuter, Stephen J., and Bamber, Jonathan L.: ICESat-2 L3 Grounding Zone for Antarctic Ice Shelves, Version 1, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/RI67B92708M9>, 2022b.
- Li, T., Dawson, G. J., Chuter, S. J., and Bamber, J. L.: Grounding line retreat and tide-modulated ocean channels at Moscow University and Totten Glacier ice shelves, East Antarctica, *The Cryosphere*, 17, 1003–1022, <https://doi.org/10.5194/tc-17-1003-2023>, 2023.
- Luthcke, S. B., Thomas, T. C., Pennington, T. A., Rebold, T. W., Nicholas, J. B., Rowlands, D. D., Gardner, A. S., and Bae, S.: ICESat-2 Pointing Calibration and Geolocation Performance, *Earth Space Sci.*, 8, e2020EA001494, <https://doi.org/10.1029/2020EA001494>, 2021.
- MacGregor, J. A., Anandakrishnan, S., Catania, G. A., and Winebrenner, D. P.: The grounding zone of the Ross Ice Shelf, West Antarctica, from ice-penetrating radar, *J. Glaciol.*, 57, 917–928, <https://doi.org/10.3189/002214311798043780>, 2011.
- Magruder, L., Brunt, K., Neumann, T., Klotz, B., and Alonzo, M.: Passive Ground-Based Optical Techniques for Monitoring the On-Orbit ICESat-2 Altimeter Geolocation and

- Footprint Diameter, *Earth Space Sci.*, 8, e2020EA001414, <https://doi.org/10.1029/2020EA001414>, 2021.
- Makinson, K., Holland, P. R., Jenkins, A., Nicholls, K. W., and Holland, D. M.: Influence of tides on melting and freezing beneath Filchner-Ronne Ice Shelf, Antarctica, *Geophys. Res. Lett.*, 38, L06601, <https://doi.org/10.1029/2010GL046462>, 2011.
- Makinson, K., King, M. A., Nicholls, K. W., and Hilmar Gudmundsson, G.: Diurnal and semidiurnal tide-induced lateral movement of Ronne Ice Shelf, Antarctica, *Geophys. Res. Lett.*, 39, L06601, <https://doi.org/10.1029/2012GL051636>, 2012.
- Markus, T., Neumann, T., Martino, A., Abdalati, W., Brunt, K., Csatho, B., Farrell, S., Fricker, H., Gardner, A., Harding, D., Jasinski, M., Kwok, R., Magruder, L., Lubin, D., Luthcke, S., Morison, J., Nelson, R., Neunschwander, A., Palm, S., Popescu, S., Shum, C., Schutz, B. E., Smith, B., Yang, Y., and Zwally, J.: The Ice, Cloud, and Land Elevation Satellite-2 (ICESat-2): Science requirements, concept, and implementation, *Remote Sens. Environ.*, 190, 260–273, <https://doi.org/10.1016/j.rse.2016.12.029>, 2017.
- Milillo, P., Rignot, E., Mouginot, J., Scheuchl, B., Morlighem, M., Li, X., and Salzer, J. T.: On the Short-term Grounding Zone Dynamics of Pine Island Glacier, West Antarctica, Observed With COSMO-SkyMed Interferometric Data: PIG Grounding Line Dynamics, *Geophys. Res. Lett.*, 44, 10436–10444, <https://doi.org/10.1002/2017GL074320>, 2017.
- Milillo, P., Rignot, E., Rizzoli, P., Scheuchl, B., Mouginot, J., Bueso-Bello, J., and Prats-Iraola, P.: Heterogeneous retreat and ice melt of Thwaites Glacier, West Antarctica, *Sci. Adv.*, 5, eaau3433, <https://doi.org/10.1126/sciadv.aau3433>, 2019.
- Milillo, P., Rignot, E., Rizzoli, P., Scheuchl, B., Mouginot, J., Bueso-Bello, J. L., Prats-Iraola, P., and Dini, L.: Rapid glacier retreat rates observed in West Antarctica, *Nat. Geosci.*, 15, 48–53, <https://doi.org/10.1038/s41561-021-00877-z>, 2022.
- Minchew, B. M., Simons, M., Riel, B., and Milillo, P.: Tidally induced variations in vertical and horizontal motion on Rutford Ice Stream, West Antarctica, inferred from remotely sensed observations, *J. Geophys. Res.-Earth*, 122, 167–190, <https://doi.org/10.1002/2016JF003971>, 2017.
- Moholdt, G., Padman, L., and Fricker, H. A.: Basal mass budget of Ross and Filchner-Ronne ice shelves, Antarctica, derived from Lagrangian analysis of ICESat altimetry: Ice shelf basal melting from altimetry, *J. Geophys. Res.-Earth*, 119, 2361–2380, <https://doi.org/10.1002/2014JF003171>, 2014.
- Mohajerani, Y., Jeong, S., Scheuchl, B., Velicogna, I., Rignot, E., and Milillo, P.: Automatic delineation of glacier grounding lines in differential interferometric synthetic-aperture radar data using deep learning, *Sci. Rep.*, 11, 4992, <https://doi.org/10.1038/s41598-021-84309-3>, 2021.
- Mosbeux, C., Padman, L., Klein, E., Bromirski, P. D., and Fricker, H. A.: Seasonal variability in Antarctic ice shelf velocities forced by sea surface height variations, *The Cryosphere*, 17, 2585–2606, <https://doi.org/10.5194/tc-17-2585-2023>, 2023.
- Mueller, R. D., Padman, L., Dinniman, M. S., Erofeeva, S. Y., Fricker, H. A., and King, M. A.: Impact of tide-topography interactions on basal melting of Larsen C Ice Shelf, Antarctica, *J. Geophys. Res.*, 117, C05005, <https://doi.org/10.1029/2011JC007263>, 2012.
- Naughten, K. A., De Rydt, J., Rosier, S. H. R., Jenkins, A., Holland, P. R., and Ridley, J. K.: Two-timescale response of a large Antarctic ice shelf to climate change, *Nat. Commun.*, 12, 1991, <https://doi.org/10.1038/s41467-021-22259-0>, 2021.
- Neumann, T. A., Martino, A. J., Markus, T., Bae, S., Bock, M. R., Brenner, A. C., Brunt, K. M., Cavanaugh, J., Fernandes, S. T., Hancock, D. W., Harbeck, K., Lee, J., Kurtz, N. T., Luers, P. J., Luthcke, S. B., Magruder, L., Pennington, T. A., Ramos-Izquierdo, L., Rebold, T., Skoog, J., and Thomas, T. C.: The Ice, Cloud, and Land Elevation Satellite – 2 mission: A global geolocated photon product derived from the Advanced Topographic Laser Altimeter System, *Remote Sens. Environ.*, 233, 111325, <https://doi.org/10.1016/j.rse.2019.111325>, 2019.
- Nicholls, K. W. and Østerhus, S.: Interannual variability and ventilation timescales in the ocean cavity beneath Filchner-Ronne Ice Shelf, Antarctica, *J. Geophys. Res.*, 109, C04014, <https://doi.org/10.1029/2003JC002149>, 2004.
- Padman, L., Fricker, H. A., Coleman, R., Howard, S., and Erofeeva, L.: A new tide model for the Antarctic ice shelves and seas, *Ann. Glaciol.*, 34, 247–254, <https://doi.org/10.3189/172756402781817752>, 2002.
- Padman, L., Siegfried, M. R., and Fricker, H. A.: Ocean Tide Influences on the Antarctic and Greenland Ice Sheets, *Rev. Geophys.*, 56, 142–184, <https://doi.org/10.1002/2016RG000546>, 2018.
- Park, J. W., Gourmelen, N., Shepherd, A., Kim, S. W., Vaughan, D. G., and Wingham, D. J.: Sustained retreat of the Pine Island Glacier, *Geophys. Res. Lett.*, 40, 2137–2142, <https://doi.org/10.1002/grl.50379>, 2013.
- Reeh, N., Mayer, C., Olesen, O. B., Christensen, E. L., and Thomsen, H. H.: Tidal movement of Nioghalvfjærdsfjorden glacier, northeast Greenland: observations and modelling, *Ann. Glaciol.*, 31, 111–117, <https://doi.org/10.3189/172756400781820408>, 2000.
- Reeh, N., Christensen, E. L., Mayer, C., and Olesen, O. B.: Tidal bending of glaciers: a linear viscoelastic approach, *Ann. Glaciol.*, 37, 83–89, <https://doi.org/10.3189/172756403781815663>, 2003.
- Rignot, E.: Tidal motion, ice velocity and melt rate of Petermann Gletscher, Greenland, measured from radar interferometry, *J. Glaciol.*, 42, 476–485, <https://doi.org/10.3189/S0022143000003464>, 1996.
- Rignot, E., Mouginot, J., and Scheuchl, B.: Antarctic grounding line mapping from differential satellite radar interferometry, *Geophys. Res. Lett.*, 38, L10504, <https://doi.org/10.1029/2011GL047109>, 2011.
- Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H., and Scheuchl, B.: Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011, *Geophys. Res. Lett.*, 41, 3502–3509, <https://doi.org/10.1002/2014GL060140>, 2014.
- Rignot, E., Mouginot, J., and Scheuchl, B.: MEaSUREs Antarctic Grounding Line from Differential Satellite Radar Interferometry, Version 2, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/IKBWW4RYHF1Q>, 2016.
- Rignot, E., Mouginot, J., and Scheuchl, B.: MEaSUREs InSAR-Based Antarctica Ice Velocity Map, Version 2, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/D7GK8F5J8M8R>, 2017.
- Robel, A. A., Tsai, V. C., Minchew, B., and Simons, M.: Tidal modulation of ice shelf buttressing stresses, *Ann. Glaciol.*, 58, 12–20, <https://doi.org/10.1017/aog.2017.22>, 2017.

- Robel, A. A., Wilson, E., and Seroussi, H.: Layered seawater intrusion and melt under grounded ice, *The Cryosphere*, 16, 451–469, <https://doi.org/10.5194/tc-16-451-2022>, 2022.
- Rosier, S. H. R. and Gudmundsson, G. H.: Tidal controls on the flow of ice streams, *Geophys. Res. Lett.*, 43, 4433–4440, <https://doi.org/10.1002/2016GL068220>, 2016.
- Rosier, S. H. R. and Gudmundsson, G. H.: Exploring mechanisms responsible for tidal modulation in flow of the Filchner–Ronne Ice Shelf, *The Cryosphere*, 14, 17–37, <https://doi.org/10.5194/tc-14-17-2020>, 2020.
- Rosier, S. H. R., Gudmundsson, G. H., and Green, J. A. M.: Temporal variations in the flow of a large Antarctic ice stream controlled by tidally induced changes in the subglacial water system, *The Cryosphere*, 9, 1649–1661, <https://doi.org/10.5194/tc-9-1649-2015>, 2015.
- Ross, N., Bingham, R. G., Corr, H. F. J., Ferraccioli, F., Jordan, T. A., Le Brocq, A., Rippin, D. M., Young, D., Blankenship, D. D., and Siegert, M. J.: Steep reverse bed slope at the grounding line of the Weddell Sea sector in West Antarctica, *Nat. Geosci.*, 5, 393–396, <https://doi.org/10.1038/ngeo1468>, 2012.
- Sayag, R. and Worster, M. G.: Elastic dynamics and tidal migration of grounding lines modify subglacial lubrication and melting, *Geophys. Res. Lett.*, 40, 5877–5881, <https://doi.org/10.1002/2013GL057942>, 2013.
- Scambos, T. A., Haran, T. M., Fahnestock, M. A., Painter, T. H., and Bohlander, J.: MODIS-based Mosaic of Antarctica (MOA) data sets: Continent-wide surface morphology and snow grain size, *Remote Sens. Environ.*, 111, 242–257, <https://doi.org/10.1016/j.rse.2006.12.020>, 2007.
- Scheick, J., Leong, W. J., Bisson, K., Arendt, A., Bhushan, S., Fair, Z., Hagen, N. R., Henderson, S., Knuth, F., Li, T., Liu, Z., Pinunno, R., Ravinder, N., Setiawan, L. D., Sutterley, T., Swinski, J. P., and Anubhav: icepyx: querying, obtaining, analyzing, and manipulating ICESat-2 datasets. In *The Journal of Open Source Software (v0.6.4_JOSS, Vol. 8, Number 84, p. 4912)*, Zenodo [code], <https://doi.org/10.5281/zenodo.7806097>, 2023.
- Schmeltz, M., Rignot, E., and MacAyeal, D. R.: Ephemeral grounding as a signal of ice-shelf change, *J. Glaciol.*, 47, 71–77, <https://doi.org/10.3189/172756501781832502>, 2001.
- Schmeltz, M., Rignot, E., and MacAyeal, D.: Tidal flexure along ice-sheet margins: comparison of InSAR with an elastic-plate model, *Ann. Glaciol.*, 34, 202–208, <https://doi.org/10.3189/172756402781818049>, 2002.
- Schmidt, B. E., Washam, P., Davis, P. E. D., Nicholls, K. W., Holland, D. M., Lawrence, J. D., Riverman, K. L., Smith, J. A., Spears, A., Dichek, D. J. G., Mullen, A. D., Clyne, E., Yeager, B., Anker, P., Meister, M. R., Hurwitz, B. C., Quartini, E. S., Bryson, F. E., Basinski-Ferris, A., Thomas, C., Wake, J., Vaughan, D. G., Anandkrishnan, S., Rignot, E., Paden, J., and Makinson, K.: Heterogeneous melting near the Thwaites Glacier grounding line, *Nature*, 614, 471–478, <https://doi.org/10.1038/s41586-022-05691-0>, 2023.
- Schoof, C.: Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *J. Geophys. Res.*, 112, F03S28, <https://doi.org/10.1029/2006JF000664>, 2007.
- Schoof, C.: Marine ice sheet dynamics. Part 2. A Stokes flow contact problem, *J. Fluid Mech.*, 679, 122–155, <https://doi.org/10.1017/jfm.2011.129>, 2011.
- Schutz, B. E., Zwally, H. J., Shuman, C. A., Hancock, D., and DiMarzio, J. P.: Overview of the ICESat Mission, *Geophys. Res. Lett.*, 32, L21S01, <https://doi.org/10.1029/2005GL024009>, 2005.
- Siegert, M., Ross, N., Corr, H., Kingslake, J., and Hindmarsh, R.: Late Holocene ice-flow reconfiguration in the Weddell Sea sector of West Antarctica, *Quaternary Sci. Rev.*, 78, 98–107, <https://doi.org/10.1016/j.quascirev.2013.08.003>, 2013.
- Smith, A. M.: The use of tiltmeters to study the dynamics of Antarctic ice-shelf grounding lines, *J. Glaciol.*, 37, 51–58, <https://doi.org/10.3189/S0022143000042799>, 1991.
- Smith, B., Fricker, H. A., Holschuh, N., Gardner, A. S., Adusumilli, S., Brunt, K. M., Csathó, B., Harbeck, K., Huth, A., Neumann, T., Nilsson, J., and Siegfried, M. R.: Land ice height-retrieval algorithm for NASA’s ICESat-2 photon-counting laser altimeter, *Remote Sens. Environ.*, 233, 111352, <https://doi.org/10.1016/j.rse.2019.111352>, 2019.
- Smith, B., Fricker, H. A., Gardner, A., Siegfried, M. R., Adusumilli, S., Csathó, B. M., Holschuh, N., Nilsson, J., Paolo, F. S., and the ICESat-2 Science Team: ATLAS/ICESat-2 L3A Land Ice Height, Version 5, Boulder Colorado USA, NASA National Snow and Ice Data Center Distributed Active Archive Center [data set], <https://doi.org/10.5067/ATLAS/ATL06.005>, 2021.
- Stephenson, S. N., Doake, C. S. M., and Horsfall, J. A. C.: Tidal flexure of ice shelves measured by tiltmeter, *Nature*, 282, 496–497, <https://doi.org/10.1038/282496a0>, 1979.
- Stubblefield, A. G., Spiegelman, M., and Creyts, T. T.: Variational formulation of marine ice-sheet and subglacial-lake grounding-line dynamics, *J. Fluid Mech.*, 919, A23, <https://doi.org/10.1017/jfm.2021.394>, 2021.
- Sutterley, T. C., Markus, T., Neumann, T. A., van den Broeke, M., van Wessem, J. M., and Ligtenberg, S. R. M.: Antarctic ice shelf thickness change from multimission lidar mapping, *The Cryosphere*, 13, 1801–1817, <https://doi.org/10.5194/tc-13-1801-2019>, 2019.
- Thomas, R. H.: The Dynamics of Marine Ice Sheets, *J. Glaciol.*, 24, 167–177, <https://doi.org/10.1017/S0022143000014726>, 1979.
- Thompson, J., Simons, M., and Tsai, V. C.: Modeling the elastic transmission of tidal stresses to great distances inland in channelized ice streams, *The Cryosphere*, 8, 2007–2029, <https://doi.org/10.5194/tc-8-2007-2014>, 2014.
- Tsai, V. C. and Gudmundsson, G. H.: An improved model for tidally modulated grounding-line migration, *J. Glaciol.*, 61, 216–222, <https://doi.org/10.3189/2015JoG14J152>, 2015.
- Vaughan, D. G.: Tidal flexure at ice shelf margins, *J. Geophys. Res.-Sol. Ea.*, 100, 6213–6224, <https://doi.org/10.1029/94JB02467>, 1995.
- Walker, R. T., Parizek, B. R., Alley, R. B., Anandkrishnan, S., Riverman, K. L., and Christianson, K.: Ice-shelf tidal flexure and subglacial pressure variations, *Earth Planet. Sc. Lett.*, 361, 422–428, <https://doi.org/10.1016/j.epsl.2012.11.008>, 2013.
- Warburton, K. L. P., Hewitt, D. R., and Neufeld, J. A.: Tidal Grounding-Line Migration Modulated by Subglacial Hydrology, *Geophys. Res. Lett.*, 47, e2020GL089088, <https://doi.org/10.1029/2020GL089088>, 2020.
- Wild, C. T., Marsh, O. J., and Rack, W.: Viscosity and elasticity: a model intercomparison of ice-shelf bending in an Antarctic grounding zone, *J. Glaciol.*, 63, 573–580, <https://doi.org/10.1017/jog.2017.15>, 2017.