

Global Ocean Monitoring: Recent Evolution, Current Status, and Predictions

Prepared by
Climate Prediction Center, NCEP/NOAA

September 10, 2024

<http://www.cpc.ncep.noaa.gov/products/GODAS/>

This project, to deliver real-time ocean monitoring products, is implemented
by CPC in cooperation with NOAA's Global Ocean Monitoring and Observing Program (GOMO)



- Overview
- Recent highlights
 - Pacific Ocean
 - Arctic & Antarctic Oceans
 - Indian Ocean
 - Atlantic Ocean
- Global SSTA Predictions

• Pacific Ocean

- ENSO neutral conditions continued with Niño3.4 = -0.1°C in Aug 2024.
- The positive SSTA and a strong negative phase of the PDO persisted in the North Pacific with PDOI = -1.9 in Aug 2024.
- Strong subsurface warming has persisted in the central North Pacific Ocean since 2020.

• Arctic & Antarctic Oceans

- The average Arctic sea ice extent for Aug 2024 was 5.2 million km^2 , ranking as the fourth lowest August since 1979.
- Antarctic sea ice extent continues to track as the 2nd historical low value.
- CPC forecasts a below-normal Arctic sea ice extent minimum in Sep 2024.

• Indian Ocean

- Positive SSTAs dominated the tropical Indian Ocean basin in Aug 2024.
- The Indian Ocean dipole (IOD) was neutral in Aug 2024.


• Atlantic Ocean

- Hurricane activity was relatively quiet in Aug 2024.
- Strong Marine heat waves persisted in the north tropical Atlantic and the central extratropical Atlantic Ocean .
- Patterns of SST, OLR and low-level wind anomalies over Jun-Aug 2024 were consistent with impact of tropical Atlantic meridional mode on local climate.


Global Oceans

Global SST Anomaly ($^{\circ}\text{C}$) and Anomaly Tendency

AUG 2024 SST Anomaly ($^{\circ}\text{C}$)
(1991–2020 Climatology)



AUG 2024 – JUL 2024 SST Anomaly ($^{\circ}\text{C}$)




- SSTs were above average in the west-central Pacific Ocean, while were near to below average SSTs were present in the eastern Pacific.
- Strong positive SSTAs continued in the mid-latitude of the North Pacific and North Atlantic Oceans.
- Positive SSTAs dominated the tropical Atlantic, Indian and Southern Oceans.

- Both positive and negative SSTA tendencies were observed in the North Pacific and Atlantic Oceans.
- Weak SSTA tendencies were present in the central-eastern equatorial Pacific Ocean.


SSTAs (top) and SSTA tendency (bottom). Data are derived from the Olv2.1 SST analysis, and anomalies are departures from the 1991-2020 base period means.

Longitude-Depth Temperature Anomaly and Anomaly Tendency in 2°S-2°N

AUG 2024 Eq. Temp Anomaly (°C)
(GODAS, Clima. 91-20)



AUG 2024 - JUL 2024 Eq. Temp Anomaly (°C)




- Negative temperature anomalies persisted along the central-eastern thermocline in the Pacific, while positive temperature anomalies continued in the western Pacific.
- Positive temperature anomalies were present in the upper 150m of the Indian Ocean.
- Positive temperature anomaly dominated the upper ocean of the Atlantic.

- Both positive and negative temperature anomaly tendencies were present along the thermocline in the Pacific Ocean.
- Negative temperature anomaly tendency emerged in the eastern Atlantic Ocean.

Equatorial depth-longitude section of ocean temperature anomalies (top) and anomaly tendency (bottom). Data is from the NCEP's GODAS. Anomalies are departures from the 1991-2020 base period means.

Global Monthly SST anomaly and Marine Heat Waves

OISSTv2.1 AUG2024 SST Anom. (°C)
Hatch area: MHW on AUG-2024-31





- MHWs were present in much of the central North Atlantic, tropical North Atlantic, north central - western extratropical Pacific and southern tropical Indian Oceans.
- MHW emerged in the Barents Sea and Kara Sea.

((Left panel) Monthly SST anomaly (shaded) and locations experiencing marine heat waves (hatched) by the end date labelled in the plot. (right panel) SST evolution at a specific location. Green line and blue line are the 90th percentile and daily climatology, respectively. Shaded area denotes the periods experiencing MHW. MHW is defined as a prolonged warming exceeding 90th percentile of daily SST for at least 14 consecutive days. Data is derived from NCEI OISSTv2.1 and the reference period is 1991-2020

Tropical Pacific Ocean and ENSO Conditions


Evolution of Pacific Niño SST Indices

Monthly Tropical Pacific SST Anomaly
(OISSTv2.1, 1991–2020 Climatology)



- Niño3.4 switched to negative phase in Aug 2024, with Niño3.4 = -0.1°C.
- Niño4 remained near 0.5 °C.
- Compared with Aug 2023, the tropical eastern Pacific was cooler in Aug 2024
- The indices may have differences if based on different SST products.


Niño region indices, calculated as the area-averaged monthly mean SSTAs (°C) for the specified region. Data are derived from the Olv2.1 SST analysis, and anomalies are departures from the 1991-2020 base period means.



SSTAs (top-left), SSTA tendency (top-right), Outgoing Long-wave Radiation (OLR) anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right; positive means heat into the ocean), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the Olv2.1 SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1991-2020 base period means.

Evolution of Equatorial Pacific Surface Zonal Current Anomaly (cm/s)

U (15m), cm/s, 2°S–2°N (Shading=Anomaly; Contour=1993–2020 Clim)




- Anomalous westward currents have been observed since mid-Dec 2023 with a weakening since May 2024.
- Anomalous westward currents strengthened in the eastern Pacific during Aug 2024 in GODAS, while were near average in OSCAR.

Equatorial Pacific Ocean Temperature Pentad Mean Anomaly

TAO


GODAS




- Negative ocean temperature anomalies in the central Pacific weakened during Aug 2024.
- Positive SSTA persisted in east of 150°W in TAO, while negative SSTAs were observed in GODAS.

Potential Impact of Missing TAO DATA on GODAS


of Daily Temp. Profiles in AUG 2024




of Daily Temp. Profiles in JUN 2011



of Daily Temp. Profiles in Upper 300m [120E-80W,10S-10N]




of Daily Temp. Profiles every 5 Days in 15-1N
(5 is 100% return rate, buoys at Eq)




- There are fewer moorings reporting in the eastern Pacific since Nov 2023.
- TAO mooring profiles number in Aug 2024 is close to the historical low since 1993.

Multiple Ocean Reanalysis Intercomparison: Temperature Anomaly and Tendency at Equator


Anomalous Temperature (C) Averaged in 1S-1N: AUG 2024



AUG 2024 - JUL 2024 1S-1N Temp Anomaly (C)











Anomalous Temperature (C) at z=5m minus OISST: AUG 2024







North Pacific & Arctic Oceans

Last 3-month North Pacific SST, SLP, and uv925 anomalies



Pacific Decadal Oscillation (PDO) Index




- The PDO has been in a negative phase since Jan 2020 and weakened slightly with PDOI = -1.9 in Aug 2024.
- Statistically, ENSO leads PDO by 3-4 months, through teleconnection via atmospheric bridge, with El Niño (La Niña) associated with positive (negative) PDO Index, but this relationship has weakened in recent years.




• PDO is defined as the 1st EOF of monthly ERSST v3b in the North Pacific for the period 1900-1993. PDO index is the standardized projection of the monthly SST anomalies onto the 1st EOF pattern.

Subsurface Temperature Anomaly in the Northcentral Pacific

Anomalous Temperature (C) in [160E-150W, 30N-45N]




Last 10 year




- Positive temperature anomaly (>0.9°C) has persisted in the upper 100m since 2020, and penetrated to 300m since Jul 2024.
- Subsurface warming in the last four years is the strongest episode since 1979.

Two Oceanic PDO indices

SST-based PDO (Wen et al. 2014: GRL)



H300-based PDO (Arun and Wen 2016: Mon. Wea. Rev.)



- The negative phase of PDO has persisted since Jan 2020 with PDOI = -1.9. in Aug 2024.

- Negative H300-based PDO index has persisted since Nov 2016, with HPDO = - 2.2 in Aug 2024.



- SST-based PDO index has considerable variability both on seasonal and decadal time scales.

- H300-based PDO index highlights the slower variability and encapsulates an integrated view of temperature variability in the upper ocean.

SST-based PDO is defined as the 1st EOF of monthly ERSST v3b in the North Pacific for the period 1900-1993. PDO index is the standardized projection of the monthly ERSSTv5 SST anomalies onto the 1st EOF pattern. H300-based Pacific Decadal Oscillation is defined as the projection of monthly mean H300 anomalies from NCEP GODAS onto their first EOF vector in the North Pacific. PDO indices are downloadable from https://www.cpc.ncep.noaa.gov/products/GODAS/ocean_briefing.shtml.




Weekly SST anomaly and MHWs in the North Pacific

Weekly OISSTv2.1 Anom. (°C)
Hatch area: MHW location





- MHWs have persisted in the central North Pacific since early May, but the coverage area varied owing to high frequency of atmosphere circulation.

(Left panel) Weekly SST anomaly (shaded) and locations experience Marine heat waves (hatched) by the date labelled in the plot. (right panel) SST evolution at a specific location. Green line and blue line denote the seasonal 90th percentile and daily climatology, respectively. Shaded area denotes the periods experiencing MHW. MHW is defined as a discrete prolonged warmer than 90th percentile of daily SST for at least 14 days. Data is derived from NCEI OISSTv2.1 and the climatology reference period is 1991-2020




- Average Arctic sea ice extent during Aug 2024 was 5.2 million square kilometers, ranking as the fourth lowest Aug in the satellite record.



- Antarctic sea ice extent continues to track as the second lowest ice extent in the satellite data record.

NCEP/CPC Arctic Sea Ice Extent (SIE) Forecast





https://www.cpc.ncep.noaa.gov/products/people/jszhu/seaice_seasonal/index.html

- CPC forecasts call for a below normal sea ice extent minimum in the Arctic in Sep 2024.


Indian Ocean

Evolution of Indian Ocean SST Indices



- Basin wide SST warming continued in the tropical Indian Ocean.
- Indian dipole mode was near neutral in Aug 2024.

Indian Ocean region indices, calculated as the area-averaged monthly mean SSTA (°C) for the SETIO [90°E–110°E, 10°S–0] and WTIO [50°E–70°E, 10°S–10°N] regions, and Dipole Mode Index, defined as differences between WTIO and SETIO. Data are derived from the OIv2.1 SST analysis, and anomalies are departures from the 1991–2020 base period means.





- Strong positive SSTAs were present across much of the tropical Indian Ocean.
- Enhanced convection was observed over Arabian Sea.
- Southwesterly wind anomaly contributed to an above-normal Indian summer monsoon rainfall.

SSTAs (top-left), SSTA tendency (top-right), OLR anomalies (middle-left), sum of net surface short- and long-wave radiation, latent and sensible heat flux anomalies (middle-right), 925-mb wind anomaly vector and its amplitude (bottom-left), 200-mb wind anomaly vector and its amplitude (bottom-right). SST are derived from the Olv2.1 SST analysis, OLR from the NOAA 18 AVHRR IR window channel measurements by NESDIS, winds and surface radiation and heat fluxes from the NCEP CDAS. Anomalies are departures from the 1991-2020 base period means.

Tropical and North Atlantic Ocean

Evolution of Tropical Atlantic SST Indices



Monthly Tropical Atlantic SST Anomaly
(OISSTv2.1, 1991–2020 Climatology)



- Meridional gradient mode index remained well above average since Jun 2024, indicating the presence of a strong positive meridional mode phase.
- ATL3 index was near average in Aug 2024.


Tropical Atlantic Variability region indices, calculated as the area-averaged monthly mean SSTAs (°C) for the TNA [60°W–30°W, 5°N–20°N], TSA [30°W–10°E, 20°S–0] and ATL3 [20°W–0, 2.5°S–2.5°N] regions, and Meridional Gradient Index, defined as differences between TNA and TSA. Data are derived from the OIv2.1 SST analysis, and anomalies are departures from the 1991–2020 base period means.

Impact of Atlantic Meridional Mode (MM) on African climate




- Patterns of SSTA, wind anomaly and OLR anomalies over the past three months are consistent with the impact of positive MM on the African and South American climate.

Tropical Atlantic: SST, SST tend., TCHP, OLR, 200 hPa wind, wind share, heat flex, & RH anom.



Top Row: SSTA (left; OI SST), SSTA tendency (central), Tropical Cyclone Heat Potential anomaly (right; GODAS).
 Middle row: OLR (left; NOAA 18 AVHRR IR), UV200 (central; NCEP CDAS), UV200-UV850 (right; NCEP CDAS) anomalies.
 Bottom row: SW+LW (left), LH+SH (central), Relative humidity at 700 hPa (right; NCEP CDAS) anomalies.
 Anomalies are departures from the 1991-2020 base period means.


Weekly OISSTv2.1 Anom. (°C)
Hatch area: MHW location



- Strong MHWs persisted in the northern tropical Atlantic and the central extratropical Atlantic Ocean.

Weekly SST anomaly (shaded) and locations experience Marine heat waves (hatched) by the date labelled in the plot. MHW is defined as a discrete prolonged warmer than 90th percentile of daily SST for at least 14 days. Data is derived from NCEI OISSTv2.1 and the climatology reference period is 1991-2020


Weekly OISSTv2.1 Anom. (°C)
Hatch area: MHW location



Hurricane Ernesto. Aug 12-Aug 20





Hurricane Debby. Aug 3-Aug 9



Evolution of SST anomaly in the North Atlantic



OISSTv2.1 Anomaly (°C) : JUN-AUG 2024



- SST warming in the hurricane main development region ranked the 2nd warmest Jun-Aug since 1982, nearly matching the record high set in 2023.



Evolution of 200mb-850mb Wind Shear Anomaly

200mb - 850mb Wind Shear Anom. (m/s): JUN-AUG 2024



- Statistically, wind shear tends to enhance (weaken) over the Hurricane Main Development region (MDR, green box) during El Niño (La Niña) events.
- Negative wind shear anomalies dominated in the MDR during Jun-Aug 2024, favouring tropical storm development.

2024 Atlantic Hurricane Season Outlook Update



- NOAA forecasters have increased the likelihood of an above-normal Atlantic hurricane season from **85%** in outlook issued in May to **90%** in August outlook update.

- Main climate factors expected to influence the 2024 Atlantic hurricane activity are :

Continuous of warm phase of AMO

Near historical high record of SST over the MDR

Below normal 200-850hPa wind shear

Enhanced west African monsoon

High chance of La Nina development during peak of the hurricane season

(<https://www.cpc.ncep.noaa.gov/products/outlooks/hurricane.shtml>)

2024 Atlantic Hurricane Season Activities






https://en.wikipedia.org/wiki/2024_Atlantic_hurricane_season

- By Sep 8 2024, five tropical storms formed, with one developing into major hurricane and two into hurricanes.
- No named storm formations since Ernesto on Aug 12.
- Dry Saharan air was considered as an important factor that prevented tropical storm development.

Atlantic	Observations (By Sep 8)	Updated Outlook (Aug) 90% above-normal	Outlook (May 23) 85% above-normal	(1991-2020)
Total storms	5	17-24	17-25	14
Hurricanes	3	8-13	8-13	7
Major hurricanes	1	4-7	4-7	3

NAO and SST Anomaly in North Atlantic




- Positive NAO decreased in Aug 2024, with NAO index=0.7.
- Strong warming continued in the eastern North Atlantic Ocean.
- The prolonged positive SSTAs in the middle latitudes were evident, due to dominance of the positive phase of NAO during the last 5-6 years.


Monthly standardized NAO index (top) derived from monthly standardized 500-mb height anomalies obtained from the NCEP CDAS in 20°N-90°N. Time-latitude section of SSTAs averaged between 80°W and 20°W (bottom). SST are derived from the Olv2.1 SST analysis, and anomalies are departures from the 1991-2020 base period means.

ENSO and Global SST Predictions

Oceanic ENSO Precursors: WWV & CTP





Red box: Warm Water Volume Precursor Blue box: Central Tropical Pacific Precursor



Warm water volume (WWV) is defined as an average of D20 anomaly across the equatorial Pacific (120° E – 80° W, 5° S-5° N) (Meinen and McPhaden 2000). Central tropical Pacific (CTP) index is calculated as the averaged D20 anomaly in the central tropical Pacific (160° W-110° W, 10° S-10° N) (Wen et al. 2014). The monthly D20 data is obtained from the Real-time Ocean Reanalysis Intercomparison Project (https://www.cpc.ncep.noaa.gov/products/GODAS/multiora93_body.html).


DJF Nino34 predictions based on ENSO precursors



- Both WWV and CTP in Aug predict La Niña condition in DJF 2025.

Prediction models are constructed using leave-one-year-out cross validation over the full period by iteratively recomputing the coefficients with the target prediction year removed. For details Wen et al. (2021) DOI: <https://doi.org/10.1175/JCLI-D-20-0648.1>


IRI/CPC Niño3.4 Forecast




- Most of models predict ENSO neutral condition through Sep-Nov, with borderline La Niña conditions developing during Oct-Dec 2024, followed by a transition to neutral conditions during JFM 2025.

- On 9 Aug 2024, CPC maintained a **“La Niña Watch”**.


- Synopsis: **“ENSO-neutral is expected to continue for the next several months, with La Niña favored to emerge during September-November (66% chance) and persist into the Northern Hemisphere winter 2024-25 (74%”**




Official NOAA CPC ENSO Probabilities (issued August 2024)




EC: Niño3.4, IC= 1 Sep 2024



UKMO: Niño3.4, Updated 11 Aug 2024




BOM: Niño3.4, Updated 31 Aug 2024



www.bom.gov.au/climate
Commonwealth of Australia 2024, Australian Bureau of Meteorology


Past analysis base period: 1991-2020
Forecast base period: 1981-2018
Model: ACCESS-S2
Model run: 31 Aug 2024

JMA: Niño3.4, Updated 10 Sep 2024




NMME forecasts from different initial conditions


NMME scaled Nino3.4, IC=202406




NMME scaled Nino3.4, IC=202407




NMME scaled Nino3.4, IC=202408



NMME scaled Nino3.4, IC=202409



NINO34 in La Nina Years following El Nino conditions




Following strong El Nino events: 1983, 1998, 2016

Following moderate El Nino events: 1988, 2007, 2010


Following weak El Nino events: 1995, 2005, 2020

Evolution of Monthly Mean Zonal Wind Stress Anomaly across [5S-5N]



Evolution of Monthly Mean D20 Anomaly across [5S-5N]

Monthly Mean D20(shaded) & SST(contour) Anomaly (5S-5N: 1991-2020 Climatology)




NMME Forecasts in the Indian Ocean


NMME Sea Surface Temperature Anomalies (DecC)

Oct2024–Dec2024

September2024 initial conditions




NMME IOD fcast, IC=202409




- NMME forecasts indicate that IOD is likely to remain neutral through spring 2025.

NMME Forecasts in the Atlantic Ocean


NMME Sea Surface Temperature Anomalies (DecC)




NMME MDR[90W–18W,9N–21N] fcst, IC=202409



NMME MGI(TNA–TSA) fcst, IC=202409



standardized PDO index



- CFSv2 predicts the negative phase of PDO will continue through winter 2024.

CFS Pacific Decadal Oscillation (PDO) index predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1991-2020 base period means. PDO is the first EOF of monthly ERSSTv3b anomaly in the region of [110°E-100°W, 20°N-60°N]. CFS PDO index is the standardized projection of CFS SST forecast anomalies onto the PDO EOF pattern.

Acknowledgement

- ❖ Drs. Arun Kumar, Zeng-Zhen Hu and Jieshun Zhu: reviewed PPT, and gave insightful suggestions and comments
- ❖ Drs. Yanjuan Guo and Pingping Xie provided the BASS/CMORPH/CFSR EVAP package
- ❖ Drs. Jieshun Zhu & Wanqiu Wang maintained the sea ice forecasts

Please send your comments and suggestions to:

Arun.Kumar@noaa.gov

Caihong.Wen@noaa.gov

Jieshun.Zhu@noaa.gov

Zeng-Zhen.Hu@noaa.gov

- **NCEP/CPC Ocean Monitoring & Briefing Operation (Hu et al., 2022, BAMS)**
- **Weekly Optimal Interpolation SST (OIv2.1 SST; Huang et al. 2021)**
- **Extended Reconstructed SST (ERSST) v5 (Huang et al. 2017)**
- **Blended Analysis of Surface Salinity (BASS) (Xie et al. 2014)**
- **CMORPH precipitation (Xie et al. 2017)**
- **CFSR evaporation adjusted to OAFlux (Xie and Ren 2018)**
- **NCEP CDAS winds, surface radiation and heat fluxes (Kalnay et al. 1996)**
- **NESDIS Outgoing Long-wave Radiation (Liebmann and Smith 1996)**
- **NCEP's GODAS temperature, heat content, currents (Behringer and Xue 2004)**
- **Aviso altimetry sea surface height from CMEMS**
- **Ocean Surface Current Analyses – Realtime (OSCAR)**
- **In situ data objective analyses (IPRC, Scripps, EN4.2.1, PMEL TAO)**
- **Operational Ocean Reanalysis Intercomparison Project**
http://www.cpc.ncep.noaa.gov/products/GODAS/multiora_body.html
http://www.cpc.ncep.noaa.gov/products/GODAS/multiora93_body.html

Backup Slides

Global Sea Surface Salinity (SSS): Anomaly for August 2024

New Update: The NCEI SST data used in the quality control procedure has been updated to version 2.1 since August 2020;


Strong positive SSS anomalies are found over certain mid-latitude regions of the Pacific and Atlantic oceans, while strong negative anomalies in some higher latitudes of Pacific and Atlantic as well as northern Indian ocean. The SSS anomaly patterns are overall consistent with the E-P flux anomaly patterns over the Pacific and Indian Oceans, which is more dominated by the precipitation anomalies over these regions. In the meantime, the consistency between the SSS and E-P flux in the southern Atlantic and Indian Oceans is less obvious.

**SSS : Blended Analysis of Surface Salinity (BASS) V0.2
(a CPC-NESDIS/NODC-NESDIS/STAR joint effort)**

<ftp.cpc.ncep.noaa.gov/precip/BASS>


Precipitation: CMORPH adjusted satellite precipitation estimates

Evaporation: Adjusted CFS Reanalysis



Global Sea Surface Salinity (SSS): Tendency for August 2024


The global SSS tendency in August 2024 is overall weak and with mixed signs. The Precipitation are generally decreasing along the equator but enhancing off equator in the Pacific, whereas strong precipitation increase is found in the equatorial Indian ocean but decrease north of the equator. The E-P flux patterns seem to be more dominated by the precipitation patterns. Overall, the SSS trend doesn't show pronounced large scale pattern for this month.



Monthly SSS Anomaly Evolution over Equatorial Pacific

NOTE: Since August 2015, the BASS SSS is from in situ, SMOS and SMAP; before August 2015, The BASS SSS is from in situ, SMOS and Aquarius.


- Hovermoller diagram for equatorial SSS anomaly (**5S-5°N**);
- Decreased SSS is found over the warm pool region, but increased over the central Pacific. SSS increasing is also found over the equatorial eastern Pacific.



Pentad SSS Anomaly Evolution over Equatorial Pacific


Figure caption:

Hovermoller diagram for equatorial (5°S - 5°N) 5-day mean SSS, SST and precipitation anomalies. The climatology for SSS is Levitus 1994 climatology. The SST data used here is the OISST V2 AVHRR only daily dataset with its climatology being calculated from 1985 to 2010. The precipitation data used here is the adjusted CMORPH dataset with its climatology being calculated from 1999 to 2013.



Equatorial Pacific SST ($^{\circ}\text{C}$), HC300 ($^{\circ}\text{C}$), u850 (m/s) Anomalies



2 $^{\circ}\text{S}$ –2 $^{\circ}\text{N}$ Average, 3 Pentad Running Mean



- Negative SSTA in the eastern Pacific enhanced slightly in Jul 2024, while positive SSTA persisted west of 150 $^{\circ}\text{W}$.
- Negative H300 anomaly increased slightly in the eastern Pacific during Aug 2024.
- Easterly wind anomalies persisted over much of central-eastern equatorial Pacific during Aug 2024, contributing to re-strengthening of subsurface cooling in the eastern Pacific.

Evolution of Pacific Niño SST Indices







Monthly Tropical Pacific SST Anomaly



- Relative Niño3.4 index is now included in ENSO monitoring, which is defined as the conventional Niño3.4 index minus the SSTA averaged in the whole tropics (0° - 360° , 20° S- 20° N), in order to remove the global warming signal. Also, to have the same variability as the conventional Niño3.4 index, the relative Niño3.4 index is renormalized (van Oldenborgh et al. 2021: ERL, 10.1088/1748-9326/abe9ed).

[Relative Niño3.4 data updated monthly at: https://www.cpc.ncep.noaa.gov/data/indices/RONI.ascii.txt](https://www.cpc.ncep.noaa.gov/data/indices/RONI.ascii.txt)


North Pacific & Arctic Ocean: SSTA, SSTA Tend., OLR, SLP, Sfc Rad, Sfc Flx Anomalies




SSTA (top-left; Olv2.1 SST Analysis), SSTA tendency (top-right), Outgoing Long-wave Radiation (OLR) (middle-left; NOAA 18 AVHRR IR), sea surface pressure (middle-right; NCEP CDAS), sum of net surface short- and long-wave radiation (bottom-left; positive means heat into the ocean; NCEP CDAS), sum of latent and sensible heat flux (bottom-right; positive means heat into the ocean; NCEP CDAS). Anomalies are departures from the 1991-2020 base period means.

North America Western Coastal Upwelling

Pentad Coastal Upwelling for West Coast North America
($\text{m}^3/\text{s}/100\text{m}$ coastline)



Standard Positions of Upwelling Index Calculations




- Both anomalous upwelling and downwelling were observed in Aug 2024.

(top) Total and (bottom) anomalous upwelling indices at the 15 standard locations for the western coast of North America. Derived from the vertical velocity of the NCEP's GODAS and are calculated as integrated vertical volume transport at 50-meter depth from each location to its nearest coast point ($\text{m}^3/\text{s}/100\text{m}$ coastline). Anomalies are departures from the 1991-2020 base period pentad means.

- Area below (above) black line indicates climatological upwelling (downwelling) season.

- Climatologically upwelling season progresses from March to July along the west coast of North America from 36°N to 57°N.

SST Configurations




- **Full2023**: 2023 Jan-Nov SST forcing
- **2023ATLONLY**: 2023 Atlantic SSTs only and climatology elsewhere
- **ELNINOPAC**: 2023 SSTs with Pacific SSTs swapped for El Niño composite (1982, 1987, 1997, 2002, 2015)
- **ATLCLIM**: Atlantic SSTs set to climatology with 2023 SSTs elsewhere







- Extremely warm Atlantic was the primary driver of the reduced vertical wind shear over the hurricane main development region.

- The unique spatial pattern of 2023 El Niño contributes to increased levels of vertical wind shear than historical moderate/strong El Niño composite.


ASO Vertical Wind Shear Anomalies



Klotzbach et al, 2024: The 2023 Atlantic Hurricane Season: An Above-Normal Season Despite Strong El Niño Conditions Bull. Amer. Meteor. Soc. <https://doi.org/10.1175/BAMS-D-23-0305.1>




NINO3.4 SST anomalies (K)



- The latest CFSv2 forecasts an La Niña will develop in Sep 2024.

CFS Niño3.4 SST prediction from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1991-2020 base period means.


Indian Ocean Dipole SST anomalies (K)



- CFSv2 predicts a positive phase of IOD in the 2nd half of 2023.

CFS Dipole Model Index (DMI) SST predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). The hindcast climatology for 1981-2006 was removed, and replaced by corresponding observation climatology for the same period. Anomalies were computed with respect to the 1991-2020 base period means.

Tropical N. Atlantic SST anomalies (K)






- Latest CFSv2 predictions call for above-normal SST in the tropical North Atlantic.
- CFSv2 has large cold bias at 0-month lead since Feb 2024.


CFS Tropical North Atlantic (TNA) SST predictions from the latest 9 initial months. Displayed are 40 forecast members (brown) made four times per day initialized from the last 10 days of the initial month (labelled as IC=MonthYear) as well as ensemble mean (blue) and observations (black). Anomalies were computed with respect to the 1991-2020 base period means. TNA is the SST anomaly averaged in the region of [60°W-30°W, 5°N-20°N].

Monthly SST Anomaly in the Atlantic Ocean

AUG 2024 Monthly SST Anomaly (°C)



Tropical N. Atlantic SSTA (°C) [60W-30W, 5N-20N]



- NSST provide SST nudging source for CFSR.
- NSST was cooler than other SST analysis in most of tropical Atlantic Ocean.
- NSST anomaly in the northern tropical Atlantic Ocean region was colder than OISST v2.1 anomaly by 0.5C in Aug 2024.
- NSST cold bias at least partially contributed to cold bias in CFSv2 short-lead forecasts.