




Food and Agriculture
Organization of the
United Nations

Global Soil Salinity Map

GSS map

LESSON 2

Software installation
and data organization



Global Soil Salinity Map

GSS map

LESSON 2

Software installation and data organization

Christian Thine Omuto
Ronald Vargas
Kostiantyn Viatkin
Yusuf Yigini

Required citation:

Omuto, C.T., Vargas, R., Viatkin, K. and Yigini, Y. 2020. *Global Soil Salinity Map – GSSmap. Lesson 2 – Software installation and data organization*. Rome, FAO.

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO.

© FAO, 2020



Some rights reserved. This work is made available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo/legalcode>).

Under the terms of this licence, this work may be copied, redistributed and adapted for non-commercial purposes, provided that the work is appropriately cited. In any use of this work, there should be no suggestion that FAO endorses any specific organization, products or services. The use of the FAO logo is not permitted. If the work is adapted, then it must be licensed under the same or equivalent Creative Commons licence. If a translation of this work is created, it must include the following disclaimer along with the required citation: "This translation was not created by the Food and Agriculture Organization of the United Nations (FAO). FAO is not responsible for the content or accuracy of this translation. The original [Language] edition shall be the authoritative edition."

Disputes arising under the licence that cannot be settled amicably will be resolved by mediation and arbitration as described in Article 8 of the licence except as otherwise provided herein. The applicable mediation rules will be the mediation rules of the World Intellectual Property Organization <http://www.wipo.int/amc/en/mediation/rules> and any arbitration will be conducted in accordance with the Arbitration Rules of the United Nations Commission on International Trade Law (UNCITRAL).

Third-party materials. Users wishing to reuse material from this work that is attributed to a third party, such as tables, figures or images, are responsible for determining whether permission is needed for that reuse and for obtaining permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

Sales, rights and licensing. FAO information products are available on the FAO website (www.fao.org/publications) and can be purchased through publications-sales@fao.org. Requests for commercial use should be submitted via: www.fao.org/contact-us/licence-request. Queries regarding rights and licensing should be submitted to: copyright@fao.org.

Contents

Figures	III
Summary	V
1 Introduction	1
1.1 Overview	1
1.2 Objective	1
1.3 Expected outcomes	1
2 GIS database for assessing salt-affected soils	1
2.1 Software for GIS database organization	1
2.1.1 Applications in QGIS	1
3 Resources	2
4 Activities	2
4.1 Software installation	2
4.1.1 Installing QGIS	2
4.1.2 Installing ILWIS	2
4.1.3 Installing R	3
4.1.4 Installing RStudio	4
4.1.5 Installing R package	4
4.2 Database organization	4
4.2.1 Organizing spreadsheet data	4
4.2.2 Organizing GIS data	5
5 Outputs	9

Figures	
Figure 4.1: Input spreadsheet data format.....	5
Figure 4.2: Reprojecting GIS layers in QGIS	6
Figure 4.3: Steps for vector-to-raster conversion in QGIS	7
Figure 4.4: Image processing using semi-automatic classification plugin	8
Figure 5.9: Steps for digital terrain analysis in SAGA.....	8
Figure 4.6: Creating spatially harmonized GIS database in ILWIS.....	9

Summary

This Lesson is the second step of the capacity-building program, which is designed to build national capacities as well as harmonize procedures for developing information of salt-affected soils at the national and global levels. The overall goal of this Lesson is to support participants in software installation and using the software to organize input data for assessing salt-affected soils at the national level. At the end of the lesson, the participants are expected to have installed software and organized database for mapping salt-affected soils in their countries.

Summary outputs for organizing input national information for assessing salt-affected soils

Item	Description		Timeline
Software	Installed software for database organization		One week
	Spreadsheet software	MS Excel	
	GIS database software	QGIS and SAGA	
	Data harmonization software	ILWIS and R	
Data	Organized input data		
	Soil indicators of salt problems	EC, ESP, pH	
	GIS layers	Boundary shapefile	
		Spatial layers	



1 Introduction

1.1 Overview

Proper data organization is an important pedometrics step in digital soil mapping and more so in identification of the true extent of salt-affected soils. Most soil profile data collection are organized in varying soil-depth steps and irregular soilscape patterns. Adequate understanding and modelling of soil variations are enhanced with well-organized input database. Furthermore, properly organized data minimizes syntax errors at the data analysis stage. The focus of this Lesson is to facilitate participants in harmonized organization of input database for assessing salt-affected soils. The input database to be organized was developed during Lesson one of the Capacity Building program. This Lesson, therefore, requires outputs from Lesson one. The Lesson is intended for participants who have completed Lesson one or participants with complete database for mapping salt-affected soils. Its output is a spatially compatible GIS database for national assessment of salt problems.

1.2 Objective

The overall objective of this Lesson is to establish a spatially compatible GIS database for national assessment of salt-affected soils in each member country.

1.3 Expected outcomes

By the end of this Lesson, the participants are expected to:

- i. Install and operationalize GIS database software for mapping salt-affected soils
- ii. Spatially harmonize input data into a GIS database for mapping salt-affected soils

2 GIS database for assessing salt-affected soils

2.1 Software for GIS database organization

Software is part of the tools and methods for digital soil mapping framework. The following software were downloaded during Lesson one: QGIS, ILWIS, and R. This section discusses some other potential applications of these software in database organization.

2.1.1 Applications in QGIS

QGIS is a versatile and relatively fast software with many GIS functionalities such as GIS database development and organization, GIS operations (layer trimming, projections, format conversion, views, etc.), remote sensing applications (such as image download, correction, pre-processing, classification, band analysis, etc.), map layouts, among others. Its documentation and tutorials for beginners are freely downloadable at <https://qgis.org/en/site/forusers/download.html> (accessed on 30th October 2019). The software also has a rich contribution of freely downloadable add-on packages known as plugins and a vibrant user-support (for example, through <https://gis.stackexchange.com/>).

QGIS has a Graphical User Interface (GUI) for easy access to its functions. The functions are easily accessible through panels or shortcut icons by simple clicks/selection.

3 Resources

The following resources are useful for implementing the activities during data organization

- (a) Software (QGIS, ILWIS, Spreadsheet software)
- (b) Own dataset (soil profile and GIS layers)
- (c) Lecture materials

4 Activities

4.1 Software installation

4.1.1 Installing QGIS

Step 1: Go to the site <https://qgis.org/en/site/forusers/download.html> and download the latest release depending on the computer OS. For example, for Windows 64 bit, download QGIS Standalone Installer Version 3.10 (64 bit) will be as shown in Figure 4.1.




Figure 4.1: Screenshot of site for downloading QGIS (source: <https://qgis.org/en/site/forusers/download.html>)

Step 2: Launch the installer

Step 3: The software (QGIS Desktop with GRASS) is available in a QGIS folder

4.1.2 Installing ILWIS

Step 1: Go to the site <https://www.itc.nl/ilwis/download/ilwis33/> and download ILWIS3.3Academic.exe (19.3 MB) and ILWIS 3.31 Update (for ILWIS 3.3 only) (Figure 4.2)

Step 2) Install ILWIS3.3Academic.exe (19.3 MB) first and do not start the software yet.

Step 3) Install ILWIS 3.31 Update (for ILWIS 3.3 only). During installation, it will request to locate ILWIS 3.3. Navigate to C:\Program Files (x86)\ILWIS 3.3 Academic and scroll down to choose the blue colored file Ilwis30.exe. Complete the installation

Step 4) Launch the software (whose icon should now be on the desktop).




Figure 4.2: Screenshot site for downloading ILWIS (source: <https://www.itc.nl/ilwis/download/ilwis33/>)

4.1.3 Installing R

Step 1: Go to the site <https://cran.r-project.org/> and download the installer suitable for your operating system (Figure 4.3). For Windows, choosing Download R for Windows takes you to another site from which you choose *Install R for the first time*. Another window for download opens.




Figure 4.3: Screenshot of site for download R (source: <https://cran.r-project.org/>)

Step 2: Install the window installer, choose the language (preferably English), bit rate for your computer, and choose *No (Accept default)* for the start-up options.

4.1.4 Installing RStudio

Step 1: Go to the site <https://www.rstudio.com/>, click download RStudio, the preferred version (like Free RStudio Desktop) and download.

Step 2. Launch the installer and install the software

4.1.5 Installing R package

Step 1: Start RStudio and click the Packages Icon in the bottom right corner, the click *Install*. A window for install Packages window pops up (Figure 4.4).

Step 2: If connected to internet, type the name of the package to install (for example sp)

Step 3: Click *Install* button and the package will be installed and ready for use




Figure 4.4 Screenshot on package installation from RStudio application

4.2 Database organization

4.2.1 Organizing spreadsheet data

Spreadsheet data should be organized to allow sequential computer scanning during data analysis.

1. Let the first column be the variable for indexing the serial order of the data.
2. Other variables in the database are consecutively ordered beginning with profile ID, Location description (if any), Latitude and Longitude, and sampling depth-range (Figure 4.5).
3. The depth-range is split into two variables: *Upper* and *Lower*. *Upper* denotes the first part of the depth-range while *Lower* denotes the last part of the depth-range. The value of *Upper* in one sample should be the same as the value for *Lower* in the preceding sample for the same profile ID (Figure 4.5).

- Other variables such as Depth Code (or Horizon number), soil texture components, organic carbon, pH, EC, ESP, and additional soil properties such soluble ions (and/or exchangeable sodium ions and CEC) are also included in that order (Figure 4.5). The Depth Code (or Horizon) should have consecutive numbers beginning from one to the last sampled depth/horizon in each profile. This arrangement creates repeated numbers/codes of the profile ID and Longitude and Latitude values (Figure 4.5).

Sample	Pits	Longitude	Latitude	Depth	Upper	Lower	Horizon	Clay	pH	EC	SAR	ESP	SolCa	SolMg	SolNa	SO4	SolCl
1	1	45.13	31.93	0-10	0	10	1	9	7.7	0.6	3	3	3.1	1.5	2.5	0	2.8
2	1	45.13	31.93	10-30	10	30	2	12	7.6	4	11	12	28	8.5	12	1.9	5.8
3	1	45.13	31.93	30-60	30	60	3	22	7.5	0.5	3	4	2.8	1.5	2		
4	1	45.13	31.93	60-100	60	100	4										
5	2	56.65	23.89	0-10	0	10	1	27	8.6	1.9	9	10	14.1	3	4.5		
6	2	56.65	23.89	10-30	10	30	2	31	7.8	0.7	4	5	4.5	1.5	2.5		
7	2	56.65	23.89	30-100	30	100	3										
8	3	55.74	39.45	0-35	0	35	1	22	7.6	0.9	5	5	6.2	2	2		
9	3	55.74	39.45	35-60	35	60	2	25	7.8	0.4	2	2	2.2	1.5	2		
10	3	55.74	39.45	60-100	60	100	3	32	7.9	0.4	2	2	2.1	1.5	2		
11	4	62.56	34.15	0-20	0	20	1	7	7.5	0.3	2	2	1.3	1	1.5		
12	4	62.56	34.15	20-60	20	60	2	10	7.7	0.2	1	1	0.7	1	1.5		
13	4	62.56	34.15	60-100	60	100	3	23	7.9	0.3	2	3	1.7	1	1.5		
14	5	70.62	-71.15	0-20	0	20	1	10	7.8	2.2	9	9	15.8	4	6		
15	5	70.62	71.15	20-50	20	50	2	12	7.7	1.6	7	6	11.4	3	5	0.1	14.2
16	5	70.62	71.15	50-100	50	100	3										
17	6	64.56	28.92	0-15	0	15	1	22	7.4	2.7	10	10	16.3	3.5	5.5		
18	6	64.56	28.92	15-30	15	30	2	25	7.5	1.2	6	7	8.3	2.5	4		
19	7	34.47	52.44	0-20	0	20	1	21	7.8	10.7	14	14	63.1	30.5	41		
20	7	34.47	52.44	20-50	20	50	2	23	7.5	2.2	9	9	14.8	4	6		
21	8	69.98	42.14	0-30	0	30	1	22	7.9	0.4	2	3	2.3	1.5	2		
22	8	69.98	42.14	30-70	30	70	2	25	7.7	0.9	6	7	6.4	1.5	2.5		
23	8	69.98	42.14	70-100	70	100	3	24	7.5	3.9	10	10	26.1	7.5	13		

Figure 4.5: Input spreadsheet data format

It is important to ensure that:

- Upper, Lower, and Horizon of each profile ID increase down the soil profile
- Latitude, Longitude and Profile ID remain constant for each Profile ID
- Upper is equivalent to the first part of the Depth range and Lower is the second part of Depth

4.2.2 Organizing GIS data

(a) Harmonizing coordinate reference system

Harmonization of coordinate reference system (CRS) is done by Reprojection, which is the terminology often used in GIS for transforming one CRS to another. The following points need consideration when *reprojecting* GIS layers:

- Identifying the CRS to use on all GIS layers (here known as harmonized CRS)
- It is important to choose UTM projection for harmonizing all layers CRS
- CRS of each GIS layer and individual need for *reprojection* should be noted before *reprojection*
- Reprojecting* vector layers requires specifying the CRS when the layers are saved anew
- When *reprojecting* raster layers, *source* CRS (current) and *target* CRS need to be specified

The steps for *reprojecting* vector and raster layers is given in Figure 4.6.





Figure 4.6: Reprojecting GIS layers in QGIS

(b) Harmonizing layer format

Not all GIS layers are available in uniform file format. Sometimes, some soil forming factors such as soil map, geology, and land cover types are only available as polygon vector files. Harmonizing GIS layer formats attempts to convert vector files to raster formats. Except for soil profile data, all GIS layers for mapping salt-affected soils need to be raster layers. Vector-to-raster conversion is a GIS operation used for harmonizing vector polygons into raster data types. Some of the pre-requisite operations before vector-to-raster harmonization include projection harmonization, identification of the target polygon attribute to use in the conversion, decision on the target pixel resolution of the final raster map, and cleaning of missing entries, topology errors, and associated errors in the data entry. Figure 4.7 illustrates the steps for vector-to-raster conversion in QGIS.



Steps for Vector-to-raster conversion

1. Select the vector layer to rasterize
2. Right-click and open attribute table
3. Inspect the attribute for rasterizing
4. An integer attribute may be created at this stage if needed
5. Activate the Processing Toolbox
6. Type *rasterize* in the search window
7. Select *Rasterize* function under SAGA
8. A *Rasterize* window will appear when the SAGA function is activated
9. Choose the vector file in the scroll-down under *Shapes* option
10. Select the attribute for rasterizing
11. Change the cellsize to the desired value. Here remember to use suitable values depending on the harmonized CRS
12. Click *Run* to launch the process
13. A *Rasterize* layer will be added into the View after successful process
14. Save the output as geoTIFF


Figure 4.7: Steps for vector-to-raster conversion in QGIS

(c) Image correction

Freely downloadable remote sensing images have varied characteristics that need harmonization before application in spatial modelling of salt-affected soils. These characteristics range from differences in spatial extent, image digital numbers, image conversion models to land surface reflectance, and downloaded file format. Most downloaded images are wrapped in compressed files such as .zip, .rar or .gz. They should be unpacked into commonly used GIS file formats such as geoTIFF, ASCII, HDF, etc. Software such as WinZip or WinRar or 7-Zip unpacks the compressed files into a preferred destination folder (such as C:/Salinity/Output). Sometimes the unpacked file may still contain compressed files and another unpacking may be necessary to extract the GIS files. It is important that different zipped files are unzipped into different folders to avoid over-writing metafiles and for processing each file independently. Semi-Automatic Classification Plugin (SCP) in QGIS provides a quick way for simultaneous geometric and radiometric correction of remote sensing images. The plugin has many functions for handling different types of remote sensing images such as Landsat, Sentinel-2, Sentinel-3, ASTER, MODIS, etc. Figure 4.4 illustrates how these types of images are corrected using SAC plugin in QGIS.

(d) Digital terrain modelling


Terrain modelling focuses on determining relief parameters influencing soil formation or translation of soil salts during leaching, runoff or wind erosion. Relief parameters are primary or secondary attributes. Primary attributes are derived from the altitude and directional location of the topographic surface. They include slope, aspect, curvature, flow accumulation and up-slope flow contributing area. Secondary attributes are obtained from the altitude and derivatives of the primary attributes. They include indices such as topographic wetness index, compound topographic index, stream power index, etc. SAGA software has a module for developing 14 terrain parameters from DEM input (Figure 4.8)



Steps for image correction using SAC

1. Select and set the working CRS for the QGIS window. Harmonized CRS is preferred
2. Launch the plugin
3. Select *Preprocessing* button
4. Select the image type (say Landsat) in front of the *Directory containing landsat bands*: Select the folder containing the unpacked Landsat images. Each folder is processed at a time
5. Select the MTL file, which is a textfile with MTL at the end of the file name. If the image type is Sentinel-2, the textfile name ends with MSI
6. Click *Run* to execute the command. A prompt for choosing the output folder comes up. The folder with the downloaded image (step 4 above) is preferred
7. If the image type is MODIS, the no need for the textfile since the information is already contained in the header-file.
8. The corrected image images will be loaded into the QGIS view window.
9. Repeat the steps for all downloaded image scenes

Figure 4.8: Image processing using semi-automatic classification plugin



Steps for deriving terrain parameters

1. Import the DEM: *Geoprocessing* then *File* then *Grid* then *Import* then *Import Raster*
2. Launch the pre-processing (fill sinks if necessary): *Geoprocessing*, *Terrain Analysis*, *Preprocessing*, *Fill Sinks* (and use option of choice, (say the 1st one))
3. Launch Basic Terrain Analysis: *Geoprocessing*, *Terrain Analysis*, *Basic Terrain Analysis*
4. Basic Terrain Analysis window opens
5. On *Grid system*: scroll to select the spatial characteristic of the loaded DEM
6. On *Elevation*: Scroll to choose the appropriate DEM
7. Click *Okay* to run the process.
8. Export the layers: *Geoprocessing*, *File*, *Grid* then *Export Raster*

Figure 4.9: Steps for digital terrain analysis in SAGA

(e) Final GIS database

All spatial data in the GIS database should have uniform CRS (i.e. harmonized CRS). In addition, all raster layers should have uniform pixel resolution. Uniform pixel resolution is obtained by resampling all the raster layers to the preferred spatial resolution. A map-list in ILWIS is one way of ensuring that all harmonized raster layers have uniform CRS and pixel resolution. Figure 4.10 gives the steps for creating a map-list of harmonized raster layers




Figure 4.9: Creating spatially harmonized GIS database in ILWIS

5 Outputs

Each participant is expected to produce the following at the end of this lesson:

1. Organized database of indicators of salt-affected soils (EC, pH, ESP)
2. Organized database of GIS layers (land cover, climate, geology, images, DEM, soil map, etc.)
3. Installed computer software for mapping salt-affected soils



Thanks to the financial support of



Ministry of Finance of the
Russian Federation



European
Commission



Rural Development
Administration

