

Efficient micromirror confinement of sub-teraelectronvolt cosmic rays in galaxy clusters

Received: 2 November 2023

Accepted: 8 November 2024

Published online: 03 January 2025

 Check for updates

Patrick Reichherzer ^{1,2}✉, Archie F. A. Bott^{1,3}, Robert J. Ewart ^{1,4},
Gianluca Gregori ^{1,5}, Philipp Kempfski⁶, Matthew W. Kunz ^{6,7} &
Alexander A. Schekochihin^{1,8}

Cosmic rays (CRs) play a pivotal role in shaping the thermal and dynamical properties of astrophysical environments, such as galaxies and galaxy clusters. Recent observations suggest a stronger confinement of CRs in certain astrophysical systems than predicted by current CR-transport theories. Here, we show that the incorporation of microscale physics into CR-transport models can account for this enhanced CR confinement. We develop a theoretical description of the effect of magnetic microscale fluctuations originating from the mirror instability on macroscopic CR diffusion. We confirm our theory with large-dynamical-range simulations of CR transport in the intracluster medium (ICM) of galaxy clusters and kinetic simulations of CR transport in micromirror fields. We conclude that sub-teraelectronvolt CR confinement in the ICM is far more effective than previously anticipated on the basis of Galactic-transport extrapolations. The transformative impact of micromirrors on CR diffusion provides insights into how microphysics can reciprocally affect macroscopic dynamics and observable structures across a range of astrophysical scales.

A good theory of cosmic-ray (CR) transport is crucial for advancing our understanding of phenomena in the Universe, including the formation and evolution of galaxies and galaxy clusters. CRs, through their transport characteristics, not only influence their environment but also modulate their own (re)acceleration and confinement efficiency as well as the observable photon and neutrino emission.

The transport characteristics of CRs in magnetic-field structures depend on the scattering efficiency and mechanism, both of which are influenced by the properties of the ambient plasma. Specifically, within a weakly collisional, high- β plasma, that is, one in which the thermal pressure greatly exceeds the magnetic pressure, deviations from local thermodynamic equilibrium provide free energy for fast-growing Larmor-scale instabilities, leading to distortions in magnetic fields on thermal-ion kinetic scales. In such a high- β plasma, two characteristic

scales are relevant for describing the global transport of CRs: the macroscale of the magnetic turbulence, characterized by the correlation length l_c or the ‘Alfvén scale’ l_A , and the microscale l_{mm} of the micromirrors created by the mirror instability^{1,2}. The prefix ‘micro’ refers to scales much smaller than the $\approx \mu\text{pc}$ gyroradii of ≥ 100 MeV CRs, and serves to distinguish the plasma-kinetic-scale ‘micromirrors’ (denoted by subscript mm) from the large-scale magnetic mirrors that also influence CR transport^{3–5}. While there are other micro-instabilities, the magnetic fluctuations created by the mirror instability are stronger and thus more influential for CR transport^{6,7}. The physics associated with these micro- and macroscales introduces three distinct transport regimes, which depend on the CR energy.

First, in the high-energy (subscript he) limit, CRs with gyroradii $r_g \gg l_c$ (energies $E \geq 100$ EeV for typical turbulence-driving scales

¹Department of Physics, University of Oxford, Oxford, UK. ²Exeter College, Oxford, UK. ³Trinity College, Oxford, UK. ⁴Balliol College, Oxford, UK.

⁵Lady Margaret Hall, Oxford, UK. ⁶Department of Astrophysical Sciences, Princeton University, Princeton, NJ, USA. ⁷Princeton Plasma Physics Laboratory, Princeton University, Princeton, NJ, USA. ⁸Merton College, Oxford, UK. ✉e-mail: patrick.reichherzer@physics.ox.ac.uk




Fig. 1 | Micromirror field generated by the PIC simulation. Colour shows fluctuations δB_{\parallel} along the field B , which is aligned with the x axis. We show two snapshots of the 3D field during its secular evolution, characterized by different $\delta B/B$, as indicated in the plots (the right-hand snapshot is later in the evolution).

in galaxy clusters) undergo scattering by small angles of order $\delta\Theta \approx l_c/r_g$ at characteristic times $\delta t \approx l_c/c$, leading to a scattering rate $\nu_{\text{he}} \approx \delta\Theta^2/\delta t \approx c l_c/r_g^2$ and, therefore, to a CR diffusion coefficient $\kappa_{\text{he}} \approx c^2/\nu_{\text{he}} \approx cr_g^2/l_c \propto E^2/l_c$ (see, for example, ref. 8). This scaling is indeed observed in both numerical simulations (for example, refs. 9–11) and (scaled) laboratory experiments¹² and serves as input for propagation models of ultrahigh-energy CRs in galaxy clusters¹³.

As the high-energy regime is, thus, believed to be understood, recent studies of CR transport have predominantly focused on the second, mesoscale regime, $r_g \lesssim l_c$, in which CRs scatter resonantly off inertial-range turbulent structures^{14–17}. This second regime is probably relevant for explaining CR spectra¹⁸; for example, the steepening of the CR spectrum in the Galactic Centre from gigaelectronvolt to petaelectronvolt energies^{11,19–21}.

We argue, and confirm numerically, that in high- β plasmas the presence of microstructures caused by plasma instabilities introduces a third regime, whose physics is similar to that of the first, but with the micromirror scale l_{mm} playing the role of l_c and the requirement that $r_g \gg l_{\text{mm}}$ ($E \gg 100$ MeV). We show that this microscale physics largely overrides the mesoscale resonant scattering and streaming. We apply our theory to the intracluster medium (ICM), a representative high- β plasma, and determine the transition between micro- and macrophysics-dominated transport to be at teraelectronvolt energies, only weakly influenced by mesoscale physics. We confirm this theory with a novel method (Methods) that incorporates the microscales ($l_{\text{mm}} \approx 100$ npc), the macroscales ($l_c \approx 100$ kpc) and the vast range in between.

Results

Effect of micromirrors on large-scale CR transport

It has long been realized that plasma instabilities may dominate the transport of low-energy CRs. However, the instabilities most often highlighted in the literature arise from the CRs themselves, rather than from the thermal plasma. One prominent example, especially for Galactic transport of CRs below 100 GeV, is the streaming instability²². This instability generates fluctuations in the magnetic field that in turn scatter the CRs and thereby reduce their streaming velocity to be comparable to the Alfvén speed in the plasma^{22–25}. In high- β , weakly collisional plasmas, there also exist a variety of instabilities that are driven by pressure anisotropies and generate magnetic fluctuations on ion Larmor scales (see ref. 2, and references therein). The pressure anisotropies arise from the (approximate) conservation of particles' adiabatic invariants during the local stretching and compression of magnetic fields²⁶. In the present context, the mirror instability^{27,28} is of particular interest because its saturated amplitude $\delta B_{\text{mm}} \approx B/3$ is of the same order of magnitude as the ambient magnetic field B (refs. 6,29,30). The other well-known instability arising in such plasmas, the firehose instability, is unlikely to affect CR transport because its expected saturation amplitude is small

under ICM conditions: $\delta B_{\text{f}} \approx (\tau\Omega_i)^{-1/4}B$ (refs. 6,7), where τ is the timescale over which a firehose-susceptible plasma evolves macroscopically, and $\Omega_i \approx 0.01(B/3 \mu\text{G}) \text{ s}^{-1}$ is the non-relativistic thermal-ion gyrofrequency. In the ICM, $\tau\Omega_i \approx 10^{11}$ (ref. 31), so $\delta B_{\text{f}} \approx 10^{-3}B$. By analogy to equation (3), it follows that the scattering rate ν_{f} of CRs off firehoses in the ICM is much smaller than that off the mirrors: $\nu_{\text{f}}/\nu_{\text{mm}} \approx 10^{-7}$, with the firehose scale taken to be comparable to the thermal-ion gyroradius^{6,7}.

To determine the impact of these micromirrors on CR transport, we begin by working out the relevant theoretical predictions for diffusion coefficients of CRs scattering at such strong fluctuations. Note that a previous assumption of weaker micromirror fluctuations led to a different, much larger diffusion coefficient based on calculations using quasilinear theory³². The velocity change $\delta\mathbf{v}$ of a relativistic CR with gamma factor $\gamma = (1 - v^2/c^2)^{-1/2}$, charge $q = Ze$ and mass m in a magnetic-field structure of scale $l_{\text{mm}} \ll r_g$ and vector amplitude $\delta\mathbf{B}_{\text{mm}}$ is given in the small-angle limit $|\delta\mathbf{v}| \ll v$ by integrating the equation of motion $\gamma m \, d\mathbf{v}/dt = q(\mathbf{v}/c) \times \delta\mathbf{B}_{\text{mm}}$ along the CR path:

$$\delta\mathbf{v} \approx \frac{q}{\gamma m c v} \int_0^{l_{\text{mm}}} d\mathbf{l} \mathbf{v} \times \delta\mathbf{B}_{\text{mm}}. \quad (1)$$

Assuming relativistic CRs with $v \approx c$, $E = \gamma mc^2$ and $r_g = \gamma mc^2/qB$ determined by $B \geq \delta B_{\text{mm}}$, the scattering angle at $\delta t \approx l_{\text{mm}}/c$ is

$$\delta\Theta \approx \frac{|\delta\mathbf{v}|}{c} \sim \frac{l_{\text{mm}}}{r_g} \frac{\delta B_{\text{mm}}}{B}. \quad (2)$$

Assuming that these small-angle deflections add up to a correlated random walk, the scattering rate is

$$\nu_{\text{mm}} \sim \frac{\delta\Theta^2}{\delta t} \approx \frac{c l_{\text{mm}}}{r_g^2} \left(\frac{\delta B_{\text{mm}}}{B} \right)^2, \quad (3)$$

which implies a spatial diffusion coefficient of

$$\kappa_{\text{mm}} \approx \frac{c^2}{\nu_{\text{mm}}} \approx \frac{cr_g^2}{l_{\text{mm}}} \left(\frac{\delta B_{\text{mm}}}{B} \right)^{-2} \propto E^2 l_{\text{mm}}^{-1}. \quad (4)$$

As usual, more energetic CRs diffuse much faster.

In arriving at equation (4), we effectively assumed that micromirrors are described by only one characteristic scale, l_{mm} . In reality, micromirrors are anisotropic (see ellipsoid-like shapes in Fig. 1) with scales perpendicular (\perp) and parallel (\parallel) to the ambient magnetic field that satisfy $l_{\perp,\text{mm}} \ll l_{\parallel,\text{mm}}$. While gyrating through this field, CRs with $r_g \gg l_{\text{mm}}$ mostly traverse micromirrors perpendicularly. Only low-energy CRs satisfying $r_g v_{\perp}/c \lesssim l_{\perp,\text{mm}} \ll l_{\parallel,\text{mm}}$ are an exception, and should be treated analogously to thermal electrons with negligible r_g being scattered^{29,33} and trapped³⁴ in the micromirrors. This subpopulation makes a negligible contribution to the overall transport of CRs with $r_g \gg l_{\perp,\text{mm}}$ considered here.

For $r_g \gg l_{\perp,\text{mm}}$, CRs will sample many different micromirrors, with deflections adding up to a correlated random walk. During one gyro-orbit, CRs will travel $\Delta l_{\parallel} \approx 2\pi r_g v_{\parallel}/c$ in the field-parallel direction. CRs with large pitch angles satisfying $\Delta l_{\parallel} \lesssim l_{\parallel,\text{mm}}$, that is, $v_{\perp}/v_{\parallel} \approx c/v_{\parallel} \geq 2\pi r_g/l_{\parallel,\text{mm}}$, that sample the same micromirror repeatedly, may become relevant only at low energies $r_g \lesssim l_{\parallel,\text{mm}}/2\pi$, not considered in this study. The scattering rate associated with the parallel micromirror perturbation $\delta B_{\parallel} \approx B_{\text{mm}}$ decreases with decreasing pitch angle, but this is overcome by scattering at the perpendicular micromirror component $\delta B_{\perp} \approx \delta B_{\parallel} l_{\perp,\text{mm}}/l_{\parallel,\text{mm}}$ for $v_{\perp}/v_{\parallel} \lesssim l_{\perp,\text{mm}}/l_{\parallel,\text{mm}} \ll 1$. Except for this cone containing CRs with small pitch angles, from which they escape quickly on the timescale $t_{\text{esc}} \approx \nu_{\text{mm}}^{-1} l_{\perp,\text{mm}}/l_{\parallel,\text{mm}}$, gyrating CRs cross micromirrors perpendicularly faster than they traverse them in the parallel direction, implying $l_{\text{mm}} \approx l_{\perp,\text{mm}}$ to be the relevant scale.

For application to the ICM, we estimate l_{mm} using an asymptotic theory of the mirror instability's nonlinear evolution³⁵ supported by previous numerical studies^{6,36}: $l_{\text{mm}} \approx (\tau\Omega)^{1/8} r_{g,i}$, where τ is the timescale over which a micromirror-susceptible plasma evolves macroscopically. Applying the theory to an ICM^{2,37} with $B \approx 3 \mu\text{G}$, the thermal-ion gyroradius $r_{g,i} \approx (2T/m_i)^{1/2}/\Omega_i \approx 1 \text{ npc}$, the mean galaxy cluster temperature $T \approx 5 \text{ keV}$, the thermal-ion mass m_i and $\tau \approx 10^{12} \text{ s}$ (refs. 31,35) yields $l_{\text{mm}} \approx l_{\perp,\text{mm}} \approx 100 r_{g,i} \approx 100 \text{ npc}$ (ref. 36), only a factor of a few smaller than the gyroradius of a giga-electronvolt CR. In combination with equation (4), this gives us the estimate

$$\begin{aligned} \kappa_{\text{mm}} &\approx 10^{30} Z^{-2} \left(\frac{l_{\text{mm}}}{100 \text{ npc}}\right)^{-1} \left(\frac{B}{3 \mu\text{G}}\right)^{-2} \left(\frac{\delta B_{\text{mm}}/B}{1/3}\right)^{-2} \left(\frac{E}{\text{TeV}}\right)^2 \text{ cm}^2 \text{ s}^{-1}, \\ &\approx 10^{30} Z^{-2} \left(\frac{T}{5 \text{ keV}}\right)^{-1/2} \left(\frac{B}{3 \mu\text{G}}\right)^{-1} \left(\frac{\delta B_{\text{mm}}/B}{1/3}\right)^{-2} \left(\frac{E}{\text{TeV}}\right)^2 \text{ cm}^2 \text{ s}^{-1}. \end{aligned} \quad (5)$$

This estimate is valid provided that $l_{\text{mm}} \ll r_g$ and $\delta t v_{\text{mm}} \approx (l_{\text{mm}}/r_g)^2 (\delta B_{\text{mm}}/B)^2 \ll 1$ ($E \gg 100 \text{ MeV}$).

We now show that the diffusion coefficient (5) is associated with parallel transport along field lines by demonstrating that the perpendicular diffusion coefficient is negligible. Each scattering at $\delta t \approx l_{\text{mm}}/c$ moves the gyrocentre by a distance $\Delta r_{\perp} \approx r_g \delta\theta$ in the plane perpendicular to the local magnetic-field line. Using the estimate (2) for the scattering angle leads to the perpendicular diffusion coefficient

$$\kappa_{\perp,\text{mm}} \approx \frac{\Delta r_{\perp}^2}{\delta t} \approx \frac{r_g^2 \delta\theta^2}{l_{\text{mm}}/c} \approx c l_{\text{mm}} \left(\frac{\delta B_{\text{mm}}}{B}\right)^2 \approx \frac{l_{\text{mm}}^2}{r_g^2} \left(\frac{\delta B_{\text{mm}}}{B}\right)^4 \kappa_{\text{mm}}. \quad (6)$$

Since $\kappa_{\perp,\text{mm}} \ll \kappa_{\text{mm}}$ for $r_g \gg l_{\text{mm}}$ and $\delta B_{\text{mm}} \lesssim B$, it is the parallel diffusion $\kappa_{\parallel,\text{mm}} \approx \kappa_{\text{mm}}$ along field lines that dominates. The smaller perpendicular diffusion coefficients arise from anisotropic scattering. The degree to which this anisotropy enhances the parallel diffusion coefficient depends on the specifics of pitch-angle scattering and the properties of B_{\perp} , both of which warrant further investigation.

To validate our theoretical prediction for the diffusion coefficients (5) and (6), we performed a numerical experiment in which a spectrum of CRs was integrated in a magnetic field containing micromirrors generated self-consistently via a particle-in-cell (PIC) simulation ('Micromirror field from PIC simulations'). For our numerical experiment, we selected two representative realizations of the three-dimensional (3D) field during its secular evolution, visualized in Fig. 1. We then determined the diffusion coefficients of the CRs in both fields by integrating the CR equation of motion. The results are shown in Fig. 2. The diffusion coefficients in the micromirror fields show good agreement with equations (5) and (6).

The micro-macro physics transition is at teraelectronvolt CR energies

In 'Effect of micromirrors on large-scale CR transport', we derived the diffusion coefficient $\kappa_{\text{mm}} \propto l_{\text{mm}}^{-1} E^2$ associated with CR scattering at micromirrors of scale l_{mm} . To determine the upper bound for the energies at which the scattering off micromirrors dominates CR transport, we need a model of the competing contribution from the resonant scattering off mesoscale magnetic turbulence. In 'Model of CR scattering at mesoscales', we present models of CR diffusion based on resonant scattering in the (mesoscale) inertial range of magnetic turbulence stirred at the macroscale l_c , leading to the diffusion coefficient

$$\kappa_{\text{res}} \sim c l_c \left(\frac{r_g}{l_c}\right)^{\delta} \propto E^{\delta} l_c^{-\delta+1}, \quad (7)$$

where the model-dependent exponent is $0 \leq \delta \leq 1/2$. The mechanism with the smallest diffusion coefficient dominates CR transport.




Fig. 2 | Diffusion coefficients of CRs in micromirror fields. Our predictions (5) (grey dashed line) and (6) (grey dash-dotted line), valid for $l_{\text{mm}} \ll r_g$, agree well with the computed diffusion coefficients. κ is dominated by the parallel diffusion; the perpendicular diffusion coefficient κ_{\perp} is negligible. Note that κ decreases with $\delta B_{\text{mm}}/B$ (triangles versus stars). We include results for CRs with gyroradii smaller than l_{mm} (vertical orange line) to highlight the change in transport regimes, which follows from our theory. The grey markers show the transport of CRs through the residual field that results after filtering out the wavenumbers associated with the micromirrors, leaving only numerical noise. The purple vertical (dash-dotted) lines show grid resolution and box sizes along the three axes. $Z = 1$ is used for the energy scale.

The transition between the micromirror and resonant-scattering transport regimes occurs when $\kappa_{\text{mm}} \approx \kappa_{\text{res}}$. Equating (4) and (7) determines the gyroradius corresponding to this transition:

$$r_g \approx l_c \left(\frac{\delta B_{\text{mm}}}{B}\right)^{2/(2-\delta)} \left(\frac{l_{\text{mm}}}{l_c}\right)^{1/(2-\delta)}. \quad (8)$$

This translates into a δ -dependent estimate for the transition energy:

$$\begin{aligned} E &\approx Z \left(\frac{B}{3 \mu\text{G}}\right) \left(\frac{l_c}{100 \text{ kpc}}\right) \\ &\times \begin{cases} 300 \left(\frac{\delta B_{\text{mm}}}{B}\right)^{2/(2-\delta)} \left(\frac{l_{\text{mm}}}{l_c}\right)^{1/(2-\delta)} \text{ EeV}, & \text{general } \delta, \\ 5 \left(\frac{\delta B_{\text{mm}}/B}{1/3}\right)^{6/5} \left(\frac{l_{\text{mm}}/l_c}{10^{-12}}\right)^{3/5} \text{ TeV}, & \delta = 1/3, \\ 600 \left(\frac{\delta B_{\text{mm}}/B}{1/3}\right)^{4/3} \left(\frac{l_{\text{mm}}/l_c}{10^{-12}}\right)^{2/3} \text{ GeV}, & \delta = 1/2. \end{cases} \end{aligned} \quad (9)$$

Below this energy, magnetic micromirrors dominate CR diffusion. The factor involving the ratio l_{mm}/l_c accounts for the scale separation between micro- and macrophysics, which is $\sim 10^{-12}$ in our fiducial ICM under the same assumptions as in 'Effect of micromirrors on large-scale CR transport'.

To test this prediction, we performed numerical simulations of CR transport in the ICM (detailed in Methods), modelling the effects of both the micromirrors ('Model of small-angle scattering in magnetic micromirrors') and of the turbulent cascade up to $l_c \approx 100 \text{ kpc}$ ('Model of CR scattering at mesoscales'). Figure 3 summarizes our results by presenting the CR diffusion coefficient as a function of energy. The vertical light-blue bar indicates our estimate (9) for the micro-macro transition assuming the most likely range of $\delta \in [1/3, 1/2]$. While this estimate of the micro-macro physics transition at teraelectronvolt CR energies




Fig. 3 | Diffusion coefficients of CRs in the ICM as functions of CR energy. The black stars show the diffusion coefficients in the micromirror field generated by a PIC simulation ('Micromirror field from PIC simulations'). The olive open crosses show the diffusion coefficients of CRs in MHD turbulence without a guide field ('Turbulence from MHD simulations'). The other open markers show the diffusion coefficients computed in isotropic synthetic turbulence with a large inertial range ('Model of synthetic magnetic turbulence'), together with our stochastic micromirror-scattering model ('Model of small-angle scattering in magnetic micromirrors'), assuming the volume-filling fraction f_{mm} of micromirrors indicated in the legend ('The case of spatially intermittent micromirrors'). The grey dash-dotted lines represent theories for CR transport depending on the macroscale l_c according to equation (7), including the most efficient (Bohm) and the least efficient (energy-independent) diffusion scenarios. The grey dotted line represents the diffusion due to streaming instability, according to equation (18) in 'Model of CR streaming'. The grey dashed line represents our prediction of the diffusion due to micromirrors according to equation (5). The vertical light-blue bar indicates our estimate (9) for the micro-macro transition for the most likely range of δ between 1/3 and 1/2. Simulation results from refs. 10 and 106 illustrate the best resolution towards the limit $r_g \ll l_c$ achieved before the present results with synthetic turbulence on a grid (2,048³ grid points) and nested grids, respectively. The effects of field-line tangling are not considered, which is expected to reduce the global CR diffusion coefficients by a factor of three.

is numerically confirmed using synthetic turbulence, we also used magnetic fields from direct PIC and magnetohydrodynamic (MHD) simulations to validate the consistency of our numerical approach at micro- and macroscales, respectively, and capture all relevant diffusion coefficients discussed in the literature, detailed as points (1)–(3) in 'Model of CR scattering at mesoscales'.

The case of spatially intermittent micromirrors

Thus far, we have effectively assumed that the micromirrors permeate the plasma uniformly. In reality, the situation is more complicated: micromirrors will most probably appear in spatially intermittent and temporally transient patches wherever turbulence leads to local amplification of the magnetic field at a rate that is sufficiently large to engender positive pressure anisotropy exceeding the mirror-instability threshold ($-1/\beta$; see, e.g., refs. 36,38 and references therein). This gives rise to an effectively two-phase plasma (see Fig. 4 for an illustration), with two different effective scattering rates: $v_{\text{mm}} \approx c^2/\kappa_{\text{mm}}$ in micromirror patches and $v_{\text{res}} \approx c^2/\kappa_{\text{res}}$ elsewhere (instead of κ_{res} , one could also use κ_{st} associated with the streaming instability—'Discussion'). For the purpose of modelling CR scattering in such a plasma, we introduce the effective micromirror fraction f_{mm} to quantify the probability of CR being scattered by the micromirrors.

By definition, the effective scattering rate in a two-phase medium is^{39,40}

$$v_{\text{eff}} = f_{\text{mm}} v_{\text{mm}} + (1 - f_{\text{mm}}) v_{\text{res}}. \quad (10)$$

The effective diffusion coefficient is then

$$\kappa_{\text{eff}} \approx \frac{\kappa_{\text{mm}}}{f_{\text{mm}} + (1 - f_{\text{mm}})\kappa_{\text{mm}}/\kappa_{\text{res}}}. \quad (11)$$

The transition at which micromirror transport takes over from resonant scattering is entirely independent of f_{mm} : $\kappa_{\text{eff}} \approx \kappa_{\text{res}}$ when $\kappa_{\text{mm}} \approx \kappa_{\text{res}}$. However, the asymptotic scaling $\kappa_{\text{eff}} \approx \kappa_{\text{mm}}/f_{\text{mm}}$ is only reached at CR energies for which $\kappa_{\text{mm}}/\kappa_{\text{res}} \lesssim f_{\text{mm}}/(1 - f_{\text{mm}})$, pulling the transition energy down by a factor of $f_{\text{mm}}^{1/(2-\delta)}$. This is not a very strong modification of our cruder ($f_{\text{mm}} = 1$) estimate (9) unless f_{mm} is extremely small.

The most intuitive interpretation of f_{mm} is that it is the fraction of the plasma volume occupied by the micromirrors. This, however, requires at least two caveats. (1) The lifetime of micromirror patches, determined by the turbulent dynamics, can be shorter than the time for a CR to diffuse through the patch. (2) If the micromirror patches form solid macroscopically extended 3D blobs, it is possible to show that CRs typically do not penetrate much farther than the mean free path $\lambda_{\text{mm}} = c/v_{\text{mm}}$ into the patches. This leaves the patch volume largely uncharted (Appendix 3 of ref. 40). The second concern obviates the first ('Model of a static two-phase inhomogeneous medium'). Under such a scenario, the effective CR diffusion in the ICM might be determined primarily by such factors as the typical size of the patches and the distance between them⁴¹. However, the scenario of micromirror-dominated transport is made more plausible as the diffusion is mostly along the field lines. In this one-dimensional (1D) problem, CRs bounce between mirror patches on the same field line until they have a lucky streak in diffusion and pass directly through a micromirror patch. This trapping effect leads to efficient confinement, if the influence of potential field-line separation and cross-field diffusion is found to be negligible, though this remains subject to further research.

Our model formula (11) proves to be a good prediction even in a simple modification of our numerical experiment with synthetic fields, designed to model micromirror patches ('Model of a static two-phase inhomogeneous medium'). Its results are shown in Fig. 5. It is a matter for future work to determine the precise dependence of f_{mm} on the morphology and dynamics of magnetic fields and micromirror-unstable patches in high- β turbulence—itsself a system that has only recently become amenable to numerical modelling^{38,42,43}. Here, we proceed to discuss the implications of dominant micromirror transport, assuming that f_{mm} is not tiny, namely, $f_{\text{mm}} \geq 0.1$, as indeed observed in recent numerical simulations^{33,43–45}. Studies of Faraday depolarization of radio emission from radio galaxies could be in principle used to constrain the volume-filling fraction of micromirrors, because depolarization increases proportionally to f_{mm} . We tested the expected Faraday depolarization arising from micromirrors within galaxy clusters in a numerical experiment using our PIC simulations ('Micromirror field from PIC simulations'). The expected depolarization angles due to micromirror fluctuations for wavelengths observed with the Very Large Telescope (VLT) and the Low-Frequency Array (LOFAR) are too small to constrain f_{mm} . Details of the resolution element in radio telescope observations may affect this result, an issue reserved for future studies.

Discussion

We have argued that CR diffusion in the ICM is determined by microscale (l_{mm}) mirrors at CR energies $\text{GeV} \lesssim E \lesssim \text{TeV}$ and by macroscale (l_c) turbulence at $E \gtrsim \text{TeV}$. Although the micro-macro physics transition mainly depends on these two scales, there is a degree of fine-tuning at mesoscales. This refers to the role played in equation (9) by the exponent δ , which depends on the details of the scattering mechanism and




Fig. 4 | Visualization of example CR trajectories through spatially intermittent micromirror patches. The numerical experiment to study the effective CR transport in a two-phase medium is described in ‘Model of a static two-phase inhomogeneous medium’. The grey surfaces are the isosurfaces of a threshold field strength B_* . In our simplified numerical experiment, these isosurfaces are assumed to enclose the micromirror patches, inside which the diffusion coefficient is much smaller than it is outside. Therefore, a given

choice of B_* corresponds to a certain value of f_{mm} (‘Model of a static two-phase inhomogeneous medium’). Example CR trajectories show increased deflections within the micromirror patches (see, for example, the lower right corner of the left panel and the zoom into a micromirror patch in the right panel). Taking the patches to be static is suitable for small f_{mm} as demonstrated in ‘Model of a static two-phase inhomogeneous medium’.




Fig. 5 | Effective diffusion coefficient of CRs in a two-phase medium versus the effective micromirror fraction f_{mm} . We model the CR transport through a two-phase medium in 3D and 1D (to reduce simulation costs), as explained in ‘Model of a static two-phase inhomogeneous medium’. The 3D case (red diamonds) shows the diffusion coefficient of 300 GeV CRs computed at their trajectory lengths of $\sim 10l_c$. The blue and grey lines show the 1D results on 400 test trajectories using our recurrent neural network (RNN) and the classical analysis method. Error bars for the 3D data represent the s.d. across different realizations of patches for each f_{mm} , while for the 1D cases the coloured contours show the s.d. across 400 CRs for a single realization. The black dash-dotted line represents expected values for averaged scattering frequencies of CRs in the two-phase plasma, formally expressed in equation (11). κ_{mm} and κ_{res} are recovered for $f_{\text{mm}} = 1$ and $f_{\text{mm}} = 0$, respectively.

of the turbulent cascade (‘Model of CR scattering at mesoscales’). While our study is tailored to the ICM, it can be adapted to other nearly collisionless high- β plasmas, such as the hot interstellar medium and the Milky Way halo. In what follows, we discuss what this revised picture of CR diffusion in the ICM implies for our understanding of (Fermi/radio) bubbles and other similar large-scale morphologies.

The reduction of CR diffusion caused by micromirrors may act as a transport barrier in the ICM, reducing the escape of sub-teraelectronvolt CRs from their sources. For example, as radio bubbles rise through the ICM, their confinement of CRs may be dominated by micromirror confinement rather than by the conventional mechanism of magnetic draping; this depends on the details of diffusion coefficients both parallel and perpendicular to the mean magnetic field (see, for example, refs. 46–48). Reference 49 considered the CR transport in such systems, and showed that the sub-teraelectronvolt CRs form a thin layer on the surface of the bubble and acquire hardened spectral-density distributions. Such sharp boundaries and hardened CR spectra, which translate into hardened photon spectra, are indeed

observed in popular morphologies such as the Fermi bubble⁵⁰ and the radio bubble in the Ophiuchus galaxy cluster⁵¹. The decreased diffusion coefficients of sub-teraelectronvolt CRs could also be relevant in high- β regions within galaxies. One possible example is the Cygnus Cocoon, a Galactic PeVatron, located within a star-forming region, where the observationally constrained suppressed CR diffusion coefficients^{52,53} may be explained by the additional collisionality due to micromirrors.

We have shown that micromirror diffusion substantially reduces the CR mean free path, thus in principle making CR coupling to the ambient motions tighter. However, to assess what this does to the efficiency (or otherwise) of the (re)acceleration^{54–60}, we must have a somewhat more detailed picture than we currently do of the nature of the ICM turbulence (that is, of turbulence in a weakly collisional, high- β plasma—a topic of active current investigations^{38,42,43}) and of how the micromirror patches might be shaped and spatially distributed in this turbulence. This will require further study before the (re)acceleration question is settled.

A further example of how micromirror diffusion may impact the surrounding plasma arises from the observation that, as the scattering of sub-teraelectronvolt CRs at micromirrors increases the effective CR collisionality in high- β environments, the effective operation of the CR streaming instability within micromirror patches is put into doubt (‘CR streaming instability’). Note that a patchy distribution of the micromirrors may allow for the existence of regions where the streaming instability remains active—indeed, possibly more so than usually expected, as those micromirror-free regions are likely to feature negative pressure anisotropies and, therefore, reduced effective Alfvén speeds. Models designed to explain the thermal balance between heating and cooling of galaxy clusters based on collisionless, resonant mechanisms^{61–63} thus become less plausible.

Finally, the micromirror scattering matters for cosmological studies of the evolution of the ICM and galaxy clusters. The suppressed CR diffusion coefficients offer a compelling justification for how CRs can be effectively ‘frozen’ within the ICM as key parameters such as gas density and magnetic-field intensity evolve—an assumption fundamental to recent models of the dynamical evolution of galaxy clusters and their surroundings^{59,64,65}. This impacts the interplay between CRs and other astrophysical processes within these massive cosmic structures^{66,67}.

While it is well established that macroscopic dynamics can trigger microscopic phenomena, the potentially transformative impact of micromirrors on CR diffusion provides a lesson that microphysics can reciprocally affect macroscopic dynamics and observable structures across a range of astrophysical scales.

Methods

Modelling CR transport across a wide energy spectrum from gigaelectronvolt to exaelectronvolt energies in a multiscale high- β plasma presents methodological and computational challenges. For our numerical

simulations, we choose parameters from the ICM, a high- β plasma. Rotation-measure data indicate magnetic-field strengths of ~ 0.1 – $1 \mu\text{G}$ averaged over a cubic megaparsec ICM volume^{37,68}, with typical field strengths of several microgauss in central regions (see, for example, ref. 69). Numerical simulations support these estimates (for example, ref. 70). We choose $B \approx 3 \mu\text{G}$, $\delta B_{\text{mm}}/B \approx 1/3$, $l_{\text{mm}} \approx 100 \text{ npc}$ and turbulence $l_c \approx 100 \text{ kpc}$ (ref. 2). Simulations are performed using the publicly available tool CRPropa 3.2 (ref. 71) with additional extensions and modelling choices described below.

Model of synthetic magnetic turbulence

Modelling the competing micro- and macrophysical transport effects requires resolving turbulence over at least ten decades in scale. Current MHD and PIC simulations are unsuitable for this as they only allow for limited scale ranges⁷². Even the current best grid resolutions of more than 10^{12} cells resolve less than four decades of scale separation. Synthetic turbulence, on the other hand, can be generated by summing over n_m plane waves at an arbitrary particle position \mathbf{r} as follows^{15,73}:

$$\delta\mathbf{B}(\mathbf{r}) = \text{Re} \left(\sum_{n=1}^{n_m} \delta\mathbf{B}_n^* e^{i\mathbf{k}_n \cdot \mathbf{r}} \right) = \sqrt{2} \delta B \sum_{n=1}^{n_m} \xi_n A_n \cos(k_n \hat{\mathbf{k}}_n \cdot \mathbf{r} + \phi_n), \quad (12)$$

with normalized amplitudes A_n determined by the assumed turbulent energy spectrum, uniformly distributed phase factors $\phi_n \in [0, 2\pi]$, unit wavevectors $\hat{\mathbf{k}}_n$ and polarizations ξ_n , satisfying $\hat{\mathbf{k}}_n \cdot \xi_n = 0$. We employ the performance-optimized method described in ref. 74. We investigated the number of wavemodes n_m needed by analysing turbulence characteristics and diffusion coefficients of CRs and found that $n_m = 1,024$ log-spaced wavemodes are sufficient for diffusion coefficients to converge.

We compared our key results on CR transport obtained with the above method with those obtained with an alternative method for synthetic turbulence, where we followed the approach proposed, for example, in refs. 75 and 72. In this alternative method, synthetic turbulence is precomputed and stored on many discrete nested grids at different scales, with magnetic fluctuations between scales $l_{\text{min},i}$ and $l_{\text{max},i}$ with individual magnetic-field strengths $\delta B_i^2 = \delta B^2 (l_{\text{max},i}^{\xi-1} - l_{\text{min},i}^{\xi-1}) / (l_{\text{max}}^{\xi-1} - l_{\text{min}}^{\xi-1})$ and $l_{\text{max}} \approx 5l_c$. We found agreement between results obtained using the two methods.

Note that CRs below petaelectronvolt energies diffuse on time-scales smaller than the typical timescale of large-scale turbulence motions, justifying our modelling turbulence as static. CRs are frozen within the ICM as the plasma evolves ('Discussion').

CR trajectories

Computing CR trajectories in magnetic fields involves solving the equation of motion for charged particles. These are then used to calculate the statistical transport characteristics of CRs. We use the Boris-push method for this task, as implemented in ref. 71. This captures the dynamics of charged particles in magnetic fields while preserving key properties, such as CR energy.

Model of small-angle scattering in magnetic micromirrors

We model the effect of magnetic micromirrors as a change of propagation after a distance s by an angle $\delta\theta$ given $v_{\text{mm}} = \delta\theta^2 c/s$. Particles travelling the mean free path $\lambda_{\text{mm}} = c/v_{\text{mm}}$ will have lost the information of their original direction. At each step s , chosen to be smaller than λ_{mm} , we introduce a small deflection

$$\delta\theta = X \sqrt{\frac{sv_{\text{mm}}}{c}}, \quad (13)$$

where the random Gaussian variable X with mean 0 and s.d. 1 represents the assumption that CRs random-walk their way through the magnetic micromirrors. Alternatively, micromirrors could be directly modelled

in the magnetic field, which would, however, necessitate step sizes $s \lesssim l_{\text{mm}} \ll \lambda_{\text{mm}}$, leading to significantly longer simulation times.

Model of CR scattering at mesoscales

There is an ongoing debate about the dominant mechanism of CR scattering off mesoscale fluctuations, with theories including 'extrinsic' (cascading) turbulence and 'self-excitation' (by kinetic CR-driven instabilities) scenarios (see refs. 76,77 for recent overviews). Here we focus on the extrinsic scenario, with the self-excitation described in 'CR streaming instability'.

The CR diffusion coefficient due to resonant scattering is, by dimensional analysis,

$$\kappa_{\text{res}} \approx \frac{c r_g}{f(r_g)}, \quad (14)$$

where $f(r_g)$ is a dimensionless model-dependent numerical factor expressing the efficiency of the resonant scattering off turbulent magnetic structures at the scale $l \approx r_g$. We assume diffusion rather than superdiffusion, which would increase diffusion coefficients with time. This serves as a conservative estimate. Quasilinear theory¹⁴ determines $f(r_g)$ for isotropic turbulence as the fraction of the parallel turbulent power located at the gyroresonant scales $l = 2\pi/k_{\parallel} \approx r_g$, namely, $f(r_g) \approx \int_{2\pi/r_g}^{\infty} dk_{\parallel} P(k_{\parallel}) / (B^2/8\pi) \leq 1$, where k_{\parallel} is the parallel wavenumber and $P(k_{\parallel})$ is the parallel magnetic-energy spectrum. Assuming an undamped turbulent cascade with $P(k_{\parallel}) \propto k_{\parallel}^{-\xi}$ gives $f(r_g) \approx (r_g/l_c)^{\xi-1}$, where l_c is the energy-containing scale. Therefore,

$$\kappa_{\text{res}} \approx c l_c \left(\frac{r_g}{l_c} \right)^{2-\xi} \propto E^{2-\xi} l_c^{\xi-1} = E^{\delta} l_c^{-\delta+1}, \quad (15)$$

where $\delta = 2 - \xi$ is defined for convenience. In 'The micro-macrophysics transition is at teraelectronvolt CR energies', we confirmed this scaling via numerical simulations with unprecedented spatial resolution for a synthetic turbulence composed of plane waves.

An important qualitative result is that the cases $\delta = 2$ ($\xi = 0$) and $\delta = 1$ ($\xi = 1$) apply only to $r_g \gg l_c$ and $r_g \approx l_c$, respectively. While there is no realistic turbulence model with $\xi = 0$, this case formally corresponds to the small-angle scattering limit of CRs with $r_g \gg l_c$: equation (7) then yields $\kappa \approx cr_g^2/l_c$, which is identical to equation (4) if we replace l_{mm} and δB_{mm} with l_c and B , respectively. This is the standard theory for the high-energy regime referred to in the introduction.

In fact, only scalings with weaker energy dependence for $r_g \ll l_c$ are typically considered. In this limit, three popular choices for this scaling have appeared in the literature: (1) $\delta = 1/3$, corresponding to isotropic turbulence with a ref. 78 spectrum ($\xi = 5/3$); (2) $\delta = 1/2$, corresponding to $\xi = 3/2$, which in the past was associated with the theory by refs. 79 and 80 for weak, isotropic Alfvénic MHD turbulence (now known not to exist); (3) $\delta = 0$, corresponding to the $\xi = 2$ Goldreich–Sridhar parallel spectrum of critically balanced Alfvénic turbulence⁸¹ (see ref. 82 for a review) by adhering to equation (7) (note that this is not observed: see, for example, ref. 83); nevertheless, this spectrum provides a simple way to estimate the decreased CR-scattering efficiency expected for the anisotropic turbulent cascade). Historically, Alfvénic turbulence was favoured until it was realized that scale-dependent anisotropy^{84–86}, damping^{77,87} and intermittency^{88,89} might lead to inefficient gyroresonant scattering. A putative $\xi = 3/2$ cascade of fast MHD modes, if isotropic and robust against steepening^{76,90} and various damping mechanisms, may help by generating fluctuations with large enough frequencies and amplitudes to scatter CRs efficiently^{91,92}. More recently, the exponents $\delta = 1/3$ and $\delta = 1/2$ were ascribed to CR scattering in intermittent distributions of sharp magnetic-field bends in Goldreich–Sridhar turbulence¹⁶ and in an MHD turbulent dynamo¹⁷. Scaling exponents in the range $0.3 \lesssim \delta \lesssim 0.5$ are in broad agreement with constraints from Galactic observations (see ref. 93, for a review).

An additional process that may contribute to the diffusion of low-energy CRs arises from CRs following diffusing magnetic-field lines. The Alfvénic scale $l_A \gtrsim 1$ kpc (ref. 94) approximates the mean free path of CRs following these field lines⁹⁵. The associated CR diffusion coefficient $\kappa_{\text{trw}} \approx cl_A \gtrsim 10^{32} \text{ cm}^2 \text{ s}^{-1}$ does not fall significantly below κ_{res} for $r_g \ll l_c$. Given this estimate, we neglect this transport process in our simulation set-up, but indicate the value of the diffusion coefficient corresponding to it as an upper boundary in Fig. 3.

CR streaming instability

Let us now explain why the streaming instability can be ignored in our multiscale model of CR transport within micromirror patches. The self-confinement of CRs due to the streaming instability is believed to play an important role in the Galaxy²⁴ and in galaxy clusters⁶². In this picture, the streaming instability generates fluctuations of the magnetic field, which in turn can scatter CRs. It is believed that this mechanism may take over at lower CR energies, with details depending on the instability's growth rate at wavenumber k . At gyroscale²²,

$$\gamma_{\text{SI}} \approx \Omega_i \frac{n_{\text{CR}}(>E)}{n_i} \left(\frac{v_{\text{st}}}{v_A} - 1 \right) \approx 10^{-14} \left(\frac{B}{3 \mu\text{G}} \right) \left(\frac{E}{\text{TeV}} \right)^{-1.6} \text{ s}^{-1}, \quad (16)$$

where $n_{\text{CR}}(>E)$ is the density of CRs with energies above energy E corresponding to the resonance condition that can interact resonantly with waves with wavenumber k , n_i is the ambient ion density, v_A is the Alfvén speed and v_{st} is the streaming speed, believed to be of the order of v_A in saturation for the \sim GeV CRs^{62,96}. In the second estimate in equation (16), we employed the common assumptions (see, for example, refs. 97,98, and references therein) that $(v_{\text{st}}/v_A - 1) \approx 1$ and $n_{\text{CR}}(>E)/n_i \approx 10^{-7} (E/\text{GeV})^{1-\alpha}$ in galaxy clusters, with $\alpha \approx 2.6$.

Scattering of sub-teraelectronvolt CRs at micromirrors increases the effective CR collisionality in high- β environments. A comparison of the gyroscale growth rate (16) with the scattering rate at micromirrors (3) gives

$$\frac{\gamma_{\text{SI}}}{\nu_{\text{mm}}} \approx 10^{-5} Z^{-2} \left(\frac{T}{5 \text{ keV}} \right)^{-1/2} \left(\frac{\delta B_{\text{mm}}/B}{1/3} \right)^{-2} \left(\frac{E}{\text{TeV}} \right)^{0.4}. \quad (17)$$

With such a large effective collisionality isotropizing and homogenizing CRs, it is doubtful that this gyroscale, resonant instability can operate.

Another way to gauge the importance of the streaming instability is to imagine that it is not suppressed and then check for self-consistency. In particular, for self-confined CRs, the CR-density scale height H is set by the properties of the ambient thermal gas and is of the order of the thermal-gas-density scale height H_ρ . If scattering by micromirrors is present with diffusion coefficient κ_{mm} , the associated diffusive flux is smaller than the minimum flux required for the streaming instability to operate if $\kappa_{\text{mm}}/H \lesssim v_A$. This corresponds to scattering by micromirrors suppressing the anisotropy in the CR distribution function to levels below $-v_A/c$, which is the threshold anisotropy for the streaming instability to operate in the first place. Thus, because $\kappa_{\text{mm}}/H \lesssim v_A$ for sub-teraelectronvolt CRs, where $H \approx H_\rho \gtrsim 10$ kpc (refs. 99,100), CRs may not self-confine in a self-consistent manner.

Model of CR streaming

Let us imagine that the instability is not suppressed, despite the arguments made in 'CR streaming instability', and estimate⁹⁵

$$\kappa_{\text{st}} \approx l_A v_{\text{st}} \gtrsim l_A v_A \gtrsim 3 \times 10^{28} \left(\frac{l_A}{1 \text{ kpc}} \right) \left(\frac{v_A}{100 \text{ km s}^{-1}} \right) \text{ cm}^2 \text{ s}^{-1}, \quad (18)$$

where we have used $v_A \approx 100 \text{ km s}^{-1}$ as the lower limit of the streaming speed. We also assumed that the magnetic-field lines stochastically tangled on the scale l_A . In super-Alfvénic turbulence, this marks the transition towards fully MHD turbulence (see, for example, ref. 95), as

it is the scale at which the turbulent velocity matches the Alfvén speed. $l_A \gtrsim 1$ kpc under typical ICM conditions⁹⁴.

Comparing this CR diffusivity with the one caused by micromirrors (5) gives

$$\frac{\kappa_{\text{mm}}}{\kappa_{\text{st}}} \lesssim Z^{-2} \left(\frac{E}{100 \text{ GeV}} \right)^2 \left(\frac{l_A}{1 \text{ kpc}} \right)^{-1} \left(\frac{T}{5 \text{ keV}} \right)^{-1/2} \left(\frac{B}{3 \mu\text{G}} \right)^{-1} \left(\frac{\delta B_{\text{mm}}/B}{1/3} \right)^{-2} \left(\frac{v_A}{100 \text{ km s}^{-1}} \right)^{-1}, \quad (19)$$

demonstrating that κ_{mm} dominates CR transport in the ICM up to almost teraelectronvolt energies. This comparison, together with the arguments for the suppression of the streaming instability ('CR streaming instability'), justifies neglecting CR streaming in our numerical experiments. Nevertheless, we show this estimate in Fig. 3.

Computation of CR diffusion coefficient

We use the CR propagation software CRPropa 3.2 (ref. 71) and extend the framework with our custom modules for the generalized nested turbulence, different turbulence geometries and micromirror scattering. We choose sufficiently small step sizes $s_{\text{step}} \approx \min\{\lambda_{\text{mm}}/10^3, \lambda_{\text{res}}/10^3, \lambda_{\text{he}}/10^3\}$, with mean free path $\lambda_i \approx \kappa_i/c$ to resolve the small-angle scattering at micromirrors, the resonant scattering in the extrinsic turbulent cascade and the small-angle scattering in the high-energy limit, respectively. The option $\lambda_{\text{he}}/10^3$ is only included for CR energies above 10 EeV, as the high-energy limit is not valid below that energy. We compute sufficiently long CR trajectories $d \approx \min\{10^3 \lambda_{\text{mm}}, 10^3 \lambda_{\text{res}}, l_{\text{max}}\}$ (min is used here to save computation resources as only the more efficient scattering process needs to be resolved) for $E \lesssim 10$ EeV and $d \approx \max\{10^3 \lambda_{\text{he}}, l_{\text{max}}\}$ otherwise. The time-dependent diffusion coefficient $\kappa(t)$ for CRs performing a correlated random walk is¹⁰¹

$$\kappa(t) = \frac{\langle \Delta r^2 \rangle}{2t} \left(1 + \frac{2 \langle \cos \Theta \rangle}{1 - \langle \cos \Theta \rangle} \right)^{t \gg \lambda/c} \approx \frac{\langle \Delta r^2 \rangle}{2t}, \quad (20)$$

with r representing the CR spatial displacement and the operation $\langle \dots \rangle$ averaging over CRs. The approximation for $t \gtrsim \lambda/c$ stems from the convergence to central-limit behaviour, assuming that the deflection angles Θ are uniformly distributed after CRs travel a distance equivalent to their mean free path λ . We compute the steady-state diffusion coefficient κ by averaging $\kappa(t)$ for 10^3 CRs. When using synthetic turbulence, we average this quantity over ten different realizations.

To compute the CR diffusion coefficients efficiently, we have developed a recurrent neural network that comprises a long short-term memory layer followed by a fully connected layer, linking the output of the long short-term memory to our output size, which is the predicted diffusion coefficient. This architecture allows us to capture the temporal dependence in our data, making the network well suited to use the CR trajectories as input for the model. For training data, we used trajectories generated by Monte Carlo simulations with the method described in 'Model of a static two-phase inhomogeneous medium', using various different effective diffusion coefficients. These effective diffusion coefficients served as labels for the training process. We chose low statistics of only 400 CRs for the 1D cases to illustrate the superiority of our neural network (trained on only 1,600 CR trajectories in less than a minute on a conventional CPU) over the classical computation of the running diffusion coefficient. We tested the convergence of the latter method to the theoretical expectation in the limit of large times and CR numbers and found good agreement. We demonstrate the capabilities of the model in Fig. 5. The good performance of the network is primarily attributable to its capacity to learn efficiently that the diffusion coefficients can be predicted accurately by the frequency of small-angle scattering, which becomes evident in a relatively small number of steps. Additionally, the ability to distinguish signal (deflection caused by fluctuations) from noise (constant

gyration) further contributes to the network's superior performance. These abilities indicate that the network could also be employed as a robust framework to assess f_{mm} in MHD turbulence by propagating charged particles through the magnetic field. Reliable computation of the mean-squared diffusion coefficient requires many trajectories, necessary to deduce f_{mm} . Our use of a neural network demonstrates a concept for efficient trajectory classification and the computation of transport characteristics in astrophysical systems.

Micromirror field from PIC simulations

The micromirror field shown in Fig. 1 was self-consistently generated using the hybrid kinetic code PEGASUS++, which models the collisionless ions using a PIC method and the electrons as an isotropic, isothermal fluid. The code can simulate a plasma's expansion or contraction using a coordinate transform method (see ref. 102 for further details), which, via double-adiabatic conservation laws, produces an ion pressure anisotropy that becomes mirror unstable.

In our simulation, we initialized a uniformly magnetized plasma ($\mathbf{B}_0 = B_0 \hat{\mathbf{x}}$) with a Maxwellian population (3,000 ion macroparticles per cell) on a cubic domain. Its size is $L_0^3 = (76.0 r_{g,i0})^3$, where $r_{g,i0}$ is the initial gyroradius of thermal ions, and the grid resolution is $\Delta x = 0.1 r_{g,i0}$, $\Delta y = \Delta z = 0.3 r_{g,i0}$. The initial ion plasma beta is $\beta_{i0} = 50$. The scale L_\perp of the plasma in the direction perpendicular to the background field then evolves as $L_\perp = L_0(1 + t/\tau_{\text{crit}})^{-2}$, with a contraction timescale of $\tau_{\text{crit}} = 5 \times 10^3 \Omega_{i0}^{-1}$, while the parallel scale remains fixed. This gives rise to an ion pressure anisotropy $\Delta_i \equiv T_{\perp,i}/T_{\parallel,i} - 1 = (1 + t/\tau_{\text{crit}})^2 - 1$ that increases with time. The mirror instability is triggered at $t \approx \tau_{\text{crit}}/2\beta_{i0} \approx 0.01\tau_{\text{crit}}$, and then back-reacts at $t \approx 0.1\tau_{\text{crit}}$, entering the secular (that is, power-law) phase of growth^{6,35}. The simulation is then run until $t_{\text{end}} \approx 0.25\tau_{\text{crit}}$. The snapshots of the field shown in Fig. 1 were taken at $t \approx 0.15\tau_{\text{crit}}$ and $t = t_{\text{end}}$, respectively.

As in all PIC simulations, the finite number of macroparticles in PEGASUS++ leads to grid-scale noise in the electromagnetic fields. To diagnose the influence of this noise on our calculation of CR propagation, we performed an experiment in which we removed the micromirrors from our PEGASUS++ simulation using a Fourier filter, and integrated CRs through the residual magnetic field. While the resulting diffusion coefficients show that PIC noise also leads to diffusive CR transport, their values are much larger than those associated with the micromirrors and so can be safely ignored (Fig. 2).

Turbulence from MHD simulations

At macroscales, we computed CR diffusion coefficients in forced incompressible MHD turbulence from the John Hopkins Turbulence Databases^{103,104} to validate the consistency of our numerical approach that relies on synthetic turbulence. The MHD turbulence was generated in a direct numerical simulation of the incompressible MHD system of equations without guide fluid using $1,024^3$ nodes, employing a pseudospectral method, with energy input from a Taylor–Green flow stirring force. CR diffusion coefficients in this field are shown in Fig. 3.

Model of a static two-phase inhomogeneous medium

We model the CR transport through a two-phase inhomogeneous medium in 1D and in 3D (see a visualization in Fig. 4). In 3D, we modified the computationally intensive numerical experiment described in the previous subsections as follows: instead of imposing an additional effective v_{mm} on all CRs propagating through our turbulent field at all times and places, we now turn this scattering on only if the CR is seeing a magnetic field above B_* . This models qualitatively the fact that micromirrors are likely to appear in regions of more vigorous magnetic-field amplification. By varying B_* between 0 and ∞ , we effectively vary the micromirror volume-filling fraction between 1 and 0, respectively. We assume the most intuitive interpretation that this volume-filling fraction is the effective micromirror fraction f_{mm} in equation (10). As our estimate of the effective diffusion coefficient works

in all dimensionalities, we also model the CR transport in a two-phase inhomogeneous medium in 1D. This allows us to compute diffusion coefficients effectively for many different values of f_{mm} . In doing so, we employ a simplified Monte Carlo model with $v(x) = v_{\text{mm}}$ for $x \bmod l_p \leq f_{\text{mm}}$ and $v(x) = v_{\text{res}}$ otherwise.

We now justify modelling micromirror patches experienced by diffusing CRs as static, based on the assumption of the short residence time of CRs in patches

$$\Delta t_p \approx \frac{l_p}{c}, \quad (21)$$

where l_p is the characteristic size of the patch. The residence time of CRs in a patch can be understood intuitively: when CRs penetrate a patch, they do so ballistically up to $\lambda_{\text{mm}} \approx \kappa_{\text{mm}}/c$ for a time $\Delta t_{\text{bal}} \approx \lambda_{\text{mm}}/c$, followed by isotropic diffusion over a time Δt_{diff} . To obtain a rough estimate for Δt_{diff} , we can consider the simplified 1D case, where CRs exit either at the point of entry or at the point on the opposite end of the patch, with probabilities $p_1 \approx (l_p - \lambda_{\text{mm}})/l_p$ and $p_2 \approx \lambda_{\text{mm}}/l_p$, respectively. The corresponding times needed to exit the patch via diffusive transport are $\Delta t_1 \approx \lambda_{\text{mm}}^2/\kappa_{\text{mm}}$ and $\Delta t_2 \approx (l_p - \lambda_{\text{mm}})^2/\kappa_{\text{mm}}$. Note that in the limit $\lambda_{\text{mm}} \ll l_p$, $\Delta t_2 \approx l_p^2/c\lambda_{\text{mm}}$ can be larger than $\Delta t_{\text{lifetime}}$, meaning that patches cannot be treated as being static anymore. However, this only affects a small fraction $p_2 \approx \lambda_{\text{mm}}/l_p \ll 1$ of CRs. The mean time duration of the diffusive transport is then given by

$$\Delta t_{\text{diff}} \approx p_1 \Delta t_1 + p_2 \Delta t_2 \approx \frac{l_p - \lambda_{\text{mm}}}{l_p} \frac{\lambda_{\text{mm}}^2}{c\lambda_{\text{mm}}} + \frac{\lambda_{\text{mm}}}{l_p} \frac{(l_p - \lambda_{\text{mm}})^2}{c\lambda_{\text{mm}}} = \frac{l_p - \lambda_{\text{mm}}}{c}, \quad (22)$$

resulting in $\Delta t_p \approx \Delta t_{\text{bal}} + \Delta t_{\text{diff}} \approx l_p/c$. Reference 40 obtained this result by a more general method of solving the time-dependent diffusion equation. It means that CRs with small diffusion coefficients, corresponding to short mean free paths, will exit the patch predominantly near their point of entry. In contrast, CRs with large diffusion coefficients traverse the patch (quasi)ballistically in time $\Delta t_p \approx \Delta t_{\text{bal}} \approx l_p/c$. As $\Delta t_{\text{diff}} \ll \Delta t_{\text{lifetime}}$, micromirrors can be treated as being static for most CRs.

Data availability

The primary data, including diffusion coefficients, energies and simulation parameters, are available in ref. 105. The magnetic-field data will be made available on reasonable request to the corresponding author.

Code availability

CR simulations were performed with CRPropa3, specifically with the version <https://github.com/reichherzerp/CRPropa3>.

References

1. Schekochihin, A. A., Cowley, S. C., Kulsrud, R. M., Hammett, G. W. & Sharma, P. Plasma instabilities and magnetic field growth in clusters of galaxies. *Astrophys. J.* **629**, 139–142 (2005).
2. Kunz, M. W., Jones, T. W. & Zhuravleva, I. in *Handbook of X-ray and Gamma-ray Astrophysics* (eds Bambi, C. & Santangelo, A.) (Springer, 2022).
3. Cesarsky, C. J. & Kulsrud, R. M. Role of hydromagnetic waves in cosmic-ray confinement in the disk. Theory of behavior in general wave spectra. *Astrophys. J.* **185**, 153–166 (1973).
4. Reichherzer, P., Becker Tjus, J., Zweibel, E. G., Merten, L. & Pueschel, M. J. Turbulence-level dependence of cosmic ray parallel diffusion. *Mon. Not. R. Astron. Soc.* **498**, 5051–5064 (2020).
5. Lazarian, A. & Xu, S. Diffusion of cosmic rays in MHD turbulence with magnetic mirrors. *Astrophys. J.* **923**, 53 (2021).
6. Kunz, M. W., Schekochihin, A. A. & Stone, J. M. Firehose and mirror instabilities in a collisionless shearing plasma. *Phys. Rev. Lett.* **112**, 205003 (2014).

7. Bott, A. F. A., Cowley, S. C. & Schekochihin, A. A. Kinetic stability of Chapman–Enskog plasmas. *J. Plasma Phys.* **90**, 975900207 (2024).
8. Aloisio, R. & Berezhinsky, V. Diffusive propagation of ultra-high-energy cosmic rays and the propagation theorem. *Astrophys. J.* **612**, 900–913 (2004).
9. Kotera, K. & Lemoine, M. Inhomogeneous extragalactic magnetic fields and the second knee in the cosmic ray spectrum. *Phys. Rev. D* **77**, 023005 (2008).
10. Subedi, P. et al. Charged particle diffusion in isotropic random magnetic fields. *Astrophys. J.* **837**, 140 (2017).
11. Reichherzer, P. et al. Regimes of cosmic-ray diffusion in Galactic turbulence. *SN Appl. Sci.* **4**, 15 (2022).
12. Chen, L. E. et al. Transport of high-energy charged particles through spatially intermittent turbulent magnetic fields. *Astrophys. J.* **892**, 114 (2020).
13. Condorelli, A., Biteau, J. & Adam, R. Impact of galaxy clusters on the propagation of ultrahigh-energy cosmic rays. *Astrophys. J.* **957**, 80 (2023).
14. Jokipii, J. R. Cosmic-ray propagation. I. Charged particles in a random magnetic field. *Astrophys. J.* **146**, 480–487 (1966).
15. Giacalone, J. & Jokipii, J. R. The transport of cosmic rays across a turbulent magnetic field. *Astrophys. J.* **520**, 204–214 (1999).
16. Lemoine, M. Particle transport through localized interactions with sharp magnetic field bends in MHD turbulence. *J. Plasma Phys.* **89**, 175890501 (2023).
17. Kempster, P. et al. Cosmic ray transport in large-amplitude turbulence with small-scale field reversals. *Mon. Not. R. Astron. Soc.* **525**, 4985–4998 (2023).
18. Amato, E. & Blasi, P. Cosmic ray transport in the Galaxy: a review. *Adv. Space Res.* **62**, 2731–2749 (2018).
19. De La Torre Luque, P. et al. Galactic diffuse gamma rays meet the PeV frontier. *Astron. Astrophys.* **672**, A58 (2023).
20. Cao, Z. et al. Measurement of ultra-high-energy diffuse gamma-ray emission of the galactic plane from 10 TeV to 1 PeV with LHAASO-KM2A. *Phys. Rev. Lett.* **131**, 151001 (2023).
21. Zhang, R., Huang, X., Xu, Z.-H., Zhao, S. & Yuan, Q. Galactic diffuse γ -ray emission from GeV to PeV energies in light of up-to-date cosmic-ray measurements. *Astrophys. J.* **957**, 43 (2023).
22. Kulsrud, R. & Pearce, W. P. The effect of wave–particle interactions on the propagation of cosmic rays. *Astrophys. J.* **156**, 445–469 (1969).
23. Skilling, J. Cosmic rays in the Galaxy: convection or diffusion?. *Astrophys. J.* **170**, 265–273 (1971).
24. Blasi, P., Amato, E. & Serpico, P. D. Spectral breaks as a signature of cosmic ray induced turbulence in the Galaxy. *Phys. Rev. Lett.* **109**, 061101 (2012).
25. Shalaby, M., Thomas, T., Pfrommer, C., Lemmerz, R. & Bresci, V. Deciphering the physical basis of the intermediate-scale instability. *J. Plasma Phys.* **89**, 175890603 (2023).
26. Chew, G. F., Goldberger, M. L. & Low, F. E. The Boltzmann equation and the one-fluid hydromagnetic equations in the absence of particle collisions. *Proc. R. Soc. Lond. A* **236**, 112–118 (1956).
27. Barnes, A. Collisionless damping of hydromagnetic waves. *Phys. Fluids* **9**, 1483–1495 (1966).
28. Hasegawa, A. Drift mirror instability of the magnetosphere. *Phys. Fluids* **12**, 2642–2650 (1969).
29. Riquelme, M. A., Quataert, E. & Verscharen, D. Particle-in-cell simulations of continuously driven mirror and ion cyclotron instabilities in high beta astrophysical and heliospheric plasmas. *Astrophys. J.* **800**, 27 (2015).
30. Ley, F., Zweibel, E. G., Miller, D. & Riquelme, M. Secondary whistler and ion-cyclotron instabilities driven by mirror modes in galaxy clusters. *Astrophys. J.* **965**, 155 (2024).
31. Rosin, M. S., Schekochihin, A. A., Rincon, F. & Cowley, S. C. A non-linear theory of the parallel firehose and gyrothermal instabilities in a weakly collisional plasma. *Mon. Not. R. Astron. Soc.* **413**, 7–38 (2011).
32. Hall, A. N. The propagation of Galactic cosmic rays. *Mon. Not. R. Astron. Soc.* **197**, 977–993 (1981).
33. Zhou, M., Zhdankin, V., Kunz, M. W., Loureiro, N. F. & Uzdensky, D. A. Magnetogenesis in a collisionless plasma: from Weibel instability to turbulent dynamo. *Astrophys. J.* **960**, 12 (2024).
34. Komarov, S. V., Churazov, E. M., Kunz, M. W. & Schekochihin, A. A. Thermal conduction in a mirror-unstable plasma. *Mon. Not. R. Astron. Soc.* **460**, 467–477 (2016).
35. Rincon, F., Schekochihin, A. A. & Cowley, S. C. Non-linear mirror instability. *Mon. Not. R. Astron. Soc.* **447**, L45–L49 (2015).
36. Melville, S., Schekochihin, A. A. & Kunz, M. W. Pressure-anisotropy-driven microturbulence and magnetic-field evolution in shearing, collisionless plasma. *Mon. Not. R. Astron. Soc.* **459**, 2701–2720 (2016).
37. Govoni, F. et al. Sardinia Radio Telescope observations of Abell 194. The intra-cluster magnetic field power spectrum. *Astron. Astrophys.* **603**, A122 (2017).
38. Squire, J. et al. Pressure anisotropy and viscous heating in weakly collisional plasma turbulence. *J. Plasma Phys.* **89**, 905890417 (2023).
39. Kalnin, J. R. & Kotomin, E. Modified Maxwell–Garnett equation for the effective transport coefficients in inhomogeneous media. *J. Phys. A* **31**, 7227–7234 (1998).
40. Kotera, K. & Lemoine, M. Optical depth of the Universe to ultrahigh energy cosmic ray scattering in the magnetized large scale structure. *Phys. Rev. D* **77**, 123003 (2008).
41. Butsky, I. S. et al. Galactic cosmic-ray scattering due to intermittent structures. *Mon. Not. R. Astron. Soc.* **528**, 4245–4254 (2024).
42. Arzamasskiy, L., Kunz, M. W., Squire, J., Quataert, E. & Schekochihin, A. A. Kinetic turbulence in collisionless high- β plasmas. *Phys. Rev. X* **13**, 021014 (2023).
43. Majeski, S., Kunz, M. W. & Squire, J. Self-organization in collisionless, high- β turbulence. Preprint at <https://arxiv.org/abs/2405.02418> (2024).
44. St-Onge, D. A. & Kunz, M. W. Fluctuation dynamo in a collisionless, weakly magnetized plasma. *Astrophys. J. Lett.* **863**, L25 (2018).
45. St-Onge, D. A., Kunz, M. W., Squire, J. & Schekochihin, A. A. Fluctuation dynamo in a weakly collisional plasma. *J. Plasma Phys.* **86**, 905860503 (2020).
46. Enßlin, T. A. On the escape of cosmic rays from radio galaxy cocoons. *Astron. Astrophys.* **399**, 409–420 (2003).
47. Dursi, L. & Pfrommer, C. Draping of cluster magnetic fields over bullets and bubbles—morphology and dynamic effects. *Astrophys. J.* **677**, 993–1018 (2008).
48. Ruszkowski, M., Enßlin, T. A., Brügggen, M., Begelman, M. C. & Churazov, E. Cosmic ray confinement in fossil cluster bubbles. *Mon. Not. R. Astron. Soc.* **383**, 1359–1365 (2008).
49. Ewart, R. J., Reichherzer, P., Bott, A. F. A., Kunz, M. W. & Schekochihin, A. A. Cosmic-ray confinement in radio bubbles by micromirrors. *Mon. Not. R. Astron. Soc.* **532**, 2098–2107 (2024).
50. Herold, L. & Malyshev, D. Hard and bright gamma-ray emission at the base of the Fermi bubbles. *Astron. Astrophys.* **625**, A110 (2019).
51. Giacintucci, S., Markevitch, M. & Clarke, T. E. uGMRT observations of the giant fossil radio lobe in the Ophiuchus galaxy cluster. *Rev. Mex. Astron. Astrofis.* **56**, 48–54 (2024).
52. Abeysekara, A. U. et al. HAWC observations of the acceleration of very-high-energy cosmic rays in the Cygnus Cocoon. *Nat. Astron.* **5**, 465–471 (2021).

53. Lhaaso Collaboration An ultrahigh-energy γ -ray bubble powered by a super PeVatron. *Sci. Bull.* **69**, 449–457 (2024).
54. Miniati, F. The Matryoshka Run. II. Time-dependent turbulence statistics, stochastic particle acceleration, and microphysics impact in a massive galaxy cluster. *Astrophys. J.* **800**, 60 (2015).
55. Brunetti, G. & Lazarian, A. Particle reacceleration by compressible turbulence in galaxy clusters: effects of a reduced mean free path. *Mon. Not. R. Astron. Soc.* **412**, 817–824 (2011).
56. Brunetti, G. & Lazarian, A. Stochastic reacceleration of relativistic electrons by turbulent reconnection: a mechanism for cluster-scale radio emission? *Mon. Not. R. Astron. Soc.* **458**, 2584–2595 (2016).
57. Bustard, C. & Oh, S. P. Turbulent reacceleration of streaming cosmic rays. *Astrophys. J.* **941**, 65 (2022).
58. Lazarian, A. & Xu, S. Mirror acceleration of cosmic rays in a high- β medium. *Astrophys. J.* **956**, 63 (2023).
59. Beduzzi, L. et al. Exploring the origins of mega radio halos. *Astron. Astrophys.* **678**, L8 (2023).
60. Nishiwaki, K. & Asano, K. Low injection rate of cosmic-ray protons in the turbulent reacceleration model of radio halos in galaxy clusters. Preprint at <https://arxiv.org/abs/2408.13846> (2024).
61. Guo, F. & Oh, S. P. Feedback heating by cosmic rays in clusters of galaxies. *Mon. Not. R. Astron. Soc.* **384**, 251–266 (2008).
62. Jacob, S. & Pfrommer, C. Cosmic ray heating in cool core clusters—I. Diversity of steady state solutions. *Mon. Not. R. Astron. Soc.* **467**, 1449–1477 (2017).
63. Jacob, S. & Pfrommer, C. Cosmic ray heating in cool core clusters—II. Self-regulation cycle and non-thermal emission. *Mon. Not. R. Astron. Soc.* **467**, 1478–1495 (2017).
64. Cuciti, V. et al. Galaxy clusters enveloped by vast volumes of relativistic electrons. *Nature* **609**, 911–914 (2022).
65. Böss, L. M., Steinwandel, U. P., Dolag, K. & Lesch, H. CRESCENDO: an on-the-fly Fokker–Planck solver for spectral cosmic rays in cosmological simulations. *Mon. Not. R. Astron. Soc.* **519**, 548–572 (2023).
66. Hopkins, P. F. et al. But what about...: cosmic rays, magnetic fields, conduction, and viscosity in galaxy formation. *Mon. Not. R. Astron. Soc.* **492**, 3465–3498 (2020).
67. Ruszkowski, M. & Pfrommer, C. Cosmic ray feedback in galaxies and galaxy clusters. *Astron. Astrophys. Rev.* **31**, 4 (2023).
68. Stuardi, C. et al. The intracluster magnetic field in the double relic galaxy cluster Abell 2345. *Mon. Not. R. Astron. Soc.* **502**, 2518–2535 (2021).
69. Bonafede, A. et al. The Coma cluster magnetic field from Faraday rotation measures. *Astron. Astrophys.* **513**, A30 (2010).
70. Steinwandel, U. P., Dolag, K., Böss, L. M. & Marin-Gilabert, T. Toward cosmological simulations of the magnetized intracluster medium with resolved coulomb collision scale. *Astrophys. J.* **967**, 125 (2024).
71. Alves Batista, R. et al. CRPropa 3.2—an advanced framework for high-energy particle propagation in extragalactic and galactic spaces. *J. Cosmol. Astropart. Phys.* **2022**, 035 (2022).
72. Mertsch, P. Test particle simulations of cosmic rays. *Astrophys. Space Sci.* **365**, 135 (2020).
73. Tautz, R. C. & Dosch, A. On numerical turbulence generation for test-particle simulations. *Phys. Plasmas* **20**, 022302 (2013).
74. Schlegel, L., Frie, A., Eichmann, B., Reichherzer, P. & Tjus, J. B. Interpolation of turbulent magnetic fields and its consequences on cosmic ray propagation. *Astrophys. J.* **889**, 123 (2020).
75. Giacinti, G., Kachelrieß, M., Semikoz, D. V. & Sigl, G. Cosmic ray anisotropy as signature for the transition from galactic to extragalactic cosmic rays. *J. Cosmol. Astropart. Phys.* **2012**, 031 (2012).
76. Kempfski, P. & Quataert, E. Reconciling cosmic-ray transport theory with phenomenological models motivated by Milky-Way data. *Mon. Not. R. Astron. Soc.* **514**, 657–674 (2022).
77. Hopkins, P. F., Squire, J., Butsky, I. S. & Ji, S. Standard self-confinement and extrinsic turbulence models for cosmic ray transport are fundamentally incompatible with observations. *Mon. Not. R. Astron. Soc.* **517**, 5413–5448 (2022).
78. Kolmogorov, A. The local structure of turbulence in incompressible viscous fluid for very large Reynolds' numbers. *Dokl. Akad. Nauk SSSR* **30**, 301–305 (1941).
79. Iroshnikov, P. S. Turbulence of a conducting fluid in a strong magnetic field. *Sov. Astron.* **7**, 566–571 (1964).
80. Kraichnan, R. H. Inertial-range spectrum of hydromagnetic turbulence. *Phys. Fluids* **8**, 1385–1387 (1965).
81. Goldreich, P. & Sridhar, S. Toward a theory of interstellar turbulence. II. Strong Alfvénic turbulence. *Astrophys. J.* **438**, 763–775 (1995).
82. Schekochihin, A. A. MHD turbulence: a biased review. *J. Plasma Phys.* **88**, 155880501 (2022).
83. Fornieri, O., Gaggero, D., Cerri, S. S., De La Torre Luque, P. & Gabici, S. The theory of cosmic ray scattering on pre-existing MHD modes meets data. *Mon. Not. R. Astron. Soc.* **502**, 5821–5838 (2021).
84. Chandran, B. D. G. Scattering of energetic particles by anisotropic magnetohydrodynamic turbulence with a Goldreich–Sridhar power spectrum. *Phys. Rev. Lett.* **85**, 4656–4659 (2000).
85. Lithwick, Y. & Goldreich, P. Compressible magnetohydrodynamic turbulence in interstellar plasmas. *Astrophys. J.* **562**, 279–296 (2001).
86. Cho, J. & Vishniac, E. T. The anisotropy of magnetohydrodynamic Alfvénic turbulence. *Astrophys. J.* **539**, 273–282 (2000).
87. Skilling, J. Cosmic ray streaming—III. Self-consistent solutions. *Mon. Not. R. Astron. Soc.* **173**, 255–269 (1975).
88. Shukurov, A., Snodin, A. P., Seta, A., Bushby, P. J. & Wood, T. S. Cosmic rays in intermittent magnetic fields. *Astrophys. J. Lett.* **839**, L16 (2017).
89. Perri, S., Pucci, F., Malara, F. & Zimbardo, G. On the power-law distribution of pitch-angle scattering times in solar wind turbulence. *Sol. Phys.* **294**, 34 (2019).
90. Kadomtsev, B. B. & Petviashvili, V. I. Acoustic turbulence. *Dokl. Phys.* **18**, 115 (1973).
91. Yan, H. & Lazarian, A. Scattering of cosmic rays by magnetohydrodynamic interstellar turbulence. *Phys. Rev. Lett.* **89**, 281102 (2002).
92. Cho, J. & Lazarian, A. Compressible sub-Alfvénic MHD turbulence in low- β plasmas. *Phys. Rev. Lett.* **88**, 245001 (2002).
93. Becker Tjus, J. & Merten, L. Closing in on the origin of Galactic cosmic rays using multimessenger information. *Phys. Rep.* **872**, 1–98 (2020).
94. Hu, Y., Lazarian, A., Li, Y., Zhuravleva, I. & Gendron-Marsolais, M.-L. Probing magnetic field morphology in galaxy clusters with the gradient technique. *Astrophys. J.* **901**, 162 (2020).
95. Brunetti, G. & Lazarian, A. Compressible turbulence in galaxy clusters: physics and stochastic particle re-acceleration. *Mon. Not. R. Astron. Soc.* **378**, 245–275 (2007).
96. Krumholz, M. R. et al. Cosmic ray transport in starburst galaxies. *Mon. Not. R. Astron. Soc.* **493**, 2817–2833 (2020).
97. Zweibel, E. G. The microphysics and macrophysics of cosmic rays. *Phys. Plasmas* **20**, 055501 (2013).
98. Kempfski, P., Quataert, E. & Squire, J. Sound-wave instabilities in dilute plasmas with cosmic rays: implications for cosmic ray confinement and the Perseus X-ray ripples. *Mon. Not. R. Astron. Soc.* **493**, 5323–5335 (2020).

99. Fabian, A. C. et al. A very deep Chandra observation of the Perseus cluster: shocks, ripples and conduction. *Mon. Not. R. Astron. Soc.* **366**, 417–428 (2006).
100. Lyskova, N. et al. X-ray surface brightness and gas density profiles of galaxy clusters up to $3 \times R_{500c}$ with SRG/eROSITA. *Mon. Not. R. Astron. Soc.* **525**, 898–907 (2023).
101. Chen, A. & Renshaw, E. The Gillis–Domb–Fisher correlated random walk. *J. Appl. Probab.* **29**, 792–813 (1992).
102. Bott, A. F. A., Arzamasskiy, L., Kunz, M. W., Quataert, E. & Squire, J. Adaptive critical balance and firehose instability in an expanding, turbulent, collisionless plasma. *Astrophys. J. Lett.* **922**, L35 (2021).
103. Eyink, G. et al. Flux-freezing breakdown in high-conductivity magnetohydrodynamic turbulence. *Nature* **497**, 466–469 (2013).
104. Aluie, H. et al. Forced MHD turbulence. *Johns Hopkins Turbulence Databases* <https://doi.org/10.7281/T1930RBS> (2019).
105. Reichherzer, P. CR diffusion in micromirrors. *Zenodo* <https://zenodo.org/records/13942018> (2024).
106. Kuhlen, M., Phan, V. H. M. & Mertsch, P. Diffusion of relativistic charged particles and field lines in isotropic turbulence. Preprint at <https://arxiv.org/abs/2211.05881> (2022).

Acknowledgements

P.R. thanks J. Becker Tjus, F. Effenberger, H. Fichtner, R. Grauer, J. Lübke and L. Schlegel for discussions on the usage of synthetic turbulence. A.A.S. thanks E. Churazov and S. Komarov for discussions of the ICM and CR physics. We thank P. Oh, E. Quataert, L. Silva, L. Turica, D. Uzdensky and especially W. Xu for discussions of CR transport in the ICM. We also acknowledge the hospitality of the Wolfgang Pauli Institute, Vienna, where these ideas were discussed during the 14th Plasma Kinetics Working Meeting (2023).

The work of P.R. was funded initially through a Gateway Fellowship and subsequently through the Walter Benjamin Fellowship by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)—grant number 518672034. A.F.A.B.'s work was supported by a UKRI Future Leaders Fellowship (grant number MR/W006723/1). R.J.E.'s work was supported by a UK EPSRC studentship. M.W.K.'s work was supported in part by NSF CAREER Award 1944972. The work of A.A.S. and G.G. was supported in part by a grant from STFC (ST/W000903/1). The work of A.A.S. was also supported by a grant from EPSRC (EP/R034737/1) and by the Simons Foundation via a Simons Investigator Award. P.K. was supported by the Lyman Spitzer, Jr. Fellowship at Princeton University. UKRI is a Plan S funder, so for the purpose of Open Access the author has applied a CC BY public copyright license to any author accepted manuscript version arising from this submission.

Author contributions

A.F.A.B., R.J.E., P.R. and A.A.S. conceptualized the study and developed the theoretical framework. A.F.A.B. and M.W.K. conducted the micromirror-field simulations, while P.R. performed the CR simulations, neural-network implementation and training, data analysis and visualization. P.R. wrote the initial draft and led the process of revision, with contributions to writing from A.F.A.B., R.J.E., M.W.K., P.K. and A.A.S. All authors provided feedback, participated in data interpretation and were involved in finalizing the manuscript. The project was supervised by A.F.A.B., G.G. and A.A.S.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Patrick Reichherzer.

Peer review information *Nature Astronomy* thanks Siang Peng Oh and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025