Stable machine-learning parameterization of subgrid processes for climate modeling at a range of resolutions

Janni Yuval and Paul O'Gorman

Support from Houghton-Lorenz Fellowship

MIT Environmental Solutions Initiative and the NSF



Outline

- Motivation: Parameterizations of small scale processes are the main cause of uncertainty in climate change temperature and precipitation projections
- Previous studies: Recent studies on machine-learning parameterization are promising (but still many problems)
- Results:
 - a. Learning from a high resolution model (SAM)
 - b. Example: Sub-grid tendencies are very important
 - c. Offline results, conservation of energy, non-negative precipitation
 - d. Online results
 - e. Is there an optimal length scale for parameterization?

Outline

- Motivation: Parameterizations of small scale processes are the main cause of uncertainty in climate change temperature and precipitation projections
- Previous studies: Recent studies on machine-learning parameterization are promising (but still many problems)
- Results:
 - a. Learning from a high resolution model (SAM)
 - b. Example: Sub-grid tendencies are very important
 - c. Offline results, conservation of energy, non-negative precipitation
 - d. Online results
 - e. Is there an optimal length scale for parameterization?

THE

LONDON, EDINBURGH, AND DUBLIN

PHILOSOPHICAL MAGAZINE

AND

JOURNAL OF SCIENCE.

[FIFTH SERIES.]

APRIL 1896.

- XXXI. On the Influence of Carbonic Acid in the Air upon the Temperature of the Ground. By Prof. SVANTE ARRHENIUS *.
 - I. Introduction: Observations of Langley on Atmospherical Absorption.

THE

LONDON, EDINBURGH, AND DUBLIN

PHILOSOPHICAL MAGAZINE

AND

JOURNAL OF SCIENCE.


[FIFTH SERIES.]

APRIL 1896.

XXXI. On the Influence of Carbonic Acid in the Air upon the Temperature of the Ground. By Prof. SVANTE ARRHENIUS *.

> 1. Introduction: Observations of Langley on Atmospherical Absorption.

Arrhenius (1896): Doubling of $CO_2 \rightarrow \text{global warming of } \sim 5^{\circ}\text{C}$



Arrhenius (1896): Doubling of $CO_2 \rightarrow \text{global warming of } \sim 5^{\circ}C$

Charney Report (1979):

"We estimate the most probable global warming for a doubling of CO₂ to be near 3°C with a probable error of ± 1.5°C"





Arrhenius (1896): Doubling of $CO_2 \rightarrow \text{global warming of } \sim 5^{\circ}C$

Charney Report (1979):

"We estimate the most probable global warming for a doubling of CO₂ to be near **3°C** with a probable error of **± 1.5°C**"


IPCC report (2013):

"Estimates of the equilibrium climate sensitivity (ECS) based on observed climate change, climate models and feedback analysis, as well as paleoclimate evidence indicate that ECS is likely in the range 1.5°C to 4.5°C with high confidence, and very unlikely greater than 6°C (medium confidence)."





IPCC AR5 **1.5-4.5°C**



Arrhenius (1896): Doubling of $CO_2 \rightarrow \text{global warming of } \sim 5^{\circ}\text{C}$

Charney Report (1979):

"We estimate the most probable global warming for a doubling of CO₂ to be near **3°C** with a probable error of **± 1.5°C**"

IPCC report **(2013)**:


"Estimates of the equilibrium climate sensitivity (ECS) based on observed climate change, climate models and feedback analysis, as well as paleoclimate evidence indicate that ECS is likely in the range 1.5°C to 4.5°C with high confidence, and very unlikely greater than 6°C (medium confidence)."






IPCC AR5 CMIP6 early

1.5-4.5°C results 2.8-5.8°C




Control climate



Extreme precipitation [mm/day]

Climate models from CMIP3
Figure adapted from O'Gorman and
Schneider 2009


There is a large spread in the intensity of extreme precipitation response to climate change



Extreme precipitation [mm/day]


Latitude
Climate models from CMIP3
Figure adapted from O'Gorman and
Schneider 2009

Climate change




Climate models from CMIP5
Figure adapted from: O'Gorman 2015

There is a large spread in the intensity of extreme precipitation response to climate change



Latitude
Climate models from CMIP3
Figure adapted from O'Gorman and
Schneider 2009

Climate change




Climate models from CMIP5


Figure adapted from: O'Gorman 2015

The large uncertainty in climate projections is mainly caused by (inaccurate) representation of subgrid physics


Law of physics (e.g., fluid dynamics)




The large uncertainty in climate projections is mainly caused by (inaccurate) representation of subgrid physics




Parameterizations represented by simplified **-** models – (e.g., convection)





Arakawa and Schubert 1974




Uncertainty in low-cloud feedback leads to large uncertainty in the equilibrium climate sensitivity









Schneider et al. (2017)







A different approach to parameterization: Use machine learning to create new parameterizations trained on high-resolution models

General Circulation Models

Law of physics (e.g., fluid dynamics)

Machine learning parameterizations (where possible)

High-resolution simulations


Outline

- Motivation: Parameterizations of small scale processes are the main cause of uncertainty in climate change temperature and precipitation projections
- Previous studies: Recent studies on machine-learning parameterization are promising (but still many problems)


Results:

- a. Learning from a high resolution model (SAM)
- b. Example: Sub-grid tendencies are very important
- c. Offline results, conservation of energy, non-negative precipitation
- d. Online results
- e. Is there an optimal length scale for parameterization?

Neural Network to learn SP-CAM (super-parameterized model) subgrid tendencies.




Neural Network to learn SP-CAM (super-parameterized model) subgrid tendencies.



SP-CAM and Neural network-CAM has similar precipitation distribution

Rasp et al. (2018)


Neural Network to learn SP-CAM (super-parameterized model) subgrid tendencies.



SP-CAM and Neural network-CAM has similar precipitation distribution

Rasp et al. (2018)

Neural Network to learn SP-CAM (super-parameterized model) subgrid tendencies.




- 1. 2D CRM with 8 gridboxes (wasn't stable with 32 gridboxes)
- 2. Changes in the Neural Network architecture can lead to unstable simulations
- 3. No energy conservation (though see Beucler et.al. 2019)


SP-CAM and Neural network-CAM has similar precipitation distribution

Rasp et al. (2018)

Neural Network to learn a 3D cloud resolving model (SAM, 4km resolution) sub-grid tendencies.




Neural Network to learn a 3D cloud resolving model (SAM, 4km resolution) sub-grid tendencies.




Neural Network to learn a 3D cloud resolving model (SAM, 4km resolution) sub-grid tendencies.

- 1. Drift of the mean climate: only short term prediction
- 2. Challenging to make it stable
- 3. No energy conservation




Random forests is a machine learning algorithm that can give a parameterization that obey conservation laws

Random Forest could emulate the relaxed Arakawa-Schubert (RAS) convection scheme



Random forests is a machine learning algorithm that can give a parameterization that obey conservation laws

Random Forest could emulate the relaxed Arakawa-Schubert (RAS) convection scheme




- 1. Stable when coupled to GCM
- 2. Conserves energy/non-negative precipitation

Goal here: to run stable and accurate climate simulations learning from a fully 3D high-resolution simulation

Outline

- Motivation: Parameterizations of small scale processes are the main cause of uncertainty in climate change temperature and precipitation projections
- Previous studies: Recent studies on machine-learning parameterization are promising (but still many problems)
- Results:
 - a. Learning from a high resolution model (SAM)
 - b. Example: Sub-grid tendencies are very important
 - c. Offline results, conservation of energy, non-negative precipitation
 - d. Online results
 - e. Is there an optimal length scale for parameterization?




We learn from a quasi-global high-resolution model


- SAM model with hypohydrostatic rescaling factor of 4 (grid spacing 12km), 48 vertical levels
- Specified SST distribution (qobs)
- Original simulation thanks to Bill Boos

SAM: *Khairoutdinov and Randall, 2003*Hypohdrostatic/DARE: e.g., *Kuang et al. 2005*


Extreme precipitation is different in high- and low-resolution simulations



Extreme precipitation is different in high- and low-resolution simulations



Privious studies: Aqua-planet simulations with different parameterization schemes lead to very different precipitation patterns



 h_L – The liquid/ice water static stability energy

 h_L – The liquid/ice water static stability energy

$$h_{\rm L} = c_p T + gz - L_c (q_c + q_r) - L_s (q_i + q_s + q_g)$$
 Liquid lce

 h_L – The liquid/ice water static stability energy

$$h_{\rm L}=c_pT+gz-L_c(q_c+q_r)-L_s(q_i+q_s+q_g)$$
 Liquid lce

 q_T – total non precipitating water mixing ratio

 h_L – The liquid/ice water static stability energy

$$h_{\rm L} = c_p T + gz - L_c (q_c + q_r) - L_s (q_i + q_s + q_g)$$
 Liquid lce

 q_T – total non precipitating water mixing ratio

 q_p – total precipitating water mixing ratio

$$\frac{\partial h_L}{\partial t}$$
 = advection + diffusion + radiation...

$$\frac{\partial q_T}{\partial t}$$
 = advection + diffusion + microphysics...

$$\frac{\partial q_p}{\partial t}$$
 = advection + precipitation + microphysics...

$$\frac{\partial h_L}{\partial t}$$
 = advection + diffusion + radiation...

$$\frac{\partial q_T}{\partial t}$$
 = advection + diffusion + microphysics...

$$\frac{\partial q_p}{\partial t}$$
 = advection + precipitation + microphysics...

$$y = \left(\frac{\partial h_L}{\partial t}, \frac{\partial q_T}{\partial t}, \frac{\partial q_p}{\partial t}\right)_{\text{subgrid}}$$

Goal: to use a Random Forest (RF) to predict the effect of subgrid processes on the resolved thermodynamic and moisture variables


$$y = \left(\frac{\partial h_L}{\partial t}, \frac{\partial q_T}{\partial t}, \frac{\partial q_p}{\partial t}\right)_{\text{subgrid}}$$


$$y = RF(x)$$


Goal: to use a Random Forest (RF) to predict the effect of subgrid processes on the resolved thermodynamic and moisture variables


$$y = \left(\frac{\partial h_L}{\partial t}, \frac{\partial q_T}{\partial t}, \frac{\partial q_p}{\partial t}\right)_{\text{subgrid}}$$


$$y = RF(x)$$












$$-rac{\overline{\partial w h_L}}{\partial z}$$

Coarse grained advection from high resolution simulation




$$-\overline{rac{\partial w h_L}{\partial z}}$$

Coarse grained advection from high resolution simulation

$$-rac{\partial \overline{w}\overline{h}_L}{\partial z}$$

Resolved advection




$$-rac{\overline{\partial w h_L}}{\partial z} - rac{\partial \overline{w} \overline{h}_L}{\partial z}$$

Coarse grained advection from high resolution simulation

Resolved advection

$$\left(\frac{\partial \overline{h}_L}{\partial t}\right)_{\text{subgrid}}^{\text{advection}} = -\left(\frac{\overline{\partial w h_L}}{\partial z} - \frac{\partial \overline{w} \overline{h}_L}{\partial z}\right)$$



$$-rac{\overline{\partial w h_L}}{\partial z} - rac{\partial \overline{w} \overline{h}_L}{\partial z}$$

Coarse grained advection from high resolution simulation

Resolved advection

$$\left(\frac{\partial \overline{h}_L}{\partial t}\right)_{\text{subgrid}}^{\text{advection}} = -\left(\frac{\overline{\partial w h_L}}{\partial z} - \frac{\partial \overline{w} \overline{h}_L}{\partial z}\right)$$

 Parameterization includes: microphysics, vertical advection, boundary-layer turbulence and radiation


Yuval & O'Gorman, Nat. Commun. (in press)

Goal: to use a Random Forest (RF) to predict the effect of subgrid processes on the resolved thermodynamic and moisture variables


$$y = \left(\frac{\partial h_L}{\partial t}, \frac{\partial q_T}{\partial t}, \frac{\partial q_p}{\partial t}\right)_{\text{subgrid}}$$

$$y = RF(x)$$

Machine-learning algorithm for parameterization: Random-forest (Breiman 2001)



Machine-learning algorithm for parameterization: Random-forest (Breiman 2001)



- Random-forest is stable and robust when implemented in a GCM
- Respects physical constraints (energy conservation, non-negative precipitation)
 O'Gorman & Dwyer 2018

$$RF(X) \rightarrow y$$

 $X = (T(z), q_T(z), q_p(z), u(z), v(z), \text{latitude})$

$$RF(X) o y$$

$$X=(T(z),q_T(z),q_p(z),u(z),v(z), ext{latitude})$$
 Latitude is a proxy for SST, albedo etc.

$$RF(X) o y$$

$$X=(T(z),q_T(z),q_p(z),u(z),v(z), ext{latitude})$$
 Latitude is a proxy for SST, albedo etc.


$$y = \left(\frac{\partial h_L(z)}{\partial t}, \frac{\partial q_T(z)}{\partial t}, \frac{\partial q_p(z)}{\partial t}, \frac{\partial$$


$$RF(X) o y$$


$$X=(T(z),q_T(z),q_p(z),u(z),v(z), {
m latitude})$$
 Latitude is a proxy for SST, albedo etc.


$$y = \left(\frac{\partial h_L(z)}{\partial t}, \frac{\partial q_T(z)}{\partial t}, \frac{\partial q_p(z)}{\partial t}, \frac{\partial$$


- Single diffusivity for all thermodynamic and moisture variables
- Flux is always down gradient















Outline


- Motivation: Parameterizations of small scale processes are the main cause of uncertainty in climate change temperature and precipitation projections
- Previous studies: Recent studies on machine-learning parameterization are promising (but still many problems)
- Results:
 - a Learning from a high resolution model (SAM)
 - b. Example: Sub-grid tendencies are very important
 - c. Offline results, conservation of energy, non-negative precipitation
 - d. Online results
 - e. Is there an optimal length scale for parameterization?

Tendency of precipitating water due to conversion of cloud water/ice and due to evaporation at 3km




subgrid tendency = coarse-grained - resolved

Tendency of precipitating water due to conversion of cloud water/ice and due to evaporation at 3km




Tendency of precipitating water due to conversion of cloud water/ice and due to evaporation at 3km




Outline

- Motivation: Parameterizations of small scale processes are the main cause of uncertainty in climate change temperature and precipitation projections
- Previous studies: Recent studies on machine-learning parameterization are promising (but still many problems)
- Results:
 - a. Learning from a high resolution model (SAM)
 - h Evample: Sub grid tandencies are very important
 - c. Offline results, conservation of energy, non-negative precipitation
 - d. Online results
 - e. Is there an optimal length scale for parameterization?


Random forest (96km horizontal grid) predicts reasonably well the subgrid tendencies in a test data set (coefficient of determination R²>0.7)




Precipitation is a diagnostic variable (we do not predict it separately in the RF)



Precipitation is a diagnostic variable (we do not predict it separately in the RF)



Random forest (almost) conserves energy in the absence of external forcing




Compared to ~92W/m² in Brenowitz and Bretherton (2019) that used neural networks (Though see Buecler et al. 2019)


Outline


- Motivation: Parameterizations of small scale processes are the main cause of uncertainty in climate change temperature and precipitation projections
- Previous studies: Recent studies on machine-learning parameterization are promising (but still many problems)
- Results:
 - a. Learning from a high resolution model (SAM)
 - b. Example: Sub-grid tendencies are very important
 - a. Offline results, conservation of energy, non-negative precipitation
 - d. Online results
 - e. Is there an optimal length scale for parameterization?

Coupling the random forest to SAM




Random-forest parameterization brings the low-resolution model simulation much closer to the high-resolution simulation







Random-forest parameterization brings the low-resolution model simulation much closer to the high-resolution simulation



Random forest parameterization leads to remarkably accurate simulation of **precipitation extremes**





Random forest parameterization leads to remarkably accurate simulation of **precipitation extremes**




Outline


- Motivation: Parameterizations of small scale processes are the main cause of uncertainty in climate change temperature and precipitation projections
- Previous studies: Recent studies on machine-learning parameterization are promising (but still many problems)
- Results:
 - a. Learning from a high resolution model (SAM)
 - b. Example: Sub-grid tendencies are very important
 - c. Offline results, conservation of energy, non-negative precipitation
 - d Opling regulta
 - e. Is there an optimal length scale for parameterization?






Conventional parameterization (non scale aware)






Conventional parameterization (non scale aware)

When increasing the grid spacing:




When increasing the grid spacing:

- Averaging over more cloud elements – subgrid tendencies more predictable
- But more of the dynamics becomes subgrid




When increasing the grid spacing:

- Averaging over more cloud elements – subgrid tendencies more predictable
- But more of the dynamics becomes subgrid




We train a different random-forest parameterization for different grid-spacing

Random-forest parameterization of subgrid tendencies becomes more accurate as grid spacing becomes coarser




But accuracy of simulations improves with decreasing grid spacing




Performance is measured by the mean precipitation distribution as compared to highresolution simulation

But accuracy of simulations improves with decreasing grid spacing




Why the discrepancy between offline and online performance?




press)

Why the discrepancy between offline and online performance?




Why the discrepancy between offline and online performance?




press)


Discrepancy between offline and online performance regardless of whether consider relative or absolute measures of error




Discrepancy between offline and online performance regardless of whether consider relative or absolute measures of error




The outputs have (a) a predictable component and (b) a stochastic component







The outputs have (a) a predictable component and (b) a stochastic component




The outputs have (a) a predictable component and (b) a stochastic component




To compare "apples-to-apples" we coarse-grain the higher resolution parameterization






To compare "apples-to-apples" we coarse-grain the higher resolution parameterization




RF with smaller coarse graining performs better in an "apples-to-apples" comparison (at the same length scale)




Coarse-grain

Conclusions

 Offline performance can be misleading in comparisons across length scales unless first coarsegrain to a reference length scale





 ML parameterization works better as we decrease grid spacing (no evidence for optimal length scale or gray zone)



Main conclusions

- Random-forest parameterization learned from fully 3-D high-resolution simulation gives stable and accurate simulations at climate-model resolution
- Machine learning can give insights into the parameterization problem (e.g., scale dependence)
- Machine learning parameterization has potential to work well in the "gray zone" of conventional parameterizations





Low resolution simulations with random forest parameterizations are faster than high resolution simulations

- 96km-RF requires x30 times less CPU time than the high resolution simulation (12km)
- 192km-RF **x120** less resources
- Without changing the time step

A different approach for Random forest parameterization

- Precipitating water (qp) is a variable that changes rapidly (due to precipitation falling with gravity)
- Cannot significantly increase the time step when correcting qp
- Want to avoid using qp as a prognostic variable

Want to avoid using precipitating water as a prognostic variable

 h_L – The liquid/ice water static stability energy

$$h_{\rm L} = c_p T + gz - L_c(q_c + q_r) - L_s(q_i + q_s + q_g)$$

 q_T – total precipitating water mixing ratio

$$y = \left(\frac{\partial h_L}{\partial t}, \frac{\partial q_T}{\partial t}, \frac{\partial q_p}{\partial t}\right)_{\text{subgrid}}$$

Want to avoid using precipitating water as a prognostic variable

 h_L – The liquid/ice water static stability energy

$$h_{\rm L} = c_p T + gz - L_c(q_e + q_r) - L_s(q_i + q_s + q_g)$$

$$H_L = c_p T + gz - L_c q_c + -L_s q_i$$

 q_T – total precipitating water mixing ratio

qp total non precipitating water mixing ratio

$$\frac{\partial H_L}{\partial t}$$
 subgrid

+ Change equation of motions in SAM

$$y = \left(\frac{\partial h_L}{\partial t}, \frac{\partial q_T}{\partial t}, \frac{\partial q_p}{\partial t}\right)_{\text{subgrid}}$$

coarse grained + <u>simplified</u> RF parameterization have similar precipitation a to a high resolution simulation

